SOCIAL INFORMATION USE SHAPES THE COEVOLUTION OF

SOCIALITY AND VIRULENCE

Ben Ashby[1,*] and Damien R Farine[2,3,4]

1. Department of Mathematical Sciences, University of Bath, Bath, BA2 7AY, UK

2. Department of Evolutionary Biology and Environmental Studies, University of Zurich, 8047 Zurich, Switzerland

3. Max Planck Institute of Animal Behavior, 78315 Radolfzell, Germany

4. Centre for the Advanced Study of Collective Behaviour, University of Konstanz, 78464 Konstanz, Germany

*Corresponding author: benashbyevo@gmail.com

Keywords: coevolution, infectious disease, social information, contact rate, sociality, virulence

ABSTRACT

Social contacts can facilitate the spread of both survival-related information and infectious disease, but little is known about how these processes combine to shape host and parasite evolution. Here, we use a theoretical model that captures both transmission processes to investigate how host sociality and parasite virulence (co)evolve. We show how selection for sociality (and in turn, virulence) depends on both the intrinsic costs and benefits of social information and disease as well as their relative prevalence in the population. Specifically, greater sociality and lower virulence evolve when the risk of infection is either low or high and social information is neither very common nor too rare. Lower sociality and higher virulence evolve when the prevalence patterns are reversed. When disease and social information are both at moderate levels, the direction of selection depends on the relative costs and benefits of being infected or informed. We also show that sociality varies inversely with virulence, and that disease may be unable to constrain the evolution of sociality. Together, these findings provide new insights for our understanding of group living and how apparently opposing ecological processes can influence the evolution of sociality and virulence in a range of ways.

INTRODUCTION

Animals exhibit strong variation in sociality (contact rates), both within (Watters and Sih 2005; Pike et al. 2008; Croft et al. 2009; Tanner and Jackson 2012; Aplin et al. 2013, 2015b; Weber et al. 2013) and between (Sachser 1986; Barton et al. 1996) populations. Some individuals have relatively few, potentially stable associations (e.g. non-gregarious species or those that form pair bonds) while others have a greater number of interactions that may be more transient (e.g. herding behaviour or roaming individuals) (Couzin 2006; Silk et al. 2014; Farine et al. 2015b). Crucially, what constitutes a good social strategy under one set of ecological conditions may be a poor choice under different circumstances (Krause and Ruxton 2002), hence it is important to understand how changes in the environment shape social behaviour. Depending on the ecological circumstances, sociality may be associated with various costs (e.g. higher risk of disease or greater competition for resources) and benefits (e.g. information exchange, increased mating opportunities, or symbiont transmission), which trade-off to determine whether selection favours increased sociality.

Intuitively, if sociality confers diminishing benefits but accelerating costs, then the population will evolve to an evolutionarily stable level of sociality. This is straightforward when costs and benefits can be considered fixed, so that a given level of sociality always incurs the same cost or provides the same benefit. For example, living in a larger groups may lead to foraging benefits (Cantor et al. 2020) or a reduced risk of predation (Kenward 1978), but these do not necessarily depend on the sociality of other individuals in the population. However, when the costs and benefits are dynamic, as is the case with infectious diseases, social information and symbiont transmission, feedbacks may exist

between sociality and the likelihood of realising an associated cost or benefit (Cantor et al.

2021). For example, more social members of a population may experience greater exposure

to disease (Lloyd-Smith et al. 2005; Eames and Keeling 2006; Ashby and Gupta 2013) and

access to social information (Aplin et al. 2012, 2015a; Snijders et al. 2021) than those that

form fewer contacts or have more stable associations (Evans et al. 2020). However, the

sociality of the population will affect the prevalence of disease or social information in the

population, and therefore the likelihood of incurring a cost or realising a benefit of sociality

(Romano et al. 2020). Thus, while we should expect infectious disease and social

information to influence the evolution of sociality, we must also consider how the evolution

of sociality shapes the dynamics of these cost/benefit processes due to eco-evolutionary

feedbacks. An outstanding question, which we address in the present study, is how dynamic

costs (in terms of disease transfer) and benefits (in terms of information transfer) mediate

the evolution of sociality.

Few theoretical studies consider how eco-evolutionary feedbacks mediate the evolution of

sociality (see Cantor et al. 2020b for a review), but those that do have thus far focused on

the role of infectious diseases as a dynamic cost. Bonds et al. (2005) showed that sociality

(contact rates) is minimised for intermediate levels of disease, and therefore higher levels of

disease can select for greater sociality. This is because there is little chance of avoiding

disease when it is very common, and so the marginal costs of greater sociality are low. In a

related study, Prado et al. (2009) used a stochastic individual-based network model to show

that coevolution between host contact number and parasite virulence can lead to

fluctuations in both traits, with the cost of sociality depending both on the virulence of the

parasite and its prevalence in the population. These studies demonstrate the importance of

infectious diseases as dynamic costs in the evolution of sociality, but both assume that the

benefits of sociality are fixed, so that each additional contact increases host fitness by a

predetermined amount. Theoretical studies have yet to consider dynamic rather than fixed

benefits of sociality.

One of the most prominent examples of a dynamic benefit of sociality is social information,

which can provide individuals with knowledge about ephemeral resources or conditions,

such as foraging or nesting sites, or predation risk (Doligez et al. 2002; Danchin et al. 2004;

Valone 2007). This information can be transmitted through active signalling (Elgar 1986) or

via inadvertent cues (Pöysä 1992; Galef and Giraldeau 2001), with highly social individuals

likely to receive information sooner than poorly connected members of the population

(Aplin et al. 2012, 2015a; Snijders et al. 2021). We should therefore expect sociality to

increase access to survival-related information, but as is the case with disease, the

prevalence of information in the population will depend on sociality, forming an eco-

evolutionary feedback that causes the evolution of sociality to diverge from the fixed-

benefit scenario.

While social information and infectious diseases are both transmission processes that share

many similarities, their dynamics may occur over similar or very different timescales, which

may affect selection on social strategies. Social information about predation or resource

availability is expected to be of high value that rapidly decays over time (between seconds

and days), whereas the risk of infectious disease may vary more slowly and over longer

periods of time (weeks, months or years). For example, the life cycle of a parasite can de-

couple the transmission from social contacts by several weeks (Grear et al. 2013). In

general, we should expect social information dynamics to be relatively fast compared to

infectious disease dynamics, although it is possible that they may occur on similar

timescales. The success of a particular social strategy is likely to depend on the relationship

between these timescales, as well as wider contact patterns in the population, the value of

social information and the prevalence and severity of disease. Environmental changes (e.g.

food availability, pathogen transmissibility or virulence) therefore have the potential to shed

light on how sociality evolves.

Here, we examine how social information mediates the evolution of sociality and virulence.

We present a mathematical model of infectious disease and social information dynamics,

allowing sociality to evolve with no inherent costs or benefits, and virulence to evolve

subject to a trade-off with transmission. Evolution is determined purely by balancing access

to social information, which reduces the mortality rate, and infectious disease, which

increases the mortality rate. We consider two versions of the model, where social

information dynamics occur on relatively fast or slow timescales, with the former allowing

for a simple approximation to social information prevalence. We show that selection

favours increased sociality when social information is neither very common nor too rare and

disease is at either high or low levels. When social information and disease are both at

intermediate levels, selection may favour increased or decreased sociality depending on the

relative costs and benefits of acquiring social information and disease, the relative

timescales of the transmission processes, as well as the sociality of other members of the

population. As a result, we show how ecological conditions can determine if: (1) the

population evolves towards an evolutionarily stable level of sociality; (2) disease is unable to

constrain selection for sociality; or (3) if selection drives social information and disease

'extinct'.

MODEL DESCRIPTION

FULL MODEL

We model the (co)evolution of sociality and virulence due to the transmission of social

information and infectious disease (Fig. 1). We assume that the environment is constantly

changing (e.g., food sites may be ephemeral, predation risk may vary over time and space)

and that individuals can learn about the environment independently and through social

cues. For simplicity, individuals either have poor (P) or good (G) information about the

environment. Those with poor information have a baseline mortality rate of d (on average),

and those with good information have mortality rate ad, where $0 \leq a < 1$. All individuals

continuously learn about the environment at a constant (arbitrary) rate, but the extent to

which an individual can learn independently is assumed to be limited, corresponding to a

relatively poor information state (P). An individual in a poor information state can only

improve to a good information state (G) through social cues from an individual who is

better informed about the environment, which they do with probability τ given social

contact. Individuals are poor at integrating information, such that multiple individuals with

poor information cannot come together to produce good information—good information

can only be socially learnt. For example, the location of a highly camouflaged predator may

(almost never) be individually discovered, but is readily socially learnt, and that information

is only valid while the individual is foraging nearby the predator. Since the environment is

constantly changing, and since individual priorities may change (e.g., an individual may seek

shelter once satiated, seek a different type of food, or simply leave the area), socially

learned information loses value to an individual over time (but may still be valuable to

others). For simplicity, individuals in the good information state revert to the poor state of

information at a constant rate, σ.

In addition to their information status, individuals are classed as either susceptible (S) or

infectious (I). Infected hosts transmit disease to susceptible social contacts with probability

β per social contact. Infected hosts recover at rate γ, at which point they become

susceptible again. Infection is detrimental and leads to an increase in the mortality rate due

to virulence, α.

We model social contacts following Bonds et al. (2005). Social contacts are assumed to be

ephemeral, random, and occur based on contact 'effort'. Each individual, i, has a contact

effort, E_i, which is assumed to be a heritable quantitative trait. An individual's contact

frequency, $C_i = E_i \bar{E}$, is the product of its own contact effort and the average contact effort

of the population $\left(\bar{E} = \frac{1}{N} \sum_{j=1}^{N} E_j \right)$. We assume that the only benefits and costs associated

with contact effort (sociality) are access to survival-related information and the risk of

infection, respectively.

In a monomorphic population with $E_i = E$ there are four classes (S_P, S_G, I_P, I_G), where S_i

and I_i refer to susceptible and infected individuals in information state $i \in \{P, G\}$ and the

total population size is $N = S_P + S_G + I_P + I_G$. The per-capita birth rate is equal to $(b -$

$qN)$, where b is the maximum birth rate and q is the strength of density-dependence

(resource competition). The per-capita information transmission rate is equal to the product

of the information transmission probability per contact, τ, the population contact

frequency, E^2, and the proportion of individuals in state G, $\frac{1}{N}(S_G + I_G)$. Similarly, the

disease transmission rate is equal to the product of the disease transmission probability per

contact β, the population contact frequency, E^2, and the proportion of individuals who are

infected, $\frac{1}{N}(I_P + I_G)$. The population dynamics are described by the following set of

ordinary differential equations (ODEs):

167
$$\frac{dS_P}{dt} = (b - qN)N - \frac{E^2 S_P}{N}\big(\beta(I_P + I_G) + \tau(S_G + I_G)\big) - dS_P + \gamma I_P + \sigma S_G \quad (1a)$$

168
$$\frac{dS_G}{dt} = \frac{E^2}{N}\big(\tau S_P(S_G + I_G) - \beta S_G(I_P + I_G)\big) - adS_G + \gamma I_G - \sigma S_G \quad (1b)$$

169
$$\frac{dI_P}{dt} = \frac{E^2}{N}\big(\beta S_P(I_P + I_G) - \tau I_G(S_G + I_G)\big) - (d + \alpha + \gamma)I_P + \sigma I_G \quad (1c)$$

170
$$\frac{dI_G}{dt} = \frac{E^2}{N}\big(\beta S_G(I_P + I_G) + \tau I_G(S_G + I_G)\big) - (ad + \alpha + \gamma + \sigma)I_G \quad (1d)$$

(a) Model schematic   (b) Social learning by B from A   (c) Information loses value to A

Figure 1. Model schematic and illustration of social information dynamics. (a) Transition diagram with per-

capita rates of transition between states, as described in the main text. (b-c) Illustration of social information

dynamics. (b) An individual, B, in the "poor" information state (black) learns socially transmitted information

about the environment from individual A, who is better informed (i.e., in the "good" information state, blue).

(c) Information loses value to individual A (expires), but remains valuable to individual B (e.g., individual A may

be satiated or may leave the area).

FAST SOCIAL INFORMATION APPROXIMATION

When the dynamics of social information are much faster than demographic ($\tau E^2, \sigma \gg$

b, d) and epidemiological processes ($\tau E^2 \gg \beta, \sigma \gg \gamma$), the proportion of individuals in the

good information state for a given value of E, G_E, rapidly reaches equilibrium. Letting $G_E =$

$\frac{1}{N}(S_G + I_G)$, we see that:

185
$$\frac{dG_E}{dt} = \frac{1}{N}\left(\frac{dS_G}{dt} + \frac{dI_G}{dt}\right)$$

186
$$\approx (\tau E^2(1 - G_E) - \sigma)G_E = 0 \tag{2}$$

187   Thus, either no individuals are in information state G, or $G_E^* = 1 - \frac{\sigma}{\tau E^2}$ when $\sigma < \tau E^2$. We

188   can therefore make the following approximation to the full model when social information

189   dynamics are relatively fast:

193
$$\frac{dS}{dt} = (b - qN)N - \frac{\beta E^2 SI}{N} - d(1 - (1 - a)G_E^*)S + \gamma I \qquad (3a)$$

194
$$\frac{dI}{dt} = \frac{\beta E^2 SI}{N} - (d(1 - (1 - a)G_E^*) + \alpha + \gamma)I \qquad (3b)$$

where $N = S + I$ and $G_E^* = \max\left(1 - \frac{\sigma}{\tau E^2}, 0\right)$ is the proportion of individuals in information

state G. Note that G_E^* is an increasing, concave downward function of E, which means that

there are diminishing effects of sociality on mortality.

ANALYSIS

We use evolutionary invasion analysis (adaptive dynamics; Geritz et al. 1998) to determine

how the host and parasite traits (co)evolve. For the full model (equation 1), we derive host

fitness analytically (see *Supplementary Material* for details), but solve the system

numerically, as it is not possible to find an expression for the coexistence equilibrium where

disease and social information are both present. For the fast social information

approximation (equation 3), it is possible to find an expression for the coexistence

equilibrium, and in turn, expressions for fitness and fitness gradients. However, we omit

these expressions as they are lengthy and provide no analytical insights, and so we present

numerical results instead. We illustrate the different dynamics using evolutionary

simulations as described in previous work (Ashby and Bruns 2018). Briefly, every T time

units of the deterministic ODE solver, a rare mutant is introduced which is phenotypically

similar to a randomly chosen member of the population. Phenotypes which fall below an

arbitrary frequency, ϵ, are removed from the population, and the process is repeated. The

simulations relax the adaptive dynamics assumptions of continuous traits and a complete

separation of ecological and evolutionary time scales (source code available in the

Supplementary Material and at github.com/ecoevogroup/Ashby_and_Farine_2020).

ECOLOGICAL DYNAMICS

In the full model, the host population has a trivial equilibrium at $S_P^* = N^* = (b - d)/q$ in

the absence of disease or social information, which exists provided $b > d$ (hereafter we

assume this condition is always satisfied). This equilibrium is stable if:

$$\hat{R}_0^I = \frac{\tau E^2}{ad + \sigma} < 1 \tag{4}$$

$$\hat{R}_0^D = \frac{\beta E^2}{d + \alpha + \gamma} < 1 \tag{5}$$

where $\hat{R}_0^I$ and $\hat{R}_0^D$ are the basic reproductive ratios for information and disease at the trivial

equilibrium, respectively. The basic reproductive ratios are the average number of

transmission events per informed or infected individual in an otherwise uninformed or

susceptible population. If $\hat{R}_0^I > 1$ then social information can spread in the absence of

disease and if $\hat{R}_0^D > 1$ then disease can spread in the absence of social information.

The full model also has three non-trivial equilibria: (i) information-only (viable if $\hat{R}_0^I > 1$), (ii)

disease-only (viable if $\hat{R}_0^D > 1$), and (iii) both information and disease present (coexistence

equilibrium). The latter is viable if $R_0^I > 1$ at the disease-only equilibrium and $R_0^D > 1$ at the

information-only equilibrium, where R_0^I and R_0^D are the full basic reproductive ratios for

information and disease:

228
$$R_0^I = \frac{\tau E^2 \left(\Gamma_\gamma^I I_P + S_P (\Gamma_\alpha^I + \gamma) \right) N + \beta E^2 I_P (I_P + S_P)}{N \left((ad + \sigma)(\Gamma_\alpha^I + \gamma) N + \beta E^2 I_P \Gamma_\alpha^I \right)} \tag{6}$$

229
$$R_0^D = \frac{\beta E^2 \left((\Gamma_D + \sigma) S_G + S_P (\Gamma_\alpha^I + \gamma) \right) N + \tau E^2 S_G (S_P + S_G))}{N (\Gamma_D (\Gamma_\alpha^I + \gamma) N + \tau E^2 S_G \Gamma_\alpha^I)} \tag{7}$$

with $\Gamma_\alpha^I = ad + \alpha + \sigma$, $\Gamma_\gamma^I = ad + \gamma + \sigma$ and $\Gamma_D = d + \alpha + \gamma$ (see *Supplementary Material*

for derivation). Note that in the special case where there is either no disease or no social

information, equations 6-7 reduce to equations 4-5. Intuitively, for the coexistence

equilibrium to be stable, we require $R_0^I > 1$ at the disease-only equilibrium and $R_0^D > 1$ at

the information-only equilibrium (i.e. social information and disease can always spread

when rare). While it is not possible to obtain an analytical expression for the coexistence

equilibrium, a numerical parameter sweep indicates that it is always stable when it is viable.

The fast social information approximation has two equilibria. There is a disease-free

equilibrium at $S^* = \frac{b - d(1 - (1-a)G_E^*)}{q}$ and a coexistence equilibrium at:

239
$$\frac{S^*}{N^*} = \frac{d(1 - (1-a)G_E^*) + \alpha + \gamma}{E^2 \beta} \tag{8}$$

240
$$\frac{I^*}{N^*} = \frac{E^2 \beta - d(1 - (1-a)G_E^*) - \alpha - \gamma}{E^2 \beta} \tag{9}$$

241   where

242
$$N^* = \frac{\alpha^2 + \alpha(d(1 - (1-a)G_E^*) + \gamma - E^2\beta) + E^2\beta(b - d(1 - (1-a)G_E^*))}{E^2 \beta q} \tag{10}$$

243 The corresponding basic reproductive ratios for the fast social information approximation

244 are:

$$\tilde{R}_0^I = \frac{\tau E^2}{\sigma} \tag{11}$$

$$\tilde{R}_0^D = \frac{\beta E^2}{d(1 - (1-a)G_E^*) + \alpha + \gamma} \tag{12}$$

(see *Supplementary Material*). The coexistence equilibrium is stable when $\tilde{R}_0^D > 1$ and is

unstable when $\tilde{R}_0^D < 1$ (the opposite is true for the disease-free equilibrium). Note that

when $\tilde{R}_0^I < 1$ we must have $G_E^* = 0$ (social information transmission cannot be sustained),

and hence $\tilde{R}_0^D = \hat{R}_0^D$.

The basic reproductive ratios for the full model and the approximation are very similar

provided social information dynamics are sufficiently fast ($\tau \gg \beta$; Fig. 2a-b). When

information dynamics occur on slower timescales, then the approximation diverges from

the full model (Fig. 2a-b). Likewise, as the information dynamics become increasingly fast

relative to the disease dynamics, the equilibrium population distributions in the full model

approach that of the approximation (Fig. 2c-d).

Figure 2. Ecological dynamics. (a)-(b) Basic reproductive ratios for (a) social information ($k = I$) and (b) disease ($k = D$) as a function of the contact effort, E. Curves correspond to the basic reproductive ratio in the full model, R_0^k (black, solid), the full model without disease or social information present, $\hat{R}_0^k$ (blue, dotted), and the approximation for fast social information dynamics, $\tilde{R}_0^k$ (red, dashed). (c)-(d) Comparison of the non-trivial equilibria in the full model (solid) and the approximation for fast social information (dashed). All parameters except for τ and σ are fixed, with $\sigma/\tau = 10$ held constant. The full model approaches the approximation when social information dynamics are fast relative to other processes. (c) Proportion of individuals in each class. (d) Proportion of individuals that are infected (green) or are in the good information state (purple). Fixed parameters (where applicable): $a = 0.2, b = 1, d = 0.5, E = 4, q = 10^{-3}, \alpha = 0.4, \beta = 0.1, \gamma = 0.2, \sigma = 1, \tau = 0.1$.

EVOLUTION OF SOCIALITY

We explore the evolution of host sociality in terms of the contact effort, E. We first investigate how the fitness gradient varies for a fixed value of E to illustrate how social information and disease prevalence influence the direction of selection. We then focus our analysis on the extent to which social information reduces mortality (a), the cost of being infected (α), the rate at which social information loses value to an individual (expires)

273 relative to its transmission probability (σ/τ), and the relative transmission probabilities of

274 social information and disease (τ/β).

275 To explore host evolution, we consider the invasion of rare host mutant with contact effort

276 $E_m \approx E$ into an established resident population at equilibrium (denoted by asterisks). In the

277 full model, the invasion dynamics of the rare host mutant are given by:

278
$$\frac{dS_P^m}{dt} = (b - qN^*)N_m - \frac{E_m E S_P^m}{N^*}\left(\beta(I_P^* + I_G^*) + \tau(S_G^* + I_G^*)\right) - dS_P^m + \gamma I_P^m + \sigma S_G^m \quad (13a)$$

279
$$\frac{dS_G^m}{dt} = \frac{E_m E}{N^*}\left(\tau S_P^m(S_G^* + I_G^*) - \beta S_G^m(I_P^* + I_G^*)\right) - adS_G^m + \gamma I_G^m - \sigma S_G^m \quad (13b)$$

280
$$\frac{dI_P^m}{dt} = \frac{E_m E}{N^*}\left(\beta S_P^m(I_P^* + I_G^*) - \tau I_P^m(S_G^* + I_G^*)\right) - (d + \alpha + \gamma)I_P^m + \sigma I_G^m \quad (13c)$$

281
$$\frac{dI_G^m}{dt} = \frac{E_m E}{N^*}\left(\beta S_G^m(I_P^* + I_G^*) + \tau I_P^m(S_G^* + I_G^*)\right) - (ad + \alpha + \gamma + \sigma)I_G^m \quad (13d)$$

282 For the fast social information approximation, we let $G_E^m = \frac{1}{N_m}(S_G^m + I_G^m)$ so that:

284
$$\frac{dG_E^m}{dt} = \frac{1}{N_m}\left(\frac{dS_G^m}{dt} + \frac{dI_G^m}{dt}\right)$$

285
$$\approx (E_m E\tau(1 - G_E^m)G_E^* - \sigma G_E^m)N_m = 0 \quad (14)$$

283 Rearranging and substituting $G_E^* = 1 - \frac{\sigma}{\tau E^2}$ gives:

286
$$G_E^m = \max\left(\frac{E_m(\tau E^2 - \sigma)}{E^2 E_m \tau + \sigma(E - E_m)}, 0\right) \quad (15)$$

287 We can therefore approximate the invasion dynamics of a rare mutant when social

288 information dynamics are fast by:

289
$$\frac{dS_m}{dt} = (b - qN^*)N_m - \frac{\beta E_m E S_m I^*}{N^*} - d(1 - (1-a)G_E^m)S_m + \gamma I_m \qquad (16a)$$

290
$$\frac{dI_m}{dt} = \frac{\beta E_m E S_m I^*}{N^*} - (d(1 - (1-a)G_E^m) + \alpha + \gamma)I_m \qquad (16b)$$

In the *Supplementary Material*, we show the derivation for invasion fitness in both versions

of the model.

HOST FITNESS GRADIENT

Fig. 3 shows how the host fitness gradient varies with social information and disease

prevalence when the benefits of sociality are relatively high (Fig. 3a) or low (Fig. 3b)

compared to the costs. Social information dynamics are assumed to be fast, but the overall

patterns hold for slow information dynamics. Note that here E is fixed, with σ and β

modulating social information and disease prevalence, but when E evolves (see *Host*

evolutionary dynamics) it will feedback to affect prevalence and hence selection for

sociality. Nevertheless, considering how the fitness gradient varies when E is fixed allows us

to qualitatively see how changes in social information and disease prevalence affect the

fitness gradient, all else being equal.

The fitness gradient is always positive when social information prevalence is intermediate

and disease is either rare or common, regardless of whether the benefits of sociality are

relatively high or low compared to the costs. Intuitively, increased sociality is likely to be

beneficial when disease prevalence is low, but only if social information is neither too rare

nor very common. This is because greater sociality does not significantly increase the rate at

which individuals obtain social information if it is readily available or unlikely to become

accessible. Somewhat counter-intuitively, increased sociality may also be beneficial when

disease prevalence is high (as shown in Bonds *et al.*, 2005). This is because most individuals

are likely to be infected anyway, but increased sociality will improve access to social

information with little increase to the risk of infection. Conversely, the fitness gradient is

always negative when social information is either rare or very common and disease is at

intermediate prevalence. This pattern mirrors the one described above, but with the roles

of social information and disease switched due to their opposing effects on the fitness

gradient. Intuitively the fitness gradient is negative when social information prevalence is

low and disease is common in the population, as greater sociality increases the risk of

infection but does not substantially increase the likelihood of obtaining social information.

But perhaps counter-intuitively the fitness gradient is also negative when social information

is widespread and disease is at moderate levels. This is because most individuals are well-

informed, and so increased sociality will only marginally improve access to social

information but may substantially increase the risk of infection.

When both social information and disease are at intermediate prevalence, the sign of the

fitness gradient depends on the relative costs and benefits of being infected or informed.

When awareness of social information is relatively more valuable than the cost of being

infected, the fitness gradient is positive whenever social information prevalence is at an

intermediate level regardless of disease prevalence (Fig. 3a). Disease may therefore not

always be sufficient to curtail the evolution of sociality. In contrast, when awareness of

social information is less valuable, and the cost of being infected is relatively high, the

fitness gradient is negative whenever disease prevalence is at an intermediate level

regardless of social information prevalence (Fig. 3b). Our qualitative analysis of the fitness

gradient suggests that the host population may experience different evolutionary outcomes

depending on the initial conditions: if disease prevalence is sufficiently low the population

may evolve towards a stable level of sociality, but if disease is initially very common then

the population may experience runaway selection for greater sociality.

Figure 3. Effects of social information and disease prevalence on the host fitness gradient. Contour plots of

the host fitness gradient, in the fast social information approximation as a function of social information and

disease prevalence with: (a) higher ($a = 0, \alpha = 0.3$) and (b) lower ($a = 0.2, \alpha = 0.4$) benefits of sociality

relative to costs. All parameters except for σ and β are fixed within each panel, with values of σ and β chosen

so that social information and disease prevalence vary accordingly. In both cases, the host fitness gradient is

positive (bluer shading) when social information prevalence is intermediate and there is either a high or low

level of disease, and is negative (redder shading) when social information prevalence is either at a high or low

level and disease is at intermediate prevalence. Fixed parameters as in Fig. 2, except $\beta = 0.2$.

HOST EVOLUTIONARY DYNAMICS

When the rate at which social information loses value to an individual (the expiry rate), σ, is

not too large relative to the information transmission probability, τ, there may exist a

continuously stable strategy or CSS, which is locally attracting (dark blue curves in Fig. 4).

Nearby trait values evolve towards the CSS by small mutations and remain there indefinitely

(Fig. 5a-b). When the costs (α) that arise from being social are relatively low compared to

the benefits of being better informed (a), the CSS increases with σ/τ until there is no

evolutionarily stable contact effort, leading to runaway selection for higher values of E (Fig.

4a, d). When the costs of sociality are higher, the CSS peaks at intermediate values of σ/τ

(e.g. Fig. 4b). This is because as σ/τ increases, social information expires more quickly and

so the overall prevalence of social information in the population falls, which increases

selection for sociality. However, as sociality increases so too does disease prevalence, which

reduces selection for sociality. Thus, if the costs of being more social are relatively low, the

population may experience runaway selection, but as the costs increase due to higher

disease prevalence, sociality is curtailed at a stable intermediate level.

Two possibilities exist for initial trait values above the CSS. For sufficiently low values of σ/τ

(social information is common) and higher costs of disease (α) or lower benefits of social

information (a), the CSS, E^*, is attracting for all values of $E > E^*$. This means that high

initial levels of sociality will always evolve to a lower level. But for sufficiently large values of

σ/τ (social information is rare) or low costs of disease (α) and high benefits of social

information (a), a *repeller* (dark red curves in Fig. 4) may exist above the CSS. Contact effort

evolves towards the CSS when the initial value is below the repeller, which reduces infection

and social information prevalence to a point where the costs and benefits of sociality are

balanced (Fig. 5). There is runaway selection for greater sociality when the contact effort

starts above the repeller, which increases infection and social information prevalence (Fig.

5c). Runaway selection occurs because most individuals in the population are infected, and

so there is little cost to increased sociality. Runaway selection for sociality is unrealistic,

however, as contact efforts would be constrained by other factors (e.g. non-disease

associated costs or physical constraints). Instead, runaway selection for sociality in the

model should be interpreted as disease being insufficient to constrain sociality.

  Figure 4. The evolution of sociality arising from social information and disease transmission. Evolution of

contact effort, E, as a function of the social information expiry rate divided by the transmission probability,

σ/τ, and for different values of the benefits of social information, a, and costs of disease, α. Results are shown

for relatively slow (solid) and fast (dotted) social information dynamics. Red and dark blue curves correspond

to repellers and continuously stable strategies (CSSs), respectively. Pink and light blue curves show thresholds

below which neither social information nor disease transmission is viable. Above the pink curve, sociality

increases and above the light blue curve sociality decreases. Arrows indicate the direction of selection when

social information dynamics are relatively slow. Grey curves do not indicate singular strategies but are

continuations of the disease and social information viability thresholds when $N = S_P$. Crosses in (e)

correspond to simulations in Fig. 5 and the asterisk in (d) corresponds to simulations in Fig. 6. Fixed

parameters as in Fig. 2, except $\beta = 0.2$.

Figure 5. Simulation trajectories when social information and disease dynamics occur on similar timescales.

Trajectories of simulations corresponding to crosses in Fig. 4e. Top row: evolutionary trajectories (solid black); singular strategies: repellers (red) and continuously stable strategies (CSSs; dark blue); and thresholds below which neither social information nor disease transmission is viable (pink and light blue). Above the pink dashed line, sociality increases, and above the light blue line sociality decreases. Bottom row: proportion of individuals that are infected (green) or are in the good information state (purple) in each simulation. Fixed parameters as in Fig. 2, except $\tau = \beta = 0.2$.

For sufficiently large values of σ/τ, the threshold for sustained social information transmission passes the threshold for sustained disease transmission ($R_0^D > R_0^I$ or $\tilde{R}_0^D > \tilde{R}_0^I$). Below these thresholds neither social information nor disease can be sustained and so the fitness gradient is zero (i.e., no selection on the contact effort). For the reasons discussed above, a repeller may also exist in this region of parameter space, where for sufficiently high contact efforts, there is runaway selection for greater sociality. Increasing the costs of disease (α) or decreasing the benefits of social information (a) limits where the repeller exists to larger values of σ/τ and greater contact efforts. This has the effect of increasing the region of attraction for the CSS. Conversely, decreasing the costs of disease or increasing

the benefits of social information may cause the repeller to collide with the CSS at

intermediate values of σ/τ (Fig. 4a, d, g). In this region, disease is less costly and

information more beneficial, so as soon as social information is able to spread there is

runaway selection for greater sociality.

Thus far, we have only discussed the evolution of sociality when social information and

disease dynamics occur on similar timescales ($\sigma \approx \gamma$). When social information dynamics

occur on a much faster timescale ($\sigma \gg \gamma$), there are some notable differences (dotted

curves in Fig. 4). First, the viability thresholds (pink and light blue curves) for disease and

social information are shifted to slightly lower contact efforts. Second, the repeller curve for

fast social information dynamics is shifted to the right and down in Fig. 4, which may

increase or decrease the domain of attraction for the CSS depending on the value of σ/τ.

Third, the collision of the CSS and repeller occurs for larger costs of disease (α) and smaller

benefits of social information (a). Finally, at low cost of disease, a CSS may exist which is

absent in the full model (e.g. Fig. 4d). This means that when social information dynamics are

relatively slow, there may be runaway selection for greater sociality, but when the social

information dynamics are relatively fast, sociality may evolve to a stable level (Fig. 6). The

timescale of social information dynamics can therefore play an important role in

determining the evolution of sociality.

Figure 6. Simulation trajectories when social information dynamics occurs on similar (a) or faster (b)

timescales than disease dynamics. Simulations corresponding to the asterisk in Fig. 4d. Top row: evolutionary

trajectories (solid black); continuously stable strategies (CSSs; dark blue); and viability threshold for social

information (pink). Bottom row: proportion of individuals that are infected (green) or are informed by social

information (purple) in each simulation. Fixed parameters as in Fig. 2, except $\alpha = 0.3$, $\beta = 0.2$, $\sigma/\tau = 2$.

EVOLUTION OF VIRULENCE

To explore parasite evolution, we assume there is a trade-off between transmission and

virulence such that $\beta = \beta(\alpha)$ (i.e. more transmissible parasites are more virulent). We

consider the invasion of rare parasite mutant with virulence $\alpha_M \approx \alpha$ into an established

resident population at equilibrium (denoted by asterisks). For simplicity we assume that

social information dynamics are fast (i.e. we use the approximation to the full model). The

invasion dynamics of the rare parasite mutant are then given by:

$$\frac{dI_M}{dt} = \frac{\beta(\alpha_M)E^2 S^* I_M}{N^*} - (d(1 - (1-a)G_E^*) + \alpha_M + \gamma)I_M \qquad (17)$$

434   In the *Supplementary Material*, we derive the invasion fitness and show that evolution

435   maximises the basic reproductive ratio, $\tilde{R}_0^D(\alpha)$:

436
$$\tilde{R}_0^D(\alpha) = \frac{\beta(\alpha)E^2}{d(1 - (1 - a)G_E^*) + \alpha + \gamma} \tag{18}$$

We now consider how the evolution of sociality might feedback to affect the evolution of

parasite virulence. In the absence of a trade-off with another life-history trait, the parasite

will evolve to the lowest level of virulence possible, as this increases the infectious period.

We therefore assume, as is common in the literature (Alizon 2008), that there is a positive

relationship between virulence and the transmission probability with diminishing returns, so

that $\frac{d\beta}{d\alpha} > 0$ and $\frac{d^2\beta}{d\alpha^2} < 0$. The optimal level of virulence, α^*, occurs when:

443
$$\left.\frac{d\beta}{d\alpha}\right|_{\alpha=\alpha^*} = \frac{\beta(\alpha^*)E^2\tau}{E^2\tau(ad + \alpha^* + \gamma) + d\sigma(1 - a)} = g(\alpha^*, E)\beta(\alpha^*) \tag{19}$$

444   To determine what happens to the optimal level of virulence when sociality changes,

445   consider $y(\alpha, E) = g(\alpha, E)\beta(\alpha) - \frac{d\beta}{d\alpha}$ and suppose $\alpha = \alpha^*$, so that $y(\alpha^*, E) = 0$. Note that

446    $g(\alpha, E)$ is a decreasing function of α and is an increasing function of E provided $0 \leq a < 1$.

447   Hence, $y(\alpha, E)$ is also an increasing function of E. Now consider the derivative of y with

448   respect to α at $\alpha = \alpha^*$:

449
$$\left.\frac{\partial y}{\partial \alpha}\right|_{\alpha=\alpha^*} = \beta(\alpha^*)\left.\frac{dg}{d\alpha}\right|_{\alpha=\alpha^*} + g(\alpha^*, E)\left.\frac{d\beta}{d\alpha}\right|_{\alpha=\alpha^*} - \left.\frac{d^2\beta}{d\alpha^2}\right|_{\alpha=\alpha^*}$$

450
$$= -\left.\frac{d^2\beta}{d\alpha^2}\right|_{\alpha=\alpha^*} > 0 \tag{20}$$

where $\left.\frac{d\beta}{d\alpha}\right|_{\alpha=\alpha^*} = g(\alpha^*, E)\beta(\alpha^*)$ from equation 19 and $\left.\frac{dg}{d\alpha}\right|_{\alpha=\alpha^*} = -g^2(\alpha^*, E)$. This means

that if sociality increases, the optimal level of virulence must decrease (Fig. 7). This may

appear to contradict classical predictions on population mixing and the evolution of

virulence (Ewald 1994), but the difference occurs due to the lack of population structure in

the present model. Note that if the host mortality rate is independent of sociality ($a = 1$),

then it follows that the optimal level of virulence occurs when:

457
$$\left.\frac{d\beta}{d\alpha}\right|_{\alpha=\alpha^*} = \frac{\beta(\alpha^*)}{d + \alpha^* + \gamma} \tag{21}$$

which does not depend on the contact effort, E. Thus, the reason optimal virulence is

inversely related to sociality in the present model is due to effects of sociality on the

mortality rate of the host (and hence the infectious period), rather than effects on

population mixing, *per se*.

Figure 7. The evolution of virulence in the fast social information approximation. Optimal virulence, α^*,

decreases with the contact effort (solid) and disease prevalence (dashed) increases. Vertical dotted lines

indicate where disease ($\tilde{R}_0^D < 1$) and ($\tilde{R}_0^I < 1$) social information transmission are not viable. Other

parameters as in Fig. 2, with $\beta(\alpha) = \sqrt{\alpha}$ and $0 \leq \alpha \leq 1$.

COEVOLUTION OF SOCIALITY AND VIRULENCE

We combine our results from the previous sections to analyse the effects of social

information on the coevolution of sociality and virulence. For tractability, we make three

simplifying assumptions: (1) social information dynamics are fast (i.e. we use the

approximation to the full model); (2) the parasite evolves much faster than the host; and (3)

the relationship between transmission and virulence is $\beta(\alpha) = \sqrt{\alpha}$ and $0 \leq \alpha \leq 1$.

Simulations indicate that relaxing these assumptions does not qualitatively affect our

results. The second assumption allows us to substitute $\alpha = \alpha^*$ into the host fitness gradient,

where:

$$\alpha^* = \begin{cases} \dfrac{d\sigma(1-a)}{E^2\tau} + (ad + \gamma) & \text{for } \tilde{R}_0^I > 1 \\ d + \gamma & \text{for } \tilde{R}_0^I < 1 \end{cases} \tag{22}$$

The range of qualitative outcomes for coevolution are the same as for one-sided adaptation

in the host (Fig. 8). The host and parasite may tend to a joint stable level of sociality and

virulence, respectively, or they may experience runaway selection for higher sociality and

lower virulence, or disease or social information may be unviable. However, note that in the

case of runaway selection virulence asymptotes to a lower bound of $\alpha^* = ad + \gamma$ (from

equation 22). Under coevolution, the evolutionary stable level of virulence increases with

the ratio of the information expiry rate to transmissibility (σ/τ), until social information

dynamics are no longer viable ($\tilde{R}_0^I < 1$).

Figure 8. Coevolution arising from social information and disease transmission dynamics. Coevolution of host

sociality (contact effort, E) and parasite virulence (α) as a function of the social information expiry rate

divided by the transmission probability, σ/τ, and for different values of the benefits of social information, a

(fast social information approximation). Black curves show the evolutionary stable level of virulence that

corresponds to the stable (dark blue) level of host sociality (or the threshold below which disease is driven

extinct, light blue). All other curves as described in Fig. 4. Other parameters as in Fig. 2, with $\tau = 1$, $\beta(\alpha) = \sqrt{\alpha}$

and $0 \leq \alpha \leq 1$.

Scenario	Variable	Description of results	Figure(s)
Host evolution	Disease prevalence	Selection against sociality at high/low disease prevalence	Fig. 3
	Social information prevalence	Selection for sociality at high/low social information prevalence	Fig. 3
	Social information expiry rate	Slow expiry: sociality either evolves to a stable level or increases without bound; fast expiry: sociality either increases without bound or decreases until disease is driven extinct	Fig. 4
	Social information timescale	Faster social information dynamics generally have a stabilising effect (prevent runaway selection for sociality)	Fig. 4, 6
Parasite evolution	Contact effort	Virulence decreases with higher contact efforts	Fig. 7
Coevolution	Social information expiry rate	Virulence increases with the social information expiry rate until social information is no longer viable	Fig. 8

492 **Table 1 – Summary of key results.**

DISCUSSION

Sociality entails both risks and benefits, but there is currently little understanding of how

dynamic ecological processes influence the evolution of sociality. Here, we explored a model

of social information and disease transmission to determine their effects on the evolution of

host sociality and parasite virulence. Although previous studies have examined the role of

parasitism in the evolution of sociality, ours is the first that captures both dynamic costs and

benefits of different social tendencies. This means that the fitness benefits of sociality, in

terms of increased access to social information, and fitness costs, in terms of increased

exposure to infection, are not imposed on the system, but instead are emergent properties

that are subject to eco-evolutionary feedbacks. Hence, changes in sociality at the individual

level affect the properties of transmission at the population level, and these population-

level properties, in turn, affect both the costs and benefits of sociality at the level of the

individual.

Our model shows how contrasting transmission processes can trade-off to mediate the

evolution of sociality and virulence. Crucially, we have shown that selection for sociality

(and hence virulence) depends non-monotonically on the prevalence of both social

information and disease (Fig. 3). Specifically, selection usually favours greater sociality (and

lower virulence) when social information is neither too common nor too rare and disease

risk is either low or high. Conversely, selection against sociality (and for greater virulence)

typically occurs when social information is either rare or very common and disease is at

moderate prevalence. The fact that these two processes mirror each other is not surprising:

the transmission processes are fundamentally the same in the model, but social information

is beneficial and disease is costly. We note, however, that in real populations transmission

processes for social information do not always mirror those of disease. In terms of social

information, individuals can use different learning rules, for example if they are conformist

(e.g. having a disproportionate preference for copying a more common behaviour, (Aplin et

al. 2015a) or if they require passing a certain threshold number of informed contacts before

becoming informed (Rosenthal et al. 2015), which can modify how transmission pathways

emerge from social contacts. From the perspective of diseases, the life cycle of pathogens,

such as the reliance of an intermediate host, can also impact how social contacts translate

to transmission events (Grear et al. 2013; Farine 2017).

In general, conditions on social information prevalence that select for greater sociality (and

lower virulence) are the same conditions on disease prevalence that select against sociality

(and for greater virulence). Naively, one might assume that selection for sociality should

increase with social information prevalence and decrease with disease prevalence. But

when social information prevalence is high, the marginal benefit of increasing sociality is

low, and similarly when disease prevalence is high, the marginal cost of being more social is

low. Slightly higher or lower sociality therefore makes little difference to the rate of

exposure to social information or disease when prevalence is high. The effects of disease

prevalence on the evolution of sociality have previously been observed by Bonds et al.

(2005), but the effects of social information prevalence on the evolution of sociality have

not previously been documented. Bonds et al. (2005) likened the relationship between

disease prevalence and selection for sociality to optimal host investment in an immune

response, which is predicted to peak when the risk of infection is neither too low, nor too

high. In fact, this pattern is quite general and occurs for a variety of behavioural and

physiological defences against parasitism, including resistance (Haldane 1949), recovery

(van Baalen 1998), mate choice (Ashby and Boots 2015) and sexual reproduction (Ashby and

King 2015).

The patterns described above for high/low prevalence are quite robust, but the picture is

more complicated when social information and disease are both at intermediate

prevalence. Selection may increase or decrease sociality and virulence depending on the

relative costs and benefits of being infected or informed, in addition to the likelihood of

exposure. As a consequence, it is possible for evolutionary trajectories to exist where the

sign of the fitness gradient is invariant to disease or social information prevalence, leading

to runaway selection for higher sociality/lower virulence or lower sociality/higher virulence

(Fig. 4, 8). Note that, while our model allowed for runaway selection for sociality, in reality

one would expect other factors to eventually constrain evolution (e.g., physiological

constraints, additional costs). We omitted additional constraints from our model so that we

could determine when disease alone is sufficient to constrain the evolution of sociality. The

results for "runaway selection" should therefore be interpreted in this light.

It is also possible for the initial conditions to determine the long-term evolutionary outcome

(stable level of sociality and virulence, or runaway selection for increased sociality and

decreased virulence) due to the existence of evolutionary repellers (Fig. 4, 8). The possibility

of more than one (co)evolutionary outcome suggests that different populations might

evolve very different social strategies for coping with infectious disease, and that variable

ecological conditions can lead to sudden shifts in sociality and virulence. How evolutionary

outcomes play out in a given population is likely to also be affected by the structure of the

social contacts. Our model assumed a well-mixed population, which can represent dynamics

in one small patch. However, at a broader scale, it is now clear that almost all animal

populations exhibit some structure (Cantor et al. 2021), and that population structure can

shape both transmission dynamics and evolution of diseases (Eames and Keeling 2006; Lion

and Boots 2010; Ashby and Gupta 2013). Structure can arise due to individual preferences

for certain relationships (Alberts 2019) or because the environment induces limitations on

who can come into contact with whom (He et al. 2019; Silk et al. 2019; Albery et al. 2021).

For example, recent work has suggested that—all else being equal—some landscapes may

be more predisposed to facilitating disease transmission than other landscapes (He et al.

2021). Some mammal (Grueter et al. 2020) and birds (Papageorgiou et al. 2019) also exhibit

multiple levels of social connections – named multilevel societies – with social contacts

being expressed within and across social units. Further, it is now becoming well-established

that individuals can change their social behaviours once they become infected (reviewed in

Romano et al. 2020; Stockmaier et al. 2021), which can introduce a more rapid feedback

between disease state and social behaviour. Thus, future research should aim to determine

how specific social strategies might evolve under different environmental conditions (e.g.

Udiani and Fefferman 2020).

An important consideration when modelling the benefits of sociality dynamically in terms of

social information, as opposed to fixed benefits (Bonds et al. 2005; Prado et al. 2009), is the

timescale of information dynamics relative to other processes. We have shown that for the

most part, evolutionary outcomes are qualitatively and quantitatively similar whether social

information dynamics are relatively fast or on a similar timescale to demographic and

epidemiological processes. However, this is not always the case. For example, increasing the

speed of social information dynamics relative to other processes has a stabilising effect on

the evolution of sociality (Fig. 6). In general, faster social information dynamics tend to have

a stabilising effect, reducing parameter ranges for runaway selection. Hence, while one can

often approximate social information dynamics by assuming the population rapidly reaches

a stable distribution, this assumption does not always hold and can fundamentally change

evolutionary outcomes.

One of the key predictions from this study and previous work (Bonds et al. 2005) is that

sociality should vary inversely with virulence, which may appear at odds with classical

predictions for virulence evolution (Ewald 1994). The contradiction is resolved by noticing

that: (1) our model considers contact rates in a randomly mixing population rather than

contact number in a structured population; and (2) higher contact rates do not directly

select for lower virulence, but do so indirectly through the spread of social information,

which reduces host mortality and increases the infectious period. Hence, greater sociality as

modelled here is not equivalent to population mixing in a structured model, and the effects

on virulence evolution are mediated via host mortality rather than contact rates. Based on

these insights, we therefore predict that sociality should covary inversely with virulence in

populations where social or spatial structuring is weak but should covary positively with

virulence when structuring is strong.

In previous studies, the fitness benefits of sociality were fixed rather than dynamic (Bonds et

al. 2005; Prado et al. 2009). While some studies have explored how social information and

infectious disease affect the contact structure of a population, this has been in the form of

host plasticity during an epidemic, rather than an evolutionary response (Funk et al. 2010),

or where the outcomes are compared across populations with different fixed social

strategies (Udiani and Fefferman 2020). Here, the two processes operate as independent

forces of selection. Our findings are closely linked to those of Bonds et al. (2005), where

sociality and virulence are shown to coevolve to stable levels. In contrast, Prado et al. (2009)

have shown that coevolutionary cycling (fluctuating selection) may occur between sociality

and virulence (Prado et al. 2009). It is not clear why Prado et al. (2009) found cycles, yet

Bonds et al. (2005) and the present study have not, but part of the reason may be due to

model structure. Prado et al. (2009) modelled disease dynamics using a stochastic

individual-based network, whereas Bonds et al. (2005) and the present study employed

deterministic models with randomly-mixing populations. A priority for future research in

this area is therefore to establish precisely when coevolutionary cycling of sociality and

virulence occurs, and whether this is driven by population structure and/or stochasticity.

The present study is relevant to empirical research on the sociality of animal populations

(Dall et al. 2004; Sih et al. 2004; Dingemanse and Réale 2005; Evans et al. 2020). In

particular, a number of studies have shown that variation in sociality influences the spread

of social information (Aplin et al. 2012; Allen et al. 2013; Farine et al. 2015a; Snijders et al.

2021). For example, the social network attributes of wild great tits (*Parus major*) can be

used to predict the discovery rate of novel food patches (Aplin et al. 2012). As in previous

theoretical studies (Bonds et al. 2005; Prado et al. 2009), we did not observe coexistence of

different social strategies. However, several empirical studies have highlighted the potential

importance of maintaining mixed strategies in social groups or local populations. For

example, Aplin et al. (2014) suggested that variation in sociality could increase group-level

efficiency in exploiting patches. Similarly, in ants, mixed colonies (in terms of aggressive

type) have more efficient task allocation and are more successful at capturing prey

(Modlmeier et al. 2012). A potential reason for the lack of mixed social strategies in current

theoretical studies is that they do not include assortative mixing (Farine 2014). Assortativity

has been shown to facilitate the coexistence of multiple parasites (Eames and Keeling 2006),

so it is possible that mixed social strategies could also evolve in the presence of non-random

mixing – a process that could be either actively generated through social preferences or

passively through exogenous drivers of social structure. It is also possible that mixed social

strategies could evolve in fluctuating environments, especially if individuals value

information differently (e.g. due to personality traits, Aplin et al. 2014), or if individuals vary

in their propensity to form social relationships (e.g. due to developmental conditions,

Boogert et al. 2014). A challenge for future theoretical research is to establish whether

selection arising from disease and information-use dynamics can drive the evolution of

mixed social strategies. Our results suggest that a number of potential mechanisms (e.g.

variation in the perceived value of information) could facilitate this process.

In summary, our study suggests that not only does the sociality of the population affect

social information and disease dynamics, but also that these processes are likely to

influence the evolution of sociality and virulence. Crucially, we predict that: (1) selection for

sociality and virulence will vary non-monotonically with both social information and disease

prevalence; and (2) if social information increases host survival, then increased sociality may

correlate with decreased virulence. The context-dependent relationships that we have

shown to exist between the ecology of information/disease dynamics and the evolution of

sociality/virulence highlights the need to understand host and parasite evolution in the

context of multiple ecological processes.

ACKNOWLEDGEMENTS

We thank Ben Sheldon for comments on an earlier version of the manuscript. BA was

supported by the Natural Environment Research Council (grant numbers NE/N014979/1 and

NE/V003909/1). DRF was funded by a grant from the European Research Council (ERC)

under the European Union's Horizon 2020 research and innovation programme (grant

agreement No. 850859) and an Eccellenza Professorship Grant of the Swiss National Science

Foundation (Grant Number PCEFP3_187058), and received additional support from the

Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's

Excellence Strategy – EXC 2117 – 422037984.

DATA AVAILABILITY

Source code is available in the *Supplementary Material* and at the following Github

repository: https://github.com/ecoevogroup/Ashby_and_Farine_2020.

COMPETING INTERESTS

The authors have no competing interests to declare.

REFERENCES

Alberts, S. C. 2019. Social influences on survival and reproduction: Insights from a long-term

   study of wild baboons. J. Anim. Ecol. 88:47–66.

Albery, G. F., L. Kirkpatrick, J. A. Firth, and S. Bansal. 2021. Unifying spatial and social

network analysis in disease ecology. J. Anim. Ecol. 90:45–61.

Alizon, S. 2008. Transmission-recovery trade-offs to study parasite evolution. Am. Nat.

172:E113-21.

Allen, J., M. Weinrich, W. Hoppitt, and L. Rendell. 2013. Network-Based Diffusion Analysis

Reveals Cultural Transmission of Lobtail Feeding in Humpback Whales. Science (80-.).

340:485–488.

Aplin, L. M., D. R. Farine, R. P. Mann, and B. C. Sheldon. 2014. Individual-level personality

influences social foraging and collective behaviour in wild birds. Proc. R. Soc. B

281:20141016.

Aplin, L. M., D. R. Farine, J. Morand-Ferron, A. Cockburn, A. Thornton, and B. C. Sheldon.

2015a. Experimentally induced innovations lead to persistent culture via conformity in

wild birds. Nature 518:538–541. Nature Publishing Group.

Aplin, L. M., D. R. Farine, J. Morand-Ferron, E. F. Cole, A. Cockburn, and B. C. Sheldon. 2013.

Individual personalities predict social behaviour in wild networks of great tits (Parus

major). Ecol. Lett. 16:1365–1372.

Aplin, L. M., D. R. Farine, J. Morand-Ferron, and B. C. Sheldon. 2012. Social networks predict

patch discovery in a wild population of songbirds. Proc. R. Soc. B 279:4199–4205.

Aplin, L. M., J. A. Firth, D. R. Farine, B. Voelkl, R. A. Crates, A. Culina, C. J. Garroway, C. A.

Hinde, L. R. Kidd, I. Psorakis, N. D. Milligan, R. Radersma, B. L. Verhelst, and B. C.

Sheldon. 2015b. Consistent individual differences in the social phenotypes of wild great

tits, Parus major. Anim. Behav. 108:117–127. Elsevier Ltd.

Ashby, B., and M. Boots. 2015. Coevolution of parasite virulence and host mating strategies.

Proc. Natl. Acad. Sci. 112:13290–13295.

Ashby, B., and E. Bruns. 2018. The evolution of juvenile susceptibility to infectious disease.

Proc. R. Soc. B Biol. Sci. 285:20180844.

Ashby, B., and S. Gupta. 2013. Sexually transmitted infections in polygamous mating

systems. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 368:20120048.

Ashby, B., and K. C. King. 2015. Diversity and the maintenance of sex by parasites. J. Evol.

Biol. 28:511–520.

Barton, R. A., R. W. Byrne, and A. Whiten. 1996. Ecology, feeding competition and social

structure in baboons. Behav. Ecol. Sociobiol. 38:321–329.

Bonds, M. H., D. C. Keenan, A. J. Leidner, and P. Rohani. 2005. Higher disease prevalence can

induce greater sociality: a game theoretic coevolutionary model. Evolution (N. Y).

59:1859–1866.

Boogert, N. J., D. R. Farine, and K. A. Spencer. 2014. Developmental stress predicts social

network position. Biol. Lett. 10:20140561.

Cantor, M., L. M. Aplin, and D. R. Farine. 2020. A primer on the relationship between group

size and group performance. Anim. Behav. 166:139–146. Elsevier Ltd.

Cantor, M., A. A. Maldonado-Chaparro, K. B. Beck, H. B. Brandl, G. G. Carter, P. He, F.

Hillemann, J. A. Klarevas-Irby, M. Ogino, D. Papageorgiou, L. Prox, and D. R. Farine.

2021. The importance of individual-to-society feedbacks in animal ecology and

evolution. J. Anim. Ecol. 90:27–44.

Couzin, I. D. 2006. Behavioral ecology: Social organization in fission-fusion societies. Curr.

Biol. 16:R169–R171.

Croft, D. P., J. Krause, S. K. Darden, I. W. Ramnarine, J. J. Faria, and R. James. 2009.

Behavioural trait assortment in a social network: patterns and implications. Behav.

Ecol. Sociobiol. 63:1495–1503.

Dall, S. R., A. I. Houston, and J. M. McNamara. 2004. The behavioural ecology of personality:

consistent individual differences from an adaptive perspective. Ecol. Lett. 7:734–739.

Danchin, E., L.-A. Giraldeau, T. J. Valone, and R. H. Wagner. 2004. Public information: from

nosy neighbors to cultural evolution. Science (80-.). 305:487–491.

Dingemanse, N. J., and D. Réale. 2005. Natural selection and animal personality. Behaviour

142:1165–1190.

Doligez, B., E. Danchin, and J. Clobert. 2002. Public information and breeding habitat

selection in a wild bird population. Science (80-.). 297:1168–1170.

Eames, K. T. D., and M. J. Keeling. 2006. Coexistence and specialization of pathogen strains

on contact networks. Am. Nat. 168:230–241.

Elgar, M. A. 1986. The establishment of foraging flocks in house sparrows: risk of predation

and daily temperature. Behav. Ecol. Sociobiol. 19:433–438.

Evans, J. C., M. J. Silk, N. J. Boogert, and D. J. Hodgson. 2020. Infected or informed? Social

structure and the simultaneous transmission of information and infectious disease.

Oikos 129:1271–1288.

Ewald, P. W. 1994. Evolution of infectious disease. Oxford Univ Press, Oxford, UK.

Farine, D. 2017. The dynamics of transmission and the dynamics of networks. J. Anim. Ecol.

86:415–418.

Farine, D. R. 2014. Measuring phenotypic assortment in animal social networks: Weighted

associations are more robust than binary edges. Anim. Behav. 89:141–153. Elsevier Ltd.

Farine, D. R., L. M. Aplin, B. C. Sheldon, and W. Hoppitt. 2015a. Interspecific social networks

promote information transmission in wild songbirds. Proc. Biol. Sci. 282:20142804.

Farine, D. R., J. A. Firth, L. M. Aplin, R. a. Crates, A. Culina, C. J. Garroway, C. a. Hinde, L. R.

Kidd, N. D. Milligan, I. Psorakis, R. Radersma, B. Verhelst, B. B. Voelkl, B. C. Sheldon, L.

M. Aplina, R. a. Crates, A. Culina, C. J. Garroway, C. a. Hinde, L. R. Kidd, N. D. Milligan, I.

Psorakis, R. Radersma, B. Verhelst, B. B. Voelkl, and B. C. Sheldon. 2015b. The role of

social and ecological processes in structuring animal populations: a case study from

automated tracking of wild birds. R. Soc. Open Sci. 2:150057.

Funk, S., M. Salathé, and V. A. A. Jansen. 2010. Modelling the influence of human behaviour

on the spread of infectious diseases: a review. J. R. Soc. Interface 7:1247–1256.

Galef, B. G., and L.-A. Giraldeau. 2001. Social influences on foraging in vertebrates: causal

mechanisms and adaptive functions. Anim. Behav. 61:3–15.

Geritz, S. A. H., E. Kisdi, G. Meszena, and J. A. J. Metz. 1998. Evolutionarily singular strategies

and the adaptive growth and branching of the evolutionary tree. Evol. Ecol. 12:35–37.

Grear, D. A., L. T. Luong, P. J. Hudson, D. A. Grear, L. T. Luong, and P. J. Hudson. 2013.

Network transmission inference : Host behavior and parasite life cycle make social

networks meaningful in disease ecology Published by : Wiley on behalf of the Ecological

Society of America Stable URL : http://www.jstor.org/stable/23596896 REFERENCES

Link. Ecol. Appl. 23:1906–1914.

Grueter, C. C., X. Qi, D. Zinner, T. Bergman, M. Li, Z. Xiang, P. Zhu, A. B. Migliano, A. Miller,

M. Krützen, J. Fischer, D. I. Rubenstein, T. N. C. Vidya, B. Li, M. Cantor, and L. Swedell.

2020. Multilevel Organisation of Animal Sociality. Trends Ecol. Evol. 35:834–847.

Elsevier Ltd.

Haldane, J. B. S. 1949. Disease and evolution. La Ric. Sci. 19:68–76.

He, P., A. A. Maldonado-Chaparro, and D. R. Farine. 2019. The role of habitat configuration

in shaping social structure: a gap in studies of animal social complexity. Behav. Ecol.

Sociobiol. 73:9. Behavioral Ecology and Sociobiology.

He, P., P.-O. Montiglio, M. Somveille, M. Cantor, and D. R. Farine. 2021. The role of habitat

configuration in shaping animal population processes: a framework to generate

quantitative predictions. Oecologia 2020.07.30.228205. Springer Berlin Heidelberg.

Kenward, R. E. 1978. Hawks and Doves : Factors Affecting Success and Selection in Goshawk

Attacks on Woodpigeons Author (s): R . E . Kenward Source : Journal of Animal Ecology

, Vol . 47 , No . 2 (Jun ., 1978), pp . 449-460 Published by : British Ecological Society

Stable. J. Anim. Ecol. 47:449–460.

Krause, J., and G. D. Ruxton. 2002. Living in groups. Oxford University Press, Oxford, UK.

Lion, S., and M. Boots. 2010. Are parasites "'prudent'" in space? Ecol. Lett. 13:1245–1255.

Lloyd-Smith, J. O., S. J. Schreiber, P. E. Kopp, and W. M. Getz. 2005. Superspreading and the

effect of individual variation on disease emergence. Nature 438:355–339.

Modlmeier, A. P., J. E. Liebmann, and S. Foitzik. 2012. Diverse societies are more productive:

a lesson from ants. Proc. R. Soc. B 279:2142–2150.

Papageorgiou, D., C. Christensen, G. E. C. Gall, J. A. Klarevas-Irby, B. Nyaguthii, I. D. Couzin,

and D. R. Farine. 2019. The multilevel society of a small-brained bird. Curr. Biol.

29:R1120–R1121. Elsevier.

Pike, T. W., M. Samanta, J. Lindström, and N. J. Royle. 2008. Behavioural phenotype affects

social interactions in an animal network. Proc. R. Soc. B 275:2515–2520.

Pöysä, H. 1992. Group foraging in patchy environments: the importance of coarse-level local

enhancement. Ornis Scand. 23:159–166.

Prado, F., A. Sheih, J. D. West, and B. Kerr. 2009. Coevolutionary cycling of host sociality and

pathogen virulence in contact networks. J. Theor. Biol. 261:561–569.

Romano, V., A. J. J. MacIntosh, and C. Sueur. 2020. Stemming the Flow: Information,

Infection, and Social Evolution. Trends Ecol. Evol. 35:849–853. Elsevier Ltd.

Rosenthal, S. B., C. R. Twomey, A. T. Hartnett, H. S. Wu, and I. D. Couzin. 2015. Revealing the

hidden networks of interaction in mobile animal groups allows prediction of complex

behavioral contagion. Proc. Natl. Acad. Sci. U. S. A. 112:4690–4695.

Sachser, N. 1986. Different forms of social organization at high and low population densities

in guinea pigs. Behaviour 97:253–272.

Sih, A., A. Bell, and J. C. Johnson. 2004. Behavioral syndromes: an ecological and

evolutionary overview. Trends Ecol. Evol. 19:372–378.

Silk, M. J., D. P. Croft, T. Tregenza, and S. Bearhop. 2014. The importance of fission–fusion

social group dynamics in birds. Ibis (Lond. 1859). 156:701–715.

Silk, M. J., D. J. Hodgson, C. Rozins, D. P. Croft, R. J. Delahay, M. Boots, and R. A. McDonald.

2019. Integrating social behaviour, demography and disease dynamics in network

models: Applications to disease management in eclining wildlife populations. Philos.

Trans. R. Soc. B Biol. Sci. 374.

Snijders, L., S. Krause, A. N. Tump, M. Breuker, C. Ortiz, S. Rizzi, I. W. Ramnarine, J. Krause,

and R. H. J. M. Kurvers. 2021. Causal evidence for the adaptive benefits of social

foraging in the wild. Commun. Biol. 4:94.

Stockmaier, S., N. Stroeymeyt, E. C. Shattuck, D. M. Hawley, L. A. Meyers, and D. I. Bolnick.

2021. Infectious diseases and social distancing in nature. Science (80-.). 371.

Tanner, C. J., and A. L. Jackson. 2012. Social structure emerges via the interaction between

local ecology and individual behaviour. J. Anim. Ecol. 81:260–277.

Udiani, O., and N. H. Fefferman. 2020. How disease constrains the evolution of social

systems. Proceedings. Biol. Sci. 287:20201284.

Valone, T. J. 2007. From eavesdropping on performance to copying the behavior of others: a

review of public information use. Behav. Ecol. Sociobiol. 62:1–14.

van Baalen, M. 1998. Coevolution of recovery ability and virulence. Proc. R. Soc. B 265:317–

325. The Royal Society.

Watters, J., and A. Sih. 2005. The mix matters: behavioural types and group dynamics in

water striders. Behaviour 142:1417–1431.

Weber, N., S. P. Carter, S. R. X. Dall, R. J. Delahay, J. L. McDonald, S. Bearhop, and R. A.

McDonald. 2013. Badger social networks correlate with tuberculosis infection. Curr.

Biol. 23:R915–R916. Elsevier.