***Arabidopsis* inositol polyphosphate kinases regulate COP9 signalosome deneddylase**

functions in phosphate-homeostasis

Yashika Walia[1], Mritunjay Kasera[1], Kishor D Ingole[1,2], Swaroop Peddiraju[1], Abhisha Roy[1],

Debabrata Laha[3], Gabriel Schaaf[4], and Saikat Bhattacharjee[1*]

[1]Laboratory of Signal Transduction and Plant Resistance, UNESCO-Regional Centre for

Biotechnology (RCB), NCR Biotech Science Cluster, 3[rd] Milestone, Faridabad-Gurgaon

Expressway, Faridabad- 121 001, Haryana, India.

[2]Kalinga Institute of Industrial Technology (KIIT) University, Bhubaneswar- 751 024, Odisha,

India.

[3]Department of Biochemistry, Indian Institute of Science, Bengaluru, Karnataka- 560 012,

India.

[4]Department of Plant Nutrition, Institute of Crop Science and Resource Conservation,

Rheinische Friedrich-Wilhelms-Universität Bonn, 53115 Bonn, Germany.

*Corresponding author:

Saikat Bhattacharjee

saikat@rcb.res.in

Tel: (+91) 0129-2848837

ORCID ID

Saikat Bhattacharjee https://orcid.org/0000-0003-1369-7700

Gabriel Schaaf https://orcid.org/0000-0001-9022-4515

Yashika Walia https://orcid.org/0000-0001-8122-8661

Mritunjay Kasera https://orcid.org/0000-0002-1110-3502

Kishor D Ingole https://orcid.org/0000-0003-1946-6639

Swaroop Peddiraju https://orcid.org/0000-0001-5290-7067

Abhisha Roy https://orcid.org/0000-0002-3431-0811

Debabrata Laha https://orcid.org/0000-0002-7823-5489

Author Contributions

30   Y.W. and S.B. conceived and designed the research. D.L. and G.S. generated transgenic line

31   used here. Y.W. and S.B. performed most of the experiments. Y.W., M.K., K.D.I, S.P., and

32   A.R. conducted and analyzed gene/protein expression changes. Y.W. and S.B. wrote the

33   manuscript.

Competing interest statement

35   The authors declare no conflict of interest for this study.

Keywords

37   Inositol polyphosphates, COP9 signalosome, CSN5, IP5K, IP6K1, IPK1, ITPK1

Abstract

Targeted protein degradation is essential for physiological development and adaptation to stress. Mammalian INOSITOL PENTAKISPHOSPHATE 2-KINASE (IP5K) and INOSITOL HEXAKISPHOSPHATE KINASE 1 (IP6K1) pair generates inositol polyphosphates (InsPs) to modulate association/dissociation equilibrium of Cullin RING Ubiquitin E3 ligases (CRLs) on the COP9 signalosome (CSN) platform. Deneddylase activity of the CSN5 subunit protects cullins from self-ubiquitination ensuring their functional continuity. In plants, similar regulations by InsP-kinases are not known. Here, we show conserved interactions of *Arabidopsis thaliana* INOSITOL PENTAKISPHOSPHATE 2-KINASE 1 (IPK1) and INOSITOL 1,3,4-TRISPHOSPHATE 5/6-KINASE 1 (ITPK1), counterparts of the above mammalian InsP-kinase pair, with selective CSN subunits. In *ipk1* or *itpk1* mutants, deneddylation deficiencies not only cause increased neddylated Cullin1 (CUL1^{Nedd8}) pools more prone to degradation but also impair CSN5 entry/exit shuttles on the CSN holo-complex. Constitutive phosphate-starvation response (PSR), previously known for these mutants are suppressed by pharmacological inhibition of neddylation thus linking CSN-CRL functions to phosphate (Pi)-sensing. Similarly, in wild-type plants exposed to compounds that impair CSN5 deneddylase function affects its dynamics and mimic PSR properties of the mutants. We further show that under Pi-deprivation more CSN5 retentions on the CSN holo-complex and the resulting enhanced CUL1^{Nedd8} pools is essential for induction of downstream Pi-starvation inducible (PSI) genes. Overall, with our data we present InsP-kinase involvements in maintenance of Pi-homeostasis in plants via CRL-CSN(5) functional synergism.

Significance

Neddylation modifications on Culling-RING E3 ligases (CRLs) while essential for their role in proteostasis, also threaten their own stability. Selective inositol polyphosphates (InsPs) aid

the constitutive photomorphogenesis 9 signalosome (CSN) functions in protecting,

deneddylating, and facilitating CRL recycling. Here we demonstrate that plant mutants

deficient in these InsPs have disturbed CSN subunit partitioning, are deficient in deneddylase

activity, and hyperaccumulate neddylated cullins that lead to constitutive phosphate-starvation

response (PSR). Inhibition of CSN functions/consequences mirror the InsP mutant properties

indicating pivotal involvement of CSN in phosphate homeostasis. These raises promising

possibilities of targeted intervention on CSN functions for nutritional benefit of plants.

Introduction:

The versatile functions of inositol polyphosphates (InsPs) in cellular processes of all eukaryotes are owed to its propensity for differential degree of phosphorylation catalyzed by relatively conserved set of InsP-kinases (Stevenson-Paulik *et al*., 2005; Monserrate and York, 2010; Shears and Wang, 2019). In plants, selective InsPs have been linked to phytohormone networks, inorganic phosphate (Pi)-homeostasis, mRNA export, (a)biotic stress responses, among others (Stevenson-Paulik *et al*., 2005; Murphy *et al*., 2008; Gillaspy, 2013; Lee *et al*., 2015; Williams *et al*., 2015; Jia *et al*., 2019). With all six-hydroxyl group of the *myo*-inositol ring phosphorylated, $InsP_6$ (myo-inositol-1,2,3,4,5,6-hexakisphosphate or phytic acid), produced by INOSITOL PENTAKISPHSPHATE 2-KINASE 1 (IPK1) is sequentially (pyro)-phosphorylated by INOSITOL 1,3,4-TRISPHOSPHATE 5/6-KINASES (ITPK1/2) and then by DIPHOSPHOINOSITOL PENTAKISPHOSPHATE KINASES (VIH1/2) to generate energy-rich phosphoanhydride bond-containing inositol pyrophosphates $InsP_7$ and $InsP_8$, respectively (Desai *et al*., 2014; Laha *et al*., 2015; Adepoju *et al*., 2019; Laha *et al*., 2019). Though *bona fide* receptors of plant InsPs remain unidentified, their co-factor functions are being increasingly elaborated (Shears *et al*., 2012; Williams *et al*., 2015). While $InsP_6$, or more recently speculated $InsP_7$, promotes auxin-dependent degradation of AUX/IAA repressors, $InsP_8$ targets JASMONATE ZIM-DOMAIN (JAZ) repressors in a jasmonic acid (JA)-dependent manner (Tan *et al*., 2007; Sheard *et al*., 2010; Laha *et al*., 2015; Laha *et al*., 2020). Interestingly, the cognate substrate receptors TRANSPORT INHIBITOR RESPONSE 1 (TIR1) or CORONATINE INSENSITIVE 1 (COI1) belong to F-box family proteins and are components of typical SCF (ASK1/SKP1-Cullin-F-Box; SCF^{TIR1} or SCF^{COI1})-type E3 ubiquitin ligases (Tan *et al*., 2007; Sheard *et al*., 2010). The *ipk1* or *itpk1* mutants with reduced $InsP_7$ or *vih2* plants with reduced $InsP_8$, are auxin- or JA-insensitive, respectively (Laha *et al*., 2015; Laha *et al*., 2020).

InsPs have also been recently implicated as sensor of phosphate (Pi)-status in plants (Kuo *et*

al., 2018; Dong *et al.*, 2019; Ried *et al.*, 2019; Zhu *et al.*, 2019). Necessity of phosphates in

central dogma processes, protein modifications, and most importantly as indicators of

adenylate currencies risk detrimental consequences when cellular Pi levels are low (Szijgyarto

et al., 2011; Wundenberg and Mayr, 2012). Under a low Pi-availability from soil, a highly

organized adaptive cascade of events collectively termed as PSR is activated and includes

increased remobilization of Pi from root to shoot, harnessing of organellar Pi-storage reserves,

accompanied by upregulated expression of PSI genes for Pi-transporters (*PHT*s), phosphatases

(*PAP*s), and several micro- (*MIR399*) or non-coding RNAs (*IPS1*) (Puga *et al.*, 2014; Ham *et*

al., 2018). In Arabidopsis, PSR-associated transcriptional changes are majorly orchestrated by

the MYB-transcription factors PHOSPHATE STARVATION RESPONSE 1 (PHR1) and its

closest homolog PHR1-LIKE (PHL1) (Bustos *et al.*, 2010). These transcription factors bind to

partially conserved sequences (P1BS elements) present on PSI gene promoters to activate their

transcriptions (Rubio *et al.*, 2001). A *phr1 phl1* double mutant is deficient in PSI-gene

expressions both basally as well as upon exposure to Pi-deplete conditions (Kuo *et al.*, 2014).

A class of stand-alone SPX (SYG1/Pho81/XPR1)-domain containing proteins SPX1/2 act as

negative regulators of PHR1/PHL1 functions in maintaining phosphate balances in a cell.

Following the first report that SPX domains can bind specific InsPs it was subsequently

demonstrated that under Pi-replete conditions, SPX1 requires $InsP_8$ co-factor to bind PHR1,

sequester it away from the nucleus, and suppress its activity (Wild *et al.*, 2016; Jung *et al.*,

2018; Dong *et al.*, 2019; Zhu *et al.*, 2019). When starved for Pi, intracellular $InsP_8$ levels reduce

because of the phosphatase activity of VIH2 relieving SPX1-tethered inhibitions on PHR1,

thus allowing successful transduction of PSR (Dong *et al.*, 2019). In $InsP_8$-deficient *ipk1-1*,

itpk1-2, or *vih1 vih2* and in the *spx* mutants liberated PHR1 causes constitutive activation of

PSI-gene expressions even under Pi-sufficient conditions accumulating high endogenous Pi

levels (Kuo *et al*., 2014; Puga *et al*., 2014; Dong *et al*., 2019; Zhu *et al*., 2019). Decreased

$InsP_8$ levels that occur upon Pi-starvation also promotes ubiquitinylation of at least two SPX

proteins (Lv *et al*., 2014; Zhong *et al*., 2018; Osorio *et al*., 2019). Recent identification of E3

ubiquitin ligases that target rice SPX4 upon Pi-starvation (Ruan *et al*., 2019) taken together

with specific InsP-requirement in determination of fate of hormonal repressors, $InsP_7/InsP_8$

clearly are also biochemical regulators of stimulus-dependent substrate stabilities.

In a typical eukaryotic cell, ~20% of targeted protein degradations are mediated by Cullin

RING ubiquitin ligases (CRLs), the largest member among the E3 ubiquitin ligase family

(Soucy *et al*., 2009; Yi Sun, 2020). A modular CRL contains a Cullin (CUL) scaffold (CUL1-

4 in Arabidopsis), an E2-interacting RING-type Rbx/Roc protein, CUL member-specified

substrate adaptor and a ubiquitination target-specified substrate receptor. CRLs require

covalent modification of a Nedd8 moiety on CULs (a process termed as neddylation) for

function (Yi Sun, 2020). Neddylations improve CRL efficiencies in complex formation with

the ubiquitin E2 enzyme for subsequent degradation of targeted substrates. In absence of

cognate substrates self-ubiquitinylation-vulnerable CRLs are protected on Constitutive

photomorphogenesis 9 signalosome (CSN), an evolutionarily conserved eight-subunit (CSN1-

8) macromolecular complex (Chamovitz and Segal, 2001; Dubiel *et al*., 2015). The CSN5

subunit in this holo-complex contains an MPN+/JAMM metalloprotease motif that

catalytically removes Nedd8 (deneddylation) from CULs (Cope *et al*., 2002) causing CRL

disassembly and promoting newer associations with substrate receptors and adaptors required

for continuum of its functions. Interaction changes between CRL-CSN affect cellular ratio of

neddylated CUL: unneddylated CUL (CUL^{Nedd8}:CUL) (Scherer *et al*., 2016). Biochemical

modes of a CSN operation including its dynamic rearrangements upon CRL binding have been

gradually unraveled at structural and functional levels (Enchev *et al*., 2012; Lingaraju *et al*.,

2014; Dubiel *et al*., 2015). From these studies, it is evident that optimal CRL activity

necessitates regulated neddylation/deneddylation cycles of CULs. Neddylated CRLs are

captured on the CSN holo-complex only in deneddylation-deficient *csn* mutants inferring that

deneddylation-dependent disassembly is very rapid (Enchev *et al*., 2012). The 'CSN paradox'

(Dubiel, 2009), an apparent contradiction between genetic requirement of a CRL for CSN-

protection at the cost of its deneddylation-dependent inactivation is now better understood at a

molecular level. Recent breakthroughs show that highly locale-specific coordinated activities

of mammalian IP5K with IP6K1, functional homologs of Arabidopsis IPK1 and ITPK1,

provide $InsP_6$ to strengthen and $InsP_7$ to loosen CRL-CSN associations (Rao *et al*., 2014;

Scherer *et al*., 2016; Lin *et al*., 2020). Through intricate orchestrations, a CRL tether-promoting

$InsP_6$ is converted to the liberator $InsP_7$ by an inert IP6K1 that gets activated at CSN only upon

a stimulus. This synchronization ensures functional demands of CRLs meet the response

requirements proportionately without a compromise on its own stabilities. With reported

interactions between Arabidopsis ITPK1 and CSN1 (Qin *et al*., 2005), direct implication of

(any) InsP-kinase on CSN-activities from a plant system has not been forthcoming.

Here we show that Arabidopsis IPK1-ITPK1 pair retains conserved CSN interactions and

modulate CUL1 deneddylation efficiencies impacting association/dissociation kinetics of

CSN5 on the CSN holo-complex. Unlike noted in animal cells, CRL-CSN associations remain

mostly unaffected in *ipk1* or *itpk1* mutants. Nevertheless, enhanced level of CUL^{Nedd8} result in

constitutive PSR in these mutants. Further, we reveal that Pi-deprivation in wild-type plants is

associated with CSN(5) perturbations and increased CUL1 neddylation that precedes

PHR1/PHL1 role in inducing PSI gene expressions. Mutational or pharmacological inhibitions

of CSN5 result in constitutive PSR even under sufficient Pi-availability. However, when

actually Pi-starved, PSR responses in these plants or in the *ipk1* or *itpk1* mutants are suppressed

indicating that equilibrium shifts of CSN5 deneddylase activity rather than its complete

inhibition is essential to execute PSR. Our investigations overall reveal activities of plant IPK1-

ITPK1 pair in regulating CSN dynamics, its consequences on CRLs functions and requirement

in maintaining Pi-homeostasis or responding to its deficiencies.

Results

Arabidopsis IPK1 and ITPK1 interact with CSN holo-complex subunits.

Mammalian IP5K co-precipitates with CSN2/5 whereas IP6K1 interacts with CSN1/2 (Rao *et*

al., 2014; Scherer *et al*., 2016). Arabidopsis ITPK1 associates with CSN1 (Qin *et al*., 2005).

Interactions of Arabidopsis IPK1 with CSN subunits have not been investigated earlier. We

immuno-enriched Myc-IPK1 or ITPK1-GFP from *ipk1-1:Myc-IPK1* and *itpk1-2:ITPK1-GFP*

transgenic lines, respectively and probed for the presence of selective CSN subunits, CUL1 or

the substrate adaptor ASK1. These transgenic lines restore high endogenous Pi and elevated

PSI gene *SPX1* expressions to wild-type (Col-0) levels implicating the functionality of the

corresponding fusion proteins (Laha *et al*., 2020) (Supplemental Figure 1). In

immunoprecipitation assays, CSN1/2/4/5, CUL1 and ASK1 co-purified with IPK1, whereas

ITPK1 enrichments detected CSN1/2 and ASK1 (Figure 1A). Through Bimolecular

Fluorescence Complementation (BiFC) assays, IPK1 or ITPK1 association with CSN1 and

CSN2 are detected as nucleocytoplasmic complexes similar to localization of the individual

proteins (Adepoju *et al*., 2019) (Figure 1B). A negative control β-Glucuronidase (GUS) protein

does not interact with IPK1, ITPK1 or CSN1/2 (Supplemental Figure 2). These data suggest

that likewise to animals, CSN associations of IPK1 and ITPK1 are conserved in Arabidopsis.

Although biochemically coupled by substrate-product relationship, interaction between IPK1

and ITPK1 has not been demonstrated *in planta*. To this end, BiFC detects IPK1-ITPK1

complexes at nucleocytoplasmic locales (Figure 1B). Further, we performed co-

immunoprecipitation assays via transient co-expression of Myc-IPK1 and GFP-ITPK1 in

Nicotiana benthamiana leaves. Enrichment of GFP-ITPK1 with anti-GFP affinity beads co-

elutes Myc-IPK1 indicating their associations *in vivo* (Figure 1C). Overall, our results

recapitulate IPK1 and ITPK1 interactions with CSN and with each other.

To determine cellular proportions of IPK1 or ITPK1 that associate with the CSN holo-complex,

we subjected Col-0, *ipk1-1:Myc-IPK1* and *itpk1-2:ITPK1-GFP* extracts to size-exclusion

chromatography followed by immunoblots for representative CSN subunits, Myc-IPK1 or

ITPK1-GFP. The elution patterns of molecular weight standards and selective CSN subunits

with demarcation of CSN holo-complex fractions, as noted in earlier reports are shown

(Gusmaroli *et al.*, 2004; Dohmann *et al.*, 2005) (Figure 2A). CSN1, exclusively associates with

the CSN holo-complex and elutes accordingly in higher molecular weight pools (>600-400

kDa). CSN4 and CSN5 fractionates both as a part of CSN holo-complex as well as in lower

molecular weight pools (monomers, 100-50 kDa). Partitioning profiles of CSN5 and CSN4 are

comparable between Col-0, *ipk1-1:Myc-IPK1* and *itpk1-2:ITPK1-GFP* (Figure 2B;

Supplemental Figure S3A). Remarkably, neither Myc-IPK1 nor ITPK1-GFP proteins are

detected in fractions corresponding to the CSN holo-complex implying that their associations

are either transient or represent only marginal proportions of these proteins (Figure 2A). Co-

elution of Myc-IPK1 and ITPK1-GFP in overlapping fractions is likely indicative of their *in*

vivo associations we noted earlier.

CSN5 deneddylase deficiencies cause enhanced CUL1^{Nedd8} pools in *ipk1* or *itpk1* plants.

Depletion of IP5K or IP6K1 reduces CRL-CSN associations in mammalian systems (Rao *et*

al., 2014; Lin *et al.*, 2020). To examine whether in *ipk1-1* or *itpk1-2* lower CUL1 pools co-

fractionate with the CSN holo-complex, we immuno-enriched CUL1 from Col-0, *ipk1-1* or

itpk1-2 extracts and probed for the presence of CSN1/2. Although slightly lower CUL1 was

immunoprecipitated from *ipk1-1* or *itpk1-2* than Col-0, nonetheless proportionately similar

amounts of CSN1/2 as in Col-0 co-eluted suggesting that CRL-CSN associations are likely not

perturbed in these mutants (Figure 2C). Surprisingly, when we compared elution profiles,

CSN5 but not CSN4 in *ipk1-1* or *itpk1-2* extracts showed increased presence in CSN holo-

complex fractions with marked decrease in monomeric elutions, relative to Col-0 (Figure 2B).

Total CSN5 levels however remained comparable between all plants.

Resident CSN5 performs the deneddylase role in a CSN holo-complex (Cope *et al*., 2002).

With changed partitioning of CSN5 noted in *ipk1-1* or *itpk1-2* we tested whether deneddylation

efficiencies on CULs are impacted. In mammalian system depleting IP5K or IP6K1 causes

strong increments in CUL^{Nedd8} levels (Rao *et al*., 2014; Scherer *et al*., 2016). Remarkably, in

ipk1-1 or *itpk1-2* lysates, enhanced $CUL1^{Nedd8}$ levels relative to Col-0 are strikingly prominent

(Figure 2D). These increases are specific to *ipk1-1* or *itpk1-2* and extracts from *ipk2β-1* with

low $InsP_6$ levels as *ipk1-1* (Stevenson-Paulik *et al*., 2005), *vih2-4* plants with elevated $InsP_7$

(Laha *et al*., 2015), or *itpk4-1* with low $InsP_6$ and $InsP_7$ levels (Kuo *et al*., 2018) do not show

elevated $CUL1^{Nedd8}$ pools (Supplemental Figure S3B). In the respective complemented lines of

ipk1-1 or *itpk1-2*, CUL^{Nedd8}:CUL ratios are restored comparable to Col-0 (Figure 2D).

Global ubiquitinylation is enhanced due to loss of mammalian IP6K1 (Rao *et al*., 2014). As a

possible consequence to increased CUL^{Nedd8} levels, we tested relative levels of ubiquitinylation

in Col-0, *ipk1-1*, or *itpk1-2* extracts. Clear increase in global ubiquitin-conjugates in

appearance of a smear when probed with anti-ubiquitin antibodies is detected in the mutant

plants than Col-0 (Figure 2D). In complemented lines, wild-type ubiquitinylation levels are

noted. Further, in-lysate deneddylation assay to determine deneddylase defects (Lin *et al*.,

2020) reveal decreased efficiencies of CUL1-deneddylation by endogenous CSN5

simultaneous with increased CUL1 turnover rates in *ipk1-1* or *itpk1-2* extracts (Figure 2E).

Decreased CUL1 stability likely accounts for the reduced immuno-enrichments from the

mutants observed earlier. Taken together, our results parallel animal studies demonstrating

impaired CSN5 deneddylase activity in *ipk1-1* or *itpk1-2* that leads to augmented $CUL1^{Nedd8}$

levels prone to increased turnover due to self-ubiquitinylation.

Pi-starvation alters dynamics of CSN5 and its deneddylase efficiencies.

Known consequences in *ipk1-1* and *itpk1-2* are auxin-insensitivity and constitutive PSR (Kuo

et al., 2014; Kuo *et al*., 2018; Laha *et al*., 2020; Riemer *et al*., 2020). Reduced $InsP_8$ in these

mutants likely disrupt SPX1-PHR1 interactions that lead to activation of PSI-genes (Ried *et*

al., 2019). MLN4924, a small molecule pharmacological inhibitor of CUL neddylation (Soucy

et al., 2009) is functional in Arabidopsis, inhibits degradation of CRL substrates and confers

auxin insensitivity (Hakenjos *et al*., 2011). To test whether increased PSI-gene expressions in

ipk1-1 or *itpk1-2* are consequences of enhanced $CUL1^{Nedd8}$ levels via augmenting CRL

functions, we treated these mutants with MLN4924. Remarkably, upregulated expression of

PSI-genes such as *IPS1, SPX1,* or *PHT1;3* as well as high endogenous Pi levels in *ipk1-1* or

itpk1-2 were suppressed by MLN4924-treatments (Figure 3A-B; Supplemental Figure 4A). On

Col-0 grown under Pi-sufficient conditions, MLN4924 treatments did not cause detectable

change in endogenous Pi-levels or *IPS1* expression (Figure 3A-B). Most surprisingly,

MLN4924 treatments on Col-0 suppressed the induction of *PHT1;3* or *IPS1* when subjected to

Pi-starvation implying that neddylation processes drive PSI-gene expressions (Figure 3C).

To test whether neddylation efficiencies of cullins are affected upon Pi-deprivation, extracts

from Col-0 grown under Pi-replete or Pi-deplete conditions were immunoblotted with anti-

CUL1 antibodies. Prominent increase in $CUL1^{Nedd8}$ levels, sensitive to MLN4924 inhibition,

was noted under Pi-starvation (Figure 3D). Interestingly, these enhancements were induced

even in the *phr1phl1* plants when subjected to Pi-deplete conditions implying that neddylation

readjustments are upstream of PHR1/PHL1 role in inducing PSI-gene expressions. Already

enhanced $CUL1^{Nedd8}$ levels in *ipk1-1* or *itpk1-2* remained unaffected under Pi-deplete

exposures (Supplemental Figure 4B). Taken together, our results identify CUL1

hyperneddylation as an early response to Pi-starvation essential for transducing downstream

signaling.

As tested earlier for *ipk1-1* or *itpk1-2*, to correlate increased CUL1^{Nedd8} levels to decreased

CSN5 deneddylase activity, we performed in-lysate deneddylation assay on Pi-replete or

starved Col-0. Diminished CSN5 deneddylase activity was clearly apparent in Pi-deprived

Col-0 extracts (Figure 4A). In the same extracts CUL1 instabilities were not evident perhaps

reflecting tight regulations between its increased utilization and turnover rates. When profiled,

CSN5 elution patterns under Pi-deplete conditions were peculiarly similar to *ipk1-1* or *itpk1-2*

with more engagements on the CSN holo-complex and lower monomeric pools (Figure 4C).

Total CSN5 protein remained comparable between Pi-replete and Pi-deplete extracts. Profiles

of CSN1 remained unaltered by the Pi-changes (Supplemental Figure 4C). Resupplied

phosphate restored both CSN5 partitioning changes and increased CUL1^{Nedd8} amounts to Pi-

replete pattern suggesting that observed changes were due to Pi-starvation (Figure 4B, C)

Curiously, unlike *ipk1-1* or *itpk1-2*, global ubiquitinylation was not enhanced upon Pi-

starvation although resupplying phosphate did cause a modest increase (Figure 4D). Together,

these results suggest intricate coordination between CSN-CRL activities to maintain specificity

in responding to reduced Pi-availability. Also, our data indicates that plants modulate CSN5

deneddylase activity and affect its dynamics during Pi-deprivation to induce PSI-gene

expressions, a phenomenon that is constitutive in *ipk1-1* or *itpk1-2*. In Pi-starved *ipk1-1:Myc-*

IPK1 or *itpk1-2:ITPK1-GFP* plants, marked increase in ITPK1-GFP, but not very prominent

for Myc-IPK1 protein, was detected in comparison to non-starved extracts (Supplemental

Figure 4D). These at least are not due to increased transcriptions as reported earlier (Riemer et

al., 2020) and may instead be in accordance with increased stability the mammalian IP6K1

acquires when associated with an active CRLs (Rao *et al*., 2014).

Functional inhibition of CSN5 phenocopy *ipk1* or *itpk1* responses.

To determine whether PSI-gene expressions are globally affected by CSN functions, we mined the available microarray data of differentially expressed transcripts in *csn* null mutants (*csn3*, *csn4* and *csn5ab*) (Dohmann *et al.*, 2008). Strong upregulation in expressions compared to wild-type plants are especially noted in genes responsible for Pi-remobilization (*PAP1, PAP10, PAP23, DFR, F3H*) and several Pi-transporters (*PHT3;2, PHT3;3, OCT1*) (Supplemental Figure 5). Expressions of *PHR1* or *PHL1* remain unaffected in these mutants similar to *ipk1-1* or *itpk1-2* (Kuo *et al.*, 2014; Kuo *et al.*, 2018). Also, neither *IPK1* nor *ITPK1* transcriptions are changed in the *csn* mutants. Thus, CSN holo-complex dysfunctions affect several PSI-gene expressions likely because of its deneddylation deficiencies. To explore CSN5 roles further, we utilized the *csn5a-2* plants harboring a T-DNA insertion in the last exon of the *CSN5A* isoform (*At1g22920*). Similar to *ipk1-1* or *itpk1-2*, *csn5a-2* extracts accumulate higher CUL1^{Nedd8} levels and display auxin-insensitivity (Gusmaroli *et al.*, 2004; Dohmann *et al.*, 2005; Dohmann *et al.*, 2008; Laha *et al.*, 2020). Under Pi-replete growth, increased basal Pi-levels accompanied by mild up-regulated expressions of *PHT1;3* or *SPX1* is detected in the *csn5a-2* plants (Figure 5A-B). Also, primary root lengths of *csn5a-2* plants are slightly, but not statistically shorter than Col-0 in this growth media (Supplemental Figure 6). These features are very similar to *ipk1-1* or *itpk1-2* (Riemer *et al.*, 2020). Growth under Pi-starvation led to strong inhibition of primary root length in Col-0, while in *csn5a-2* (and in *ipk1-1* or *itpk1-2*) plants these were prominently less. With exposure to Pi-deplete conditions, *csn5a-2* plants are completely impaired in inducing *PHT1;3* or *SPX1* (Figure 5B). These results resemble PSI gene induction deficiencies in *itpk1-2* when subjected to Pi-starvation (Riemer *et al.*, 2020) suggesting that CSN5 functions are necessary in maintenance of Pi-homeostasis and successful transduction of PSR.

To directly relate CSN5 deneddylase role in Pi-homeostasis, we tested the effects of recently

reported CSN5 inhibitors Azaindole#6 and CSN5i-3 on Arabidopsis (Schlierf *et al.*, 2016;

Altmann *et al.*, 2017). In animal cell lines, these inhibitors interfere with the CSN5 deneddylase

activity, and trap CRLs in their neddylated states. Because they remain untested on any plant

systems, we first determined their efficacy on Col-0 seedlings. Parallel MLN4924 treatments

were used as neddylation-inhibition controls. Dose-dependent increments in CUL1^{Nedd8} levels

with collateral decrease in unneddylated CUL1 was detected at 24hr post-treatment with

CSN5i-3 (Figure 5C). At 48-hrs post-treatment, efficiency of CSN5i-3 inhibitions intensified

(Supplemental Figure 7A). At all doses tested, Azaindole#6 remained ineffective in changing

CUL1^{Nedd8}:CUL1 ratios. As expected, MLN4924 treatments in a dose-dependent manner

reduced CUL1^{Nedd8} levels. For subsequent experiments, we continued with 10μM dose of

CSN5i-3 and the same for MLN4924, as used in our earlier assays. On Pi-replete Col-0, CSN5i-

3 caused a modest increase in basal Pi levels and upregulation in expression of *PHT1;3*, but

not significantly for *SPX1* or *IPS1* (Figure 5D-E; Supplemental Figure 7B-C). Under Pi-deplete

exposure, CSN5i-3 suppressed the induction of *PHT1;3, SPX1* or *IPS1* in Col-0 and mirrored

MLN4924 effects (Figure 5D-E; Supplemental Figure 7B-C). CSN5i-3-treated Col-0 extracts

when fractionated, CSN5 was markedly enriched in the CSN holo-complex pools with

negligible detection in monomeric elutions, a pattern resembling *ipk1-1*, *itpk1-2,* or Pi-starved

profiles (Figure 6A). With these results, we reason strong correlation between CSN5

deneddylation deficiencies to its enhanced retention on the CSN holo-complex, and trigger of

PSR as observed in *ipk1-1* or *itpk1-2* plants. Most interestingly, *ipk1-1* or *itpk1-2* seedlings

grown under sufficient Pi when treated with CSN5i-3 had reduced *IPS1* expressions than

untreated plants indicating that skewed kinetics of CSN5 activity rather than its complete

inhibition were attributed to these mutants (Supplemental Figure 7D). MLN4924-treatments

on *ipk1-1* or *itpk1-2* in Pi-deplete medium offered relief by elevating *IPS1* expressions further

supporting our hypothesis that inadequacies in these mutants to support PSR is related to

dynamics of CSN5-regulated neddylation/deneddylation cycles of CRLs (Supplemental Figure

7E). From these data, it is suggestive that IPK1/ITPK1 functions influence deneddylase

efficiencies and shuttling modes of CSN5 thereby affecting CRL requirements in executing

PSR.

IP6K1-like activity, but not InsP$_6$ regulates CSN5 deneddylase functions of CSN.

InsP$_6$ is implicated as the physiological glue in potentiating CSN2-CRL associations (Scherer

et al., 2016). That a kinase-dead IP5K interacts weakly with CSN2/5 further indicates that InsP$_6$

augmenting these interactions is produced locally. Supplementing InsP$_6$ to IP5K-depleted cell

lysates strengthens CRL-CSN2 associations, and increases deneddylation activities of

recombinant CSN on purified neddylated cullins (Scherer *et al.*, 2016). Apparently, the

externally added metabolite gets access its designated binding pocket at the CRL-CSN2

interface. Even though CRL-CSN associations are not affected in *ipk1-1* or *itpk1-2*, we

investigated whether adding InsP$_6$ to lysates recover deneddylation deficiencies in these

mutants, or augments it in Col-0. In lysate deneddylation assays showed that supplemented

InsP$_6$ did not alter deneddylation deficiencies or CUL1 instability in *ipk1-1* or *itpk1-2*

(Supplemental Figure 8). In similar assays on Col-0 extracts, InsP$_6$ addition did not change

deneddylation or CUL1 turnover rates indicating that unlike mammalian systems, CSN

functions in plants are not influenced by InsP$_6$.

As a CRL-CSN release factor, InsP$_7$ when supplemented to mammalian cell extracts does not

cause complex disassembly meaning that IP6K1 activity is very tightly regulated

spatiotemporally (Rao *et al.*, 2014). With observed ITPK1-CSN2 interactions, this instigated

us to investigate for a similar mechanism in plants. The cell-permeable compound N2-(m-

(trifluoromethyl)benzyl) N6-(p-nitrobenzyl) purine (TNP) inhibits animal IP6K1 functions and

reduce $InsP_7$ levels (Padmanabhan *et al*., 2009). TNP effects so far have remained untested on

Arabidopsis InsPs. In extracts of 10μM TNP-treated Col-0, distinct increase in $CUL1^{NEDD8}$

levels were observed although less potent than a similar dose of CSN5i-3 (Figure 6B). To

improve TNP efficacies, we used 25μM doses for subsequent assays. In Col-0 grown in Pi-

sufficient media, TNP presence replicated CSN5i-3 results in augmenting basal Pi levels,

upregulation of *IPS1* or *PHT1;3* expressions, and more retention of CSN5 in the CSN holo-

complex pools (Figure 6A-C-D). TNP also suppressed endogenous deneddylation rates with

increased CUL1 instability (Figure 6E). And likewise, to CSN5i-3, TNP also suppressed PSI-

gene inductions in Col-0 under Pi-deplete circumstances (Figure 6D). Curiously, CSN5i-3,

TNP or MLN4924 treatments on *itpk1-2:ITPK1-GFP* resulted in increased ITPK1-GFP levels,

similar to observed in Pi-deplete media (Supplemental Figure 9). Although direct inhibitions

of TNP on Arabidopsis ITPK1 or other $InsP_{(6)}$-kinases still remains to be tested, nevertheless

our results imply that similar to mammals an IP6K1-like activity affects CSN5 performance on

regulating CUL1 deneddylation rates and its consequences on PSR. Overall, with our data we

present localized coordinated involvement of InsP-kinase pair in modulating CSN dynamics,

defining molecular concept of the CSN paradox, and underlining CSN role in phosphate stress

responses of plants.

Discussion

In animal systems, CRL-CSN tether is augmented by $InsP_6$ and released via stimulus-driven

$InsP_7$ synthesis. Here, we show that although Arabidopsis IPK1 and ITPK1 retain conserved

CSN interactions, $InsP_6/InsP_7$ changes in the corresponding mutants do not overall affect

CRL1-CSN associations. Thus, either the residual levels of these InsPs suffice at the CRL-

CSN junctures or the association/dissociation cycles involve an altogether different InsP(s) not

locally altered in the above mutants. Unlike mammalian results, that added $InsP_6$ does not

rescue *ipk1-1* or even alter CUL1 deneddylation rates/stability in Col-0 supports our

speculation. Further, an *ITPK4* mutant (*itpk4-1*) with reduced $InsP_6/InsP_7$ as in *ipk1-1* or *itpk1-*

2 has normal PSR (Kuo *et al*., 2018) and matches Col-0 ratio of $CUL1^{Nedd8}$:CUL1 also lends

credence to our hypothesis. One striking distinction in *ipk1-1* or *itpk1-2* is that these mutants

accumulate increased levels of $Ins(3,4,5,6)P_4$ which neither the *itpk4-1* nor Col-0 subjected to

PSR does (Kuo *et al*., 2018). Whether this increased abundance of $Ins(3,4,5,6)P_4$ or another

unidentified InsP altered in *ipk1-1* or *itpk1-2* satisfy CRL-CSN interaction requirements

remains a possibility to explore further. Nevertheless, analogous to mammals our observations

overall parallel that decreased CSN5 deneddylase performance with lower CUL1

deneddylation rates associate with the loss of IPK1 or ITPK1 and implicates their distinct roles

on CSN functions.

Mammalian ITPK1 over-expression partitions more monomeric CSN5 free from the CSN

holo-complex (Sun *et al*., 2002). We demonstrate here that absence of Arabidopsis ITPK1 or

TNP inhibitions antagonize CSN5 release from the CSN holo-complex. Even with yet to be

tested TNP effects on ITPK1 (or another InsP-kinase) functions in plants, nonetheless a

mammalian IP6K1-like activity in choreographing CSN5 entry-exit shuttles seems to be

conserved in Arabidopsis. Overall structural integrity of an inert CSN holo-complex is

remarkably maintained without CSN5, albeit for deneddylation CSN5 incorporation is

mandatory (Lingaraju *et al*., 2014). When CSN5 is integrated into the holo-complex, its

interfaces with CSN4-CSN6 inhibit deneddylase activation in the absence of a bound CRL

(Enchev *et al*., 2012; Lingaraju *et al*., 2014). Upon binding a neddylated CRL, a series of

conformational change prompted by CSN4 orients CSN5 active site towards the neddylated

cullins. Closely proximal to this active site, conserved lysines in CSN2 alpha helices coordinate

bound $InsP_6$ to bridge for more stable associations with CRL. The bound $InsP_6$ in this

electropositive environment is uniquely oriented with its 5'-phosphate exposed, amenable for

subsequent pyrophosphorylation. Contrastingly, when an $InsP_7$ is modeled into the same active

site strong steric clashes especially at CRL-CSN junctures are noted (Lin *et al.*, 2020). These

steric intolerances may underlie $InsP_7$-triggered CSN5 deneddylase activation, followed by

CRL and perhaps CSN5 release from the CSN holo-complex. When added to cell lysates, since

free $InsP_7$ fails to enhance cullin neddylation or disrupt CRL-CSN associations (Rao *et al.*,

2014), its localized synthesis achieved through site-specific availability of $InsP_6$, and its

accessibility to CSN5 active site is hence crucial for its proposed role. Thus, similar to the

mammalian IP6K1 a plant ITPK1 activity (via $InsP_7$ or an unknown InsP) may possess

propensity to reconfigure CSN, initiating deneddylation coupled to CRLs and CSN5

disassembly from the holo-complex. These roles may be supportive of the recent implication

that a plant ITPK1 has evolutionarily integrated functional signatures of both mammalian

ITPK1 and IP6K1 (Whitfield *et al.*, 2020).

With obvious $InsP_8$ reductions in *ipk1-1*, *itpk1-2* or upon VIH2 phosphatase activity during Pi-

starvation (Dong *et al.*, 2019; Zhu *et al.*, 2019), PHR1 liberated from SPX1 causes activation

of PSI-gene expressions. Here we demonstrate that a mode of perception of Pi-deprivation in

addition to the above also occurs at CSN, involving changes in associated InsP-kinase

activities, and is responded by decreased deneddylation rates with pronounced CSN5 retentions

on the CSN holo-complex. Since the resulting increase in $CUL1^{Nedd8}$ levels remain independent

of PHR1/PHL1 presence, these observations likely indicate parallel responsive routes of a Pi-

starved plant to facilitate more available pools of neddylated CRLs to ubiquitinylate substrates

including the recently identified SPX4, released from PHR1-hold (Osorio *et al.*, 2019). These

features we show are constitutive in *ipk1-1*, *itpk1-2*, and mimicked in wild-type plants with

CSN5i-3 or TNP treatment, and suppressed by the neddylation inhibitor MLN4924. Our

observation that phosphate re-supplementation post-starvation restores CSN5 partitioning and

CUL^{Nedd8}:CUL1 ratios to Pi-replete patterns in wild-type plants implies that the noted shifts

are only transitory adaptations. That, unlike in *ipk1-1* or *itpk1-2*, global polyubiquitin-

conjugates remains unchanged in Col-0 under low-Pi availability additionally emphasize on

the strictly regulatory PSR mechanisms to controls ubiquitinylation-specificities on intended

targets. Not the least, observed failure in *itpk1-2* (Riemer *et al.*, 2020) or *csn5a-2* to respond to

low-Pi by anticipated augmentation of already upregulated PSI-gene expressions, or in CSN5i-

3 or TNP treated Col-0 indicates that CSN5 partitioning re-adjustments are only shifts in

neddylation/deneddylation equilibrium and not a complete inhibition of the CSN5

deneddylase. Also, that MLN4924 presence causes sub-optimal induction of PSI-gene

expressions in Pi-depleted Col-0, suggests continued requirements of CRL

neddylation/deneddylation cycles to successfully transduce PSR. Thus, sustained and

successful response to Pi-starvation is likely achieved by a shift in equilibrium of CSN5-

mediated deneddylation dynamics on CUL1 and not complete inhibition of its activities as

described in our model (Figure 7). Similar phenomenon is mirrored by the shuttling

requirements of CSN5 in cell proliferation wherein its perturbations cause complete arrest

(Tomoda *et al.*, 2002; Yoshida *et al.*, 2010).

Distinct increase in ITPK1 protein but not gene expression under Pi-starvation (Riemer *et al.*,

2020) and as shown here with CSN5i-3, TNP or MLN4924 treatments is conceivably similar

to the gain of stability the mammalian IP6K1 acquires upon CRL binding (Rao *et al.*, 2014).

Thus, similar to CUL1 vulnerability for self-ubiquitinylation, ITPK1 fates may also be

regulated in the absence of CRL substrates. Increased stabilities hence may therefore reflect

ITPK1's priority to accommodate stimulus-appropriate co-factor requirements in downstream

responses not limited only for modulating CRL functions. The mammalian phosphate exporter

Xenotropic and polytropic retrovirus receptor 1 (XPR1) relies on $InsP_7$ synthesized by IP6K1/2

for function (Wilson *et al.*, 2019). Arabidopsis XPR1 ortholog PHO1 does not complement the

pho1 mutant if mutations are introduced in its InsP$_7$-binding pocket (Wild *et al.*, 2016). And as

is known during PSR, augmented auxin responses involving SCF^{TIR1} also require InsP$_7$ (Laha

et al., 2020). Whether these new engagements of ITPK1 are adaptation strategies to transiently

deprive CSN (and CSN5 release from the holo-complex) and shift balances towards more

neddylated CRL availability and/or for providing InsP$_7$ for other cellular processes are

encouraging possibility to explore. In conclusion, with our data we highlight remarkable

conservation, albeit with slight distinction, in plant InsP-kinase roles in regulating CRL-CSN

functions and its implications in responding to Pi-imbalances.

References

Adepoju O, Williams SP, Craige B, Cridland CA, Sharpe AK, Brown AM, Land E, Perera IY,
Mena D, Sobrado P, Gillaspy GE (2019) bioRxiv**:724914**; doi:
https://doi.org/10.1101/724914
Altmann E, Erbel P, Renatus M, Schaefer M, Schlierf A, Druet A, Kieffer L, Sorge M, Pfister K,
Hassiepen U, Jones M, Ruedisser S, Ostermeier D, Martoglio B, Jefferson AB,
Quancard J (2017) Azaindoles as Zinc-Binding Small-Molecule Inhibitors of the JAMM
Protease CSN5. Angew Chem Int Ed Engl **56:** 1294-1297
Bhattacharjee S, Halane MK, Kim SH, Gassmann W (2011) Pathogen effectors target
Arabidopsis EDS1 and alter its interactions with immune regulators. Science **334:** 1405-
1408
Bustos R, Castrillo G, Linhares F, Puga MI, Rubio V, Perez-Perez J, Solano R, Leyva A, Paz-
Ares J (2010) A central regulatory system largely controls transcriptional activation and
repression responses to phosphate starvation in Arabidopsis. PLoS Genet **6:** e1001102
Chakrabarty R, Banerjee R, Chung SM, Farman M, Citovsky V, Hogenhout SA, Tzfira T,
Goodin M (2007) pSITE vectors for stable integration or transient expression of
autofluorescent protein fusions in plants: probing *Nicotiana benthamiana*-virus
interactions. Mol Plant Microbe Interact **20:** 740-750
Chamovitz DA, Segal D (2001) JAB1/CSN5 and the COP9 signalosome. A complex situation.
EMBO Reports **2:** 96-101
Chiou TJ, Aung K, Lin SI, Wu CC, Chiang SF, Su CL (2006) Regulation of phosphate homeostasis
by MicroRNA in Arabidopsis. Plant Cell **18:** 412-421
Cope GA, Suh GS, Aravind L, Schwarz SE, Zipursky SL, Koonin EV, Deshaies RJ (2002) Role of
predicted metalloprotease motif of Jab1/Csn5 in cleavage of Nedd8 from Cul1. Science
298: 608-611
Desai M, Rangarajan P, Donahue JL, Williams SP, Land ES, Mandal MK, Phillippy BQ, Perera
IY, Raboy V, Gillaspy GE (2014) Two inositol hexakisphosphate kinases drive inositol
pyrophosphate synthesis in plants. Plant J **80:** 642-653
Dohmann EM, Kuhnle C, Schwechheimer C (2005) Loss of the CONSTITUTIVE
PHOTOMORPHOGENIC9 signalosome subunit 5 is sufficient to cause the cop/det/fus
mutant phenotype in Arabidopsis. Plant Cell **17:** 1967-1978

Dohmann EM, Levesque MP, Isono E, Schmid M, Schwechheimer C (2008) Auxin responses in
mutants of the Arabidopsis CONSTITUTIVE PHOTOMORPHOGENIC9 signalosome. Plant
Physiol **147**: 1369-1379

Dong J, Ma G, Sui L, Wei M, Satheesh V, Zhang R, Ge S, Li J, Zhang TE, Wittwer C, Jessen HJ,
Zhang H, An GY, Chao DY, Liu D, Lei M (2019) Inositol Pyrophosphate InsP$_8$ Acts as an
Intracellular Phosphate Signal in Arabidopsis. Mol Plant **12**: 1463-1473

Dubiel D, Rockel B, Naumann M, Dubiel W (2015) Diversity of COP9 signalosome structures
and functional consequences. FEBS Lett **589**: 2507-2513

Dubiel W (2009) Resolving the CSN and CAND1 paradoxes. Mol Cell **35**: 547-549

Enchev RI, Scott DC, da Fonseca PC, Schreiber A, Monda JK, Schulman BA, Peter M, Morris
EP (2012) Structural basis for a reciprocal regulation between SCF and CSN. Cell Rep **2**:
616-627

Gillaspy GE (2013) The role of phosphoinositides and inositol phosphates in plant cell signaling.
Adv Exp Med Biol **991**: 141-157

Gusmaroli G, Feng S, Deng XW (2004) The Arabidopsis CSN5A and CSN5B subunits are present
in distinct COP9 signalosome complexes, and mutations in their JAMM domains exhibit
differential dominant negative effects on development. Plant Cell **16**: 2984-3001

Hakenjos JP, Richter R, Dohmann EM, Katsiarimpa A, Isono E, Schwechheimer C (2011)
MLN4924 is an efficient inhibitor of NEDD8 conjugation in plants. Plant Physiol **156**: 527-
536

Ham BK, Chen J, Yan Y, Lucas WJ (2018) Insights into plant phosphate sensing and signaling.
Curr Opin Biotechnol **49**: 1-9

Huang H, Quint M, Gray WM (2013) The *eta7/csn3-3* auxin response mutant of Arabidopsis
defines a novel function for the CSN3 subunit of the COP9 signalosome. PLoS One **8**:
e66578

Jia Q, Kong D, Li Q, Sun S, Song J, Zhu Y, Liang K, Ke Q, Lin W, Huang J (2019) The Function of
Inositol Phosphatases in Plant Tolerance to Abiotic Stress. Int J Mol Sci **20**

Jia X, Chanda B, Zhao M, Brunner AM, Beers EP (2015) Instability of the Arabidopsis mutant
csn5a-2 caused by epigenetic modification of intronic T-DNA. Plant Sci **238**: 53-63

Jung JY, Ried MK, Hothorn M, Poirier Y (2018) Control of plant phosphate homeostasis by
inositol pyrophosphates and the SPX domain. Curr Opin Biotechnol **49**: 156-162

Kim SH, Gao F, Bhattacharjee S, Adiasor JA, Nam JC, Gassmann W (2010) The Arabidopsis
resistance-like gene SNC1 is activated by mutations in SRFR1 and contributes to
resistance to the bacterial effector AvrRps4. PLoS Pathog **6**: e1001172

Kuo HF, Chang TY, Chiang SF, Wang WD, Charng YY, Chiou TJ (2014) Arabidopsis inositol
pentakisphosphate 2-kinase, AtIPK1, is required for growth and modulates phosphate
homeostasis at the transcriptional level. Plant J **80**: 503-515

Kuo HF, Hsu YY, Lin WC, Chen KY, Munnik T, Brearley CA, Chiou TJ (2018) Arabidopsis
inositol phosphate kinases IPK1 and ITPK1 constitute a metabolic pathway in
maintaining phosphate homeostasis. Plant J **95**:613-630

Laha D, Johnen P, Azevedo C, Dynowski M, Weiss M, Capolicchio S, Mao H, Iven T,
Steenbergen M, Freyer M, Gaugler P, de Campos MK, Zheng N, Feussner I, Jessen HJ,
Van Wees SC, Saiardi A, Schaaf G (2015) VIH2 Regulates the Synthesis of Inositol
Pyrophosphate InsP$_8$ and Jasmonate-Dependent Defenses in Arabidopsis. Plant Cell **27**:
1082-1097

Laha D, Parvin N, Hofer A, Giehl RFH, Fernandez-Rebollo N, von Wiren N, Saiardi A, Jessen
HJ, Schaaf G (2019) Arabidopsis ITPK1 and ITPK2 Have an Evolutionarily Conserved
Phytic Acid Kinase Activity. ACS Chem Biol **14**: 2127-2133

Laha NP, Dhir YW, Giehl RFH, Schäfer EM, Gaugler P, Shishavan ZH, Gulabani H, Mao H,
Zheng N, von Wirén N, Jessen HJ, Saiardi A, Bhattacharjee S, Laha D, Schaaf G (2020)
bioRxiv: 2020.04.23.058487

Lee HS, Lee DH, Cho HK, Kim SH, Auh JH, Pai HS (2015) InsP$_6$-sensitive variants of the Gle1
mRNA export factor rescue growth and fertility defects of the *ipk1* low-phytic-acid
mutation in Arabidopsis. Plant Cell **27**: 417-431

Lin H, Zhang X, Liu L, Fu Q, Zang C, Ding Y, Su Y, Xu Z, He S, Yang X, Wei X, Mao H, Cui Y, Wei Y, Zhou C, Du L, Huang N, Zheng N, Wang T, Rao F (2020) Basis for metabolite-dependent Cullin-RING ligase deneddylation by the COP9 signalosome. Proc Natl Acad Sci U S A **117:** 4117-4124

Lingaraju GM, Bunker RD, Cavadini S, Hess D, Hassiepen U, Renatus M, Fischer ES, Thoma NH (2014) Crystal structure of the human COP9 signalosome. Nature **512:** 161-165

Lv Q, Zhong Y, Wang Y, Wang Z, Zhang L, Shi J, Wu Z, Liu Y, Mao C, Yi K, Wu P (2014) SPX4 Negatively Regulates Phosphate Signaling and Homeostasis through Its Interaction with PHR2 in Rice. Plant Cell **26:** 1586-1597

Monserrate JP, York JD (2010) Inositol phosphate synthesis and the nuclear processes they affect. Curr Opin Cell Biol **22:** 365-373

Mosadeghi R, Reichermeier KM, Winkler M, Schreiber A, Reitsma JM, Zhang Y, Stengel F, Cao J, Kim M, Sweredoski MJ, Hess S, Leitner A, Aebersold R, Peter M, Deshaies RJ, Enchev RI (2016) Structural and kinetic analysis of the COP9-Signalosome activation and the cullin-RING ubiquitin ligase deneddylation cycle. Elife **5:** e12102

Murphy AM, Otto B, Brearley CA, Carr JP, Hanke DE (2008) A role for inositol hexakisphosphate in the maintenance of basal resistance to plant pathogens. Plant J **56:** 638-652

Osorio MB, Ng S, Berkowitz O, De Clercq I, Mao C, Shou H, Whelan J, Jost R (2019) SPX4 Acts on PHR1-Dependent and -Independent Regulation of Shoot Phosphorus Status in Arabidopsis. Plant Physiol **181:** 332-352

Padmanabhan U, Dollins DE, Fridy PC, York JD, Downes CP (2009) Characterization of a selective inhibitor of inositol hexakisphosphate kinases: use in defining biological roles and metabolic relationships of inositol pyrophosphates. J Biol Chem **284:** 10571-10582

Puga MI, Mateos I, Charukesi R, Wang Z, Franco-Zorrilla JM, de Lorenzo L, Irigoyen ML, Masiero S, Bustos R, Rodriguez J, Leyva A, Rubio V, Sommer H, Paz-Ares J (2014) SPX1 is a phosphate-dependent inhibitor of Phosphate Starvation Response 1 in Arabidopsis. Proc Natl Acad Sci U S A **111:** 14947-14952

Qin ZX, Chen QJ, Tong Z, Wang XC (2005) The Arabidopsis inositol 1,3,4-trisphosphate 5/6 kinase, AtItpk-1, is involved in plant photomorphogenesis under red light conditions, possibly via interaction with COP9 signalosome. **43:** 947-954

Rao F, Xu J, Khan AB, Gadalla MM, Cha JY, Xu R, Tyagi R, Dang Y, Chakraborty A, Snyder SH (2014) Inositol hexakisphosphate kinase-1 mediates assembly/disassembly of the CRL4-signalosome complex to regulate DNA repair and cell death. Proc Natl Acad Sci U S A **111:** 16005-16010

Ried MK, Wild R, Zhu J, Broger L, Harmel RK, Hothorn LA, Fiedler D, Hothorn M (2019) Inositol pyrophosphates promote the interaction of SPX domains with the coiled-coil motif of PHR transcription factors to regulate plant phosphate homeostasis. bioRxiv 2019.12.13.875393

Riemer E, Laha D, Harmel RK, Gaugler P, Pries V, Frei M, Hajirezaei M-R, Laha NP, Krusenbaum L, Schneider R, Jessen HJ, Saiardi A, Fiedler D, Schaaf G, Giehl RFH (2020) ITPK1 is an $InsP_6$/ADP phosphotransferase that controls systemic phosphate homeostasis in Arabidopsis. bioRxiv 2020.05.18.100297

Ruan W, Guo M, Wang X, Guo Z, Xu Z, Xu L, Zhao H, Sun H, Yan C, Yi K (2019) Two RING-Finger Ubiquitin E3 Ligases Regulate the Degradation of SPX4, An Internal Phosphate Sensor, for Phosphate Homeostasis and Signaling in Rice. Mol Plant **12:** 1060-1074

Rubio V, Linhares F, Solano R, Martin AC, Iglesias J, Leyva A, Paz-Ares J (2001) A conserved MYB transcription factor involved in phosphate starvation signaling both in vascular plants and in unicellular algae. Genes Dev **15:** 2122-2133

Scherer PC, Ding Y, Liu Z, Xu J, Mao H, Barrow JC, Wei N, Zheng N, Snyder SH, Rao F (2016) Inositol hexakisphosphate (IP6) generated by IP5K mediates cullin-COP9 signalosome interactions and CRL function. Proc Natl Acad Sci U S A **113:** 3503-3508

Schlierf A, Altmann E, Quancard J, Jefferson AB, Assenberg R, Renatus M, Jones M, Hassiepen U, Schaefer M, Kiffe M, Weiss A, Wiesmann C, Sedrani R, Eder J, Martoglio

B (2016) Targeted inhibition of the COP9 signalosome for treatment of cancer. Nat
Commun **7:** 13166

Sheard LB, Tan X, Mao H, Withers J, Ben-Nissan G, Hinds TR, Kobayashi Y, Hsu FF, Sharon
M, Browse J, He SY, Rizo J, Howe GA, Zheng N (2010) Jasmonate perception by inositol-
phosphate-potentiated COI1-JAZ co-receptor. Nature **468:** 400-405

Shears SB, Ganapathi SB, Gokhale NA, Schenk TM, Wang H, Weaver JD, Zaremba A, Zhou Y
(2012) Defining signal transduction by inositol phosphates. Subcell Biochem **59:** 389-412

Shears SB, Wang H (2019) Inositol phosphate kinases: Expanding the biological significance of
the universal core of the protein kinase fold. Adv Biol Regul **71:** 118-127

Soucy TA, Smith PG, Milhollen MA, Berger AJ, Gavin JM, Adhikari S, Brownell JE, Burke KE,
Cardin DP, Critchley S, Cullis CA, Doucette A, Garnsey JJ, Gaulin JL, Gershman RE,
Lublinsky AR, McDonald A, Mizutani H, Narayanan U, Olhava EJ, Peluso S, Rezaei M,
Sintchak MD, Talreja T, Thomas MP, Traore T, Vyskocil S, Weatherhead GS, Yu J,
Zhang J, Dick LR, Claiborne CF, Rolfe M, Bolen JB, Langston SP (2009) An inhibitor of
NEDD8-activating enzyme as a new approach to treat cancer. Nature **458:** 732-736

Stevenson-Paulik J, Bastidas RJ, Chiou ST, Frye RA, York JD (2005) Generation of phytate-free
seeds in Arabidopsis through disruption of inositol polyphosphate kinases. Proc Natl
Acad Sci U S A **102:** 12612-12617

Sun Y, Wilson MP, Majerus PW (2002) Inositol 1,3,4-trisphosphate 5/6-kinase associates with
the COP9 signalosome by binding to CSN1. J Biol Chem **277:** 45759-45764

Szijgyarto Z, Garedew A, Azevedo C, Saiardi A (2011) Influence of inositol pyrophosphates on
cellular energy dynamics. Science **334:** 802-805

Tan X, Calderon-Villalobos LI, Sharon M, Zheng C, Robinson CV, Estelle M, Zheng N (2007)
Mechanism of auxin perception by the TIR1 ubiquitin ligase. Nature **446:** 640-645

Tomoda K, Kubota Y, Arata Y, Mori S, Maeda M, Tanaka T, Yoshida M, Yoneda-Kato N, Kato
JY (2002) The cytoplasmic shuttling and subsequent degradation of p27[Kip1] mediated by
Jab1/CSN5 and the COP9 signalosome complex. J Biol Chem **277:** 2302-2310

Whitfield H, White G, Sprigg C, Riley AM, Potter BVL, Hemmings AM, Brearley CA (2020) An
ATP-responsive metabolic cassette comprised of inositol tris/tetrakisphosphate kinase 1
(ITPK1) and inositol pentakisphosphate 2-kinase (IPK1) buffers diphosphoinositol
phosphate levels. Biochem J **477:** 2621-2638

Wild R, Gerasimaite R, Jung JY, Truffault V, Pavlovic I, Schmidt A, Saiardi A, Jessen HJ,
Poirier Y, Hothorn M, Mayer A (2016) Control of eukaryotic phosphate homeostasis by
inositol polyphosphate sensor domains. Science **352:** 986-990

Williams SP, Gillaspy GE, Perera IY (2015) Biosynthesis and possible functions of inositol
pyrophosphates in plants. Front Plant Sci **6:** 67

Wilson MS, Jessen HJ, Saiardi A (2019) The inositol hexakisphosphate kinases IP6K1 and -2
regulate human cellular phosphate homeostasis, including XPR1-mediated phosphate
export. J Biol Chem **294:** 11597-11608

Wundenberg T, Mayr GW (2012) Synthesis and biological actions of diphosphoinositol
phosphates (inositol pyrophosphates), regulators of cell homeostasis. Biol Chem **393:**
979-998

Yi Sun WW, Jianping Jin (2020) CRL Book Sun et al 2020. Advances in Experimental Medicine
and Biology Biology and Therapeutics **1217**

Yoshida A, Yoneda-Kato N, Panattoni M, Pardi R, Kato JY (2010) CSN5/Jab1 controls multiple
events in the mammalian cell cycle. FEBS Lett **584:** 4545-4552

Zhong Y, Wang Y, Guo J, Zhu X, Shi J, He Q, Liu Y, Wu Y, Zhang L, Lv Q, Mao C (2018) Rice SPX6
negatively regulates the phosphate starvation response through suppression of the
transcription factor PHR2. **219:** 135-148

Zhu J, Lau K, Puschmann R, Harmel RK, Zhang Y, Pries V, Gaugler P, Broger L, Dutta AK,
Jessen HJ, Schaaf G, Fernie AR, Hothorn LA, Fiedler D, Hothorn M (2019) Two
bifunctional inositol pyrophosphate kinases/phosphatases control plant phosphate
homeostasis. Elife **8:** e43582

 Figure legends

 Fig. 1. IPK1 and ITPK1 interact with each other and with selective CSN or CRLs

 components *in planta*. *(A)* Immuno-enrichments of IPK1 *(top panel)* or ITPK1 *(bottom panel)*

 from corresponding complemented lines (*ipk1-1:Myc-IPK1* or *itpk1-2:ITPK1-GFP*,

 respectively) was probed with anti-CSN1/2/4/5, anti-CUL1, or anti-ASK1 antibodies. IgG-

 agarose enrichments were used as negative controls. *(B)* Bimolecular Fluorescence

 Complementation (BiFC) assays showing *in planta* IPK1 or ITPK1 interaction with CSN1/2

 and with each other. Indicated binary vectors combinations were co-expressed via

 Agrobacterium in *Nicotiana benthamiana* leaves and fluorescence observed under a confocal

 microscope with YFP filter. Scale bar=50 μm. *(C)* Co-immunoprecipitation of transiently

 expressed GFP-ITPK1 with Myc-IPK1 from *N. benthamiana* leaves. Anti-GFP pulldown was

 probed with anti-GFP or anti-Myc antibodies.

 Fig. 2. CSN5 deneddylase deficiencies with increased holo-complex retention in *ipk1-1* or

 ***itpk1-2* cause enhanced CUL1^{Nedd8} levels.** *(A)* IPK1 or ITPK1 does not co-elute with the CSN

 holo-complex. *(B)* Enhanced CSN5 engagements on the CSN holo-complex in *ipk1-1* and

 itpk1-2. Extracts from marked plants fractionated by size-exclusion chromatography were

 immuno-blotted with indicated antibodies. Fraction numbers and elution positions of molecular

 weight standards (in kDa) are shown. Fractions corresponding to CSN holo-complex and CSN

 subunit monomeric elutions are marked. *(C)* Immuno-enrichment of CUL1 detects

 stoichiometrically similar amounts of CSN1 and CSN2 in *ipk1-1* or *itpk1-2* to Col-0. Pulldowns

 with either IgG or anti-CUL1 antibodies were probed with anti-CUL1, anti-CSN1 or anti-

 CSN2 antibodies. *(D)* Increased CUL1^{Nedd8} levels (*top panel*) and enhanced global

 polyubiquitin-conjugates (*bottom panel*) in *ipk1-1* or *itpk1-2*. Total protein extracts were

 probed with anti-CUL1, anti-actin, or anti-Ubiquitin antibodies as indicated. *(E)* CSN5-

 deneddylation deficiencies and lower CUL1 half-life in *ipk1-1* or *itpk1-2*. In-lysate CUL1

deneddylation assay (Lin *et al*., 2020) at the indicated time points (minutes post-incubation;

mpi) probed with anti-CUL1 antibodies. Migration position of molecular weight standards (in

kDa) and neddylated (CUL1^{Nedd8}) or unneddylated (CUL1) Cullin1 are shown. Anti-actin or

Ponceau S staining demonstrates loadings controls.

Fig. 3. Enhanced CUL1^{Nedd8} levels associated with PSR is essential for PSI-gene

inductions. MLN4924 suppresses *(A)* elevated PSI-gene expressions, and *(B)* higher

endogenous Pi-levels in *ipk1-1* or *itpk1-2*. *(C)* Induction of *PHT1;3* or *IPS1* expressions in Col-

0 upon Pi-starvation is inhibited by MLN4924. *(D)* Increments in CUL1^{Nedd8} levels during Pi-

deprivation is PHR1/PHL1-independent. Plants grown in Pi-replete medium or transferred to

Pi-deplete medium were treated with or without MLN4924. Gene expression changes were

relative to *MON1* transcripts and reported as fold-change to Col-0 levels. Data is mean $\pm$ SD

(n=3 for qPCRs; n=6-7 for Pi-estimations). Statistical significance indicated with different

alphabets is according to post-hoc Tukey's test (p<0.05). For *PHT1;3* or *SPX1* expression data,

statistical comparisons were to their respective DMSO-treated Col-0. For immunoblots, anti-

CUL1 antibodies were used. Ponceau S staining shows loadings controls.

Fig. 4. Pi-starvation reduces endogenous CSN5 deneddylase activity and is restored upon

phosphate resupply. *(A)* Extracts at indicated time points (minutes post-incubation; mpi) from

in-lysate CUL1 deneddylation assay on Col-0 from Pi-replete or Pi-deplete media were probed

with anti-CUL1 antibodies. *(B)* Restoration of enhanced CUL1^{Nedd8} levels occur upon Pi-

resupply post-starvation. *(C)* Increased retention of CSN5 on the CSN holo-complex under Pi-

deplete exposure is reversed with Pi-resupply. *(D)* Global levels of polyubiquitin-conjugates

remain unchanged during PSR. For these assays, plants were subjected to Pi-starvation for 3

days and phosphate re-supplementation was done for 12hrs. Comparable loadings are shown

by Ponceau S staining. Elution positions of molecular weight standards (in kDa) are indicated.

Fig. 5. Inhibition of CSN5 functions activates PSR under Pi-replete but confers non-

responsiveness under Pi-deplete growth. *csn5a-2* plants have *(A)* elevated endogenous Pi

levels, *(B)* increased *PHT1;3* or *SPX1* transcript levels under Pi-replete growth but suppressed

expression upon Pi-starvation. *(C)* CSN5i-3 inhibits Arabidopsis CUL1 deneddylation, *(D)*

elevates endogenous Pi-levels, and *(E)* suppresses *PHT1;3* induction during phosphate-

deprivation. Immunoblots were probed with anti-CUL1 antibodies. Ponceau S staining shows

loading controls. Inhibitor treatments were done for 24 hrs and doses used are indicated.

Expression levels of *MON1* transcripts was used as an internal reference for qPCRs and

reported as fold-change relative to DMSO-treated Col-0 under Pi-replete growth. Error bars

are mean $\pm$ SD (n=3 for qPCRs; n=6-7 for Pi-estimations). Different alphabets denote statistical

significance determined by post-hoc Tukey's test (p<0.05). *PHT1;3* and *SPX1* data are

compared to respective Pi-replete Col-0.

Fig. 6. CSN5i-3 or TNP treatments on Col-0 mirrors *ipk1-1* or *itpk1-2* properties. *(A)*

CSN5 partition shifts to CSN holo-complex pools upon CSN5i-3 or TNP treatment. TNP *(B)*

enhances pools of CUL1^{Nedd8}, *(C)* increases endogenous Pi-content, *(D)* upregulates *IPS1* or

PHT1;3 expressions under Pi-replete, but suppresses them under Pi-deplete growth, and *(E)*

decreases endogenous deneddylation rates with increased CUL1 instability. Immunoblots were

probed with anti-CUL1 or anti-CSN5 antibodies as indicated. Ponceau S staining shows

loading controls. Expression levels of *MON1* transcripts was used as internal reference for

qPCRs. Error bars are mean $\pm$ SD (n=3 for qPCRs; n=6-7 for Pi-estimations). Statistical

significance according post-hoc Tukey's test (p<0.05) is shown with different alphabets.

Fig. 7. Schematic representation of CRL-CSN dynamics in plants. CSN holo-complex

platform protects CRLs in the absence of bound ubiquitinylation substrate (steric hindrance

prevents E2 enzyme binding (Enchev *et al*., 2012; Mosadeghi *et al*., 2016) utilizing coordinated

activities (bi-directional white arrows) of the IPK1-ITPK1 pair. Equilibrium is maintained

between CSN-mediated deneddylation cycles and $CUL1^{Nedd8}$ requirements in steady-state CRL

functions. Balanced assembly/disassembly cycles of CSN5 on the CSN holo-complex are also

linked to these regulations. Perception (via unknown mode; question mark) of a stimulus, such

as Pi-starvation, reduces CSN5 deneddylase functions, likely via more facilitation of the InsP-

kinase functions to CRL-SCF complex roles in targeted ubiquitinylation (Ub^n) and degradation

of specific substrates (such as SPX4 or Aux/IAAs). IPK1-ITPK1 pair provides necessary InsP

co-factors to these activities. Inhibition modes of TNP, CSN5i-3 and MLN4924 are indicated

at the respective processes. CSN subunits are presented as numbers; CUL1, Cullin1; RBX,

RING H2 protein1; E2, Ubiquitin cojugating enzyme E2; SA, substrate adapter; SR, substrate

receptor; N8, Nedd8; IPK1, Inositol pentakisphosphate 2-kinase1; ITPK1, Inositol 1,3,4-

trisphosphate 5/6-kinase 1.

Materials and Methods

Plant material and growth conditions. Details of *Arabidopsis thaliana* (ecotype Col-0) T-

DNA insertion lines *ipk1-1, itpk1-2, ipk2β-1, itpk4-1, csn5a-2, vih2-4, itpk1-2:ITPK1-GFP,*

and *phr1 phl1* plants have been described earlier (Kuo *et al*., 2014; Jia *et al*., 2015; Kuo *et al*.,

2018; Laha *et al*., 2020). T-DNA and respective gene-specific primers are listed in

Supplementary Table 1. Seeds were stratified for 2-days at $4^{o}C$ in dark and then sterilized using

30% bleach solution. They were then germinated either directly on soil or on ½ strength

Murashige and Skoog (MS) Pi-replete (625 μM KH_2PO_4) agar plates in growth chambers

maintained at $22^{o}C$ with 70% Relative humidity (RH) and 16hrs: 8hrs; light: dark having light

intensity 100 μmol $μm^{-2}s^{-1}$. For fractionation, Pi-starvation assays, and inhibitor treatments,

plants grown on media plates as above were transferred 7-days post-germination to liquid ½

MS medium in 12-well sterile culture plates are proceeded accordingly as indicated in

respective experiments. For immuno-precipitation assays 4-week-old soil-grown plants were

used.

Size-exclusion chromatography of plant extracts. Gel-filtration protocol was according to

Huang *et al* (2013). Briefly, 14-day-old plants ground to a fine powder with liquid nitrogen

were homogenized in Gel filtration buffer (50mM Tris-HCl pH 7.5, 150mM NaCl, 1mM

EDTA, 10% Glycerol, 10mM $MgCl_2$, 1mM DTT, 1mM PMSF and 1X Protease inhibitor

cocktail). Homogenates were centrifuged for 10 mins at $4^{o}C$ and supernatant passed through

0.2 µM syringe filters. Total protein extracts were fractionated through Superdex 10/300 GL

Column (GE- Healthcare, USA) and eluted in the fractionation buffer at 0.2mL/min flow rate.

The eluants were TCA (Trichloroacetic acid) precipitated, washed with acetone and

resuspended in 1X Laemmli loading dye and used for indicated immunoblots.

In-lysate deneddylation assay. In-lysate deneddylayion assay was according to Lin *et al*

(2020). Briefly, 14-day-old seedlings were quickly homogenized in deneddylayion buffer

(10mM Tris-HCl pH7.5, 150mM NaCl and 0.5% NP-40) in absence of protease inhibitors. The

extracts were incubated at $37^{o}C$ for indicated times, reaction stopped by adding Laemmli

loading dye to 1X, and then used for immunoblots.

Phosphate-starvation/re-supplementation assays. After germination of seeds as described

earlier, seedlings were transferred to liquid ½ MS medium in 12-well sterile culture plates with

Pi-replete conditions (625 µM KH_2PO_4) and placed in the growth chamber. After 4-days, the

plants were either maintained with Pi-replete or transferred to Pi-deplete (10 µM KH_2PO_4)

liquid media and continued for another 3 days (total 14-days post-germination) before being

used for Pi-estimation, qRT-PCR, total protein extraction, or fractionation assays. For Pi-

resupply, plants in Pi-deplete medium were transferred to Pi-replete medium for 12 hrs before

analysis.

Inhibitor treatments. Stock solutions of MLN4924 (Cayman Chemicals, USA), CSN5i-3 (Novartis Inc, Switzerland), or TNP (Sigma-Aldrich, USA) was prepared in DMSO. For treatments, 10μM final concentration of MLN4924 or CSN5i-3 were used. TNP was used at 25μM final unless otherwise indicated. For assays on Col-0, *ipk1-1*, or *itpk1-2* or for fractionations, inhibitors were treated for 24hrs. For effects on PSR, inhibitors were added during the shift to Pi-deplete medium.

Ubiquitination assays. For determining global ubiquitination levels, extracts from indicated plants prepared in Gel filtration buffer in the presence of 10μM proteasome inhibitor MG132 were incubated on ice for 2hrs. Immunoblotting was then performed with anti-ubiquitin antibodies (P4D1; SantaCruz Biotechnology, USA).

Statistical analysis. All Data presented here are representative of at least three biological and technical replicates for each. Post-hoc Tukey's test ($p < 0.05$) were applied for statistical analysis.

Additional details of cloning methodologies for generation of constructs, and transgenic lines, BiFC assays, immuno-enrichments, immunoblotting, Pi-estimations, and qRT-PCRs are described in *Supplementary Materials and Methods*.

Acknowledgements

We thank Regional Centre for Biotechnology (RCB), Faridabad, for financial support, lab infrastructure, and central instrumental facilities. S.B. expresses gratitude to Department of Biotechnology (DBT), Govt. of India for the award of Ramalingaswami re-entry fellowship and Grant (No. BT/PR23666/AGIII/103/1039/2018) that supported this research. The *phr1 phl1* double mutant seeds were from Prof. Tzyy-Jen Chiou, Academia Sinica, Taipei. A special

thanks to Dr. Eva Altmann, Novartis Institutes for BioMedical Research, Switzerland for the

CSN5i-3 gift. Also to Dr. Divya Chandran (RCB) for help with transcriptomic data analysis of

csn mutants. Y.W. thanks DBT for Young Investigator award. M.K. and K.D.I thank UGC and

DBT for fellowships, respectively. D.L. acknowledges the Indian Institute of Science for start-

up funds. All authors thanks Dr. Nripendra Singh and his team at Advanced Technology

Platform Centre (ATPC), Faridabad, India for constant support with sample fractionations.

Fig. 1. IPK1 and ITPK1 interact with each other and with selective CSN or CRLs components *in planta*. *(A)* Immuno-enrichments of IPK1 *(top panel)* or ITPK1 *(bottom panel)* from corresponding complemented lines (*ipk1-1:Myc-IPK1* or *itpk1-2:ITPK1-GFP*, respectively) was probed with anti-CSN1/2/4/5, anti-CUL1, or anti-ASK1 antibodies. IgG-agarose enrichments were used as negative controls. *(B)* Bimolecular Fluorescence Complementation (BiFC) assays showing *in planta* IPK1 or ITPK1 interaction with CSN1/2 and with each other. Indicated binary vectors combinations were co-expressed via Agrobacterium in *Nicotiana benthamiana* leaves and fluorescence observed under a confocal microscope with YFP filter. Scale bar=50 μm. *(C)* Co-immunoprecipitation of transiently expressed GFP-ITPK1 with Myc-IPK1 from *N. benthamiana* leaves. Anti-GFP pulldown was probed with anti-GFP or anti-Myc antibodies.

Fig. 2. CSN5 deneddylase deficiencies with increased holo-complex retention in *ipk1-1* or
itpk1-2 **cause enhanced CUL1^{Nedd8} levels.** *(A)* IPK1 or ITPK1 does not co-elute with the CSN
holo-complex. *(B)* Enhanced CSN5 engagements on the CSN holo-complex in *ipk1-1* and
itpk1-2. Extracts from marked plants fractionated by size-exclusion chromatography were
immuno-blotted with indicated antibodies. Fraction numbers and elution positions of molecular
weight standards (in kDa) are shown. Fractions corresponding to CSN holo-complex and CSN
subunit monomeric elutions are marked. *(C)* Immuno-enrichment of CUL1 detects
stoichiometrically similar amounts of CSN1 and CSN2 in *ipk1-1* or *itpk1-2* to Col-0. Pulldowns
with either IgG or anti-CUL1 antibodies were probed with anti-CUL1, anti-CSN1 or anti-
CSN2 antibodies. *(D)* Increased CUL1^{Nedd8} levels (*top panel*) and enhanced global
polyubiquitin-conjugates (*bottom panel*) in *ipk1-1* or *itpk1-2*. Total protein extracts were
probed with anti-CUL1, anti-actin, or anti-Ubiquitin antibodies as indicated. *(E)* CSN5-
deneddylation deficiencies and lower CUL1 half-life in *ipk1-1* or *itpk1-2*. In-lysate CUL1

deneddylation assay (Lin *et al.*, 2020) at the indicated time points (minutes post-incubation;
mpi) probed with anti-CUL1 antibodies. Migration position of molecular weight standards (in
kDa) and neddylated (CUL1^{Nedd8}) or unneddylated (CUL1) Cullin1 are shown. Anti-actin or
Ponceau S staining demonstrates loadings controls. Migration position of molecular weight
standards (in kDa) and neddylated (CUL1^{Nedd8}) or unneddylated (CUL1) cullin1 are shown.
Anti-actin or Ponceau S staining demonstrate loadings controls.

Fig. 3. Enhanced CUL1^{Nedd8} levels associated with PSR is essential for PSI-gene inductions. MLN4924 suppresses *(A)* elevated PSI-gene expressions, and *(B)* higher endogenous Pi-levels in *ipk1-1* or *itpk1-2*. *(C)* Induction of *PHT1;3* or *IPS1* expressions in Col-0 upon Pi-starvation is inhibited by MLN4924. *(D)* Increments in CUL1^{Nedd8} levels during Pi-deprivation is PHR1/PHL1-independent. Plants grown in Pi-replete medium or transferred to Pi-deplete medium were treated with or without MLN4924. Gene expression changes were relative to *MON1* transcripts and reported as fold-change to Col-0 levels. Data is mean $\pm$ SD (n=3 for qPCRs; n=6-7 for Pi-estimations). Statistical significance indicated with different alphabets is according to post-hoc Tukey's test (p<0.05). For *PHT1;3* or *SPX1* expression data, statistical comparisons were to respective DMSO-treated Col-0. For immunoblots, anti-CUL1 antibodies were used. Ponceau S staining shows loadings controls.

Fig. 4. Pi-starvation reduces endogenous CSN5 deneddylase activity and is restored upon phosphate resupply. *(A)* Extracts at indicated time points (minutes post-incubation; mpi) from in-lysate CUL1 deneddylation assay on Col-0 from Pi-replete or Pi-deplete media were probed with anti-CUL1 antibodies. *(B)* Restoration of enhanced CUL1^{Nedd8} levels occur upon Pi-resupply post-starvation. *(C)* Increased retention of CSN5 on the CSN holo-complex under Pi-deplete exposure is reversed with Pi-resupply. *(D)* Global levels of polyubiquitin-conjugates remain unchanged during PSR. For these assays, plants were subjected to Pi-starvation for 3 days and phosphate re-supplementation was done for 12hrs. Comparable loadings are shown by Ponceau S staining. Elution positions of molecular weight standards (in kDa) are indicated.

Fig. 5. Inhibition of CSN5 functions activates PSR under Pi-replete but confers non-responsiveness under Pi-deplete growth. *csn5a-2* plants have *(A)* elevated endogenous Pi levels, *(B)* increased *PHT1;3* or *SPX1* transcript levels under Pi-replete growth but suppressed expression upon Pi-starvation. *(C)* CSN5i-3 inhibits Arabidopsis CUL1 deneddylation, *(D)* elevates endogenous Pi-levels, and *(E)* suppresses *PHT1;3* induction during phosphate-deprivation. Immunoblots were probed with anti-CUL1 antibodies. Ponceau S staining shows loading controls. Inhibitor treatments were done for 24 hrs and doses used are indicated. Expression levels of *MON1* transcripts was used as an internal reference for qPCRs and reported as fold-change relative to DMSO-treated Col-0 under Pi-replete growth. Error bars are mean $\pm$ SD (n=3 for qPCRs; n=6-7 for Pi-estimations). Different alphabets denote statistical significance determined by post-hoc Tukey's test (p<0.05). *PHT1;3* and *SPX1* data are compared to respective Pi-replete Col-0.

Fig. 6. CSN5i-3 or TNP treatments on Col-0 mirrors *ipk1-1* or *itpk1-2* properties. *(A)* CSN5 partition shifts to CSN holo-complex pools upon CSN5i-3 or TNP treatment. TNP *(B)* enhances pools of CUL1^Nedd8, *(C)* increases endogenous Pi-content, *(D)* upregulates *IPS1* or *PHT1;3* expressions under Pi-replete, but suppresses them under Pi-deplete growth, and *(E)* decreases endogenous deneddylation rates with increased CUL1 instability. Immunoblots were probed with anti-CUL1 or anti-CSN5 antibodies as indicated. Ponceau S staining shows loading controls. Expression levels of *MON1* transcripts was used as internal reference for qPCRs. Error bars are mean $\pm$ SD (n=3 for qPCRs; n=6-7 for Pi-estimations). Statistical significance according post-hoc Tukey's test (p<0.05) is shown with different alphabets.

Fig. 7. Schematic representation of CRL-CSN dynamics in plants. CSN holo-complex platform protects CRLs in the absence of bound ubiquitinylation substrate (steric hindrance prevents E2 enzyme binding (Enchev *et al*., 2012; Mosadeghi *et al*., 2016) utilizing coordinated activities (bi-directional white arrows) of the IPK1-ITPK1 pair. Equilibrium is maintained between CSN-mediated deneddylation cycles and $CUL1^{Nedd8}$ requirements in steady-state CRL functions. Balanced assembly/disassembly cycles of CSN5 on the CSN holo-complex are also linked to these regulations. Perception (via unknown mode; question mark) of a stimulus, such as Pi-starvation, reduces CSN5 deneddylase functions, likely via more facilitation of the InsP-kinase functions to CRL-SCF complex roles in targeted ubiquitinylation (Ub^n) and degradation of specific substrates (such as SPX4 or Aux/IAAs). IPK1-ITPK1 pair provides necessary InsP co-factors to these activities. Inhibition modes of TNP, CSN5i-3 and MLN4924 are indicated at the respective processes. CSN subunits are presented as numbers; CUL1, Cullin1; RBX, RING H2 protein1; E2, Ubiquitin cojugating enzyme E2; SA, substrate adapter; SR, substrate receptor; N8, Nedd8; IPK1, Inositol pentakisphosphate 2-kinase1; ITPK1, Inositol 1,3,4-trisphosphate 5/6-kinase 1.