Trait Anxiety Effects on Late Phase Threatening Speech Processing: Evidence from Evidence from Electroencephalography

Simon Busch-Moreno[a], Jyrki Tuomainen[a], David Vinson[a]

[a] Division of Psychology and Language Sciences, University College London, London, United Kingdom

Abstract:

The effects of threatening stimuli, including threatening language, on trait anxiety have been widely studied. However, whether anxiety levels have a direct effect on language processing has not been so consistently explored. The present study focuses on event-related potential (ERP) patterns resulting from electroencephalographic (EEG) measurement of participants' (n = 36) brain activity while they perform a dichotic listening task. Participants' anxiety level was measured via a behavioural inhibition system scale (BIS). Later, participants listened to dichotically paired sentences, one neutral and the other threatening, and indicated at which ear they heard the threatening stimulus. Threatening sentences expressed threat semantically-only, prosodically-only, or both combined (congruent threat). ERPs showed a late positivity, interpreted as a late positive complex (LPC). Results from Bayesian hierarchical models provided strong support for an association between LPC and BIS score. This was interpreted as an effect of trait anxiety on deliberation processes. We posit that verbal repetitive thinking, as associated with anxious rumination and worry, can be the mechanism disrupting late phase deliberation processes. Instantiated by inner speech, verbal repetitive thinking might be the vehicle of anxiety-related reappraisal and/or rehearsal, which would make processing more effortful and slower.

Keywords: prosody, semantics, anxiety, emotion, cognition, electrophysiology

Graphical Abstract:

Introduction

Both anxiety and language processing are characterized by well-defined lateralization patterns. Anxiety has been widely associated to a dual processing pattern: while attention-related arousal and fear-related responses tend to involve greater right hemisphere (RH) processing, evaluation-related apprehension and inhibition-related responses tend to involve greater left hemisphere (LH) processing (Heller et al., 1997; Nietschke et al., 2000; Spielberg et al, 2013). According to current models of speech processing (i.e. dual stream model), speech comprehension recruits both LH and RH (Kemmerer, 2015). While more RH involvement for the slow rate suprasegmental processing of prosody and/or affect/attitude recognition is required, more LH involvement is required for fast rate segmental processing and/or lexical categorization (Belin et al., 2004; Liebenthal et al., 2010; Poeppel et al., 2008; Zatorre et al., 2002). These observations speak not only of general hemispheric activity, but also of very specific activity patterns and brain anatomical structures that, in many cases, are shared by language and anxiety processing. This opens the question of whether these processes simply co-occur, showing superficial similarities, or they actually interact with each other.

However, anatomical differences are not necessarily absolute or static, but they may depend on the time-course of language processing, as evidenced by emotional prosody and semantics lateralization patterns varying across time (Kotz and Paulmann, 2011). Similarly, over-attention to and over-engagement with threat may be modulated by anxiety at early and late processing phases respectively (Bar-Haim et al., 2007). Here, over-attention implies the facilitated attention to threatening stimuli, and over-engagement implies difficulty in disengaging from threatening stimuli (Cisler and Koster, 2010). Therefore, understanding these anxiety-related attention processes as multistage processes might be crucial. A two-stage model proposes that trait anxious personality is characterized by an over-active behavioural inhibition system (BIS), when attempting to resolve approach avoidance conflicts. This would develop in two stages: 1) valuation of stimuli immediately after stimulus input, and 2) motivation for behaviour as approach or withdrawal (Corr and McNaughton, 2012). However, this model does not give an account of processes that might occur between valuation and motivation, as proposed in a multistage model of anxiety (Bar-Haim et al, 2007). In this model, anxiety develops through four processing stages: 1) pre-attentive threat

evaluation, 2) resource allocation, 3) guided threat evaluation, 4) goal engagement system. Here, a valuation input can respond to pre-attentive stages, which can account for fear responses inducing rapid withdrawal (output), or fight-flight-freeze (FFFS) system responses (Corr and McNaughton, 2012); or can respond to resource allocation (output), which can lead to behavioural inhibition (or BIS engagement) in a threat evaluation stage. Likewise, a goal engagement system can account for a motivation output that occurs after BIS activation, for either approaching or withdrawing.

Interestingly, a multistep model has also been proposed for characterizing emotional language processing (Kotz and Paulmann, 2011). This model proposes three main stages: 1) early stage perceptual processing, 2) mid stage recognition processing, 3) late stage evaluation processing. An important aspect of this model is that it takes into account hemispheric differences of language and emotional processing. Early stages would respond to the processing of acoustic properties, where greater right hemisphere (RH) engagement would be associated to prosodic slower rate spectral information and more left hemisphere engagement (LH) would be associated to phonological (and by extension semantic) faster rate temporal information (Poeppel et al., 2008; Zatorre et al., 2002). Mid stages might involve the emotional recognition of stimuli, implying greater involvement of RH or LH depending on stimulus type and/or conveyed emotion (Schirmer and Kotz, 2006). Late stages would be associated to informational integration and evaluation of emotional stimuli (Kotz and Paulmann, 2011). If indeed the similarities between the multistage model of anxiety and the multistep model of language are more than coincidental, the latter should also have a fourth stage, which can account for motivation/goal-orientation processes associated to behavioural outputs in parallel to the multistage model of anxiety.

Nevertheless, to develop such a model, more evidence is necessary about how informational properties of language interact with intrinsic affect. Given that lexical items and prosody can convey emotional meaning at the same time (Nygaard et al., 2009; Schirmer and Kotz, 2003), the simultaneous recognition and integration of diverse informational properties is required, including the processing of segmental and spectral information, tonal and temporal information, and identification and categorization processes. As mentioned before, these aspects of language processing show different hemispheric asymmetries, which might be linked to anxiety related lateralisation in terms of how this information is recognized, integrated, and evaluated.

Thus, a model of anxious lateralization (Heller et al., 1997) could be extended into a multistage model of anxiety (Bar-Haim et al., 2007), given early arousal-related over-attention to threat and later (and sustained) worry-associated over-engagement with threat differences (Spielberg et al., 2013).

Recent electroencephalography (EEG) evidence indicates that a hyperactive BIS, signalled by higher scores in BIS psychometric scales, is characterised by a right frontal hemispheric pattern (Gable et al., 2018; Neal and Gable, 2017). However, some key theoretical notions cannot be overlooked. On the theoretical side, the proposed anatomic-physiological basis of BIS/anxiety is the amygdala-hippocampal-septal route (Gray and McNaughton, 2000), which can modulate diverse cortical patterns depending on environmental conditions. Indeed, it has been proposed that the BIS limbic system might modulate higher processes such as worry and rumination through language systems (in particular left lateralized production/rehearsal), which could induce additional inhibitory control (Gray and McNaughton, 2000). Hence, while RH may induce initial inhibitory control (McNaughton et al., 2013), there might be later LH influence involving evaluation or deliberation processes. Previous EEG studies have observed left or bilateral frontal alpha activity associated with anxious apprehension as measured by worry (Heller et al., 1997; Nitschke et al., 1999), bilateral alpha for rumination-correlated BIS (Keune et al., 2012), and no hemisphericity patterns of delta or theta waves associated with BIS (De Pascalis et al., 2013). Indeed, recent models of anxiety (stated very generally), propose activation of dorsolateral pre-frontal cortex (dLPFC) inducing medial prefrontal cortex (mPFC) activity, together with hippocampus and insula, that will induce control over amygdala; where structures such as dLPFC show different lateralisation patterns (Robinson et al., 2019).

Given this, manipulating hemispheric input could reveal possible effects of anxiety on threatening speech processing at different processing phases. In other words, stimuli that are processed first by LH or RH might be affected differently depending on: 1) their language informational properties (i.e. semantics or prosody) and/or 2) participants' intrinsic lateralization differences when processing threatening stimuli (i.e. anxiety). Considering this, dichotic listening (DL) stands out as an ideal behavioural test to observe how anxiety-related hemispheric asymmetries might influence emotional language processing hemispheric asymmetries. This relates to the frequently observed phenomenon of a right ear advantage (REA) for non-

prosodic language stimuli (Hugdahl, 2011); this implies participants answering faster and/or more accurately to stimuli presented at their right ear when compared to stimuli presented at their left ear. On the other hand, prosody, in particular emotional prosody, has been observed to present either a left ear advantage (LEA) or a diminished REA (Godfrey and Grimshaw, 2015; Grimshaw et al., 2003). Few DL experiments have researched the effects of anxiety on emotional speech processing (Gadea et al., 2011). Instead, they either use speech/prosody as an emotion-eliciting stimulus or use DL mainly as an attentional manipulation technique (Bruder et al., 2004; Leshem, 2018; Peschard et al., 2016; Sander et al., 2005; but see Busch-Moreno et al., 2020).

In our previous research (Busch-Moreno et al., 2020), we investigated possible laterality differences via a two DL experiments. Participants had to complete two tasks per experiment. In one task (direct-threat) they answered by pressing a keyboard's left or right arrows indicating at which ear (left or right) they heard a threatening sentence (paired with a neutral sentence at the contralateral ear). In the second task, they answered at which ear they heard the neutral sentence of the dichotic pair (indirect-threat). In both tasks, they heard three possible types of sentence: Prosody (threatening prosody but neutral content), Semantic (threatening content but neutral prosody), and Congruent (both prosodic and semantic threat). In the first experiment, participants answered only after sentences finished playing, while in the second they had to answer before sentences' end. Both experiments showed a similar pattern of results, where reaction times (RTs) increased as a function of worry level (trait anxiety) and accuracy did not relevantly change, irrespective of ear, sentence-type or task. We interpreted this as a more general effect of anxiety on responses to threatening speech, indicating that longer stimuli may be sufficient for generating over-engagement with threat (see also Cisler and Koster, 2010). In particular we propose that verbal repetitive thinking could affect later processing stages (over 500ms) and create particularly high increase in RT when anxious participants answer to threatening speech.

However, these behavioural tasks cannot provide evidence of effects on specific time-windows, as required by multistep models (i.e. Kotz and Paulmann, 2011). EEG measurements, via the event-related potential (ERP) technique, could help to elucidate when anxiety, of the apprehensive type (i.e. Heller et al., 1997), has stronger effects. In

other words, whether early and/or late ERPs differ given ear, sentence-type and task; and how this may relate to over-attention or over-engagement. To our knowledge, no EEG experiment has integrated trait anxiety measures with DL to investigate the effects of anxiety on threatening semantics and prosody. Presently, we aim to implement our previous study's first experiment (response after sentence's offset), including both direct- and indirect-threat tasks. This would help to make more consistent replication of our previous research and to see whether direct- or indirect-threat can differently affect the ERP patterns at early (over-attention) or later (over-engagement) time-windows. In the present behavioural replication, we included some changes. We used a newly recorded set sentences to improve audio quality. Also, we used a BIS scale from the Reinforcement Sensitivity Theory Personality Questionnaire (RST-PQ) (Corr and Cooper, 2016) to measure BIS levels as a proxy of trait anxiety. This psychometric measure, well grounded in physiological research, accommodates better to our present theoretical framework and experimental aims. Finally, via EEG measurement, we can provide fine-grained temporal information via the ERP technique, crucial for testing our theoretical model.

Given this setup and the previously discussed theory and research, we expect that higher levels of anxiety (higher BIS scores) will induce early over-attention to threat but mid-late over-engagement with threat. The first should be associated with anxiety induced arousal, which has been observed to be right lateralized (Heller et al., 1997; Neo et al., 2011). The second should be associated to a left lateralized or bilateral pattern (Nitschke et al, 1999), which might be due to mid-stage LH thought-induced (e.g. worry or rumination) evaluation and inhibition (Spielberg et al., 2013), but a later RH involvement associated to arousal induced by sustained state anxiety (McNaughton and Gray, 2000). We established four a priori time windows where differences may be present: 1) pre-attentive (50-150ms), 2) attentive (150-250ms), 3) evaluative (250-500ms), 4) orientative (500-750ms). These time-windows are selected to span over periods of activity for which theoretically relevant ERPs may be present around 100ms (sensorial perceptual stage), 200ms (recognition stage), ~300-400ms (evaluation stage), and ~600ms (evaluation/deliberation stage) (Kotz and Paulmann, 2011). We propose that most noticeable effects of anxiety should be present at the deliberation/orientation phase (~600ms), which due to effects of repetitive thinking (Busch-Moreno et al., 2020) should make language processing more effortful.

Following previous predictions, we specify the following hypotheses: 1) Early (~100ms) and Mid-early (~200ms) effects of anxiety on speech processing (Pell et al., 2015), where over-attention to threat will show stronger RH involvement for prosody, facilitating detection, noting that this should imply contralateral left ear (RH) greater activity. 2) Mid-late stage (~400ms) anxiety effects on LH due to over-engagement with threat (Spielberg et al., 2013); this should particularly affect semantic stimuli by slowing down their processing, noting contralateral right ear (LH) increased activity. 3) A late stage (~600ms) effect of anxiety on goal orientation (Bar-Haim, 2007), where both LH and RH (more strongly) should be involved (no ear differences). This could be the results of a worry-arousal loop (McNaughton and Gray, 2000) due to continued exposure to threatening stimuli. In other words, verbal repetitive thinking could take prevalence at this stage due to over-engagement with threat (Busch-Moreno et al., 2020). No specific ERPs are predicted, as present research has no direct precedents in the literature. Hence, it is not clear whether observed ERPs and their amplitude differences by condition will be coincidental with those observed in previous research, which regards DL, emotional speech, or anxiety, but not all three combined.

Methods

Participants

Participants (n = 36, mean age = 28.6, age range = [19, 54], 19 females) were recruited using Sona Systems (sona-systems.com). Only participants reporting being right-handed, having English as first language, without hearing problems and with no history of neurological/psychiatric disorders were recruited. Participants were remunerated at a £7.5/hour rate. All participants were informed about their rights, including those regarding their data (protected under GDPR protocols), and gave their informed consent before participating. They received a debriefing statement at the end of the experiment. Present experiment was carried out under approval from the UCL Research Ethics Committee in accord with the Declaration of Helsinki.

Materials

Four types of sentences were recorded: Prosody (neutral-semantics and threatening-prosody), Semantic (threatening-semantics and neutral-prosody),

Congruent (threatening-semantics and threatening-prosody), and Neutral (neutral-semantics and neutral-prosody). Sentences were extracted from movie subtitles by matching them with a list of normed threatening words from the extended Affective Norms for English Words (ANEW) (Warriner et al., 2013). For the present study, any word over 5 points in the arousal scale, and below 5 points in the valence and dominance scales was considered semantically threatening. Every word with less than 5 arousal points and between 4 and 6 valence points was considered semantically neutral. Words' frequencies were extracted from SUBTLEX-UK (van Heuven et al., 2014), only sentences containing words with Zipf log frequencies over 3 were included.

Sentences were recorded in an acoustically isolated chamber with a RODE NT1-A1 microphone. The speaker was instructed to speak in what he considered his own threatening/angry or neutral voice for recording Prosody/Congruent and Semantic/Neutral sentences respectively. Table 1 summarises sentences word length and duration. Figure 1 shows oscillograms and spectrograms of four sentences used as stimuli. All threatening sentences were paired with a different Neutral sentence of similar length, minor increases of sentences' silences (up to 40ms) were applied to make paired sentences' durations match perfectly. This resulted in 54 recorded sentences per threatening category (Prosody, Semantic, Congruent), each one paired with a different sentence from the Neutral category. This provided 162 pairings of two sentences (one threatening and one neutral). We created two versions of each pair: one with threat at the right ear, and one with threat at the left ear: 324 distinct stimuli in total.

Table 1. Average number of words, duration and reaction time per stimulus type

Type	Words Threat	Words Neutral	Stimulus Duration	Average RT
Congruent	4.63	5.63	1838.63	522.37
Prosody	4.24	4.93	1558.25	601.20
Semantic	4.43	4.89	1543.53	593.40

Figure 1. Examples of sentences with different semantic content and prosody. Top of each image: oscillogram. Bottom of each image: spectrograms. Top left: neutral prosody and neutral semantics (Neutral). Top right: threatening prosody and threatening semantics (Congruent). Bottom left: neutral prosody and threatening semantics (Semantic). Bottom right: threatening prosody and neutral semantics (Prosody). Green dots indicate fundamental frequency (F0) contours.

In addition, we assessed the prosodic properties of threatening sentences as compared to neutral sentences. Threatening prosody has been observed to align with hot anger (Banse and Scherer, 1996; Hammerschmidt and Jürgens, 2007) and can be defined mainly by pitch and roughness (voice quality). Thus, we focused only on Median Pitch (median each sentence's fundamental frequency) and Hammarberg Index (difference between 0Hz to 2000Hz and 2000Hz to 5000Hz frequency ranges), as these acoustic measures can capture pitch (fundamental frequency) and voice quality variation respectively and remain in the spectral domain. Measures were extracted from sentences with Python's Parselmouth interface to Praat (Jadoul et al., 2018). Acoustic measures were compared by using BEST (Kruschke, 2013), but using $SD \sim Exponential\ (\lambda = 2)$ instead of uniform distributions. All models were sampled with MCM and used 1000 tuning samples and 1000 samples, convergence and sampling were excellent for both models ($\hat{R} \cong 1$, ESS > 400, BFMI > 0.8). As very clear

in Table 2, sentences containing threatening prosody have higher pitch (median pitch) and lower voice quality (Hammarberg index, i.e. more roughness) than sentences without threatening prosody. For both measures, Prosody and Congruent distributions greatly overlap and Semantic and Neutral sentences almost completely overlap.

Table 2. Estimates from BEST Model

Arousal					Median Pitch				
Category	Mean	SD	HDI 5%	HDI 95%	Category	Mean	SD	HDI 5%	HDI 95%
Neutral	3.55	0.08	3.41	3.68	Neutral	101.38	0.7	100.29	102.62
Congruent	6.13	0.09	5.97	6.28	Congruent	167.25	2.55	163.06	171.35
Prosody	3.8	0.07	3.68	3.92	Prosody	175.85	2.16	172.17	179.23
Semantic	6.26	0.09	6.1	6.4	Semantic	102.01	0.7	100.89	103.15
Valence					Hammarberg Index				
Category	Mean	SD	HDI 5%	HDI 95%	Category	Mean	SD	HDI 5%	HDI 95%
Neutral	5.43	0.05	5.34	5.5	Neutral	21.09	0.52	20.32	22.04
Congruent	2.88	0.13	2.68	3.11	Congruent	7.93	0.54	6.96	8.79
Prosody	5.5	0.05	5.42	5.58	Prosody	7.85	0.41	7.25	8.61
Semantic	2.75	0.12	2.55	2.94	Semantic	21.2	0.57	20.28	22.2

Note: All estimates are taken from comparisons between means resulting posterior distributions.

Finally, we collected sentences threat ratings post-experimentally, for further confirming effects of both acoustic measures and ANEW norms. Ratings were given by 10 participants for 54 Prosody and 54 Neutral sentences (total number of Prosody and Neutral) in a 9-points Likert scale (0 = no threatening at all, 8 = very threatening). The same procedure was followed for the 54 Semantic sentences together with the same 54 Neutral sentences in a second rating session by 10 participants. Ratings were analysed by using a hierarchical Ordered-logistic model, with subjects as a varying intercept, eight scale cutpoints and normal priors (equivalent to L2 regularisation) for non-varying slopes for Pitch, Roughness, Arousal and Valence. The full model and plots can be found in our OSF repository (https://osf.io/n5b6h/). Figure 2 summarises posterior distributions as probabilities derived from the ordered-logistic model. For the Prosody task, these indicate that when Median Pitch (MP) is high and Hammarberg Index (HI) low, the probability of rating sentences over 6 threat scores increases over 40%. For the Semantic task, when Arousal is high and Valence is low the probability of rating sentences over 7 threat scores increases over 60%. In both tasks when MP/Arousal are low and HI/Valence are high the probability of giving 1 or less threat scores increases by around 60%. The models also indicate that Arousal/Valence had

no relevant effects on Prosody and MP/HI had no relevant effects on Semantic. See supplementary materials for tables with more details (Supplement 1, supplementary tables 1 and 2). All relevant materials used and their measures can also be found in supplementary materials (Supplement 1, tables 3 and 4).

Figure 2. Probability of threat ratings. Left panel: probability of rating a Prosody sentence as threatening in a 0-8 Likert scale. Orange solid line: probability increase when a sentence has the Max Median Pitch (MP) and Min Hammarberg Index (HI). Violet dashed line: probability decrease when a sentence has the Min MP and Max HI. Left panel: probability of rating a Semantic sentence as threatening in a 0-8 Likert scale. Red solid line: probability increase when a sentence has the Max Arousal and Min Valence. Blue dashed line: probability decrease when a sentence has the Min Arousal and Max Valence. Faded lines are random samples from the posterior indicating uncertainty.

Procedure

Participants were introduced to the recording room (electrically and sound isolated chamber), signed consent, and sat at 1m distance from a 20" screen used to display all tasks via PsychoPy2 (Peirce et al., 2019). Next, participants provided their demographic information (age and sex) and completed the BIS scale (m = 23, SD = 13) and FFFS scale (m = 29, SD = 9.5) (for more details see Corr and Cooper, 2016). Participants' head dimensions were measured and EEG caps of appropriate size were placed and centered, conductivity gel was placed and the Biosemi 64 Ag/Cl electrode system (biosemi.com) was connected. Impedance levels were kept below 20Ω and electrodes were checked to be working properly. While EEG recording, participants completed a direct-threat task where they answered at which ear (left or right) they heard a threatening sentence (always paired with a neutral sentence) by pressing the left or right arrows on a keyboard (as fast and accurately as possible) only after the stimulus finished playing (sentence offset) and a target symbol appeared on screen.

Trials (324 per task) started at sentence onset and were separated by a 1.5s inter-stimulus interval (ISI). In the indirect-threat task they answered to the neutral sentence of the pair in the same manner as in direct-threat. Order of task was counterbalanced across participants. Participants heard sentences twice such that a threatening sentence was presented once at each ear. Participants were requested to swap response hand every other block. Each block consisted of 18 sentences (trials), after which participants were allowed to rest. Starting hand and starting ear were also counterbalanced across participants. Due to a coding issue, in the indirect-threat condition for half of the participants (list B: starting with right ear only) stimuli were not exactly balanced for ear presentation. This issue was not noticed by participants during debriefing, it did not affect more than 4 sentences per condition per subject, and did not systematically affect the results. See supplementary materials for follow-up analyses to address this issue (Supplement 3).

EEG Data Processing

EEG recordings were pre-processed using Python's MNE package (Gramfort et al., 2014). A completely automated pre-processing pipeline was implemented, based on Jas et al.'s (2018) method. This consisted in the following steps:

1) Importing data, previously down-sampled to 512Hz using Biosemi decimation tools, checking that all events were correctly placed and fixing if necessary.

2) Preparing data for independent component analysis (ICA): set data to average reference (Dien, 1998; Lei and Liao, 2017), low-pass filter at 40hz to avoid aliasing artifacts, down sample to 256hz, high pass filter at 1hz for better artifact detection (both FIR filters, -6dB cut-off), and automatic rejection of noisy segments based on the Autoreject MNE package (Jas et al., 2017; Winkler et al., 2015).

3) Compute ICA components by using python Picard package (Ablin et al., 2017) for EOG artifact detection. ICA components that are highly correlated with noise (> 0.35, using MNE component correlation method) at Fpz channel were marked for later rejection.

5) Reload raw data to apply average reference to raw data (Dien, 1998; Lei & Liao, 2017); to use more relaxed filters: first a high pass (0.1Hz, FIR, -6dB cut-off) and later a low pass (100Hz, FIR, -6dB cut-off) filter (Luck, 2014; VanRullen, 2011; Widmann et

al., 2015); and to exclude EOG artifacts detected in previous step (1 to 2 components were removed from each participant recording.).

6) Create epochs from 0ms (sentence onset) to 1000ms using a 100ms pre-onset period (10% of epoch). This 1s epoch allows sufficient time for observing the proposed 4 time-windows and safely avoids possible contamination from later behavioural responses (analysis of later response preparation and/or response is out of present scope and would require dynamic epoching respect to RTs).

7) Applying automatic detection, repairing and rejection of noisy epochs by using Autoreject's Bayesian optimization method (Jas et al., 2017). 4 to 8 channels were interpolated by participant. Between 0 and 206 trials were dropped per participant (32.75 on average), and between 29 and 54 trials were retained per condition. 54 being the total number of trials per condition (but see minor technical issue at the end of Procedure section, see also Supplement 3). So, trials are generally balanced.

8) Applying baseline correction (baseline subtraction) (Luck, 2014; Tanner et al., 015) by using the mean of the baseline period established in the previous step.

9) Extracting trial by trial mean amplitudes at 4 a priori defined time windows: 50-150ms, 150-250ms, 250-500ms, 500-750ms; these time windows cover the time-windows proposed by the multistep model (Kotz and Paulmann, 2011), plus our proposed fourth time-window. ERP and scalp plots of processed data were produced using the MNE package.

Data Analysis

Figure 3 shows the model used for EEG data, the diagram is based on Kruschke's (2015) and Martin's (2018) model specification and presentation, and their guidelines on robust regression. The model was sampled using Markov Chain Monte Carlo (MCMC) No U-turn Sampling (NUTS) as provided by PyMC3 (Salvatier et al., 2016). All models were sampled with two chains of 2000 tuning steps and 2000 samples, and initialised using automatic differentiation variational inference (ADVI) as provided by PyMC3. Plots of results were produced using Arviz (Kumar et al., 2019) and Matplotlib (Hunter, 2007). Results were assessed using a region of practical equivalence (ROPE) method (Kruschke, 2015; Martin, 2018), where high posterior density (HDI) intervals were considered as presenting a considerable difference when far away from ROPEs defined as 1SD to 2SDs around zero.

The core idea of the model is to un-pool data from the individual level of subjects and items, so we can focus on group level slopes at each single interaction point, trial-by-trial. Given this, pooling at the electrode (Channel in Figure 3) level through a normal non-varying prior helped us to define an offset for the other reparametrized priors (e.g. McElreath, 2020), which do not contain individual location parameters. In this way we could obtain intercepts from each electrode and improve sampling and convergence in a substantial manner. To be consistent with previous research, we did not include FFFS rating in the analyses, as our main focus is on trait anxiety and not on trait fear. Fear measures (FFFS) were collected as they could be required for comparison in future analyses. See our OSF repository for behavioural models (https://osf.io/n5b6h/).

In short, the model (Figure 3) comprises one non-varying intercept Channel (64 electrodes matrix); two varying intercept Subjects (36 participants) and Sentences (162 stimuli); and one varying slope (matrix: 2 ears by 3 sentence-types by 64 electrodes), multiplied by the BIS (x in Figure 3) continuous variable. All varying parameters are re-parametrised by using a scale (half-normal) by the main parameter (normal distribution). The likelihood is a Student-t distribution (see parameters in Figure 3), ideal for mean amplitude data, which tends to cluster around zero but leave long negative and positive tails; in this way possible outliers are included into the model instead of being discarded (Kruschke, 2015; Martin, 2018). The present model will be able to tell whether BIS varies differently given ear, sentence-type and electrodes parameters. In this way, we will obtain a posterior distribution with a respective HDI indicating the slope of BIS, namely how much amplitude increase/decreases as a function of BIS per electrode at each ear (left or right) per type (Congruent, Prosody, Semantic). This model is implemented separately at each time-window, thus allowing to observe whether BIS has an effect at early (50-150ms), mid-early (150-250ms), mid-late (250-500ms) or late (500-750ms) time-windows according to present hypotheses. This implies that when there is an important increase/decrease in amplitude at a given condition (e.g. Congruent at Left Ear at Cz Channel), HDIs of BIS slopes should fall apart from zero and ROPE and the specific amplitude change at a given electrode could predicted simply by obtaining the product of any BIS score and the estimated posterior distribution of the slope.

Figure 3. The diagram shows the prior distributions used for robust regression on amplitude. Arrows indicate the relationship between a parameter and priors or hyperpriors, where tilde (~) indicates a stochastic relationship and equal (=) indicates a deterministic relationship. Observations (Obs.) in the likelihood distribution are equivalent to mean amplitude at each single trial. Channels: 64 electrodes. Ear: 2 ears (left and right). Type: 3 sentence-type (Congruent, Prosody, Semantic).

Results

Behavioural Results

All models sampled well, showing good convergence ($\hat{R} \cong 1$, ESS > 200, BFMIs > 0.6); all summaries and plots can be found in present study's Open Science Framework (OSF) repository (https://osf.io/n5b6h/). Accuracy results largely replicate results reported in our previous research (Busch-Moreno et al., 2020), showing uncertain accuracy increases which are not consistent across condition (direct- or indirect-threat), type (prosody or semantic threat) or ear; generally suggesting no accuracy advantages or disadvantages. Table 4 and Table 5 summarise accuracy results from direct- and indirect-threat conditions. Note that these results consider BIS slopes (in log-odds) across all conditions, and all posterior distributions overlap zero, as clearly indicated by HDIs (see OSF repository for more details).

Table 4. Direct-threat Accuracy Estimates

Type	Ear	BIS Score	Posterior Mean	Posterior SD	HDI 5%	HDI 95%
Semantic	Left	1	-0.01	0.02	-0.04	0.03
Prosody	Left	1	-0.01	0.02	-0.05	0.02
Semantic	Right	1	0.00	0.02	-0.03	0.03
Prosody	Right	1	0.01	0.02	-0.02	0.04
Semantic	Left	55	-0.42	1.08	-2.03	1.45
Prosody	Left	55	-0.74	1.06	-2.48	0.94
Semantic	Right	55	-0.15	0.94	-1.61	1.39
Prosody	Right	55	0.63	0.92	-0.88	2.07

Note: *All estimates are in log-odds. Estimates come from slope's posteriors.*

Table 5. Indirect-threat Accuracy Estimates

Type	Ear	BIS Score	Posterior Mean	Posterior SD	HDI 5%	HDI 95%
Semantic	Left	1	-0.008	0.017	-0.037	0.018
Prosody	Left	1	0.026	0.017	0.000	0.056
Semantic	Right	1	-0.006	0.014	-0.030	0.015
Prosody	Right	1	0.014	0.014	-0.009	0.038
Semantic	Left	55	-0.444	0.925	-2.059	1.013
Prosody	Left	55	1.441	0.947	-0.009	3.089
Semantic	Right	55	-0.344	0.782	-1.647	0.852
Prosody	Right	55	0.793	0.786	-0.474	2.103

Note: *All estimates are in log-odds. Estimates come from slope's posteriors.*

Results from RT data also replicate previous findings from Busch-Moreno et al. (2020). Table 6 and Table 7 summarise results from direct- and indirect-threat conditions respectively. Note the big increase in estimates from lowest BIS scores to highest BIS scores, with HDIs well apart from each other. When intercepts are taken into account, the Direct-threat estimates portray an average effective increase of ~175ms from lowest (1 point) to highest (55 points) BIS score by Prosody at the left ear, and ~164ms at the right ear. While for Semantic the increase is ~172ms at the left ear and ~150ms at the right. The apparent left ear advantage for Semantic can be discounted due to HDI overlap. Indirect-threat results are similar, excepting that responses tend to be slightly slower. However, overall, RTs were the shortest for Congruent; more details on these additional patterns can be found in the OSF repository (https://osf.io/n5b6h/).

Table 6. Direct-threat Reaction Time Estimates

Type	Ear	BIS Score	Posterior Mean	Posterior SD	HDI 5%	HDI 95%
Semantic	Left	1	3.20	0.40	2.54	3.84
Prosody	Left	1	3.25	0.39	2.62	3.90
Semantic	Right	1	2.79	0.33	2.25	3.35
Prosody	Right	1	3.04	0.34	2.49	3.60
Semantic	Left	55	176.09	21.76	139.95	211.20
Prosody	Left	55	178.50	21.64	143.94	214.60
Semantic	Right	55	153.21	18.25	123.59	184.04
Prosody	Right	55	167.04	18.87	137.08	198.26

Note: *All estimates indicate reaction times in milliseconds (*ms*). Estimates come from slope's posteriors.*

Table 7. Indirect-threat Reaction time estimates

Type	Ear	BIS Score	Posterior Mean	Posterior SD	HDI 5%	HDI 95%
Semantic	Left	1	4.37	0.41	3.71	5.06
Prosody	Left	1	4.29	0.41	3.64	4.96
Semantic	Right	1	4.23	0.35	3.65	4.79
Prosody	Right	1	3.79	0.35	3.22	4.35
Semantic	Left	55	240.48	22.47	204.31	278.46
Prosody	Left	55	235.85	22.29	200.14	272.72
Semantic	Right	55	232.38	19.09	200.92	263.32
Prosody	Right	55	208.63	19.03	177.11	239.50

Note: *All estimates indicate reaction times in milliseconds (*ms*). Estimates come from slope's posteriors.*

ERP Results

Although global field power (GFP) at EEG channels does not drift and EEG polarity is aligned (see: Luck, 2014), visual inspection of grand average ERP waves for direct- and indirect-threat conditions also indicate a potentially artifactual drift at EOG activity (Figure 4). This also shows that ERP waves are similar across tasks (direct- and indirect-threat), indicating a systematic amplitude increase after 500ms. Figure 5 and Figure 6 show mean amplitude peaks selected by maximum local GFP, which indicates a spatial standard deviation quantifying electrical activity from all electrodes at specific time-points (Skrandies, 1990), reliably revealing peaks of greater amplitude. As an initial descriptive inspection, these peaks may suggest the lack of earlier phases effects of BIS, as sufficiently strong amplitudes mostly appear after 400ms for higher BIS levels (divided into terciles for illustration only).

Figure 4. General ERP grand averages summary. Left panels show EOG and EEG waves from the direct-threat task, and right panels from the indirect-threat task. Bottom panels grand averages across all 64 electrodes. EOG: two electrooculogram channels. *N*ave: total number of trials averaged. GFP: global field power (local maximum).

Figure 5. Direct-threat topographies. Images show raw mean amplitude across epoch by BIS tercile (for illustration purposes only). Top: upper BIS score tercile. Middle: mid BIS score tercile. Bottom: lower BIS score tercile. Time-points were selected as the local maxim Global Field Power (GFP).

Figure 6. Indirect-threat topographies. Images show raw mean amplitude across epoch by BIS tercile (for illustration purposes only). Top: upper BIS score tercile. Middle: mid BIS score tercile. Bottom: lower BIS score tercile. Time-points were selected as the local maxim Global Field Power (GFP).

Hierarchical models indicate that excepting window4 (500-750ms), no other time-window showed relevant effects (HIDs overlapping zero or widely inside ROPE). All models sampled well, showing excellent convergence ($\hat{R} \cong 1$, ESS > 300, BFMIs > 0.6). Tables 8 and 9 summarise estimates that represent posteriors from BIS slope of the regression. Table 8 summarise direct-threat results from window4, showing some example electrodes with the most consistent amplitude increases as a function of BIS. This indicates that within this group of electrodes amplitude increases between 0.030 and 0.055μV per BIS score. At the electrode with maximum increase (TP7) amplitude estimated increase ranges from 0.055μV to 3.025μV; that is, from 1 BIS point to 55 BIS points amplitude is estimated to increase ~2.5μV. Table 9 summarises the same electrodes from the indirect-threat task (window4), indicating very similar results. Similar increases were registered at for electrodes such as P7, P8, P9 and P10. Supplementary materials contain forest plots of estimates of all relevant electrodes (Supplement 2, supplementary figures 1 and 2).

Note that in both tables HDIs (5%) remain well above zero and do not touch (in some cases barely touch) the 2SD ROPE. This implies that the magnitude of effects shows good certainty, indicating a sustained increase in amplitude from the lowest BIS level (1 point) to the highest BIS level (55 points), with good precision (HDI widths do not surpass ROPE width). This indicates that the greatest amplitude increases in window4 (500-750ms) occur at bilateral temporo-parietal electrodes, but with a clearer LPC pattern at temporal sites (e.g. TP7, TP8). Figures 7 and 8 show average amplitude waves from example electrodes from the direct- and indirect-threat tasks respectively. Both tasks show a similar pattern and are consistent with models' estimates. In short, irrespective of task (answer to threat or neutral), ear presentation (left or right) or sentence-type (Congruent, Prosody or Semantic), after 500ms amplitude increases as a function of BIS score at bilateral tempo-parietal electrodes.

Table 8. Direct-threat Estimates of Selected Electrodes

Channel	Type	Ear	Mean	SD	HDI 5%	HDI 95%	ESS	$\hat{R}$
TP8	Prosody	left	0.035	0.008	0.019	0.049	2551.103	1.001
TP8	Semantic	left	0.041	0.008	0.027	0.057	2408.821	1.000
TP8	Congruent	left	0.038	0.008	0.023	0.053	2461.246	1.001
TP8	Prosody	right	0.030	0.008	0.014	0.044	2486.100	1.000
TP8	Semantic	right	0.044	0.008	0.029	0.059	2345.198	1.000
TP8	Congruent	right	0.040	0.008	0.024	0.055	2600.853	1.000
TP7	Prosody	left	0.050	0.008	0.035	0.066	2273.178	1.002
TP7	Semantic	left	0.051	0.008	0.036	0.067	2138.450	1.001
TP7	Congruent	left	0.043	0.008	0.028	0.059	2161.740	1.001
TP7	Prosody	right	0.049	0.008	0.032	0.064	2140.949	1.001
TP7	Semantic	right	0.055	0.009	0.038	0.070	2168.432	1.002
TP7	Congruent	right	0.045	0.009	0.028	0.061	2314.087	1.001
T8	Prosody	left	0.020	0.008	0.004	0.035	2848.618	1.000
T8	Semantic	left	0.032	0.008	0.017	0.047	2640.524	1.001
T8	Congruent	left	0.028	0.008	0.012	0.043	2599.219	1.000
T8	Prosody	right	0.029	0.008	0.014	0.044	2645.124	1.000
T8	Semantic	right	0.038	0.008	0.023	0.053	2823.160	1.001
T8	Congruent	right	0.046	0.008	0.031	0.061	2678.274	1.001
T7	Prosody	left	0.037	0.008	0.021	0.052	2201.530	1.003
T7	Semantic	left	0.043	0.008	0.028	0.059	2376.424	1.000
T7	Congruent	left	0.039	0.008	0.024	0.055	2428.522	1.000
T7	Prosody	right	0.043	0.009	0.027	0.059	2446.879	1.000
T7	Semantic	right	0.048	0.008	0.033	0.064	2510.958	1.000
T7	Congruent	right	0.041	0.008	0.026	0.057	2470.028	1.001

Note: All estimates indicate amplitude in microvolts (μV). Mean: BIS slope's posterior mean. SD: BIS slope's posterior standard deviation. HDI: BIS slope's posterior highest density interval. ESS: estimated sample size. $\hat{R}$: convergence statistic.

Table 9. Indirect-threat Estimates of Selected Electrodes

Channel	Type	Ear	Mean	SD	HDI 5%	HDI 95%	ESS	R̂
TP8	Prosody	left	0.030448	0.009	0.015	0.047	2656.280	1.000
TP8	Semantic	left	0.041245	0.009	0.025	0.057	2646.560	1.001
TP8	Congruent	left	0.043562	0.009	0.027	0.060	2358.431	1.000
TP8	Prosody	right	0.035502	0.009	0.020	0.051	2467.430	1.002
TP8	Semantic	right	0.039979	0.008	0.025	0.056	2492.759	1.002
TP8	Congruent	right	0.046347	0.008	0.031	0.062	2128.460	1.000
TP7	Prosody	left	0.047278	0.009	0.030	0.063	2761.056	1.000
TP7	Semantic	left	0.067855	0.008	0.053	0.084	2702.787	1.000
TP7	Congruent	left	0.035341	0.008	0.020	0.051	2826.380	1.000
TP7	Prosody	right	0.052123	0.008	0.037	0.069	3238.207	1.000
TP7	Semantic	right	0.065843	0.009	0.050	0.082	2677.620	1.000
TP7	Congruent	right	0.038637	0.008	0.024	0.054	2537.611	1.000
T8	Prosody	left	0.037257	0.009	0.020	0.053	2664.166	1.001
T8	Semantic	left	0.032783	0.008	0.016	0.048	2494.279	1.000
T8	Congruent	left	0.028632	0.009	0.013	0.046	2626.227	1.001
T8	Prosody	right	0.027142	0.008	0.011	0.042	2573.088	1.000
T8	Semantic	right	0.033297	0.008	0.019	0.050	2664.163	1.000
T8	Congruent	right	0.036383	0.008	0.021	0.051	2412.832	1.001
T7	Prosody	left	0.041609	0.009	0.024	0.059	2627.236	1.000
T7	Semantic	left	0.046354	0.009	0.030	0.063	2631.563	1.001
T7	Congruent	left	0.030891	0.009	0.014	0.047	2303.773	1.000
T7	Prosody	right	0.03706	0.009	0.020	0.053	2262.723	1.000
T7	Semantic	right	0.059059	0.009	0.041	0.075	2539.533	1.002
T7	Congruent	right	0.030344	0.009	0.014	0.047	2352.111	1.000

Note: All estimates indicate amplitude in microvolts (μV). Mean: BIS slope's posterior mean. SD: BIS slope's posterior standard deviation. HDI: BIS slope's posterior higher density interval. ESS: estimated sample size. R̂: convergence statistic.

Figure 7. Direct-threat example electrodes. Plots show grand averages (-100-1000ms epoch), low-pass filtered at 40Hz and divided into BIS terciles for display purposes only. High, red solid line: BIS points > 28. Mid, yellow dashed line: 29 > BIS points > 15. Low, blue dotted line: BIS points < 16. Black dotted vertical line indicates sentence onset, period before the dotted line is the 100ms baseline.

Figure 8. Indirect-threat example electrodes. Plots show grand averages (-100-1000ms epoch), low-pass filtered at 40Hz and divided into BIS terciles for display purposes only. High, red solid line: BIS points > 28. Mid, yellow dashed line: 29 > BIS points > 15. Low, blue dotted line: BIS points < 16. Black dotted vertical line indicates sentence onset, period before the dotted line is the 100ms baseline.

Discussion

Present behavioural results indicate that threatening stimuli in both direct- and indirect-threat tasks induce longer RTs as a function of BIS (trait anxiety), but have minor effects on accuracy. These results replicate our previous experiment very closely (Busch-Moreno et al., 2020), consistent with the interpretation of trait anxiety disrupting orientation phases, as accuracy is minorly affected but RTs are strongly affected by trait anxiety levels. ERP results indicate that after 500ms (window4: 500-750ms) amplitude increases as a function of BIS, which can be interpreted as BIS-related late positive complex (LPC). This is consistent with part of present hypothesis indicating late phase effects of trait anxiety on threatening language processing during orientative (deliberation) phases. However, no earlier effects can be observed as a function of BIS, which goes contrary to our hypotheses predicting the influence of trait anxiety at all time-windows from sentence's onset.

An important aspect of our results contradicting our predictions is the lack of laterality effects. Although some EEG studies show lateralization effects associated with emotional semantic and prosody variation (Kotz and Paulmann, 2007), recent research does not show much evidence for these effects (Chen et al., 2011; Paulmann et al., 2012). One proposed explanation is that concurrent multi-channel (prosody and semantic) information obscures laterality effects (Paulmann et al., 2012). In addition, the present LPC is not modulated by stimulus-type. This differs from previous research, which did not include anxiety measures, finding late positive amplitudes in association with prosody/semantic emotional variations and/or congruency effects (Astésano et al., 2004; Chen et al., 2011; Zhao et al., 2015). Given this, the present LPC, which shows a different distribution and different preceding ERPs from the aforementioned studies, could be understood as signalling something particular to anxiety rather than being directly associated with prosody or semantics.

Previous EEG studies have observed that worry is associated with a parietal late positive potential (LPP) increases and related with negative reappraisal of visual stimuli (Moser et al., 2014). This is understood to be associated with sustained attention inducing increased late-phase processing (Hajcak et al., 2010). This may correspond to delayed disengagement from threat (Cisler and Koster, 2010), which in the present paper we understand as over-engagement with threat during deliberation phases, occurring after 500ms (i.e. LPC time-window). However, studies using a cross-modal paradigm, priming words with emotional faces to observe implicit reappraisal

have observed early-phase (e.g. N170) but not late-phase (i.e. LPP) effects (Liu et al., 2018), suggesting implicit rather than explicit engagement with threat when stimuli are short and/or fear-inducing. EEG studies focusing on social anxiety have shown that abusive words (short duration stimuli) induce effects on early- or mid-phase ERPs (i.e. P1, N400) which could be better related with over-attention to threat or threat evaluation issues (Wabnitz et al., 2015). Other studies have shown that angry vocalisations (prosody) are associated with early (i.e. P1) but not late (i.e. LPC) amplitude increases associated with anxiety (Pell et al., 2015). So, for anxious participants to over-engage with threat longer stimuli/ISIs and less directly threatening (i.e. not fear inducing or directly harming) stimuli may be required (Cisler and Koster, 2010).

This could imply that the present LPC, peaking ~600ms after sentence onset, requires the deliberation time provided by long duration sentences. For the present experiment, sentences had a duration of ~1646ms on average, which gives plenty of time (during sentence) for assessing and re-assessing stimuli before responses are executed after sentence's offset. In other words, with sufficient time to over-engage with threatening stimuli in a decision task, effects were mainly driven by trait anxiety; namely higher BIS levels induced over-engagement with threat, resulting in increased processing during late-phase deliberation (after 500ms). This is consistent with LPC evidencing decision modulation in relation to evaluation through memory processes (Finningan, 2002; Yang et al., 2019), or LPP/LPC evidencing late-phase processing associated with anxious negative re-appraisal (Hajcak et al., 2010; Moser et al., 2014). In this line, trait anxiety has been strongly associated with patterns of repetitive thinking (McEvoy et al., 2010; McLaughlin et al., 2007), which is a mechanism for re-appraisal and over-engagement. Presently, we propose that in the present experiment, threatening speech may have induced anxious participants to engage in verbal repetitive thinking, which can be associated with both memory-related evaluation and re-appraisal. This is theoretically sound respect to BIS and anxiety models (Bar-Haim et al., 2007; Cisler and Koster, 2010; McNaughton, 2011; Robinson et al., 2019); although verbal repetitive thinking is not the only possible mechanism for explaining over-engagement with threat. Here, as in our previous research (Busch-Moreno et al., 2020), we propose verbal repetitive thinking due to its possible relationship with threatening speech processing, but we acknowledge that present evidence is not enough as to confirm this explanation.

Although present research provides evidence of both behavioural and EEG effects of trait anxiety (BIS) on responses to threatening speech, we need to address some relevant limitations. Firstly, we do not have a direct neutral-speech control, as the only alternative task requiring participants to answer to neutral sentences (indirect-threat), usually intended to control for attentional differences (e.g. Peschard et al., 2016; Sander et al., 2005), still exposes participants to threatening speech. Secondly, although precision of the estimates indicates that RT and ERP models are sufficiently powered, accuracy analyses indicate that logistic regression models may need more participants and responses to provide more precise and certain estimates. Thirdly, even though we have applied an automated pre-processing and cleaning procedure to EEG data, rigorously based on previous literature, we acknowledge that there may still be a positive artefactual drift present in our epoched data. However, if the observed LPC component is not really an LPC (i.e. derived from brain activity), its association with BIS (trait anxiety) still needs to be explained. For instance, it may be the case that increased base muscle tension (e.g. jaw clenching) or rumination-related oro-facial muscle activity, characteristic of anxiety (Nalborczyk, 2017), induced artifacts on the ERP waves. However, muscular activity should induce higher frequency activity rather than lower frequency drifts, especially if related with temporalis muscles affecting temporal electrodes (Luck, 2014). Such artifacts, nonetheless, may have been detrimental for earlier activity (before 400-500ms), which may also explain the difficulty in observing earlier ERP effects. Despite this, excessive muscle tension could have indirectly induced eye artifacts at lower frequencies and amplitudes, which may have not been caught by ICA and other cleaning procedures. However, here we cannot provide direct evidence for this possible anxiety-related EOG activity or find an explicit account of this possibility in the literature.

Besides this, there are some design limitations that should be acknowledged. Presently, the high variability of type and sentence position of threatening lexical items in Semantic and Congruent conditions may have led participants to primarily carry out threat recognition by focusing on threatening prosody, which always started at sentence onset. Thus, possible ear differences may have become very difficult to observe at early processing stages and/or quickly receded before later stages. In addition, this possibly strategic response of participants (focus on prosody) may have obscured sentence-type effects. Better controlled semantic stimuli could help to addressed this issue, albeit at the cost of the semi-naturalness of stimuli. Also, it may

be useful to employ methods that can directly induce response inhibition to differentiated stimuli (i.e. responses to threatening content or sound), rather than decision tasks as employed here. Previous research has indeed proposed that tasks such as go/no-go are better for understanding BIS processes, because they directly involve inhibiting responses as part of their design (e.g. McNaughton et al., 2013; Neo et al., 2011). Future research could implement such tasks by, for instance, including responses to stimuli type instead to threat in general. Finally, it would be relevant to implement tasks which can compare language stimuli with other type of stimuli and have a proper neutral-stimuli control; this could help to determine whether anxiety effects are specifically related to threatening speech or, instead, to threat in general. In other words, to determine whether the observed slower RTs and increasing LPC are task-induced rather than threat-induced.

In conclusion, present experimental results provide strong evidence for trait anxiety effects, replicating our previous behavioural results (Busch-Moreno et al., 2020). ERP analyses show that these responses are preceded by a clear positive amplitude deflection in temporal electrodes peaking at around 600ms. We interpreted this ERP as an LPC associated with trait anxiety affecting deliberation processes through verbal repetitive thinking. As BIS scores increased, the LPC became more positive, suggesting a disruptive deliberation process, such as induced by over-engagement with threat, during an orientative stage. In other words, higher levels of anxiety due to their association with repetitive thinking (i.e. worry and rumination) induce a more effortful decision-making process after evaluation of threatening stimuli. In addition, the observed relationship between anxiety and language is not specific to speech information type. Hence, the effects of anxiety on prosody and semantics remain to be further explored. Furthermore, verbal repetitive thinking remains a tentative explanation, as it may be the case that a more general or fundamental mechanism associated with anxiety/BIS induces the observed effects. Overall, this experiment paves the way for future research on the relationship between speech, individual differences and emotional language in terms of information channels, anxiety and threatening language. This outlines an improved basis for reshaping models of emotional language processing by providing clear evidence on how anxiety can affect late processing phases.

Authors Contributions

Contributions to the study were as follows. **Busch-Moreno**: Conceptualization, Methodology, Software, Analysis, Investigation, Resources, Curation, Writing, Visualization, Administration. **Vinson**: Conceptualization, Methodology, Revision, Supervision, Administration. **Tuomainen**: Conceptualization, Resources.

Data statement

Declaration of interests

None.

Funding Sources

This research is supported by Becas Chile, ANID.

References

Ablin, P.; Cardoso, J.; Gramfort, A. (2017). Faster independent component analysis by preconditioning with Hessian approximations. ArXiv Preprint, https://arxiv.org/abs/1706.08171

Astésano, C., Besson, M., & Alter, K. (2004). Brain potentials during semantic and prosodic processing in French. *Cognitive Brain Research*, *18*(2), 172–184. https://doi.org/10.1016/j.cogbrainres.2003.10.002

Banse, R., & Scherer, K. R. (1996). Acoustic profiles in vocal emotion expression. *Journal of Personality and Social Psychology*, *70*(3), 614–636. https://doi.org/10.1037//0022-3514.70.3.614

Bar-Haim, Y., Lamy, D., Pergamin, L., Bakermans-Kranenburg, M. J., & van IJzendoorn, M. H. (2007). Threat-related attentional bias in anxious and nonanxious individuals: A meta-analytic study. *Psychological Bulletin*, *133*(1), 1–24. https://doi.org/10.1037/0033-2909.133.1.1

Belin, P., Fecteau, S., & Bédard, C. (2004). Thinking the voice: neural correlates of voice perception. *Trends in Cognitive Sciences*, *8*(3), 129–135. https://doi.org/10.1016/j.tics.2004.01.008

Bruder, G. E., Schneier, F. R., Stewart, J. W., McGrath, P. J., & Quitkin, F. (2004). Left Hemisphere Dysfunction During Verbal Dichotic Listening Tests in Patients Who Have Social Phobia With or Without Comorbid Depressive Disorder. *American Journal of Psychiatry*, *161*(1), 72–78. https://doi.org/10.1176/appi.ajp.161.1.72

Busch-Moreno, S., Vinson, D., & Tuomainen, J. (2020). Semantic and Prosodic Threat Processing in Trait Anxiety: Is Repetitive Thinking Influencing Responses? *Cognition and Emotion*. In press. https://doi.org/10.1080/02699931.2020.1804329

Chen, X., Zhao, L., Jiang, A., & Yang, Y. (2011). Event-related potential correlates of the expectancy violation effect during emotional prosody processing. *Biological Psychology*, *86*(3), 158–167. https://doi.org/10.1016/j.biopsycho.2010.11.004

Cisler, J. M., & Koster, E. H. W. (2010). Mechanisms of attentional biases towards threat in anxiety disorders: An integrative review. *Clinical Psychology Review*, *30*(2), 203–216. https://doi.org/10.1016/j.cpr.2009.11.003

Corr, P. J., & Cooper, A. J. (2016). The Reinforcement Sensitivity Theory of Personality Questionnaire (RST-PQ): Development and validation. *Psychological Assessment*, *28*(11), 1427–1440. https://doi.org/10.1037/pas0000273

Corr, P. J., & McNaughton, N. (2012). Neuroscience and approach/avoidance personality traits: A two stage (valuation–motivation) approach. *Neuroscience & Biobehavioral Reviews*, *36*(10), 2339–2354. https://doi.org/10.1016/j.neubiorev.2012.09.013

De Pascalis, V., Cozzuto, G., Caprara, G. V., & Alessandri, G. (2013). Relations among EEG-alpha asymmetry, BIS/BAS, and dispositional optimism. *Biological Psychology*, *94*(1), 198–209. https://doi.org/10.1016/j.biopsycho.2013.05.016

Dien, J. (1998). Issues in the application of the average reference: Review, critiques, and recommendations. Behavior Research Methods, Instruments, & Computers, 30(1), 34-43. doi: 10.3758/bf03209414

Finnigan, S. (2002). ERP "old/new" effects: memory strength and decisional factor(s). *Neuropsychologia*, *40*(13), 2288–2304. https://doi.org/10.1016/s0028-3932(02)00113-6

Gadea, M., Espert, R., Salvador, A., & Martí-Bonmatí, L. (2011). The sad, the angry, and the asymmetrical brain: Dichotic Listening studies of negative affect and depression. *Brain and Cognition*, *76*(2), 294–299. https://doi.org/10.1016/j.bandc.2011.03.003

Godfrey, H. K., & Grimshaw, G. M. (2015). Emotional language is all right: Emotional prosody reduces hemispheric asymmetry for linguistic processing. *Laterality: Asymmetries of Body, Brain and Cognition*, *21*(4–6), 568–584. https://doi.org/10.1080/1357650x.2015.1096940

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., … Hämäläinen, M. S. (2014). MNE software for processing MEG and EEG data. *NeuroImage*, *86*(2014), 446–460. https://doi.org/10.1016/j.neuroimage.2013.10.027

Grimshaw, G. M., Kwasny, K. M., Covell, E., & Johnson, R. A. (2003). The dynamic nature of language lateralization: effects of lexical and prosodic factors. *Neuropsychologia*, *41*(8), 1008–1019. https://doi.org/10.1016/s0028-3932(02)00315-9

Hajcak, G., MacNamara, A., & Olvet, D. M. (2010). Event-Related Potentials, Emotion, and Emotion Regulation: An Integrative Review. *Developmental Neuropsychology*, *35*(2), 129–155. https://doi.org/10.1080/87565640903526504

Hammerschmidt, K., & Jürgens, U. (2007). Acoustical Correlates of Affective Prosody. *Journal of Voice*, *21*(5), 531–540. https://doi.org/10.1016/j.jvoice.2006.03.002

Heller, W., Nitschke, J. B., & Lindsay, D. L. (1997). Neuropsychological Correlates of Arousal in Self-reported Emotion. *Cognition & Emotion*, *11*(4), 383–402. https://doi.org/10.1080/026999397379854

Hugdahl, K. (2011). Fifty years of dichotic listening research – Still going and going and…. *Brain and Cognition*, *76*(2), 211–213. https://doi.org/10.1016/j.bandc.2011.03.006

Hunter, J. D. (2007). Matplotlib: A 2D Graphics Environment. *Computing in Science & Engineering*, *9*(3), 90–95. https://doi.org/10.1109/mcse.2007.55

Jadoul, Y., Thompson, B., & de Boer, B. (2018). Introducing Parselmouth: A Python interface to Praat. *Journal of Phonetics*, *71*(71), 1–15. https://doi.org/10.1016/j.wocn.2018.07.001

Jas, M., Engemann, D., Bekhti, Y., Raimondo, F., & Gramfort, A. (2017). Autoreject: Automated artifact rejection for MEG and EEG data. Neuroimage, 159, 417-429. doi: 10.1016/j.neuroimage.2017.06.030

Jas, M., Larson, E., Engemann, D., Leppäkangas, J., Taulu, S., Hämäläinen, M., & Gramfort, A. (2018). A Reproducible MEG/EEG Group Study With the MNE Software: Recommendations, Quality Assessments, and Good Practices. Frontiers In Neuroscience, 12. doi: 10.3389/fnins.2018.00530

Keune, P. M., Bostanov, V., Kotchoubey, B., & Hautzinger, M. (2012). Mindfulness versus rumination and behavioral inhibition: A perspective from research on frontal brain asymmetry. *Personality and Individual Differences*, *53*(3), 323–328. https://doi.org/10.1016/j.paid.2012.03.034

Kotz, S. A., & Paulmann, S. (2007). When emotional prosody and semantics dance cheek to cheek: ERP evidence. *Brain Research*, *1151*(2007), 107–118. https://doi.org/10.1016/j.brainres.2007.03.015

Kotz, S. A., & Paulmann, S. (2011). Emotion, Language, and the Brain. *Language and Linguistics Compass*, *5*(3), 108–125. https://doi.org/10.1111/j.1749-818x.2010.00267.x

Kruschke, J. K. (2013). Bayesian estimation supersedes the t test. *Journal of Experimental Psychology: General*, *142*(2), 573–603. https://doi.org/10.1037/a0029146

Kruschke, J. K. (2015). *Doing Bayesian data analysis: a tutorial with R, JAGS, and stan*. Amsterdam Etc: Elsevier, Academic Press, Cop.

Kumar, R., Carroll, C., Hartikainen, A., & Martin, O. (2019). ArviZ a unified library for exploratory analysis of Bayesian models in Python. *Journal of Open Source Software*, *4*(33), 1143. https://doi.org/10.21105/joss.01143

Lei, X., & Liao, K. (2017). Understanding the Influences of EEG Reference: A Large-Scale Brain Network Perspective. Frontiers In Neuroscience, 11. doi: 10.3389/fnins.2017.00205

Leshem, R. (2018). Trait Anxiety and Attention: Cognitive Functioning as a Function of Attentional Demands. *Current Psychology*. https://doi.org/10.1007/s12144-018-9884-9

Liebenthal, E., Desai, R., Ellingson, M. M., Ramachandran, B., Desai, A., & Binder, J. R. (2010). Specialization along the Left Superior Temporal Sulcus for Auditory Categorization. *Cerebral Cortex*, *20*(12), 2958–2970. https://doi.org/10.1093/cercor/bhq045

Luck, S. J. (2014). *An introduction to the event-related potential technique* (Second). MIT Press.

Martin, O. (2018). *Bayesian analysis with Python : introduction to statistical modeling and probabilistic programming using PyMC3 and ArviZ*. Birmingham, Uk: Packt Publishing.

McElreath, R. (2020). *Statistical Rethinking: A Bayesian Course with Examples in R and STAN* (Second). CRC Press.

McEvoy, P. M., Mahoney, A. E. J., & Moulds, M. L. (2010). Are worry, rumination, and post-event processing one and the same? *Journal of Anxiety Disorders*, *24*(5), 509–519. https://doi.org/10.1016/j.janxdis.2010.03.008

McLaughlin, K. A., Borkovec, T. D., & Sibrava, N. J. (2007). The Effects of Worry and Rumination on Affect States and Cognitive Activity. *Behavior Therapy*, *38*(1), 23–38. https://doi.org/10.1016/j.beth.2006.03.003

McNaughton, N., & Gray, J. A. (2000). Anxiolytic action on the behavioural inhibition system implies multiple types of arousal contribute to anxiety. *Journal of Affective Disorders*, *61*(3), 161–176. https://doi.org/10.1016/s0165-0327(00)00344-x

McNaughton, N., Swart, C., Neo, P., Bates, V., & Glue, P. (2013). Anti-anxiety drugs reduce conflict-specific "theta"—A possible human anxiety-specific biomarker. *Journal of Affective Disorders*, *148*(1), 104–111. https://doi.org/10.1016/j.jad.2012.11.057

Moser, J. S., Hartwig, R., Moran, T. P., Jendrusina, A. A., & Kross, E. (2014). Neural markers of positive reappraisal and their associations with trait reappraisal and worry. *Journal of Abnormal Psychology*, *123*(1), 91–105. https://doi.org/10.1037/a0035817

Nalborczyk, L., Perrone-Bertolotti, M., Baeyens, C., Grandchamp, R., Polosan, M., Spinelli, E., Koster, E. H. W., & Lœvenbruck, H. (2017). Orofacial electromyographic correlates of induced verbal rumination. *Biological Psychology*, *127*(127), 53–63. https://doi.org/10.1016/j.biopsycho.2017.04.013

Neal, L. B., & Gable, P. A. (2017). Regulatory control and impulsivity relate to resting frontal activity. *Social Cognitive and Affective Neuroscience*, *12*(9), 1377–1383. https://doi.org/10.1093/scan/nsx080

Neo, P. S.-H., Thurlow, J. K., & McNaughton, N. (2011). Stopping, goal-conflict, trait anxiety and frontal rhythmic power in the stop-signal task. *Cognitive, Affective, & Behavioral Neuroscience*, *11*(4), 485–493. https://doi.org/10.3758/s13415-011-0046-x

Nitschke, J. B., Heller, W., Palmieri, P. A., & Miller, G. A. (1999). Contrasting patterns of brain activity in anxious apprehension and anxious arousal. *Psychophysiology*, *36*(5), 628–637. https://doi.org/10.1111/1469-8986.3650628

Nygaard, L. C., Herold, D. S., & Namy, L. L. (2009). The Semantics of Prosody: Acoustic and Perceptual Evidence of Prosodic Correlates to Word Meaning. *Cognitive Science*, *33*(1), 127–146. https://doi.org/10.1111/j.1551-6709.2008.01007.x

Paulmann, S., Jessen, S., & Kotz, S. A. (2012). It's special the way you say it: An ERP investigation on the temporal dynamics of two types of prosody. *Neuropsychologia*, *50*(7), 1609–1620. https://doi.org/10.1016/j.neuropsychologia.2012.03.014

Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., … Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior Research Methods*, *51*(1), 195–203. https://doi.org/10.3758/s13428-018-01193-y

Pell, M. D., Rothermich, K., Liu, P., Paulmann, S., Sethi, S., & Rigoulot, S. (2015). Preferential decoding of emotion from human non-linguistic vocalizations versus speech prosody. *Biological Psychology*, *111*(2015), 14–25. https://doi.org/10.1016/j.biopsycho.2015.08.008

Peschard, V., Gilboa-Schechtman, E., & Philippot, P. (2016). Selective attention to emotional prosody in social anxiety: a dichotic listening study. *Cognition and Emotion*, *31*(8), 1749–1756. https://doi.org/10.1080/02699931.2016.1261012

Poeppel, D., Idsardi, W. J., & van Wassenhove, V. (2008). Speech perception at the interface of neurobiology and linguistics. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *363*(1493), 1071–1086. https://doi.org/10.1098/rstb.2007.2160

Robinson, O. J., Pike, A. C., Cornwell, B., & Grillon, C. (2019). The translational neural circuitry of anxiety. *Journal of Neurology, Neurosurgery & Psychiatry*, jnnp-2019-321400. https://doi.org/10.1136/jnnp-2019-321400

Salvatier, J., Wiecki, T. V., & Fonnesbeck, C. (2016). Probabilistic programming in Python using PyMC3. *PeerJ Computer Science*, *2*, e55. https://doi.org/10.7717/peerj-cs.55

Sander, D., Grandjean, D., Pourtois, G., Schwartz, S., Seghier, M. L., Scherer, K. R., & Vuilleumier, P. (2005). Emotion and attention interactions in social cognition: Brain regions involved in processing anger prosody. *NeuroImage*, *28*(4), 848–858. https://doi.org/10.1016/j.neuroimage.2005.06.023

Schirmer, A., & Kotz, S. A. (2006). Beyond the right hemisphere: brain mechanisms mediating vocal emotional processing. *Trends in Cognitive Sciences*, *10*(1), 24–30. https://doi.org/10.1016/j.tics.2005.11.009

Spielberg, J. M., De Leon, A. A., Bredemeier, K., Heller, W., Engels, A. S., Warren, S. L., … Miller, G. A. (2013). Anxiety type modulates immediate versus delayed engagement of attention-related brain regions. *Brain and Behavior*, *3*(5), 532–551. https://doi.org/10.1002/brb3.157

Tanner, D., Morgan-Short, K., & Luck, S. J. (2015). How inappropriate high-pass filters can produce artifactual effects and incorrect conclusions in ERP studies of language and cognition. *Psychophysiology*, *52*(8), 997–1009. https://doi.org/10.1111/psyp.12437

van Heuven, W. J. B., Mandera, P., Keuleers, E., & Brysbaert, M. (2014). Subtlex-UK: A New and Improved Word Frequency Database for British English. *Quarterly Journal of Experimental Psychology*, *67*(6), 1176–1190. https://doi.org/10.1080/17470218.2013.850521

VanRullen, R. (2011). Four Common Conceptual Fallacies in Mapping the Time Course of Recognition. *Frontiers in Psychology*, *2*. https://doi.org/10.3389/fpsyg.2011.00365

Wabnitz, P., Martens, U., & Neuner, F. (2015). Written threat: Electrophysiological evidence for an attention bias to affective words in social anxiety disorder. *Cognition and Emotion*, *30*(3), 516–538. https://doi.org/10.1080/02699931.2015.1019837

Warriner, A. B., Kuperman, V., & Brysbaert, M. (2013). Norms of valence, arousal, and dominance for 13,915 English lemmas. *Behavior Research Methods*, *45*(4), 1191–1207. https://doi.org/10.3758/s13428-012-0314-x

Widmann, A., Schröger, E., & Maess, B. (2015). Digital filter design for electrophysiological data – a practical approach. Journal Of Neuroscience Methods, 250, 34-46. doi: 10.1016/j.jneumeth.2014.08.002

Winkler, I., Debener, S., Muller, K., & Tangermann, M. (2015). On the influence of high-pass filtering on ICA-based artifact reduction in EEG-ERP. 2015 37Th Annual International Conference Of The IEEE Engineering In Medicine And Biology Society (EMBC). doi: 10.1109/embc.2015.7319296

Yang, H., Laforge, G., Stojanoski, B., Nichols, E. S., McRae, K., & Köhler, S. (2019). Late positive complex in event-related potentials tracks memory signals when they are decision relevant. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598-019-45880-y

Zatorre, R. J., Belin, P., & Penhune, V. B. (2002). Structure and function of auditory cortex: music and speech. *Trends in Cognitive Sciences*, *6*(1), 37–46. https://doi.org/10.1016/s1364-6613(00)01816-7

Zhao, J., Liang, W.-K., Juan, C.-H., Wang, L., Wang, S., & Zhu, Z. (2015). Dissociated stimulus and response conflict effect in the Stroop task: Evidence from evoked brain potentials and brain oscillations. *Biological Psychology*, *104*(2015), 130–138. https://doi.org/10.1016/j.biopsycho.2014.12.001