Title: Sucrose transport and Sucrose Synthase activity control the source-sink

relationship for biomass and grain yield in rice

Running Title: Source-sink relationship in cultivated and wild rice

Jyotirmaya Mathan, Anuradha Singh, Aashish Ranjan[*]

National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi-110067,

India.

*Corresponding Author

e-mail: aranjan@nipgr.ac.in

Phone no. +91-11-26735117

Authors' email

Jyotirmaya Mathan: jyotirmaya@nipgr.ac.in

Anuradha Singh: annusingh1206@gmail.com

Date of submission: 19[th] October 2020

Number of figures: 7 figures

The word count: 5,085

Supplementary data: 10 supplementary figures and 1 supplementary table

Title: Sucrose transport and Sucrose Synthase activity control the source-sink
relationship for biomass and grain yield in rice

Running Title: Source-sink relationship in cultivated and wild rice

Highlight

Differences in vascular features, sucrose transport, and Sucrose Synthase activity contribute
to the differential source-sink relationship between the selected cultivated and wild rice that
differ in biomass and grain yield.

Abstract

The source-sink relationship is key to overall crop performance. Detailed understanding of
the factors that determine source-sink dynamics is imperative for the balance of biomass and
grain yield in a crop plant. We investigated the differences in the source-sink relationship
between a cultivated rice *Oryza sativa* cv. Nipponbare and a wild rice *Oryza australiensis*
that show striking differences in biomass and grain yield. The wild rice, accumulating higher
biomass, was not only photosynthetically efficient but also had efficient sucrose export from
leaves. However, sucrose mobilization to the grains was limited due to impaired vasculature
and sucrose transporter functions at the reproductive tissues of the wild rice. High cleavage
activity of Sucrose Synthase followed by higher expression of Cellulose Synthase genes in
the wild rice stem efficiently utilized photosynthates for the synthesis of structural
carbohydrates, resulting in high biomass. In contrast, the source-sink relationship favored
high grain yield in Nipponbare via accumulation of transitory starch in the stem, due to
higher expression of starch biosynthetic genes, which is mobilized to panicles at the grain
filling stage. Thus, sucrose transport along with functions of key sugar metabolic enzymes
explained the differences in the source-sink relationship between the selected cultivated and
wild rice.

Keywords: Biomass; Grain yield; Photosynthesis; Source-sink relationship; Starch;
Structural carbohydrates; Sucrose synthase; Sucrose transport; Vascular features

Introduction

Source-sink relationship is critical for determining overall growth and performance of crop plants (White *et al*., 2016). Photosynthetic carbon assimilation in source leaves, sucrose transport through vasculature, type and strength of sink tissues, as well as metabolic status of the tissues are the key features that determine the source-sink dynamics of a plant. Photosynthetic leaves are the major source tissues of a plant, whereas types and efficiency of sink organs vary depending upon the growth stages and environmental conditions. For example, grains are the primary sink in the reproductive stage of crop plants, while roots, stem/internodes, and growing leaves function as the sink in the vegetative stage. Natural genetic variation for differences in the source-sink dynamics, resulting in changes in biomass and yield, has been evident (Burnett *et al*., 2016; White *et al*., 2016; Fabre *et al.,* 2020). For example, many of the wild relatives of rice accumulate high biomass with poor grain filling, whereas cultivated varieties produce higher grain yield (Sanchez *et al*., 2013).

Leaf morphological and anatomical features along with photosynthesis per unit leaf area largely determine the source strength of a plant (Mathan *et al*., 2016). Functions of the key carbohydrate metabolic enzymes, such as ADP-glucose pyrophosphorylase (AGPase), sucrose phosphate synthase (SPS), sucrose phosphate phosphatase (SPP) and Sucrose synthase (SUS), are known to be key determinants of the metabolic status of source leaves (Osorio *et al*., 2014; Ruan, 2014). Similarly, the capacity to utilize photosynthates efficiently largely determines the sink strength. Sink size, influenced by tiller number, spikelet number per panicle, and seed size; and sink activity, determined by functions of various metabolic enzymes such as cell wall invertase (cwINV) and sucrose synthase (SUS), dictate the photosynthates utilization at a sink organ (Smith *et al*., 2018; Stein and Granot, 2019). Photosynthates are primarily partitioned from source leaves to different sink tissues in the form of sucrose via phloem with the aid of SWEET and SUT transporters (Chen *et al*., 2012; Braun, 2012; Julius *et al*., 2017). Mutations or overexpression of genes encoding sucrose transporters affect plant yield and biomass via modulation of source-sink dynamics (Scofield *et al*., 2002; Chen *et al*., 2012; Yang *et al*., 2018). Besides the sucrose transporter functions, vascular features not only determine the rate of sucrose export from leaf tissues to phloem sap but also control mobilization of photosynthates to the panicles at the reproductive stage (Qi *et al*., 2008; Fujita *et al*., 2013; Sack and Scoffoni, 2013).

Photosynthates assimilated in leaves are either used to meet immediate cellular needs or mobilized into different sinks. Starch stored in leaves during the day, over and above the immediate utilization and transport of photosynthates, is degraded during the night to be

transported to different sink organs for growth (Smith and Stitt, 2007; Chen *et al.*, 2012).
Mutants defective in sucrose transport, usually, accumulate starch in leaves at the end of the
night, limiting plant growth (Chen *et al.*, 2012). The utilization and storage of sugars are
regulated diurnally as well as developmentally. Stem serves as the main storage organ for
carbohydrates in crop plants, including rice, during vegetative growth. Sucrose, after getting
unloaded into the stem, could possibly be converted to different non-structural carbohydrates
(NSCs), such as glucose, fructose, and starch. Starch is reported to be the major storage form
in rice stem during the vegetative stage, which is mobilized to developing grains with the
onset of the reproductive stage, contributing up to half of the final grain yield (Wang *et al.*,
2017). A portion of the photosynthates is also used as a structural component of the cell wall
for the growth and development of new organs as well as elongation and thickening of
existing organs. High biomass accumulation is usually associated with accumulation of
cellulose and other structural carbohydrates in crop plants, with cellulose alone contributing
around 25-50% of plant biomass (Haigler *et al.*, 2001). Thus, the balance of structural and
non-structural carbohydrates derived from the sucrose in the stem is critical for determining
biomass and grain yield in cereal crops.

Sucrose, after reaching to the stem, is hydrolysed by two classes of enzymes,
Invertases (INV) and Sucrose Synthase (SUS). Invertases cleave sucrose into glucose and
fructose, while Sucrose Synthase (SUS) reversibly catalyses the formation of UDP-glucose
and fructose. The UDP-glucose, generated by the cleavage activity of SUS, is used as a
precursor for cellulose synthesis. Further, UDP-glucose can be converted to ADP-glucose for
usage in starch synthesis, making SUS a key enzyme for biomass and grain yield (Smith *et*
al., 2012; Ruan, 2014; Stein and Granot, 2019). The functions of SUS is important for
phloem loading and unloading as well as for determining sink strength (Smith *et al.*, 2012;
Fan *et al.*, 2017; Yao *et al.*, 2020). Among the seven genes encoding SUS enzymes in rice,
OsSUS1 and *OsSUS3* contribute to seed starch biosynthesis (Cho *et al.*, 2011). Higher
expression of *OsSUS3* also led to increased levels of structural carbohydrates, cellulose and
hemicellulose, in rice (Fan *et al.*, 2017). *OsSUS1* is highly expressed in rice internodes, with
a strong correlation with the expression pattern of genes encoding Cellulose Synthase,
OsCES4, *OsCES7*, and *OsCES9* (Hirose *et al.*, 2008; Guevara *et al.*, 2014). Among Cellulose
Synthase enzymes, OsCES4, OsCES7, and OsCES9 have been reported to function in
secondary cell wall formation (Tanaka *et al.*, 2003; Wang *et al.*, 2016). Thus, functions of
enzymes, such as INV, SUS and CES, are important for determining source-sink dynamics
for biomass and grain yield.

Despite the key importance of sucrose partitioning and metabolism in determining

source-sink relationship, this aspect is largely unexplored for optimization of grain yield and

biomass in rice. Cultivated rice varieties and their wild relatives, with striking differences in

yield and biomass traits, provide an excellent system to investigate the mechanistic basis of

the source-sink relationship. We investigated the limitations related to sucrose transport and

metabolism for grain yield in wild rice *O. australiensis*, resulting in higher biomass,

compared to the cultivated variety *O. sativa* cv. Nipponbare. We report higher accumulation

of structural carbohydrates, such as cellulose and hemicellulose, in the wild rice is due to

higher SUS cleavage activity together with higher expression of genes encoding Cellulose

Synthases. Cultivated rice, in contrast, accumulated more starch in the stem due to higher

expression of genes encoding starch biosynthesis enzymes. We also established the

contribution of specific sucrose transporters and vascular features towards impaired

mobilization of photosynthates to *O. australiensis* panicles.

Materials and methods

Phenotyping of the selected rice accessions for biomass and yield-related traits

Two cultivated varieties, *Oryza sativa* ssp. *indica* cv IR 64 and *Oryza sativa* ssp. *japonica* cv

Nipponbare, and three wild rice species, *Oryza rufipogon* (IRGC 99562)*, Oryza latifolia*

(IRGC 99596)*,* and *Oryza australiensis* (IRGC 105272), were grown in the experimental

field of National Institute of Plant Genome Research, New Delhi (*latitude* 28°36

N; *longitude* 77°12 E; *altitude* 216 m) under natural growing conditions. At least ten

individual plants of each genotype were used for phenotyping. Plant architectural traits were

quantified at the milk-stage of grain-filling. Seed traits, and fresh and dry weight were

quantified at grain maturity stage. Plant height was measured as the height from the soil

surface to the tip of central panicle; total leaves per plant, tiller number per plant, and

internode number per the main stem were counted manually; flag leaf width was quantified as

the width of the widest portion of the flag leaf; and internode length was calculated using the

distance between two consecutive nodes on the main stem. Stem width and circumference

were quantified using Fiji-Image J software from images of hand-cut sections of the same

region of the main stem. Fresh weight was quantified by directly weighing the entire plant

mass, and dry weight was quantified by weighing the dried plant mass after drying the entire

plant mass at 60°C for five days. Spikelet per panicle was counted manually, whereas seed

length and seed width (50 seeds per replicate) were measured using a digital Vernier calliper.

For seed weight analysis, 500-seeds were randomly selected and weighed using an electronic

balance. Raw data of biomass and grain yield-related traits for each genotype were

normalized to median from the entire sample set for principal Component analysis. Then, all

the data was converted on a log base 2 scale on JMP software from SAS

(https://www.jmp.com/en_us/home.html).

Quantification of leaf photosynthesis rate and related physiological traits

Leaf photosynthesis rate (A), stomatal conductance to water (g_sH_2O), and intercellular CO_2

concentration (C_i) were recorded from flag leaves at booting, and milk- and dough- stage of

grain filling for *O. sativa* ssp. *indica* cv. Nipponbare and *O. australiensis* using a portable

photosynthesis system LI-6400XT (LI-COR Biosciences). Data were recorded from at least

15 independent plants of each genotype on a clear day between 9.00 h to 11.00 h under

ambient CO_2 level (C_a; 400 μmol mol^{-1}), air temperature (30 ºC), leaf temperature (27 ºC),

and photosynthetic photon flux density (PPFD, 1,500 μmol m^{-2} s^{-1}) conditions.

Quantification of glucose, fructose, and sucrose

The non-structural carbohydrates (NSCs: glucose, fructose, and sucrose) present in the flag

leaf and stem were quantified for *O. sativa* ssp. *indica* cv. Nipponbare and *O. australiensis* at

the milk-stage of grain-filling by Gas Chromatography-Mass Spectrometry (GC/MS). Flag

leaf and stem tissues were harvested from field-grown plants of both the genotypes at 9:00-

11:00 am. For the quantification of NSCs in flag leaves at the end of the day (EOD) and end

of the night, tissue samples were collected at 6.00 pm and 6.00 am, respectively. The tissues

were sampled from at least four individual plants per genotypes and immediately frozen into

liquid nitrogen after harvesting. A 100 mg fresh weight (FW) of the tissues were extracted

using 1.0 ml water: chloroform: methanol (1:1:2.5) spiked with 10 μl internal standard (1 mg

ml^{-1} ribitol in water). The mixture was vortexed, centrifuged and 200 μl of polar phase were

dried for 3 hours in a speed-vac concentrator. Next, dried samples were derivatized with 50

μl of 20 mg ml^{-1} methoxyamine in pyridine warmed for 37°C for 120 mins with shaking.

Metabolites were further derivatized by adding 70 μl of N-methyl N-trimethylsilyl-

trifluoroacetamide (MSTFA) at 37°C for 30 min. Once derivatization was completed,

samples were transferred to GC compatible vials and 1μl of samples were injected on GC-

MS-QP2010 (Shimadzu) as described in Lisec *et al.* (2006). For metabolite identification, the

mass spectra were matched with commercially available spectral libraries NIST08, NIST11

and Wiley08 for automated peak identification and validated by retention index (RI) libraries

(http://gmd.mpimp-golm.mpg.de/). The relative amount of metabolites was presented by total
ion current signal which was normalized to Ribitol and tissue weight.

Starch quantification

Starch was quantified from flag leaves, stems, matured seeds, and different internodes of *O.*
sativa ssp. *indica* cv. Nipponbare and *O. australiensis*. The quantification of starch from flag
leaves and stems of the two genotypes were performed at the milk-stage of grain-filling at the
time of phenotyping of photosynthesis. In addition, flag leaves samples were used for starch
quantification at EOD and EON. Starch was also quantified from leaves at vegetative and
booting stage, as well as in different internodes of the two species at the milk-stage of grain-
filling. Starch was quantified using Mega-Calc Total starch determination kit (K-TSTA;
Megazyme) according to the manufacturer's protocol. Four independent biological replicates
were analysed for each tissue type for each genotype.

[14]C labelled sucrose loading assay

The phloem loading capacity of [14]C labelled sucrose was quantified by following the method
as descried by Yadav *et al.* (2017). Briefly, rice flag leaves at milk-stage of grain-filling were
cut, and the leaf bases were immediately transferred to petri-dish submerged with $MES/CaCl_2$
buffer (20 mM MES, 2mM $CaCl_2$, pH 5.5 with KOH). Next, leaf discs (3.6 × 1.0 cm) were
excised using a cork borer, and immediately placed abaxial side down in the 24-well
microtiter plate pre-filled with 1 ml $MES/CaCl_2$ buffer spiked with 1mM sucrose solution
(1mM unlabelled Sucrose supplemented with 0.81µCi ml[-1] [14]C Sucrose). Leaf discs
immersed in the above solution were vacuum-infiltrated for at least 20 min with gentle
shaking at room temperature. Labelled leaf discs were transferred to a fresh microtiter plate
and washed twice with 1.0 ml $MES/CaCl_2$ buffer, blot dried on absorbent filter paper, and
immediately frozen on dry ice and lyophilized. Next, [14]C sucrose loading capacity into leaves
was quantified by scintillation counting (PerkinElmer). Four leaf discs were used for each
biological replicate, and six biological replicates were used for each genotype per experiment.

Sucrose quantification in phloem sap

The amount of sucrose in the phloem was quantified using the EDTA-facilitated exudation
technique described by King and Zeevaart (1974). Flag leaves of the cultivated and wild rice
were excised at the milk-stage of grain-filling at around 9-10 am (when leaf photosynthesis
was quantified), leaf bases were immediately recut under exudation buffer (10 mM, Hepes,

10 mM EDTA, pH 7.5) to allow dispersion of any leaf contaminants, and placed in a 15 ml
tube containing 5 ml of the exudation buffer. Samples were incubated in a humid chamber
(relative humidity >90%) in the dark to prevent evapotranspiration, and exudates were
collected after 5 h of incubation. 2 ml of sample aliquots were spiked with 50 µl internal
standard (1 mg ml^{-1} ribitol in water), and freeze-dried for sucrose quantification using GC-
MS based method as described earlier. Individual flag leaves from different plants were
considered as separate biological replicates, and four biological replicates were analyzed for
each genotype.
246

RNA isolation, cDNA synthesis and qRT-PCR analysis

Total RNA from flag leaf, stem, and panicle base at milk-stage of grain-filling was extracted
using TRIzol reagent (Invitrogen), whereas RNA from spikelet at milk-stage was extracted
using plant RNA purification kit (Sigma-Aldrich) according to manufacturer's protocol. One
microgram (1µg) of total RNA was reverse transcribed to first strand-cDNA by anchored
oligodT priming using Thermo Scientific RevertAid cDNA synthesis kit following
manufacturer's instructions. The primer pairs (Table S1) of selected targeted genes used in
this study produced a single product as viewed through dissociation curve analysis. Further,
the efficiency of each primers was also evaluated using a standard curve method where five
cDNA quantities (1, 25, 50, 75, and 100 ng) from each genotype was used to construct a
standard curve, and efficiency were calculated using the formula E = $(10^{[-1/slope]}-1)*100$.
Expression analysis of each gene was performed at least in three biological replicates for each
genotype. The transcript levels of genes were normalized independently to two stable internal
controls actin (LOC_Os03g50885) and ubiquitin (LOC_Os03g03920) with similar results.
Relative expressions of target genes are presented using actin as the internal control applying
$2^{-\Delta Ct}$ method by Livak and Schmittgen (2001). *In sillico* expression analysis of the genes was
performed using OS_AFFY_RICE_6 dataset present in Genevestigator database
(https://genevestigator.com/gv/).
265

Visualization and quantification of anatomical traits

267 The vascular features of flag leaves as well as vascular bundles in the inner and outer ring of
268 the stem, and panicle base (0.5 cm above the panicle node) were visualized and quantified
269 through hand-cut transverse sections. Flag leaves, stems and panicle bases were hand-
270 sectioned using a razor-blade at the milk-stage of grain-filling, at least from five independent
271 plants for each genotype, and stained with toluidine blue O (0.02% toluidine blue O in water)

as described by Mitra and Loqué, (2014). Photographs were taken under a bright field microscope (LMI). Quantification of the different anatomical traits were made using Fiji-Image J software (Wayne Rasband, National Institute of Health).

Cellulose staining and quantification

For histochemical visualisation of cellulose, transverse hand-sections of rice stem (approximately 20 μm thickness) at the milk-stage of grain-filling were stained using calcofluor white (0.2% calcofluor white M2R in water) staining solution as described by Ambavaram *et al.* (2011). Stained sections were photographed under UV light range using a fluorescence microscope (excitation filter: 340-380 nm; Carl Zeiss Microscope). The quantification of cellulose and hemicellulose was made using National Renewable Energy Laboratory (NREL) protocols as described in Mund *et al.* (2016). The quantifications were made in four biological replications for each genotype.

Estimation of Sucrose synthase synthesis and cleavage activity

The sucrose synthase (SUS) synthesis and cleavage activity were performed in stem tissue of the two genotypes at the milk-stage of grain-filling. Tissue samples were homogenized in extraction buffer (50 mM HEPES/NaOH (pH 7.5), 7.5 mM $MgCl_2$, 2 mM EDTA, 2% (w/v) PEG 8000, 2% (w/v) PVP and 5 mM DDT) at 4 °C. The homogenate was centrifuged for 1 min at ~16,000 × g, the pellet was discarded and the crude extract was used for enzyme assay. To quantify OsSUS synthesis activity, 50 μl of crude extract was incubated with 20 mM HEPES/NaOH buffer (pH 7.5), 5 mM $MgCl_2$, 20 mM KCl, 12 mM fructose, 0.4 mM phosphoenolpyruvate, 2 mM uridine diphosphate UDP-glucose, 20 U pyruvate kinase, 20 U lactate dehydrogenase, and 0.15 mM NADH in total 1.0 ml final reaction volume. Similarly, to quantify OsSUS cleavage activity, 50 μl of crude extract was incubated with 20 mM HEPES/NaOH buffer (pH 7.5), 100 mM sucrose, 2 mM UDP, 2mM $MgCl_2$, 0.005 UDP-glucose dehydrogenase, and 1.5 mM NAD^+ in total 1.0 ml final reaction volume. The reaction mixture was mixed gently, and decrease and increase in absorbance were recorded for synthesis and cleavage activity, respectively, at 340 nm continuously for 0 to 300 sec using a UV–Vis spectrophotometer. The OsSUS activity was calculated in terms of μmol NADH oxidized (synthesis activity) and NAD+ reduced (cleavage activity) per min per mg protein as described in Qazi *et al.* (2012).

Subcellular localization of *OsSUS1*

Full length of *OsSUS1* CDS without stop codon was cloned under constitutive 35S promoter with a C-terminal eYFP fusion into binary vector pEG10 (Earley *et al.*, 2006). The recombinant plasmid along with empty vector control were then, introduced into EHA105 strain of *Agrobacterium tumefaciens*. Each transformant was co-infiltrated with a plasma membrane-localized marker (PM-mCherry; Nelson *et al.*, 2007) into the leaves of *Nicotiana benthamiana*. Confocal laser scanning microscopy TCS SP5 (Leica Microsystems) was used to take the images with appropriate lasers.

Results

Biomass and grain yield differences among the selected wild and cultivated rice accessions

We initiated the study with two cultivated rice varieties, *Oryza sativa* ssp. *japonica* cv. Nipponbare and *Oryza sativa* ssp. *indica* cv. IR 64, and three wild relatives of rice, *Oryza rufipogon*, *Oryza latifolia,* and *Oryza australiensis* that show remarkable variations in overall growth and architectural features at both vegetative and reproductive stages (Fig. 1A). All the selected wild rice species grew taller compared to the two cultivated varieties (Fig 1B). Among all the accessions, *O. sativa* cv. Nipponbare produced lowest number of total leaves, while *O. rufipogon* showed the highest number of leaves per plant (Fig. 1C). Nipponbare showed a significantly lower number of leaves compared to all the selected wild species. In addition, Nipponbare also showed significantly lower tiller number per plant compared to all the selected accessions (Supplementary Fig. S1B). Selected wild rice species showed wider leaves compared to the cultivated varieties, with *O. latifolia* showing widest leaves (Fig. 1D). We, then, quantified number and length of internodes in the main stem, as internodes are important contributors to the biomass of a plant. Cultivated rice varieties had fewer, smaller, and slender internodes in the main stem compared to the wild accessions (Fig. 1E; and Supplementary Fig. S1A, C). In addition, stem width and circumference were also lower for Nipponbare and IR 64 compared to all selected wild species (Supplementary Fig. S1C, D). Thus, plant architectural traits, such as height, branching, and internode features, as well as leaf traits attributed to high biomass of the selected wild rice. Consistent with this, the selected wild rice also showed higher fresh and dry weight compared to the cultivated varieties. (Fig. 1F, G).

We, then, quantified seed traits of the selected cultivated and wild rice. The cultivated rice varieties produced higher numbers of spikelets per panicle as well as higher seed weight

compared to the wild relatives (Fig. 1H, I). Higher seed weight of cultivated rice varieties resulted due to longer and wider grains compared to the wild species (Supplementary Fig. S1E, F). Principal Component analysis for all the quantified biomass and seed traits showed a clear separation of wild and cultivated genotypes (Supplementary Fig. S2). Taken together, these results confirmed the higher biomass accumulation at the expanse of grain yield in the selected wild rice compared to the cultivated varieties. Wild rice *O. australiensis* had higher phenotypic value for all the traits contributing to biomass, including tiller number, compared to Nipponbare. Therefore, we investigated the source strength, sink features, sucrose translocation as well as the fate of the photosynthates towards biomass and yield using representative wild rice and cultivated variety, *O. sativa* cv. Nipponbare and *O. australiensis*, respectively.

Comparison of source efficiency and photosynthates utilization

In order to compare source efficiency of the two selected accessions, we quantified leaf photosynthesis rate and related physiological traits as well as soluble sugars/non-structural carbohydrates (NSCs) content in leaves of the two accessions. A significant difference in the leaf photosynthesis rate A was observed between the two species at booting, and milk- and dough-stage of grain filling. Both the species showed increasing trend of leaf photosynthesis from booting to milk stage, followed by reduction at the dough stage of grain filling (Supplementary Fig. S3A). The higher photosynthesis in *O. australiensis* at the three stages were also associated with higher g_sH_2O and Ci compared to Nipponbare (Supplementary Fig. S3B, C). Next, we measured the different soluble sugars/NSCs, such as sucrose, fructose, glucose and starch, at the time of photosynthesis quantification in leaves at the milk stage of grain filling to evaluate the levels of primary carbon metabolites, the major outcome of photosynthesis. The abundance of all four NSCs were significantly higher in the leaves of *O. australiensis* than Nipponbare (Fig. 2A, B). Interestingly, *O. australiensis* also accumulated higher levels of starch in leaves compared to Nipponbare, suggesting abundance of residual sugar in the wild rice that is converted to starch. In contrast, the starch content was found to be significantly lower in *O. australiensis* grains compared to Nipponbare, despite starch being the major reserve product of seeds (Fig. 2C). Higher starch and soluble sugar content in *O. australiensis* leaf and lower starch levels in seed suggested possible bottlenecks in the mobilization of photosynthates to grains in the wild rice.

We, then, quantified end of the day (EOD) and end of the night (EON) carbon status in the leaves of the two accessions. Glucose and fructose level were significantly lower in *O.*

australiensis than *O. sativa* cv. Nipponbare at the EOD and EON (Supplementary Fig. S4A).

Sucrose content was marginally, but significantly, higher in *O. australiensis* compared to *O.*

sativa cv. Nipponbare at the EOD as well as EON. At the EOD, *O. australiensis* accumulated

more starch in leaves than *O. sativa* cv. Nipponbare (Supplementary Fig. S4B). However, *O.*

australiensis displayed a significantly lower leaf starch level than *O. sativa* cv. Nipponbare at

the EON, suggesting efficient utilization of stored starch in *O. australiensis* during the night

(Supplementary Fig. S4B). This is in contrast to lower starch content in the grains of *O.*

australiensis as explained earlier (Fig. 2C). These results clearly indicated the efficient

utilization of the photosynthates in the wild rice, however not towards grain filling,

suggesting the possibility of the alternative sink and alternative fate of photosynthates.

Phloem loading and export of sucrose from leaves

Since sucrose is the major transportable form of photosynthates from source leaves to

sink organs, we investigated the sucrose transport differences between the two accessions.

We performed [^{14}C] labelled sucrose assay in leaf discs of *O. sativa* cv. Nipponbare and *O.*

australiensis. The higher amount of ^{14}C per unit leaf disc area was detected *O. australiensis*

(5598.83 CPM per cm^2) compared to that of in Nipponbare (3446.66 CPM per cm^2) (Fig.

2D). We, then, quantified sucrose in the phloem sap of the two accessions. Phloem sap of *O.*

australiensis had four times higher sucrose content than *O. sativa* cv. Nipponbare (Fig. 2E).

Together, ^{14}C-assay and analysis of phloem sap showed the efficient phloem loading and

sucrose export from the leaf in the wild rice *O. australiensis* compared to Nipponbare.

Expression pattern of genes encoding sucrose transporters

We examined the contribution of sucrose transporters for differences in phloem

loading and sucrose export from leaves. Transcript levels of genes encoding clade III SWEET

transporters, as well as SUT transporters were quantified. Expression analysis of *OsSWEET*s

using publicly available datasets showed preferentially source-specific expression of

OsSWEET13 (Supplementary Fig. S5A). *OsSWEET14* also followed a similar source-specific

expression pattern, but at a relatively lower expression level than *OsSWEET13*. *OsSWEET15*

and *OsSWEET11* showed preferentially sink (panicle and endosperm) specific expression

patterns. The expression levels of *OsSWEET12* was observed to be very low in all the tissues.

Consistent with this, quantitative Real Time-PCR results showed higher transcript levels of

OsSWEET13 in flag leaf and stem; *OsSWEET11* in the spikelet; and *OsSWEET15* in the stem,

spikelet, and flag leaf of Nipponbare (Fig. 3A-D). Interestingly, the transcript level of

OsSWEET13, encoding preferentially source-specific SWEET transporter, was significantly
higher in flag leaf and stem of *O. australiensis* compared to Nipponbare (Fig. 3A, B). *In*
silico expression analysis showed higher expression of *OsSUT1* and *OsSUT2* in different
tissue compared to other *OsSUT*s (Supplementary Fig. S5B). qRT-PCR analysis confirmed
the generally higher expression of *OsSUT1* and *OsSUT2* in different tissues of Nipponbare
(Fig. 3E-H). We detected higher transcript levels of *OsSUT1* in the flag leaf, and *OsSUT2* in
the stem of *O. australiensis* compared to Nipponbare (Fig. 3E, F). The expression pattern of
*OsSWEET*s and *OsSUT*s suggested the potential involvement of selected SWEET and SUT
transporters, such as OsSWEET13, OsSUT1, and OsSUT2, in efficient phloem loading and
sucrose export from leaf to stem in the wild rice.

Despite higher photosynthetic efficiency and efficient sucrose transport from source
leaves, *O. australiensis* produced smaller and lighter grains with less starch content compared
to the cultivated variety Nipponbare. Therefore, we also examined the transcript levels of
different sucrose transporters in developing spikelets. The expression levels of all the
*OsSWEET*s were significantly lower in spikelets of *O. australiensis* compared to Nipponbare,
with almost no expression of preferentially sink-specific *OsSWEET15* and *OsSWEET11* (Fig.
3D). In addition, the transcript levels of *OsSUT1* was significantly lower in developing
spikelet of *O. australiensis* than Nipponbare (Fig. 3H). Thus, reduced grain filling in *O.*
australiensis could potentially be associated with reduced expression of *OsSWEET15*,
OsSWEET11, and *OsSUT1*.

Differences in vascular features associated with sucrose transport

Vascular features are also key to photoassimilate partitioning from source to sink
tissues. Therefore, we quantified the vein number and vein width in the fully matured flag
leaves of the two accessions. *O. australiensis* leaves had a greater number of wider veins
compared to Nipponbare (Fig. 4A, B). Moreover, *O. australiensis* also exhibited higher vein
density with lesser interveinal distance compared to Nipponbare (Supplementary Fig. S6A,
B). We, then, checked the organization of vascular bundles at the panicle base of the two
accessions, as panicle base is the site of attachment of spikelets to the main plant (Zhang *et*
al., 2002; Zhai *et al.,* 2018). *O. australiensis* had a fewer number of vascular bundles with the
reduced area than Nipponbare at the panicle base (Fig. 4C, D). We also quantified the
vascular features in the stem of the two accessions (Fig. 4E, F). Interestingly, we observed
~1.7 times larger vascular bundles in the *O. australiensis* stem compared to Nipponbare.

Further, *O. australiensis* stem also had approximately twice the number of vascular bundles
than Nipponbare stem due to wider stem (Fig. 4F). Taken together, the sucrose might not be
efficiently mobilized to panicle in *O. australiensis*, likely due to defects in the vascular
bundle at the panicle base and minimal expression of *OsSWEET*s in developing spikelets,
strengthening the hypothesis of alternative utilization of photosynthates in the *O.*
australiensis stem towards higher biomass.

Starch accumulation and expression levels of starch-biosynthesis genes in stem

The leaf photosynthesis, efficient phloem loading in leaves, vascular features as well
as the expression pattern of sucrose transporter genes indicated stem to be a major sink organ
utilizing sucrose received from leaves in *O. australiensis*. Therefore, we quantified the
content of NSCs in stems of the two species at milk-stage of grain filling (Fig. 5A, B).
Significantly lower amounts of glucose, fructose and sucrose were detected in the stem of *O.*
australiensis as compared to Nipponbare (Fig. 5A). This was in contrast to more amount of
sucrose being exported from leaves of *O. australiensis*. Therefore, we suspected that *O.*
australiensis might reserve a high amount of starch in their stem. To our surprise, a negligible
amount of starch was present in the stem of *O. australiensis* compared to Nipponbare (Fig.
5B). Starch content was, further, found to be very low in the *O. australiensis* stems at
different stages and in different internodes (Fig. S7A, B). We, then, checked the expression
of key genes involved in starch biosynthesis, *OsAPL3*, *OsAPS1* and *OsSSIIb*. Expectedly,
very high expressions of these genes were detected in internode and stem of the cultivated
variety Nipponbare by both *in silico* expression analysis as well as qRT-PCR analysis (Fig.
5C; Supplementary Fig. S8A). Consistent with very low starch levels in *O. australiensis* stem
and internode, the transcript abundance of genes encoding two key enzymes, *OsAPL3* and
OsAPS1, was drastically less in *O. australiensis* compared to Nipponbare (Fig. 5C). Lower
amount of soluble non-structural carbohydrates, in particular remarkably low starch content,
despite efficient sucrose transport from leaves to stem, confirmed an alternative fate to
photosynthates in *O. australiensis* stem.

Expression levels of sucrose metabolism genes, Sucrose Synthase enzyme activity, and
structural carbohydrate levels in stem

Invertases (INV) and Sucrose Synthase (SUS) are the key enzymes for degradation of
sucrose. Expression of genes encoding Invertases have been shown to be strongly correlated
with starch synthesis (Ruan, 2014; Bahaji *et al.*, 2014). Therefore, we checked the expression

pattern of different genes encoding cell wall Invertase (*OsCIN1*), cytoplasmic Invertase
(*OsNIN7*), and vacuolar Invertase (*OsINV2*) (Supplementary Fig. S9). The expression of
OsCIN1 was drastically lower in *O. australiensis* indicating limited breakdown of sucrose
that remained in stem apoplast during the transport.

Among all the *OsSUS*s, *OsSUS1* was shown to be highly expressed in internodes
(Hirose *et al.*, 2008). *In silico* expression analysis also showed high expression of *OsSUS1* in
internodes (Fig. S8A). Therefore, we checked the intracellular localization of OsSUS1 and
expression levels of the corresponding gene. The OsSUS1-YFP signal overlapped with the
plasma membrane-localized marker (PM-mCherry), confirming OsSUS1 localization in
plasma membrane (Supplementary Fig. S10). *OsSUS1* was expressed four times higher in *O.*
australiensis stem than Nipponbare, indicating a possibility of cellulose accumulation in *O.*
australiensis stem (Fig. 6A). Since SUS catalyzes a reversible cleavage of sucrose, it has a
synthesis activity promoting sucrose synthesis and a cleavage activity promoting sucrose
degradation. The enzyme activity assay revealed higher OsSUS1 synthesis activity in
Nipponbare, while higher cleavage activity in *O. australiensis* (Fig. 6B, C). Lesser
accumulation of starch, more transcript levels of *OsSUS1*, and higher cleavage activity of
OsSUS1 in *O. australiensis* stem compared to Nipponbare along with localization of OsSUS1
to plasma membrane prompted us to check the cellulose content in the stems of the two
accessions. Cellulose synthase genes are usually expressed at the highest levels in rice
internodes compared to other tissues as evident from *in silico* expression analysis (Fig. S8B).
Interestingly, transcript levels of *OsCES4*, *OsCES7*, and *OsCES9*, key cellulose synthesis
genes, were significantly higher in the stems of *O. australiensis* than in Nipponbare (Fig.
6D). Calcofluor-white staining along with cellulose quantification confirmed more cellulose
deposition in *O. australiensis* stem compared to Nipponbare (Fig. 6E, F). Like cellulose,
hemicellulose content was also significantly higher in *O. australiensis* stem compared to
Nipponbare (Fig. 6F). Taken together, higher expression of *OsSUS1* and higher cleavage
activity of OsSUS1 coupled with higher expression of Cellulose Synthase genes, at least in
part, led to efficient utilization of photosynthates in the *O. australiensis* stem.

Discussion

We investigated the differences in source-sink dynamics between a cultivated rice
variety *O. sativa* cv. Nipponbare, which is optimized for high grain yield, and a wild relative
of rice *O. australiensis*, which accumulates high biomass with poor grain yield. *O.*
australiensis had higher source strength, as evident by consistently higher leaf photosynthesis

rate compared to Nipponbare, at least in part due to vascular features of the leaf

(Supplementary Fig. S3A and Fig.4A, B). High leaf photosynthesis, usually, results in high

biomass or yield depending upon preferential sink tissues, provided efficient transport of

photoassimilates from leaves (Burnett *et al.*, 2016; Fabre *et al.*, 2020; Fernie *et al.*, 2020).

Limitations in sucrose export accumulate sugars in leaves, thereby inhibiting leaf

photosynthesis. Therefore, an efficient sucrose export system from leaf would be warranted

for the realization of higher source strength of the wild rice to high biomass and/or yield. [14]C

sucrose uptake assay, as well as sucrose content in phloem sap, confirmed the efficient

sucrose export from wild rice leaves (Fig. 2D, E). Larger vascular bundles and fewer number

of mesophyll cells between two consecutive veins potentially mediated efficient sucrose

export from leaves in wild rice, as reported in different plant system (Qi *et al.*, 2008; Fujita *et*

al., 2013; Mathan *et al.*, 2016). In addition, higher expression of genes encoding sucrose

transporters, OsSWEET13 and OsSUT1, would also facilitate the efficient phloem loading in

the wild rice (Fig. 3). High expression of *SbSWEET8-1* of sorghum, a close homolog of

OsSWEET13, in leaf and its function in phloem loading supports our idea of the important

role of *OsSWEET13* for phloem loading (Mizuno *et al.,* 2016). Similarly, *OsSUT1* not only

functions for enhancing phloem loading for sucrose transport but also for retrieval of sucrose

from the apoplasm along the transport pathway (Scofield *et al.*, 2007). Higher amount of

soluble sugars in wild rice leaves, despite having higher sucrose export from leaves,

compared to the cultivated variety, further supported the better source strength of the wild

rice (Fig. 2 A, B).

The better grain filling and higher grain yield of cultivated rice, despite of the

relatively lower source strength compared to *O. australiensis*, could be explained by optimal

vascular features and desirable expression sucrose transporter genes at the panicle base and

spikelets of cultivated rice (Fig. 3, 4). Panicle architecture, a key determinant of rice grain

yield, is reported to be shaped by the vascular pattern (Sasahara *et al.*, 1999; Terao *et al.*,

2010). Reduced expression of *OsSWEET13* and *OsSWEET15* in the developing spikelet, and

of *OsSUT1* at the panicle base and developing spikelet, along with reduced supply of smaller

vascular bundles at the panicle base limited mobilization of photosynthates to grains in the

wild rice compared to the cultivated. In agreement with this, *ossweet11 ossweet15* double

mutant has been reported to show smaller seed size (Yang *et al.*, 2018). Similarly, RNA

antisense lines of *OsSUT1* showed reduced grain filling and grain weight (Scofield *et al.*,

2007). Increased supply of larger vascular bundles to rice rachis has been shown to promote

more grains per panicle (Zhang *et al.*, 2002; Terao *et al.*, 2010; Zhai *et al.*, 2018). Altogether,

higher source strength and limited grain sink strength in the wild rice, attributed to vascular

features and sucrose transporter functions, indicated preferential utilization of photosynthates

in stem and internodes, contributing to source-sink relationship differences between the

selected cultivated and wild rice. In conjunction with this, larger vascular bundles would

facilitate efficient sucrose transport and unloading in the wild rice stem. Such vascular

features in the stem would, in turn, also provide mechanical strength to support the high

biomass of the wild rice (Aohara *et al.*, 2009). Our expression results indicated the potential

involvement of OsSWEET13 and OsSUT2 in unloading larger amount of sucrose being

received in the wild stem.

There is a competition among the sink tissues for utilization of photosynthates

(Patrick *et al.*, 2013; Durand *et al.*, 2018). According to the EcoMeristem model, the final

plant architecture is an outcome of competition for resources among different plant parts that

depends on photoassimilate-partitioning patterns (Luquet *et al.*, 2006). More number of

longer internodes and thicker stem with larger leaves, driving high biomass in the wild rice

compared to the cultivated variety, suggested higher utilization of photosynthates for

vegetative growth. The higher biomass and larger organ size of wild rice would require more

photosynthates for general respiration and maintenance. However, this would likely not

explain the utilization of photosynthates in total as wild rice showed remarkably higher

amount of sucrose in phloem sap compared to the cultivated variety. Altogether, wild rice

efficiently utilized photoassimilates reaching to stem towards larger vegetative organ size and

number, cell-wall thickening as well as the overall maintenance of higher biomass as

compared to the cultivated rice variety.

The preferential utilization of photosynthates in the stem/internodes of wild rice in

contrast to grains in the cultivated rice suggested differences in the photosynthates

metabolism in the stem of the two species. Cultivated rice variety clearly accumulated higher

starch content in the stem, which is mobilized to panicle during grain filling (Fig. 5B).

OsSWEET11 and *OsSWEET15* have been suggested to be important for remobilization of

carbon reserve from stem to grain, and OsSUT1 in the retrieval of sucrose from apoplasmic

space to stem for conversion to transitory starch (Wang *et al.*, 2020). Indeed, Nipponbare

stem showed higher expression of *OsSWEET11*, *OsSWEET15*, and *OsSUT1* compared to the

wild rice along with desirable vascular features at the panicle base. High expression of starch

biosynthesis genes, further, contributed to the roles of the stem as an effective source in the

cultivated variety at the grain filling stage (Fig. 7). The differences in the stem starch content

between the two genotypes projected the stored transitory starch in the stem as a key for source-sink dynamics favoring high grain yield.

O. australiensis stem showed higher expression of *OsSUS1* and higher cleavage activity OsSUS producing higher amount of UDP-glucose and fructose compared to cultivated rice. UDP-glucose has been shown to serve as the prime source for cellulose synthesis (Smith *et al*., 2012). UDP-glucose can possibly also be converted to starch (Asano *et al*., 2002; Koch, 2004). However, the possibility is minimized in the wild rice due to very limited expression of *OsAPL3* and *OsAPS1*. Thus, most of the SUS-generated UDP-glucose was converted to cellulose, supported by high expression of cellulose synthase genes, *OsCES4*, *OsCES7*, and *OsCES9* in the wild stem (Fig. 6D-F). OsSUS1 has been reported to be located in SE-CC complexes of phloem, and localization of SUS in phloem is important for cellulose synthesis (Smith *et al*., 2012; Regmi *et al*., 2016). Plasma membrane localization of OsSUS1 in this study together with earlier reported association of cellulose synthase complexes to plasma membrane highlighted the key importance of OsSUS1 for cellulose synthesis in rice (Supplementary Fig. S10, Lei *et al*., 2012). In addition, SUS functions in companion cells of phloem has been suggested to enhance the sucrose unloading at sink tissues, further helping in unloading photosynthates in wild rice stem (Nolte and Koch, 1993; Stein and Granot, 2019, Yao *et al*., 2020). The potential role of SUS in cellulose synthesis, as well as in secondary cell wall thickening, has been investigated in the past (Coleman *et al*., 2009; Baroja-Fernández *et al*., 2012; Wei *et al*., 2015). Consistent with the higher expression of *OsSUS1* in wild rice, overexpression of *SUS* has been shown to result in increased vegetative growth rate, plant height and biomass in different plant species (Coleman *et al*., 2006; Nguyen *et al*., 2016; Stein and Granot, 2019). Contrary to the wild rice, higher synthesis activity of OsSUS in Nipponbare stem would facilitate the formation of sucrose from the stored starch at the heading stage to be mobilized for grain filling. Thus, differential functions in OsSUS1 between wild and cultivated rice would, at least in part, explain the differential fate of photosynthates in the stem (Fig. 7).

In summary, differences in vascular features and sucrose transporter functions led to a differential source-sink relationship between wild rice *O. australiensis* and cultivated variety *O. sativa* cv. *Nipponbare*. Wild rice showed source-sink dynamics favoring high biomass through accumulation of structural carbohydrates, mediated by higher cleavage activity of SUS enzyme together with higher expression of Cellulose Synthase genes (Fig. 7). In contrast, source-sink dynamics favored higher grain yield in Nipponbare via accumulation of transitory starch in the stem, to be mobilized to panicles with the onset of grain filling. Taken

together, vascular features and sucrose transporter functions along with transitory starch
storage mechanism and SUS enzyme activity can potentially be targeted for source-sink
dynamics favoring either biomass accumulation in fodder crops or higher grain yield in cereal
crops.

Supplementary data

Fig. S1. Biomass and yield-related traits of the selected cultivated and wild rice genotypes.

Fig. S2. Principal component analysis of biomass and seed traits of the selected cultivated
and wild rice species.

Fig. S3. Leaf photosynthesis and related physiological differences between the cultivated rice
O. sativa cv. Nipponbare and wild rice *O. australiensis*.

Fig. S4. Sugar content in leaves of cultivated rice *O. sativa* cv. Nipponbare and wild rice *O.*
australiensis at End of Day (EOD) and End of Night (EON).

Fig. S5. *In silico* organ-specific expression analysis of rice genes encoding SWEET and SUT
transporters.

Fig. S6. Quantification of leaf interveinal distance and vein density for cultivated rice *O.*
sativa cv. Nipponbare and wild rice *O. australiensis*.

Fig. S7. Total starch content in stem and internodes of cultivated rice *O. sativa* cv.
Nipponbare and wild rice *O. australiensis*.

Fig. S8. *In silico* expression pattern of key genes involved in sugar metabolism in rice.

Fig. S9. Transcript levels of genes encoding cell wall invertase (*OsCIN1*), cytoplasmic
invertase (*OsNIN7*), and vacuolar invertase (*OsINV2*).

Fig. S10. Localization of OsSUS1-YFP in plasma membrane.

Table S1. List of primer pairs used in the study

Acknowledgements

This work was supported by the Innovative Young Biotechnologist Award

(BT/09/IYBA/2015/01) and Ramalingaswamy Re-entry Fellowship

(BT/RLF/reentry/05/2013) to AR from the Department of Biotechnology, Ministry of

Science and Technology, India. JM and AS acknowledge their CSIR-JRF and SERB-NPDF

fellowships, respectively. We also acknowledge Central Instrument Facility, NIPGR;

Advanced Instrumentation Research Facility, JNU; Confocal Microscopy Facility, NIPGR;

and DBT-eLibrary Consortium (DeLCON) for providing access to e-resources.

Author Contributions

JM and AR conceptualized the study and designed experiments. JM performed experiments.

JM and AS analysed data. JM, AS and AR wrote the manuscript. All the authors have read

and edited the final manuscript.

Data availability statement

Data sharing is not applicable to this article as all created data is already contained within this

article or in the supplementary material.

References

Ambavaram, M.M.R., Krishnan, A., Trijatmiko, K.R. and Pereira, A. 2011. Coordinated activation of cellulose and repression of lignin biosynthesis pathways in rice. Plant Physiology **155,** 916–931.

Asano, T., Kunieda, N., Omura, Y., et al. 2002. Rice SPK, a calmodulin-like domain protein kinase, is required for storage product accumulation during seed development: Phosphorylation of sucrose synthase is a possible factor. Plant Cell **14,** 619–628.

Aohara T, Kotake T, Kaneko Y, Takatsuji H, Tsumuraya Y, Kawasaki S. 2009. Rice brittle culm 5 (brittle node) is involved in secondary cell wall formation in the sclerenchyma tissue of nodes. Plant and Cell Physiology **50**, 1886–1897.

Bahaji A, Li J, Sánchez-López ÁM, *et al.* 2014. Starch biosynthesis, its regulation and biotechnological approaches to improve crop yields. Biotechnology Advances **32**, 87–106.

Baroja-Fernández E, Muñoz FJ, Li J, Bahaji A, Almagro G, Montero M, Etxeberria E, Hidalgo M, Sesma MT, Pozueta-Romero J. 2012. Sucrose synthase activity in the sus1/sus2/sus3/sus4 Arabidopsis mutant is sufficient to support normal cellulose and starch production. Proceedings of the National Academy of Sciences of the United States of America **109**, 321–326.

Braun DM. 2012. Sweet! The pathway is complete. Science **335**, 173–174.

Burnett AC, Rogers A, Rees M, Osborne CP. 2016. Carbon source–sink limitations differ between two species with contrasting growth strategies. Plant Cell and Environment **39**, 2460–2472.

Chen LQ, Qu XQ, Hou BH, Sosso D, Osorio S, Fernie AR, Frommer WB. 2012. Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. Science **335**, 207–211.

Cho J Il, Kim HB, Kim CY, Hahn TR, Jeon JS. 2011. Identification and characterization of the duplicate rice sucrose synthase genes OsSUS5 and OsSUS7 which are associated with the plasma membrane. Molecules and Cells **31**, 553–561.

Coleman HD, Ellis DD, Gilbert M, Mansfield SD. 2006. Up-regulation of sucrose synthase and UDP-glucose pyrophosphorylase impacts plant growth and metabolism. Plant Biotechnology Journal **4**, 87–101.

Coleman, H.D., Yan, J. and Mansfield, S.D. 2009. Sucrose synthase affects carbon partitioning to increase cellulose production and altered cell wall ultrastructure. Proceedings of the National Academy of Sciences of the United States of America **106**, 13118–13123.

Durand M, Mainson D, Porcheron B, Maurousset L, Lemoine R, Pourtau N. 2018. Carbon source–sink relationship in Arabidopsis thaliana: the role of sucrose transporters. Planta **247**, 587–611.

Earley KW, Haag JR, Pontes O, Opper K, Juehne T, Song K, Pikaard CS. 2006. Gateway-compatible vectors for plant functional genomics and proteomics. The Plant Journal **45**, 616–629.

Fabre D, Dingkuhn M, Yin X, Clément-Vidal A, Roques S, Soutiras A, Luquet D. 2020. Genotypic variation in source and sink traits affects the response of photosynthesis and growth to elevated atmospheric CO2. Plant Cell and Environment **43**, 579–593.

Fan C, Feng S, Huang J, *et al.* 2017. AtCesA8-driven OsSUS3 expression leads to largely enhanced biomass saccharification and lodging resistance by distinctively altering lignocellulose features in rice. Biotechnology for Biofuels **10**, 1–12.

Fernie AR, Bachem CWB, Helariutta Y, *et al.* 2020. Metabolic Aspects of Source – Sink Interactions. Nature Plants **6**.

Fujita D, Trijatmiko KR, Tagle AG, *et al.* 2013. NAL1 allele from a rice landrace greatly increases yield in modern indica cultivars. Proceedings of the National Academy of Sciences of the United States of America **110**, 20431–20436.

Guevara DR, El-Kereamy A, Yaish MW, Mei-Bi Y, Rothstein SJ. 2014. Functional characterization of the rice UDP-glucose 4-epimerase 1, OsUGE1: A potential role in cell wall carbohydrate partitioning during limiting nitrogen conditions. PLoS ONE **9**.

Haigler CH, Ivanova-Datcheva M, Hogan PS, Salnikov V V., Hwang S, Martin K, Delmer DP. 2001. Carbon partitioning to cellulose synthesis. Plant Molecular Biology **47**,

29–51.

Hirose T, Scofield GN, Terao T. 2008. An expression analysis profile for the entire sucrose synthase gene family in rice. Plant Science **174**, 534–543.

Julius BT, Leach KA, Tran TM, Mertz RA, Braun DM. 2017. Sugar transporters in plants: New insights and discoveries. Plant and Cell Physiology **58**, 1442–1460.

King, R.W. and Zeevaart, J.A.D. 1974. Enhancement of Phloem Exudation from Cut Petioles by Chelating Agents. Plant Physiology **53**, 96–103.

Koch, K. 2004. Sucrose metabolism: Regulatory mechanisms and pivotal roles in sugar sensing and plant development. Current Opinion in Plant Biology **7**, 235–246.

Lei, L., Li, S., and Gu, Y. 2012. Cellulose synthase complexes: composition and regulation. Frontiers in plant science **3**, 75.

Lisec J, Schauer N, Kopka J, Willmitzer L, Fernie AR. 2006. Gas chromatography mass spectrometry-based metabolite profiling in plants. Nature Protocols **1**, 387–396.

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2-$\Delta\Delta$CT method. Methods **25**, 402–408.

Luquet D, Dingkuhn M, Kim H, Tambour L, Clement-Vidal A. 2006. EcoMeristem, a model of morphogenesis and competition among sinks in rice. 1. Concept, validation and sensitivity analysis. Functional Plant Biology **33**, 309–323.

Mathan J, Bhattacharya J, Ranjan A. 2016. Enhancing crop yield by optimizing plant developmental features. Development (Cambridge) **143**, 3283-3294.

Mitra, P.P. and Loqué, D. 2014. Histochemical staining of Arabidopsis thaliana secondary cell wall elements. Journal of Visualized Experiments **13**, 51381.

Mizuno, H., Kasuga, S. and Kawahigashi, H. 2016. The sorghum *SWEET* gene family: stem sucrose accumulation as revealed through transcriptome profiling. Biotechnology for Biofuels **9,** 127.

Mund, N.K., Dash, D., Barik, C.R., Goud, V. V., Sahoo, L., Mishra, P. and Nayak, N.R. 2016. Chemical composition, pretreatments and saccharification of Senna siamea (Lam.) H.S.

Irwin & Barneby: An efficient biomass producing tree legume. Bioresource Technology **207**, 205–212.

Nelson, B.K., Cai, X. and Nebenführ, A. 2007. A multicolored set of in vivo organelle markers for co-localization studies in Arabidopsis and other plants. Plant Journal **51**, 1126–1136.

Nguyen QA, Luan S, Wi SG, Bae H, Lee DS, Bae HJ. 2016. Pronounced phenotypic changes in transgenic tobacco plants overexpressing sucrose synthase may reveal a novel sugar signaling pathway. Frontiers in Plant Science **6**, 1216.

Nolte KD, Koch KE. 1993. Companion-cell specific localization of sucrose synthase in zones of phloem loading and unloading. Plant Physiology **101**, 899–905.

Osorio S, Ruan YL, Fernie AR. 2014. An update on source-to-sink carbon partitioning in tomato. Frontiers in Plant Science **5**, 1–11.

Patrick JW, Botha FC, Birch RG. 2013. Metabolic engineering of sugars and simple sugar derivatives in plants. Plant Biotechnology Journal **11**, 142–156.

Qazi, H.A., Paranjpe, S. and Bhargava, S. 2012. Stem sugar accumulation in sweet sorghum - Activity and expression of sucrose metabolizing enzymes and sucrose transporters. Journal of Plant Physiology **169**, 605–613.

Qi J, Qian Q, Bu Q, *et al.* 2008. Mutation of the rice narrow leaf1 gene, which encodes a novel protein, affects vein patterning and polar auxin transport. Plant Physiology **147**, 1947–1959.

Regmi KC, Zhang S, Gaxiola RA. 2016. Apoplasmic loading in the rice phloem supported by the presence of sucrose synthase and plasma membrane-localized proton pyrophosphatase. Annals of Botany **117**, 257–268.

Ruan Y-L. 2014. Sucrose Metabolism: Gateway to Diverse Carbon Use and Sugar Signaling. Annual Review of Plant Biology **65**, 33–67.

Sack L, Scoffoni C. 2013. Leaf venation : structure , function , development , evolution , ecology and applications in the past, present and future. New Phytologist **4**, 983–1000.

Sanchez, P.L., Wing, R.A. and Brar, D.S. 2013. The wild relative of rice: genomes and genomics. In *Genetics and genomics of rice* (Zhang, Q. and Wing, R.A., eds). Springer, pp. 10–25.

Sasahara H, Fukuta Y, Fukuyama T. 1999. Mapping of QTLs for Vascular Bundle System and Spike Morphology in Rice, Oryza satva L. Breeding Science **49**, 75–81.

Scofield GN, Hirose T, Aoki N, Furbank RT. 2007. Involvement of the sucrose transporter, OsSUT1, in the long-distance pathway for assimilate transport in rice. Journal of Experimental Botany **58**, 3155–3169.

Scofield GN, Hirose T, Gaudron JA, Upadhyaya NM, Ohsugi R, Furbank RT. 2002. Antisense suppression of the rice sucrose transporter gene, OsSUT1, leads to impaired grain filling and germination but does not affect photosynthesis. Functional Plant Biology **29**, 815–826.

Smith AM, Kruger NJ, Lunn JE. 2012. Source of sugar nucleotides for starch and cellulose synthesis. Proceedings of the National Academy of Sciences of the United States of America **109**, E776–E776.

Smith AM, Stitt M. 2007. Coordination of carbon supply and plant growth. Plant, Cell and Environment **30**, 1126–1149.

Smith, M.R., Rao, I.M. and Merchant, A. 2018. Source-sink relationships in crop plants and their influence on yield development and nutritional quality. Frontiers in Plant Science **871**, 1–10.

Stein O, Granot D. 2019. An overview of sucrose synthases in plants. Frontiers in Plant Science **10**, 1–14.

Tanaka K, Murata K, Yamazaki M, Onosato K, Miyao A, Hirochika H. 2003. Three distinct rice cellulose synthase catalytic subunit genes required for cellulose synthesis in the secondary wall. Plant Physiology **133**, 73–83.

Terao T, Nagata K, Morino K, Hirose T. 2010. A gene controlling the number of primary rachis branches also controls the vascular bundle formation and hence is responsible to increase the harvest index and grain yield in rice. Theoretical and Applied Genetics **120**, 875–

893.

Wang DR, Han R, Wolfrum EJ, McCouch SR. 2017. The buffering capacity of stems: genetic architecture of nonstructural carbohydrates in cultivated Asian rice, Oryza sativa. New Phytologist **215**, 658–671.

Wang G, Li H, Gong Y, Yang J, Yi Y, Zhang J, Ye N. 2020*a*. Expression profile of the carbon reserve remobilization from the source to sink in rice in response to soil drying during grain filling. Food and Energy Security **3**, 1–14.

Wang D, Qin Y, Fang J, Yuan S, Peng L, Zhao J, Li X. 2016*a*. A missense mutation in the zinc finger domain of OsCESA7 deleteriously affects cellulose biosynthesis and plant growth in rice. PLoS ONE **11**, 1–16.

Wei Z, Qu Z, Zhang L, Zhao S, Bi Z, Ji X, Wang X, Wei H. 2015. Overexpression of poplar xylem sucrose synthase in tobacco leads to a thickened cell wall and increased height. PLoS ONE **10**, 1–20.

White, A.C., Rogers, A., Rees, M. and Osborne, C.P. 2016. How can we make plants grow faster? A source-sink perspective on growth rate. Journal of Experimental **67**, 31–45.

Yadav, U., Khadilkar, A., Shaikh, M., Turgeon, R. and Ayre, B. (2017) Quantifying the Capacity of Phloem Loading in Leaf Disks with [^{14}C]Sucrose. Bio-Protocol **7**, 1–12.

Yang J, Luo D, Yang B, Frommer WB, Eom JS. 2018. SWEET11 and 15 as key players in seed filling in rice. New Phytologist **218**, 604–615.

Yao D, Gonzales-Vigil E, Mansfield SD. 2020. Arabidopsis sucrose synthase localization indicates a primary role in sucrose translocation in phloem. Journal of Experimental Botany **71**, 1858–1869.

Zhai L, Zheng T, Wang X, Wang Y, Chen K, Wang S, Wang Y, Xu J, Li Z. 2018. QTL mapping and candidate gene analysis of peduncle vascular bundle related traits in rice by genome-wide association study. Rice **11**, 13.

Zhang ZH, Li P, Wang LX, Tan CJ, Hu ZL, Zhu YG, Zhu LH 2002. Identification of quantitative trait loci (QTLs) for the characters of vascular bundles in peduncle related to indica -japonica differentiation in rice (*Oryza sativa L.*) Euphytica **128**, 279 –284.

Figure legends

Fig. 1. Comparison of biomass and yield-related traits of the selected cultivated and wild rice genotypes.

(A) Shown are the photographs of the selected rice genotypes at the reproductive stage (scale bar = 30 cm).

(B–I) Quantification of plant height (B), total leaves per plant (C), flag leaf width (D), total internode number per main stem (E), fresh weight (F), dry weight (G), spikelet per panicle (H), and 500-seed weight (I) of the selected rice genotypes. Significance of difference among the genotypes was calculated using One-Way ANOVA with Tukey's post-hoc test (n = 10 data point from different plants, $p < 0.05$).

Fig. 2. Sugar accumulation, phloem loading, and sucrose export from a leaf for the cultivated rice *O. sativa* cv. Nipponbare and wild rice *O. australiensis*.

(A) Shown are glucose, fructose and sucrose content in the flag leaves of the cultivated and wild rice at the time of photosynthesis quantification.

(B–C) Starch content in the flag leaves (B), and in the matured seeds (C) of the cultivated and wild rice.

(D) Uptake of [^{14}C] sucrose in the flag leaf discs of the two species expressed as counts per minute per square centimeter of leaf area.

(E) Sucrose quantification in the phloem sap of the two species.

Data represents the mean and standard error of mean (SE), and significance of difference between the genotype was calculated using student t-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Fig. 3. Tissue-specific expression pattern of genes encoding SWEET and SUT transporters in cultivated rice *O. sativa* cv. Nipponbare and wild rice *O. australiensis*.

(A–D) Transcript levels of *SWEET* genes in the flag leaf (A), stem (B) panicle base (C), and spikelet (D) at the milk-stage of grain-filling in the two rice species.

(E–H) Transcript levels of *SUT* genes in the flag leaf (E), stem (F) panicle base (G), and spikelet (H) at the milk-stage of grain-filling in the two rice species.

Data represents the mean and standard error of mean (SE) from three biological replicate per genotype (n = 3). Significance of difference between the genotype was calculated using student t-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Fig. 4. Vascular features in flag leaf, panicle base, and stem of cultivated rice *O. sativa* cv. Nipponbare and wild rice *O. australiensis*.

(A-B) Cross-sections of flag leaves of the two species (A, scale bar = 100 μm), and quantification of minor vein width and the total number of veins (B).

(C-D) Transverse sections at the panicle base (0.5 cm above panicle node) of the two species (C, scale bar = 250 μm), and quantification of number and area of vascular bundles (D).

(E-F) Transverse sections of stems of the two species (E, scale bar = 100 μm), and quantification of number and area of vascular bundles (F).

The peripheral concentric ring of the panicle base and stem is represented as outer ring, and the inner wider circle is represented as inner ring in figures.

Data represents the mean and standard error of mean (SE) from fifteen cross-section images collected from five different plants and significance difference between the genotype was based on student t-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Fig. 5. Quantification of soluble sugars and expression pattern of starch-biosynthesis genes in stems of cultivated rice *O. sativa* cv. Nipponbare and wild rice *O. australiensis* at milk-stage of grain-filling.

(A-B) Glucose, fructose, and sucrose content (A), and total starch content (B) in the stem of the two species ($n = 4$).

(C) Transcript levels of starch-biosynthesis genes, *OsAPL3* (encoding ADP-glucose pyrophosphorylase large subunit), *OsAPS1* (encoding ADP-glucose pyrophosphorylase small subunit), and *OsSSIIB* (encoding Starch synthase), in the stem of the two species.

Data represents the mean and standard error of mean (SE), and significance difference between the genotype was calculated using student t-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Fig. 6. Expression levels of sucrose metabolism genes, Sucrose Synthase enzyme activity, and structural carbohydrates levels in stem tissue of cultivated rice *O. sativa* cv. Nipponbare and wild rice *O. australiensis* at milk-stage of grain-filling.

(A) Transcript levels of *Sucrose Synthase* 1 (*OsSUS1*).

(B-C) OsSUS enzyme activity in synthesis (B) and cleavage (C) direction.

(D) Transcript levels of *Cellulose Synthase* genes, *OsCES4*, *OsCES7*, and *OsCES9*.

(E) Calcofluor-white staining for cellulose deposition in the transverse stem sections of the two species. Red arrowhead shows higher deposition of cellulose in *O. australiensis* stem. Scale bar represents 500 μm.

(F) Quantification of cellulose and hemicellulose in the stems of the two species.

Data in a- d and f represents the mean and standard error of mean (SE), and significance of difference between the genotype was calculated using student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 7. A model explaining the contribution of sucrose transport, Sucrose Synthase activity, starch biosynthesis, and synthesis of structural carbohydrates towards yield and biomass differences between wild rice *O. australiensis* and cultivated rice *O. sativa* cv. Nipponbare. High leaf photosynthesis rate coupled with higher expression of selected *OsSWEET* and *OsSUT* genes lead to efficient export of sucrose from leaf in *O. australiensis* compared to *O. sativa* cv. Nipponbare. Sucrose, after getting unloaded to stem due to functions of *OsSWEET13*, *OsSUT1*, and *OsSUT2*, gets converted to UDP-G more efficiently in the wild rice due to higher expression of *OsSUS1* and more cleavage activity of OsSUS compared to the cultivated rice. Higher expression of *OsCES4, OsCES7,* and *OsCES9*, then, leads to more synthesis of cellulose in the stem of the wild rice. In contrast, higher expression of starch-biosynthesis genes, *AGPase* (small and large subunit), leads to higher starch content in the stem of the cultivated rice. Higher synthesis activity of OsSUS along with more expression of *OsSWEET11* and *OsSWEET15* in the cultivated rice facilitate efficient remobilisation of sucrose from stem to panicles at the grain-filling stage. SUC, sucrose (yellow colour circle); F, fructose; G, glucose; UDP-G, UDP-glucose; ADP-G, ADP-glucose; *INV*, Invertase; cwINV, cell wall Invertase; SUS, Sucrose Synthase.

Fig. 1. Comparison of biomass and yield-related traits of the selected cultivated and wild rice genotypes.
(A) Shown are the photographs of the selected rice genotypes at the reproductive stage (scale bar = 30 cm).
(B–I) Quantification of plant height (B), total leaves per plant (C), flag leaf width (D), total internode number per main stem (E), fresh weight (F), dry weight (G), spikelet per panicle (H), and 500-seed weight (I) of the selected rice genotypes. Significance of difference among the genotypes was calculated using One-Way ANOVA with Tukey's post-hoc test (n = 10 data point from different plants, p < 0.05).

Fig. 2. Sugar accumulation, phloem loading, and sucrose export from a leaf for the cultivated rice *O. sativa* cv. Nipponbare and wild rice *O. australiensis*.

(A) Shown are glucose, fructose and sucrose content in the flag leaves of the cultivated and wild rice at the time of photosynthesis quantification.

(B–C) Starch content in the flag leaves (B), and in the matured seeds (C) of the cultivated and wild rice.

(D) Uptake of [^{14}C] sucrose in the flag leaf discs of the two species expressed as counts per minute per square centimeter of leaf area.

(E) Sucrose quantification in the phloem sap of the two species.

Data represents the mean and standard error of mean (SE), and significance of difference between the genotype was calculated using student t-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Fig. 3. Tissue-specific expression pattern of genes encoding SWEET and SUT transporters in cultivated rice *O. sativa* cv. Nipponbare and wild rice *O. australiensis*.

(A–D) Transcript levels of *SWEET* genes in the flag leaf (A), stem (B) panicle base (C), and spikelet (D) at the milk-stage of grain-filling in the two rice species.

(E–H) Transcript levels of *SUT* genes in the flag leaf (E), stem (F) panicle base (G), and spikelet (H) at the milk-stage of grain-filling in the two rice species.

Data represents the mean and standard error of mean (SE) from three biological replicate per genotype (n = 3). Significance of difference between the genotype was calculated using student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 4. Vascular features in flag leaf, panicle base, and stem of cultivated rice *O. sativa* cv. Nipponbare and wild rice *O. australiensis*.

(A-B) Cross-sections of flag leaves of the two species (A, scale bar = 100 µm), and quantification of minor vein width and the total number of veins (B).

(C-D) Transverse sections at the panicle base (0.5 cm above panicle node) of the two species (C, scale bar = 250 µm), and quantification of number and area of vascular bundles (D).

(E-F) Transverse sections of stems of the two species (E, scale bar = 100 µm), and quantification of number and area of vascular bundles (F).

The peripheral concentric ring of the panicle base and stem is represented as outer ring, and the inner wider circle is represented as inner ring in figures.

Data represents the mean and standard error of mean (SE) from fifteen cross-section images collected from five different plants and significance difference between the genotype was based on student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 5. Quantification of soluble sugars and expression pattern of starch-biosynthesis genes in stems of cultivated rice *O. sativa* cv. Nipponbare and wild rice *O. australiensis* at milk-stage of grain-filling.
(A-B) Glucose, fructose, and sucrose content (A), and total starch content (B) in the stem of the two species (n = 4).
(C) Transcript levels of starch-biosynthesis genes, *OsAPL3* (encoding ADP-glucose pyrophosphorylase large subunit), *OsAPS1* (encoding ADP-glucose pyrophosphorylase small subunit), and *OsSSIIB* (encoding Starch synthase), in the stem of the two species.
Data represents the mean and standard error of mean (SE), and significance difference between the genotype was calculated using student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 6. Expression levels of sucrose metabolism genes, Sucrose Synthase enzyme activity, and structural carbohydrates levels in stem tissue of cultivated rice *O. sativa* cv. Nipponbare and wild rice *O. australiensis* at milk-stage of grain-filling.

(A) Transcript levels of *Sucrose Synthase* 1 (*OsSUS1*).

(B-C) OsSUS enzyme activity in synthesis (B) and cleavage (C) direction.

(D) Transcript levels of *Cellulose Synthase* genes, *OsCES4*, *OsCES7*, and *OsCES9*.

(E) Calcofluor-white staining for cellulose deposition in the transverse stem sections of the two species. Red arrowhead shows higher deposition of cellulose in *O. australiensis* stem. Scale bar represents 500 μm.

(F) Quantification of cellulose and hemicellulose in the stems of the two species.

Data in a- d and f represents the mean and standard error of mean (SE), and significance of difference between the genotype was calculated using student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 7. A model explaining the contribution of sucrose transport, Sucrose Synthase activity, starch biosynthesis, and synthesis of structural carbohydrates towards yield and biomass differences between wild rice *O. australiensis* and cultivated rice *O. sativa* cv. Nipponbare.

High leaf photosynthesis rate coupled with higher expression of selected *OsSWEET* and *OsSUT* genes lead to efficient export of sucrose from leaf in *O. australiensis* compared to *O. sativa* cv. Nipponbare. Sucrose, after getting unloaded to stem due to functions of *OsSWEET13*, *OsSUT1*, and *OsSUT2*, gets converted to UDP-G more efficiently in the wild rice due to higher expression of *OsSUS1* and more cleavage activity of OsSUS compared to the cultivated rice. Higher expression of *OsCES4, OsCES7,*and *OsCES9*, then, leads to more synthesis of cellulose in the stem of the wild rice. In contrast, higher expression of starch-biosynthesis genes, *AGPase* (small and large subunit), leads to higher starch content in the stem of the cultivated rice. Higher synthesis activity of OsSUS along with more expression of *OsSWEET11* and *OsSWEET15* in the cultivated rice facilitate efficient remobilisation of sucrose from stem to panicles at the grain-filling stage. SUC, sucrose (yellow colour circle); F, fructose; G, glucose; UDP-G, UDP-glucose; ADP-G, ADP-glucose; *INV*, Invertase; cwINV, cell wall Invertase; SUS, Sucrose Synthase.