1   **Title: Sucrose transport and metabolism control carbon partitioning between stem and**
2   **grain in rice**
3
4
5   **Running Title:** Source-sink relationship in cultivated and wild rice
6
7

8   Jyotirmaya Mathan, Anuradha Singh, Aashish Ranjan[*]
9   National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi-110067,
10 India.
11
12 **Corresponding Author*
13 **e-mail:** aranjan@nipgr.ac.in
14 Phone no. +91-11-26735117
15
16 Authors' email
17 Jyotirmaya Mathan: jyotirmaya@nipgr.ac.in
18 Anuradha Singh: annusingh1206@gmail.com

19
20 Date of submission: 19[th] October 2020
21
22
23 Number of figures: 7 figures
24
25
26 The word count: 6,121
27
28
29 Supplementary data: 13 supplementary figures and 1 supplementary table

Title: Sucrose transport and metabolism control carbon partitioning between stem and grain in rice

Running Title: Source-sink relationship in cultivated and wild rice

Highlight

Vascular features, sucrose transport, and sugar metabolic enzyme activity contribute to the differential source-sink relationship between the selected cultivated and wild rice that differ in biomass and grain yield.

Abstract

The source-sink relationship is key to overall crop performance. Detailed understanding of the factors that determine source-sink dynamics is imperative for the balance of biomass and grain yield in crop plants. We investigated the differences in the source-sink relationship between a cultivated rice *Oryza sativa* cv. Nipponbare and a wild rice *Oryza australiensis* that show striking differences in biomass and grain yield. *Oryza australiensis*, accumulating higher biomass, not only showed higher photosynthesis per unit leaf area but also exported more sucrose from leaves than Nipponbare. However, grain features and sugar levels suggested limited sucrose mobilization to the grains in the wild rice due to vasculature and sucrose transporter functions. Low cell wall Invertase activity and high Sucrose Synthase cleavage activity followed by higher expression of Cellulose Synthase genes in *Oryza australiensis* stem utilized photosynthates preferentially for the synthesis of structural carbohydrates, resulting in high biomass. In contrast, the source-sink relationship favored high grain yield in Nipponbare via accumulation of transitory starch in the stem, due to higher expression of starch biosynthetic genes, which is mobilized to panicles at the grain filling stage. Thus, vascular features, sucrose transport, and functions of sugar metabolic enzymes explained the differences in the source-sink relationship between Nipponbare and *Oryza australiensis*.

Keywords: Biomass; Grain yield; Photosynthesis; Source-sink relationship; Starch; Structural carbohydrates; Sucrose synthase; Sucrose transport; Vascular features

Introduction

Source-sink relationship is critical for determining the overall growth and
performance of crop plants (White *et al*., 2016). Photosynthetic carbon assimilation in source
leaves, sucrose transport through the vasculature, type and strength of sink tissues, as well as
the metabolic status of the tissues are the key features that determine the source-sink
dynamics of a plant. Photosynthetic leaves are the major source tissues of a plant, whereas
types and efficiency of sink organs vary depending upon the growth stages and
environmental conditions. For example, grains are the primary sink in the reproductive stage
of crop plants, while roots, stem/internodes, and growing leaves function as the sink in the
vegetative stage. Natural genetic variation for differences in the source-sink dynamics,
resulting in changes in biomass and yield, has been evident (Yin *et al.*, 2009; Burnett *et al.*,
2016; White *et al.*, 2016; Fabre *et al.,* 2020). For example, many of the wild relatives of rice
accumulate high biomass with poor grain filling, whereas cultivated varieties produce higher
grain yield (Sanchez *et al.*, 2013).

Leaf morphological and anatomical features along with photosynthesis per unit leaf
area largely determine the source strength of a plant (Mathan *et al.*, 2016). Functions of the
key carbohydrate metabolic enzymes, such as ADP-glucose pyrophosphorylase (AGPase),
sucrose phosphate synthase (SPS), sucrose phosphate phosphatase (SPP), and Sucrose
synthase (SUS), are known to be key determinants of the metabolic status of source leaves
(Osorio *et al.*, 2014; Ruan, 2014). Similarly, the capacity to utilize photosynthates towards
storage and maintenance largely determines the sink strength. Sink size, influenced by tiller
number, spikelet number per panicle, and seed size; and sink activity, determined by
functions of various metabolic enzymes such as cell wall invertase (cwINV) and sucrose
synthase (SUS), dictate the photosynthates utilization at a sink organ (Smith *et al.*, 2018;
Stein and Granot, 2019). Photosynthates are primarily partitioned from source leaves to
different sink tissues in the form of sucrose via phloem with the aid of SWEET and SUT
transporters (Chen *et al.*, 2012; Braun, 2012; Julius *et al.*, 2017). Mutations or overexpression
of genes encoding sucrose transporters affect plant yield and biomass via modulation of
source-sink dynamics (Scofield *et al.*, 2002; Chen *et al.*, 2012; Yang *et al.*, 2018). Besides
the sucrose transporter functions, vascular features not only determine the rate of sucrose
export from leaf tissues to phloem sap but also control mobilization of photosynthates to the
panicles at the reproductive stage (Qi *et al.*, 2008; Fujita *et al.*, 2013; Sack and Scoffoni,
2013).

Photosynthates assimilated in leaves are either used to meet immediate cellular needs
or mobilized into different sinks. Starch stored in leaves during the day, over and above the
immediate utilization and transport of photosynthates, is degraded during the night to be
transported to different sink organs for growth (Smith and Stitt, 2007; Chen *et al.*, 2012).
Mutants defective in sucrose transport, usually, accumulate starch in leaves at the end of the
night, limiting plant growth (Chen *et al.*, 2012). The utilization and storage of sugars are
regulated diurnally as well as developmentally. Stem serves as the main storage organ for
carbohydrates in crop plants, including rice, during vegetative growth. Sucrose, after getting
unloaded into the stem, could possibly be converted to different non-structural carbohydrates
(NSCs), such as glucose, fructose, and starch. Starch is reported to be the major storage form
in rice stem during the vegetative stage, which is mobilized to developing grains with the
onset of the reproductive stage, contributing up to half of the final grain yield (Wang *et al.*,
2017). A portion of the photosynthates is also used as a structural component of the cell wall
for the growth and development of new organs as well as elongation and thickening of
existing organs. High biomass accumulation is usually associated with the accumulation of
cellulose and other structural carbohydrates in crop plants, with cellulose alone contributing
around 25-50% of plant biomass (Haigler *et al.*, 2001). Thus, the balance of structural and
non-structural carbohydrates derived from the sucrose in the stem is critical for determining
biomass and grain yield in cereal crops.

Sucrose, after reaching the stem, is hydrolysed by two classes of enzymes, Invertases
(INV) and Sucrose Synthase (SUS). Invertases cleave sucrose into glucose and fructose,
while Sucrose Synthase (SUS) reversibly catalyses the formation of UDP-glucose and
fructose. The UDP-glucose, generated by the cleavage activity of SUS, is used as a precursor
for cellulose synthesis. Further, UDP-glucose can be converted to ADP-glucose for usage in
starch synthesis, making SUS a key enzyme for biomass and grain yield (Smith *et al.*, 2012;
Ruan, 2014; Stein and Granot, 2019). The functions of SUS are important for phloem loading
and unloading as well as for determining sink strength (Smith *et al.*, 2012; Fan *et al.*, 2017;
Yao *et al.*, 2020). Among the seven genes encoding SUS enzymes in rice, *OsSUS1* and
OsSUS3 contribute to seed starch biosynthesis (Cho *et al.*, 2011). Higher expression of
OsSUS3 also led to increased levels of structural carbohydrates, cellulose and hemicellulose,
in rice (Fan *et al.*, 2017). *OsSUS1* is highly expressed in rice internodes, with a strong
correlation with the expression pattern of genes encoding Cellulose Synthase, *OsCES4*,
OsCES7, and *OsCES9* (Hirose *et al.*, 2008; Guevara *et al.*, 2014). Among Cellulose Synthase
enzymes, OsCES4, OsCES7, and OsCES9 have been reported to function in secondary cell

wall formation (Tanaka *et al.*, 2003; Wang *et al.*, 2016). Thus, functions of enzymes, such as

INV, SUS, and CES, are important for determining source-sink dynamics for biomass and

grain yield.

Despite the key importance of sucrose partitioning and metabolism in determining the

source-sink relationship, this aspect is largely unexplored for optimization of grain yield and

biomass in rice. Cultivated rice varieties and their wild relatives, with striking differences in

yield and biomass traits, provide an excellent system to investigate the mechanistic basis of

the source-sink relationship. Comparative characterization of source and sink features along

with physiological and biochemical attributes for the selected wild and cultivated rice, which

differ in grain yield and biomass traits, might provide key information related to bottlenecks

for higher grain yield. In addition, such a study would also underline the factors contributing

to high biomass in wild rice. The knowledge can be utilized for streamlining the sucrose

transport and metabolism towards optimizing source-sink dynamics for higher yield and/or

biomass. Here, we investigated the limitations related to sucrose transport and metabolism for

grain yield in wild rice *O. australiensis*, resulting in higher biomass, compared to the

cultivated variety *O. sativa* cv. Nipponbare. We report a higher accumulation of structural

carbohydrates, cellulose and hemicellulose, in *O. australiensis* is due to lower cell wall

Invertase activity and higher SUS cleavage activity together with higher expression of genes

encoding Cellulose Synthases. Cultivated rice Nipponbare, in contrast, accumulated more

starch in the stem due to higher expression of genes encoding starch biosynthesis enzymes.

We also established the contribution of specific sucrose transporters and vascular features

towards limited mobilization of photosynthates to *O. australiensis* panicles. Results from

additional cultivated and wild rice accessions, *O. sativa* cv. IR 64 and *O. latifolia*, further

corroborated our model.

Materials and methods

Phenotyping of the selected rice accessions for biomass and yield-related traits

Two cultivated varieties, *Oryza sativa* ssp. *indica* cv IR 64 and *Oryza sativa* ssp. *japonica* cv

Nipponbare, and three wild rice species, *Oryza rufipogon* (IRGC 99562)*, Oryza latifolia*

(IRGC 99596)*,* and *Oryza australiensis* (IRGC 105272) were investigated for the study.

Seeds were germinated on the germination paper in a petri dish for 5-7 days. Germinated

seedlings were grown in half-strength Yoshida media for two weeks, and then transplanted in

the experimental field of National Institute of Plant Genome Research, New Delhi

(*latitude* 28°36 N; *longitude* 77°12 E; *altitude* 216 m). Plants were grown under natural

growing conditions with average air temperature >30°C, 70-80% humidity, and more than 100 cm annual rainfall. All the phenotyping was performed after the transition to the reproductive stage. At least ten individual plants of each genotype were used for phenotyping. Plant architectural traits were quantified at the milk-stage of grain-filling. Seed traits and fresh and dry weight were quantified at the grain maturity stage. Plant height was measured as the height from the soil surface to the tip of the central panicle; total leaves per plant, tiller number per plant, and internode number per the main stem were counted manually; and internode length was calculated using the distance between two consecutive nodes on the main stem. Leaf surface area was quantified for fully expanded top four leaves, including flag leaf, using a portable leaf area meter (LI-3000C, LI-COR Biosciences). Stem thickness and stem circumference were quantified using Fiji-Image J software from images of hand-cut sections of the main stem. Fresh weight was quantified by directly weighing the entire plant mass, and dry weight was quantified by weighing the dried plant mass after drying the entire plant mass at 60°C for five days. Spikelet per panicle was counted manually. For grain weight analysis, 1000-grains were randomly selected and weighed using an electronic balance. For grain yield per plant, physiologically dried grains per plant were harvested and weighed. Raw data of biomass and yield-related traits for each genotype were normalized to median from the entire sample set for Principal Component analysis. Then, all the data was converted on a log base 2 scale on JMP software from SAS (https://www.jmp.com/en_us/home.html).

Quantification of leaf photosynthesis rate and related physiological traits

Leaf photosynthesis rate (A), stomatal conductance (g_s), and intercellular CO_2 concentration (C_i) were recorded from flag leaves at booting, milk- and dough- stage of grain filling for *O. sativa* ssp. *japonica* cv. Nipponbare and *O. australiensis* using a portable photosynthesis system LI-6400XT (LI-COR Biosciences). A, gs, and C_i were quantified from the flag leaf regions fully exposed to sunlight at the booting stage, as leaves were not fully expanded. The values were measured from the middle widest part of fully expanded flag leaves at the milk- and dough- stage of grain filling. Data were recorded from at least 15 independent plants of each genotype on a clear day between 9.00 h to 11.00 h at ambient CO_2 level (C_a; 400 µmol mol^{-1}), photosynthetic photon flux density (PPFD, 1,500 µmol m^{-2} s^{-1}), 300 µmol s^{-1} flow rate, and 70% relative humidity. During measurements, the leaf chamber air temperature was set at 30ºC with maximum vapor pressure deficit in the range of 1.0-1.5 kPa.

Quantification of glucose, fructose, and sucrose

The non-structural carbohydrates (NSCs: glucose, fructose, and sucrose) present in the flag leaf and stem were quantified at the milk-stage of grain-filling by Gas Chromatography-Mass Spectrometry (GC/MS). Flag leaf and stem tissues were harvested from field-grown plants of both the genotypes at 9:00-11:00 am. For the quantification of NSCs in flag leaves at the end of the day (EOD) and end of the night (EON), tissue samples were collected at 6.00 pm and 6.00 am, respectively. The tissues were sampled from at least four individual plants per genotype and immediately frozen into liquid nitrogen after harvesting. A 100 mg fresh weight (FW) of the tissues was extracted using 1.0 ml water: chloroform: methanol (1:1:2.5) spiked with 10 µl internal standard (1 mg ml^{-1} ribitol in water). The mixture was vortexed, centrifuged and 200 µl of polar phase were dried for 3 hours in a speed-vac concentrator. Next, dried samples were derivatized with 50 µl of 20 mg ml^{-1} methoxyamine in pyridine warmed for 37°C for 120 mins with shaking. Metabolites were further derivatized by adding 70 µl of N-methyl N-trimethylsilyl-trifluoroacetamide (MSTFA) at 37°C for 30 min. Once derivatization was completed, samples were transferred to GC compatible vials and 1µl of samples were injected on GC-MS-QP2010 (Shimadzu) as described in Lisec *et al.* (2006). For metabolite identification, the mass spectra were matched with commercially available spectral libraries NIST08, NIST11, and Wiley08 for automated peak identification and validated by retention index (RI) libraries (http://gmd.mpimp-golm.mpg.de/). The relative amount of metabolites was presented by the total ion current signal which was normalized to Ribitol and tissue weight.

Starch quantification

Starch was quantified from flag leaves, stems, different internodes, and mature seeds of the selected wild and cultivated accessions. The quantification of starch from flag leaves and stems was performed at the milk-stage of grain-filling at the time of phenotyping of photosynthesis. In addition, flag leaves samples were used for starch quantification at EOD and EON. Starch was also quantified from matured leaves at the vegetative and booting stage, as well as in different internodes of *O. sativa* ssp. *japonica* cv. Nipponbare and *O. australiensis* at the milk-stage of grain-filling. Starch was quantified using Mega-Calc Total starch determination kit (K-TSTA; Megazyme) according to the manufacturer's protocol. Four independent biological replicates were analysed for each tissue type for each genotype.

^{14}C labelled sucrose loading assay

The phloem loading capacity of ^{14}C labelled sucrose was quantified by following the method

as described by Yadav *et al.* (2017). Briefly, rice flag leaves at the milk-stage of grain-filling

were cut, and the leaf bases were immediately transferred to petri-dish submerged with

MES/CaCl$_2$ buffer (20 mM MES, 2mM CaCl$_2$, pH 5.5 with KOH). Next, leaf discs (3.6×1.0

cm) were excised using a cork borer, and immediately placed abaxial side down in the 24-

well microtiter plate pre-filled with 1 ml MES/CaCl$_2$ buffer spiked with 1mM sucrose

solution (1mM unlabelled Sucrose supplemented with 0.81µCi ml^{-1} ^{14}C Sucrose). Leaf discs

immersed in the above solution were vacuum-infiltrated for at least 20 min with gentle

shaking at room temperature. Labelled leaf discs were transferred to a fresh microtiter plate

and washed twice with 1.0 ml MES/CaCl$_2$ buffer, blot dried on absorbent filter paper, and

immediately frozen on dry ice and lyophilized. Next, ^{14}C sucrose loading capacity into leaves

was quantified by scintillation counting (PerkinElmer). Four leaf discs were used for each

biological replicate, and six biological replicates were used for each genotype per experiment.

Sucrose quantification in phloem sap

The amount of sucrose in the phloem was quantified using the EDTA-facilitated exudation

technique described by King and Zeevaart (1974). Flag leaves of the cultivated and wild rice

were excised at the milk-stage of grain-filling at around 9-10 am (when leaf photosynthesis

was quantified), leaf bases were immediately recut under exudation buffer (10 mM, Hepes,

10 mM EDTA, pH 7.5) to allow dispersion of any leaf contaminants, and placed in a 15 ml

tube containing 5 ml of the exudation buffer. Samples were incubated in a humid chamber

(relative humidity > 90%) in the dark to prevent evapotranspiration, and exudates were

collected after 5 h of incubation. 2 ml of sample aliquots were spiked with 50 µl internal

standard (1 mg ml^{-1} ribitol in water), and freeze-dried for sucrose quantification using GC-

MS based method as described earlier. Individual flag leaves from different plants were

considered as separate biological replicates, and four biological replicates were analyzed for

each genotype.

RNA isolation, cDNA synthesis, and qRT-PCR analysis

Tissue samples for RNA isolation and qRT-PCR analysis were collected from the same

developmental stage of *O. sativa* ssp. *japonica* cv. Nipponbare and *O. australiensis*. Samples

were collected at the milk-stage of grain filling (ten days after heading) for the two species

from the same regions of flag leaves, stems, panicle bases, and spikelets. Total RNA from

flag leaf, stem, and panicle base was extracted using TRIzol reagent (Invitrogen), whereas

RNA from spikelet was extracted using plant RNA purification kit (Sigma-Aldrich) according to manufacturer's protocol. One microgram (1µg) of total RNA was reverse transcribed to first strand-cDNA by anchored oligodT priming using Thermo Scientific RevertAid cDNA synthesis kit following manufacturer's instructions. The primer pairs (Table S1) of selected targeted genes used in this study produced a single product as viewed through dissociation curve analysis. Further, the efficiency of each primer pair was also evaluated using a standard curve method where five cDNA quantities (1, 25, 50, 75, and 100 ng) from each genotype were used to construct a standard curve, and efficiency were calculated using the formula $E = (10^{[-1/slope]}-1)*100$. Expression analysis of each gene was performed in four biological replicates for each genotype. The transcript levels of genes were normalized independently to two stable internal controls actin (LOC_Os03g50885) and ubiquitin (LOC_Os03g03920) with similar results. Relative expressions of target genes are presented using actin as the internal control applying $2^{-\Delta Ct}$ method by Livak and Schmittgen (2001). *In silico* expression analysis of the genes was performed using OS_AFFY_RICE_6 dataset present in Genevestigator database (https://genevestigator.com/gv/).

Visualization and quantification of anatomical traits

The vascular features of flag leaves as well as vascular bundles in the inner and outer ring of the stem, and panicle base (0.5 cm above the panicle node) were visualized and quantified through hand-cut transverse sections. Flag leaves, stems, and panicle bases were hand-sectioned using a razor-blade at the milk-stage of grain-filling, at least from five independent plants for each genotype, and stained with toluidine blue O (0.02% toluidine blue O in water) as described by Mitra and Loqué, (2014). Photographs were taken under a bright field microscope (LMI). Quantification of the different anatomical traits was made using Fiji-Image J software (Wayne Rasband, National Institute of Health).

Cellulose staining and quantification

For histochemical visualisation of cellulose, transverse hand-sections of rice stem (approximately 20 µm thickness) at the milk-stage of grain-filling were stained using calcofluor white (0.2% calcofluor white M2R in water) staining solution as described by Ambavaram *et al.* (2011). Stained sections were photographed under UV light range using a fluorescence microscope (excitation filter: 340-380 nm; Carl Zeiss Microscope). The quantification of cellulose and hemicellulose was made using National Renewable Energy

306 Laboratory (NREL) protocols as described in Mund *et al.* (2016). The quantifications were
307 made in four biological replications for each genotype.
308

Extraction of stem tissue for quantification of sucrose metabolic enzyme activity
The enzyme activity was performed in the stem tissues of *O. sativa* ssp. *japonica* cv.
Nipponbare and *O. australiensis*. Tissue samples (100 mg) was ground in liquid nitrogen and
homogenized in extraction buffer (50 mM HEPES/NaOH (pH 7.5), 7.5 mM $MgCl_2$, 2 mM
EDTA, 2% (w/v) PEG 8000, 2% (w/v) PVP and 5 mM DDT) at 4 °C. The homogenate was
centrifuged for 1 min at ~16,000 × *g*, the pellet was discarded and the crude extract was used
for enzyme activity assay.
316

Estimation of cell wall Invertase activity
Cell wall Invertase activity was performed as described in Tomlinson *et al.* (2004). Briefly,
50 µl of crude enzyme extract was added to 150 µl assay mix, containing 0.1 M sucrose in 50
mM sodium acetate at pH 4.7, on ice. The assay reaction was incubated at 37°C for 30 min.
The reaction was alkalinized by the addition of 50 µl 1 M TRIS-HCl (pH 8.0) and then
heated at 85 °C for 3 min. To measure acid hydrolysis of sucrose and endogenous glucose
levels, two blanks were set up. The amount of released hexoses was measured enzymatically
using Sucrose, D-Fructose, D-Glucose assay kit (Megazyme), and Invertase activity was
expressed as described in Nishanth *et al.* (2018).
326

Estimation of Sucrose Synthase synthesis and cleavage activity
To quantify OsSUS synthesis activity, 50 µl of crude extract was incubated with 20 mM
HEPES/NaOH buffer (pH 7.5), 5 mM $MgCl_2$, 20 mM KCl, 12 mM fructose, 0.4 mM
phosphoenolpyruvate, 2 mM uridine diphosphate UDP-glucose, 20 U pyruvate kinase, 20 U
lactate dehydrogenase, and 0.15 mM NADH in total 1.0 ml final reaction volume. Similarly,
to quantify OsSUS cleavage activity, 50 µl of crude extract was incubated with 20 mM
HEPES/NaOH buffer (pH 7.5), 100 mM sucrose, 2 mM UDP, 2mM $MgCl_2$, 0.005 UDP-
glucose dehydrogenase, and 1.5 mM NAD^+ in total 1.0 ml final reaction volume. The
reaction mixture was mixed gently, and a decrease and increase in absorbance were recorded
for synthesis and cleavage activity, respectively, at 340 nm continuously for 0 to 300 sec
using a UV–Vis spectrophotometer. The OsSUS activity was calculated in terms of µmol
NADH oxidized (synthesis activity) and NAD+ reduced (cleavage activity) per min per mg
protein as described in Qazi *et al.* (2012).

Subcellular localization of *OsSUS1*

Full length of *OsSUS1* CDS without stop codon was cloned under constitutive 35S promoter with a C-terminal eYFP fusion into binary vector pEG10 (Earley *et al.*, 2006). The recombinant plasmid along with empty vector control was then, introduced into EHA105 strain of *Agrobacterium tumefaciens*. Each transformant was co-infiltrated with a plasma membrane-localized marker (PM-mCherry; Nelson *et al.*, 2007) into the leaves of *Nicotiana benthamiana*. Confocal laser scanning microscopy TCS SP5 (Leica Microsystems) was used to take the images with appropriate lasers.

Results

Biomass and grain yield differences among the selected wild and cultivated rice accessions

We initiated the study with two cultivated rice varieties, *Oryza sativa* ssp. *japonica* cv. Nipponbare and *Oryza sativa* ssp. *indica* cv. IR 64, and three wild relatives of rice, *Oryza rufipogon*, *Oryza latifolia,* and *Oryza australiensis* that show remarkable variations in overall growth and architectural features at both vegetative and reproductive stages (Fig. 1A). All the selected wild rice species grew taller compared to the two cultivated varieties (Fig. 1B). Among all the accessions, *O. sativa* cv. Nipponbare produced the lowest number of total leaves, while *O. rufipogon* showed the highest number of leaves per plant (Fig. 1C). In addition, Nipponbare showed a significantly lower tiller number per plant compared to all the selected accessions (Supplementary Fig. S1B). Selected wild rice species had a larger leaf surface area compared to the cultivated varieties (Fig. 1D). We, then, quantified the number and length of internodes in the main stem, as internodes are important contributors to the biomass of a plant. Cultivated rice varieties had fewer, smaller, and slender internodes in the main stem compared to the wild accessions (Fig. 1E; and Supplementary Fig. S1A). In addition, stem thickness and circumference were also lower for Nipponbare and IR 64 compared to all selected wild species (Supplementary Fig. S1C, D). Thus, plant architectural traits, such as height, branching, and internode features, as well as leaf traits attributed to the high biomass of the selected wild rice. Consistent with this, the selected wild rice also showed higher fresh and dry weight compared to the cultivated varieties. (Fig. 1F, G).

We, then, quantified seed traits of the selected cultivated and wild rice. The cultivated rice varieties produced higher numbers of spikelets per panicle as well as higher seed weight, resulting in higher grain yield per plant compared to the wild relatives (Fig. 1H-J). Principal

Component analysis for all the quantified biomass and grain-yield traits showed a clear
separation of wild and cultivated genotypes (Supplementary Fig. S2). Taken together, these
results confirmed the higher biomass accumulation at the expense of grain yield in the
selected wild rice compared to the cultivated varieties. Wild rice *O. australiensis* had higher
phenotypic values for all the traits contributing to biomass, including tiller number, compared
to Nipponbare. Therefore, we investigated the source strength, sink features, sucrose
translocation as well as the fate of the photosynthates towards biomass and yield using
representative wild rice and cultivated variety, *O. sativa* cv. Nipponbare and *O. australiensis*,
respectively.
383

Comparison of source efficiency and photosynthates utilization
In order to compare source efficiency, we quantified leaf photosynthesis rate and
related physiological traits as well as soluble sugars/non-structural carbohydrates (NSCs)
content in the leaves of the two selected accessions. A significant difference in the leaf
photosynthesis rate *A* was observed between the two species at booting, and milk- and dough-
stage of grain filling. Both the species showed an increasing trend of leaf photosynthesis from
booting to the milk stage, followed by reduction at the dough stage of grain filling (Fig. 2A).
The higher photosynthesis in *O. australiensis* at the three stages was also associated with
higher g_s and *Ci* compared to Nipponbare (Supplementary Fig. S3A, B). Next, we measured
the different soluble sugars/NSCs, such as sucrose, fructose, glucose, and starch, at the time
of photosynthesis quantification in leaves at the milk stage of grain filling to evaluate the
levels of primary carbon metabolites, the major outcome of photosynthesis. The abundance of
all four NSCs was significantly higher in the leaves of *O. australiensis* than Nipponbare (Fig.
2B, C). Interestingly, *O. australiensis* also accumulated higher levels of starch in leaves
compared to Nipponbare, suggesting an abundance of residual sugar in the wild rice that is
converted to starch. In contrast, the glucose, fructose, and sucrose content as well as starch
content were found to be significantly lower in *O. australiensis* grains compared to
Nipponbare, despite starch being the major reserve product of seeds (Supplementary Fig. S4).
Similarly, seed glucose, fructose, sucrose, and starch content were also found to be
significantly lower in two other wild species *O. latifolia* and *O. rufipogon* compared to the
cultivated varieties IR 64 and Nipponbare. Higher starch and soluble sugar content in *O.*
australiensis leaf and lower starch and soluble sugar levels in seed suggested possible
bottlenecks in the mobilization of photosynthates to grains in the wild rice.

We, then, quantified end of the day (EOD) and end of the night (EON) carbon status

in the leaves of Nipponbare and *O. australiensis*. Glucose and fructose level were

significantly lower in *O. australiensis* than *O. sativa* cv. Nipponbare at the EOD and EON

(Fig. 2D). Sucrose content was marginally, but significantly, higher in *O. australiensis*

compared to *O. sativa* cv. Nipponbare at the EOD as well as EON. At the EOD, *O.*

australiensis accumulated more starch in leaves than *O. sativa* cv. Nipponbare (Fig. 2E).

However, *O. australiensis* displayed a significantly lower leaf starch level than *O. sativa* cv.

Nipponbare at the EON, suggesting efficient utilization of stored starch in *O. australiensis*

during the night (Fig. 2E). This is in contrast to lower starch content in the grains of *O.*

australiensis as explained earlier (Supplementary Fig. S4). These results indicated the proper

utilization of the photosynthates in *O. australiensis*, however not towards grain filling,

suggesting the possibility of the alternative sink and alternative fate of photosynthates.

Phloem loading and export of sucrose from leaves

Since sucrose is the major transportable form of photosynthates from source leaves to

sink organs, we investigated the sucrose transport differences between the two accessions.

We performed [^{14}C] labelled sucrose assay in leaf discs of *O. sativa* cv. Nipponbare and *O.*

australiensis. A higher amount of ^{14}C per unit leaf disc area was detected in *O. australiensis*

(5598.83 CPM per cm^2) compared to that of in Nipponbare (3446.66 CPM per cm^2) (Fig. 2F).

We, then, quantified sucrose in the phloem sap of the two accessions. Phloem sap of *O.*

australiensis had four times higher sucrose content than *O. sativa* cv. Nipponbare (Fig. 2G).

Together, ^{14}C-assay and analysis of phloem sap showed better phloem loading and sucrose

export from the leaf in the wild rice *O. australiensis* compared to Nipponbare.

Expression pattern of genes encoding sucrose transporters

We examined the contribution of sucrose transporters for differences in phloem

loading and sucrose export from leaves. Since clade III SWEET transporters and SUT

transporters are the major sucrose transporters in plants, we preferentially quantified the

transcript levels of genes encoding those transporters. Expression analysis of *OsSWEET*s

using publicly available datasets showed preferentially source-specific expression of

OsSWEET13 (Supplementary Fig. S5A). *OsSWEET14* also followed a similar source-specific

expression pattern, but at a relatively lower expression level than *OsSWEET13*. *OsSWEET15*

and *OsSWEET11* showed preferentially sink (panicle and endosperm) specific expression

patterns. The expression levels of *OsSWEET12* was observed to be very low in all the tissues.

Consistent with this, quantitative Real Time-PCR results showed higher transcript levels of
OsSWEET13 in flag leaf and stem; *OsSWEET11* in the spikelet; and *OsSWEET15* in the stem,
spikelet, and flag leaf of Nipponbare (Fig. 3A-D). Interestingly, the transcript level of
OsSWEET13, encoding preferentially source-specific SWEET transporter, was significantly
higher in flag leaf and stem of *O. australiensis* compared to Nipponbare (Fig. 3A, B). *In*
silico expression analysis showed higher expression of *OsSUT1* and *OsSUT2* in different
tissue compared to other *OsSUT*s (Supplementary Fig. S5B). qRT-PCR analysis confirmed
the generally higher expression of *OsSUT1* and *OsSUT2* in different tissues of Nipponbare
(Fig. 3E-H). We detected higher transcript levels of *OsSUT1* in the flag leaf, and *OsSUT2* in
the stem of *O. australiensis* compared to Nipponbare (Fig. 3E, F). The expression pattern of
*OsSWEET*s and *OsSUT*s suggested the potential involvement of selected transporters, such as
OsSWEET13, OsSUT1, and OsSUT2, for higher phloem loading and sucrose export from
leaf to stem in the wild rice.

Despite higher leaf photosynthesis rate and more sucrose export from source leaves,
O. australiensis produced smaller and lighter grains with less starch content compared to the
cultivated variety Nipponbare. Therefore, we also examined the transcript levels of different
sucrose transporters in developing spikelets. The expression levels of all the *OsSWEET*s were
significantly lower in spikelets of *O. australiensis* compared to Nipponbare, with almost no
expression of preferentially sink-specific *OsSWEET15* and *OsSWEET11* (Fig. 3D). In
addition, the transcript levels of *OsSUT1* was significantly lower in developing spikelet of *O.*
australiensis than Nipponbare (Fig. 3H). Thus, reduced grain filling in *O. australiensis* could
potentially be associated with reduced expression of *OsSWEET15*, *OsSWEET11*, and
OsSUT1.

Differences in vascular features associated with sucrose transport

Vascular features are also key to photoassimilate partitioning from source to sink
tissues. Therefore, we quantified the vein number and vein width in the fully matured flag
leaves of *O. sativa* cv. Nipponbare and *O. australiensis*. *O. australiensis* leaves had more
number of wider veins compared to Nipponbare (Fig. 4A, B). Moreover, *O. australiensis* also
exhibited higher vein density with lesser interveinal distance compared to Nipponbare
(Supplementary Fig. S6A, B). We, then, checked the organization of vascular bundles at the
panicle base of the two accessions, as the panicle base is the site of attachment of spikelets to
the main plant (Zhang *et al.,* 2002; Zhai *et al.,* 2018). *O. australiensis* had a fewer number of
vascular bundles with a reduced area than Nipponbare at the panicle base (Fig. 4C, D). We

also quantified the vascular features in the stem of the two accessions (Fig. 4E, F).

Interestingly, we observed ~1.7 times larger vascular bundles in the *O. australiensis* stem

compared to Nipponbare. Further, *O. australiensis* stem also had approximately twice the

number of vascular bundles than Nipponbare stem due to wider stem (Fig. 4F). We also

checked the vascular features in additional cultivated and wild rice accessions, *O. sativa* cv.

IR 64 and *O. latifolia*, respectively. The results were found to be consistent with the

differences between Nipponbare and *O. australiensis* for flag leaves, stem, as well as panicle

base (Supplementary Fig. S7). Taken together, the sucrose mobilization to a panicle in *O.*

australiensis might likely be hindered due to defects in the vascular bundle at the panicle

base and minimal expression of *OsSWEET*s in developing spikelets, strengthening the

hypothesis of alternative utilization of photosynthates in the *O. australiensis* stem towards

higher biomass.

Starch accumulation and expression levels of starch-biosynthesis genes in stem

The higher leaf photosynthesis rate and phloem loading in leaves, along with vascular

features and the expression pattern of sucrose transporter genes indicated stem to be a major

sink organ utilizing sucrose received from leaves in *O. australiensis*. Therefore, we

quantified the content of NSCs in stems of the two species at the milk-stage of grain filling

(Fig. 5A, B). Significantly lower amounts of glucose, fructose, and sucrose were detected in

the stem of *O. australiensis* as compared to Nipponbare (Fig. 5A). This was in contrast to

more amount of sucrose being exported from leaves of *O. australiensis*. Therefore, we

suspected that *O. australiensis* might reserve a high amount of starch in their stem. To our

surprise, a negligible amount of starch was present in the stem of *O. australiensis* compared

to Nipponbare (Fig. 5B). Starch content was, further, found to be very low in the *O.*

australiensis stems at different stages and in different internodes (Fig. 5C, D). Starch content

in the stems of another cultivated variety, *O. sativa* cv. IR 64, and wild species, *O. latifolia*

and *O. rufipogon*, also showed a similar pattern. Similar to *O. australiensis*, a negligible

amount of starch was present in the stems of *O. latifolia* and *O. rufipogon* compared to IR 64

(Supplementary Fig. 8). We, then, checked the expression of genes involved in starch

biosynthesis. Since we were hypothesizing differential fate of sucrose in the stem, we initially

checked the expression of members of ADP-glucose pyrophosphorylase large subunit

(*OsAPL*) and small subunit (*OsAPS*), and starch synthase (*OsSS*) gene families in internode,

node, and culm of Nipponbare using publicly available data at Genevestigator

(https://genevestigator.com/gv/). We observed generally high expression of these genes in the
internode, node, and stem of the cultivated variety (Supplementary Fig. 9A, B). Since
OsAPL3, *OsAPS1,* and *OsSSIIb* from the starch biosynthesis gene families showed the
highest expression levels in the internode, we selected these genes for expression analysis at
multiple tissues using Genevestigator as well as qRT-PCR validation in the stem. Expectedly,
very high expressions of these genes were detected in the internode and stem of the cultivated
variety Nipponbare by both *in silico* expression analysis as well as qRT-PCR analysis (Fig.
5E; Supplementary Fig. S11A). Consistent with very low starch levels in *O. australiensis*
stem and internode, the transcript abundance of genes encoding two key enzymes, *OsAPL3*
and *OsAPS1*, was drastically less in *O. australiensis* compared to Nipponbare (Fig. 5E). A
lower amount of soluble non-structural carbohydrates, in particular remarkably low starch
content, despite more sucrose transport from leaves to stem, confirmed an alternative fate to
photosynthates in *O. australiensis* stem.
521

Expression levels of sucrose metabolism genes, Sucrose Synthase and cell wall Invertase
enzyme activity, and structural carbohydrate levels in stem

Invertases (INV) and Sucrose Synthase (SUS) are the key enzymes for the
degradation of sucrose. Expression of genes encoding Invertases has been shown to be
strongly correlated with starch synthesis (Bahaji *et al.*, 2014; Ruan, 2014). An *in silico*
expression analysis of members of Invertase gene-family, including cell wall Invertase
(*OsCIN*), cytoplasmic Invertase (*OsNIN*), and vacuolar Invertase (*OsINV*), showed high
expression of many of those genes in the stem, node, and internode tissues of the cultivated
rice (Supplementary Fig. S10A). Previously, *OsCIN1* and *OsINV2* were shown to be highly
abundant in rice stem as compared to other members of the gene-family (Ji et al 2005). We,
then, checked the expression pattern of representative genes encoding cell wall Invertase
(*OsCIN1*), cytoplasmic Invertase (*OsNIN8*), and vacuolar Invertase (*OsINV2*) in stem tissues
by qRT-PCR (Fig. 6A). The expression of *OsCIN1* was drastically lower in *O. australiensis*
stem compared to Nipponbare. In addition, *O. australiensis* also showed lower cell wall
invertase activity in the stem (Fig. 6B). Together, limited expression of a cell wall Invertase
gene along with lower cell wall Invertase enzyme activity would limit the breakdown of
sucrose in *O. australiensis* stem apoplast.

Among all the *OsSUS*s, *OsSUS1* was shown to be highly expressed in internodes
(Hirose *et al.*, 2008). *In silico* expression analysis also showed high expression of *OsSUS1* in
internode, node, and culm (Supplementary Fig. S10B, S11A). Therefore, we checked the

intracellular localization of OsSUS1 and the expression levels of the corresponding gene. The

OsSUS1-YFP signal overlapped with the plasma membrane-localized marker (PM-mCherry),

confirming OsSUS1 localization in the plasma membrane (Supplementary Fig. S12). *OsSUS1*

was expressed four times higher in *O. australiensis* stem than Nipponbare, indicating a

possibility of cellulose accumulation in *O. australiensis* stem (Fig. 6C). Since SUS catalyzes

a reversible cleavage of sucrose, it has a synthesis activity promoting sucrose synthesis and a

cleavage activity promoting sucrose degradation. The enzyme activity assay revealed a higher

OsSUS1 synthesis activity in Nipponbare, while higher cleavage activity in *O. australiensis*

(Fig. 6D, E). Lesser accumulation of starch, more transcript levels of *OsSUS1*, and higher

cleavage activity of OsSUS1 in *O. australiensis* stem compared to Nipponbare along with

localization of OsSUS1 to plasma membrane prompted us to check the cellulose content in

the stems of the two accessions. Cellulose synthase genes are usually expressed at the highest

levels in rice internodes compared to other tissues as evident from *in silico* expression

analysis (Supplementary Fig. S10C, S11B). Interestingly, transcript levels of *OsCES4*,

OsCES7, and *OsCES9*, key cellulose synthesis genes, were significantly higher in the stems

of *O. australiensis* than in Nipponbare (Fig. 6F). Calcofluor-white staining along with

cellulose quantification confirmed more cellulose deposition in *O. australiensis* stem

compared to Nipponbare (Fig. 6G, H). Like cellulose, hemicellulose content was also

significantly higher in *O. australiensis* stem compared to Nipponbare (Fig. 6H). We also

compared cultivated rice *O. sativa* cv. IR 64 and wild rice *O. latifolia*, and observed more

cellulose deposition in *O. latifolia* stem compared to IR 64 (Supplementary Fig. S13). Taken

together, lower expression and activity of cell wall Invertase, higher expression and cleavage

activity of OsSUS1 coupled with higher expression of Cellulose Synthase genes, at least in

part, led to the utilization of photosynthates in the *O. australiensis* stem towards cellulose

deposition.

Discussion

We investigated the differences in source-sink dynamics between a cultivated rice

variety *O. sativa* cv. Nipponbare, which is optimized for high grain yield, and a wild relative

of rice *O. australiensis*, which accumulates high biomass with poor grain yield. *O.*

australiensis had higher source strength, as evident by consistently higher leaf photosynthesis

rate compared to Nipponbare, at least in part due to vascular features of the leaf (Fig. 2A and

Fig. 4A, B). High leaf photosynthesis, usually, results in high biomass or yield depending

upon preferential sink tissues, provided efficient transport of photoassimilates from leaves

(Burnett *et al.*, 2016; Fabre *et al.*, 2020; Fernie *et al.*, 2020). Limitations in sucrose export

promote accumulation of sugars in leaves, thereby inhibiting leaf photosynthesis. Therefore,

an efficient sucrose export system from leaf would be warranted for the realization of higher

source strength of the wild rice to high biomass and/or yield. ^{14}C sucrose uptake assay, as

well as sucrose content in phloem sap, confirmed a better sucrose export system from wild

rice *O. australiensis* leaves (Fig. 2F, G). Larger vascular bundles and fewer number of

mesophyll cells between two consecutive veins potentially led to more export of sucrose from

leaves in *O. australiensis*, as reported in different plant systems (Qi *et al.*, 2008; Fujita *et al.*,

2013; Mathan *et al.*, 2016). In addition, higher expression of genes encoding sucrose

transporters, OsSWEET13 and OsSUT1, would also facilitate the phloem loading in the wild

rice (Fig. 3). High expression of *SbSWEET8-1* of sorghum, a close homolog of *OsSWEET13*,

in leaf and its function in phloem loading supports our idea of the important role of

OsSWEET13 for phloem loading (Mizuno *et al.,* 2016). Similarly, *OsSUT1* not only functions

for enhancing phloem loading for sucrose transport but also for retrieval of sucrose from the

apoplasm along the transport pathway (Scofield *et al.*, 2007). A higher amount of soluble

sugars in *O. australiensis* leaves, despite having higher sucrose export from leaves, compared

to the cultivated variety, further supported the better source strength of the wild rice (Fig. 2

B, C).

The better grain filling and higher grain yield of cultivated rice Nipponbare, despite

the relatively lower source strength compared to *O. australiensis*, could be explained by more

number of larger vascular bundles at the panicle base as well as desirable expression of

sucrose transporter genes at the panicle base and spikelets of the cultivated rice (Fig. 3, 4).

Panicle architecture, a key determinant of rice grain yield, is reported to be shaped by the

vascular pattern (Sasahara *et al.*, 1999; Terao *et al.*, 2010). Reduced expression of

OsSWEET13 and *OsSWEET15* in the developing spikelet, and of *OsSUT1* at the panicle base

and developing spikelet, along with a reduced supply of smaller vascular bundles at the

panicle base limited mobilization of photosynthates to grains in *O. australiensis* compared to

the cultivated. In agreement with this, *ossweet11 ossweet15* double mutant has been reported

to show a smaller seed size (Yang *et al.*, 2018). Similarly, RNA antisense lines of *OsSUT1*

showed reduced grain filling and grain weight (Scofield *et al.*, 2007). Increased supply of

larger vascular bundles to rice rachis has been shown to promote more grains per panicle

(Zhang *et al.*, 2002; Terao *et al.*, 2010; Zhai *et al.*, 2018). Altogether, higher source strength

and limited grain sink strength in *O. australiensis*, attributed to vascular features and sucrose

transporter functions, indicated preferential utilization of photosynthates in stem and

internodes, contributing to source-sink relationship differences between the selected cultivated and wild rice. In conjunction with this, larger vascular bundles would facilitate efficient sucrose transport and unloading in the wild rice stem. Such vascular features in the stem would, in turn, also provide mechanical strength to support the high biomass of *O. australiensis* (Aohara *et al.*, 2009). Our expression results indicated the potential involvement of OsSWEET13 and OsSUT2 in unloading a larger amount of sucrose being received in the wild stem.

There is a competition among the sink tissues for utilization of photosynthates (Patrick *et al.*, 2013; Durand *et al.*, 2018). According to the EcoMeristem model, the final plant architecture is an outcome of competition for resources among different plant parts that depends on photoassimilate-partitioning patterns (Luquet *et al.*, 2006). More number of longer internodes and thicker stem with larger leaves, driving high biomass in *O. australiensis* compared to the cultivated variety, suggested higher utilization of photosynthates for vegetative growth. The higher biomass and larger organ size of the wild rice would require more photosynthates for general respiration and maintenance. However, this would likely not explain the utilization of photosynthates in total as *O. australiensis* showed a remarkably higher amount of sucrose in phloem sap compared to the cultivated variety. Altogether, *O. australiensis* utilized photoassimilates reaching to stem towards larger vegetative organ size and number, cell-wall thickening as well as the overall maintenance of higher biomass.

The preferential utilization of photosynthates in the stem/internodes of wild rice *O. australiensis* in contrast to grains in the cultivated rice Nipponbare suggested differences in the photosynthates metabolism in the stem of the two species. Cultivated rice variety clearly accumulated higher starch content in the stem, which is mobilized to panicle during grain filling (Fig. 5B). *OsSWEET11* and *OsSWEET15* have been suggested to be important for the remobilization of carbon reserve from stem to grain, and OsSUT1 in the retrieval of sucrose from apoplasmic space to stem for conversion to transitory starch (Wang *et al.*, 2020). Indeed, the Nipponbare stem showed higher expression of *OsSWEET11*, *OsSWEET15*, and *OsSUT1* compared to the wild rice along with desirable vascular features at the panicle base. High expression of starch biosynthesis genes, further, contributed to the roles of the stem as an effective source in the cultivated variety at the grain filling stage (Fig. 7). The differences in the stem starch content between the two genotypes projected the stored transitory starch in the stem as a key for source-sink dynamics favoring high grain yield.

643   Cell wall Invertases mediate the breakdown of sucrose into glucose and fructose, which enter the cytoplasm through H+/hexose symporters (HXTs) (Ruan et al., 2010). Lower expression and activity of Cell wall Invertase in *O. australiensis* stem, which receives more amount of sucrose from the leaves, would lead to sucrose accumulation in the apoplasmic space of *O. australiensis* stem. In contrast, higher expression and activity of Cell wall Invertase in Nipponbare would generate relatively more hexoses that may facilitate the hexose transport pathway to promote starch biosynthesis. The excess amount of sucrose present in the apoplasmic space of *O. australiensis* can be cleaved by the plasma membrane-associated SUS enzyme to produce UDP-glucose, which eventually would promote biomass accumulation through cellulose deposition (Stein and Granot, 2019). Consistent with this, *O. australiensis* stem showed a higher expression of *OsSUS1* and higher cleavage activity SUS producing a higher amount of UDP-glucose and fructose compared to Nipponbare. UDP-glucose has been shown to serve as the prime source for cellulose synthesis (Smith *et al*., 2012). UDP-glucose can also be converted to starch (Asano *et al*., 2002; Koch, 2004). However, the possibility is minimized in *O. australiensis* due to the very limited expression of *OsAPL3* and *OsAPS1*. Thus, most of the SUS-generated UDP-glucose was converted to cellulose, supported by high expression of cellulose synthase genes, *OsCES4*, *OsCES7*, and *OsCES9* in *O. australiensis* stem (Fig. 6F). OsSUS1 has been reported to be located in SE-CC complexes of phloem, and localization of SUS in phloem is important for cellulose synthesis (Smith *et al*., 2012; Regmi *et al.*, 2016). Plasma membrane localization of OsSUS1 in this study together with earlier reported association of cellulose synthase complexes to plasma membrane highlighted the key importance of OsSUS1 for cellulose synthesis in rice (Supplementary Fig. S12, Lei *et al*., 2012). In addition, SUS functions in companion cells of phloem have been suggested to enhance the sucrose unloading at sink tissues, further helping in unloading photosynthates in wild rice stem (Nolte and Koch, 1993; Stein and Granot, 2019, Yao *et al.*, 2020). The potential role of SUS in cellulose synthesis, as well as in secondary cell wall thickening, has been investigated in the past (Coleman *et al.*, 2009; Baroja-Fernández *et al.*, 2012; Wei *et al.*, 2015). Consistent with the higher expression of *OsSUS1* in *O. australiensis*, overexpression of *SUS* has been shown to result in increased vegetative growth rate, plant height, and biomass in different plant species (Coleman *et al.*, 2006; Nguyen *et al.*, 2016; Stein and Granot, 2019). Contrary to the *O. australiensis*, higher synthesis activity of SUS in Nipponbare stem would facilitate the formation of sucrose from the stored starch at the heading stage to be mobilized for grain filling.

Given the key importance of synthesis and cleavage activities of SUS for carbon
partitioning differences between the two species, factors contributing to the difference in
enzyme activity are intriguing. pH and metabolic status of the tissue, in addition to cellular
localization, phosphorylation and oligomerization status of the enzyme, are known to be
important factors determining the synthesis or cleavage activity of SUS. Acidic pH (in the
range of 5.5 – 7.5) has been shown to promote cleavage activity, whereas pH in the range of
7.5 – 9.5 promotes the synthesis activity of the enzyme (Schmolzer et al., 2016; Stein and
Granot, 2019). SUS activity also depends on the carbohydrate status, particularly on sucrose
metabolism (Takeda et al., 2017). As discussed earlier, a higher amount of sucrose reaching
the stem of *O. australiensis* accumulated in the apoplasm, due to sucrose transporters and
lower cell wall invertase activity. The acidic pH of apoplasm would then promote higher
cleavage activity of SUS in *O. australiensis* stem, subsequently driving cellulose
accumulation. In contrast, more sucrose and/or hexoses would enter into the cytoplasm of
Nipponbare to be converted to starch. The metabolism of the transitory starch for subsequent
mobilization to grains in the form of sucrose might promote the synthesis activity of SUS in
Nipponbare. Therefore, apoplasmic levels of sucrose and metabolic status of the stem
appeared to be important determinants for the differences in synthesis and cleavage activity
of the enzyme between the two species. Nonetheless, differential functions of cell wall
Invertase and Sucrose Synthase between *O. australiensis* and Nipponbare would, at least in
part, explain the differential fate of photosynthates in the stem (Fig. 7).

In summary, differences in vascular features and sucrose transporter functions led to a
differential source-sink relationship between wild rice *O. australiensis* and cultivated variety
O. sativa cv. *Nipponbare*. *O. australiensis* showed source-sink dynamics favoring high
biomass through the accumulation of structural carbohydrates, mediated by lower cell wall
Invertase activity, higher cleavage activity of SUS enzyme together with higher expression of
Cellulose Synthase genes (Fig. 7). In contrast, source-sink dynamics favored higher grain
yield in Nipponbare via accumulation of transitory starch in the stem, to be mobilized to
panicles with the onset of grain filling. Taken together, vascular features and sucrose
transporter functions along with transitory starch storage mechanism and Invertase and SUS
enzyme activity can potentially be targeted for source-sink dynamics favoring either biomass
accumulation in fodder crops or higher grain yield in cereal crops.

Supplementary data

Fig. S1. Internode length, tiller number, and stem features of the selected cultivated and wild rice genotypes.

Fig. S2. Principal component analysis of biomass and yield traits of the selected cultivated and wild rice species.

Fig. S3. Stomatal conductance (g_s) and intercellular CO2 concentration (C_i) of a cultivated rice *O. sativa* cv. Nipponbare and a wild rice *O. australiensis* at different stages during booting and grain-filling.

Fig. S4. Quantification of glucose, fructose, sucrose, and starch content in matured seeds of the selected cultivated and wild rice genotypes.

Fig. S5. *In silico* organ-specific expression analysis of rice genes encoding clade III *SWEET* and *SUT* transporters using publicly available data at Genevestigator (https://genevestigator.com/gv/).

Fig. S6. Quantification of leaf interveinal distance and vein density for a cultivated rice *O. sativa* cv. Nipponbare and a wild rice *O. australiensis*.

Fig. S7. Vascular features in flag leaf, panicle base, and stem of a cultivated rice *O. sativa* cv. IR 64 and a wild rice *O. latifolia*.

Fig. S8. Total starch content in the stem of a cultivated rice *O. sativa* cv. IR 64 and two wild rice species, *O. rufipogon* and *O. latifolia*.

Fig. S9. *In silico* expression analysis of rice starch biosynthesis genes using publicly available data at Genevestigator (https://genevestigator.com/gv/) in culm (stem), node, and internode.

Fig. S10. *In silico* expression analysis of genes encoding rice Invertases (A), Sucrose synthases (B), and Cellulose synthases (C) in culm (stem), node, and internode using publicly available data at Genevestigator (https://genevestigator.com/gv/).

Fig. S11. *In silico* expression pattern of the selected key genes involved in sugar metabolism in rice using Genevestigator database (https://genevestigator.com/gv/) across multiple different tissues.

Fig. S12. Localization of OsSUS1-YFP in the plasma membrane of leaf epidermal cells of *Nicotiana benthamiana.*

Fig. S13. Calcofluor-white staining for cellulose deposition in the transverse stem sections of a cultivated rice *O. sativa* cv. IR 64 and a wild rice *O. latifolia*.

Table S1. List of primer pairs used in the study

Acknowledgements

This work was supported by the Innovative Young Biotechnologist Award (BT/09/IYBA/2015/01) and Ramalingaswamy Re-entry Fellowship (BT/RLF/reentry/05/2013) to AR from the Department of Biotechnology, Ministry of Science and Technology, India. JM and AS acknowledge their CSIR-JRF and SERB-NPDF fellowships, respectively. We also acknowledge Central Instrument Facility, NIPGR; Advanced Instrumentation Research Facility, JNU; Confocal Microscopy Facility, NIPGR; and DBT-eLibrary Consortium (DeLCON) for providing access to e-resources.

Author Contributions

JM and AR conceptualized the study and designed experiments. JM performed experiments. JM and AS analysed data. JM, AS and AR wrote the manuscript. All the authors have read and edited the final manuscript.

Data availability statement

Data sharing is not applicable to this article as all created data is already contained within this article or in the supplementary material.

References

Ambavaram MMR, Krishnan A, Trijatmiko KR, Pereira A. 2011. Coordinated activation of cellulose and repression of lignin biosynthesis pathways in rice. Plant Physiology **155,** 916–931.

Asano T, Kunieda N, Omura Y, et al. 2002. Rice SPK, a calmodulin-like domain protein kinase, is required for storage product accumulation during seed development: Phosphorylation of sucrose synthase is a possible factor. Plant Cell **14,** 619–628.

Aohara T, Kotake T, Kaneko Y, Takatsuji H, Tsumuraya Y, Kawasaki S. 2009. Rice brittle culm 5 (brittle node) is involved in secondary cell wall formation in the sclerenchyma tissue of nodes. Plant and Cell Physiology **50**, 1886–1897.

Bahaji A, Li J, Sánchez-López ÁM, *et al.* 2014. Starch biosynthesis, its regulation and biotechnological approaches to improve crop yields. Biotechnology Advances **32**, 87–106.

Baroja-Fernández E, Muñoz FJ, Li J, Bahaji A, Almagro G, Montero M, Etxeberria E, Hidalgo M, Sesma MT, Pozueta-Romero J. 2012. Sucrose synthase activity in the sus1/sus2/sus3/sus4 Arabidopsis mutant is sufficient to support normal cellulose and starch production. Proceedings of the National Academy of Sciences of the United States of America **109**, 321–326.

Braun DM. 2012. Sweet! The pathway is complete. Science **335**, 173–174.

Burnett AC, Rogers A, Rees M, Osborne CP. 2016. Carbon source–sink limitations differ between two species with contrasting growth strategies. Plant Cell and Environment **39**, 2460–2472.

Chen LQ, Qu XQ, Hou BH, Sosso D, Osorio S, Fernie AR, Frommer WB. 2012. Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. Science **335**, 207–211.

Cho J Il, Kim HB, Kim CY, Hahn TR, Jeon JS. 2011. Identification and characterization of the duplicate rice sucrose synthase genes OsSUS5 and OsSUS7 which are associated with the plasma membrane. Molecules and Cells **31**, 553–561.

Coleman HD, Ellis DD, Gilbert M, Mansfield SD. 2006. Up-regulation of sucrose synthase and UDP-glucose pyrophosphorylase impacts plant growth and metabolism. Plant Biotechnology Journal **4**, 87–101.

Coleman, H.D., Yan, J. and Mansfield, S.D. 2009. Sucrose synthase affects carbon partitioning to increase cellulose production and altered cell wall ultrastructure. Proceedings of the National Academy of Sciences of the United States of America **106**, 13118–13123.

Durand M, Mainson D, Porcheron B, Maurousset L, Lemoine R, Pourtau N. 2018. Carbon source–sink relationship in Arabidopsis thaliana: the role of sucrose transporters. Planta **247**, 587–611.

Earley KW, Haag JR, Pontes O, Opper K, Juehne T, Song K, Pikaard CS. 2006. Gateway-compatible vectors for plant functional genomics and proteomics. The Plant Journal **45**, 616–629.

Fabre D, Dingkuhn M, Yin X, Clément-Vidal A, Roques S, Soutiras A, Luquet D. 2020. Genotypic variation in source and sink traits affects the response of photosynthesis and growth to elevated atmospheric CO2. Plant Cell and Environment **43**, 579–593.

Fan C, Feng S, Huang J, *et al.* 2017. AtCesA8-driven OsSUS3 expression leads to largely enhanced biomass saccharification and lodging resistance by distinctively altering lignocellulose features in rice. Biotechnology for Biofuels **10**, 221.

Fernie AR, Bachem CWB, Helariutta Y, *et al.* 2020. Metabolic Aspects of Source – Sink Interactions. Nature Plants **6, 55-66**.

Fujita D, Trijatmiko KR, Tagle AG, *et al.* 2013. NAL1 allele from a rice landrace greatly increases yield in modern indica cultivars. Proceedings of the National Academy of Sciences of the United States of America **110**, 20431–20436.

Guevara DR, El-Kereamy A, Yaish MW, Mei-Bi Y, Rothstein SJ. 2014. Functional characterization of the rice UDP-glucose 4-epimerase 1, OsUGE1: A potential role in cell wall carbohydrate partitioning during limiting nitrogen conditions. PLoS ONE **9,** e96158.

Haigler CH, Ivanova-Datcheva M, Hogan PS, Salnikov V V., Hwang S, Martin K, Delmer DP. 2001. Carbon partitioning to cellulose synthesis. Plant Molecular Biology **47**,

29–51.

Hirose T, Scofield GN, Terao T. 2008. An expression analysis profile for the entire sucrose synthase gene family in rice. Plant Science **174**, 534–543.

Ji X, Ende WV, Laere AV, Cheng S, Bennett J. 2008. Structure, evolution, and expression of the two Invertase gene families of rice. Journal of Molecular Evolution **60**, 615–634.

Julius BT, Leach KA, Tran TM, Mertz RA, Braun DM. 2017. Sugar transporters in plants: New insights and discoveries. Plant and Cell Physiology **58**, 1442–1460.

King, R.W. and Zeevaart, J.A.D. 1974. Enhancement of Phloem Exudation from Cut Petioles by Chelating Agents. Plant Physiol*o*gy **53**, 96–103.

Koch, K. 2004. Sucrose metabolism: Regulatory mechanisms and pivotal roles in sugar sensing and plant development. Current Opinion in Plant Biology **7**, 235–246.

Lei, L., Li, S., and Gu, Y. 2012. Cellulose synthase complexes: composition and regulation. Frontiers in plant science **3**, 75.

Lisec J, Schauer N, Kopka J, Willmitzer L, Fernie AR. 2006. Gas chromatography mass spectrometry-based metabolite profiling in plants. Nature Protocols **1**, 387–396.

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2-ΔΔCT method. Methods **25**, 402–408.

Luquet D, Dingkuhn M, Kim H, Tambour L, Clement-Vidal A. 2006. EcoMeristem, a model of morphogenesis and competition among sinks in rice. 1. Concept, validation and sensitivity analysis. Functional Plant Biology **33**, 309–323.

Mathan J, Bhattacharya J, Ranjan A. 2016. Enhancing crop yield by optimizing plant developmental features. Development (Cambridge) **143**, 3283-3294.

Mitra, P.P. and Loqué, D. 2014. Histochemical staining of Arabidopsis thaliana secondary cell wall elements. Journal of Visualized Experiments **13**, e51381.

Mizuno, H., Kasuga, S. and Kawahigashi, H. 2016. The sorghum *SWEET* gene family: stem sucrose accumulation as revealed through transcriptome profiling. Biotechnology for Biofuels **9,** 127.

Mund, N.K., Dash, D., Barik, C.R., Goud, V. V., Sahoo, L., Mishra, P. and Nayak, N.R. 2016. Chemical composition, pretreatments and saccharification of Senna siamea (Lam.) H.S. Irwin & Barneby: An efficient biomass producing tree legume. Bioresource Technology **207**, 205–212.

Nelson, B.K., Cai, X. and Nebenführ, A. 2007. A multicolored set of in vivo organelle markers for co-localization studies in Arabidopsis and other plants. Plant Journal **51**, 1126–1136.

Nguyen QA, Luan S, Wi SG, Bae H, Lee DS, Bae HJ. 2016. Pronounced phenotypic changes in transgenic tobacco plants overexpressing sucrose synthase may reveal a novel sugar signaling pathway. Frontiers in Plant Science **6**, 1216.

Nishanth MJ, Sheshadri SA, Rathore SS, Srinidhi S, Simon B. 2018. Expression analysis of *Cell wall invertase* under abiotic stress conditions influencing specialized metabolism in *Catharanthus roseus*. Scientific Reports **8,** 15059.

Nolte KD, Koch KE. 1993. Companion-cell specific localization of sucrose synthase in zones of phloem loading and unloading. Plant Physiology **101**, 899–905.

Osorio S, Ruan YL, Fernie AR. 2014. An update on source-to-sink carbon partitioning in tomato. Frontiers in Plant Science **5**, 516.

Patrick JW, Botha FC, Birch RG. 2013. Metabolic engineering of sugars and simple sugar derivatives in plants. Plant Biotechnology Journal **11**, 142–156.

Qazi, H.A., Paranjpe, S. and Bhargava, S. 2012. Stem sugar accumulation in sweet sorghum - Activity and expression of sucrose metabolizing enzymes and sucrose transporters. Journal of Plant Physiology **169**, 605–613.

Qi J, Qian Q, Bu Q, *et al.* 2008. Mutation of the rice narrow leaf1 gene, which encodes a novel protein, affects vein patterning and polar auxin transport. Plant Physiology **147, 1947–1959.

Regmi KC, Zhang S, Gaxiola RA. 2016. Apoplasmic loading in the rice phloem supported by the presence of sucrose synthase and plasma membrane-localized proton pyrophosphatase. Annals of Botany **117**, 257–268.

Ruan Y-L. 2014. Sucrose Metabolism: Gateway to Diverse Carbon Use and Sugar Signaling. Annual Review of Plant Biology **65**, 33–67.

Sack L, Scoffoni C. 2013. Leaf venation□: structure , function , development , evolution , ecology and applications in the past, present and future. New Phytologist **4**, 983–1000.

Sanchez, P.L., Wing, R.A. and Brar, D.S. 2013. The wild relative of rice: genomes and genomics. In *Genetics and genomics of rice* (Zhang, Q. and Wing, R.A., eds). Springer, pp. 10–25.

Sasahara H, Fukuta Y, Fukuyama T. 1999. Mapping of QTLs for Vascular Bundle System and Spike Morphology in Rice, Oryza sativa L. Breeding Science **49**, 75–81.

Schmolzer K, Gutmann A, Diricks M, Desmet T, Nidetzky B. 2016. Sucrose synthase: a unique glycosyltransferase for biocatalytic glycosylation process development. Biotechnology Advances **34**, 88–111.

Scofield GN, Hirose T, Aoki N, Furbank RT. 2007. Involvement of the sucrose transporter, OsSUT1, in the long-distance pathway for assimilate transport in rice. Journal of Experimental Botany **58**, 3155–3169.

Scofield GN, Hirose T, Gaudron JA, Upadhyaya NM, Ohsugi R, Furbank RT. 2002. Antisense suppression of the rice sucrose transporter gene, OsSUT1, leads to impaired grain filling and germination but does not affect photosynthesis. Functional Plant Biology **29**, 815–826.

Smith AM, Kruger NJ, Lunn JE. 2012. Source of sugar nucleotides for starch and cellulose synthesis. Proceedings of the National Academy of Sciences of the United States of America **109**, E776–E776.

Smith AM, Stitt M. 2007. Coordination of carbon supply and plant growth. Plant, Cell and Environment **30**, 1126–1149.

Smith, M.R., Rao, I.M. and Merchant, A. 2018. Source-sink relationships in crop plants and their influence on yield development and nutritional quality. Frontiers in Plant Science **871**, 1889.

Stein O, Granot D. 2019. An overview of sucrose synthases in plants. Frontiers in Plant

Science **10**, 95.

Takeda H, Niikura M, Narumi A, Aoki H, Sasaki T, Shimada H. 2017. Phosphorylation of rice sucrose synthase isoforms promotes the activity of sucrose degradation. Plant Biotechnology **34(2)**,107-113.

Tanaka K, Murata K, Yamazaki M, Onosato K, Miyao A, Hirochika H. 2003. Three distinct rice cellulose synthase catalytic subunit genes required for cellulose synthesis in the secondary wall. Plant Physiology **133**, 73–83.

Terao T, Nagata K, Morino K, Hirose T. 2010. A gene controlling the number of primary rachis branches also controls the vascular bundle formation and hence is responsible to increase the harvest index and grain yield in rice. Theoretical and Applied Genetics **120**, 875–893.

Tomlinson KL, McHugh S, Labbe H, Grainger JL, James LE, Pomeroy KM, Mullin JW, Miller SS, Dennis DT, Miki BLA. 2004. Evidence that the hexose-to-sucrose ratio does not control the switch to storage product accumulation in oilseeds: analysis of tobacco seed development and effects of overexpressing apoplastic invertase. Journal of Experimental Botany **55**, 2291–2303.

Wang DR, Han R, Wolfrum EJ, McCouch SR. 2017. The buffering capacity of stems: genetic architecture of nonstructural carbohydrates in cultivated Asian rice, Oryza sativa. New Phytologist **215**, 658–671.

Wang G, Li H, Gong Y, Yang J, Yi Y, Zhang J, Ye N. 2020*a*. Expression profile of the carbon reserve remobilization from the source to sink in rice in response to soil drying during grain filling. Food and Energy Security **3**, e204.

Wang LQ, Guo K, Li Y, Tu YY, Hu HZ, Wang BR, Cui XC, Peng LC. 2010. Expression profiling and integrative analysis of the CESA/CSL superfamily in rice. BMC Plant Biology **10**, 282–297

Wang D, Qin Y, Fang J, Yuan S, Peng L, Zhao J, Li X. 2016*a*. A missense mutation in the zinc finger domain of OsCESA7 deleteriously affects cellulose biosynthesis and plant growth in rice. PLoS ONE **11**, e0153993.

Wei Z, Qu Z, Zhang L, Zhao S, Bi Z, Ji X, Wang X, Wei H. 2015. Overexpression of poplar xylem sucrose synthase in tobacco leads to a thickened cell wall and increased height. PLoS ONE **10**, e0120669.

White, A.C., Rogers, A., Rees, M. and Osborne, C.P. 2016. How can we make plants grow faster? A source-sink perspective on growth rate. Journal of Experimental Botany **67**, 31–45.

Yadav, U., Khadilkar, A., Shaikh, M., Turgeon, R. and Ayre, B. (2017) Quantifying the Capacity of Phloem Loading in Leaf Disks with [^{14}C]Sucrose. Bio-Protocol **7**, e2658.

Yang J, Luo D, Yang B, Frommer WB, Eom JS. 2018. SWEET11 and 15 as key players in seed filling in rice. New Phytologist **218**, 604–615.

Yao D, Gonzales-Vigil E, Mansfield SD. 2020. Arabidopsis sucrose synthase localization indicates a primary role in sucrose translocation in phloem. Journal of Experimental Botany **71**, 1858–1869.

Yin X, Guo W, Spiertz JH. 2009. A quantitative approach to characterize sink-source relationships during grain filling in contrasting wheat genotypes. Field Crops Research **114**, 119-126.

Zhai L, Zheng T, Wang X, Wang Y, Chen K, Wang S, Wang Y, Xu J, Li Z. 2018. QTL mapping and candidate gene analysis of peduncle vascular bundle related traits in rice by genome-wide association study. Rice **11**, 13.

Zhang ZH, Li P, Wang LX, Tan CJ, Hu ZL, Zhu YG, Zhu LH 2002. Identification of quantitative trait loci (QTLs) for the characters of vascular bundles in peduncle related to indica -japonica differentiation in rice (*Oryza sativa L.*) Euphytica **128**, 279 –284.

Figure legends

Fig. 1. Comparison of biomass and yield-related traits of the selected cultivated and wild rice genotypes.

(A) Shown are the photographs of the selected rice genotypes at the reproductive stage (scale bar = 30 cm).

(B–J) Quantification of plant height (B), total leaves per plant (C), average leaf surface area (D), total internode number per main stem (E), fresh weight (F), dry weight (G), spikelet per panicle (H), 1000-grain weight (I), and grain yield per plant (J) of the selected rice genotypes. Significance of difference among the genotypes was calculated using One-Way ANOVA with Tukey's post-hoc test (n = 10 data points from different plants, $p \leq 0.05$).

Fig. 2. Leaf photosynthesis, sugar accumulation, phloem loading, and sucrose export from leaves of a cultivated rice *O. sativa* cv. Nipponbare and a wild rice *O. australiensis*.

(A) Quantification of flag leaf photosynthesis per unit area (*A)* at different stages during booting and grain-filling.

(B-C) Shown are glucose, fructose, and sucrose content (B), and total starch content (C) in the flag leaves of the cultivated and wild rice at the time of photosynthesis quantification.

(D-E) Quantification of glucose, fructose, and sucrose content (D), and total starch content (E) in the flag leaves of the cultivated and wild rice at the End Of Day (EOD) and End Of Night (EON).

(F) Uptake of [^{14}C] sucrose in the flag leaf discs expressed as counts per minute (CPM) per square centimeter of leaf area.

(G) Sucrose quantification in the phloem sap of the two species.

Data represents the mean and standard error of mean (SE), and significance of difference between the genotypes was calculated using student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 3. Tissue-specific expression pattern of genes encoding SWEET and SUT transporters in a cultivated rice *O. sativa* cv. Nipponbare and a wild rice *O. australiensis*.

(A–D) Transcript levels of *SWEET* genes in the flag leaf (A), stem (B), panicle base (C), and spikelet (D) at the milk-stage of grain-filling in the two rice species.

(E–H) Transcript levels of *SUT* genes in the flag leaf (E), stem (F), panicle base (G), and spikelet (H) at the milk-stage of grain-filling in the two rice species.

Data represents the mean and standard error of mean (SE) from four biological replicate per genotype. Significance of difference between the genotypes was calculated using student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 4. Vascular features in flag leaf, panicle base, and stem of a cultivated rice *O. sativa* cv. Nipponbare and a wild rice *O. australiensis*.

(A-B) Cross-sections of flag leaves of the two species (A, scale bar = 100 µm), and quantification of minor vein width and the total number of veins (B).

(C-D) Transverse sections at the panicle base (0.5 cm above panicle node) of the two species (C, scale bar = 250 µm), and quantification of number and area of vascular bundles (D).

(E-F) Transverse sections of stems of the two species (E, scale bar = 100 µm), and quantification of number and area of vascular bundles (F).

The peripheral concentric ring of the panicle base and stem is represented as the outer ring, and the inner wider circle is represented as the inner ring in figures.

Data represents the mean and standard error of mean (SE) from fifteen cross-section images collected from five different plants and significance difference between the genotype was based on student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 5. Quantification of soluble sugars and expression pattern of starch-biosynthesis genes in stems of a cultivated rice *O. sativa* cv. Nipponbare and a wild rice *O. australiensis*.

(A-B) Glucose, fructose, and sucrose content (A), and total starch content (B) in the stem of the two species at the milk-stage of grain-filling.

(C) Quantification of total starch in the stem at the vegetative and booting stage of the two species.

(D) Total starch content in the flag leaf sheath and different internodes at milk-stage of grain-filling of the two species.

(E) Transcript levels of starch-biosynthesis genes, *OsAPL3* (encoding ADP-glucose pyrophosphorylase large subunit), *OsAPS1* (encoding ADP-glucose pyrophosphorylase small

subunit), and *OsSSIIB* (encoding Starch synthase) in the stem of the two species at milk-stage of grain-filling.

Data represents the mean and standard error of mean (SE), and significance difference between the genotype was calculated using student t-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Fig. 6. Gene expression and activities of sugar metabolic enzymes, and structural carbohydrates levels in stem tissue of a cultivated rice *O. sativa* cv. Nipponbare and a wild rice *O. australiensis* at milk-stage of grain-filling.

(A) Transcript levels of genes encoding cell wall invertase (*OsCIN1*), cytoplasmic invertase (*OsNIN8*), and vacuolar invertase (*OsINV2*).

(B) Cell wall Invertase enzyme activity.

(C) Transcript levels of *Sucrose Synthase* 1 (*OsSUS1*).

(D-E) OsSUS enzyme activity in synthesis (D) and cleavage (E) direction.

(F) Transcript levels of *Cellulose Synthase* genes, *OsCES4*, *OsCES7*, and *OsCES9.*

(G) Calcofluor-white staining for cellulose deposition in the transverse stem sections of the two species. Red arrowhead shows higher deposition of cellulose in *O. australiensis* stem. Scale bar represents 500 µm.

(H) Quantification of cellulose and hemicellulose in the stems of the two species.

Data in barplots represent the mean and standard error of mean (SE), and significance of difference between the genotypes was calculated using student t-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Fig. 7. A model explaining the contribution of sucrose transport, sucrose metabolic enzyme activity, starch biosynthesis, and synthesis of structural carbohydrates towards yield and biomass differences between a wild rice *O. australiensis* and a cultivated rice *O. sativa* cv. Nipponbare.

High leaf photosynthesis rate coupled with higher expression of selected *OsSWEET* and *OsSUT* genes mediate export of more sucrose from a leaf in *O. australiensis* compared to *O. sativa* cv. Nipponbare. Sucrose, after getting unloaded into stem due to the functions of *OsSWEET13*, *OsSUT1,* and *OsSUT2*, accumulates in the wild rice apoplast due to lower expression of a gene encoding Cell wall Invertase (*OsCIN1*) along with the lower activity of Cell wall Invertase. Sucrose gets converted to UDP-G more efficiently in the wild rice *O. australiensis* due to higher expression of *OsSUS1* and more cleavage activity of OsSUS

compared to the cultivated rice Nipponbare. Higher expression of *OsCES4, OsCES7,* and *OsCES9*, then, leads to more synthesis of cellulose in the stem of the wild rice. In contrast, higher expression of starch-biosynthesis genes (*AGPase*, small and large subunit) leads to higher starch content in the stem of the cultivated rice. Higher synthesis activity of OsSUS along with more expression of *OsSWEET11* and *OsSWEET15* in the cultivated rice facilitate efficient remobilisation of sucrose from stem to panicles at the grain-filling stage. SUC, sucrose (yellow colour circle); F, fructose; G, glucose; UDP-G, UDP-glucose; ADP-G, ADP-glucose; *INV*, Invertase; cwINV, cell wall Invertase; SUS, Sucrose Synthase.

Fig. 1. Comparison of biomass and yield-related traits of the selected cultivated and wild rice genotypes.
(A) Shown are the photographs of the selected rice genotypes at the reproductive stage (scale bar = 30 cm).
(B–J) Quantification of plant height (B), total leaves per plant (C), average leaf surface area (D), total internode number per main stem (E), fresh weight (F), dry weight (G), spikelet per panicle (H), 1000-grain weight (I), and grain yield per plant (J) of the selected rice genotypes. Significance of difference among the genotypes was calculated using One-Way ANOVA with Tukey's post-hoc test (n = 10 data points from different plants, $p \leq 0.05$).

Fig. 2. Leaf photosynthesis, sugar accumulation, phloem loading, and sucrose export from leaves of a cultivated rice _O. sativa_ cv. Nipponbare and a wild rice _O. australiensis_.

(A) Quantification of flag leaf photosynthesis per unit area (_A_) at different stages during booting and grain-filling.

(B-C) Shown are glucose, fructose, and sucrose content (B), and total starch content (C) in the flag leaves of the cultivated and wild rice at the time of photosynthesis quantification.

(D-E) Quantification of glucose, fructose, and sucrose content (D), and total starch content (E) in the flag leaves of the cultivated and wild rice at the End Of Day (EOD) and End Of Night (EON).

(F) Uptake of [^{14}C] sucrose in the flag leaf discs expressed as counts per minute (CPM) per square centimeter of leaf area.

(G) Sucrose quantification in the phloem sap of the two species.

Data represents the mean and standard error of mean (SE), and significance of difference between the genotypes was calculated using student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 3. Tissue-specific expression pattern of genes encoding SWEET and SUT transporters in a cultivated rice *O. sativa* cv. Nipponbare and a wild rice *O. australiensis*.
(A–D) Transcript levels of *SWEET* genes in the flag leaf (A), stem (B), panicle base (C), and spikelet (D) at the milk-stage of grain-filling in the two rice species.
(E–H) Transcript levels of *SUT* genes in the flag leaf (E), stem (F), panicle base (G), and spikelet (H) at the milk-stage of grain-filling in the two rice species.
Data represents the mean and standard error of mean (SE) from four biological replicate per genotype. Significance of difference between the genotypes was calculated using student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 4. Vascular features in flag leaf, panicle base, and stem of a cultivated rice *O. sativa* cv. Nipponbare and a wild rice *O. australiensis*.

(A-B) Cross-sections of flag leaves of the two species (A, scale bar = 100 µm), and quantification of minor vein width and the total number of veins (B).

(C-D) Transverse sections at the panicle base (0.5 cm above panicle node) of the two species (C, scale bar = 250 µm), and quantification of number and area of vascular bundles (D).

(E-F) Transverse sections of stems of the two species (E, scale bar = 100 µm), and quantification of number and area of vascular bundles (F).

The peripheral concentric ring of the panicle base and stem is represented as the outer ring, and the inner wider circle is represented as the inner ring in figures.

Data represents the mean and standard error of mean (SE) from fifteen cross-section images collected from five different plants and significance difference between the genotype was based on student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 5. Quantification of soluble sugars and expression pattern of starch-biosynthesis genes in stems of a cultivated rice *O. sativa* cv. Nipponbare and a wild rice *O. australiensis*.
(A-B) Glucose, fructose, and sucrose content (A), and total starch content (B) in the stem of the two species at the milk-stage of grain-filling.
(C) Quantification of total starch in the stem at the vegetative and booting stage of the two species.
(D) Total starch content in the flag leaf sheath and different internodes at milk-stage of grain-filling of the two species.
(E) Transcript levels of starch-biosynthesis genes, *OsAPL3* (encoding ADP-glucose pyrophosphorylase large subunit), *OsAPS1* (encoding ADP-glucose pyrophosphorylase small subunit), and *OsSSIIB* (encoding Starch synthase) in the stem of the two species at milk-stage of grain-filling.
Data represents the mean and standard error of mean (SE), and significance difference between the genotype was calculated using student t-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Fig. 6. Gene expression and activities of sugar metabolic enzymes, and structural carbohydrates levels in stem tissue of a cultivated rice *O. sativa* cv. Nipponbare and a wild rice *O. australiensis* at milk-stage of grain-filling.

(A) Transcript levels of genes encoding cell wall invertase (*OsCIN1*), cytoplasmic invertase (*OsNIN8*), and vacuolar invertase (*OsINV2*).
(B) Cell wall Invertase enzyme activity.
(C) Transcript levels of *Sucrose Synthase* 1 (*OsSUS1*).
(D-E) OsSUS enzyme activity in synthesis (D) and cleavage (E) direction.
(F) Transcript levels of *Cellulose Synthase* genes, *OsCES4*, *OsCES7*, and *OsCES9.*
(G) Calcofluor-white staining for cellulose deposition in the transverse stem sections of the two species. Red arrowhead shows higher deposition of cellulose in *O. australiensis* stem. Scale bar represents 500 µm.
(H) Quantification of cellulose and hemicellulose in the stems of the two species.
Data in barplots represent the mean and standard error of mean (SE), and significance of difference between the genotypes was calculated using student t-test (*p < 0.05, **p < 0.01, ***p < 0.001).

Fig. 7. A model explaining the contribution of sucrose transport, sucrose metabolic enzyme activity, starch biosynthesis, and synthesis of structural carbohydrates towards yield and biomass differences between a wild rice *O. australiensis* and a cultivated rice *O. sativa* cv. Nipponbare.

High leaf photosynthesis rate coupled with higher expression of selected *OsSWEET* and *OsSUT* genes mediate export of more sucrose from a leaf in *O. australiensis* compared to *O. sativa* cv. Nipponbare. Sucrose, after getting unloaded into stem due to the functions of *OsSWEET13*, *OsSUT1,* and *OsSUT2*, accumulates in the wild rice apoplast due to lower expression of a gene encoding Cell wall Invertase (*OsCIN1*) along with the lower activity of Cell wall Invertase. Sucrose gets converted to UDP-G more efficiently in the wild rice *O. australiensis* due to higher expression of *OsSUS1* and more cleavage activity of OsSUS compared to the cultivated rice Nipponbare. Higher expression of *OsCES4, OsCES7,*and *OsCES9*, then, leads to more synthesis of cellulose in the stem of the wild rice. In contrast, higher expression of starch-biosynthesis genes (*AGPase*, small and large subunit) leads to higher starch content in the stem of the cultivated rice. Higher synthesis activity of OsSUS along with more expression of *OsSWEET11* and *OsSWEET15* in the cultivated rice facilitate efficient remobilisation of sucrose from stem to panicles at the grain-filling stage. SUC, sucrose (yellow colour circle); F, fructose; G, glucose; UDP-G, UDP-glucose; ADP-G, ADP-glucose; *INV*, Invertase; cwINV, cell wall Invertase; SUS, Sucrose Synthase.