***In silico* characterization and expression profiles of zinc transporter-like (LOC100037509)**

gene of tomato

Ahmad Humayan Kabir

Department of Botany, University of Rajshahi, Rajshahi 6205, Bangladesh

E-mail: ahmad.kabir@ru.ac.bd

Key message:

♦ ZIP protein homologs are localized in the plasma membrane and are linked to ZIP zinc

transporter (PF02535) domain at chromosome 7.

♦ ZIP protein motifs are associated with the ZIP zinc transporter, N-glycosylation site, and

phosphorylation site.

♦ Phylogenetic studies reveal a closet relationship of Solyc07g065380 with *Solanum pennellii*

homolog.

♦ ZIP protein homologs of *S. lycopersicum* and *S. pennellii* show unique signal peptide along

with extended alpha-helix.

ABSTRACT

Zinc (Zn) is an essential microelement for plants. ZIP transporters play a critical role in Zn

homeostasis in plants. This *in silico* study characterizes different characteristics of putative Zn

transporter of tomato (Solyc07g065380) and its homologs. A total of 10 ZIP protein homologs

were identified across nine plant species by protein BLAST. All these ZIP protein homologs

located at chromosome 7 showed 305-350 amino acid residues, 7-8 transmembrane helices, and

stable instability index. Further, these ZIP protein homologs are localized in the plasma

membrane at the subcellular level corresponding to the ZIP zinc transporter (PF02535) domain.

Gene organization analysis reveals the presence of 3 exon along with the position of the

promoter, TATA-box, transcriptional start site, and splice sites in these ZIP transporter

homologs, in which tomato ZIP transporter (NM_001247420.1) contains a promoter, TATA-box,

transcriptional start site at 500, 911 and 946 bp, respectively along with several splice sites,

which may be useful for targeting binding sites and transcription factor analysis. Further, the

cutting sites and restriction enzymes of each ZIP gene homologs might be helpful for future

transgenic studies underlying Zn homeostasis. MEMO displayed five conserved motifs

associated with the ZIP zinc transporter, N-glycosylation site, and phosphorylation site.
Phylogenetic studies reveal a closet relationship of Solyc07g065380 with *Solanum pennellii*
homolog, while ZIP transporter of *Nicotiana sylvestris* and *Nicotiana tabacum* predicted to be in
close connection. The Solyc07g065380 transporter is predominantly linked to several
uncharacterized zinc metal ion transporters and expressed in diverse anatomical part,
developmental stage, and subjected to pathogen and heat stress. The secondary structural
prediction reveals unique signal peptide in the ZIP protein homologs of *S. lycopersicum* and *S.*
pennellii along with extended alpha-helix. These bioinformatics analyses might provide essential
background to perform wet-lab experiments and to understand Zn homeostasis for the
development of Zn-biofortified crops.

Keywords: Zn transporter, homologs, interactome analysis, transmembrane helix, tomato.

Running head: *In silico* characterization of zinc transporter-like gene
Introduction
Zinc (Zn) shortage, a plant nutritional deficit, has adverse effects on vegetable (Mattiello et al.
2015). There are lands with high pH or alkaline stress in many parts of the world (Alloway 2009;
Cakmak 2008), the primary explanation for Zn-deficient soil. The soil of lower pH, sandy
texture, and lower phosphorus are often deficient in Zn (Marschner, 1995). Zn-starved plants
often show impressive growth, tiny stems, and mild chlorosis (Marschner, 1995). Zn functions in
photosynthetic and gene expression processes in in addition to enzymatic and catalytic activities
(Welch 2001). Zn deficiency also resulted in a decline in plant stomatal activity (Mattiello et al.
2015). In response to pathogenic attacks on plants, Zn proteins also play a significant role (Cabot
et al. 2019). Zn's insufficiency in plant-based foods contributes to global human malnutrition
(Hotz et al. 2004).

While generally there are genotypic disparities in Zn-deficiency responses, lots of plants are
susceptible to Zn deprivation (Hacisalihoglu et al. 2001; Khatun et al. 2018). While Zn
efficiency mechanisms are fairly complex, research found several adaptive characteristics in Zn
deficient plants (Höller et al. 2014; Hacisalihoglu et al. 2003). As Zn cannot penetrate the root
cell membranes, Zn's absorption is achieved primarily via transporters (Kabir et al. 2017). The

IRT, known as Fe-regulated transporter is often induced to enhance translocation as well as Zn

absorption whenever the plant has possibly Zn or Fe shortcomings (Lin et al. 2009; Eckhardt et

al. 2001). *IRT1* is mainly responsible for transporting different metals like Zn, Mn, and Fe,

despite the fact that the pattern of expression varies depending on mineral stresses (Durmaz et al.

2011; Eckhardt et al. 2001). Till today, progress is pronounced to imparting the performance of

ZIP (Zn transporter) in Fe and Zn uptake (Morea et al. 2002; Xu et al. 2010), though the proof in

tomato continues to be restricted. Pavithra et al. (2016) classified low-affinity (*SlZIP5L2*,

SlZIP5L1, *SlZIP4*, and *SlZIP2*) and high-affinity (*SlZIPL*) Zn transporters in tomato. In a recent

study (Akther et al. 2020) demonstrated that upregulation of SlZIPL and SlZIP2 is involved with

the increase of Zn in Zn-deficient tomato. However, there is a putative zinc transporter

(Solyc07g065380) provisionally submitted to NCBI.

Tomato (*Solanum lycopersicum* L.) is a high-value crop worldwide (Wilcox et al., 2003). The

deficiency of the essential micronutrient Zn harms tomato production worldwide. Therefore, the

characterization of Zn transporter possesses immense potential to combat Zn-deficiency in

plants. Therefore, the molecular characterization of ZIP homologs may provide in-depth insight

into these genes/proteins. In this study, we have searched for ZIP transporter homologs based on

putative zinc transporter of tomato (Solyc07g065380/LOC100037509) in different plant species.

The CDS, mRNA, and protein sequences of these ZIP homologs were taken into different

computational analysis with advanced bioinformatics and an online-based platform along with

the wet-lab expression of mRNA transcript of putative zinc transporter in tomato under

differential Zn-conditions.

Materials and Methods

Retrieval of ZIP genes/proteins

Zn transporter-like (LOC100037509) gene named as Solyc07g065380 in Uniprort/Aramene

database (protein accession: NP_001234349.1 and gene accession: NM_001247420.1) was

obtained from NCBI to use as a reference for homology search (Stephen et al. 1997). The search

is filtered to match records with expect value between 0 and 0. The corresponding FASTA

sequences of gene and protein were retrieved from NCBI database.

Analyses of ZIP genes/proteins

Physico-chemical features of MTP protein sequences were analyzed by the ProtParam tool

(https://web.expasy.org/protparam) as previously described. Transmembrane (TM) helix

prediction was carried out by TMHMM Server (http://www.cbs.dtu.dk/services/TMHM). The

CELLO (http://cello.life.nctu.edu.tw) server predicted the subcellular localization of proteins

(Yu et al. 2006). Chromosomal locations, protein domain families, and functions were searched

in ARAMEMNON (http://aramemnon.uni-koeln.de/).

Organization of ZIP genes

The organization of exon and intron was analyzed by the ARAMEMNON web tool.

Organization of coding sequence along with the position of the transcriptional start site, TATA-

box, and the splice sites were predicted by FGENESH, TSSPlant, and FSPLICE tools

(http://www.softberry.com/berry.phtml). Further, the promoter site was predicted by the

Promoter 2.0 Prediction Server (http://www.cbs.dtu.dk/services/Promoter/). The analysis of

restriction   sites   and   enzymes   were   predicted   by   Restriction   Analyzer

(http://molbiotools.com/restrictionanalyzer.html) and Restriction Enzyme Picker Online v.1.3

(https://rocaplab.ocean.washington.edu/tools/repk).

Phylogenetic relationships and identification of conserved protein motifs

Multiple sequence alignments of MTP1 proteins were performed to identify conserved residues

by using Clustal Omega and MView. Furthermore, the five conserved protein motifs of the

proteins were characterized by MEME Suite 5.1.1 (http://meme-suite.org/tools/meme) with

default parameters, but five maximum numbers of motifs to find. Motifs were further scanned by

MyHits (https://myhits.sib.swiss/cgi-bin/motif_scan) web tool to identify the matches with

different domains (Sigrist et al. 2010). The MEGA (V. 6.0) developed the phylogenetic tree with

the maximum likelihood (ML) method for 1000 bootstraps using ZIP homologs (Tamura et al.

2013).

Interactions and co-expression of tomato ZIP protein

The interactome network of Solyc07g065380 was generated using the STRING server

(http://string-db.org) visualized in Cytoscape (Szklarczyk et al. 2019). Further, gene co-

occurrence and neighborhood pattern were also retried from the STRING server. Additionally,

the expression data of Solyc07g065380 was retrieved from Genevestigator software and

analyzed at hierarchical clustering and co-expression levels on the basis of the

SL_mRNASeq_Tomato_GL-0.

Structural analysis of ZIP protein homologs:

Analysis of the outer membrane of ZIP proteins was performed by Philius Transmembrane

Prediction Server (http://www.yeastrc.org/philius/pages/philius/runPhilius.jsp) as previously

described (Reynolds et al. 2008). The secondary structure was described by the SOPMA tool

(https://npsa-prabi.ibcp.fr/cgi-bin/npsa_automat.pl?page=npsa_sopma.html).

Results

Retrieval of Zn transporter-like genes/proteins:

The blast analysis of the Zn transporter-like (LOC100037509) gene of tomato showed 10

homologs by filtering the E-value to 0.0. The retrieved gene/proteins include *Solanum*

lycopersicum, Solanum pennellii, Solanum tuberosum, Nicotiana sylvestris, Nicotiana tabacum,

Capsicum annuum, Nicotiana tomentosiformis, Nicotiana attenuate and *Camellia sinensis* (Table

1).

Physiochemical features and localization of MTP proteins

In total, 10 ZIP homolog proteins encoded a protein with residues of 305–350 amino acid having

residues having 32994.85-37677.87 (Da) molecular weight. The pI value, instability index, and

Grand average of hydropathicity ranged from 5.51-7.07, 27.36-31.19 (stable), 0.555-0.689,

respectively (Table 1). Notably, all these ZIP proteins showed 7-8 transmembrane helices

(TMH). All of these ZIP protein homologs belonging to the ZIP zinc transporter (PF02535)

domain localized in the plasma membrane having membrance as cellular components (Table 2).

These ZIP proteins are located in chromosome 7 involved in metal ion transmembrane

transporter activity (Table 2).

Organizational features of ZIP transporter genes:

The structural analysis of the ZIP genes showed the presence of 1 exon the coding sequence,

which varies in 1-1203 position (Table 3, Fig. 1). The most of the promoters (NM_001247420.1,

XM_015224935.2, XM_015308168.1, XM_016725928.1,) of ZIP genes are positioned at 500 bp

while some other are at 1200 bp (XM_019395528.1), 400 bp (XM_016650095.1,

XM_009604423.3) and 200 bp (XM_028258694.1) position. However, no promoter was

predicted in two ZIP protein (XM_016644574.1, XM_016650095.1) homologs (Table 3). The

position of transcriptional start site (TSS) ranged from 201-1245, whereas the TATA-box was

located from 884-1212 base positions (Table 3). We also found several AG and GT splice sites

among the ZIP homologs across the plant species analyzed (Table 3).

Analysis of restriction enzyme and cutting position (Fig. 6)

We have searched for cutting positions and a list of restriction enzymes to explore the genetic

information for future genome editing programs. All these ZIP transporters contain several

cutting positions for several restriction enzymes (Fig. 2). Out of these restriction enzymes, *Bglll*,

Sacl and *Nsil* are generally common across the ZIP transporter genes. The cutting position was

ranged from 182- 1442 base positions (Fig. 2).

Conserved motif, sequence similarities, and phylogenetic analysis

We have used the MEME tool to search for the five most conserved motifs in identified 10 ZIP

protein homologs (Fig. 3). Motif 1, 2, 3, 4, and 5 showed 50, 42, 50, 41, and 41 long residues of

amino   acids,   respectively.   These   five   motifs   were

CFHSVFEGIAIGVADSKADAWRALWTVCLHKIFAAIAMGIALLRMIPNRP,

SPYFMKWNEGFLVLGTQFAGGVFLGTALMHFLSDSNETFGDL,

TTQGVVADWIFAISMGLACGVFIYVSINHLLSKGYKPQKMVLVDKPHFKF,

TNKEYPFAFMLACAGYLLTMLADCVICFVYAKQNNSNBVQL,   and

KSNGIVAQGQSQVCDGREBYFSKAPLATASSLGDSILLIVA in the consecutive arrangement

(Fig. 3). Out of these 5 motifs, motif 1-3 is best matched to PF02535.14 (ZIP), while motif 4 and

5 showed the best matching to PS00001 (N-glycosylation site) and PS0006 (phosphorylation

site) sites (Fig. 3).

All also aligned all these 10 ZIP protein homologs to classify the regions of preserved

(consensus) protein sequences. The ZIP protein homologs showed 94.6-100% sequence

similarities among the species. The consensus sequence ranged from 70%-100% (Supplementary

Fig. S1.). The phylogenetic was divided into two main groups based on tree topologies, such as

A, B, and C (Fig. 4). In group A, ZIP transporter of *Solanum lycopersicum* showed 98%, 88%

and 89% bootstrap with *Solanum pennellii, Solanum tuberosum* and *Capsicum annuum*,

respectively (Fig. 4). Cluster B consisted of the 5 ZIP protein homologs, among which,

Nicotiana sylvestris and *Nicotiana tabacum* showed highest 92% bootstrap value. The cluster C

is solely placed with ZIP transporter of *Camelia sinensis* (Fig. 4).

Predicted interaction partner analysis

Predicted interaction partner analysis was performed for tomato ZIP transporter (100037509).

STRING showed ten interaction partners of 100037509 (350 aa), which include 101266778

(uncharacterized protein; Ralf-like 34), 101247486 (uncharacterized protein; copper transporter

5), 101245988 (annotation not available), Solyc01g087530.2.1 (uncharacterized protein; ZIP

metal ion transporter family), Solyc04g008340.2.1 (uncharacterized protein; ZIP metal ion

transporter family), 101253965 (uncharacterized protein; ZIP metal ion transporter family),

101255936 (uncharacterized protein; WCRKC thioredoxin 1, GGPSI (uncharacterized protein;

geranylgeranyl pyrophosphate synthase 1), 101253075 (uncharacterized protein; ATPase EI),

101262011 (zinc-binding dehydrogenase family protein) proteins (Fig. 5a). Further, ZIP protein

of *S. lycopersicum* showed co-expression with other protein members of the same species and

other organisms (*S. aureus, C. albicans, S. cerevisiae, S. pombe, A. thaliana*) (Fig. 45).

The genvestigator analysis against SL_mRNASeq_Tomato_GL-0 showed co-expression data of

tomato ZIP transporter (Solyc07g065380) in different anatomical parts, developmental stages

and perturbations (Fig. 6a-c). In the anatomical part, the Solyc07g065380 showed the maximum

potential of expression in the ovule, septum, columella, placenta, locular tissue, and vascular

tissue (Fig. 6a). Also, Solyc07g065380 is highly predicted to be expressed during flower, fruit

maturation, fruit ripening, fruit maturation, fruit initiation, and seedlings stages of development

(Fig. 6b). Under perturbation analysis, we predicted the effect of biotic and abiotic stress

interactions on Solyc07g065380 expression. It showed that Solyc07g065380 is highly

upregulated (2-4 fold) in response to several pathogens, including *P. putida, B. cockerelli, P.*

fluorescens and *P. syringae* (Fig. 6c). Although the data of abiotic stress is limited,

Solyc07g065380 seems to be induced due to heat stress (Fig. 6d).

Analysis of transmembrane helix in MTP1 proteins in different plant species

The outer membrane analysis of ZIP protein homologs exhibited variations in the localization of

transmembrane strands and topology loops across the proteins as predicted by Philius

transmembrane server (Fig. 7). Philius server displayed that only NP_001234349.1 and

XP_015080421.1 proteins contain a signal peptide at the C-terminus end. In addition to these

two ZIP proteins, XP_016581414.1 also displayed similar localization of 8 transmembrane

helices in the protein sequences (Fig. 7). Further, XP_015163654.1, XP_009764644.1,

XP_016500060.1 and XP_016505581.1 protein homologs without signal peptide showed

identical localization of 7 transmembrane helices 3 (Fig. 7). The XP_009602718.1,

XP_019251073.1 and XP_028114495.1 protein contain 7 transmembrane helices without any

signal peptide (Fig. 7). Among the ZIP protein homologs, NP_001234349.1 and

XP_015080421.1 showed the highest alpha helix, which is consistent with the presence of signal

peptide (Supplementary Fig. S2). On average, the alpha helix of all these ZIP proteins ranged

from 24.06% to 31.12% of the protein structure. In addition, the extended strand contains

21.60% to 28.12% of the structural organization of ZIP proteins (Supplementary Fig. S2). As

expected, random coil contains stands roughly around 50% locations of all ZIP proteins. None of

these ZIP protein homologs displayed the presence of 3_{10} helix, Pi helix, beta bridge, beta turn,

bend region and ambiguous state in structure (Supplementary Fig. S2).

Discussion

Characterization of a gene is of great interest to further accelerate the wet-lab experiments on

plant stress. This *in silico* work led to the identification of 10 ZIP protein homologs among nine

plant species. MSA shows these proteins are 98.5-100% coverage of similarity with 94.6-100%

matching of consensus sequences among the different ZIP proteins. Most ZIP proteins are

predicted to have eight potential transmembrane domains and a similar membrane topology in

which the amino- and carboxy-terminal ends of the protein are located on the outside surface of the plasma membrane (Guerinot 2000). Our *in silico* analysis showed the existence of 7-8 transmembrane helices in all 10 ZIP protein homologs connected to ZIP Zinc transporter (PF02535). All of these ZIP protein homologs are localized in chromosome 7 and the plasma membrane.

The position and organization of the coding sequence of a gene is considered to be a critical factor in predicting evolutionary relations and functional genomics potentialities. In this study, all ZIP gene homologs among the nine plant species showed 3 exons, suggesting that these ZIP genes are phylogenetically closer to each other. Localization of promoter plays an essential part in CRISPR-Cas9 and other genome editing studies in plant science. Our analysis suggests that the promoter of *ZIP* gene homologs are highly positioned at 400-500 bp. Additionally, we explored the position of TSS and splice sites of ZIP homologs, which are crucial in understanding the transcriptional and post-transcriptional modification of mRNA. Restriction site and specificity of restriction enzymes are essential before transformation studies on a particular gene of interest. In this study, several cutting sites and respective restriction enzymes have been revealed, which will be useful to develop transgenic tomato or other plants associated with Zn homeostasis. In general, it is known that genes without intron have recently evolved (Deshmukh et al. 2015), which is not the case for ZIP gene homologs, as evident in our analysis. ZIP transporters are not involved in cellular Zn homeostasis but also regulate the adaptive growth response in plants (Coninx et al. 2019).

Conserved motifs are identical sequences across species that are maintained by natural selection. A highly conserved sequence is of functional roles in plants and can be a useful start point to start research on a particular topic of interest (Wong et al. 2015). Among the 10 ZIP protein homologs, we searched for five motifs using the MEME tool. Out of the five motifs, three motifs mainly matched with the ZIP zinc transporter family. We also observed the N-glycosylation site in motif 4, which is in agreement with the previous report on the conserved features of NRAMP transporter (Curie et al. 2000). The relation of motif 5 with the phosphorylation site may be attributed to the release of Zn from intracellular stores leading to phosphorylation kinases and activation of signaling pathways as previously reported (Thingholm et al. 2020). The presence of

common and long conserved residues pinpoints that ZIP homologs may possess highly

conserved structures between species. Additionally, this information can be targeted for

sequence-specific binding sites and transcription factor analysis.

In phylogenetic analysis, we clustered the tree in 3 sub-groups. According to the tree, ZIP protein

of *S. lycopersicum* showed the closest relationship (98% bootstraps) with *S. pennellii*, while *S.*

tuberosum and *C. annuum* also existed with the cluster of A. Consistently, ZIP protein homologs

of five tobacco species clustered within B, suggesting the close evolutionary emergence from a

common ancestor. It also appears that the ZIP transporter of *S. lycopersicum* is relatively

distantly related to *C. sinensis* over the evolutional trends. However, it is not yet reported

whether this Solyc07g065380 putative Zn transporter is also involved in Zn uptake in other

closely related plant species. Thus, our results might infer a functional relationship ZIP

sequences in Zn or other metal uptake across different plant species.

Interactome map and co-expresion analysis were performed using the Solyc07g065380 in the

STRING platform. In the interactome map, tomato putative zinc transporter (100037509) seemed

to be associated with several interaction partners linked to uncharacterized ZIP metal ion

transporter family. As a result, these findings might be useful to characterize several ZIP metal

transporters in plants. In plants, ZIP family members have also been characterized in plants

involved in metal uptake and transport, including Zn (Kavitha et al., 2015). However, studies

also reported that Zn homeostasis is closed associated with P-type ATPase heavy metal

transporters (HMA). Both HMA2 and HMA4 were reported to be involved with Zn homeostasis

in Arabidopsis (Hussain et al., 2004). Our further search on the coexpression of Solyc07g065380

suggests that this particular ZIP transporter is highly possible to coexpressed with several genes

of *S. lycopersicum* as well as *A. thalina*. Overall, this interactome findings might provide

essential background for functional genomics studies of Zn uptake and transport in plants.

The potentiality of expression of a gene in different conditions is a crucial factor in the genome

editing program. The *in silico* analysis of expression in the Genevestigator platform showed

interesting outputs concerning the expression of Solyc07g065380 in different anatomical,

perturbations, and developmental stages. Although root is the primary focus of ZIP transporter,

this Genevestigator platform suggested the maximum potential of expression in ovule, septum,

columella, placenta, locular tissue, and vascular tissue. In wet lab experiences, several CDF and

ATPase family transporters were shown root-specific expression regulating Zn and Cu

homeostasis in plants (Desbrosses-Fonrouge et al. 2005). Further, Solyc07g065380 is

predominantly expressed in several development stages during maturation, although the

expression potential during seedlings development is crucial for Zn uptake and Zn-deficiency

tolerance in tomato. Besides, the expression of Solyc07g065380 seemed to be upregulated in

response to several pathogens and heat stress. Although the data of stress is limited in the

Genevestigator platform, it implies that Solyc07g065380 is highly active in response to

perturbation. Given this potentiality of Solyc07g065380, this study further advances our

knowledge to elucidate the uptake and mobilization of Zn and other metals in plants.

In this study, all of the identified sequences of ZIP proteins demonstrated acidic character having

the pI value of around 6 along with the stable instability index and positive hydropathicity. The

protein length of 10 ZIP proteins was 305-350. Several studies reported the length of amino acid

residues in the ZIP family transporter from 309–476 (Guerinot, 2000). Computational prediction

of the secondary structure further reveals the presence of the signal peptide in ZIP protein

homologs of *S. lycopersicum* and *S. pennellii*. Signal peptides function to prompt a cell to

translocate the protein, usually to the cellular membrane. Thus, it suggests that the ZIP protein of

these two Solanum species carries enormous potentiality in the Zn uptake system in plants. The

highest % of the alpha helix in these two ZIP protein homologs is in accordance with the higher

transmembrane domains as integral membrane proteins are predominantly located in alpha-

helices. These findings may provide insights into the protein architecture and particular function.

Conclusion

In conclusion, these computational studies analyzed 10 ZIP protein homologs in different plant

species. The analysis showed similar physicochemical properties, gene organization, and

conserved motifs related to the Zn ion transport. Sequence homology and phylogenetic tree of

ZIP proteins showed the closest evolutionary relationship of *S. lycopersicum* with *S. pennellii*.

In addition, the interactome map displayed the co-expression of Solyc07g065380 with a number

of closely related genes involved with the ZIP metal ion transporter family. In addition,

Solyc07g065380 does have a high potentiality to express in several organs at different developmental stages subjected to biotic and heat stress in the Genevestigator platform. This *in silico* analysis reveals several gene features, such as promoter position, TSS, restriction sites, and enzymes, which will be useful for functional genomics and transgenic studies related to Zn uptake in plants. The unique feature of having signal peptide in Solyc07g065380 and other structural organization of protein further confirm the potentiality of this putative Zn transporter. These findings will provide basic theoretical knowledge for future studies to better understand the function and features of gene/protein related to Zn homeostasis in various plants.

Acknowledgment

I would like to acknowledge the accompaniment of my little daughters, Arya and Alina without whom it was hard to get rid of the depression and to concentrate on this work in COVID-19 pandemic.

References

Akther MS, Das U, Tahura S, Prity SA, Islam M, Kabir AH. 2020. Regulation of Zn uptake and redox status confers Zn deficiency tolerance in tomato. Scientia Horticulturae 273, 109624.

Alloway, B.J., 2009. Soil factors associated with zinc deficiency in crops and humans. Environ Geochem Heal. 31, 537–548.

Cabot, C., Martos, S., Llugany, M., Gallego, B., Tolra, R., Poschenrieder, C., 2019. A role for zinc in plant defense against pathogens and herbivores. Front. Plant Sci. 4, 1171.

Cakmak, I., 2000. Enrichment of cereal grains with zinc: Agronomic or genetic biofortification? Plant Soil 302, 1–17.

Coninx, L., Smisdom, N., Kohler, A., Arnauts, N., Ameloot, M., Rineau, F., Colpaert, J. V., Ruytinx, J. (2019). SlZRT2 Encodes a ZIP family Zn transporter with dual localization in the ectomycorrhizal fungus *Suillus luteus*. Frontiers in microbiology, 10, 2251.

Curie C, Alonso J, Le Jean M, Ecker J, Briat J (2000) Involvement of NRAMP1 from Arabidopsis thaliana in iron transport. Biochem J 347:749–755.

Desbrosses-Fonrouge A.G., Voigt K., Schroder A., Arrivault S., Thomine S., Kramer U. 2005.
Arabidopsis thaliana *MTP1* is a Zn transporter in the vacuolar membrane which mediates
Zn detoxification and drives leaf Zn accumulation. FEBS Lett. 579, 4165–4174.

Deshmukh, R.K., Sonah, H., Singh, N.K., 2015. Intron gain, a dominant evolutionary process
supporting high level of gene expression in rice. J. Plant Biochem. Biotechnol.
25(2),142–6.

Durmaz, E., Coruh, C., Dinler, G., Grusak, M., Peleg, Z., Saranga, Y., Fahima, T., Yazici, A.,
Ozturk, L., Cakmak, I., 2011. Expression and cellular localization of *ZIP1* transporter
under zinc deficiency in wild emmer wheat. Plant Mol. Biol. Rep. 29(3), 582–596.

Eckhardt, U., Marques, A.M., Buckhout, T.J., 2001. Two iron-regulated cation transporters from
tomato complement metal uptake-deficient yeast mutants. Plant Mol. Biol. 45, 437–448.

Guerinot ML. 2000. The ZIP family of metal transporters. Biochimica et Biophysica Acta (BBA)
- Biomembranes. 1465, 190-198.

Hacisalihoglu G., Hart J.J., Wang Y.H., Cakmak I., Kochian L.V., 2003. Zinc efficiency is
correlated with enhanced expression and activity of zinc-requiring enzymes in wheat.
Plant Physiol. 131, 595-602.

Hacisalihoglu, G., Hart, J.J., Kochian, L.V., 2001. High- and low-affinity zinc transport systems
and their possible role in zinc efficiency in bread wheat. Plant Physiol. 125, 456–463.

Höller, S., Meyer, A., Frei, M., 2014. Zinc deficiency differentially affects redox homeostasis of
rice genotypes contrasting in ascorbate level. J Plant Physiol. 171, 1748-1756.

Hussain, D., Haydon, M.J., Wang, Y., Sherson, S.M., Young, J., Camakaris, J., Harper, J.F.,
Cobbett, C.S. 2004. P-type ATPase heavy metal transporters with roles in essential zinc
homeostasis in Arabidopsis. The Plant Cell. 16(5), 1327-1339.

Hotz, C., Brown, K.H., 2004. Assessment of the risk of zinc deficiency in populations and
options for its control. Food Nutr Bull. 25, 94–204.

Kabir, A.H., Hossain, M.M., Khatun, M.A., Haque, M.N., Sarkar, M.M.R., Haider, S.A., 2017.
Biochemical and molecular mechanisms associated with Zn deficiency tolerance and
signaling in rice (*Oryza sativa* L.). J. Plant Interact. 12, 447–456.

Kavitha, P., Kuruvilla, S., and Mathew, M. 2015. Functional characterization of a transition
metal ion transporter, OsZIP6 from rice (*Oryza sativa* L.). Plant Physiol. Biochem. 97,
165–174.

Khatun, M.A., Hossain, M.M., Bari, M.A., Abdullahil, K.M., Parvez, M.S., Alam, M.F., Kabir,
405   A.H., 2018. Zinc deficiency tolerance in maize is associated with the up-regulation of Zn
406   transporter genes and antioxidant activities. Plant Biol. 20, 765-770.

407   Lin, Y.F., Liang, H.M., Yang, S.Y., Boch, A., Clemens, S., Chen, C.C., Wu, J.F., Huang, J.L.,
408   Yeh, K.C., 2009. Arabidopsis IRT3 is a zinc-regulated and plasma membrane localized
409   zinc/iron transporter. New Phytol 182(2), 392–404.

410   Marschner, H., 1995. Mineral nutrition of higher plants. Boston: USA: Academic Press.

411   Mattiello, E.M., Ruiz, H.A., Neves, J.C., Ventrella, M.C., Araujo, W.L., 2015. Zinc deficiency
412   affects physiological and anatomical characteristics in maize leaves. J Plant Physiol. 183,
413   138–43.

414   Moreau, S., Thomson, R.M., Kaiser, B.N., Trevaskis, B., Guerinot, M.L., Udvardi, M.K., Puppo,
415   A., Day, D.A., 2002. *GmZIP1* encodes a symbiosis-specific zinc transporter in soybean.
416   J. Biol. Chem. 277(7), 4738–4746.

417   Reynolds SM, Käll L, Riffle ME, Bilmes JA, Noble WS (2008) Transmembrane Topology and
418   Signal Peptide Prediction Using Dynamic Bayesian Networks. PLoS Comput Biol 4(11):
419   e1000213.

420   Sigrist, C.J., Cerutti, L., de Castro, E., Langendijk-Genevaux, P.S., Bulliard, V., Bairoch, A.,
421   Hulo, N. 2010. ROSITE, a protein domain database for functional characterization and
422   annotation. Nucleic Acids Res. 38, D161-6.

423   Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M.,
424   Doncheva, N.T., Morris, J.H., Bork, P., Jensen, L.J., von Mering, C. 2019. STRING v11:
425   protein-protein association networks with increased coverage, supporting functional
426   discovery in genome-wide experimental datasets. Nucleic Acids Res. 47, D607-613.

427   Tamura K., Stecher G., Peterson D., Filipski A., Kumar S. 2013. MEGA6: Molecular
428   Evolutionary Genetics Analysis version 6.0. Molecular Biol. Evol. 30, 12, 2725– 2729.

429   Thingholm TE, Rönnstrand L, Rosenberg PA. 2020. Why and how to investigate the role of
430   protein phosphorylation in ZIP and ZnT zinc transporter activity and regulation. Cellular
431   and Molecular Life Sciences (2020) 77:3085–3102

432   Welch, R.M., 2001. Impact of mineral nutrients in plants on human nutrition on a worldwide
433   scale. In: Plant Nutrition – Food Security and Sustainability of Agroecosystems. The
434   Netherlands, pp. 284–285.

Wong, A., Gehring, C., Irving, H.R. 2015. Conserved Functional Motifs and Homology

Modeling to Predict Hidden Moonlighting Functional Sites. Frontiers Bioeng. Biotech. 3,

82.

Wilcox, J.K., Catignani, G.L., Lazarus, C., 2003. Tomatoes and cardiovascular health. Crit. Rev.

Food Sci. Nutri. 43(1), 1-18.

Xu, Y., Wang, B., Yu, J., Ao, G., Zhao, Q., 2010. Cloning and characterisation of *ZmZLP1*, a

gene encoding an endoplasmic reticulum-localised zinc transporter in *Zea mays*. Func.

Plant Biol. 37(3), 194–205.

Table 1. Physiochemical properties of ZIP transporter homologs.

Sl.	Protein Accession	Species	Protein length	MW (Da)	pI	transmembrane helices (TMH)	Instability index	Grand average of hydropathicity (GRAVY)
1	NP_001234349.1	*Solanum lycopersicum*	350	37677.87	5.75	8	31.08	0.555
2	XP_015080421.1	*Solanum pennellii*	347	37422.57	5.51	8	31.07	0.566
3	XP_015163654.1	*Solanum tuberosum*	315	33985.68	6.06	7	32.65	0.566
4	XP_009764644.1	*Nicotiana sylvestris*	324	34632.36	5.87	7	29.84	0.576
5	XP_016500060.1	*Nicotiana tabacum*	324	34642.40	5.87	7	31.19	0.573
6	XP_016505581.1	*Nicotiana tabacum*	308	33316.18	6.56	7	28.54	0.668
7	XP_016581414.1	*Capsicum annuum*	317	34266.01	5.47	8	27.36	0.574
8	XP_009602718.1	*Nicotiana tomentosiformis*	306	33070.99	7.07	7	29.66	0.688
9	XP_019251073.1	*Nicotiana attenuata*	305	32994.85	6.62	7	29.21	0.689
10	XP_028114495.1	*Camellia sinensis*	320	34154.71	6.23	7	29.75	0.567

447 Table 2. Domain, localization, cellular component, chromosome position and molecular function

448 of ZIP proteins.

Sl.	Protein Accession	Species	Domain	Localization	Cellular component	Chromosome number	Molecular function
1	NP_001234349.1	*Solanum lycopersicum*	*ZIP Zinc transporter (PF02535)*	*Plasma Membrane*	*membrane*	7	metal ion transmembrane transporter activity
2	XP_015080421.1	*Solanum pennellii*	*ZIP Zinc transporter (PF02535)*	*Plasma Membrane*	*membrane*	7	metal ion transmembrane transporter activity
3	XP_015163654.1	*Solanum tuberosum*	*ZIP Zinc transporter (PF02535)*	*Plasma Membrane*	*membrane*	7	metal ion transmembrane transporter activity
4	XP_009764644.1	*Nicotiana sylvestris*	*ZIP Zinc transporter (PF02535)*	*Plasma Membrane*	*membrane*	7	metal ion transmembrane transporter activity
5	XP_016500060.1	*Nicotiana tabacum*	*ZIP Zinc transporter (PF02535)*	*Plasma Membrane*	*membrane*	7	metal ion transmembrane transporter activity
6	XP_016505581.1	*Nicotiana tabacum*	*ZIP Zinc transporter (PF02535)*	*Plasma Membrane*	*membrane*	7	metal ion transmembrane transporter activity
7	XP_016581414.1	*Capsicum annuum*	*ZIP Zinc transporter (PF02535)*	*Plasma Membrane*	*membrane*	7	metal ion transmembrane transporter activity
8	XP_009602718.1	*Nicotiana tomentosiformis*	*ZIP Zinc transporter (PF02535)*	*Plasma Membrane*	*membrane*	7	metal ion transmembrane transporter activity
9	XP_019251073.1	*Nicotiana attenuata*	*ZIP Zinc transporter (PF02535)*	*Plasma Membrane*	*membrane*	7	metal ion transmembrane transporter activity

10	XP_028114495.1	*Camellia sinensis*	*ZIP Zinc transporter (PF02535)*	*Plasma Membrane*	*membrane*	7	metal ion transmembrane transporter activity

450 **Table 3.** Organization of ZIP genes and position features.

No.	Gene Accession	Exon number	Coding region	Promoter position	TATA-box position	Position of transcriptional start site (TSS)	Splice sites
1	NM_001247420.1	3	1-1053	500 (marginal prediction)	911	946	AG: 409, 444, 468, 658, 1012 GT: 694
2	XM_015224935.2	3	1-1044	500 (marginal prediction)	902	936	AG: 397, 432, 659, 1003 GT: 685
3	XM_015308168.1	3	1-1050	500 (marginal prediction)	908	942	AG: 397, 432, 655, 1009 GT: treshold
4	XM_009766342.1	3	37-1074	No promoter predicted	1212	1245	AG: 229, 322, 679, 802, 1033, 1097, 1221 GT: 311, 547, 1359
5	XM_016644574.1	3	163-975	No promoter predicted	-	261	AG: 130, 223, 580, 703, 934 GT: 212, 448
6	XM_016650095.1	3	122-984	400 (marginal prediction)	-	252	AG: 121, 130, 223, 375, 589, 712, 943 GT: 457
7	XM_016725928.1	3	33-1024	500 (marginal prediction)	884	918	AG: 409 GT: Treshold
8	XM_009604423.3	3	163-984	400 (marginal prediction)	-	-	AG: 121, 223, 375, 589, 712 GT: 457
9	XM_019395528.1	3	166-1203	1200 (highly likely prediction)	-	201	AG: 79, 349, 358, 451, 808, 931, 1162, 1216, 1332 GT: 440, 676, 1278
10	XM_028258694.1	3	75-1088	200 (marginal prediction)	1126	1159	AG: 475, 784, 819, 983 GT: 514, 564, 732, 1234, 1365

456
457 **Fig. 1**. Gene organization of ZIP transporter homologs.

NM_001247420.1	
Enzyme	Cutting Positions
AseI ATTAAT	973
BglII AGATCT	192
BspHI TCATGA	277
ClaI ATCGAT	886
MfeI CAATTG	859
NsiI ATGCAT	355
SacI GAGCTC	731

XM_015224935.2	
Enzyme	Cutting Positions
BglII AGATCT	182
BspHI TCATGA	267
Bsu36I CCTNAGG	782
ClaI ATCGAT	879
EagI CGGCCG	188
MfeI CAATTG	852
NsiI ATGCAT	345
SacI GAGCTC	724

XM_015308168.1	
Enzyme	Cutting Positions
AseI ATTAAT	972
BglII AGATCT	182
BspHI TCATGA	267
ClaI ATCGAT	885
EagI CGGCCG	188
MfeI CAATTG	858
NsiI ATGCAT	345
SacI GAGCTC	730

XM_009766342.1	
Enzyme	Cutting Positions
BglII AGATCT	227
BsrGI TGTACA	1435
BspHI TCATGA	1107
MfeI CAATTG	885
NdeI CATATG	856
NsiI ATGCAT	390
SacI GAGCTC	757
ScaI AGTACT	1442

XM_016644574.1	
Enzyme	Cutting Positions
BglII AGATCT	129
BspHI TCATGA	1009
MfeI CAATTG	787
NdeI CATATG	758
NsiI ATGCAT	292
SacI GAGCTC	659

XM_016650095.1	
Enzyme	Cutting Positions
AseI ATTAAT	910
BglII AGATCT	129
MfeI CAATTG	796
NdeI CATATG	767
NsiI ATGCAT	292

XM_016725928.1	
Enzyme	Cutting Positions
AflII CTTAAG	324
AseI ATTAAT	944
BglII AGATCT	157
ClaI ATCGAT	904
MfeI CAATTG	830
NdeI CATATG	801
NsiI ATGCAT	320
SacI GAGCTC	702

XM_009604423.3	
Enzyme	Cutting Positions
AseI ATTAAT	914
BglII AGATCT	133
MfeI CAATTG	800
NdeI CATATG	771
NsiI ATGCAT	296
SacI GAGCTC	72, 672

XM_019395528.1	
Enzyme	Cutting Positions
AseI ATTAAT	1127
BglII AGATCT	355
BsrGI TGTACA	1353
MfeI CAATTG	1013
NdeI CATATG	984
NsiI ATGCAT	518
PstI CTGCAG	87
SacI GAGCTC	885

XM_028258694.1	
Enzyme	Cutting Positions
BstBI TTCGAA	406
EcoRI GAATTC	408
MfeI CAATTG	731
MscI TGGCCA	910
NcoI CCATGG	183
NsiI ATGCAT	392
SacI GAGCTC	771

459 **Fig. 2**. Restriction site and enzymes of ZIP transporter homologs.

Fig. 3. Most conserved five motifs of ZIP homologs detected by MEME tool.

465
466 **Fig. 4**. Phylogenetic tree of ZIP homologs using Mega 6. Statistical method: Maximum
467 likelihood phylogeny test, test of phylogeny: bootstrap method, No. of bootstrap replications:
468 1000.
469
470
471
472

(a)

(b)

473
474
Fig. 5. Predicted interaction partners of tomato ZIP transporter protein. Interactome was
generated using Cytoscape for STRING data.

Fig. 6. Co-expression of tomato ZIP transporter gene in different anatomical part, developmental stage and perturbations.

Fig. 7. Prediction of transmembrane helix of ZIP protein homologs.

486 **Supplementary Fig. S1**. Multiple sequence alignment of ZIP protein homologs.

NP_001234349.1

```
Alpha helix      (Hh) :   110 is   31.43%
3₁₀ helix        (Gg) :     0 is    0.00%
Pi helix         (Ii) :     0 is    0.00%
Beta bridge      (Bb) :     0 is    0.00%
Extended strand  (Ee) :    73 is   20.86%
Beta turn        (Tt) :     0 is    0.00%
Bend region      (Ss) :     0 is    0.00%
Random coil      (Cc) :   167 is   47.71%
Ambiguous states (?)  :     0 is    0.00%
Other states          :     0 is    0.00%
```

XP_016505581.1

```
Alpha helix      (Hh) :    82 is   26.62%
3₁₀ helix        (Gg) :     0 is    0.00%
Pi helix         (Ii) :     0 is    0.00%
Beta bridge      (Bb) :     0 is    0.00%
Extended strand  (Ee) :    75 is   24.35%
Beta turn        (Tt) :     0 is    0.00%
Bend region      (Ss) :     0 is    0.00%
Random coil      (Cc) :   151 is   49.03%
Ambiguous states (?)  :     0 is    0.00%
Other states          :     0 is    0.00%
```

XP_015080421.1

```
Alpha helix      (Hh) :   108 is   31.12%
3₁₀ helix        (Gg) :     0 is    0.00%
Pi helix         (Ii) :     0 is    0.00%
Beta bridge      (Bb) :     0 is    0.00%
Extended strand  (Ee) :    73 is   21.04%
Beta turn        (Tt) :     0 is    0.00%
Bend region      (Ss) :     0 is    0.00%
Random coil      (Cc) :   166 is   47.84%
Ambiguous states (?)  :     0 is    0.00%
Other states          :     0 is    0.00%
```

XP_016581414.1

```
Alpha helix      (Hh) :    81 is   25.55%
3₁₀ helix        (Gg) :     0 is    0.00%
Pi helix         (Ii) :     0 is    0.00%
Beta bridge      (Bb) :     0 is    0.00%
Extended strand  (Ee) :    91 is   28.71%
Beta turn        (Tt) :     0 is    0.00%
Bend region      (Ss) :     0 is    0.00%
Random coil      (Cc) :   145 is   45.74%
Ambiguous states (?)  :     0 is    0.00%
Other states          :     0 is    0.00%
```

XP_015163654.1

```
Alpha helix      (Hh) :    76 is   24.13%
3₁₀ helix        (Gg) :     0 is    0.00%
Pi helix         (Ii) :     0 is    0.00%
Beta bridge      (Bb) :     0 is    0.00%
Extended strand  (Ee) :    88 is   27.94%
Beta turn        (Tt) :     0 is    0.00%
Bend region      (Ss) :     0 is    0.00%
Random coil      (Cc) :   151 is   47.94%
Ambiguous states (?)  :     0 is    0.00%
Other states          :     0 is    0.00%
```

XP_009602718.1

```
Alpha helix      (Hh) :    82 is   26.80%
3₁₀ helix        (Gg) :     0 is    0.00%
Pi helix         (Ii) :     0 is    0.00%
Beta bridge      (Bb) :     0 is    0.00%
Extended strand  (Ee) :    74 is   24.18%
Beta turn        (Tt) :     0 is    0.00%
Bend region      (Ss) :     0 is    0.00%
Random coil      (Cc) :   150 is   49.02%
Ambiguous states (?)  :     0 is    0.00%
Other states          :     0 is    0.00%
```

XP_009764644.1

```
Alpha helix      (Hh) :    87 is   26.85%
3₁₀ helix        (Gg) :     0 is    0.00%
Pi helix         (Ii) :     0 is    0.00%
Beta bridge      (Bb) :     0 is    0.00%
Extended strand  (Ee) :    70 is   21.60%
Beta turn        (Tt) :     0 is    0.00%
Bend region      (Ss) :     0 is    0.00%
Random coil      (Cc) :   167 is   51.54%
Ambiguous states (?)  :     0 is    0.00%
Other states          :     0 is    0.00%
```

XP_019251073.1

```
Alpha helix      (Hh) :    77 is   25.25%
3₁₀ helix        (Gg) :     0 is    0.00%
Pi helix         (Ii) :     0 is    0.00%
Beta bridge      (Bb) :     0 is    0.00%
Extended strand  (Ee) :    77 is   25.25%
Beta turn        (Tt) :     0 is    0.00%
Bend region      (Ss) :     0 is    0.00%
Random coil      (Cc) :   151 is   49.51%
Ambiguous states (?)  :     0 is    0.00%
Other states          :     0 is    0.00%
```

XP_016500060.1

```
Alpha helix      (Hh) :    87 is   26.85%
3₁₀ helix        (Gg) :     0 is    0.00%
Pi helix         (Ii) :     0 is    0.00%
Beta bridge      (Bb) :     0 is    0.00%
Extended strand  (Ee) :    70 is   21.60%
Beta turn        (Tt) :     0 is    0.00%
Bend region      (Ss) :     0 is    0.00%
Random coil      (Cc) :   167 is   51.54%
Ambiguous states (?)  :     0 is    0.00%
Other states          :     0 is    0.00%
```

XP_028114495.1

```
Alpha helix      (Hh) :    77 is   24.06%
3₁₀ helix        (Gg) :     0 is    0.00%
Pi helix         (Ii) :     0 is    0.00%
Beta bridge      (Bb) :     0 is    0.00%
Extended strand  (Ee) :    90 is   28.12%
Beta turn        (Tt) :     0 is    0.00%
Bend region      (Ss) :     0 is    0.00%
Random coil      (Cc) :   153 is   47.81%
Ambiguous states (?)  :     0 is    0.00%
Other states          :     0 is    0.00%
```

488   **Supplementary Fig. S2**. Description of secondary structure of ZIP protein homologs.