1 **Force-dependent remodeling of cytoplasmic ZO-1 condensates contributes to**
2 **robust cell-cell adhesion**
3
4
5
6

7 Noriyuki Kinoshita*[1,3], Takamasa S. Yamamoto[1], Naoko Yasue[1], Chiyo Takagi[1],
8 Toshihiko Fujimori[2,3], and Naoto Ueno*[1,3,4,5]
9
10 Running title: Force-dependent remodeling of ZO-1 by phase separation
11
12
13 1: Division of Morphogenesis, Department of Developmental Biology, National Institute
14 for Basic Biology, 38 Nishigonaka, Myodaiji, Okazaki, Aichi, 444-8585, Japan
15
16 2: Division of Embryology, Department of Developmental Biology, National Institute for
17 Basic Biology, 5-1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan
18
19 3: School of Life Science, SOKENDAI (The Graduate University for Advanced Studies),
20 Okazaki, Aichi, 444-8585, Japan
21
22 4. Unit of Quantitative and Imaging Biology, International Research Collaboration Center,
23 National Institutes of Natural Sciences, 38 Nishigonaka, Myodaiji, Okazaki, Aichi,
24 444-8585, Japan
25
26 5: Lead Contact
27
28 * corresponding authors:
29 nueno@nibb.ac.jp (NU), nkinoshi@nibb.ac.jp (NK.)
30

**Summary (149 words)**

Although the physiological importance of biomolecular condensates is widely recognized,

how it is controlled in time and space during development is largely unknown. Here we

show that a tight junction protein ZO-1 forms cytoplasmic condensates in the

trophectoderm (TE) of the mouse embryo before E4.0. These disappear via dissolution,

and ZO-1 accumulates at the cell junction as the blastocyst cavity grows and internal

pressure on TE cells increases. In contrast, this dissolution was less evident in TE cells

attached to the inner cell mass, as they receive weaker tensile forces. Furthermore,

analyses using MDCK cells have demonstrated that the ZO-1 condensates are generated

and maintained by liquid-liquid phase separation. Our study also highlights that the

dynamics of these condensates depends on the physical environment via an interaction

between ZO-1 and F-actin. We propose that the force-dependent regulation of ZO-1

condensation contributes to establishing robust cell-cell adhesion during early

development.

**Keywords**

Cell-to-cell adhesion, ZO-1, tight junction, mechanical force, mouse development,

condensate, liquid-liquid phase separation (LLPS), intrinsically disordered region (IDR).

## Introduction

Mechanical forces are generated at different times and in different tissues during the development of organisms due to dynamic movements and changes in the shape of cells and tissues (Bodor et al., 2020), growth of a cell mass by proliferation (Godard and Heisenberg, 2019), the removal of cells by apoptosis (Teng et al., 2017), luminal pressure (Chan et al., 2019), and shear stress of body fluids such as blood (Paolini and Abdelilah-Seyfried, 2018), etc. All of these forces can contribute to the morphogenesis of organs through mechanochemical feedback mechanisms (Hannezo and Heisenberg, 2019). Although the presence and physiological importance of such forces have long been implicated in the homeostasis as well as morphogenesis of living organisms (Hallou and Brunet, 2020; Thompson, 1917), how cells sense and respond to these forces is not fully understood at the molecular level. This study demonstrates that mechanical forces control the remodeling of ZO-1 through the regulation of liquid-liquid phase separation (LLPS), a physical process that facilitates the demixing of proteins through protein condensation.

In our previous study using *Xenopus laevis* embryos, we demonstrated that among a number of identified phosphoproteins, cell junction- as well as focal adhesion-related proteins are highly phosphorylated immediately after the application of mechanical forces by centrifugation or compression (Hashimoto et al., 2019). Intriguingly, a global analysis of the phosphoproteome suggested that mechanical stimuli induced the embryonic cells to adopt an epithelial state rather than a mesenchymal state, known as the mesenchymal-epithelial transition (MET), which is opposite to the epithelial-mesenchymal transition (EMT), a well-characterized phenomenon in some developmental contexts and in cancer pathogenesis in which cells compromise their cell-to-cell adhesiveness (Baum et al., 2008). In fact, we confirmed that the adherens and tight junction, revealed by the accumulation of C-cadherin (Choi et al., 1990; Ginsberg et al., 1991) and ZO-1 (Anderson et al., 1988), respectively, became enhanced after the application of force. We also found that F-actin, which is normally localized in the basal and lateral domain of early embryonic cells, accumulated at the apical domain, reinforcing the stiffness of the cell cortex (Hashimoto et al., 2019). These observations suggest that these junctions were remodeled following force stimuli and that the cells adopted more epithelialized states. This led us to propose that force-induced epithelialization is an important cellular response to physical forces to maintain tissue

integrity, allowing it to acquire robustness against forces that might perturb normal
morphogenesis.

More recently, we showed that FGFR1 is activated and that Erk2 is consequently
phosphorylated and translocated into the nucleus in ectodermal cells (Kinoshita et al.,
2020) to achieve MET during gastrulation. In that study, we repeatedly observed that
ZO-1 condensates were found as cytoplasmic puncta in ectoderm cells prior to, but not
after, gastrulation (Kinoshita et al., 2020). Based on previous reports that showed that
ZO-1 could undergo LLPS in zebrafish embryos (Schwayer et al., 2019) and cultured
cells (Beutel et al., 2019), we speculated that the ZO-1 LLPS is mechanically regulated,
particularly by dynamic morphogenesis known as epiboly during gastrulation which
imposes a tensile force on the ectoderm (Hernández-Vega et al., 2017).

To extend the above study that employed *Xenopus laevis* and to examine whether our
working model could be extrapolated to other species such as mammals, we focused on
mouse embryogenesis in this study. Basically, we obtained similar results to those with
*Xenopus* and found that cytoplasmic condensates of ZO-1 exist in the trophectoderm (TE)
cells of early mouse embryos before E4.0, although the condensate disappears upon
stretching of TE cells due to the expansion of blastocyst cavity (blastocoel).

To confirm that ZO-1 condensates are the product of cellular LLPS (Brangwynne,
2013), we used cultured cells in which GFP-ZO-1 condensates were observed in various
conditions. Following both the treatment of a cell-permeabilizing agent digitonin (Shiina,
2019) and analyses with fluorescent recovery after photobleaching (FRAP), ZO-1
condensates showed to have typical properties of liquid droplets generated by LLPS. We
also identified an intrinsically disordered region (IDR) (Kato et al., 2012; Li et al., 2012),
an unstructured stretch of an amino acid sequence of low complexity that resides in the
C-terminal half, and showed that it is necessary for phase separation of ZO-1. Finally, we
highlighted the importance of cytoskeletal actin in the regulation of ZO-1 assembly and
proposed that ZO-1 protein became liberated from droplets and was deployed to enhance
the tight junction in a force-dependent manner. This was supported by observations of
MDCK cells in different culture conditions, including wound healing and substrates of
different stiffness.

The present study, using early mouse embryos and cultured cells, reveals a previously
unknown role of physical force that regulates the condensation of ZO-1, which
contributes to the enhancement of cell-to-cell adhesion during early development.

## Results

### *ZO-1 condensates in early mouse embryos*

In our previous report (Kinoshita et al., 2020), we demonstrated that the stretch of ectodermal tissue in the *Xenopus* gastrula embryo induces a MET-like cellular response, including a reduction of cytoplasmic ZO-1 puncta and its accumulation at the cell junction. To examine whether the behavior of ZO-1 protein is conserved across species, especially in mammals, we first immunostained mouse embryo ZO-1. We focused on E3.5 and E4.5 embryos since they hatch out of the zona pellucida (ZP) and expand their shape (Figure 1A). We fixed E3.5 and E4.5 embryos with paraformaldehyde and immunostained them with a ZO-1 antibody (Figure 1B). We found that the cells of E3.5 embryos showed a significantly higher number of ZO-1 puncta in the cytoplasm relative to E4.5 embryos. As development proceeded, the surface area of TE cells, particularly on the mural side, expanded and became thinner (data not shown). We found that the number of cytoplasmic ZO-1 puncta were reduced, and ZO-1 signal intensity at the plasma membrane in E4.5 embryos became much higher than those of E3.5 embryos at the expense of their cytoplasmic pool (Figure 1C, D), without changing the total fluorescence intensity of the protein (Figure 1E left). Importantly, this change coincides well with the accumulation of F-actin at the cell cortex in E4.5 embryos (Figure 1B and 1E right). This result suggests that the shuttling of ZO-1 protein from the cytoplasmic puncta to cell junctions occurs as development progressed. We observed a similar behavior of ZO-1 protein using ZO-1-EGFP-expressing mouse embryos by live-imaging (Katsunuma et al., 2016) (Figure 1F, G and Video S1). These results are consistent with our previous finding, where ZO-1 puncta decreased in *Xenopus* ectodermal cells that were exposed to a higher tensile force (Hashimoto et al., 2019; Kinoshita et al., 2020).

Interestingly, however, in the polar TE cells attached to the inner cell mass (ICM) of E4.5 embryos, a significant number of condensates still remained in the cytoplasm (Figure S1A and B). Consistently, the fluorescent intensity of both anti-ZO-1 immunostaining and F-actin accumulation at the plasma membrane in polar TE cells was lower than in mural cells (Figure S1C). Since polar TE cells attach to the crowded ICM, these cells might be exposed to lower tensions.

### *ZO-1 condensate is regulated by a force-dependent mechanism*

We next examined whether the increased tensile force reduced ZO-1 puncta and

accumulated it to the junction during hatching. It is known that developing mouse

embryos from E4.0 to E4.5 experience a gradually increasing magnitude of luminal

pressure due to expansion of the blastocyst cavity in the presence of the ZP (Chan et al.,

2019; Leonavicius et al., 2018). Accordingly, the cortical tension of TE increases from

E3.5 to E4.5 (Chan et al., 2019). To confirm that the disappearance of ZO-1 condensates

from the cytoplasm is dependent on a mechanical force (inner hydraulic pressure of

blastocyst cavity), using ouabain, we inhibited $Na^+/K^+$ ATPase, which promotes the

influx of water and therefore increases the volume of the blastocyst cavity (Figure 2A and

B). In these ouabain-treated embryos, a significant number of ZO-1 condensates was

retained in the cytoplasm of mural cells of E4.5 embryos (Figures 2C and D),

demonstrating that the cells released from this tension failed to trigger the dissolution of

ZO-1 condensates.

Next, we mechanically reduced tension by releasing the blastocoel fluid of the

ZO-1-EGFP-expressing embryos by piercing embryos with a glass needle. Immediately

after they were pierced, embryos shrank, and membrane localization of ZO-1 was

reduced, although it recovered within several hours after the wound site had healed and

the embryo regained its original shape (Figure 2E, F and Video S2). Immunostaining with

the anti-ZO-1 antibody confirmed that endogenous ZO-1 also behaves similarly,

observing more cytoplasmic puncta in pierced E4.5 embryos than in control E4.5

embryos (Figure 2G-I). Together, these results suggest that the inner pressure of the

growing blastocyst cavity increases the tensile force on TE cells in E4.5 embryos, thereby

reducing the number and volume of ZO-1 puncta in the cytoplasm.

This dynamic behavior of ZO-1 protein led us to test whether the ZO-1 puncta in E3.5

embryos were formed by phase separation, treating embryos with 1,6-hexanediol, which

is known to dissolve the LLPS assembly (Kroschwald et al., 2015; Shulga and Goldfarb,

2003). As shown in Figures 3A and B, 5% 1,6-hexanediol reduced the number of particles

within a few minutes, suggesting that phase separation forms these droplets in the mouse

embryos.

## *LLPS forms ZO-1 droplets in MDCK cells*

In order to analyze the nature of the ZO-1 condensate in various cell environments, we

performed a series of studies using canine MDCK cells. The results using mouse embryos

suggested that tensile force to the TE cells negatively regulates ZO-1 phase separation. In
MDCK cells, it had been demonstrated that increased substrate stiffness increases tension
acting on ZO-1 (Haas et al., 2020). We thus examined whether substrate stiffness affects
ZO-1 droplet formation. We plated MDCK cells on a polyacrylamide gel substrate of
different stiffnesses and immunostained ZO-1. As shown in Figures 4A and B, cells on
the soft substrates formed more cytoplasmic ZO-1 puncta than cells on the rigid
substrates. This result supports our notion that sensing mechanical stimuli, including
through focal adhesion, triggers the remodeling of ZO-1.

We also queried whether ZO-1 localization would change in an interaction with
surrounding cells, which might affect the mechanical environment (Zhu et al., 2000). We
grew MDCK cells to form islands in a culture dish and transfected a plasmid harboring
human ZO-1 fused to GFP (GFP-ZO-1). It is notable that cells in the central region lost
condensates while peripheral cells tended to have condensates (Figure 4C and D).
However, we occasionally observed cells that harbored relatively large condensates in the
central region, but those cells were not well attached to, and were free from, neighboring
cells (data not shown). This tendency was confirmed by immunofluorescence using the
anti-ZO-1 antibody (Figure 4E). Quantitatively, cells in the forefront row (most outer
layer) and the second row had 15 to 20 condensates per cell on average. In contrast, cells
in the inner or central region had only a few condensates (Figure 4F). One feature that
distinguishes the inner cells from peripheral cells is that after cells reach confluency, the
former adopts polygonal shapes and has higher densities than the latter. These results
suggest that the inner cells establish robust cell-to-cell contact by tight and adherens
junctions, which are physically supported by the actin cytoskeleton, and an individual cell
rigidly contacts with surrounding cells, resembling embryonic cells under a tensile force.

To confirm further that the ZO-1 droplets observed in MDCK cells are indeed the
products of LLPS, we treated cells harboring GFP-ZO-1 droplets with digitonin, a
cell-permeabilizing reagent that disrupts the equilibrium between a condensate and the
cytoplasm. That treatment shrank the droplets within several minutes in MDCK cells
(Figure 5A, B and Video S3). Furthermore, we performed FRAP analyses on the
GFP-ZO-1 droplets in MDCK cells (Figure 5C and D). The fluorescent signal of
GFP-ZO-1 droplets recovered by almost 50% within 2 min. These results demonstrate
that the ZO-1 puncta are droplet-like condensates produced by LLPS and are in
equilibrium with the cytoplasm.

It has been reported that junctional ZO-1 is also assembled by the LLPS (Beutel et al.,

2019). We thus conducted a FRAP assay using inner cells with junctional ZO-1 in the

same culture dish (Figure 5E and F) and found that ZO-1 at the junction showed a similar

recovery rate. These results indicate that both ZO-1 in the cytoplasmic droplets and ZO-1

at the junction are liquid-like condensates and that ZO-1 changes its localization

depending on the cellular environment.

*Cell-cell interaction regulates ZO-1 droplet formation*

Our finding that peripheral cells have more droplets than inner cells suggested that robust

cell-cell contact negatively regulates ZO-1 LLPS. To verify this idea, we reduced

cell-to-cell contact by treating MDCK cells expressing GFP-ZO-1 with PBS containing

trypsin-EDTA. As expected, when trypsin-EDTA solution was added, ZO-1 detached

from the cell membrane and formed puncta as cells lost their contact (Figure 5G and

Video S4).

The robustness of cell-cell contact was also examined by laser ablation. MDCK cells

were grown so that they formed islands in one culture dish. After the GFP-ZO-1

expression plasmid was transfected, cells were irradiated with laser light. When cells

inside the colonies were irradiated at the cell junction, cells stably stayed after irradiation

due to cell-cell contact with neighboring cells (Figure S2A, C and Video S5). When

peripherally-located cells with ZO-1 puncta were irradiated, those cells immediately

erupted and disappeared after irradiation (Figure S2B, C, and Video S6), indicating that

those cells are weakly attached to surrounding cells. These results suggest that in MDCK

cells, the formation of ZO-1 condensate is negatively associated with the establishment of

robust cell-to-cell contact.

*ZO-1 forms cytoplasmic condensates in migrating cells during wound healing*

To further examine whether cell-to-cell contact negatively regulates ZO-1 condensate

formation in the cytoplasm of MDCK cells, we conducted a wound-healing assay using

GFP-ZO-1–expressing cells (Figure 6A and B and Video S7). Initially, most of the cells

had few condensates in a confluent cell sheet. After scratching, cells started active

migration toward the wound site, forming a significant number of droplets. By longer

incubation in this assay, cells migrating from both sides eventually met in the middle of

the gap, re-establishing a new cell-cell interaction. During this process, ZO-1 droplets

disappeared and accumulated at the cell junction (Figure S3 and Video S8). These results

indicate that ZO-1 condensates shuttle between cytoplasmic droplets and the cell junction,

depending on cell-cell interaction.

The FRAP assay revealed that the fluorescent signals of GFP-ZO-1 condensates that

had formed during the wound-healing assay recovered to about 40% within 10 min,

suggesting that these condensates were formed by LLPS (Figure 6C-E, Video S9, and

S10). This result again indicates that the formation of ZO-1 condensate is regulated by

intercellular and extracellular environments such as cell-to-cell contact.

In the wound-healing assay, we found that the distribution and intensity of F-actin

were different between inner cells and edge cells. As shown in Figures 6F and G, cortical

F-actin was well-developed in the non-migrating polygonal cells with junctional ZO-1

("inner" cells in Figure 6F). In contrast, migrating cells around the wound site, which

form ZO-1 droplets, had reduced cortical actin and developed more stress fibers. This

correlation between ZO-1 and F-actin is consistent with our previous finding in which the

application of force to *Xenopus* embryonic cells induced a MET-like response, including

the accumulation of cortical F-actin, and reduced ZO-1 condensates as cell-cell junctions

were enhanced (Hashimoto et al., 2019 ; Kinoshita et al., 2020).

*Interaction with F-actin regulates ZO-1 phase separation*

We further investigated the relationship between ZO-1 phase separation and F-actin

organization by expressing GFP-ZO-1 in non-confluent *Xenopus* A6 cells. In A6 cells,

GFP-ZO-1 localized mostly in the cell periphery and colocalized with F-actin bundles

(Figure 7A). After treating the cells with latrunculin B, a potent inhibitor of actin

polymerization, ZO-1 granules rapidly emerged in the cytoplasm, which disappeared

soon after washing out the drug (Figure 7B and Video S11). These results suggest that the

ZO-1 granules, which were highly mobile in the FRAP assay (Figure 7C and D), are

negatively regulated by the interaction with F-actin in A6 cells. We also treated MDCK

cells expressing GFP-ZO-1 with latrunculin B, but no detectable change in the

localization of ZO-1 was observed (data not shown). Therefore, we speculate that the

contribution of F-actin to suppress cytoplasmic ZO-1 condensation is more significant in

A6 cells than in MDCK cells, which develop more rigid junctional structures.

To understand the F-actin-driven mechanism, we constructed GFP-ZO-1ΔABD

(Figure 7E), which lacks the actin-binding domain (ABD) (Fanning et al., 2002), and

expressed it in A6 cells. As shown in Figure 7F, GFP-ZO-1ΔABD formed droplets with a
smooth surface. This suggests that cytoplasmic droplet formation is restricted by the
binding of ZO-1 to F-actin (Beutel et al., 2019; Schwayer et al., 2019) and that
GFP-ZO-1ΔABD lost its capacity for the interaction. Supporting this notion,
co-expression with a red fluorescent protein fused to F-actin-binding peptides, Lifeact,
Utrophin, and the actin-binding domain of Moesin (MoesinABD), similarly induced
condensates that were indistinguishable from those induced by GFP-ZO-1ΔABD (Figure
S3A). It is likely that the expression of these actin-binding proteins competes binding to
F-actin with ZO-1.
We also observed that the MoesinABD-induced ZO-1 condensates moved dynamically,
sometimes fused and underwent fission (Figure S4B and Video S12), reminiscent of
typical liquid droplets. In addition, the size of these droplets was rapidly reduced by cell
permeabilization following digitonin treatment (Figure S4C and Video S13) and
exhibited a rapid recovery of fluorescence in the FRAP assay (Figure S4D). These results
confirm that these cytoplasmic droplets are formed by phase separation.
297

***The C-terminal intrinsically disordered region is required for ZO-1 phase separation***
It is known that intrinsically disordered regions (IDRs) are required for proteins to
undergo phase separation (Kato et al., 2012; Li et al., 2012). Four IDRs are predicted in
human ZO-1 (Figure S5A) using the Protein DisOrder prediction System
(https://prdos.hgc.jp/cgi-bin/top.cgi) (Ishida and Kinoshita, 2007). In order to address
which IDR is critical for ZO-1 in A6 cells, we expressed a series of deletion mutants
(Figure S5B). As a positive control, we expressed GFP-ZO1-ΔABD and confirmed that it
formed droplets. Then, we deleted IDR1, IDR2, and IDR3, which reside in the N-terminal
half, from GFP-ZO1-ΔABD (GFP-ZO1-ΔABD-ΔIDR123). When it was expressed, it
also formed droplets (Figure S5C), indicating that these regions are not required for phase
separation in A6 cells. In contrast, deletion of the C-terminal IDR4
(GFP-ZO1-ΔABD-ΔIDR4) abolished ZO-1 droplets, and it became dispersed in the
cytoplasm. The result demonstrates that IDR4 is critical for cytoplasmic liquid droplet
formation by LLPS. Since the actin-binding domain overlaps with IDR4, it is possible
that F-actin binding may affect the LLPS-inducing activity of IDR4.
313

**Discussion**

Our present study using mouse embryos as well as cultured cells, together with a previous

study that employed *Xenopus* embryos (Kinoshita et al., 2020), indicates that similar

mechanisms may govern cellular responses to mechanical forces across species. Briefly,

mechanical stresses caused by dynamic morphogenesis and/or vigorous cell movements

during development, such as gastrulation, activate the force-dependent pathway leading

to the remodeling of cell-to-cell junctions. The present study also suggests that the

process involves the negative regulation of cytoplasmic ZO-1 condensate formation by

LLPS, leading to its translocation to the cytoplasm. Since embryonic tissues, as well as

some adult tissues are constantly exposed to various forces, and since ZO-1 is a

structurally conserved protein among species, we propose that this mechanism may be

employed to maintain the robustness and integrity of tissues against various forces in

general, and might also have been an evolutionary innovation when multicellular animal

species such as placozoa emerged (González-Mariscal et al., 2011). Although our

previous study using *Xenopus* embryos identified that the function of the FGF receptor is

essential for the force-induced pathway, it also suggested that FGF ligands are not

required for signal activation, shedding light onto the as yet unrevealed activation

mechanisms for FGFRs and other receptor tyrosine kinases (RTKs) by force. In fact, the

epidermal growth factor receptor (EGFR), which is an upstream RTK for Erk, was

implicated in the mechanical force-dependent activation of Erk in the collective

migration of wound-healing cells (Hino et al., 2020). In both cases, cytoskeletal

remodeling by tensile forces appears to trigger these pathways, supporting the idea that

mechanoresponsive signaling pathways may be highly conserved.

Interestingly, our present study using cultured cells demonstrated that the efficacy of

ZO-1 condensation as droplets largely depends on cell type. In MDCK cells, cytoplasmic

ZO-1 condensates are formed when they are cultured sparsely, or cell-cell interaction is

weakened, for example, at the periphery of the colonies, during the wound healing

process, or by the trypsin/EDTA treatment. In contrast, in cells with a robust cell-cell

interaction, ZO-1 condensates appear to be localized at the cell junction. In the case of A6

cells, blocking the interaction between ZO-1 and F-actin induces ZO-1 droplet formation,

suggesting that ZO-1 droplet formation is negatively regulated by the interaction with

F-actin. This suggests that the difference in the amount or dynamics of F-actin may

distinguish these two cell types and the revelation of ZO-1 from F-actin is critical for

maintaining large-sized ZO-1 condensate.

It is tempting to speculate that Erk might regulate F-actin polymerization through
small G proteins such as GEF-H1 and its downstream target RhoA (Fujishiro et al., 2008;
Itoh et al., 2014; Ren et al., 1998). Recently, Haas et al. (2020) reported that substrate
stiffness regulates GEF-H1, thereby increasing tension, by acting on ZO-1. Since we
reported in this study that substrate stiffness changes ZO-1 droplet formation, it would be
intriguing to investigate whether the GEF-H1 pathway is involved in the regulation of
ZO-1 phase separation. Second, as described above, it was reported that Erk is activated
in collectively migrating MDCK cells by EGFR and regulates F-actin organization and
cellular contraction (Hino et al., 2020). It is also possible that the cascade of
phosphorylation initiated by RTKs might eventually result in the remodeling of F-actin
and ZO-1 localization.

It has been reported that the assembly of ZO-1 at the cell junction is also driven by
phase separation (Beutel et al., 2019). We also confirmed that both the cell junction and
the cytoplasmic ZO-1 granules show typical characteristics of phase-separated
condensates, which suggests that ZO-1 localization may be dynamically switched
depending on the mechanical environments and F-actin plays a key role in the switching
in A6 cells. We also mapped the region essential for cytoplasmic granule formation to the
fourth IDR (IDR4 in Figure S4). As IDR4 covers the actin-binding domain, the
interaction with F-actin may directly affect its phase separation. It has been reported that
F-actin binding to ZO-1 induces its conformational change (Spadaro et al., 2017). Such
structural changes may regulate the phase separation of IDR4 to form cytoplasmic
droplets. Beutel et al. (2019) demonstrated that the PDZ-SH3-GuK domain in ZO-1 plays
an essential role in phase separation at the cell junction. Since ZO-1 is a scaffold protein
that interacts with many proteins (Fanning and Anderson, 2009), such a protein-protein
interaction may also change the state of phase separation, resulting in the difference
between the cytoplasmic and junctional forms of condensates. Investigating the precise
molecular mechanisms that could explain the two distinct states of ZO-1 will be
important.

It has been reported that tension mediates the remodeling of cell junctions (Ito et al.,
2017). Phase separation of ZO-1 may be directly involved in this regulatory system. In
this scenario, LLPS may act as an on-demand ON/OFF switch of ZO-1 supply to the cell
membrane from the cytoplasm, and this mechanism may be shared and employed by
multicellular animal species. Therefore, its evolutionary origin is an intriguing problem

worth studying in the future.

ZO-1 is implicated in various pathologies, including cancer. Particularly, in EMT, the

loss of cell-to-cell adhesion is known to drive the progress and metastasis of cancer. It is

also known that in neuronal cells, some condensates which contain proteins, such as the

RNA-binding protein FUS (Murray et al., 2017) and tau implicated in Alzheimer's

disease (Wegmann et al., 2018), become aggregated depending possibly on their

concentrations, sometime after their formation and/or their chemical modifications.

Those proteins are irreversibly aggregated and become dysfunctional, attenuating the

neuronal activity of the cells. Collectively considering these observations, we speculate

that impairment of the normal regulation of the LLPS of ZO-1 due to changes in

intracellular or extracellular conditions could be a possible cause of cancer pathology.

Therefore, in addition to embryogenesis, investigating the behavior of ZO-1 in adult

tissues, especially in pathological conditions, would deepen our understanding of the

physiological significance of the ZO-1 reservoir in cells.

**Limitations of this study**

We propose in this paper that F-actin remodeling is one of the key factors to regulate

ZO-1 shuttling between cytoplasmic and junctional condensates. However, particularly

in MDCK cells, disruption of F-actin did not change junctional ZO-1 localization,

suggesting that F-actin remodeling may not be sufficient for re-localizing ZO-1 in

MDCK cells. Since reducing cell-cell contact-induced cytoplasmic ZO-1 droplet

formation, a mechanism that releases junctional ZO-1 to the cytoplasm must exist. The

interactions with their junctional components, such as claudin, occludin, JAM-A, and

cingulin (Fanning and Anderson, 2009; Haas et al., 2020; Vasileva et al., 2017) remains to

be clarified.

**Acknowledgments**

R26-ZO-1-GFP mice were provided by LARGE, BDR (accession # CDB026K), RIKEN,

Japan. We thank Dr. Fumio Matsuzaki, RIKEN, for human ZO-1 cDNA, Dr. Makoto

Suzuki, Hiroshima University, for the GFP-ZO-1/pCS2 construct, and Dr. Lance

Davidson, University of Pittsburgh, for the mCherry-Utrophin construct. We thank Dr.

Yuko Mimori-Kiyosue, RIKEN, for A6 cells and Dr. Mitsuru Nishita, Fukushima

Medical University, and Dr. Kensaku Mizuno, Tohoku University, for MDCK I cells. We

thank the BioImaging Facility, Core Research Facilities of the National Institute for
Biology (NIBB), for technical support for live imaging and FRAP analysis. We also
thank Dr. Nobuyuki Shiina, NIBB, and Dr. Michael Levine, Princeton University, for
technical advice and valuable discussion on LLPS, and insightful comments on this
work. This research was supported by JSPS KAKENHI 20K06663 to NK, JSPS
KAKENHI 17H03689 and 16H06280 to TF, and MEXT KAKENHI 22127007 and JSPS
KAKENHI 15H05865 to NU. TF and NU were also supported by JST CREST
JPMJCR1654 and Joint Research of the Exploratory Research Center on Life and
Living Systems (ExCELLS), respectively.

**Authors' Contributions**

N.K., T.S.Y., N.Y. and C.T. conducted experiments using tissue culture cells; T.F., N.K.,
and N.Y. conducted experiments using mouse embryos; N.K. analyzed the data and
co-wrote the manuscript. N.U. supervised the research, reviewed the data and
conclusions, and wrote the manuscript.

**Declaration of Interests**

The authors declare no competing interests.

 **Figure legends**

**Figure 1. Change of ZO-1 localization in mouse hatching embryos.**

**A**. Schematic diagram of the hatching process of the mouse embryo. At E3.5, the

embryo is covered with the zona pellucida (ZP). During hatching, the embryo is

enlarged with an expansion of the trophectoderm (TE), and emerges from the ZP. **B**.

Immunofluorescence of mouse E3.5 and E4.5 embryos. Embryos were stained with an

anti-ZO-1 antibody, Hoechst 33342, and Alexa Fluor 546 phalloidin. Scale bar, 20 μm.

The inset in the E3.5 phalloidin image was acquired with higher laser power,

demonstrating that the structure of cortical F-actin is formed at E3.5 even though the

signal intensity was weaker than that in the E4.5 embryo. The bottom panels are

enlarged images from the upper panels indicated by the dashed squares B' and B". **C**.

The number of particles in one TE cell was counted. **D**. The signal intensities of ZO-1 at

the cell periphery of the TE at E3.5 and E4.5 were quantified. **E**. Fluorescence intensity

of whole embryos stained with the anti-ZO-1 antibody and phalloidin was measured.

For comparison, intensities were normalized by Hoechst staining intensity. n = 5

embryos. n.s.: not significant. *p < 0.01 (*t*-test). **F**. Embryos from mice expressing

ZO-1-EGFP by EGFP-knock-in were observed. Before hatching (E3.5), ZO-1 puncta

were observed around the cell periphery, but cell-cell boundaries were unclear. After

hatching (E4.5), membrane localization of ZO-1-GFP became evident. **G**. Fluorescence

intensity around cell boundaries in ZO-1-EGFP–expressing embryos were quantified.

After hatching, ZO-1-EGFP localized to the plasma membrane and formed sharp

boundaries with ZO-1-EGFP. Scale bars, 20 μm.

**Figure 2. Expansion of TE regulates ZO-1 localization.**

**A and B**. Embryos were obtained at E3.5 and cultured for 24 h in the absence (A) or

presence of 0.3 mM ouabain (+Ob) (B). Embryos were immunostained with the

anti-ZO-1 antibody (left), and the images were analyzed using particle analysis in Image

J (right). **C and D**. The percentage of the particle area (C) and the number of particles

(D) of E4.5 mural TE cells were quantified. **E**. E4.5 mouse embryos expressing

ZO-1-GFP were pierced with a glass needle. Pierced embryos shrank immediately, and

the recovery process was imaged. **F**. The fluorescence intensity of GFP-ZO-1 at the cell

periphery at 0 h and 6 h were quantified. The X-axis indicates distance from the cell

membrane. *p < 0.01 (*t*-test). **G**. Mouse E4.5 embryos expressing ZO-1-GFP were

pierced with a glass needle, then the volume of the blastocyst cavity was reduced, and

embryos shrank. Those embryos were fixed and stained with the anti-ZO-1 antibody

and fluorescent phalloidin. **H**. The area of particles and cytoplasm of one mural TE cell

were measured by Image J software, and the ratio of those values was plotted. **I**. The

number of particles was counted using Image J particle analysis. *p < 0.01 (*t*-test).

Scale bars, 20 μm.

**Figure 3. 1,6-Hexanediol treatment dissolved ZO-1 droplets in the mouse E3.5**

**embryo.**

**A**. Mouse E3.5 embryos were treated with 5 % 1,6-hexanediol. The images were

analyzed by particle analysis in Image J (right panels). **B**. The number of particles was

counted. n = 5 (hexanediol) and 7 (control) embryos.

**Figure 4. 1,6-Hexanediol treatment dissolved ZO-1 droplets in the mouse E3.5**

**embryo.**

**A**. MDCK cells were grown on glass or on a polyacrylamide gel with 303, 138, and 19

kPa elasticity. **B**. The ratio of the particle area in each cell was counted by Particle

Analysis in Image J. **C** and **D**. GFP-ZO-1 was expressed in MDCK cells, forming

islands. In cells located inside the islands (indicated by arrowheads), GFP-ZO-1 was

localized mainly at the plasma membrane. In contrast, in cells at the periphery or

outside of the colony (indicated by arrows), GFP-ZO-1 formed puncta. **E**. MDCK cells

that formed a colony were immunostained with the anti-ZO-1 antibody. Images were

analyzed using Particle Analysis in Image J. **F**. Cells in the island were divided into

three groups with respect to their location, namely the first and second rows from the

edge and the central region of the island. The number of particles in each cell was

counted. n = 20 cells. *p < 0.01 (*t*-test).

**Figure 5. Dynamics of ZO-1 puncta in MDCK cells.**

**A** and **B.** MDCK cells expressing GFP-ZO-1 were semi-permeabilized with 0.012%

digitonin. The size of puncta was quantified with Image J. n = 12 puncta. **C** - **F**.

Fluorescent recovery after a photobleaching (FRAP) assay. **C** and **D**. The recovery of

fluorescence of puncta was quantified. n = 20 puncta. **E** and **F**. The recovery of

junctional ZO-1 was quantified. **G**. MDCK cells were treated with trypsin-EDTA solution. As cells lost their cell-cell contact and their shape became spherical, ZO-1 detached from the plasma membrane and formed puncta. Scale bars, 20 μm.

**Figure 6. Wound-healing assay using MDCK cells expressing GFP-ZO-1.**

**A** and **B**. The confluent cell sheet was scratched and observed. Immediately after scratching (0 h), GFP-ZO-1 was at the cell periphery, whereas after 3 hours (3 h), it formed puncta in many of the cells close to the scratch (within 200 μm). The right panels were magnified from #1 and #2 cells in **A**. Scale bar, 50 μm. **B**. The number of cells with GFP-ZO-1 particles was counted. Cells located within 200 μm from the scratch site were defined as "edge" cells. Cells more than 200 μm away from the scratch site were defined as "inner" cells. Data were obtained from three experiments. **C-E**. Wound-healing assay of MDCK cells expressing GFP-ZO-1 followed by a FRAP assay. Cells formed GFP-ZO-1 puncta in the wound-healing assay were subjected to the FRAP assay. Three cells were analyzed. Scale bars, 20 μm. **F**. Inner and edge cells were fixed 10 h after scratching and stained with phalloidin. Arrow indicates the wound site: scale bar, 20 μm. **G**. Fluorescence intensity of phalloidin staining at the plasma membrane was measured. *$p < 0.01$ (*t*-test).

**Figure 7. GFP-ZO-1 binding to F-actin in *Xenopus* A6 cells.**

**A**. GFP ZO-1 was transiently expressed in *Xenopus laevis* A6 kidney cells. Cells were fixed and stained with fluorescent phalloidin. **B**. A6 cells expressing GFP-ZO-1 were treated with 0.5 μM latrunculin B for 20 min, then washed out. **C**. A6 cells were treated with 0.5 μM latrunculin B for 30 min. Then, a FRAP assay was performed. Granules, indicated by arrowheads, were bleached. **D**. The recovery rate was quantified (n = 14). **E**. Domain structure of ZO-1 protein and expression constructs used here: GFP-tagged full-length human ZO-1 (GFP-ZO-1), actin-binding domain (ABD)-deletion mutant (GFP-ZO-1ΔABD). **F**. Localization of GFP-ZO-1 and GFP-ZO-1ΔABD. Scale bars, 20 μm.

**Supplementary Figure and Video Legends**

**Figure S1. Comparison of ZO-1 and F-actin localization between polar and mural TE cells.**

**A**. The nuclear-dense inner cell mass (ICM) region was determined by Hoechst33342
staining. 'Polar' indicates TE attaching ICM, and 'mural' includes TE not attaching the
ICM. The upper images are the same as in Figure 1B right images (E4.5). The indicated
areas of the middle and the lower images were magnified. Scale bars, 20 μm. **B**. The
number of particles in one TE cell was counted. **C**. Signal intensities of ZO-1 and
phalloidin at the cell periphery of the mural and polar TE were quantified. 4 - 5 embryos
were analyzed for each condition.
540

**Figure S2. Laser ablation of MDCK cells expressing GFP-ZO-1.**
**A**. Cells inside an island with GFP-ZO-1 around the cell membrane. About 80% of
these cells remained stable after ablation (Video S7). **B**. A cell located close to the edge
of an island had GFP-ZO-1 puncta. About 80% of these cells rapidly erupted and
detached from the bottom of the dish (Video S8). Arrowheads indicate laser ablation
sites. **C**. A statistical summary of the laser ablation experiment. "attached" indicates
cells that stayed attached to the dish after ablation, as shown in **A**. "erupted" indicates
cells that erupted after ablation and detached from the dish, as shown in **B**. Data were
obtained from two experiments. The total number of cells for the laser ablation were 26
inside and 24 outside cells. Scale bars, 20 μm.
551

**Figure S3. Shuttling GFP-ZO-1 between droplets and cell junction.**
In the wound healing assay, GFP-ZO-1 was initially localized at the cell junction (left).
When cells started the migration, GFP-ZO1 formed droplets (middle). When the wound
gap was filled, and a new cell-cell interaction was established, GFP-ZO-1 was
re-localized to the cell junction (right). White lines indicate the wound site and yellow
lines indicate the new cell-cell contact site. BF: bright field. Scale bar, 20 μm.
558

**Figure S4. Dynamics of GFP-ZO-1 in A6 cells.**
**A**. GFP-ZO-1 was co-expressed with Lifeact-RFP, mCherry-Utrophin, and
RFP-MoesinABD (actin-binding domain). **B**. Dynamics of GFP-ZO-1 droplets induced
by co-expression with MoesinABD. Droplets often underwent fission and fusion, as
indicated by arrowheads. These images are from Video S3. **C**. A6 cells expressing
GFP-ZO-1 were semi-permeabilized with 0.012% digitonin. The size of puncta was
quantified with Image J. **D**. Fluorescent recovery after a photobleaching (FRAP) assay

of GFP-ZO-1 droplets. n = 20 puncta. Scale bars, 20 μm.

**Figure S5. Dynamics of GFP-ZO-1 in A6 cells.**

**A**. IDRs (intrinsically disordered regions) of ZO-1 were predicted by the Protein

Disorder Prediction System (http://prodos.hgc.jp/cgi-bin/top.cgi). Four IDRs were

designated as IDR1-4. **B**. IDR deletion mutants were constructed. **C**. These mutants

were expressed in A6 cells. Scale bar, 20 μm.

**Video S1.** Mouse embryos expressing ZO-1-EGFP were flushed from uteri at one-day

post-coitum and observed by time-lapse microscopy for 92 h (related to Figure S1).

**Video S2**. Mouse E4.5 embryos expressing ZO-1-EGFP were pierced by a glass needle

and observed by time-lapse microscopy (related to Figure 2F).

**Video S3.** An MDCK cell expressing GFP-ZO-1 was treated with digitonin

("+digitonin" at 50 sec in the video) and imaged by time-lapse microscopy (related to

Figure 4D).

**Video S4.** MDCK cells expressing GFP-ZO-1 were treated with a trypsin-EDTA

solution. The video starts immediately after treatment (related to Figure S4).

**Video S5.** Laser ablation of MDCK cells expressing GFP-ZO-1 inside the island. The

laser was irradiated at 4 sec at the membrane indicated by arrowheads (related to Figure

5E).

**Video S6.** Laser ablation of MDCK cells expressing GFP-ZO-1 at the periphery of the

island. The laser was irradiated at 4 sec at the membrane indicated by the arrowhead.

The cell erupted and disappeared immediately after irradiation (related to Figure 5F).

**Video S7.** A wound-healing assay of MDCK cells. A sheet of MDCK cells transfected

with the GFP-ZO-1 construct was scratched and observed by time-lapse microscopy

(related to Figure 6).

**Video S8.** Shuttling GFP-ZO-1 between droplets and cell junction. In the wound
healing assay, GFP-ZO-1 was initially localized at the cell junction. When cells started
the migration, GFP-ZO1 formed droplets. When the wound gap was filled, and new
cell-cell interaction was established, GFP-ZO-1 was re-localized to the cell junction.
603
604 **Video S9.** A wound-healing assay of MDCK cells, followed by a FRAP assay (1). A
605 sheet of MDCK cells transfected with the GFP-ZO-1 construct was scratched and
606 observed by time-lapse microscopy (related to Figure S5A). The cell for the following
607 FRAP assay was indicated by an arrowhead at the first and last frames.
608
609 **Video S10.** A wound-healing assay of MDCK cells, followed by a FRAP assay (2). The
610 FRAP assay was conducted using a cell in Video S11, which formed ZO-1 puncta
611 (related to Figure S5B). Arrowhead indicates the granule bleached in the FRAP assay.
612
613 **Video S11.** Latrunculin B treatment of an A6 cell co-expressing GFP-ZO-1 and
614 RFP-MoesinABD (related to Figure 3F).
615
616 **Video S12.** An A6 cell co-expressing GFP-ZO-1 and RFP-MoesinABD was imaged by
617 time-lapse microscopy for 885 min (related to Figure S3).
618
619 **Video S13.** A6 cells co-expressing GFP-ZO-1 and RFP-MoesinABD were treated with
620 digitonin ("+digitonin" at 50 sec in the video) and imaged by time-lapse microscopy
621 (related to Figure 4C).
622
623

624    **STAR★Methods**

625    **KEY RESOURCES TABLE**

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---|---|---|
| Antibodies | | |
| anti-ZO-1 antibody | Thermo Fisher Scientific | cat. # 33-9100; RRID:AB_2533147 |
| goat anti-mouse IgG antibody, Alexa Fluor 488 | Thermo Fisher Scientific | cat. # A28175; RRID:AB_2536161 |
| Chemicals, Peptides, and Recombinant Proteins | | |
| Blocking One | Nacalai Tesque | cat.# 03953 |
| KSOM | Millipore | cat. # MR-106D |
| ouabain | Sigma | cat.# O3125 |
| Alexa Fluor 546 Phalloidin | Thermo Fisher Scientific | cat.# A22283; RRID:AB_2632953 |
| Hoechst 33342 | Thermo Fisher Scientific | cat.# H3570 |
| Lipofectamine 2000 | Invitrogen | cat. # 11668-027 |
| Effectene | Qiagen | cat. # 301425 |
| Latrunculin B | Enzo Life Science | cat. # BML-T110-0001 |
| digitonin | WAKO | cat. # 048-02124 |
| trypsin | Nacalai Tesque | cat # 35555-54 |
| 3-aminopropyl) triethoxysilane | Sigma-Aldrich | cat. # 440140 |
| Sulfo-SANPAH | Pierce | cat. # 22589 |
| 1.6-hexanediol | WAKO | cat. # 087-00432 |
| Experimental Models: Organisms/Strains | | |
| *Xenopus laevis* (J-strain) | N/A | N/A |
| R26-ZO-1EGFP | RIKEN | accession # CDB026K |
| wild type ICR mice | Japan CREA | N/A |
| Experimental Models: Cell Lines | | |
| MDCK I cells | Fujiwara et al. (2016) | N/A |
| A6 cells | Mimori-Kiyosue et al. (2007) | N/A |

| Recombinant DNA | | |
|---|---|---|
| human ZO-1 | Konno et al. (2008) | N/A |
| GFP-ZO-1/pCS2 | This paper | N/A |
| GFP-ZO-1ΔABD/pCS2 (Δ1151-1371) | This paper | N/A |
| GFP-ZO-1ΔABD-ΔIDR123/pCS2 (Δ114-118, Δ275-415, Δ588-627, Δ1151-1371) | This paper | N/A |
| GFP-ZO-1ΔIDR4/pCS2 (Δ858-1580) | This paper | N/A |
| GFP-ZO-1-PSG/pCS2 (424-794) | This paper | N/A |
| GFP-ZO-1ΔABDΔPSG/pCS2 (Δ424--794, Δ1151-1371) | This paper | N/A |
| Lifeact-RFP | Iioka et al. (2007) | N/A |
| RFP-MoesinABD | Iioka et al. (2004) | N/A |
| mCherry-Utrophin | Kim and Davidson (2013) | N/A |
| Software and Algorithms | | |
| Fiji | NIH | https://fiji.sc |
| Python 3.6 | Python Core Team | https://www.python.org |
| Pandas | v0.22 | http://pandas.pydata.org |
| seaborn | v0.8 | https://seaborn.pydata.org |
| Matplotlib | v2.2 | https://matplotlib.org |
| Andor IQ2 | Oxford Instruments | https://andor.oxinst.com/products/iq-live-cell-imaging-software/ |
| PrDOS (protein disorder prediction system) | (Ishida and Kinoshita, 2007) | http://prdos.hgc.jp/cgi-bin/top.cgi |

**Lead Contact and Materials Availability**

Further information and requests for resources and reagents should be directed to and

will be fulfilled by the Lead Contact, Naoto Ueno (nueno@nibb.ac.jp). All

unique/stable reagents generated in this study are available from the Lead Contact with

a completed Material Transfer Agreement.

**Method details**

**Mouse embryo collection**

Animal care and experiments were conducted in accordance with the Guidelines of

Animal Experimentation of the National Institutes for Natural Sciences. All animal

experiments were approved by the Animal Research Committee of the National

Institutes for Natural Sciences. Mice were maintained in a light- and

temperature-controlled room using a 12 h light:12 h dark cycle at 23+/-2 °C.

Males of R26-ZO-1EGFP (Katsunuma et al., 2016) or wild-type ICR (Japan CREA)

were mated with wild-type ICR females to obtain blastocysts. Preimplantation E3.5

embryos were flushed from uteri with KSOM (MR-106D, Millipore). Embryos were

cultured in KSOM covered with mineral oil at 37 °C and 5% $CO_2$. Fluorescent signals

were monitored using a confocal microscope (A1, Nikon) or a spinning disc confocal

microscope (CV1000, Yokogawa, Japan). For immunofluorescence, embryos were fixed

at 4 °C in 4% paraformaldehyde (PFA) in PBS overnight. For ouabain treatment, 10

mM ouabain (cat. # O3125, Sigma) in DMSO was diluted to the indicated

concentrations.

**Manipulation of mouse embryos**

To remove the zona pellucida, embryos were treated with Tyrode's solution (Bradley,

1987). Embryos were transferred to a drop of Tyrode's solution. The zona was removed

typically within 1 min. Then, embryos were washed three times in KSOM. To pierce the

mural trophectoderm, microinjection needles were made from a 1 mm diameter glass

capillary (GD1, Narishige, Japan), the needle was set to a micromanipulator (Leica),

and embryos were pierced with the needle by manual manipulation.

For hexanediol treatment, 1.6-hexanediol (cat. # 087-00432, WAKO, Japan) was

dissolved in water at 50% w/v, and a 1/10 volume was added to the culture medium.

Fluorescent images were captured using CV1000 (Yokogawa, Japan).

**Immunofluorescence with mouse embryos**

Fixed embryos were washed three times with 1% bovine serum albumin (BSA) in PBS

and then soaked in a blocking solution, 0.1% Triton X-100 in Blocking One (cat. #

03953, Nacalai, Japan), for 1 h at room temperature. Embryos were then incubated in

primary antibody solution containing an anti-ZO-1 antibody (cat. # 33-9100, Thermo)

diluted 100-times in Blocking One at 4 °C overnight. After washing three times with the

blocking solution, embryos were incubated in a secondary antibody solution containing

100-times diluted goat anti-mouse IgG antibody Alexa Flour 488 (cat. # A27185,

Thermo), 100-times diluted Alexa Fluor 546 Phalloidin (cat. # A22283, Thermo Fisher),

and 10 µg/ml Hoechst 33342 (cat. # H3570, Thermo Fisher) in Blocking One.

Fluorescent signals were monitored using Nikon A1 or Leica SP8 confocal

microscopes.

**Measurement of fluorescent intensities and particle analyses**

Fiji/Image J was used to measure fluorescent intensities. Particle analyses in Fiji/Image

J were used to count particles and measure particle areas. A large mass signal in the

cavity was removed for quantitative analyses. Images were inverted and converted to an

8-bit type. Cytoplasmic regions were manually determined. The "Analyze Particles"

program was run with the threshold signal value: 230, circularity: 0.3-1.0, and size: 3 -

infinity.

**GFP-ZO-1 expression constructs**

Human ZO-1 cDNA was a gift from Dr. Fumio Matsuzaki (Konno et al., 2008).

EGFP-human ZO-1/pCS2 was constructed by Dr. Makoto Suzuki. ZO-1ΔABD lacks

amino acid (aa) # 1151-1371. Lifeact-RFP is described in Iioka et al. (2007).

MoesinABD is described in Iioka et al. (2004). The mCherry-Utrophin construct was a

gift from Lance A. Davidson (Kim and Davidson, 2013).

**Cell culture and transfection of plasmids**

A6 cells, established from a normal *X. laevis* kidney, were a gift from Dr. Yuko

Mimori-Kiyosue (Mimori-Kiyosue et al., 2007). A6 cells were grown at 24 °C without

$CO_2$ in Leibovitz's L-15 medium (50% L-15 medium, 10% fetal bovine serum (FBS),

200 mg/l kanamycin). MDCK cells were a gift from Dr. Mitsuru Nishita and Dr.

Kensaku Mizuno. MDCK cells were cultured in Dulbecco's modified Eagle medium

(DMEM) containing 10% FBS. Transfection was conducted using Lipofectamine 2000

(Invitrogen # 11668-027) for MDCK cells and Effectene (Qiagen # 301425) for A6 cells

following the manufacturers' instructions.

**Immunofluorescence of the tissue culture cells**

Cells were fixed in 4% PFA in PBS at 4 °C overnight, washed with 0.1% Triton X-100

in PBS. Blocking One (cat. # 03953, Nacalai Tesque, Japan) was used to block for 1 h at

room temperature, followed by incubation in the primary antibody solution, a 100-times

diluted anti-ZO-1 antibody (cat. # 33-9100, Thermo) in Blocking One at 4 °C overnight.

After washing three times in Blocking One, cells were incubated in the secondary

antibody solution containing 200-times diluted goat anti-mouse IgG antibody Alexa

Fluor 488 (cat. # A27185, Thermo Fisher) and 200-times diluted Alexa Fluor 546

Phalloidin (cat. # A22283, Thermo Fisher) in Blocking One at 4 °C overnight. After

washing three times in PBS, cells were observed using a Leica SP8 confocal

microscope.

**Laser ablation**

Laser ablation was conducted using an Olympus IX 81 inverted microscope (20 x /0.70

NA dry objective lens), equipped with a spinning-disc confocal unit Yokogawa CSUX-1

and iXon3 897 EM-CCD camera (Andor), controlled with Andor IQ2 software. An N2

Micropoint laser (16 Hz, 365 nm, 2.0 µW, Photonic Instruments) was focused on the

membrane at the cell membrane. Time-lapse images were acquired every 200 msec

before and during the laser ablation process and analyzed with Fiji/ImageJ software.

**Fluorescence recovery after photobleaching (FRAP)**

FRAP assays in the tissue culture cells were conducted with a Leica SP8 confocal

microscope and software equipped in the microscope operating system. Regions of

interest (ROI) were bleached using a 488 nm laser. Pre-bleach and post-bleach images

were acquired with a 488 nm laser. Fluorescence recovery of GFP-ZO-1 was monitored.

Recovery data were background corrected and normalized to the ROI intensity before

bleaching. A reference ROI outside the bleached area was processed in the same way.

**Observation of GFP-ZO-1 dynamics in the tissue culture cells**

For the wound-healing assay, MDCK cells almost confluently grown on a glass-bottom

dish were transfected with GFP-ZO-1/pCS2. After one day, the cell layer was scratched

with a Pipetman tip, incubated at 37 °C, 5% $CO_2$ in a stage-top incubator (cat. # STXG,

Tokai HIT, Japan), and observed by a Leica SP8 confocal microscope.

For the latrunculin B treatment, 2.5 µM latrunculin B in the culture medium was

added to the culture dish of A6 cells transiently expressing GFP-ZO-1 so that the final

concentration of latrunculin B was 0.5 µM. For the digitonin treatment, 0.06% digitonin

in the culture medium was added to the culture dish so that the final concentration of

digitonin was 0.012%. For the hyperosmolarity treatment of A6 cells, $10 \times$ PBS was

added so that the concentration of PBS changed from $0.5 \times$ to $1.5 \times$. Cells were

observed at room temperature. For the trypsin-EDTA treatment, the culture medium of

MDCK cells was replaced with PBS containing 0.05% trypsin and 1 mM EDTA.

**Preparation of polyacrylamide-based plates**

Glass bottom dishes (27-mm diameter, cat.# 3910-035, IWAKI, Japan) were treated

with 2% (v/v) (3-aminopropyl) triethoxysilane (cat.# 440140 Sigma-Aldrich)/acetone.

Air-dried plates were washed with sterilized water three times and dried at 50 °C for 2 h.

45 µl acrylamide solution was loaded, and then a 24-mm diameter coverslip was loaded

onto the solution. After gelation, the coverslip was removed, and the surface of gelled

polyacrylamide was washed with 50 mM HEPES (pH 8.5). 200 µl of 0.5 mg/ml

Sulfo-SANPAH (cat.# 22589, Pierce) was loaded and irradiated with UV for 30 min

using Stratalinker 2400 (Stratagene). Then, it was washed twice with 50 mM HEPES

(pH 8.5), and 200 µl of 100 µg/ml fibronectin was loaded and incubated overnight at

4 °C. It was washed with the MDCK culture medium to block residual reactive groups,

and cells were plated.

**Atomic force microscopy (AFM) measurements**

AFM measurements were conducted with a JPK Nanowizard Cellhesion 200 (JPK

Instruments AG, Germany) fitted to an Axio Zoom V. 16 system (Zeiss). For a

cantilever, CP-CONT-BSG-A (sQUBE, Germany) was used. Young's modulus was

calculated using JPK data processing software (Bruker). Measurement conditions were

as follows: setpoint 50-200 nN, approach speed 2 µm/s, data rate 1000 Hz, measured

spring constant 0.342 N/m.

**References**

Anderson, J.M., Stevenson, B.R., Jesaitis, L.A., Goodenough, D.A., and Mooseker, M.S.

(1988). Characterization of ZO-1, a protein component of the tight junction from mouse

liver and Madin-Darby canine kidney cells. J Cell Biol *106*, 1141-1149.

Baum, B., Settleman, J., and Quinlan, M.P. (2008). Transitions between epithelial and

mesenchymal states in development and disease. Semin Cell Dev Biol *19*, 294-308.

Beutel, O., Maraspini, R., Pombo-García, K., Martin-Lemaitre, C., and Honigmann, A.

(2019). Phase separation of zonula occludens proteins drives formation of tight

junctions. Cell *179*, 923-936.

Bodor, D.L., Pönisch, W., Endres, R.G., and Paluch, E.K. (2020). Of cell shapes and

motion: The physical basis of animal cell migration. Dev Cell *52*, 550-562.

Bradley, A. (1987). Production and analysis of chimeric mice. In Teratocarcinoma and

Embryonic Stem Cells, E.J. Robertson, ed. (Oxford: IRL Press), pp. 113-151.

Brangwynne, C.P. (2013). Phase transitions and size scaling of membrane-less

organelles. J Cell Biol *203*, 875-881.

Chan, C.J., Costanzo, M., Ruiz-Herrero, T., Mönke, G., Petrie, R.J., Bergert, M.,

Diz-Muñoz, A., Mahadevan, L., and Hiiragi, T. (2019). Hydraulic control of mammalian

embryo size and cell fate. Nature *571*, 112-116.

Choi, Y.S., Sehgal, R., McCrea, P., and Gumbiner, B. (1990). A cadherin-like protein in

eggs and cleaving embryos of *Xenopus laevis* is expressed in oocytes in response to

progesterone. J Cell Biol *110*, 1575-1582.

Fanning, A.S., and Anderson, J.M. (2009). Zonula occludens-1 and -2 are cytosolic

scaffolds that regulate the assembly of cellular junctions. Ann N Y Acad Sci *1165*,

113-120.

Fanning, A.S., Ma, T.Y., and Anderson, J.M. (2002). Isolation and functional

characterization of the actin binding region in the tight junction protein ZO-1. FASEB J

*16*, 1835-1837.

Fujishiro, S.H., Tanimura, S., Mure, S., Kashimoto, Y., Watanabe, K., and Kohno, M.

(2008). ERK1/2 phosphorylate GEF-H1 to enhance its guanine nucleotide exchange

activity toward RhoA. Biochem Biophys Res Commun *368*, 162-167.

Fujiwara, S., Ohashi, K., Mashiko, T., Kondo, H., and Mizuno, K. (2016). Interplay

between Solo and keratin filaments is crucial for mechanical force-induced stress fiber

reinforcement. Mol Biol Cell *27*, 954-966.

Ginsberg, D., DeSimone, D., and Geiger, B. (1991). Expression of a novel cadherin

(EP-cadherin) in unfertilized eggs and early *Xenopus* embryos. Development *111*,

315-325.

Godard, B.G., and Heisenberg, C.P. (2019). Cell division and tissue mechanics. Curr

Opin Cell Biol *60*, 114-120.

González-Mariscal, L., Quirós, M., and Díaz-Coránguez, M. (2011). ZO proteins and

redox-dependent processes. Antioxid Redox Signal *15*, 1235-1253.

Haas, A.J., Zihni, C., Ruppel, A., Hartmann, C., Ebnet, K., Tada, M., Balda, M.S., and

Matter, K. (2020). Interplay between extracellular matrix stiffness and JAM-A regulates

mechanical load on ZO-1 and tight junction assembly. Cell Rep *32*, 107924.

Hallou, A., and Brunet, T. (2020). On growth and force: mechanical forces in

development. Development *147*, dev187302.

Hannezo, E., and Heisenberg, C.P. (2019). Mechanochemical feedback loops in

development and disease. Cell *178*, 12-25.

Hashimoto, Y., Kinoshita, N., Greco, T.M., Federspiel, J.D., Jean Beltran, P.M., Ueno,

N., and Cristea, I.M. (2019). Mechanical force induces phosphorylation-mediated

signaling that underlies tissue response and robustness in *Xenopus* embryos. Cell

Systems *8*, 226-241.

Hernández-Vega, A., Marsal, M., Pouille, P.A., Tosi, S., Colombelli, J., Luque, T.,

Navajas, D., Pagonabarraga, I., and Martín-Blanco, E. (2017). Polarized cortical tension

drives zebrafish epiboly movements. EMBO J *36*, 25-41.

Hino, N., Rossetti, L., Marín-Llauradó, A., Aoki, K., Trepat, X., Matsuda, M., and

Hirashima, T. (2020). ERK-mediated mechanochemical waves direct collective cell

polarization. Dev Cell *53*, 646-660.

Iioka, H., Iemura, S., Natsume, T., and Kinoshita, N. (2007). Wnt signalling regulates

paxillin ubiquitination essential for mesodermal cell motility. Nat Cell Biol *9*, 813-821.

Iioka, H., Ueno, N., and Kinoshita, N. (2004). Essential role of MARCKS in cortical

actin dynamics during gastrulation movements. J Cell Biol *164*, 169-174.

Ishida, T., and Kinoshita, K. (2007). PrDOS: prediction of disordered protein regions

from amino acid sequence. Nucleic Acids Res *35*, W460-464.

Ito, S., Okuda, S., Abe, M., Fujimoto, M., Onuki, T., Nishimura, T., and Takeichi, M.

(2017). Induced cortical tension restores functional junctions in adhesion-defective

carcinoma cells. Nat Commun *8*, 1834.

Itoh, K., Ossipova, O., and Sokol, S.Y. (2014). GEF-H1 functions in apical constriction

and cell intercalations and is essential for vertebrate neural tube closure. J Cell Sci *127*,

2542-2553.

Kato, M., Han, T.W., Xie, S., Shi, K., Du, X., Wu, L.C., Mirzaei, H., Goldsmith, E.J.,

Longgood, J., Pei, J., *et al.* (2012). Cell-free formation of RNA granules: low

complexity sequence domains form dynamic fibers within hydrogels. Cell *149*,

753-767.

Katsunuma, S., Honda, H., Shinoda, T., Ishimoto, Y., Miyata, T., Kiyonari, H., Abe, T.,

Nibu, K., Takai, Y., and Togashi, H. (2016). Synergistic action of nectins and cadherins

generates the mosaic cellular pattern of the olfactory epithelium. J Cell Biol *212*,

561-575.

Kim, H.Y., and Davidson, L.A. (2013). Investigating morphogenesis in *Xenopus*

embryos: imaging strategies, processing, and analysis. Cold Spring Harb Protoc *2013*,

298-304.

Kinoshita, N., Hashimoto, Y., Yasue, N., Suzuki, M., Cristea, I.M., and Ueno, N. (2020).

Mechanical stress regulates epithelial tissue integrity and stiffness through the

FGFR/Erk2 signaling pathway during embryogenesis. Cell Rep *30*, 3875-3888.

Konno, D., Shioi, G., Shitamukai, A., Mori, A., Kiyonari, H., Miyata, T., and Matsuzaki,

F. (2008). Neuroepithelial progenitors undergo LGN-dependent planar divisions to

maintain self-renewability during mammalian neurogenesis. Nat Cell Biol *10*, 93-101.

Kroschwald, S., Maharana, S., Mateju, D., Malinovska, L., Nüske, E., Poser, I., Richter,

D., and Alberti, S. (2015). Promiscuous interactions and protein disaggregases

determine the material state of stress-inducible RNP granules. eLife *4*, e06807.

Leonavicius, K., Royer, C., Preece, C., Davies, B., Biggins, J.S., and Srinivas, S. (2018).

Mechanics of mouse blastocyst hatching revealed by a hydrogel-based

microdeformation assay. Proc Natl Acad Sci U S A *115*, 10375-10380.

Li, P., Banjade, S., Cheng, H.C., Kim, S., Chen, B., Guo, L., Llaguno, M.,

Hollingsworth, J.V., King, D.S., Banani, S.F*., et al.* (2012). Phase transitions in the

assembly of multivalent signalling proteins. Nature *483*, 336-340.

Mimori-Kiyosue, Y., Matsui, C., Sasaki, H., and Tsukita, S. (2007). Adenomatous

polyposis coli (APC) protein regulates epithelial cell migration and morphogenesis via

PDZ domain-based interactions with plasma membranes. Genes Cells *12*, 219-233.

Murray, D.T., Kato, M., Lin, Y., Thurber, K.R., Hung, I., McKnight, S.L., and Tycko, R.

(2017). Structure of FUS protein fibrils and its relevance to self-assembly and phase

separation of low-complexity domains. Cell *171*, 615-627.

Paolini, A., and Abdelilah-Seyfried, S. (2018). The mechanobiology of zebrafish cardiac

valve leaflet formation. Curr Opin Cell Biol *55*, 52-58.

Ren, Y., Li, R., Zheng, Y., and Busch, H. (1998). Cloning and characterization of

GEF-H1, a microtubule-associated guanine nucleotide exchange factor for Rac and Rho

GTPases. J Biol Chem *273*, 34954-34960.

Schwayer, C., Shamipour, S., Pranjic-Ferscha, K., Schauer, A., Balda, M., Tada, M.,

Matter, K., and Heisenberg, C.P. (2019). Mechanosensation of tight junctions depends

on ZO-1 phase separation and flow. Cell *179*, 937-952.

Shiina, N. (2019). Liquid- and solid-like RNA granules form through specific scaffold

proteins and combine into biphasic granules. J Biol Chem *294*, 3532-3548.

Shulga, N., and Goldfarb, D.S. (2003). Binding dynamics of structural nucleoporins

govern nuclear pore complex permeability and may mediate channel gating. Mol Cell

Biol *23*, 534-542.

Spadaro, D., Le, S., Laroche, T., Mean, I., Jond, L., Yan, J., and Citi, S. (2017).

Tension-dependent stretching activates ZO-1 to control the junctional localization of its

interactors. Curr Biol *27*, 3783-3795 e3788.

Teng, X., Qin, L., Le Borgne, R., and Toyama, Y. (2017). Remodeling of adhesion and

modulation of mechanical tensile forces during apoptosis in *Drosophila* epithelium.

Development *144*, 95-105.

Thompson, D.W. (1917). On Growth and Form (Cambridge Univ. Press).

Vasileva, E., Sluysmans, S., Bochaton-Piallat, M.L., and Citi, S. (2017). Cell-specific

diversity in the expression and organization of cytoplasmic plaque proteins of apical

junctions. Ann N Y Acad Sci *1405*, 160-176.

Wegmann, S., Eftekharzadeh, B., Tepper, K., Zoltowska, K.M., Bennett, R.E., Dujardin,

S., Laskowski, P.R., MacKenzie, D., Kamath, T., Commins, C.*, et al.* (2018). Tau

protein liquid-liquid phase separation can initiate tau aggregation. EMBO J *37*, e98049.

Zhu, C., Bao, G., and Wang, N. (2000). Cell mechanics: mechanical response, cell

adhesion, and molecular deformation. Annu Rev Biomed Eng *2*, 189-226.

# Kinoshita et al. Figure 1

# Kinoshita et al. Figure 2

# Kinoshita et al. Figure 3

**A**

**B**

# Kinoshita et al. Figure 4

# Kinoshita et al. Figure 5

# Kinoshita et al. Figure 6

# Kinoshita et al. Figure S3