A meta-analysis of the effect of organic and mineral fertilizers on soil microbial diversity

Daniel P. Bebber[1] & Victoria R. Richards

[1] Department of Biosciences, College of Life and Environmental Sciences, Geoffrey Pope Building,
University of Exeter, Stocker Road, Exeter, EX4 4QD, UK.

Author for correspondence: Daniel P. Bebber (d.bebber@exeter.ac.uk)

ABSTRACT

Organic agriculture, employing manures or composts, has been proposed as a way of mitigating
undesirable impacts of mineral fertilizer use. Of particular interest is the effect of fertilizer regime on
soil microbes, which are key to nutrient cycling, plant health and soil structure. However, the effect of
fertilizers on soil microbial diversity remains poorly understood. Since biological diversity is an
important determinant of ecosystem function and a fundamental metric in community ecology, the
effects of fertilizer regimes on soil microbial diversity are of theoretical and applied interest. Here, we
conduct a meta-analysis of 31 studies reporting microbial diversity metrics in mineral fertilized (NPK),
organically fertilized (ORG) and unfertilized control (CON) soils. Of these studies, 26 reported
taxonomic diversity derived from sequencing, gradient gel electrophoresis, or RFLP. Functional
diversity, derived from Biolog Ecoplate^{TM} measures of carbon substrate metabolism, was reported in 8
studies, with 3 studies reporting both diversity metrics. We found that functional diversity was greater
in NPK compared with CON, greater in ORG vs CON, and greater in ORG vs NPK. Prokaryote
taxonomic diversity was not significantly different between NPK and CON, but greater in ORG vs
CON, in ORG vs. NPK. Fungal taxonomic diversity was not significantly different between any
treatment pairs. There was very high residual heterogeneity in all meta-analyses of soil diversity,
suggesting that a large amount of further research with detailed analysis of soil properties is required to
fully understand the influence of fertilizer regimes on microbial diversity and ecosystem function.

Keywords:

Soil; Microbiome; Diversity; Organic; Agriculture; Nitrogen

INTRODUCTION

Diversity plays a key role in the resilience and adaptability of complex systems [1], and biological

diversity has been central to understanding of the structure and function of ecological communities [2].

Human activities are rapidly eroding global biodiversity [3], hence understanding how human activities

influence diversity and how negative impacts can be avoided is an important goal of applied ecology

[4, 5]. Recent advances in environmental DNA sequencing and metabarcoding have revealed enormous

and unexpected microbial diversity in all habitats, particularly soils [6, 7]. Given the fundamental

importance of soils in terrestrial ecosystems, agriculture and food security, there is growing interest in

the role of microbial diversity in processes such as nutrient cycling [8], and how soil diversity can be

managed for maintenance of ecosystem services [9]. In agriculture, soil microbes are critical to desirable

functions such as nutrient cycling, carbon storage, erosion control via soil aggregation, and disease

suppression [10–12]. A large fraction of the world's agriculture soils are in poor and deteriorating

condition largely due to agricultural activities [13]. Therefore, understanding how to manage soils for

microbial diversity could help to prevent further deterioration.

Modern intensive, or conventional, agricultural methods, including application of mineral and chemical

fertilisers, regular tillage, and use of synthetic pesticides and herbicides, aim to increase soil nutrition

and suppress harmful species to produce higher crop yields, but these methods are environmentally

damaging (Robertson, 2000). For example, growing pressure on agricultural land has resulted in soil

nutrient depletion and soil erosion (Wopereis, Tamèlokpo, & Ezui, 2006; Lal, 2009). Mineral fertilizers

supply nitrogen (N) as ammonium nitrate or urea, phosphorus (P) and potassium (K), with around 100

Tg N applied globally each year [14]. The nitrogen sources are derived from the Haber-Bosch process

which reduces dinitrogen to ammonia, accounting for around one third of the nitrogen fixed from the

atmosphere [15]. Soil and plant-symbiotic microbes fix another third, marine cyanobacteria another

third, with small contributions from lightning and fossil fuel combustion [15]. Typically less than half

the N applied is taken up by crops, the remainder contributing to water pollution and the release of

NOX greenhouse gases [16]. Organic agriculture, which applies manure- or compost-based fertilizers

and soil conditioners and does not employ chemical pest controls (with certain exceptions), has been

proposed as a means of both increasing soil nutrition and reducing environmental impacts [17], but

evidence for organic benefits remains equivocal. For example, eutrophication potential appears greater

in many organic than conventional crop systems [18]

Soil health, the capacity of soils to function as a living system and sustain and promote plant and animal

communities, is linked to microbial diversity [19, 20]. Microbial diversity is correlated with soil

ecosystem multifunctionality including plant productivity, microbial biomass, availability of nitrate,

ammonium and phosphorus, and nitrogen mineralisation rates [8]. Hence, management to enhance soil

microbial diversity could improve soil health. Soils are exceedingly complex and biologically-diverse

ecosystems, varying in physico-chemical and biological composition across spatial scales [7, 21]. Meta-
analyses of multiple individual studies can reveal general trends and patterns in this complexity and
determine the effects of management interventions. For example, observational data suggest that
biological richness (measured by DNA sequences) is greatest at neutral pH and at a mean temperature
of 10 °C [7]. Meta-analysis of experimental studies shows that pH has the greatest influence on soil
microbial diversity among global change factors, alpha diversity rising with pH [22]. Nitrogen and NPK
have negative or non-significant effects on alpha diversity depending on microbial group, with
significant negative influences of N alone on diversity in agricultural soils. Soil functionality, defined
as the range of biogeochemical processes carried out by soils, increases with N and NPK. Changes in
diversity are negatively correlated with changes in functionality, perhaps due to functional redundancy
[22]. Other meta-analyses have demonstrated that microbial diversity [23] and biomass [24] increase in
crop rotations compared with monocultures. Organic agriculture greatly increases microbial biomass
carbon, microbial biomass nitrogen, and enzymatic activity compared with conventional systems [25].
Previously, a small meta-analysis found no significant effect of organic agriculture on soil organism
diversity, but this only included five studies and occurred before widespread use of sequencing to soil
microbial diversity [26]. However, despite growing application of sequencing technologies in
microbial ecology, results on the effects of organic and mineral fertilizers on measures of soil microbial
diversity remain unsynthesized.

Here, we conduct a meta-analysis quantifying the effects of organic fertilizers (manures and manure
composts) and mineral fertilizers (NPK) on soil microbial diversity, in comparison with unfertilized
controls. We compare results for taxonomic and functional diversity, and investigate the influence of
factors such as soil chemistry and duration of organic treatment on these effects. We hypothesise that
organic inputs will increase microbial diversity as compared to mineral fertilizer inputs, based on the
observation that organic fertilizers increase microbial biomass and enzymatic activity compared with
NPK [25].

MATERIALS AND METHODS

Criteria for meta-analysis

We used meta-analysis to test for differences in soil microbial functional (H_{fun}) and taxonomic (H_{tax}) diversity between soils treated with manure-based organic fertilizer (ORG), mineral fertilizer (NPK) and control (CON). Here, H_{tax} is defined as Shannon's diversity index calculated from the relative abundance of species, operational taxonomic units (OTUs) or amplicon sequence variants [ASVs; 27] in a sample. Shannon's diversity index was reported in all valid studies, hence we do not consider any other diversity indices (e.g. Simpson's). H_{fun} is defined as Shannon's diversity calculated from the relative conversion rates of various carbon sources in Biolog Ecoplates [e.g. 28]. We conducted a literature search using combinations of the terms 'soil', 'organic', 'agriculture', 'microbial', 'diversity', 'bacteria*', 'fung*', 'communit*', 'fertilizer', 'manure' and 'compost' (where * indicates a wildcard search, where appropriate) on Google Scholar, Scopus and Web of Science between August 2018 and June 2019. Additionally, the reference lists of the papers were browsed to find potentially appropriate studies which were not identified during the online literature search. All potentially appropriate papers identified for the meta-analysis came from peer-reviewed journals.

Papers were considered eligible for the analysis if they met the following criteria, adapted from Lori et al. [25]: Comparisons of the farming systems should be pairwise, meaning the organic- and mineral-fertilised treatments were subject to the same climatic conditions before and during sampling; The organic treatment must have been applied for a minimum of two consecutive years prior to soil sampling, and the treatments must be defined by the study; Results must report the mean Shannon diversity index (H) per treatment, uncertainty of the mean (either standard error of the mean or standard deviation of the sample distribution) and sample size (n); experiments were conducted in open fields or under cover, not in pots or containers; the study must have been published no earlier than the year 2000; the mineral and organic treatments must be clearly described (description of fertilizer, application rate). If there were several different organic or conventional treatments within the same study, all appropriate combinations were reported and treated as individual comparisons but with non-independent errors (see meta-analysis methods). Results of combination treatments (e.g. manure plus NPK fertilizer) were not included.

For all valid studies, we extracted mean, SE or SD, and sample size for H, for control, NPK and organic treatments. Where SE was reported, this was converted to SD by multiplying by the square root of sample size. When data were presented in graphical form, we extracted means and errors using the online tool WebPlotDigitalizer (https://automeris.io/WebPlotDigitizer/). Additionally we abstracted the following variables: study location; reported soil type, duration of organic treatment, organic treatment type (i.e. source of manure), organic treatment application rate (mass per hectare per year or equivalent nitrogen mass per hectare per year); mineral fertilizer application rate (nitrogen mass per hectare per

year); crops grow; taxonomic group of microbes analysed (e.g. all, bacteria, fungi); median soil depth

sampled (usually the mid-point of the range of depths sampled); brief summary of the methodology

used (e.g. 16S sequencing, 16S DGGE, Biolog Ecoplate). Where reported, we abstracted mean and SE

of soil pH, and mean and SE of available nitrogen content (mg kg^{-1}) which was often reported as the

concentration of ammonium and nitrate.

Meta-analysis

Meta-analysis was conducted using the *metafor* package v. 2.4–0 for R v. 4.0.2 [29]. Effect size was

calculated as the log–transformed ratio of means, $\log(y_{exp}/y_{con})$ where y_{exp} and y_{con} are the means of the

experimental treatment and control, respectively [30]. There was at least one pairwise comparison of H

in each study. However, several studies reported data from more than one pairwise comparison, with

multiple organic and/or conventional fertilisation treatments investigated. We fitted multi-level

random– and mixed–effects (i.e. testing for the effects of moderator variables) models by restricted

maximum likelihood, with random intercepts per study. Other similar meta-analyses have ignored

potential non-independence among results from the same study and treated each reported mean as an

independent replicate [e.g. 25]. Standard errors and confidence limits for parameters were estimated

from the *t*–distribution. We report the I^2 statistic and *Q*-test for heterogeneity [31], and provide funnel

plots rank correlation tests for funnel plot asymmetry [29].

RESULTS

We abstracted data from 32 studies which met our requirements (Table S1). Our search criteria provided

a long-list of 259 studies, of which 195 were rejected on the basis of the abstract, leaving a shortlist of

64. A further 32 were rejected after detailed reading of the full text. Reasons for the rejection of studies

included: no reporting of diversity metrics (e.g. reporting of species richness); no or unclear reporting

of standard error or standard deviation for diversity metrics; no manure used in organic treatment; no

organic treatment alone (e.g. manure only applied in combination with NPK); no NPK treatment; no

experimental treatments (e.g. comparison of different farms under organic and conventional

treatments); lack of detail in description of treatment types or levels.

Around half (17) of the included studies reported one set of comparisons, with either or both of

unfertilized control (CON) and mineral fertilizer (NPK), 10 reported two comparisons, and five studies

reported up to six comparisons, giving 58 comparisons in total. The majority of treatments were

replicated three or four times. The functional diversity studies, and a single study reporting soil dilution

plate assays [32], were taxonomically non-specific. The single soil dilution plate assay was omitted

from the meta-analysis due to the low sensitivity of the method. In that study, poultry manure treatment

resulted in significantly greater microbial diversity than inorganic fertilizer treatment [32]. In the

taxonomic diversity studies, bacteria were the most commonly analysed group (25 studies, including

one of actinomycetes only, one of bacteria and archaea, and one of nitrogen-fixing bacteria), followed

by fungi (6 studies, including one of arbuscular mycorrhizal fungi only). Most studies were conducted

in China (20), followed by India (6), with single studies from Austria, Canada, Denmark, Kenya, the

Netherlands, and the USA.

Soil types were not reported according to any standard taxonomy. Soils were sampled to a median depth

of up to 15 cm (median 10.0 cm, IQR 7.5 - 10.0 cm). Soil nitrogen content in unfertilized plots was

reported in 16 studies (mean mg N kg^{-1} median 34.9, IQR 26.5 - 90.7). Manure-based organic fertilizers

were applied in all studies, derived variously from cattle, pigs, poultry, horses, sheep, or mixtures of

these. Composted manures were employed in 9 studies, fresh manures in 24 studies (one study

employed both). A small number of studies reported more than one manure treatment or level [32–34].

Application rates were reported as either total mass per area per year (25 studies, median 11.4 t ha^{-1} y^{-1}

^{1}, IQR 5.0 - 18.6 t ha^{-1} y^{-1}) and/or as nitrogen addition (15 studies, median 91.6 kg ha^{-1} y^{-1}, IQR 58.8 -

148.8 kg ha^{-1} y^{-1}). We did not attempt to estimate nitrogen content in manures where this was not

reported, due to the large variability in the nitrogen fraction among different treatments. The reported

nitrogen fraction varied from 0.4 % in horse manure to 2.7 % in pig manure, with overall median 1.0

% (IQR 0.7 - 2.0 %) among the treatments reported in the studies. The level of nitrogen in the manure

treatment was similar to that in corresponding NPK fertilizer treatments, where reported (absolute

percentage difference median 13.4 %, IQR 0.0 - 56.0 %). The duration of organic treatment was right-

skewed, but many studies reported treatment periods of several decades (median 20.5 y, IQR 11.5 -

34.3 y). A variety of crops were grown, either as monocultures or in mixtures, the most common being

maize (10 studies), wheat (8 studies), and rice (6 studies). Soil pH in unfertilized plots was reported in

18 studies (mean pH median 6.4, interquartile range 5.8 - 7.4). Meta-regression showed that soil pH

declined under NPK treatment compared to control soil (Fig. S1a). Manure-based fertilizers increased

pH in acid soils and reduced pH in alkaline soils, (Fig. S1b). Funnel plots showed no asymmetry (Fig.

S2). There was a large degree of residual heterogeneity in the meta-analyses of soil pH (Table S2).

H_{fun} was reported by 8 studies, all employing Biolog Ecoplates (Table S1). H_{fun} varied between 1.12 and

4.61 with values in three clusters comprising a single study from Austria with $H_{fun} \sim 1.2$, four studies

from China and one from India with $H_{fun} \sim 2.9$, and two studies from India with $H_{fun} \sim 4.5$ (Fig. 1a-c).

H_{fun} increased by an average of 2.8 % in NPK vs. CON, 7.0 % in ORG vs. CON, and 3.8 % in ORG vs.

NPK (Fig. 2, Table 1). We found no influence of the duration of organic amendment on the change in

H_{fun} compared with control (LRM change per year of organic treatment -0.0008 ± 0.0013 y^{-1}, t = -0.58,

p = 0.56) or NPK (0.0006 ± 0.0007 y^{-1}, t = 0.84, p = 0.40) treatments. We found no effect of the level

of NPK addition relative to control (-0.0013 ± 0.0009, t = -1.42, p = 0.15), nor of level of organic

fertilizer addition quantity (0.0000 ± 0.0000, t = -0.35, p = 0.72) or nitrogen equivalent relative to

control (0.0001 ± 0.0005, t = 0.18, p = 0.86). We found a marginally-significant effect of control soil

pH on the effect size of ORG vs. CON (-0.052 ± 0.025, $t = -2.10$, $p = 0.036$), but not of soil pH on NPK

vs. CON (0.0045 ± 0.0527, $t = 0.085$, $p = 0.93$). Soil N did not influence the effect size of ORG vs.

CON (0.000 ± 0.001, $t = -0.01$, $p = 0.99$). Funnel plots did not exhibit significant asymmetry (Fig. S3).

There was a large degree of residual heterogeneity in all meta-analyses of H_{fun} ($I^2 > 75\%$, Table S2).

H_{tax} was reported in 26 studies, employing DNA amplicon sequencing (16 studies), various forms of

gradient gel electrophoresis (9 studies), and T-RFLP (1 study). H_{tax} tended to be higher in prokaryotes

than fungi, and in sequencing compared with DGGE methods (Fig. 1d-f). Median H_{tax} for prokaryotes

was 6.36 using sequencing and 2.85 using DGGE. Median H_{tax} for fungi was 3.36 using sequencing and

1.40 using DGGE. H_{tax} of NPK-fertilized soils did not differ from control for prokaryotes or fungi, nor

in the single study using dilution plate assays (Fig. 3, Table 2). H_{tax} of prokaryotes was greater in ORG

soils than both NPK and CON. There was no significant difference in fungal H_{tax} between ORG and

CON or NPK. We found no evidence that duration of organic amendments (0.0006 ± 0.0009, $t = 0.66$,

$p = 0.51$), pH (-0.0054 ± 0.0038, $t = -1.42$, $p = 0.15$) or soil N (-0.0008 ± 0.0004, $t = -2.1$, $p = 0.051$)

influenced the effect size of ORG vs. CON. Funnel plots did not exhibit significant asymmetry (Fig.

S4). There was a large degree of residual heterogeneity in meta-analyses of H_{tax} ($I^2 > 95\%$, Table S2).

DISCUSSION

We found significant, but varying, effects of both NPK and organic fertilizers on soil microbe

taxonomic and functional diversity. Taxonomic diversity in organic treatments compared with NPK

was greater for prokaryotes but not significantly different for fungi. We found no significant difference

in taxonomic diversity between NPK and control. Functional diversity was significantly greater in NPK

compared with control, and in organic treatment compared with NPK. However, the diversity

differences we found were small, indicating that choice of fertilization has marginal effects on this

measure of soil microbial community structure. We did not detect any significant influence of factors

such as duration or type of organic treatment, soil chemistry or experimental methodology on effect

sizes. Importantly, the residual heterogeneity in all meta-analyses was very high.

Several factors must be considered when interpreting the effects of fertilizers on soil microbial

functional and taxonomic diversity (Fig. 4). First, while in many studies the amount of nitrogen applied

in organic and NPK treatments was similar, farmyard manure is far more physically, chemically and

biologically complex. In addition to NPK, manures contain undigested plant matter (lignin, cellulose,

hemicellulose), lipids, carbohydrates, proteins, and nutritive elements (e.g. magnesium, iron,

manganese, zinc, copper) [35]. If structural complexity and heterogeneity are enhanced by the addition

of organic matter across spatial scales [36], this could in itself increase microbial diversity through

provision of ecospace that reduces interspecific competition [37]. Provision of additional energy

(organic carbon) and micronutrients in manures could sustain a greater diversity of microbes via the

species-energy hypothesis which predicts that more species can be sustained in ecosystems supporting

more individuals [38], since manure increases soil microbial biomass more than NPK [25]. Second,

microbial communities may be indirectly influenced by changes in crop plants. While organic

agriculture tends to be less productive in terms of crop yields [39, 40], there is some evidence that root

development is greater in organic farming [41]. Plant rhizospheres tend to harbour the greatest microbial

diversity of all terrestrial ecosystems [7], hence it is possible that enhanced root development and

exudation of organic compounds influence the soil diversity measured in these studies. Third, there is

the gut microbial community residing within the manure itself. Both fresh and composted manures have

high fungal and prokaryote diversity, with community composition changing as composting proceeds

[42]. Bacterial diversity has been found to peak during the thermophilic phase of aerobic composting

during which temperatures exceeded 55 °C, while fungal diversity was lowest during this stage [42].

Composting can reduce the presence of undesirable microbes in manure, for example those carrying

antimicrobial resistance genes [43], but gut microbes (including human pathogens) can survive for long

periods in compost-amended soils [44]. Hence, increased soil microbial diversity in organic systems

may be due to persistence of gut microbes.

Few studies reported both taxonomic and functional diversity, hence we were unable to determine a

relationship between these two metrics. Functional diversity has been found to increase with species

diversity in plant communities [45, 46], but less clearly in bird communities [47]. The relationship

between soil taxonomic diversity and ecosystem function (as opposed to functional diversity) tends to

be positive [48–50]. Given that ecosystem functions like nitrogen cycling require multiple interacting

processes, a positive relationship between taxonomic diversity and functional diversity might be

expected. Soil ecosystem multifunctionality (the capacity of soils to sustain many functions

simultaneously) increases with microbial diversity [8], while taxonomic and functional gene diversity

are closely correlated [51]. However, taxonomic and functional diversity can be decoupled by the

process of horizontal gene transfer, because different taxa can perform similar tasks through shared

genes [51]. In addition, the presence of physiologically-distinct subgroups within OTUs commonly

defined by > 97 % sequence similarity (so-called 'microdiversity') means that functional diversity can

be large in groups of apparently identical microbial taxa [52]. Microbial functional diversity tends to

be considered from the perspective of potential functions inferred from gene sequences [53]. In contrast,

we report results from studies of the diversity of actual functions (metabolization of carbon sources)

carried out by the soil microbial community. These genotypic vs. phenotypic measures of functional

diversity are analogous to the fundamental and realized niches of species, whereby the potential

functions identified in gene sequences are likely to exceed the observed functions carried out by the

community under particular environmental conditions [53]. Discovering the relationships between

potential and observed functions of microbial communities is a key goal of microbial ecology.

Organic farming has been promoted as a more sustainable alternative to conventional agriculture,
because of greater energy efficiency, potentially closed nutrient cycles and increased biodiversity [17].
Certainly, the rapid decline of insect populations around the world due to intensive agriculture and
pesticides, is an urgent impetus for reconsidering the way we produce food [54]. These potential
benefits come at a cost of reduced productivity compared with convential farming [39, 40], though there
is some evidence that organic production can eventually catch up with conventional yields and provides
greater spatial and temporal stability [55]. Environmental impacts of organic agriculture include greater
land use and eutrophication potential per unit of food produced, contradicting the aim of closed nutrient
cycles [18]. Our results suggest that, when considering organic crop fertilization alone and as a strict
alternative to NPK (rather than a mixed system of manure-derived and NPK fertilizer), microbial
functional diversity and bacterial taxonomic diversity are slightly, but significantly greater in organic
systems while the effect on fungal taxonomic diversity is unclear. The enhanced eutrophication
potential of organic farming, due to temporal mismatching between fertilizer addition and plant
demand, is much larger than the increases in taxonomic or functional diversity we found [18]. Hence,
it appears that increased soil microbial diversity under organic farming does not translate to improved
nutrient cycling and retention. While we did not compare combined treatments (e.g. NPK with manure
or compost) with single treatments, farmers commonly apply a diversity of fertilizers. For example, a
comparison of low-input organic systems with conventional mixed (NPK and manure) and conventional
NPK-fertilized systems in Switzerland found the greatest bacterial α-diversity in the organic systems,
followed by the mixed and NPK-only systems [56]. Some studies which were not included in the meta-
analysis employed manure-free organic treatments, finding no effect [57] or variables effects [58] on
soil microbial diversity.

Many studies addressing soil microbial diversity in response to fertilization could not be included in
our meta-analysis because of incomplete or unclear reporting. Without giving specific examples, we
found that many studies did not provide uncertainties (standard error or standard deviation) for
parameter estimates, gave unclear descriptions of the fertilisation routines and inputs, or reported
alternative measures of diversity. Of the studies we included, several did not report basic soil chemistry
metrics such as pH and nitrogen content. Given the importance of pH in determining soil microbial
diversity [22], and the central relevance of nitrogen in these experiments, any future research should
report these variables at minimum. We were only able to include a relatively small number of studies
in our meta-analysis, though we note that in the Cochrane Library of medical meta-analyses, the median
number of studies is seven or below [59]. We found very large residual heterogeneity among effect
sizes, i.e. most of the variation in effect sizes among studies remains unexplained. Further research will
reveal whether the mean effects we detected are general, and what other factors help explain variation
among soils, climates, locations and experimental treatments. To achieve this, it is critical that detailed
and complete data on key variables are reported. In addition, better understanding of soil ecosystem

functioning will be achieved via analysis of the taxa and functional genes identified within samples, in

addition to summary metrics like diversity [56]. In this way, the powerful new bioinformatics tools at

our disposal can be harnessed to fully understand the relationships between agriculture and soil health.

DECLARATIONS

FUNDING

VR was funded by an undergraduate project studentship at Exeter University, UK. DB was funded by

EC Horizon 2020 project ID 727624.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest

AVAILABILITY OF DATA

The data used in the meta-analysis will be made available on publication of the manuscript.

CODE AVAILABILITY

Code used in the meta-analysis is available from the author by reasonable request.

AUTHORS CONTRIBUTIONS

DB developed the research. DB and VR collected data, analysed results and wrote the paper.

ETHICS APPROVAL

Not applicable

CONSENT TO PARTICIPATE

Not applicable

CONSENT FOR PUBLICATION

Not applicable

REFERENCES

1.   Page SE (2011) Diversity and Complexity. Princeton University Press, Princeton

2.   Ricklefs RE, Schluter D (1994) Species Diversity in Ecological Communities. University of
Chicago Press, Chicago

3.   Ceballos G, Ehrlich PR, Barnosky AD, et al (2015) Accelerated modern human–induced species
losses: Entering the sixth mass extinction. Science Advances 1:e1400253.
https://doi.org/10.1126/sciadv.1400253

4.   Rudd MA, Beazley KF, Cooke SJ, et al (2011) Generation of Priority Research Questions to
Inform Conservation Policy and Management at a National Level. Conserv Biol 25:476–484.
https://doi.org/10.1111/j.1523-1739.2010.01625.x

5.   Sutherland WJ, Armstrong-Brown S, Armsworth PR, et al (2006) The identification of 100
ecological questions of high policy relevance in the UK. Journal of Applied Ecology 43:617–627.
https://doi.org/10.1111/j.1365-2664.2006.01188.x

6.   Lloyd KG, Steen AD, Ladau J, et al (2018) Phylogenetically Novel Uncultured Microbial Cells
Dominate Earth Microbiomes. mSystems 3:. https://doi.org/10.1128/mSystems.00055-18

7.   Thompson LR, Sanders JG, McDonald D, et al (2017) A communal catalogue reveals Earth's
multiscale microbial diversity. Nature 551:457. https://doi.org/10.1038/nature24621

8.   Delgado-Baquerizo M, Maestre FT, Reich PB, et al (2016) Microbial diversity drives
multifunctionality in terrestrial ecosystems. Nature Communications 7:10541.
https://doi.org/10.1038/ncomms10541

9.   Lemanceau P, Maron P-A, Mazurier S, et al (2015) Understanding and managing soil
biodiversity: a major challenge in agroecology. Agron Sustain Dev 35:67–81.
https://doi.org/10.1007/s13593-014-0247-0

10.   Mazzola M (2004) Assessment and Management of Soil Microbial Community Structure for
Disease Suppression. Annual Review of Phytopathology 42:35–59.
https://doi.org/10.1146/annurev.phyto.42.040803.140408

11.   Rillig MC, Wright SF, Eviner VT (2002) The role of arbuscular mycorrhizal fungi and glomalin
in soil aggregation: comparing effects of five plant species. Plant and Soil 238:325–333.
https://doi.org/10.1023/A:1014483303813

12.   Sahu N, Vasu D, Sahu A, et al (2017) Strength of microbes in nutrient cycling: A key to soil
health. In: Agriculturally Important Microbes for Sustainable Agriculture

13.   Wuepper D, Borrelli P, Finger R (2020) Countries and the global rate of soil erosion. Nature
Sustainability 3:51–55. https://doi.org/10.1038/s41893-019-0438-4

14.   FAO (2020) FAOSTAT. Food and Agriculture Organization of the United Nations, Rome

15.   Fowler D, Coyle M, Skiba U, et al (2013) The global nitrogen cycle in the twenty-first century.
Phil Trans R Soc B 368:20130164. https://doi.org/10.1098/rstb.2013.0164

16.   Zhang X, Davidson EA, Mauzerall DL, et al (2015) Managing nitrogen for sustainable
development. Nature 528:51–59. https://doi.org/10.1038/nature15743

17.   Reganold JP, Wachter JM (2016) Organic agriculture in the twenty-first century. Nature Plants
371   2:1–8. https://doi.org/10.1038/nplants.2015.221

372   18.   Clark M, Tilman D (2017) Comparative analysis of environmental impacts of agricultural
373   production systems, agricultural input efficiency, and food choice. Environ Res Lett 12:064016.
374   https://doi.org/10.1088/1748-9326/aa6cd5

375   19.   Kibblewhite MG, Ritz K, Swift MJ (2008) Soil health in agricultural systems. Philosophical
376   Transactions   of   the   Royal   Society   B: Biological   Sciences   363:685–701.
377   https://doi.org/10.1098/rstb.2007.2178

378   20.   van Bruggen AHC, Goss EM, Havelaar A, et al (2019) One Health - Cycling of diverse microbial
379   communities as a connecting force for soil, plant, animal, human and ecosystem health. Science
380   of the Total Environment 664:927–937. https://doi.org/10.1016/j.scitotenv.2019.02.091

381   21.   Howe AC, Jansson JK, Malfatti SA, et al (2014) Tackling soil diversity with the assembly of
382   large, complex metagenomes. PNAS 111:4904–4909. https://doi.org/10.1073/pnas.1402564111

383   22.   Zhou Z, Wang C, Luo Y (2020) Meta-analysis of the impacts of global change factors on soil
384   microbial   diversity   and   functionality.   Nature   Communications   11:3072.
385   https://doi.org/10.1038/s41467-020-16881-7

386   23.   Venter ZS, Jacobs K, Hawkins HJ (2016) The impact of crop rotation on soil microbial diversity:
387   A meta-analysis. Pedobiologia

388   24.   McDaniel MD, Tiemann LK, Grandy AS (2014) Does agricultural crop diversity enhance soil
389   microbial biomass and organic matter dynamics? A meta-analysis. Ecological Applications.
390   https://doi.org/10.1890/13-0616.1

391   25.   Lori M, Symnaczik S, Mäder P, et al (2017) Organic farming enhances soil microbial abundance
392   and   activity—A   meta-analysis   and   meta-regression.   PLOS   ONE   12:e0180442.
393   https://doi.org/10.1371/journal.pone.0180442

394   26.   Bengtsson J, AhnströM J, Weibull A-C (2005) The effects of organic agriculture on biodiversity
395   and abundance: a meta-analysis: Organic agriculture, biodiversity and abundance. Journal of
396   Applied Ecology 42:261–269. https://doi.org/10.1111/j.1365-2664.2005.01005.x

397   27.   Callahan BJ, McMurdie PJ, Holmes SP (2017) Exact sequence variants should replace operational
398   taxonomic   units   in   marker-gene   data   analysis.   The   ISME   Journal   11:2639–2643.
399   https://doi.org/10.1038/ismej.2017.119

400   28.   Ros M, Klammer S, Knapp B, et al (2006) Long-term effects of compost amendment of soil on
401   functional and structural diversity and microbial activity. Soil Use and Management 22:209–218.
402   https://doi.org/10.1111/j.1475-2743.2006.00027.x

403   29.   Viechtbauer W (2010) Conducting Meta-Analyses in R with the metafor Package. Journal of
404   Statistical Software 36:1–48. https://doi.org/10.18637/jss.v036.i03

405   30.   Hedges LV, Gurevitch J, Curtis PS (1999) The Meta-Analysis of Response Ratios in Experimental
406   Ecology.   Ecology   80:1150–1156.   https://doi.org/10.1890/0012-
407   9658(1999)080[1150:TMAORR]2.0.CO;2

408   31.   Higgins JPT, Thompson SG (2002) Quantifying heterogeneity in a meta-analysis. Statistics in
409   Medicine 21:1539–1558. https://doi.org/10.1002/sim.1186

32. Mahanta D, Bhattacharyya R, Mishra PK, et al (2017) Influence of a six-year organic and
inorganic fertilization on the diversity of the soil culturable microrgansims in the Indian mid-
Himalayas. Applied Soil Ecology 120:229–238. https://doi.org/10.1016/j.apsoil.2017.08.012

33. Hu X, Liu J, Wei D, et al (2018) Soil Bacterial Communities Under Different Long-Term
Fertilization Regimes in Three Locations Across the Black Soil Region of Northeast China.
Pedosphere 28:751–763. https://doi.org/10.1016/S1002-0160(18)60040-2

34. Liu W, Zhang Y, Jiang S, et al (2019) Spatiotemporal differences in the arbuscular mycorrhizal
fungi communities in soil and roots in response to long-term organic compost inputs in an
intensive agricultural cropping system on the North China Plain. J Soils Sediments 19:2520–2533.
https://doi.org/10.1007/s11368-019-02244-3

35. Levi-Minzi R, Riffaldi R, Saviozzi A (1986) Organic matter and nutrients in fresh and mature
farmyard manure. Agricultural Wastes 16:225–236. https://doi.org/10.1016/0141-
4607(86)90068-5

36. Lehmann J, Solomon D, Kinyangi J, et al (2008) Spatial complexity of soil organic matter forms
at nanometre scales. Nature Geoscience 1:238–242. https://doi.org/10.1038/ngeo155

37. Vos M, Wolf AB, Jennings SJ, Kowalchuk GA (2013) Micro-scale determinants of bacterial
diversity in soil. FEMS Microbiology Reviews 37:936–954. https://doi.org/10.1111/1574-
6976.12023

38. Clarke A, Gaston KJ (2006) Climate, energy and diversity. Proceedings of the Royal Society B:
Biological Sciences 273:2257–2266. https://doi.org/10.1098/rspb.2006.3545

39. de Ponti T, Rijk B, van Ittersum MK (2012) The crop yield gap between organic and conventional
agriculture. Agricultural Systems 108:1–9. https://doi.org/10.1016/j.agsy.2011.12.004

40. Hijbeek R, van Ittersum MK, ten Berge HFM, et al (2017) Do organic inputs matter – a meta-
analysis of additional yield effects for arable crops in Europe. Plant Soil 411:293–303.
https://doi.org/10.1007/s11104-016-3031-x

41. Hu T, Sørensen P, Wahlström EM, et al (2018) Root biomass in cereals, catch crops and weeds
can be reliably estimated without considering aboveground biomass. Agriculture, Ecosystems &
Environment 251:141–148. https://doi.org/10.1016/j.agee.2017.09.024

42. Meng Q, Yang W, Men M, et al (2019) Microbial Community Succession and Response to
Environmental Variables During Cow Manure and Corn Straw Composting. Front Microbiol 10:.
https://doi.org/10.3389/fmicb.2019.00529

43. Gou M, Hu H-W, Zhang Y-J, et al (2018) Aerobic composting reduces antibiotic resistance genes
in cattle manure and the resistome dissemination in agricultural soils. Science of The Total
Environment 612:1300–1310. https://doi.org/10.1016/j.scitotenv.2017.09.028

44. Sharma M, Reynnells R (2016) Importance of Soil Amendments: Survival of Bacterial Pathogens
in Manure and Compost Used as Organic Fertilizers. Microbiology Spectrum 4:PFS-0010-2015.
https://doi.org/10.1128/microbiolspec.PFS-0010-2015

45. Mensah S, Veldtman R, Assogbadjo AE, et al (2016) Tree species diversity promotes
aboveground carbon storage through functional diversity and functional dominance. Ecology and
Evolution 6:7546–7557. https://doi.org/10.1002/ece3.2525

46. Zhang Q, Buyantuev A, Li FY, et al (2017) Functional dominance rather than taxonomic diversity
and functional diversity mainly affects community aboveground biomass in the Inner Mongolia
grassland. Ecology and Evolution 7:1605–1615. https://doi.org/10.1002/ece3.2778

47. Morelli F, Benedetti Y, Perna P, Santolini R (2018) Associations among taxonomic diversity,
functional diversity and evolutionary distinctiveness vary among environments. Ecological
Indicators 88:8–16. https://doi.org/10.1016/j.ecolind.2018.01.022

48. Bardgett RD, Putten WH van der (2014) Belowground biodiversity and ecosystem functioning.
Nature 515:505. https://doi.org/10.1038/nature13855

49. Maron P-A, Sarr A, Kaisermann A, et al (2018) High Microbial Diversity Promotes Soil
Ecosystem Functioning. Appl Environ Microbiol 84:e02738-17.
https://doi.org/10.1128/AEM.02738-17

50. Philippot L, Spor A, Hénault C, et al (2013) Loss in microbial diversity affects nitrogen cycling
in soil. The ISME Journal 7:1609–1619. https://doi.org/10.1038/ismej.2013.34

51. Zhang X, Johnston ER, Wang Y, et al (2019) Distinct Drivers of Core and Accessory Components
of Soil Microbial Community Functional Diversity under Environmental Changes. mSystems 4:.
https://doi.org/10.1128/mSystems.00374-19

52. Larkin AA, Martiny AC (2017) Microdiversity shapes the traits, niche space, and biogeography
of microbial taxa. Environmental Microbiology Reports 9:55–70. https://doi.org/10.1111/1758-
2229.12523

53. Escalas A, Hale L, Voordeckers JW, et al (2019) Microbial functional diversity: From concepts
to applications. Ecology and Evolution 9:12000–12016. https://doi.org/10.1002/ece3.5670

54. Sánchez-Bayo F, Wyckhuys KAG (2019) Worldwide decline of the entomofauna: A review of its
drivers. Biological Conservation 232:8–27. https://doi.org/10.1016/j.biocon.2019.01.020

55. Schrama M, de Haan JJ, Kroonen M, et al (2018) Crop yield gap and stability in organic and
conventional farming systems. Agriculture, Ecosystems and Environment.
https://doi.org/10.1016/j.agee.2017.12.023

56. Hartmann M, Frey B, Mayer J, et al (2015) Distinct soil microbial diversity under long-term
organic and conventional farming. The ISME Journal 9:1177–1194.
https://doi.org/10.1038/ismej.2014.210

57. Wu T, Milner H, Díaz-Pérez JC, Ji P (2015) Effects of soil management practices on soil microbial
communities and development of southern blight in vegetable production. Applied Soil Ecology
91:58–67. https://doi.org/10.1016/j.apsoil.2015.02.011

58. Wang L, Yang F, E Y, et al (2016) Long-Term Application of Bioorganic Fertilizers Improved
Soil Biochemical Properties and Microbial Communities of an Apple Orchard Soil. Front
Microbiol 7:. https://doi.org/10.3389/fmicb.2016.01893

59. von Hippel PT (2015) The heterogeneity statistic I2 can be biased in small meta-analyses. BMC
Med Res Methodol 15:35. https://doi.org/10.1186/s12874-015-0024-z

60. Balachandar D, Doud MS, Schneper L, et al (2014) Long-Term Organic Nutrient Management
Fosters the Eubacterial Community Diversity in the Indian Semi-arid Alfisol as Revealed by
Length Heterogeneity–PCR. Communications in Soil Science and Plant Analysis 45:189–203.
https://doi.org/10.1080/00103624.2013.841919

61.   Chaudhry V, Rehman A, Mishra A, et al (2012) Changes in Bacterial Community Structure of
492   Agricultural Land Due to Long-Term Organic and Chemical Amendments. Microb Ecol 64:450–
493   460. https://doi.org/10.1007/s00248-012-0025-y

494   62.   Chen X, Li Z, Liu M, et al (2015) Microbial community and functional diversity associated with
495   different aggregate fractions of a paddy soil fertilized with organic manure and/or NPK fertilizer
496   for 20 years. J Soils Sediments 15:292–301. https://doi.org/10.1007/s11368-014-0981-6

497   63.   Chen C, Zhang J, Lu M, et al (2016) Microbial communities of an arable soil treated for 8 years
498   with organic and inorganic fertilizers. Biol Fertil Soils 52:455–467.
499   https://doi.org/10.1007/s00374-016-1089-5

500   64.   Cui X, Zhang Y, Gao J, et al (2018) Long-term combined application of manure and chemical
501   fertilizer sustained higher nutrient status and rhizospheric bacterial diversity in reddish paddy soil
502   of Central South China. Sci Rep 8:1–11. https://doi.org/10.1038/s41598-018-34685-0

503   65.   Ding J, Jiang X, Ma M, et al (2016) Effect of 35 years inorganic fertilizer and manure amendment
504   on structure of bacterial and archaeal communities in black soil of northeast China. Applied Soil
505   Ecology 105:187–195. https://doi.org/10.1016/j.apsoil.2016.04.010

506   66.   Ding J, Jiang X, Guan D, et al (2017) Influence of inorganic fertilizer and organic manure
507   application on fungal communities in a long-term field experiment of Chinese Mollisols. Applied
508   Soil Ecology 111:114–122. https://doi.org/10.1016/j.apsoil.2016.12.003

509   67.   Ge Y, Zhang J, Zhang L, et al (2008) Long-term fertilization regimes affect bacterial community
510   structure and diversity of an agricultural soil in northern China. J Soils Sediments 8:43–50.
511   https://doi.org/10.1065/jss2008.01.270

512   68.   Gu Y, Zhang X, Tu S, Lindström K (2009) Soil microbial biomass, crop yields, and bacterial
513   community structure as affected by long-term fertilizer treatments under wheat-rice cropping.
514   European Journal of Soil Biology 45:239–246. https://doi.org/10.1016/j.ejsobi.2009.02.005

515   69.   Guanghua W, Junjie L, Xiaoning Q, et al (2008) Effects of fertilization on bacterial community
516   structure and function in a black soil of Dehui region estimated by Biolog and PCR-DGGE
517   methods. Acta Ecologica Sinica 28:220–226. https://doi.org/10.1016/S1872-2032(08)60023-2

518   70.   Hamm AC, Tenuta M, Krause DO, et al (2016) Bacterial communities of an agricultural soil
519   amended with solid pig and dairy manures, and urea fertilizer. Applied Soil Ecology 103:61–71.
520   https://doi.org/10.1016/j.apsoil.2016.02.015

521   71.   Han S, Zeng L, Luo X, et al (2018) Shifts in Nitrobacter- and Nitrospira-like nitrite-oxidizing
522   bacterial communities under long-term fertilization practices. Soil Biology and Biochemistry
523   124:118–125. https://doi.org/10.1016/j.soilbio.2018.05.033

524   72.   Ji L, Wu Z, You Z, et al (2018) Effects of organic substitution for synthetic N fertilizer on soil
525   bacterial diversity and community composition: A 10-year field trial in a tea plantation.
526   Agriculture, Ecosystems & Environment 268:124–132.
527   https://doi.org/10.1016/j.agee.2018.09.008

528   73.   Kamaa M, Mburu H, Blanchart E, et al (2011) Effects of organic and inorganic fertilization on
529   soil bacterial and fungal microbial diversity in the Kabete long-term trial, Kenya. Biol Fertil Soils
530   47:315–321. https://doi.org/10.1007/s00374-011-0539-3

531   74.   Kumar U, Shahid M, Tripathi R, et al (2017) Variation of functional diversity of soil microbial
532   community in sub-humid tropical rice-rice cropping system under long-term organic and

inorganic   fertilization.   Ecological   Indicators   73:536–543.
https://doi.org/10.1016/j.ecolind.2016.10.014

75.   Kumar U, Kumar Nayak A, Shahid M, et al (2018) Continuous application of inorganic and
organic fertilizers over 47 years in paddy soil alters the bacterial community structure and its
influence on rice production. Agriculture, Ecosystems & Environment 262:65–75.
https://doi.org/10.1016/j.agee.2018.04.016

76.   Li J, Cooper JM, Lin Z, et al (2015) Soil microbial community structure and function are
significantly affected by long-term organic and mineral fertilization regimes in the North China
Plain. Applied Soil Ecology 96:75–87. https://doi.org/10.1016/j.apsoil.2015.07.001

77.   Liu W, Wang Q, Wang B, et al (2015) Changes in the abundance and structure of bacterial
communities under long-term fertilization treatments in a peanut monocropping system. Plant
Soil 395:415–427. https://doi.org/10.1007/s11104-015-2569-3

78.   Nautiyal CS, Chauhan PS, Bhatia CR (2010) Changes in soil physico-chemical properties and
microbial functional diversity due to 14 years of conversion of grassland to organic agriculture in
semi-arid   agroecosystem.   Soil   and   Tillage   Research   109:55–60.
https://doi.org/10.1016/j.still.2010.04.008

79.   Qiu S-L, Wang L-M, Huang D-F, Lin X-J (2014) Effects of fertilization regimes on tea yields,
soil fertility, and soil microbial diversity. Chilean journal of agricultural research 74:333–339.
https://doi.org/10.4067/S0718-58392014000300012

80.   Sun HY, Deng SP, Raun WR (2004) Bacterial Community Structure and Diversity in a Century-
Old   Manure-Treated   Agroecosystem.   Appl   Environ   Microbiol   70:5868–5874.
https://doi.org/10.1128/AEM.70.10.5868-5874.2004

81.   van Bruggen AHC, Sharma K, Kaku E, et al (2015) Soil health indicators and Fusarium wilt
suppression in organically and conventionally managed greenhouse soils. Applied Soil Ecology
86:192–201. https://doi.org/10.1016/j.apsoil.2014.10.014

82.   van der Bom F, Nunes I, Raymond NS, et al (2018) Long-term fertilisation form, level and
duration affect the diversity, structure and functioning of soil microbial communities in the field.
Soil Biology and Biochemistry 122:91–103. https://doi.org/10.1016/j.soilbio.2018.04.003

83.   Wang Q, Jiang X, Guan D, et al (2018) Long-term fertilization changes bacterial diversity and
bacterial communities in the maize rhizosphere of Chinese Mollisols. Applied Soil Ecology
125:88–96. https://doi.org/10.1016/j.apsoil.2017.12.007

84.   Wei D, Yang Q, Zhang J-Z, et al (2008) Bacterial Community Structure and Diversity in a Black
Soil as Affected by Long-Term Fertilization*1 *1Project supported by the Heilongjiang
Provincial Natural Science Funds for Distinguished Young Scholars, China (No. JC200622), the
Heilongjiang Provincial Natural Science Foundation of China (No. C2004-17), and the National
"Eleventh Five Years Plan" Key Project on Science and Technology of China (No.
2006BAD25B05). Pedosphere 18:582–592. https://doi.org/10.1016/S1002-0160(08)60052-1

85.   Wen Y, Xiao J, Liu F, et al (2018) Contrasting effects of inorganic and organic fertilisation
regimes on shifts in Fe redox bacterial communities in red soils. Soil Biology and Biochemistry
117:56–67. https://doi.org/10.1016/j.soilbio.2017.11.003

86.   Yao L, Wang D, Kang L, et al (2018) Effects of fertilizations on soil bacteria and fungi
communities in a degraded arid steppe revealed by high through-put sequencing. PeerJ 6:e4623.
https://doi.org/10.7717/peerj.4623

87.   Zhong W, Gu T, Wang W, et al (2010) The effects of mineral fertilizer and organic manure on
soil microbial community and diversity. Plant Soil 326:511–522. https://doi.org/10.1007/s11104-
009-9988-y

TABLES

Table 1. Summary statistics for meta-analysis of functional diversity (H_{fun}) differences between

fertilizer treatments and control. One study [28] reported three comparisons, hence a multi-level random

model was employed to control for within-study effects. Studies (N) refers to the number of studies and

number of comparisons. LRM is the log-transformed ratio of means ± SE and 95 % Confidence Interval.

Difference (%) is the effect-size back-transformed to percentage difference.

Comparison	Studies (N)	LRM	LRM (95% CI)	Difference (%)	Difference (95% CI)
NPK vs. CON	6 (8)	0.027 ± 0.004	0.018, 0.037	2.8	2.0, 3.6
ORG vs. CON	7 (9)	0.068 ± 0.012	0.041, 0.095	7.0	4.6, 9.5
ORG vs. NPK	7 (9)	0.037 ± 0.014	0.004, 0.070	3.8	0.9, 6.7

589   Table 2. Summary statistics for meta-analysis of taxonomic diversity. Statistics as in Table 1.

Comparison	Class	Studies (N)	LRM	LRM (95% CI)	Change (%)	Change % (95 % CI)
NPK vs. CON	Fungi	6 (7)	0.014 ± 0.025	-0.038, 0.066	1.4	-3.5, 6.6
	Prokaryotes	19 (24)	-0.025 ± 0.022	-0.071, 0.021	-2.4	-6.7, 2.0
ORG vs. CON	Fungi	6 (8)	0.018 ± 0.022	-0.027, 0.062	1.8	-2.5, 6.2
	Prokaryotes	19 (24)	0.043 ± 0.014	0.015, 0.072	4.4	1.6, 7.4
ORG vs. NPK	Fungi	6 (8)	-0.042 ± 0.027	-0.098, 0.012	-4.2	-9.2, 1.1
	Prokaryotes	24 (34)	0.066 ± 0.022	0.021, 0.110	7.0	2.3, 11.5

592   FIGURES

594   Fig. 1. Comparison of H_{fun} (a-c) and H_{tax} (d-f) in (a,d) NPK vs. CON, (b,e) ORG vs. CON, (c,f) ORG

595   vs. NPK. Points show reported values with size proportional to log(1/variance). Results for Fungi are

596   in blue, Bacteria in red. Gel electrophoresis results are circles, sequencing results are triangles, TRFLP

597   are diamonds.

Fig. 2. Functional soil diversity differences. a) NPK vs. CON, b) ORG vs. CON, c) ORG vs. NPK. Horizontal bars and ticks show 95% confidence intervals and means for effect sizes from the log-transformed ratio of means, back-transformed to give percentage differences. Summary estimates are given at the bottom of each plot. Random intercepts were fitted per Study.

Fig. 3. Taxonomic diversity effect sizes. a) NPK vs. CON, b) ORG vs. CON, c) ORG vs. NPK. Horizontal bars and ticks show 95% confidence intervals and means for effect sizes from the log-transformed ratio of means, back-transformed to give percentage differences. Summary estimates are given at the bottom of each plot, for Fungi (blue) and Bacteria (red). Random intercepts were fitted per Study.

Fig. 4. Representation of processes influencing organic fertilizer affects on soil microbial diversity. See

Discussion for details.

SUPPLEMENTARY MATERIAL

SUPPLEMENTARY FIGURES

Fig. S1. Effect of fertilization on soil pH. a) Mean difference between NPK and CON vs. mean CON

pH. Blue solid and dashed lines show mean difference (-0.59 ± 0.19 SE, Z = -3.09, p = 0.002) and 95

% CI (-0.97, -0.22) of meta-analysis fit. b) Mean difference between ORG and CON vs. mean CON

pH. Blue solid and dashed lines show mean and 95% CI of meta-regression fit. Slope mean -0.27 ±

0.087, Z = -3.08, p = 0.002, 95% CI = -0.44, -0.10. Points show means from individual studies with

size proportional to log(1/variance).

Fig. S2. Funnel plots for soil pH meta-analysis. Kendall's τ and p-values are given for rank correlation

tests for funnel plot asymmetry. a) NPK vs. CON (τ = 0.14, p = 0.50), b) ORG vs. CON (τ = 0.25, p =

0.24).

Fig. S3. Funnel plots for functional diversity meta-analysis. Kendall's τ and p-values are given for rank
correlation tests for funnel plot asymmetry. a) NPK vs. CON (τ = 0.00, p = 1.00), b) ORG vs. CON (τ
= -0.11, p = 0.76), c) ORG vs. NPK (τ = -0.16, p = 0.91).

Fig. S4. Funnel plots for taxonomic diversity meta-analysis. Kendall's τ and p-values are given for rank
correlation tests for funnel plot asymmetry. a) NPK vs. CON (τ = -0.10, p = 0.44), b) ORG vs. CON (τ
= 0.02, p = 0.91), c) ORG vs. NPK (τ = 0.12, p = 0.26).

SUPPLEMENTARY TABLES

Table S1. Summary of studies included in the meta-analysis (alphabetical order).

Reference	Location	Measure	Methodology
[60]	Tamil Nadu, India	Taxonomic	16S seq
[61]	Maharashtra, India	Functional	Biolog Ecoplate
[62]	Jiangxi, China	Functional	Biolog Ecoplate
[63]	Jiangsu, China	Taxonomic	18S seq
[64]	Hunan, China	Taxonomic	16S seq
[65]	Heilongjiang, China	Taxonomic	16S seq
[66]	Heilongjiang, China	Taxonomic	ITS seq
[67]	Fengqiu, China	Taxonomic	16S seq
[68]	Sichuan, China	Taxonomic	16S DGGE
[69]	Jilin, China	Taxonomic	16S DGGE
		Functional	Biolog Ecoplate
[70]	Manitoba, Canada	Taxonomic	16S seq
[71]	Hunan, China	Taxonomic	nxr seq
[33]	Northeast China	Taxonomic	16S seq
[72]	Fujian, China	Taxonomic	16S seq
[73]	Kabete, Kenya	Taxonomic	16S and 28S DGGE
[74]	Odisha, India	Functional	Biolog Ecoplate
[75]	Odisha, India	Taxonomic	16S seq
[76]	Shandong, China	Taxonomic	16S DGGE
[77]	Jiangxi, China	Taxonomic	16S seq
[34]	Hebei, China	Taxonomic	TRFLP
[32]	Uttarakhand, India	Taxonomic	Soil dilution plate
[78]	Maharashtra, India	Functional	Biolog Ecoplate
[79]	Fujian, China	Taxonomic	16S and 18S TGGE
[28]	Linz, Austria	Taxonomic	16S PGGE
		Functional	Biolog Ecoplate
[80]	Oklahoma, USA	Taxonomic	16S DGGE
[81]	Netherlands	Taxonomic	16S DGGE
[82]	Denmark	Taxonomic	16S seq
[83]	Heilongjiang, China	Taxonomic	16S seq
[84]	Heilongjiang, China	Taxonomic	16S DGGE
		Functional	Biolog Ecoplate
[85]	Hunan, China	Taxonomic	16S seq
[86]	Inner Mongolia, China	Taxonomic	16S and 18S seq
[87]	Jiangxi, China	Functional	Biolog Ecoplate

648   Table S2. Meta-analysis diagnostics.

Test	I^2 (%)	Q	df	P
H_{fun} NPK vs. CON	75.8	16.5	7	0.0210
H_{fun} ORG vs. CON	77.8	29.1	8	0.0003
H_{fun} ORG vs. NPK	76.6	42.3	8	<0.0001
H_{tax} NPK vs. CON	99.1	570.0	29	<0.0001
H_{tax} ORG vs. CON	97.7	256.6	30	<0.0001
H_{tax} ORG vs. NPK	99.3	874.6	40	<0.0001
pH_{NPK} - pH_{CON} vs. pH_{CON}	99.0	1353.5	11	<0.0001
pH_{ORG} - pH_{CON} vs. pH_{CON}	94.5	10.9	10	0.0009