A meta-analysis of the effect of organic and mineral fertilizers on soil microbial diversity

Daniel P. Bebber[1] & Victoria R. Richards

[1] Department of Biosciences, College of Life and Environmental Sciences, Geoffrey Pope Building,

University of Exeter, Stocker Road, Exeter, EX4 4QD, UK.

Author for correspondence: Daniel P. Bebber (d.bebber@exeter.ac.uk)

ABSTRACT

Organic agriculture, employing manures or composts, has been proposed as a way of mitigating

undesirable impacts of mineral fertilizer use. Of particular interest is the effect of fertilizer regime on

soil microbes, which are key to nutrient cycling, plant health and soil structure. However, the effect of

fertilizers on soil microbial diversity remains poorly understood. Since biological diversity is an

important determinant of ecosystem function and a fundamental metric in community ecology, the

effects of fertilizer regimes on soil microbial diversity are of theoretical and applied interest. Here, we

conduct a meta-analysis of 37 studies reporting microbial diversity metrics in mineral fertilized (NPK),

organically fertilized (ORG) and unfertilized control (CON) soils. Of these studies, 32 reported

taxonomic diversity derived from sequencing, gradient gel electrophoresis, or RFLP. Functional

diversity, derived from Biolog Ecoplate^{TM} measures of carbon substrate metabolism, was reported in 8

studies, with 3 studies reporting both diversity metrics. Bacterial and archaeal diversity was reported in

28 taxonomic studies, and fungal diversity in 8 taxonomic studies. We found that functional diversity

was 2.8 % greater in NPK compared with CON, 7.0 % greater in ORG vs CON, and 3.8 % greater in

ORG vs NPK. Bacterial and archaeal taxonomic diversity was not significantly different between NPK

and CON, but on average 2.9% greater in ORG vs CON, and 2.4 % greater in ORG vs. NPK. Fungal

taxonomic diversity was not significantly different between any treatment pairs. There was very high

residual heterogeneity in all meta-analyses of soil diversity, suggesting that a large amount of further

research is required to fully understand the influence of fertilizer regimes on microbial diversity and

ecosystem function.

Keywords: Soil; Microbiome; Diversity; Organic; Agriculture; Nitrogen

INTRODUCTION

Diversity plays a key role in the resilience and adaptability of complex systems (Page, 2011), and

biological diversity has been central to understanding of the structure and function of ecological

communities (Ricklefs and Schluter, 1994). Human activities are rapidly eroding global biodiversity

(Ceballos et al., 2015), hence understanding how human activities influence diversity and how negative

impacts can be avoided is an important goal of applied ecology (Rudd et al., 2011; Sutherland et al.,

2006). Recent advances in environmental DNA sequencing and metabarcoding have revealed enormous

and unexpected microbial diversity in all habitats, particularly soils (Lloyd et al., 2018; Thompson et

al., 2017). Given the fundamental importance of soils in terrestrial ecosystems, agriculture and food

security, there is growing interest in the role of microbial diversity in processes such as nutrient cycling

(Delgado-Baquerizo et al., 2016), and how soil diversity can be managed for maintenance of ecosystem

services (Lemanceau et al., 2015). Soil health, the capacity of soils to function as a living system and

sustain and promote plant and animal communities, is linked to microbial diversity (Kibblewhite et al.,

2008; van Bruggen et al., 2019). Microbial diversity is correlated with soil ecosystem multifunctionality

including plant productivity, microbial biomass, availability of nitrate, ammonium and phosphorus, and

nitrogen mineralisation rates (Delgado-Baquerizo et al., 2016). In agriculture, soil microbes are critical

to desirable functions such as nutrient cycling, carbon storage, erosion control via soil aggregation, and

disease suppression (Mazzola, 2004; Rillig et al., 2002; Sahu et al., 2017). A large fraction of the

world's agricultural soils are in poor and deteriorating condition largely due to agricultural activities

(Wuepper et al., 2020). Therefore, understanding how to manage soils for microbial diversity could

help to prevent further deterioration.

Modern intensive, or conventional, agricultural methods, including application of mineral and chemical

fertilisers, regular tillage, and use of synthetic pesticides and herbicides, aim to increase soil nutrition

and suppress harmful species to produce higher crop yields, but these methods are environmentally

damaging (Reganold and Wachter, 2016). For example, growing pressure on agricultural land has

resulted in soil nutrient depletion and soil erosion (Wuepper et al., 2020). Mineral fertilizers supply

nitrogen (N) as ammonium nitrate or urea, phosphorus (P) and potassium (K), with around 100 Tg N

applied globally each year (FAO, 2020). Typically less than half the N applied is taken up by crops, the

remainder contributing to water pollution and the release of NOX greenhouse gases (Zhang et al., 2015).

Organic agriculture, which applies manure- or compost-based fertilizers and soil conditioners and does

not employ chemical pest controls (with certain exceptions), has been proposed as a means of both

increasing soil nutrition and reducing environmental impacts (Luo et al., 2018; Reganold and Wachter,

2016), but evidence for organic benefits remains equivocal. For example, eutrophication potential

appears greater in many organic than conventional crop systems (Clark and Tilman, 2017)

Soils are exceedingly complex and biologically-diverse ecosystems, varying in physico-chemical and

biological composition across spatial scales (Howe et al., 2014; Thompson et al., 2017). Meta-analyses

of multiple individual studies can reveal general trends and patterns in this complexity and determine

the effects of management interventions. For example, meta-analysis of observational data suggest that

biological richness (measured by DNA sequences) is greatest at neutral pH and at a mean temperature

of 10 °C (Thompson et al., 2017). Meta-analysis of experimental studies suggests that pH has the

greatest influence on soil microbial diversity among global change factors, alpha diversity rising with

pH; that nitrogen and NPK have negative or non-significant effects on alpha diversity depending on

microbial group, with significant negative influences of N alone on diversity in agricultural soils; that

soil functionality, defined as the range of biogeochemical processes carried out by soils, increases with

N and NPK; and that changes in diversity are negatively correlated with changes in functionality,

perhaps due to functional redundancy (Zhou et al., 2020). Other meta-analyses have demonstrated that

microbial diversity (Venter et al., 2016) and biomass (McDaniel et al., 2014) increase in crop rotations

compared with monocultures. A meta-analysis of experimental studies showed that organic agriculture

greatly increases microbial biomass carbon, microbial biomass nitrogen, and enzymatic activity

compared with conventional systems (Lori et al., 2017). Another meta-analysis found that organic

amendments increase crop yields by supporting soil microbial activity, but this did not investigate

effects on soil microbial diversity (Luo et al., 2018). A small meta-analysis found no significant effect

of organic agriculture on soil organism diversity, but this only included five studies and occurred before

widespread use of sequencing to soil microbial diversity (Bengtsson et al., 2005). However, despite

growing application of sequencing technologies in microbial ecology, the effects of organic and mineral

fertilizers on measures of soil microbial diversity remain unsynthesized.

Here, we conduct a meta-analysis quantifying the effects of organic fertilizers (manures and manure

composts) and mineral fertilizers (NPK) on soil microbial diversity, in comparison with unfertilized

controls. We compare results for taxonomic and functional diversity, and investigate the influence of

factors such as soil chemistry and duration of organic treatment on these effects. We hypothesise that

organic inputs will increase microbial diversity as compared to mineral fertilizer inputs, based on the

observation that organic fertilizers increase microbial biomass and enzymatic activity compared with

NPK (Lori et al., 2017). Ecological theory and observational data suggest that soil microbial diversity

has a hump-shaped relationship with microbial biomass, with fungal and bacterial diversity increasing

with biomass for all but the highest biomass levels (Bastida et al., 2021).

MATERIALS AND METHODS

Criteria for meta-analysis

We used meta-analysis to test for differences in soil microbial functional (H_{fun}) and taxonomic (H_{tax})

diversity between soils treated with manure-based organic fertilizer (ORG), mineral fertilizer (NPK)

and control (CON). Here, H_{tax} is defined as Shannon's diversity index calculated from the relative

abundance of species, operational taxonomic units (OTUs) or amplicon sequence variants (ASVs;

Callahan et al., 2017) in a sample. Shannon's diversity index was reported in all valid studies, hence

we do not consider any other diversity indices (e.g. Simpson's) that were reported additionally in a

small number of studies (e.g. Ma et al., 2018). H_{fun} is defined as Shannon's diversity calculated from

the relative conversion rates of various carbon sources in Biolog Ecoplates (e.g. Ros et al., 2006). We

conducted a literature search using combinations of the terms 'soil', 'organic', 'agriculture',

'microbial', 'diversity', 'bacteria*', 'fung*', 'communit*', 'fertilizer', 'manure' and 'compost' (where

* indicates a wildcard search, where appropriate) on Google Scholar, Scopus and Web of Science

between June 2021 and October 2021. Additionally, the reference lists of the papers were browsed to

find potentially appropriate studies which were not identified during the online literature search. All

potentially appropriate papers identified for the meta-analysis came from peer-reviewed journals.

Papers were considered eligible for the analysis if they met the following criteria, adapted from Lori et

al. (2017): Comparisons of the farming systems should be pairwise, meaning the organic- and mineral-

fertilised treatments were subject to the same climatic conditions before and during sampling; The

organic treatment must have been applied for a minimum of two consecutive years prior to soil

sampling, and the treatments must be defined by the study; Results must report the mean Shannon

diversity index (H) per treatment, uncertainty of the mean (either standard error of the mean or standard

deviation of the sample distribution) and sample size (n); experiments were conducted in open fields or

under cover, not in pots or containers; the study must have been published no earlier than the year 2000

because of the lack of comparable methods before this date; the mineral and organic treatments must

be clearly described (description of fertilizer, application rate). Non-manure organic fertilizers were not

included (e.g. Wang et al., 2016). We only included experimental studies in which the only variable

was fertilizer treatment, and NPK and organic treatments were applied. If there were several different

organic or conventional treatments within the same study, all appropriate combinations were reported

and treated as individual comparisons but with non-independent errors (see meta-analysis methods).

Results of combination treatments (e.g. manure plus NPK fertilizer) were not included (e.g. Zhao et al.,

2014).

For all valid studies, we extracted mean, SE or SD, and sample size for H, for control, NPK and organic

treatments. Where SE was reported, this was converted to SD by multiplying by the square root of

sample size. When data were presented in graphical form, we extracted means and errors using the

online tool WebPlotDigitalizer (https://automeris.io/WebPlotDigitizer/). Additionally we abstracted the

following variables: study location; reported soil type, duration of organic treatment, organic treatment

type (i.e. source of manure), organic treatment application rate (mass per hectare per year or equivalent

nitrogen mass per hectare per year); mineral fertilizer application rate (nitrogen mass per hectare per

year); crops grown; taxonomic group of microbes analysed (e.g. all, bacteria, fungi); median soil depth

sampled (usually the mid-point of the range of depths sampled); brief summary of the methodology

used (e.g. 16S sequencing, 16S DGGE, Biolog Ecoplate). Where reported, we abstracted mean and SE

of soil pH, and mean and SE of available soil nitrogen content (mg kg^{-1}) which was often reported as

the concentration of ammonium and nitrate.

Meta-analysis

Meta-analysis was conducted using the *metafor* package v. 2.4–0 for R v. 4.0.2 (Viechtbauer, 2010).

Effect size was calculated as the log−transformed ratio of means, $\log(y_{exp}/y_{con})$ where y_{exp} and y_{con} are

the means of the experimental treatment and control, respectively (Hedges et al., 1999). There was at

least one pairwise comparison of H in each study. However, several studies reported data from more

than one pairwise comparison, with multiple organic and/or conventional fertilisation treatments

investigated. In addition, several studies were conducted at the same research site. We fitted multi-level

random− and mixed−effects (i.e. testing for the effects of moderator variables) models by restricted

maximum likelihood, with random intercepts per location. Other similar meta-analyses have ignored

potential non-independence among results from the same study or location and treated each reported

mean as an independent replicate (e.g. Lori et al., 2017). Standard errors and confidence limits for

parameters were estimated from the t–distribution. We report the I^2 statistic and Q-test for heterogeneity

(Higgins and Thompson, 2002), and provide funnel plots rank correlation tests for funnel plot

asymmetry (Viechtbauer, 2010).

RESULTS

We abstracted data from 37 studies which met our requirements (Table S1). Our search criteria provided

a long-list of 267 studies, of which 199 were rejected on the basis of the abstract, leaving a shortlist of

68. A further 31 were rejected after detailed reading of the full text. Reasons for the rejection of studies

included: no reporting of diversity metrics (e.g. reporting of species richness); no or unclear reporting

of standard error or standard deviation for diversity metrics; no manure used in organic treatment; no

organic treatment alone (e.g. manure only applied in combination with NPK); no NPK treatment; no

experimental treatments (e.g. comparison of different farms under organic and conventional

treatments); lack of detail in description of treatment types or levels.

Around half (21) of the included studies reported one set of comparisons, with either or both of

unfertilized control (CON) and mineral fertilizer (NPK), 10 reported two comparisons, and five studies

reported up to six comparisons, giving 65 comparisons in total. The majority of treatments were

replicated three or four times. The functional diversity studies, and a single study reporting soil dilution

plate assays (Mahanta et al., 2017), did not differentiate between fungi, bacteria and archaea. The single

soil dilution plate assay was omitted from the meta-analysis due to the low sensitivity of the method

and is not considered further here. In that study, poultry manure treatment resulted in significantly

greater microbial diversity than inorganic fertilizer treatment (Mahanta et al., 2017). In the taxonomic

diversity studies, bacteria were the most commonly analysed group (26 studies, including one of

actinomycetes only, one of bacteria and archaea, and one of nitrogen-fixing bacteria), followed by fungi

(8 studies, including one of arbuscular mycorrhizal fungi only). Most studies were conducted in China

(24), followed by India (6), with single studies from Austria, Canada, Denmark, Kenya, Korea, the

Netherlands, and the USA.

Soil types were not reported according to any standard taxonomy. Soils were sampled to a median depth

of up to 15 cm (median 10.0 cm, IQR 7.5 - 10.0 cm). Soil nitrogen content in unfertilized plots was

reported in 16 studies (mean mg N kg^{-1} median 36.3, IQR 27.7 – 86.3). Manure-based organic fertilizers

were applied in all studies, derived variously from cattle, pigs, poultry, horses, sheep, or mixtures of

these. Composted manures were employed in 9 studies, fresh manures in 26 studies, and one study

employed a mix. A small number of studies reported more than one manure treatment or level (X. Hu

et al., 2018; Liu et al., 2019; Mahanta et al., 2017). Application rates were reported as either total mass

per area per year (30 studies, median 16.8 t ha^{-1} y^{-1}, IQR 5.1 – 19.6 t ha^{-1} y^{-1}) and/or as nitrogen addition

(18 studies, median 103.3 kg ha^{-1} y^{-1}, IQR 75.0 – 223.8 kg ha^{-1} y^{-1}). We did not attempt to estimate

nitrogen content in manures where this was not reported, due to the large vari ability in the nitrogen

fraction among different treatments. The reported nitrogen fraction varied from 0.4 % in horse manure

to 2.7 % in pig manure, with overall median 1.0 % (IQR 0.7 – 2.0 %) among the treatments reported in

the studies. The level of nitrogen in the manure treatment was similar to that in corresponding NPK

fertilizer treatments, where reported (absolute percentage difference median 0.1 %, IQR 0.0 – 29.9 %).

The duration of organic treatment was right-skewed, but many studies reported treatment periods of

several decades (median 22 y, IQR 13 – 34 y). A variety of crops were grown, either as monocultures

or in mixtures, the most common being maize (12 studies), wheat (10 studies), and rice (6 studies). Soil

pH in unfertilized plots was reported in 18 studies (mean pH median 6.4, interquartile range 5.9 - 7.5).

Meta-regression showed that soil pH declined under NPK treatment compared to control soil (Fig. S1a).

Manure-based fertilizers increased pH in acid soils and reduced pH in alkaline soils (Fig. S1b). Funnel

plots showed no asymmetry (Fig. S2). There was a large degree of residual heterogeneity in the meta-

analyses of soil pH (Table S2).

H_{fun} was reported by 8 studies, all employing Biolog Ecoplates (Table S1). H_{fun} varied between 1.12 and

4.61 with values in three clusters comprising a single study from Austria with $H_{fun} \sim 1.2$, four studies

from China and one from India with $H_{fun} \sim 2.9$, and two studies from India with $H_{fun} \sim 4.5$ (Fig. 1a-c).

H_{fun} increased by an average of 2.8 % in NPK vs. CON, 7.0 % in ORG vs. CON, and 3.8 % in ORG vs.

NPK (Fig. 2, Table 1). We found no influence of the duration of organic amendment on the change in

H_{fun} compared with control (LRM change per year of organic treatment -0.0008 ± 0.0013 y^{-1}, t = -0.58,

p = 0.56) or NPK (0.0006 ± 0.0007 y^{-1}, t = 0.84, p = 0.40) treatments. We found no effect of the level

of NPK addition relative to control (-0.0013 ± 0.0009, t = -1.42, p = 0.15), nor of level of organic

fertilizer addition quantity (0.0000 ± 0.0000, t = -0.35, p = 0.72) or nitrogen equivalent relative to

control (0.0001 ± 0.0005, t = 0.18, p = 0.86). We found marginally-significant effect of control soil pH

on the effect size of ORG vs. CON (-0.057 ± 0.018, t = -3.2, df = 2, p = 0.088), but not of soil pH on

NPK vs. CON (0.0055 ± 0.0432, t = 0.128, df = 2, p = 0.91). Funnel plots did not exhibit significant

asymmetry (Fig. S3). There was a large degree of residual heterogeneity in all meta-analyses of H_{fun} (I^2

> 75 %, Table S2).

H_{tax} was reported in 31 studies, employing DNA amplicon sequencing (21 studies), various forms of

gradient gel electrophoresis (9 studies), and T-RFLP (1 study). H_{tax} tended to be higher in bacteria and

archaea (hereafter, prokaryotes) than fungi, and in sequencing compared with DGGE methods (Fig. 1d-

f). Median H_{tax} for prokaryotes was 6.40 using sequencing and 2.63 using DGGE, in controls. Median

H_{tax} for fungi was 3.95 using sequencing and 1.62 using DGGE, in controls. H_{tax} of NPK-fertilized soils

did not differ from control for prokaryotes or fungi (Fig. 3, Table 2). H_{tax} of prokaryotes was

significantly greater in ORG soils than both NPK and CON (Fig. 3, Table 2). There was no significant

difference in fungal H_{tax} between ORG and CON or NPK (Fig. 3, Table 2). Funnel plots did not exhibit

significant asymmetry (Fig. S4). There was a large degree of residual heterogeneity in meta-analyses

of H_{tax} ($I^2 > 95$ %, Table S2).

DISCUSSION

We found significant, but varying, effects of both NPK and organic fertilizers on soil microbe

taxonomic and functional diversity. Taxonomic diversity in organic treatments compared with NPK

was greater for prokaryotes but not significantly different for fungi. We found no significant difference

in taxonomic diversity between NPK and control. Functional diversity was significantly greater in NPK

compared with control, and in organic treatment compared with NPK. However, the diversity

differences we found were small, indicating that choice of fertilization has marginal effects on this

measure of soil microbial community structure. A quantitative review on the effects of fertilizers on

soil fungal diversity found that mineral fertilizers tended to reduce diversity while organic fertilizers

had no significant effect (Ye et al., 2020), although the analysis did not consider sample sizes or

variances of study data and so the findings are difficult to interpret.

A large meta-analysis of the impacts of global change factors found that N addition alone significantly

reduced soil bacterial but not fungal Shannon diversity, particularly in agricultural systems (Zhou et al.,

2020). The same meta-analysis did not detect significant effects of NPK overall, but the effects of

organic fertilizers were not investigated. The most important moderator was found to be pH, which

strongly controlled the response ratio of diversity in response to global change factors. In line with our

findings, N and NPK treatments were found to reduce pH. In contrast, we found a buffering effect of

manure fertilizers on soil pH, as reported in earlier studies (reviewed in Köninger et al., 2021). This

buffering effect, along with nutrient and organic matter content, are considered the main benefits of

manure fertilizers for soil biodiversity (Köninger et al., 2021).

Several factors must be considered when interpreting the effects of fertilizers on soil microbial

functional and taxonomic diversity (Fig. 4). First, while in many studies the amount of nitrogen applied

in organic and NPK treatments was similar, farmyard manure is far more physically, chemically and

biologically complex. In addition to NPK, manures contain undigested plant matter (lignin, cellulose,

hemicellulose), lipids, carbohydrates, proteins, and nutritive elements (e.g. magnesium, iron,

manganese, zinc, copper) (Levi-Minzi et al., 1986). If structural complexity and heterogeneity are

enhanced by the addition of organic matter across spatial scales (Lehmann et al., 2008), this could in

itself increase microbial diversity through provision of ecospace (Vos et al., 2013). Provision of

additional energy (organic carbon) and micronutrients in manures could sustain a greater diversity of

microbes via the species-energy hypothesis which predicts that more species can be sustained in

ecosystems supporting more individuals (Clarke and Gaston, 2006), since manure increases soil

microbial biomass more than NPK (Lori et al., 2017). However, one potential risk of manure fertilizers

is that high concentrations of heavy metals such as zinc and copper can lead to soil accumulation and

reductions in microbial diversity (Köninger et al., 2021).

Second, microbial communities may be indirectly influenced by changes in crop plants. While organic

agriculture tends to be less productive in terms of crop yields (de Ponti et al., 2012; Hijbeek et al.,

2017), there is some evidence that root development is greater in organic farming (T. Hu et al., 2018).

Plant rhizospheres tend to harbour the greatest microbial diversity of all terrestrial ecosystems

(Thompson et al., 2017), hence it is possible that enhanced root development and exudation of organic

compounds influence the soil diversity measured in these studies. Third, there is the gut microbial

community residing within the manure itself. Both fresh and composted manures have high fungal and

prokaryote diversity, with community composition changing as composting proceeds (Meng et al.,

2019). Composting can reduce the presence of undesirable microbes in manure, for example those

carrying antimicrobial resistance genes (Gou et al., 2018), but gut microbes (including human

pathogens) can survive for long periods in compost-amended soils (Sharma and Reynnells, 2016).

Hence, increased soil microbial diversity in organic systems may be due to persistence of gut microbes.

Few studies reported both taxonomic and functional diversity, hence we were unable to determine a

relationship between these two metrics. Microbial functional diversity tends to be considered from the

perspective of potential functions inferred from gene sequences (Escalas et al., 2019). In contrast, we

report results from studies of the diversity of actual functions (metabolization of carbon sources) carried

out by the soil microbial community. The relationship between soil taxonomic diversity and ecosystem

function (as opposed to functional diversity) tends to be positive (Bardgett and Putten, 2014; Maron et

al., 2018; Philippot et al., 2013). Soil ecosystem multifunctionality (the capacity of soils to sustain many

functions simultaneously) increases with microbial diversity (Delgado-Baquerizo et al., 2016), while

taxonomic and functional gene diversity are closely correlated (Zhang et al., 2019). However,

taxonomic and functional diversity can be decoupled by the process of horizontal gene transfer, because

different taxa can perform similar tasks through shared genes (Zhang et al., 2019). In addition, the

presence of physiologically-distinct subgroups within OTUs commonly defined by > 97 % sequence

similarity (so-called 'microdiversity') means that functional diversity can be large in groups of

apparently identical microbial taxa (Larkin and Martiny, 2017).

Organic farming has been promoted as a more sustainable alternative to conventional agriculture,

because of greater energy efficiency, potentially closed nutrient cycles and increased biodiversity

(Reganold and Wachter, 2016). These potential benefits come at a cost of reduced productivity

compared with convential farming (de Ponti et al., 2012; Hijbeek et al., 2017), though there is some

evidence that organic production can eventually catch up with conventional yields and provides greater

spatial and temporal stability (Schrama et al., 2018). Environmental impacts of organic agriculture

include greater land use and eutrophication potential per unit of food produced, contradicting the aim

of closed nutrient cycles (Clark and Tilman, 2017). Our results suggest that, when considering organic

crop fertilization alone and as a strict alternative to NPK (rather than a mixed system of manure-derived

and NPK fertilizer), microbial functional diversity and bacterial taxonomic diversity are slightly, but

significantly greater in organic systems while the effect on fungal taxonomic diversity is unclear.

Despite the increases in microbial diversity under organic vs. conventional fertilization we found,

organic farming systems still often exhibit higher eutrophication potential than conventional systems

due to temporal mismatching between fertilizer addition and plant demand (Clark and Tilman, 2017).

Therefore, it appears that the diversity of the microbial community is likely not the most important

factor governing rates of nutrient loss from organic vs. conventional systems. While we did not compare

combined treatments (e.g. NPK with manure or compost) with single treatments, farmers commonly

apply a diversity of fertilizers. For example, a comparison of low-input organic systems with

conventional mixed (NPK and manure) and conventional NPK-fertilized systems in Switzerland found

the greatest bacterial α-diversity in the organic systems, followed by the mixed and NPK-only systems

(Hartmann et al., 2015). Some studies which were not included in the meta-analysis employed manure-

free organic treatments, finding no effect (Wu et al., 2015) or variable effects (Wang et al., 2016) on

soil microbial diversity.

Many studies addressing soil microbial diversity in response to fertilization could not be included in

our meta-analysis because of incomplete or unclear reporting. Without giving specific examples, we

found that many studies did not provide uncertainties (standard error or standard deviation) for

parameter estimates, gave unclear descriptions of the fertilisation routines and inputs, or reported

alternative measures of diversity. Of the studies we included, several did not report basic soil chemistry

metrics such as pH and nitrogen content. Given the importance of pH in determining soil microbial

diversity (Zhou et al., 2020), and the central relevance of nitrogen in these experiments, any future

research should report these variables at minimum. We were only able to include a relatively small

number of studies in our meta-analysis, though we note that in the Cochrane Library of medical meta-

analyses, the median number of studies is seven or below (von Hippel, 2015). The Cochrane

Collaboration has been instrumental in developing methods of systematic review and meta-analyses for

evidence-based medicine(Higgins et al., 2011). One reason for the lack of experimental comparisons of

organic and mineral fertilizer treatments could be that organic agriculture comprises only a tiny fraction,

around 1 per cent by area, of agriculture globally (Meemken and Qaim, 2018). We found very large

residual heterogeneity among effect sizes, i.e. most of the variation in effect sizes among studies

remains unexplained. Further research will reveal whether the mean effects we detected are general,

and what other factors help explain variation among soils, climates, locations and experimental

treatments. To achieve this, it is critical that detailed and complete data on key variables are reported.

In addition, better understanding of soil ecosystem functioning will be achieved via analysis of the taxa

and functional genes identified within samples, in addition to summary metrics like diversity (Hartmann

et al., 2015). In this way, the powerful new bioinformatics tools at our disposal can be harnessed to

fully understand the relationships between microbial communities and soil health in agricultural

systems.

DECLARATIONS

FUNDING

VR was funded by an undergraduate project studentship at Exeter University, UK. DB was funded by

EC Horizon 2020 project ID 727624.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest

AVAILABILITY OF DATA

The data used in the meta-analysis will be made available on publication of the manuscript.

CODE AVAILABILITY

Code used in the meta-analysis is available from the author by reasonable request.

AUTHORS CONTRIBUTIONS

DB developed the research. DB and VR collected data, analysed results and wrote the paper.

ETHICS APPROVAL

Not applicable

CONSENT TO PARTICIPATE

Not applicable

CONSENT FOR PUBLICATION

Not applicable

REFERENCES

Balachandar, D., Doud, M.S., Schneper, L., Mills, D., Mathee, K., 2014. Long-Term Organic Nutrient
Management Fosters the Eubacterial Community Diversity in the Indian Semi-arid Alfisol as
Revealed by Length Heterogeneity–PCR. Commun. Soil Sci. Plant Anal. 45, 189–203.
https://doi.org/10.1080/00103624.2013.841919
Bardgett, R.D., Putten, W.H. van der, 2014. Belowground biodiversity and ecosystem functioning.
Nature 515, 505. https://doi.org/10.1038/nature13855
Bastida, F., Eldridge, D.J., García, C., Kenny Png, G., Bardgett, R.D., Delgado-Baquerizo, M., 2021.
Soil microbial diversity–biomass relationships are driven by soil carbon content across global
biomes. ISME J. 15, 2081–2091. https://doi.org/10.1038/s41396-021-00906-0
Bengtsson, J., AhnströM, J., Weibull, A.-C., 2005. The effects of organic agriculture on biodiversity
and abundance: a meta-analysis: Organic agriculture, biodiversity and abundance. J. Appl.
Ecol. 42, 261–269. https://doi.org/10.1111/j.1365-2664.2005.01005.x
Callahan, B.J., McMurdie, P.J., Holmes, S.P., 2017. Exact sequence variants should replace operational
taxonomic   units   in   marker-gene   data   analysis.   ISME   J.   11,   2639–2643.
https://doi.org/10.1038/ismej.2017.119
Ceballos, G., Ehrlich, P.R., Barnosky, A.D., García, A., Pringle, R.M., Palmer, T.M., 2015. Accelerated
modern human–induced species losses: Entering the sixth mass extinction. Sci. Adv. 1,
e1400253. https://doi.org/10.1126/sciadv.1400253
Chaudhry, V., Rehman, A., Mishra, A., Chauhan, P.S., Nautiyal, C.S., 2012. Changes in Bacterial
Community Structure of Agricultural Land Due to Long-Term Organic and Chemical
Amendments. Microb. Ecol. 64, 450–460. https://doi.org/10.1007/s00248-012-0025-y
Chen, C., Zhang, J., Lu, M., Qin, C., Chen, Y., Yang, L., Huang, Q., Wang, J., Shen, Z., Shen, Q., 2016.
Microbial communities of an arable soil treated for 8 years with organic and inorganic
fertilizers. Biol. Fertil. Soils 52, 455–467. https://doi.org/10.1007/s00374-016-1089-5
Chen, X., Li, Z., Liu, M., Jiang, C., Che, Y., 2015. Microbial community and functional diversity
associated with different aggregate fractions of a paddy soil fertilized with organic manure
and/or   NPK   fertilizer   for   20 years.   J.   Soils   Sediments   15,   292–301.
https://doi.org/10.1007/s11368-014-0981-6
Clark, M., Tilman, D., 2017. Comparative analysis of environmental impacts of agricultural production
systems, agricultural input efficiency, and food choice. Environ. Res. Lett. 12, 064016.
https://doi.org/10.1088/1748-9326/aa6cd5
Clarke, A., Gaston, K.J., 2006. Climate, energy and diversity. Proc. R. Soc. B Biol. Sci. 273, 2257–
2266. https://doi.org/10.1098/rspb.2006.3545
Cui, X., Zhang, Y., Gao, J., Peng, F., Gao, P., 2018. Long-term combined application of manure and
chemical fertilizer sustained higher nutrient status and rhizospheric bacterial diversity in
reddish paddy soil of Central South China. Sci. Rep. 8, 1–11. https://doi.org/10.1038/s41598-
018-34685-0
de Ponti, T., Rijk, B., van Ittersum, M.K., 2012. The crop yield gap between organic and conventional
agriculture. Agric. Syst. 108, 1–9. https://doi.org/10.1016/j.agsy.2011.12.004
Delgado-Baquerizo, M., Maestre, F.T., Reich, P.B., Jeffries, T.C., Gaitan, J.J., Encinar, D., Berdugo,
M., Campbell, C.D., Singh, B.K., 2016. Microbial diversity drives multifunctionality in
terrestrial ecosystems. Nat. Commun. 7, 10541. https://doi.org/10.1038/ncomms10541
Ding, J., Jiang, X., Guan, D., Zhao, B., Ma, M., Zhou, B., Cao, F., Yang, X., Li, L., Li, J., 2017.
Influence of inorganic fertilizer and organic manure application on fungal communities in a
long-term field experiment of Chinese Mollisols. Appl. Soil Ecol. 111, 114–122.
https://doi.org/10.1016/j.apsoil.2016.12.003
Ding, J., Jiang, X., Ma, M., Zhou, B., Guan, D., Zhao, B., Zhou, J., Cao, F., Li, L., Li, J., 2016. Effect
of 35 years inorganic fertilizer and manure amendment on structure of bacterial and archaeal
communities in black soil of northeast China. Appl. Soil Ecol. 105, 187–195.
https://doi.org/10.1016/j.apsoil.2016.04.010
Ding, J., Ma, M., Jiang, X., Liu, Y., Zhang, J., Suo, L., Wang, L., Wei, D., Li, J., 2019. Effects of
applying inorganic fertilizer and organic manure for 35 years on the structure and diversity of

408   ammonia-oxidizing archaea communities in a Chinese Mollisols field. MicrobiologyOpen 9,
409    e00942. https://doi.org/10.1002/mbo3.942

410 Escalas, A., Hale, L., Voordeckers, J.W., Yang, Y., Firestone, M.K., Alvarez-Cohen, L., Zhou, J., 2019.
411    Microbial functional diversity: From concepts to applications. Ecol. Evol. 9, 12000–12016.
412    https://doi.org/10.1002/ece3.5670

413 FAO, 2020. FAOSTAT. Food and Agriculture Organization of the United Nations, Rome.

414 Ge, Y., Zhang, J., Zhang, L., Yang, M., He, J., 2008. Long-term fertilization regimes affect bacterial
415    community structure and diversity of an agricultural soil in northern China. J. Soils Sediments
416    8, 43–50. https://doi.org/10.1065/jss2008.01.270

417 Gou, M., Hu, H.-W., Zhang, Y.-J., Wang, J.-T., Hayden, H., Tang, Y.-Q., He, J.-Z., 2018. Aerobic
418    composting reduces antibiotic resistance genes in cattle manure and the resistome
419    dissemination in agricultural soils. Sci. Total Environ. 612, 1300–1310.
420    https://doi.org/10.1016/j.scitotenv.2017.09.028

421 Gu, Y., Zhang, X., Tu, S., Lindström, K., 2009. Soil microbial biomass, crop yields, and bacterial
422    community structure as affected by long-term fertilizer treatments under wheat-rice cropping.
423    Eur. J. Soil Biol. 45, 239–246. https://doi.org/10.1016/j.ejsobi.2009.02.005

424 Guanghua, W., Junjie, L., Xiaoning, Q., Jian, J., Yang, W., Xiaobing, L., 2008. Effects of fertilization
425    on bacterial community structure and function in a black soil of Dehui region estimated by
426    Biolog and PCR-DGGE methods. Acta Ecol. Sin. 28, 220–226. https://doi.org/10.1016/S1872-
427    2032(08)60023-2

428 Hamm, A.C., Tenuta, M., Krause, D.O., Ominski, K.H., Tkachuk, V.L., Flaten, D.N., 2016. Bacterial
429    communities of an agricultural soil amended with solid pig and dairy manures, and urea
430    fertilizer. Appl. Soil Ecol. 103, 61–71. https://doi.org/10.1016/j.apsoil.2016.02.015

431 Han, S., Zeng, L., Luo, X., Xiong, X., Wen, S., Wang, B., Chen, W., Huang, Q., 2018. Shifts in
432    Nitrobacter- and Nitrospira-like nitrite-oxidizing bacterial communities under long-term
433    fertilization practices. Soil Biol. Biochem. 124, 118–125.
434    https://doi.org/10.1016/j.soilbio.2018.05.033

435 Hartmann, M., Frey, B., Mayer, J., Mäder, P., Widmer, F., 2015. Distinct soil microbial diversity under
436    long-term organic and conventional farming. ISME J. 9, 1177–1194.
437    https://doi.org/10.1038/ismej.2014.210

438 Hedges, L.V., Gurevitch, J., Curtis, P.S., 1999. The Meta-Analysis of Response Ratios in Experimental
439    Ecology. Ecology 80, 1150–1156. https://doi.org/10.1890/0012-
440    9658(1999)080[1150:TMAORR]2.0.CO;2

441 Higgins, J.P.T., Altman, D.G., Gøtzsche, P.C., Jüni, P., Moher, D., Oxman, A.D., Savović, J., Schulz,
442    K.F., Weeks, L., Sterne, J.A.C., 2011. The Cochrane Collaboration's tool for assessing risk of
443    bias in randomised trials. BMJ 343, d5928. https://doi.org/10.1136/bmj.d5928

444 Higgins, J.P.T., Thompson, S.G., 2002. Quantifying heterogeneity in a meta-analysis. Stat. Med. 21,
445    1539–1558. https://doi.org/10.1002/sim.1186

446 Hijbeek, R., van Ittersum, M.K., ten Berge, H.F.M., Gort, G., Spiegel, H., Whitmore, A.P., 2017. Do
447    organic inputs matter – a meta-analysis of additional yield effects for arable crops in Europe.
448    Plant Soil 411, 293–303. https://doi.org/10.1007/s11104-016-3031-x

449 Howe, A.C., Jansson, J.K., Malfatti, S.A., Tringe, S.G., Tiedje, J.M., Brown, C.T., 2014. Tackling soil
450    diversity with the assembly of large, complex metagenomes. Proc. Natl. Acad. Sci. 111, 4904–
451    4909. https://doi.org/10.1073/pnas.1402564111

452 Hu, T., Sørensen, P., Wahlström, E.M., Chirinda, N., Sharif, B., Li, X., Olesen, J.E., 2018. Root biomass
453    in cereals, catch crops and weeds can be reliably estimated without considering aboveground
454    biomass. Agric. Ecosyst. Environ. 251, 141–148. https://doi.org/10.1016/j.agee.2017.09.024

455 Hu, X., Liu, J., Wei, D., Zhu, P., Cui, X., Zhou, B., Chen, X., Jin, J., Liu, X., Wang, G., 2018. Soil
456    Bacterial Communities Under Different Long-Term Fertilization Regimes in Three Locations
457    Across the Black Soil Region of Northeast China. Pedosphere 28, 751–763.
458    https://doi.org/10.1016/S1002-0160(18)60040-2

459 Ji, L., Wu, Z., You, Z., Yi, X., Ni, K., Guo, S., Ruan, J., 2018. Effects of organic substitution for
460    synthetic N fertilizer on soil bacterial diversity and community composition: A 10-year field

trial in a tea plantation. Agric. Ecosyst. Environ. 268, 124–132.
https://doi.org/10.1016/j.agee.2018.09.008
Kamaa, M., Mburu, H., Blanchart, E., Chibole, L., Chotte, J.-L., Kibunja, C., Lesueur, D., 2011. Effects
of organic and inorganic fertilization on soil bacterial and fungal microbial diversity in the
Kabete long-term trial, Kenya. Biol. Fertil. Soils 47, 315–321. https://doi.org/10.1007/s00374-
011-0539-3
Kibblewhite, M.G., Ritz, K., Swift, M.J., 2008. Soil health in agricultural systems. Philos. Trans. R.
Soc. B Biol. Sci. 363, 685–701. https://doi.org/10.1098/rstb.2007.2178
Köninger, J., Lugato, E., Panagos, P., Kochupillai, M., Orgiazzi, A., Briones, M.J.I., 2021. Manure
management and soil biodiversity: Towards more sustainable food systems in the EU. Agric.
Syst. 194, 103251. https://doi.org/10.1016/j.agsy.2021.103251
Kumar, U., Kumar Nayak, A., Shahid, M., Gupta, V.V.S.R., Panneerselvam, P., Mohanty, S., Kaviraj,
M., Kumar, A., Chatterjee, D., Lal, B., Gautam, P., Tripathi, R., Panda, B.B., 2018. Continuous
application of inorganic and organic fertilizers over 47 years in paddy soil alters the bacterial
community structure and its influence on rice production. Agric. Ecosyst. Environ. 262, 65–75.
https://doi.org/10.1016/j.agee.2018.04.016
Kumar, U., Shahid, M., Tripathi, R., Mohanty, S., Kumar, A., Bhattacharyya, P., Lal, B., Gautam, P.,
Raja, R., Panda, B.B., Jambhulkar, N.N., Shukla, A.K., Nayak, A.K., 2017. Variation of
functional diversity of soil microbial community in sub-humid tropical rice-rice cropping
system under long-term organic and inorganic fertilization. Ecol. Indic. 73, 536–543.
https://doi.org/10.1016/j.ecolind.2016.10.014
Larkin, A.A., Martiny, A.C., 2017. Microdiversity shapes the traits, niche space, and biogeography of
microbial taxa. Environ. Microbiol. Rep. 9, 55–70. https://doi.org/10.1111/1758-2229.12523
Lehmann, J., Solomon, D., Kinyangi, J., Dathe, L., Wirick, S., Jacobsen, C., 2008. Spatial complexity
of soil organic matter forms at nanometre scales. Nat. Geosci. 1, 238–242.
https://doi.org/10.1038/ngeo155
Lemanceau, P., Maron, P.-A., Mazurier, S., Mougel, C., Pivato, B., Plassart, P., Ranjard, L., Revellin,
C., Tardy, V., Wipf, D., 2015. Understanding and managing soil biodiversity: a major challenge
in agroecology. Agron. Sustain. Dev. 35, 67–81. https://doi.org/10.1007/s13593-014-0247-0
Levi-Minzi, R., Riffaldi, R., Saviozzi, A., 1986. Organic matter and nutrients in fresh and mature
farmyard manure. Agric. Wastes 16, 225–236. https://doi.org/10.1016/0141-4607(86)90068-5
Li, J., Cooper, J.M., Lin, Z., Li, Y., Yang, X., Zhao, B., 2015. Soil microbial community structure and
function are significantly affected by long-term organic and mineral fertilization regimes in the
North China Plain. Appl. Soil Ecol. 96, 75–87. https://doi.org/10.1016/j.apsoil.2015.07.001
Liu, W., Wang, Q., Wang, B., Wang, X., Franks, A.E., Teng, Y., Li, Z., Luo, Y., 2015. Changes in the
abundance and structure of bacterial communities under long-term fertilization treatments in a
peanut monocropping system. Plant Soil 395, 415–427. https://doi.org/10.1007/s11104-015-
2569-3
Liu, W., Zhang, Y., Jiang, S., Murray, P.J., Liao, L., Li, X., Zhang, J., 2019. Spatiotemporal differences
in the arbuscular mycorrhizal fungi communities in soil and roots in response to long-term
organic compost inputs in an intensive agricultural cropping system on the North China Plain.
J. Soils Sediments 19, 2520–2533. https://doi.org/10.1007/s11368-019-02244-3
Lloyd, K.G., Steen, A.D., Ladau, J., Yin, J., Crosby, L., 2018. Phylogenetically Novel Uncultured
Microbial Cells Dominate Earth Microbiomes. mSystems 3.
https://doi.org/10.1128/mSystems.00055-18
Lori, M., Symnaczik, S., Mäder, P., Deyn, G.D., Gattinger, A., 2017. Organic farming enhances soil
microbial abundance and activity—A meta-analysis and meta-regression. PLOS ONE 12,
e0180442. https://doi.org/10.1371/journal.pone.0180442
Luo, G., Li, L., Friman, V.-P., Guo, J., Guo, S., Shen, Q., Ling, N., 2018. Organic amendments increase
crop yields by improving microbe-mediated soil functioning of agroecosystems: A meta-
analysis. Soil Biol. Biochem. 124, 105–115. https://doi.org/10.1016/j.soilbio.2018.06.002
Ma, M., Jiang, X., Wang, Q., Ongena, M., Wei, D., Ding, J., Guan, D., Cao, F., Zhao, B., Li, J., 2018.
Responses of fungal community composition to long-term chemical and organic fertilization

strategies in Chinese Mollisols. MicrobiologyOpen 7, e00597.
https://doi.org/10.1002/mbo3.597

Mahanta, D., Bhattacharyya, R., Mishra, P.K., Gopinath, K.A., Channakeshavaih, C., Krishnan, J.,
Raja, A., Tuti, M.D., Varghese, E., Pandey, B.M., Bisht, J.K., Bhatt, J.C., 2017. Influence of a
six-year organic and inorganic fertilization on the diversity of the soil culturable microrgansims
in the Indian mid-Himalayas. Appl. Soil Ecol. 120, 229–238.
https://doi.org/10.1016/j.apsoil.2017.08.012

Maron, P.-A., Sarr, A., Kaisermann, A., Lévêque, J., Mathieu, O., Guigue, J., Karimi, B., Bernard, L.,
Dequiedt, S., Terrat, S., Chabbi, A., Ranjard, L., 2018. High Microbial Diversity Promotes Soil
Ecosystem Functioning. Appl Env. Microbiol 84, e02738-17.
https://doi.org/10.1128/AEM.02738-17

Mazzola, M., 2004. Assessment and Management of Soil Microbial Community Structure for Disease
Suppression. Annu. Rev. Phytopathol. 42, 35–59.
https://doi.org/10.1146/annurev.phyto.42.040803.140408

McDaniel, M.D., Tiemann, L.K., Grandy, A.S., 2014. Does agricultural crop diversity enhance soil
microbial biomass and organic matter dynamics? A meta-analysis. Ecol. Appl.
https://doi.org/10.1890/13-0616.1

Meemken, E.-M., Qaim, M., 2018. Organic Agriculture, Food Security, and the Environment. Annu.
Rev. Resour. Econ. 10, 39–63. https://doi.org/10.1146/annurev-resource-100517-023252

Meng, Q., Yang, W., Men, M., Bello, A., Xu, X., Xu, B., Deng, L., Jiang, X., Sheng, S., Wu, X., Han,
Y., Zhu, H., 2019. Microbial Community Succession and Response to Environmental Variables
During Cow Manure and Corn Straw Composting. Front. Microbiol. 10.
https://doi.org/10.3389/fmicb.2019.00529

Mensah, S., Veldtman, R., Assogbadjo, A.E., Kakaï, R.G., Seifert, T., 2016. Tree species diversity
promotes aboveground carbon storage through functional diversity and functional dominance.
Ecol. Evol. 6, 7546–7557. https://doi.org/10.1002/ece3.2525

Morelli, F., Benedetti, Y., Perna, P., Santolini, R., 2018. Associations among taxonomic diversity,
functional diversity and evolutionary distinctiveness vary among environments. Ecol. Indic.
88, 8–16. https://doi.org/10.1016/j.ecolind.2018.01.022

Nautiyal, C.S., Chauhan, P.S., Bhatia, C.R., 2010. Changes in soil physico-chemical properties and
microbial functional diversity due to 14 years of conversion of grassland to organic agriculture
in semi-arid agroecosystem. Soil Tillage Res. 109, 55–60.
https://doi.org/10.1016/j.still.2010.04.008

Page, S.E., 2011. Diversity and Complexity. Princeton University Press, Princeton.

Philippot, L., Spor, A., Hénault, C., Bru, D., Bizouard, F., Jones, C.M., Sarr, A., Maron, P.-A., 2013.
Loss in microbial diversity affects nitrogen cycling in soil. ISME J. 7, 1609–1619.
https://doi.org/10.1038/ismej.2013.34

Qiu, S.-L., Wang, L.-M., Huang, D.-F., Lin, X.-J., 2014. Effects of fertilization regimes on tea yields,
soil fertility, and soil microbial diversity. Chil. J. Agric. Res. 74, 333–339.
https://doi.org/10.4067/S0718-58392014000300012

Reganold, J.P., Wachter, J.M., 2016. Organic agriculture in the twenty-first century. Nat. Plants 2, 1–
8. https://doi.org/10.1038/nplants.2015.221

Ricklefs, R.E., Schluter, D. (Eds.), 1994. Species Diversity in Ecological Communities, 1. International
developments in the administration of publicly-funded forest research: Challenges and
opportunities 2. Payments for ecosystem services and their institutional dimensions:
Institutional frameworks and governance structures of PES schemes. University of Chicago
Press, Chicago.

Rillig, M.C., Wright, S.F., Eviner, V.T., 2002. The role of arbuscular mycorrhizal fungi and glomalin
in soil aggregation: comparing effects of five plant species. Plant Soil 238, 325–333.
https://doi.org/10.1023/A:1014483303813

Ros, M., Klammer, S., Knapp, B., Aichberger, K., Insam, H., 2006. Long-term effects of compost
amendment of soil on functional and structural diversity and microbial activity. Soil Use
Manag. 22, 209–218. https://doi.org/10.1111/j.1475-2743.2006.00027.x

Rudd, M.A., Beazley, K.F., Cooke, S.J., Fleishman, E., Lane, D.E., Mascia, M.B., Roth, R., Tabor, G., Bakker, J.A., Bellefontaine, T., Berteaux, D., Cantin, B., Chaulk, K.G., Cunningham, K., Dobell, R., Fast, E., Ferrara, N., Findlay, C.S., Hallstrom, L.K., Hammond, T., Hermanutz, L., Hutchings, J.A., Lindsay, K.E., Marta, T.J., Nguyen, V.M., Northey, G., Prior, K., Ramirez-Sanchez, S., Rice, J., Sleep, D.J.H., Szabo, N.D., Trottier, G., Toussaint, J.-P., Veilleux, J.-P., 2011. Generation of Priority Research Questions to Inform Conservation Policy and Management at a National Level. Conserv. Biol. 25, 476–484. https://doi.org/10.1111/j.1523-1739.2010.01625.x

Sahu, N., Vasu, D., Sahu, A., Lal, N., Singh, S.K., 2017. Strength of microbes in nutrient cycling: A key to soil health, in: Agriculturally Important Microbes for Sustainable Agriculture. https://doi.org/10.1007/978-981-10-5589-8_4

Sánchez-Bayo, F., Wyckhuys, K.A.G., 2019. Worldwide decline of the entomofauna: A review of its drivers. Biol. Conserv. 232, 8–27. https://doi.org/10.1016/j.biocon.2019.01.020

Schrama, M., de Haan, J.J., Kroonen, M., Verstegen, H., Van der Putten, W.H., 2018. Crop yield gap and stability in organic and conventional farming systems. Agric. Ecosyst. Environ. https://doi.org/10.1016/j.agee.2017.12.023

Sharma, M., Reynnells, R., 2016. Importance of Soil Amendments: Survival of Bacterial Pathogens in Manure and Compost Used as Organic Fertilizers. Microbiol. Spectr. 4, PFS-0010-2015. https://doi.org/10.1128/microbiolspec.PFS-0010-2015

Sun, H.Y., Deng, S.P., Raun, W.R., 2004. Bacterial Community Structure and Diversity in a Century-Old Manure-Treated Agroecosystem. Appl. Environ. Microbiol. 70, 5868–5874. https://doi.org/10.1128/AEM.70.10.5868-5874.2004

Sutherland, W.J., Armstrong-Brown, S., Armsworth, P.R., Tom, B., Brickland, J., Campbell, C.D., Chamberlain, D.E., Cooke, A.I., Dulvy, N.K., Dusic, N.R., Fitton, M., Freckleton, R.P., Godfray, H.C.J., Grout, N., Harvey, H.J., Hedley, C., Hopkins, J.J., Kift, N.B., Kirby, J., Kunin, W.E., Macdonald, D.W., Marker, B., Naura, M., Neale, A.R., Oliver, T., Osborn, D., Pullin, A.S., Shardlow, M.E.A., Showler, D.A., Smith, P.L., Smithers, R.J., Solandt, J.-L., Spencer, J., Spray, C.J., Thomas, C.D., Thompson, J., Webb, S.E., Yalden, D.W., Watkinson, A.R., 2006. The identification of 100 ecological questions of high policy relevance in the UK. J. Appl. Ecol. 43, 617–627. https://doi.org/10.1111/j.1365-2664.2006.01188.x

Thompson, L.R., Sanders, J.G., McDonald, D., Amir, A., Ladau, J., Locey, K.J., Prill, R.J., Tripathi, A., Gibbons, S.M., Ackermann, G., Navas-Molina, J.A., Janssen, S., Kopylova, E., Vázquez-Baeza, Y., González, A., Morton, J.T., Mirarab, S., Xu, Z.Z., Jiang, L., Haroon, M.F., Kanbar, J., Zhu, Q., Song, S.J., Kosciolek, T., Bokulich, N.A., Lefler, J., Brislawn, C.J., Humphrey, G., Owens, S.M., Hampton-Marcell, J., Berg-Lyons, D., McKenzie, V., Fierer, N., Fuhrman, J.A., Clauset, A., Stevens, R.L., Shade, A., Pollard, K.S., Goodwin, K.D., Jansson, J.K., Gilbert, J.A., Knight, R., Consortium, T.E.M.P., Rivera, J.L.A., Al-Moosawi, L., Alverdy, J., Amato, K.R., Andras, J., Angenent, L.T., Antonopoulos, D.A., Apprill, A., Armitage, D., Ballantine, K., Bárta, J., Baum, J.K., Berry, A., Bhatnagar, A., Bhatnagar, M., Biddle, J.F., Bittner, L., Boldgiv, B., Bottos, E., Boyer, D.M., Braun, J., Brazelton, W., Brearley, F.Q., Campbell, A.H., Caporaso, J.G., Cardona, C., Carroll, J., Cary, S.C., Casper, B.B., Charles, T.C., Chu, H., Claar, D.C., Clark, R.G., Clayton, J.B., Clemente, J.C., Cochran, A., Coleman, M.L., Collins, G., Colwell, R.R., Contreras, M., Crary, B.B., Creer, S., Cristol, D.A., Crump, B.C., Cui, D., Daly, S.E., Davalos, L., Dawson, R.D., Defazio, J., Delsuc, F., Dionisi, H.M., Dominguez-Bello, M.G., Dowell, R., Dubinsky, E.A., Dunn, P.O., Ercolini, D., Espinoza, R.E., Ezenwa, V., Fenner, N., Findlay, H.S., Fleming, I.D., Fogliano, V., Forsman, A., Freeman, C., Friedman, E.S., Galindo, G., Garcia, L., Garcia-Amado, M.A., Garshelis, D., Gasser, R.B., Gerdts, G., Gibson, M.K., Gifford, I., Gill, R.T., Giray, T., Gittel, A., Golyshin, P., Gong, D., Grossart, H.-P., Guyton, K., Haig, S.-J., Hale, V., Hall, R.S., Hallam, S.J., Handley, K.M., Hasan, N.A., Haydon, S.R., Hickman, J.E., Hidalgo, G., Hofmockel, K.S., Hooker, J., Hulth, S., Hultman, J., Hyde, E., Ibáñez-Álamo, J.D., Jastrow, J.D., Jex, A.R., Johnson, L.S., Johnston, E.R., Joseph, S., Jurburg, S.D., Jurelevicius, D., Karlsson, A., Karlsson, R., Kauppinen, S., Kellogg, C.T.E., Kennedy, S.J., Kerkhof, L.J., King, G.M., Kling, G.W., Koehler, A.V., Krezalek, M., Kueneman, J., Lamendella, R., Landon, E.M., Lane-deGraaf, K., LaRoche, J., Larsen, P.,

Laverock, B., Lax, S., Lentino, M., Levin, I.I., Liancourt, P., Liang, W., Linz, A.M., Lipson,
D.A., Liu, Y., Lladser, M.E., Lozada, M., Spirito, C.M., MacCormack, W.P., MacRae-Crerar,
A., Magris, M., Martín-Platero, A.M., Martín-Vivaldi, M., Martínez, L.M., Martínez-Bueno,
M., Marzinelli, E.M., Mason, O.U., Mayer, G.D., McDevitt-Irwin, J.M., McDonald, J.E.,
McGuire, K.L., McMahon, K.D., McMinds, R., Medina, M., Mendelson, J.R., Metcalf, J.L.,
Meyer, F., Michelangeli, F., Miller, K., Mills, D.A., Minich, J., Mocali, S., Moitinho-Silva, L.,
Moore, A., Morgan-Kiss, R.M., Munroe, P., Myrold, D., Neufeld, J.D., Ni, Y., Nicol, G.W.,
Nielsen, S., Nissimov, J.I., Niu, K., Nolan, M.J., Noyce, K., O'Brien, S.L., Okamoto, N.,
Orlando, L., Castellano, Y.O., Osuolale, O., Oswald, W., Parnell, J., Peralta-Sánchez, J.M.,
Petraitis, P., Pfister, C., Pilon-Smits, E., Piombino, P., Pointing, S.B., Pollock, F.J., Potter, C.,
Prithiviraj, B., Quince, C., Rani, A., Ranjan, R., Rao, S., Rees, A.P., Richardson, M., Riebesell,
U., Robinson, C., Rockne, K.J., Rodriguezl, S.M., Rohwer, F., Roundstone, W., Safran, R.J.,
Sangwan, N., Sanz, V., Schrenk, M., Schrenzel, M.D., Scott, N.M., Seger, R.L., Seguin-
Orlando, A., Seldin, L., Seyler, L.M., Shakhsheer, B., Sheets, G.M., Shen, C., Shi, Y., Shin,
H., Shogan, B.D., Shutler, D., Siegel, J., Simmons, S., Sjöling, S., Smith, D.P., Soler, J.J.,
Sperling, M., Steinberg, P.D., Stephens, B., Stevens, M.A., Taghavi, S., Tai, V., Tait, K., Tan,
C.L., Taş, N., Taylor, D.L., Thomas, T., Timling, I., Turner, B.L., Urich, T., Ursell, L.K.,
Lelie, D. van der, Treuren, W.V., Zwieten, L. van, Vargas-Robles, D., Thurber, R.V.,
Vitaglione, P., Walker, D.A., Walters, W.A., Wang, S., Wang, T., Weaver, T., Webster, N.S.,
Wehrle, B., Weisenhorn, P., Weiss, S., Werner, J.J., West, K., Whitehead, A., Whitehead, S.R.,
Whittingham, L.A., Willerslev, E., Williams, A.E., Wood, S.A., Woodhams, D.C., Yang, Y.,
Zaneveld, J., Zarraonaindia, I., Zhang, Q., Zhao, H., 2017. A communal catalogue reveals
Earth's multiscale microbial diversity. Nature 551, 457. https://doi.org/10.1038/nature24621

van Bruggen, A.H.C., Goss, E.M., Havelaar, A., van Diepeningen, A.D., Finckh, M.R., Morris, J.G.,
Jr, 2019. One Health - Cycling of diverse microbial communities as a connecting force for soil,
plant, animal, human and ecosystem health. Sci. Total Environ. 664, 927–937.
https://doi.org/10.1016/j.scitotenv.2019.02.091

van Bruggen, A.H.C., Sharma, K., Kaku, E., Karfopoulos, S., Zelenev, V.V., Blok, W.J., 2015. Soil
health indicators and Fusarium wilt suppression in organically and conventionally managed
greenhouse soils. Appl. Soil Ecol. 86, 192–201. https://doi.org/10.1016/j.apsoil.2014.10.014

van der Bom, F., Nunes, I., Raymond, N.S., Hansen, V., Bonnichsen, L., Magid, J., Nybroe, O., Jensen,
L.S., 2018. Long-term fertilisation form, level and duration affect the diversity, structure and
functioning of soil microbial communities in the field. Soil Biol. Biochem. 122, 91–103.
https://doi.org/10.1016/j.soilbio.2018.04.003

Venter, Z.S., Jacobs, K., Hawkins, H.J., 2016. The impact of crop rotation on soil microbial diversity:
A meta-analysis. Pedobiologia. https://doi.org/10.1016/j.pedobi.2016.04.001

Viechtbauer, W., 2010. Conducting Meta-Analyses in R with the metafor Package. J. Stat. Softw. 36,
1–48. https://doi.org/10.18637/jss.v036.i03

von Hippel, P.T., 2015. The heterogeneity statistic I2 can be biased in small meta-analyses. BMC Med.
Res. Methodol. 15, 35. https://doi.org/10.1186/s12874-015-0024-z

Vos, M., Wolf, A.B., Jennings, S.J., Kowalchuk, G.A., 2013. Micro-scale determinants of bacterial
diversity in soil. FEMS Microbiol. Rev. 37, 936–954. https://doi.org/10.1111/1574-6976.12023

Wang, L., Yang, F., E, Y., Yuan, J., Raza, W., Huang, Q., Shen, Q., 2016. Long-Term Application of
Bioorganic Fertilizers Improved Soil Biochemical Properties and Microbial Communities of
an Apple Orchard Soil. Front. Microbiol. 7. https://doi.org/10.3389/fmicb.2016.01893

Wang, Q., Jiang, X., Guan, D., Wei, D., Zhao, B., Ma, M., Chen, S., Li, L., Cao, F., Li, J., 2018. Long-
term fertilization changes bacterial diversity and bacterial communities in the maize
rhizosphere of Chinese Mollisols. Appl. Soil Ecol. 125, 88–96.
https://doi.org/10.1016/j.apsoil.2017.12.007

Wang, Y., Ji, H., Hu, Y., Wang, R., Rui, J., Guo, S., 2018. Different Selectivity in Fungal Communities
Between Manure and Mineral Fertilizers: A Study in an Alkaline Soil After 30 Years
Fertilization. Front. Microbiol. 9, 2613. https://doi.org/10.3389/fmicb.2018.02613

Wei, D., Yang, Q., Zhang, J.-Z., Wang, S., Chen, X.-L., Zhang, X.-L., Li, W.-Q., 2008. Bacterial
Community Structure and Diversity in a Black Soil as Affected by Long-Term Fertilization*1

*1Project supported by the Heilongjiang Provincial Natural Science Funds for Distinguished
Young Scholars, China (No. JC200622), the Heilongjiang Provincial Natural Science
Foundation of China (No. C2004-17), and the National "Eleventh Five Years Plan" Key Project
on Science and Technology of China (No. 2006BAD25B05). Pedosphere 18, 582–592.
https://doi.org/10.1016/S1002-0160(08)60052-1

Wen, Y., Xiao, J., Liu, F., Goodman, B.A., Li, W., Jia, Z., Ran, W., Zhang, R., Shen, Q., Yu, G., 2018.
Contrasting effects of inorganic and organic fertilisation regimes on shifts in Fe redox bacterial
communities in red soils. Soil Biol. Biochem. 117, 56–67.
https://doi.org/10.1016/j.soilbio.2017.11.003

Wu, T., Milner, H., Díaz-Pérez, J.C., Ji, P., 2015. Effects of soil management practices on soil microbial
communities and development of southern blight in vegetable production. Appl. Soil Ecol. 91,
58–67. https://doi.org/10.1016/j.apsoil.2015.02.011

Wuepper, D., Borrelli, P., Finger, R., 2020. Countries and the global rate of soil erosion. Nat. Sustain.
3, 51–55. https://doi.org/10.1038/s41893-019-0438-4

Yao, L., Wang, Dangjun, Kang, L., Wang, Dengke, Zhang, Y., Hou, X., Guo, Y., 2018. Effects of
fertilizations on soil bacteria and fungi communities in a degraded arid steppe revealed by high
through-put sequencing. PeerJ 6, e4623. https://doi.org/10.7717/peerj.4623

Ye, G., Lin, Y., Luo, J., Di, H.J., Lindsey, S., Liu, D., Fan, J., Ding, W., 2020. Responses of soil fungal
diversity and community composition to long-term fertilization: Field experiment in an acidic
Ultisol and literature synthesis. Appl. Soil Ecol. 145, 103305.
https://doi.org/10.1016/j.apsoil.2019.06.008

Zhang, Q., Buyantuev, A., Li, F.Y., Jiang, L., Niu, J., Ding, Y., Kang, S., Ma, W., 2017. Functional
dominance rather than taxonomic diversity and functional diversity mainly affects community
aboveground biomass in the Inner Mongolia grassland. Ecol. Evol. 7, 1605–1615.
https://doi.org/10.1002/ece3.2778

Zhang, X., Davidson, E.A., Mauzerall, D.L., Searchinger, T.D., Dumas, P., Shen, Y., 2015. Managing
nitrogen for sustainable development. Nature 528, 51–59. https://doi.org/10.1038/nature15743

Zhang, X., Johnston, E.R., Wang, Y., Yu, Q., Tian, D., Wang, Z., Zhang, Y., Gong, D., Luo, C., Liu,
W., Yang, J., Han, X., 2019. Distinct Drivers of Core and Accessory Components of Soil
Microbial Community Functional Diversity under Environmental Changes. mSystems 4.
https://doi.org/10.1128/mSystems.00374-19

Zhang, Y., Shen, H., He, X., Thomas, B.W., Lupwayi, N.Z., Hao, X., Thomas, M.C., Shi, X., 2017.
Fertilization Shapes Bacterial Community Structure by Alteration of Soil pH. Front. Microbiol.
8, 1325. https://doi.org/10.3389/fmicb.2017.01325

Zhao, J., Ni, T., Li, Y., Xiong, W., Ran, W., Shen, B., Shen, Q., Zhang, R., 2014. Responses of Bacterial
Communities in Arable Soils in a Rice-Wheat Cropping System to Different Fertilizer Regimes
and Sampling Times. PLOS ONE 9, e85301. https://doi.org/10.1371/journal.pone.0085301

Zhong, W., Gu, T., Wang, W., Zhang, B., Lin, X., Huang, Q., Shen, W., 2010. The effects of mineral
fertilizer and organic manure on soil microbial community and diversity. Plant Soil 326, 511–
522. https://doi.org/10.1007/s11104-009-9988-y

Zhou, Z., Wang, C., Luo, Y., 2020. Meta-analysis of the impacts of global change factors on soil
microbial diversity and functionality. Nat. Commun. 11, 3072. https://doi.org/10.1038/s41467-
020-16881-7

TABLES

Table 1. Summary statistics for meta-analysis of functional diversity (H_{fun}) differences between

fertilizer treatments and control. One study (Ros et al., 2006) reported three comparisons, hence a multi-

level random model was employed to control for within-study effects. Studies (N) refers to the number

of studies and number of comparisons. LRM is the log-transformed ratio of means ± SE and 95 %

Confidence Interval. Difference (%) is the effect-size back-transformed to percentage difference.

Comparison	Studies (N)	LRM	LRM (95% CI)	Difference (%)	Difference (95% CI)
NPK vs. CON	6 (8)	0.027 ± 0.004	0.018, 0.037	2.8	2.0, 3.6
ORG vs. CON	7 (9)	0.068 ± 0.012	0.041, 0.095	7.0	4.6, 9.5
ORG vs. NPK	7 (9)	0.037 ± 0.014	0.004, 0.070	3.8	0.9, 6.7

728   Table 2. Summary statistics for meta-analysis of taxonomic diversity. Statistics as in Table 1.

Comparison	Class	Studies (N)	LRM	LRM (95% CI)	Change (%)	Change % (95 % CI)
NPK vs. CON	Fungi	8 (10)	0.026 ± 0.016	-0.007, 0.058	2.6	-0.6, 5.9
	Prokaryotes	22 (29)	-0.011 ± 0.014	-0.017, 0.040	1.1	-1.7, 4.0
ORG vs. CON	Fungi	8 (10)	0.028 ± 0.014	-0.002, 0.055	2.7	-0.2, 5.7
	Prokaryotes	22 (29)	0.029 ± 0.008	0.011, 0.047	2.9	1.1, 4.8
ORG vs. NPK	Fungi	8 (10)	-0.017 ± 0.015	-0.048, 0.014	-4.2	-4.7, 1.4
	Prokaryotes	27 (39)	0.024 ± 0.011	0.001, 0.046	2.4	0.1, 4.7

731   FIGURES

733   Fig. 1. Comparison of H$_{fun}$ (a-c) and H$_{tax}$ (d-f) in (a,d) NPK vs. CON, (b,e) ORG vs. CON, (c,f) ORG

734   vs. NPK. Results for fungi are in blue, prokaryotes in red. Points show reported values with size

735   proportional to log(1/variance). Gel electrophoresis results are circles, sequencing results are triangles,

736   TRFLP are diamonds.

Fig. 2. Functional soil diversity differences. a) NPK vs. CON, b) ORG vs. CON, c) ORG vs. NPK. Horizontal bars and ticks show 95% confidence intervals and means for effect sizes from the log-transformed ratio of means, back-transformed to give percentage differences. Summary estimates are given at the bottom of each plot. Random intercepts were fitted per Study.

Fig. 3. Taxonomic diversity effect sizes. a) NPK vs. CON, b) ORG vs. CON, c) ORG vs. NPK. Horizontal bars and ticks show 95% confidence intervals and means for effect sizes from the log-transformed ratio of means, back-transformed to give percentage differences. Summary estimates are given at the bottom of each plot, for fungi (blue) and prokaryotes (red). Random intercepts were fitted per Study.

Fig. 4. Representation of processes potentially influencing fertilizer affects on soil microbial diversity. See Discussion for details. Graphical elements from https://svg-clipart.com/ and http://clipart-library.com/

SUPPLEMENTARY MATERIAL

SUPPLEMENTARY FIGURES

Fig. S1. Effect of fertilization on soil pH. a) Mean difference between NPK and CON vs. mean CON

pH. Blue solid and dashed lines show mean difference (-0.59 ± 0.19 SE, Z = -3.09, p = 0.002) and 95

% CI (-0.97, -0.22) of meta-analysis fit. b) Mean difference between ORG and CON vs. mean CON

pH. Blue solid and dashed lines show mean and 95% CI of meta-regression fit. Slope mean -0.27 ±

0.087, Z = -3.08, p = 0.002, 95% CI = -0.44, -0.10. Points show means from individual studies with

size proportional to log(1/variance).

Fig. S2. Funnel plots for soil pH meta-analysis. Kendall's τ and p-values are given for rank correlation

tests for funnel plot asymmetry. a) NPK vs. CON (τ = 0.14, p = 0.50), b) ORG vs. CON (τ = 0.25, p =

0.24).

Fig. S3. Funnel plots for functional diversity meta-analysis. Kendall's τ and p-values are given for rank

correlation tests for funnel plot asymmetry. a) NPK vs. CON (τ = 0.00, p = 1.00), b) ORG vs. CON (τ

= -0.11, p = 0.76), c) ORG vs. NPK (τ = -0.16, p = 0.91).

Fig. S4. Funnel plots for taxonomic diversity meta-analysis. Kendall's τ and p-values are given for rank

correlation tests for funnel plot asymmetry. a) NPK vs. CON (τ = -0.10, p = 0.44), b) ORG vs. CON (τ

= 0.02, p = 0.91), c) ORG vs. NPK (τ = 0.12, p = 0.26).

SUPPLEMENTARY TABLES

Table S1. Summary of studies included in the meta-analysis (alphabetical order).

Reference	Location	Measure	Methodology
(Balachandar et al., 2014)	Tamil Nadu, India	Taxonomic	16S seq
(Chaudhry et al., 2012)	Maharashtra, India	Functional	Biolog Ecoplate
(Chen et al., 2015)	Jiangxi, China	Functional	Biolog Ecoplate
(Chen et al., 2016)	Jiangsu, China	Taxonomic	18S seq
(Cui et al., 2018)	Hunan, China	Taxonomic	16S seq
(Ding et al., 2016)	Heilongjiang, China	Taxonomic	16S seq
(Ding et al., 2017)	Heilongjiang, China	Taxonomic	ITS seq
(Ding et al., 2019)	Heilongjiang, China	Taxonomic	*amoA* seq
(Ge et al., 2008)	Fengqiu, China	Taxonomic	16S seq
(Gu et al., 2009)	Sichuan, China	Taxonomic	16S DGGE
(Guanghua et al., 2008)	Jilin, China	Taxonomic	16S DGGE
		Functional	Biolog Ecoplate
(Hamm et al., 2016)	Manitoba, Canada	Taxonomic	16S seq
(Han et al., 2018)	Hunan, China	Taxonomic	nxr seq
(X. Hu et al., 2018)	Northeast China	Taxonomic	16S seq
(Ji et al., 2018)	Fujian, China	Taxonomic	16S seq
(Kamaa et al., 2011)	Kabete, Kenya	Taxonomic	16S and 28S DGGE
(Kumar et al., 2017)	Odisha, India	Functional	Biolog Ecoplate
(Kumar et al., 2018)	Odisha, India	Taxonomic	16S seq
(Li et al., 2015)	Shandong, China	Taxonomic	16S DGGE
(Liu et al., 2015)	Jiangxi, China	Taxonomic	16S seq
(Liu et al., 2019)	Hebei, China	Taxonomic	TRFLP
(Ma et al., 2018)	Heilongjiang, China	Taxonomic	ITS seq

(Mahanta et al., 2017)	Uttarakhand, India	Taxonomic	Soil dilution plate
(Nautiyal et al., 2010)	Maharashtra, India	Functional	Biolog Ecoplate
(Qiu et al., 2014)	Fujian, China	Taxonomic	16S and 18S TGGE
(Ros et al., 2006)	Linz, Austria	Taxonomic	16S PGGE
		Functional	Biolog Ecoplate
(Sun et al., 2004)	Oklahoma, USA	Taxonomic	16S DGGE
(van Bruggen et al., 2015)	Netherlands	Taxonomic	16S DGGE
(van der Bom et al., 2018)	Denmark	Taxonomic	16S seq
(Q. Wang et al., 2018)	Heilongjiang, China	Taxonomic	16S seq
(Y. Wang et al., 2018)	Shaanxi, China	Taxonomic	ITS seq
(Wei et al., 2008)	Heilongjiang, China	Taxonomic	16S DGGE
		Functional	Biolog Ecoplate
(Wen et al., 2018)	Hunan, China	Taxonomic	16S seq
(Yao et al., 2018)	Inner Mongolia, China	Taxonomic	16S and 18S seq
(Y. Zhang et al., 2017)	Chongqing, China	Taxonomic	16S seq
(Zhong et al., 2010)	Jiangxi, China	Functional	Biolog Ecoplate

787   Table S2. Meta-analysis diagnostics.

Test	I^2 (%)	Q	df	P
H_{fun} NPK vs. CON	73.5	16.5	7	0.0210
H_{fun} ORG vs. CON	84.6	29.1	8	0.0003
H_{fun} ORG vs. NPK	82.3	42.3	8	<0.0001
H_{tax} NPK vs. CON	96.6	570.0	29	<0.0001
H_{tax} ORG vs. CON	91.4	256.6	30	<0.0001
H_{tax} ORG vs. NPK	96.4	874.6	40	<0.0001
$pH_{NPK} - pH_{CON}$ vs. pH_{CON}	99.0	1353.5	11	<0.0001
$pH_{ORG} - pH_{CON}$ vs. pH_{CON}	94.5	10.9	10	0.0009