A meta-analysis of the effect of organic and mineral fertilizers on soil microbial diversity

Daniel P. Bebber[1] & Victoria R. Richards

[1] Department of Biosciences, College of Life and Environmental Sciences, Geoffrey Pope Building,

University of Exeter, Stocker Road, Exeter, EX4 4QD, UK.

Author for correspondence: Daniel P. Bebber (d.bebber@exeter.ac.uk)

ABSTRACT

The Green Revolution of agriculture was in part driven by application of synthetic mineral fertilizers,

largely supplanting organic manure as a source of the major nutrients nitrogen, phosphorous and

potassium (NPK). Though enhancing crop production and global food security, fertilizers have

contributed to soil acidification, eutrophication of water bodies, and greenhouse gas emissions. Organic

agriculture, employing manures or composts, has been proposed as a way of mitigating these

undesirable effects. Of particular interest is the effect of fertilizer regime on soil microbes, which are

key to nutrient cycling, plant health and soil structure. Meta-analyses of experimental studies indicate

that mineral fertilizer increases soil microbial biomass over unfertilized controls, and that organic

fertilizers increase microbial biomass and activity over mineral fertilizers. However, the effect of

fertilizers on soil microbial diversity remains poorly understood. Since biological diversity is an

important determinant of ecosystem function and a fundamental metric in community ecology, the

effects of fertilizer regimes on soil microbial diversity are of theoretical and applied interest. Here, we

conduct a meta-analysis of 31 studies reporting microbial diversity metrics in mineral fertilized (NPK),

organically fertilized (ORG) and unfertilized control (CON) soils. Of these studies, 26 reported

taxonomic diversity derived from sequencing, gradient gel electrophoresis, RFLP, or dilution plate

assay. Functional diversity, derived from Biolog Ecoplate™ measures of carbon substrate metabolism,

was reported in 8 studies, with 3 studies reporting both diversity metrics. We found that functional

diversity was on average 2.6 % greater in NPK compared with CON, 6.8 % greater in ORG vs CON

and 3.6 % greater in ORG vs NPK. Prokaryote taxonomic diversity was not significantly different

between NPK and CON, 4.2 % greater in ORG vs CON and 4.6 % greater in ORG vs. NPK. Fungal

taxonomic diversity was not significantly different between NPK or ORG vs CON, but 5.4 % lower

between ORG and NPK. There was very high residual heterogeneity in all meta-analyses of soil

diversity, suggesting that a large amount of further research with detailed analysis of soil properties is

required to fully understand the influence of fertilizer regimes on microbial diversity and ecosystem

function.

INTRODUCTION

Diversity plays a key role in the resilience and adaptability of complex systems (Page, 2011), and
biological diversity has been central to understanding of the structure and function of ecological
communities (Ricklefs & Schluter, 1994). Human activities are rapidly eroding global biodiversity
(Ceballos *et al*., 2015), hence understanding how human activities influence diversity and how negative
impacts can be avoided is an important goal of applied ecology (Sutherland *et al*., 2006; Rudd *et al*.,
2011). Recent advances in environmental DNA sequencing and metabarcoding have revealed enormous
and unexpected microbial diversity in all habitats, particularly soils (Thompson *et al*., 2017; Lloyd *et*
al., 2018). Given the fundamental importance of soils in terrestrial ecosystems, agriculture and food
security, there is growing interest in the role of microbial diversity in processes such as nutrient cycling
(Delgado-Baquerizo *et al*., 2016), and how soil diversity can be managed for maintenance of ecosystem
services (Lemanceau *et al*., 2015). In agriculture, soil microbes are critical to desirable functions such
as nutrient cycling, carbon storage, erosion control via soil aggregation, and disease suppression (Rillig
et al., 2002; Mazzola, 2004; Sahu *et al*., 2017). A large fraction of the world's agriculture soils are in
poor and deteriorating condition largely due to agricultural activities (Wuepper *et al*., 2020). Therefore,
understanding how to manage soils for microbial diversity could help to prevent further deterioration.

Modern intensive, or conventional, agricultural methods, including application of mineral and chemical
fertilisers, regular tillage, and use of synthetic pesticides and herbicides, aim to increase soil nutrition
and suppress harmful species to produce higher crop yields, but these methods are environmentally
damaging (Robertson, 2000). For example, growing pressure on agricultural land has resulted in soil
nutrient depletion and soil erosion (Wopereis, Tamèlokpo, & Ezui, 2006; Lal, 2009). Mineral fertilizers
supply nitrogen (N) as ammonium nitrate or urea, phosphorus (P) and potassium (K), with around 100
Tg N applied globally each year (FAO, 2020). The nitrogen sources are derived from the Haber-Bosch
process which reduces dinitrogen to ammonia, accounting for around one third of the nitrogen fixed
from the atmosphere (Fowler *et al*., 2013). Soil and plant-symbiotic microbes fix another third, marine
cyanobacteria another third, with small contributions from lightning and fossil fuel combustion (Fowler
et al., 2013). Typically less than half the N applied is taken up by crops, the remainder contributing to
water pollution and the release of NOX greenhouse gases (Zhang *et al*., 2015). Organic agriculture,
which applies manure- or compost-based fertilizers and soil conditioners and does not employ chemical
pest controls (with certain exceptions), has been proposed as a means of both increasing soil nutrition
and reducing environmental impacts (Reganold & Wachter, 2016), but evidence for organic benefits
remains equivocal. For example, eutrophication potential appears greater in many organic than
conventional crop systems (Clark & Tilman, 2017)

Soil health, the capacity of soils to function as a living system and sustain and promote plant and animal
communities, is linked to microbial diversity (Kibblewhite *et al*., 2008; van Bruggen *et al*., 2019).

Microbial diversity is correlated with soil ecosystem multifunctionality including plant productivity,
microbial biomass, availability of nitrate, ammonium and phosphorus, and nitrogen mineralisation rates
(Delgado-Baquerizo *et al*., 2016). Hence, management to enhance soil microbial diversity could
improve soil health. Soils are exceedingly complex and biologically-diverse ecosystems, varying in
physico-chemical and biological composition across spatial scales (Howe *et al*., 2014; Thompson *et al*.,
2017). Meta-analyses of multiple individual studies can reveal general trends and patterns in this
complexity and determine the effects of management interventions. For example, observational data
suggest that biological richness (measured by DNA sequences) is greatest at neutral pH and at a mean
temperature of 10 °C (Thompson *et al*., 2017). Meta-analysis of experimental studies shows that pH
has the greatest influence on soil microbial diversity among global change factors, alpha diversity rising
with pH (Zhou *et al*., 2020). Nitrogen and NPK have negative or non-significant effects on alpha
diversity depending on microbial group, with significant negative influences of N alone on diversity in
agricultural soils. Soil functionality, defined as the range of biogeochemical processes carried out by
soils, increases with N and NPK. Changes in diversity are negatively correlated with changes in
functionality, perhaps due to functional redundancy (Zhou *et al*., 2020). Other meta-analyses have
demonstrated that microbial diversity (Venter *et al*., 2016) and biomass (McDaniel *et al*., 2014) increase
in crop rotations compared with monocultures. Organic agriculture greatly increases microbial biomass
carbon, microbial biomass nitrogen, and enzymatic activity compared with conventional systems (Lori
et al., 2017). Previously, a small meta-analysis found no significant effect of organic agriculture on soil
organism diversity, but this only included five studies and occurred before widespread use of
sequencing to soil microbial diversity (Bengtsson *et al*., 2005). However, despite growing application
of sequencing technologies in microbial ecology, results on the effects of organic and mineral fertilizers
on measures of soil microbial diversity remain unsynthesized.

Here, we conduct a meta-analysis quantifying the effects of organic fertilizers (manures and manure
composts) and mineral fertilizers (NPK) on soil microbial diversity, in comparison with unfertilized
controls. We compare results for taxonomic and functional diversity, and investigate the influence of
factors such as soil chemistry and duration of organic treatment on these effects. We hypothesise that
organic inputs will increase microbial diversity as compared to mineral fertilizer inputs, based on the
observation that organic fertilizers increase microbial biomass and enzymatic activity compared with
NPK (Lori *et al*., 2017).

MATERIALS AND METHODS

Criteria for meta-analysis

We used meta-analysis to test for differences in soil microbial functional (H_{fun}) and taxonomic (H_{tax})

diversity between soils treated with manure-based organic fertilizer (ORG), mineral fertilizer (NPK)

and control (CON). Here, H_{tax} is defined as Shannon's diversity index calculated from the relative

abundance of species, operational taxonomic units (OTUs) or amplicon sequence variants (ASVs;

Callahan *et al*., 2017) in a sample. H_{fun} is defined as Shannon's diversity calculated from the relative

conversion rates of various carbon sources in Biolog Ecoplates (e.g. Ros *et al*., 2006). We conducted a

literature search using combinations of the terms 'soil', 'organic', 'agriculture', 'microbial', 'diversity',

'bacteria*', 'fung*', 'communit*', 'fertilizer', 'manure' and 'compost' (where * indicates a wildcard

search, where appropriate) on Google Scholar, Scopus and Web of Science between August 2018 and

June 2019. Additionally, the reference lists of the papers were browsed to find potentially appropriate

studies which were not identified during the online literature search. All potentially appropriate papers

identified for the meta-analysis came from peer-reviewed journals.

Papers were considered eligible for the analysis if they met the following criteria, adapted from Lori et

al. (2017): Comparisons of the farming systems should be pairwise, meaning the organic- and mineral-

fertilised treatments were subject to the same climatic conditions before and during sampling; The

organic treatment must have been applied for a minimum of two consecutive years prior to soil

sampling, and the treatments must be defined by the study; Results must report the mean Shannon

diversity index (H) per treatment, uncertainty of the mean (either standard error of the mean or standard

deviation of the sample distribution) and sample size (n); experiments were conducted in open fields or

under cover, not in pots or containers; the study must have been published no earlier than the year 2000;

the mineral and organic treatments must be clearly described (description of fertilizer, application rate).

If there were several different organic or conventional treatments within the same study, all appropriate

combinations were reported and treated as individual comparisons but with non-independent errors (see

meta-analysis methods). Results of combination treatments (e.g. manure plus NPK fertilizer) were not

included.

For all valid studies, we extracted mean, SE or SD, and sample size for H, for control, NPK and organic

treatments. Where SE was reported, this was converted to SD by multiplying by the square root of

sample size. When data were presented in graphical form, we extracted means and errors using the

online tool WebPlotDigitalizer (https://automeris.io/WebPlotDigitizer/). Additionally we abstracted the

following variables: study location; reported soil type, duration of organic treatment, organic treatment

type (i.e. source of manure), organic treatment application rate (mass per hectare per year or equivalent

nitrogen mass per hectare per year); mineral fertilizer application rate (nitrogen mass per hectare per

year); crops grow; taxonomic group of microbes analysed (e.g. all, bacteria, fungi); median soil depth

sampled (usually the mid-point of the range of depths sampled); brief summary of the methodology

used (e.g. 16S sequencing, 16S DGGE, Biolog Ecoplate). Where reported we abstracted mean and SE

of soil pH; and mean and SE of soil nitrogen content (mg kg^{-1}).

Meta-analysis

Meta-analysis was conducted using the *metafor* package v. 2.4–0 for R v. 4.0.2 (Viechtbauer, 2010).

Effect size was calculated as the log–transformed ratio of means (Hedges *et al*., 1999). There was at

least one pairwise comparison of H in each study. However, several studies reported data from more

than one pairwise comparison, with multiple organic and/or conventional fertilisation treatments

investigated. We fitted multi-level random– and mixed–effects (i.e. testing for the effects of moderator

variables) models by restricted maximum likelihood, with random intercepts per study. Other similar

meta-analyses have ignored potential non-independence among results from the same study and treated

each reported mean as an independent replicate (Lori *et al*., 2017). Standard errors and confidence limits

for parameters were estimated from the *t*–distribution. We report the I^2 statistic and *Q*-test for

heterogeneity (Higgins & Thompson, 2002), and provide funnel plots rank correlation tests for funnel

plot asymmetry (Viechtbauer, 2010).

RESULTS

We abstracted data from 31 studies which met our requirements (Table S1). Our search criteria provided

a long-list of 259 studies, of which 195 were rejected on the basis of the abstract, leaving a shortlist of

64. A further 33 were rejected after detailed reading of the full text. Around half (16) of the included

studies reported one set of comparisons, with either or both of unfertilized control (CON) and mineral

fertilizer (NPK), 10 reported two comparisons, and five studies reported up to six comparisons, giving

57 comparisons in total. The majority of treatments were replicated three or four times. Bacteria were

the most commonly analysed group (24 studies, including one of actinomycetes only, one of bacteria

and archaea, and one of nitrogen-fixing bacteria), followed by fungi (6 studies, including one of

arbuscular mycorrhizal fungi only). The functional diversity studies, and the single study reporting soil

dilution plate assays (Mahanta *et al*., 2017), were taxonomically non-specific. Most studies were

conducted in China (19), followed by India (6), with single studies from Austria, Canada, Denmark,

Kenya, the Netherlands, and the USA.

Soil types were not reported according to any standard taxonomy. Soils were sampled to a median depth

of up to 15 cm (median 10.0 cm, IQR 7.5 - 10.0 cm). Soil nitrogen content in unfertilized plots was

reported in 14 studies (mean mg N kg^{-1} median 77.9, IQR 28.5 - 99.6). Manure-based organic fertilizers

were applied in all studies, derived variously from cattle, pigs, poultry, horses, sheep, or mixtures of

these. Composted manures were employed in 8 studies, fresh manures in 24 studies (one study

employed both). A small number of studies reported more than one manure treatment or level (Mahanta

et al., 2017; Hu *et al*., 2018b; Liu *et al*., 2019). Application rates were reported as either total mass per

area per year (24 studies, median 10 t ha^{-1} y^{-1}, IQR 4.9 - 18.6 t ha^{-1} y^{-1}) and/or as nitrogen addition (15

studies, median 102.0 kg ha^{-1} y^{-1}, IQR 60.0 - 140.0 kg ha^{-1} y^{-1}). We did not attempt to estimate nitrogen

content in manures where this was not reported, due to the large variability in the nitrogen fraction

among different treatments. The reported nitrogen fraction varied from 0.4 % in horse manure to 2.7 %

in pig manure, with overall median 1.0 % (IQR 0.7 - 2.0 %) among the treatments reported in the

studies. The level of nitrogen in the manure treatment was similar to that in corresponding NPK

fertilizer treatments, where reported (absolute percentage difference median 13.4 %, IQR 0.0 - 56.0 %).

The duration of organic treatment was right-skewed, but many studies reported treatment periods of

several decades (median 20 y, IQR 11 - 34.5 y). A variety of crops were grown, either as monocultures

or in mixtures, the most common being maize (9 studies), wheat (7 studies), and rice (6 studies). Soil

pH in unfertilized plots was reported in 16 studies (mean pH median 6.5, interquartile range 6.0 - 7.7).

Meta-regression showed that soil pH declined under NPK treatment by -0.53 ± 0.20 compared to control

soil (Fig. S1a). Manure-based fertilizers increased pH in acid soils and reduced pH in alkaline soils

(Fig. S1b). Funnel plots showed no asymmetry (Fig. S2). There was a large degree of residual

heterogeneity in the meta-analyses of soil pH (Table S2).

H_{fun} was reported by 8 studies, all employing Biolog Ecoplates (Table S1). H_{fun} varied between 1.12 and

4.61 with values in three clusters comprising a single study from Austria with $H_{fun} \sim 1.2$, four studies

from China and one from India with $H_{fun} \sim 2.9$, and two studies from India with $H_{fun} \sim 4.5$ (Fig. 1a-c).

H_{fun} increased by an average of 2.6 % in NPK vs. CON, 6.8 % in ORG vs. CON, and 3.6 % in ORG vs.

NPK (Fig. 2, Table 1). We found no influence of the duration of organic amendment on the change in

H_{fun} compared with control (LRM change per year of organic treatment -0.0008 ± 0.0013 y^{-1}, t = -0.58,

p = 0.56) or NPK (0.0006 ± 0.0007 y^{-1}, t = 0.84, p = 0.40) treatments. We found no effect of the level

of NPK addition relative to control (-0.0013 ± 0.0009, t = -1.42, p = 0.15), nor of level of organic

fertilizer addition quantity (0.0000 ± 0.0000, t = -0.35, p = 0.72) or nitrogen equivalent relative to

control (0.0001 ± 0.0005, t = 0.18, p = 0.86). We found a marginally-significant effect of control soil

pH on the effect size of ORG vs. CON (-0.052 ± 0.025, t = -2.10, p = 0.036), but not of soil pH on NPK

vs. CON (0.0045 ± 0.0527, t = 0.085, p = 0.93). Soil N did not influence the effect size of ORG vs.

CON (0.000 ± 0.001, t = -0.01, p =0.99). Funnel plots did not exhibit significant asymmetry (Fig. S3).

There was a large degree of residual heterogeneity in all meta-analyses of H_{fun} ($I^2 > 75$ %, Table S2).

H_{tax} was reported in 31 studies, employing DNA amplicon sequencing (17 studies), various forms of

gradient gel electrophoresis (11 studies), soil dilution plate assay (1 study) and T-RFLP (1 study). H_{tax}

tended to be higher in prokaryotes than fungi, and in sequencing compared with DGGE methods (Fig.

1d-f). Median H_{tax} for prokaryotes was 6.36 using sequencing and 2.85 using DGGE. Median H_{tax} for

fungi was 3.36 using sequencing and 1.40 using DGGE. H_{tax} of NPK-fertilized soils did not differ from

control for prokaryotes or fungi, nor in the single study using dilution plate assays (Fig. 3, Table 2). H_{tax}

of prokaryotes was greater in ORG soils than both NPK-fertilized soils and control. Fungi were less

taxonomically diverse in ORG than NPK, but there was no significant difference in fungal H_{tax} between

ORG and CON. In the soil dilution plate assay, organically fertilized soils were more diverse than NPK

or CON. We found no evidence that duration of organic amendments (0.0006 ± 0.0009, t = 0.66, p =

0.51), pH (-0.0054 ± 0.0038, t = -1.42, p = 0.15) or soil N (-0.0008 ± 0.0004, t = -2.1, p = 0.051)

influenced the effect size of ORG vs. CON. Funnel plots did not exhibit significant asymmetry (Fig.

S4). There was a large degree of residual heterogeneity in meta-analyses of H_{tax} ($I^2 > 95$ %, Table S2).

DISCUSSION

We found significant, but varying, effects of both NPK and organic fertilizers on soil microbe

taxonomic and functional diversity. Taxonomic diversity in organic treatments compared with NPK

was greater for prokaryotes but lower for fungi, and was greater for organic treatments compared with

control for prokaryotes with no significant difference in fungi. We found no significant difference in

taxonomic diversity between NPK and control for either prokaryotes or microbes. Functional diversity

was significantly greater in NPK compared with control, and in organic treatment compared with NPK.

We did not detect any significant influence of factors such as duration or type of organic treatment, soil

chemistry or experimental methodology on effect sizes. Importantly, the residual heterogeneity in all

meta-analyses was very high.

Several factors must be considered when interpreting the effects of fertilizers on soil microbial

functional and taxonomic diversity (Fig. 4). First, while in many studies the amount of nitrogen applied

in organic and NPK treatments was similar, farmyard manure is far more physically, chemically and

biologically complex. In addition to NPK, manures contain undigested plant matter (lignin, cellulose,

hemicellulose), lipids, carbohydrates, proteins, and nutritive elements (e.g. magnesium, iron,

manganese, zinc, copper) (Levi-Minzi *et al*., 1986). If structural complexity and heterogeneity are

enhanced by the addition of organic matter across spatial scales (Lehmann *et al*., 2008), this could in

itself increase microbial diversity through provision of ecospace that reduces interspecific competition

(Vos *et al*., 2013). Provision of additional energy (organic carbon) and micronutrients in manures could

sustain a greater diversity of microbes via the species-energy hypothesis which predicts that more

species can be sustained in ecosystems supporting more individuals (Clarke & Gaston, 2006), since

manure increases soil microbial biomass more than NPK (Lori *et al*., 2017). Second, microbial

communities may be indirectly influenced by changes in crop plants. While organic agriculture tends

to be less productive in terms of crop yields (de Ponti *et al*., 2012; Hijbeek *et al*., 2017), there is some

evidence that root development is greater in organic farming (Hu *et al*., 2018a). Plant rhizospheres tend

to harbour the greatest microbial diversity of all terrestrial ecosystems (Thompson *et al*., 2017), hence

it is possible that enhanced root development and exudation of organic compounds influence the soil

diversity measured in these studies. Third, there is the gut microbial community residing within the

manure itself. Both fresh and composted manures have high fungal and prokaryote diversity, with

community composition changing as composting proceeds (Meng *et al*., 2019). Bacterial diversity has

been found to peak during the thermophilic phase of aerobic composting during which temperatures

exceeded 55 °C, while fungal diversity was lowest during this stage (Meng *et al*., 2019). Composting

can reduce the presence of undesirable microbes in manure, for example those carrying antimicrobial

resistance genes (Gou *et al*., 2018), but gut microbes (including human pathogens) can survive for long

periods in compost-amended soils (Sharma & Reynnells, 2016). Hence, increased soil microbial

diversity in organic systems may be due to persistence of gut microbes.

Few studies reported both taxonomic and functional diversity, hence we were unable to determine a

relationship between these two metrics. Functional diversity has been found to increase with species

diversity in plant communities (Mensah *et al*., 2016; Zhang *et al*., 2017), but less clearly in bird

communities (Morelli *et al*., 2018). The relationship between soil taxonomic diversity and ecosystem

function (as opposed to functional diversity) tends to be positive (Philippot *et al*., 2013; Bardgett &

Putten, 2014; Maron *et al*., 2018). Given that ecosystem functions like nitrogen cycling require multiple

interacting processes, a positive relationship between taxonomic diversity and functional diversity

might be expected. Soil ecosystem multifunctionality (the capacity of soils to sustain many functions

simultaneously) increases with microbial diversity (Delgado-Baquerizo *et al*., 2016), while taxonomic

and functional gene diversity are closely correlated (Zhang *et al*., 2019). However, taxonomic and

functional diversity can be decoupled by the process of horizontal gene transfer, because different taxa

can perform similar tasks through shared genes (Zhang *et al*., 2019). In addition, the presence of

physiologically-distinct subgroups within OTUs commonly defined by > 97 % sequence similarity (so-

called 'microdiversity') means that functional diversity can be large in groups of apparently identical

microbial taxa (Larkin & Martiny, 2017). Microbial functional diversity tends to be considered from

the perspective of potential functions inferred from gene sequences (Escalas *et al*., 2019). In contrast,

we report results from studies of the diversity of actual functions (metabolization of carbon sources)

carried out by the soil microbial community. These genotypic vs. phenotypic measures of functional

diversity are analogous to the fundamental and realized niches of species, whereby the potential

functions identified in gene sequences are likely to exceed the observed functions carried out by the

community under particular environmental conditions (Escalas *et al*., 2019). Discovering the

relationships between potential and observed functions of microbial communities is a key goal of

microbial ecology.

Organic farming has been promoted as a more sustainable alternative to conventional agriculture,

because of greater energy efficiency, potentially closed nutrient cycles and increased biodiversity

(Reganold & Wachter, 2016). Certainly, the rapid decline of insect populations around the world due

to intensive agriculture and pesticides, is an urgent impetus for reconsidering the way we produce food

(Sánchez-Bayo & Wyckhuys, 2019). These potential benefits come at a cost of reduced productivity

compared with convential farming (de Ponti *et al*., 2012; Hijbeek *et al*., 2017), though there is some

evidence that organic production can eventually catch up with conventional yields and provides greater

spatial and temporal stability (Schrama *et al*., 2018). Environmental impacts of organic agriculture

include greater land use and eutrophication potential per unit of food produced, contradicting the aim

of closed nutrient cycles (Clark & Tilman, 2017). Our results suggest that, when considering organic

crop fertilization alone and as a strict alternative to NPK (rather than a mixed system of manure-derived

and NPK fertilizer), microbial functional diversity and bacterial taxonomic diversity are slightly, but

significantly greater in organic systems while fungal taxonomic diversity is slightly but significantly

lower. The enhanced eutrophication potential of organic farming, due to temporal mismatching between

fertilizer addition and plant demand, is much larger than the increases in taxonomic or functional

diversity we found (Clark & Tilman, 2017). Hence, it appears that increased soil microbial diversity

under organic farming does not translate to improved nutrient cycling and retention. While we did not

compare combined treatments (e.g. NPK with manure or compost) with single treatments, farmers

commonly apply a diversity of fertilizers. For example, a comparison of low-input organic systems with

conventional mixed (NPK and manure) and conventional NPK-fertilized systems in Switzerland found

the greatest bacterial α-diversity in the organic systems, followed by the mixed and NPK-only systems

(Hartmann *et al*., 2015).

Many studies addressing soil microbial diversity in response to fertilization could not be included in

our meta-analysis because of incomplete or unclear reporting. Without giving specific examples, we

found that many studies did not provide uncertainties (standard error or standard deviation) for

parameter estimates, gave unclear descriptions of the fertilisation routines and inputs, or reported

alternative measures of diversity. Of the studies we included, several did not report basic soil chemistry

metrics such as pH and nitrogen content. Given the importance of pH in determining soil microbial

diversity (Zhou *et al*., 2020), and the central relevance of nitrogen in these experiments, any future

research should report these variables at minimum. We were only able to include a small number of

studies in our meta-analysis, which revealed very large residual heterogeneity among effect sizes, i.e.

most of the variation in effect sizes among studies remains unexplained. Further research will reveal

whether the mean effects we detected are general, and what other factors help explain variation among

soils, climates, locations and experimental treatments. To achieve this, it is critical that detailed and

complete data on key variables are reported. In addition, better understanding of soil ecosystem

functioning will be achieved via analysis of the taxa and functional genes identified within samples, in

addition to summary metrics like diversity (Hartmann *et al*., 2015). In this way, the powerful new

bioinformatics tools at our disposal can be harnessed to fully understand the relationships between

agriculture and soil health.

DATA AVAILABILITY

The data used in the meta-analysis will be made available on publication of the manuscript.

AUTHORS CONTRIBUTIONS

DB developed the research. DB and VR collected data, analysed results and wrote the paper.

FUNDING

VR was funded by an undergraduate project studentship at Exeter University, UK. DB was funded by

EC Horizon 2020 project ID 727624.

REFERENCES

Balachandar, D., Doud, M.S., Schneper, L., Mills, D. & Mathee, K. (2014) Long-Term Organic Nutrient
Management Fosters the Eubacterial Community Diversity in the Indian Semi-arid Alfisol as
Revealed by Length Heterogeneity–PCR. *Communications in Soil Science and Plant Analysis*,
45, 189–203.
Bardgett, R.D. & Putten, W.H. van der (2014) Belowground biodiversity and ecosystem functioning.
Nature, **515**, 505.
Bengtsson, J., AhnströM, J. & Weibull, A.-C. (2005) The effects of organic agriculture on biodiversity
and abundance: a meta-analysis: Organic agriculture, biodiversity and abundance. *Journal of*
Applied Ecology, **42**, 261–269.
van der Bom, F., Nunes, I., Raymond, N.S., Hansen, V., Bonnichsen, L., Magid, J., Nybroe, O. &
Jensen, L.S. (2018) Long-term fertilisation form, level and duration affect the diversity,
structure and functioning of soil microbial communities in the field. *Soil Biology and*
Biochemistry, **122**, 91–103.
van Bruggen, A.H.C., Goss, E.M., Havelaar, A., van Diepeningen, A.D., Finckh, M.R. & Morris, J.G.,
Jr (2019) One Health - Cycling of diverse microbial communities as a connecting force for soil,
plant, animal, human and ecosystem health. *Science of the Total Environment*, **664**, 927–937.
van Bruggen, A.H.C., Sharma, K., Kaku, E., Karfopoulos, S., Zelenev, V.V. & Blok, W.J. (2015) Soil
health indicators and Fusarium wilt suppression in organically and conventionally managed
greenhouse soils. *Applied Soil Ecology*, **86**, 192–201.
Callahan, B.J., McMurdie, P.J. & Holmes, S.P. (2017) Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. *The ISME Journal*, **11**, 2639–2643.
Ceballos, G., Ehrlich, P.R., Barnosky, A.D., García, A., Pringle, R.M. & Palmer, T.M. (2015)
Accelerated modern human–induced species losses: Entering the sixth mass extinction. *Science*
Advances, **1**, e1400253.
Chaudhry, V., Rehman, A., Mishra, A., Chauhan, P.S. & Nautiyal, C.S. (2012) Changes in Bacterial
Community Structure of Agricultural Land Due to Long-Term Organic and Chemical
Amendments. *Microbial Ecology*, **64**, 450–460.

Chen, C., Zhang, J., Lu, M., Qin, C., Chen, Y., Yang, L., Huang, Q., Wang, J., Shen, Z. & Shen, Q. (2016) Microbial communities of an arable soil treated for 8 years with organic and inorganic fertilizers. *Biology and Fertility of Soils*, **52**, 455–467.

Chen, X., Li, Z., Liu, M., Jiang, C. & Che, Y. (2015) Microbial community and functional diversity associated with different aggregate fractions of a paddy soil fertilized with organic manure and/or NPK fertilizer for 20 years. *Journal of Soils and Sediments*, **15**, 292–301.

Clark, M. & Tilman, D. (2017) Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. *Environmental Research Letters*, **12**, 064016.

Clarke, A. & Gaston, K.J. (2006) Climate, energy and diversity. *Proceedings of the Royal Society B: Biological Sciences*, **273**, 2257–2266.

Cui, X., Zhang, Y., Gao, J., Peng, F. & Gao, P. (2018) Long-term combined application of manure and chemical fertilizer sustained higher nutrient status and rhizospheric bacterial diversity in reddish paddy soil of Central South China. *Scientific Reports*, **8**, 1–11.

Delgado-Baquerizo, M., Maestre, F.T., Reich, P.B., Jeffries, T.C., Gaitan, J.J., Encinar, D., Berdugo, M., Campbell, C.D. & Singh, B.K. (2016) Microbial diversity drives multifunctionality in terrestrial ecosystems. *Nature Communications*, **7**, 10541.

Ding, J., Jiang, X., Guan, D., Zhao, B., Ma, M., Zhou, B., Cao, F., Yang, X., Li, L. & Li, J. (2017) Influence of inorganic fertilizer and organic manure application on fungal communities in a long-term field experiment of Chinese Mollisols. *Applied Soil Ecology*, **111**, 114–122.

Ding, J., Jiang, X., Ma, M., Zhou, B., Guan, D., Zhao, B., Zhou, J., Cao, F., Li, L. & Li, J. (2016) Effect of 35 years inorganic fertilizer and manure amendment on structure of bacterial and archaeal communities in black soil of northeast China. *Applied Soil Ecology*, **105**, 187–195.

Escalas, A., Hale, L., Voordeckers, J.W., Yang, Y., Firestone, M.K., Alvarez-Cohen, L. & Zhou, J. (2019) Microbial functional diversity: From concepts to applications. *Ecology and Evolution*, **9**, 12000–12016.

FAO (2020) *FAOSTAT*, Food and Agriculture Organization of the United Nations, Rome.

Fowler, D., Coyle, M., Skiba, U., Sutton, M.A., Cape, J.N., Reis, S., Sheppard, L.J., Jenkins, A., Grizzetti, B., Galloway, J.N., Vitousek, P., Leach, A., Bouwman, A.F., Butterbach-Bahl, K., Dentener, F., Stevenson, D., Amann, M. & Voss, M. (2013) The global nitrogen cycle in the twenty-first century. *Phil. Trans. R. Soc. B*, **368**, 20130164.

Ge, Y., Zhang, J., Zhang, L., Yang, M. & He, J. (2008) Long-term fertilization regimes affect bacterial community structure and diversity of an agricultural soil in northern China. *Journal of Soils and Sediments*, **8**, 43–50.

Gou, M., Hu, H.-W., Zhang, Y.-J., Wang, J.-T., Hayden, H., Tang, Y.-Q. & He, J.-Z. (2018) Aerobic composting reduces antibiotic resistance genes in cattle manure and the resistome dissemination in agricultural soils. *Science of The Total Environment*, **612**, 1300–1310.

Gu, Y., Zhang, X., Tu, S. & Lindström, K. (2009) Soil microbial biomass, crop yields, and bacterial community structure as affected by long-term fertilizer treatments under wheat-rice cropping. *European Journal of Soil Biology*, **45**, 239–246.

Guanghua, W., Junjie, L., Xiaoning, Q., Jian, J., Yang, W. & Xiaobing, L. (2008) Effects of fertilization on bacterial community structure and function in a black soil of Dehui region estimated by Biolog and PCR-DGGE methods. *Acta Ecologica Sinica*, **28**, 220–226.

Hamm, A.C., Tenuta, M., Krause, D.O., Ominski, K.H., Tkachuk, V.L. & Flaten, D.N. (2016) Bacterial communities of an agricultural soil amended with solid pig and dairy manures, and urea fertilizer. *Applied Soil Ecology*, **103**, 61–71.

Han, S., Zeng, L., Luo, X., Xiong, X., Wen, S., Wang, B., Chen, W. & Huang, Q. (2018) Shifts in Nitrobacter- and Nitrospira-like nitrite-oxidizing bacterial communities under long-term fertilization practices. *Soil Biology and Biochemistry*, **124**, 118–125.

Hartmann, M., Frey, B., Mayer, J., Mäder, P. & Widmer, F. (2015) Distinct soil microbial diversity under long-term organic and conventional farming. *The ISME Journal*, **9**, 1177–1194.

Hedges, L.V., Gurevitch, J. & Curtis, P.S. (1999) The Meta-Analysis of Response Ratios in Experimental Ecology. *Ecology*, **80**, 1150–1156.

Higgins, J.P.T. & Thompson, S.G. (2002) Quantifying heterogeneity in a meta-analysis. *Statistics in*
Medicine, **21**, 1539–1558.
Hijbeek, R., van Ittersum, M.K., ten Berge, H.F.M., Gort, G., Spiegel, H. & Whitmore, A.P. (2017) Do
organic inputs matter – a meta-analysis of additional yield effects for arable crops in Europe.
Plant and Soil, **411**, 293–303.
Howe, A.C., Jansson, J.K., Malfatti, S.A., Tringe, S.G., Tiedje, J.M. & Brown, C.T. (2014) Tackling
soil diversity with the assembly of large, complex metagenomes. *Proceedings of the National*
Academy of Sciences, **111**, 4904–4909.
Hu, T., Sørensen, P., Wahlström, E.M., Chirinda, N., Sharif, B., Li, X. & Olesen, J.E. (2018a) Root
biomass in cereals, catch crops and weeds can be reliably estimated without considering
aboveground biomass. *Agriculture, Ecosystems & Environment*, **251**, 141–148.
Hu, X., Liu, J., Wei, D., Zhu, P., Cui, X., Zhou, B., Chen, X., Jin, J., Liu, X. & Wang, G. (2018b) Soil
Bacterial Communities Under Different Long-Term Fertilization Regimes in Three Locations
Across the Black Soil Region of Northeast China. *Pedosphere*, **28**, 751–763.
Ji, L., Wu, Z., You, Z., Yi, X., Ni, K., Guo, S. & Ruan, J. (2018) Effects of organic substitution for
synthetic N fertilizer on soil bacterial diversity and community composition: A 10-year field
trial in a tea plantation. *Agriculture, Ecosystems & Environment*, **268**, 124–132.
Kamaa, M., Mburu, H., Blanchart, E., Chibole, L., Chotte, J.-L., Kibunja, C. & Lesueur, D. (2011)
Effects of organic and inorganic fertilization on soil bacterial and fungal microbial diversity in
the Kabete long-term trial, Kenya. *Biology and Fertility of Soils*, **47**, 315–321.
Kibblewhite, M.G., Ritz, K. & Swift, M.J. (2008) Soil health in agricultural systems. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, **363**, 685–701.
Kumar, U., Kumar Nayak, A., Shahid, M., Gupta, V.V.S.R., Panneerselvam, P., Mohanty, S., Kaviraj,
M., Kumar, A., Chatterjee, D., Lal, B., Gautam, P., Tripathi, R. & Panda, B.B. (2018)
Continuous application of inorganic and organic fertilizers over 47 years in paddy soil alters
the bacterial community structure and its influence on rice production. *Agriculture, Ecosystems*
& Environment, **262**, 65–75.
Kumar, U., Shahid, M., Tripathi, R., Mohanty, S., Kumar, A., Bhattacharyya, P., Lal, B., Gautam, P.,
Raja, R., Panda, B.B., Jambhulkar, N.N., Shukla, A.K. & Nayak, A.K. (2017) Variation of
functional diversity of soil microbial community in sub-humid tropical rice-rice cropping
system under long-term organic and inorganic fertilization. *Ecological Indicators*, **73**, 536–
543.
Larkin, A.A. & Martiny, A.C. (2017) Microdiversity shapes the traits, niche space, and biogeography
of microbial taxa. *Environmental Microbiology Reports*, **9**, 55–70.
Lehmann, J., Solomon, D., Kinyangi, J., Dathe, L., Wirick, S. & Jacobsen, C. (2008) Spatial complexity
of soil organic matter forms at nanometre scales. *Nature Geoscience*, **1**, 238–242.
Lemanceau, P., Maron, P.-A., Mazurier, S., Mougel, C., Pivato, B., Plassart, P., Ranjard, L., Revellin,
C., Tardy, V. & Wipf, D. (2015) Understanding and managing soil biodiversity: a major
challenge in agroecology. *Agronomy for Sustainable Development*, **35**, 67–81.
Levi-Minzi, R., Riffaldi, R. & Saviozzi, A. (1986) Organic matter and nutrients in fresh and mature
farmyard manure. *Agricultural Wastes*, **16**, 225–236.
Li, J., Cooper, J.M., Lin, Z., Li, Y., Yang, X. & Zhao, B. (2015) Soil microbial community structure
and function are significantly affected by long-term organic and mineral fertilization regimes
in the North China Plain. *Applied Soil Ecology*, **96**, 75–87.
Liu, W., Wang, Q., Wang, B., Wang, X., Franks, A.E., Teng, Y., Li, Z. & Luo, Y. (2015) Changes in
the abundance and structure of bacterial communities under long-term fertilization treatments
in a peanut monocropping system. *Plant and Soil*, **395**, 415–427.
Liu, W., Zhang, Y., Jiang, S., Murray, P.J., Liao, L., Li, X. & Zhang, J. (2019) Spatiotemporal
differences in the arbuscular mycorrhizal fungi communities in soil and roots in response to
long-term organic compost inputs in an intensive agricultural cropping system on the North
China Plain. *Journal of Soils and Sediments*, **19**, 2520–2533.
Lloyd, K.G., Steen, A.D., Ladau, J., Yin, J. & Crosby, L. (2018) Phylogenetically Novel Uncultured
Microbial Cells Dominate Earth Microbiomes. *mSystems*, **3**.

Lori, M., Symnaczik, S., Mäder, P., Deyn, G.D. & Gattinger, A. (2017) Organic farming enhances soil
microbial abundance and activity—A meta-analysis and meta-regression. *PLOS ONE*, **12**,
e0180442.
Mahanta, D., Bhattacharyya, R., Mishra, P.K., Gopinath, K.A., Channakeshavaih, C., Krishnan, J.,
Raja, A., Tuti, M.D., Varghese, E., Pandey, B.M., Bisht, J.K. & Bhatt, J.C. (2017) Influence of
a six-year organic and inorganic fertilization on the diversity of the soil culturable
microrgansims in the Indian mid-Himalayas. *Applied Soil Ecology*, **120**, 229–238.
Maron, P.-A., Sarr, A., Kaisermann, A., Lévêque, J., Mathieu, O., Guigue, J., Karimi, B., Bernard, L.,
Dequiedt, S., Terrat, S., Chabbi, A. & Ranjard, L. (2018) High Microbial Diversity Promotes
Soil Ecosystem Functioning. *Appl. Environ. Microbiol.*, **84**, e02738-17.
Mazzola, M. (2004) Assessment and Management of Soil Microbial Community Structure for Disease
Suppression. *Annual Review of Phytopathology*, **42**, 35–59.
McDaniel, M.D., Tiemann, L.K. & Grandy, A.S. (2014) Does agricultural crop diversity enhance soil
microbial biomass and organic matter dynamics? A meta-analysis. *Ecological Applications*.
Meng, Q., Yang, W., Men, M., Bello, A., Xu, X., Xu, B., Deng, L., Jiang, X., Sheng, S., Wu, X., Han,
Y. & Zhu, H. (2019) Microbial Community Succession and Response to Environmental
Variables During Cow Manure and Corn Straw Composting. *Frontiers in Microbiology*, **10**.
Mensah, S., Veldtman, R., Assogbadjo, A.E., Kakaï, R.G. & Seifert, T. (2016) Tree species diversity
promotes aboveground carbon storage through functional diversity and functional dominance.
Ecology and Evolution, **6**, 7546–7557.
Morelli, F., Benedetti, Y., Perna, P. & Santolini, R. (2018) Associations among taxonomic diversity,
functional diversity and evolutionary distinctiveness vary among environments. *Ecological
Indicators*, **88**, 8–16.
Nautiyal, C.S., Chauhan, P.S. & Bhatia, C.R. (2010) Changes in soil physico-chemical properties and
microbial functional diversity due to 14 years of conversion of grassland to organic agriculture
in semi-arid agroecosystem. *Soil and Tillage Research*, **109**, 55–60.
Page, S.E. (2011) *Diversity and Complexity*, Princeton University Press, Princeton.
Philippot, L., Spor, A., Hénault, C., Bru, D., Bizouard, F., Jones, C.M., Sarr, A. & Maron, P.-A. (2013)
Loss in microbial diversity affects nitrogen cycling in soil. *The ISME Journal*, **7**, 1609–1619.
de Ponti, T., Rijk, B. & van Ittersum, M.K. (2012) The crop yield gap between organic and conventional
agriculture. *Agricultural Systems*, **108**, 1–9.
Qiu, S.-L., Wang, L.-M., Huang, D.-F. & Lin, X.-J. (2014) Effects of fertilization regimes on tea yields,
soil fertility, and soil microbial diversity. *Chilean journal of agricultural research*, **74**, 333–
339.
Reganold, J.P. & Wachter, J.M. (2016) Organic agriculture in the twenty-first century. *Nature Plants*,
2, 1–8.
Ricklefs, R.E. & Schluter, D. eds. (1994) *Species Diversity in Ecological Communities*, University of
Chicago Press, Chicago.
Rillig, M.C., Wright, S.F. & Eviner, V.T. (2002) The role of arbuscular mycorrhizal fungi and glomalin
in soil aggregation: comparing effects of five plant species. *Plant and Soil*, **238**, 325–333.
Ros, M., Klammer, S., Knapp, B., Aichberger, K. & Insam, H. (2006) Long-term effects of compost
amendment of soil on functional and structural diversity and microbial activity. *Soil Use and
Management*, **22**, 209–218.
Rudd, M.A., Beazley, K.F., Cooke, S.J., Fleishman, E., Lane, D.E., Mascia, M.B., Roth, R., Tabor, G.,
Bakker, J.A., Bellefontaine, T., Berteaux, D., Cantin, B., Chaulk, K.G., Cunningham, K.,
Dobell, R., Fast, E., Ferrara, N., Findlay, C.S., Hallstrom, L.K., Hammond, T., Hermanutz, L.,
Hutchings, J.A., Lindsay, K.E., Marta, T.J., Nguyen, V.M., Northey, G., Prior, K., Ramirez-
Sanchez, S., Rice, J., Sleep, D.J.H., Szabo, N.D., Trottier, G., Toussaint, J.-P. & Veilleux, J.-P.
(2011) Generation of Priority Research Questions to Inform Conservation Policy and
Management at a National Level. *Conservation Biology*, **25**, 476–484.
Sahu, N., Vasu, D., Sahu, A., Lal, N. & Singh, S.K. (2017) *Strength of microbes in nutrient cycling: A
key to soil health. Agriculturally Important Microbes for Sustainable Agriculture*,.
Sánchez-Bayo, F. & Wyckhuys, K.A.G. (2019) Worldwide decline of the entomofauna: A review of its
drivers. *Biological Conservation*, **232**, 8–27.

Schrama, M., de Haan, J.J., Kroonen, M., Verstegen, H. & Van der Putten, W.H. (2018) Crop yield gap and stability in organic and conventional farming systems. *Agriculture, Ecosystems and Environment*.

Sharma, M. & Reynnells, R. (2016) Importance of Soil Amendments: Survival of Bacterial Pathogens in Manure and Compost Used as Organic Fertilizers. *Microbiology Spectrum*, **4**, PFS-0010-2015.

Sun, H.Y., Deng, S.P. & Raun, W.R. (2004) Bacterial Community Structure and Diversity in a Century-Old Manure-Treated Agroecosystem. *Applied and Environmental Microbiology*, **70**, 5868–5874.

Sutherland, W.J., Armstrong-Brown, S., Armsworth, P.R., Tom, B., Brickland, J., Campbell, C.D., Chamberlain, D.E., Cooke, A.I., Dulvy, N.K., Dusic, N.R., Fitton, M., Freckleton, R.P., Godfray, H.C.J., Grout, N., Harvey, H.J., Hedley, C., Hopkins, J.J., Kift, N.B., Kirby, J., Kunin, W.E., Macdonald, D.W., Marker, B., Naura, M., Neale, A.R., Oliver, T., Osborn, D., Pullin, A.S., Shardlow, M.E.A., Showler, D.A., Smith, P.L., Smithers, R.J., Solandt, J.-L., Spencer, J., Spray, C.J., Thomas, C.D., Thompson, J., Webb, S.E., Yalden, D.W. & Watkinson, A.R. (2006) The identification of 100 ecological questions of high policy relevance in the UK. *Journal of Applied Ecology*, **43**, 617–627.

Thompson, L.R., *et al*. (2017) A communal catalogue reveals Earth's multiscale microbial diversity. *Nature*, **551**, 457.

Venter, Z.S., Jacobs, K. & Hawkins, H.J. (2016) The impact of crop rotation on soil microbial diversity: A meta-analysis. *Pedobiologia*.

Viechtbauer, W. (2010) Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical Software*, **36**, 1–48.

Vos, M., Wolf, A.B., Jennings, S.J. & Kowalchuk, G.A. (2013) Micro-scale determinants of bacterial diversity in soil. *FEMS Microbiology Reviews*, **37**, 936–954.

Wang, Q., Jiang, X., Guan, D., Wei, D., Zhao, B., Ma, M., Chen, S., Li, L., Cao, F. & Li, J. (2018) Long-term fertilization changes bacterial diversity and bacterial communities in the maize rhizosphere of Chinese Mollisols. *Applied Soil Ecology*, **125**, 88–96.

Wei, D., Yang, Q., Zhang, J.-Z., Wang, S., Chen, X.-L., Zhang, X.-L. & Li, W.-Q. (2008) Bacterial Community Structure and Diversity in a Black Soil as Affected by Long-Term Fertilization*1 *1Project supported by the Heilongjiang Provincial Natural Science Funds for Distinguished Young Scholars, China (No. JC200622), the Heilongjiang Provincial Natural Science Foundation of China (No. C2004-17), and the National "Eleventh Five Years Plan" Key Project on Science and Technology of China (No. 2006BAD25B05). *Pedosphere*, **18**, 582–592.

Wuepper, D., Borrelli, P. & Finger, R. (2020) Countries and the global rate of soil erosion. *Nature Sustainability*, **3**, 51–55.

Yao, L., Wang, D., Kang, L., Wang, D., Zhang, Y., Hou, X. & Guo, Y. (2018) Effects of fertilizations on soil bacteria and fungi communities in a degraded arid steppe revealed by high through-put sequencing. *PeerJ*, **6**, e4623.

Zhang, Q., Buyantuev, A., Li, F.Y., Jiang, L., Niu, J., Ding, Y., Kang, S. & Ma, W. (2017) Functional dominance rather than taxonomic diversity and functional diversity mainly affects community aboveground biomass in the Inner Mongolia grassland. *Ecology and Evolution*, **7**, 1605–1615.

Zhang, X., Davidson, E.A., Mauzerall, D.L., Searchinger, T.D., Dumas, P. & Shen, Y. (2015) Managing nitrogen for sustainable development. *Nature*, **528**, 51–59.

Zhang, X., Johnston, E.R., Wang, Y., Yu, Q., Tian, D., Wang, Z., Zhang, Y., Gong, D., Luo, C., Liu, W., Yang, J. & Han, X. (2019) Distinct Drivers of Core and Accessory Components of Soil Microbial Community Functional Diversity under Environmental Changes. *mSystems*, **4**.

Zhong, W., Gu, T., Wang, W., Zhang, B., Lin, X., Huang, Q. & Shen, W. (2010) The effects of mineral fertilizer and organic manure on soil microbial community and diversity. *Plant and Soil*, **326**, 511–522.

Zhou, Z., Wang, C. & Luo, Y. (2020) Meta-analysis of the impacts of global change factors on soil microbial diversity and functionality. *Nature Communications*, **11**, 3072.

564   TABLES

565   Table 1. Summary statistics for meta-analysis of functional diversity (H_{fun}) differences between fertilizer treatments and control. One study (Ros *et al*., 2006)

566   reported three comparisons, hence a multi-level random model was employed to control for within-study effects. Studies (N) refers to the number of studies

567   and number of comparisons in this estimate. LRM is the log-transformed ratio of means, giving mean ± SE and 95 % Confidence Interval. t and P are a t-test

568   of the effect size vs. zero. Change (%) is the effect-size back-transformed to percentage change.

Comparison	Studies (N)	LRM	LRM (95% CI)	t	P	Change (%)	Change 95 % CI
NPK vs. CON	6 (8)	0.026 ± 0.004	0.016, 0.035	6.51	0.0003	2.6	1.7, 3.6
ORG vs. CON	7 (9)	0.066 ± 0.011	0.041, 0.090	6.21	0.0003	6.8	4.2, 9.4
ORG vs. NPK	7 (9)	0.035 ± 0.012	0.008, 0.063	3.00	0.0171	3.6	0.8, 6.5

572 Table 2. Summary statistics for meta-analysis of taxonomic diversity differences between fertilizer treatments and control. Statistics as in Table 1.

Comparison	Class	Studies (N)	LRM	LRM (95% CI)	t	P	Change (%)	Change % (95 % CI)
NPK vs. CON	All*	1 (1)	-0.042 ± 0.100	-0.239, 0.155	-0.42	0.67	-4.1	-21.1, 16.7
	Fungi	6 (7)	0.031 ± 0.020	-0.007, 0.069	1.58	0.11	3.1	-0.73, 7.5
	Prokaryotes	18 (23)	-0.006 ± 0.016	-0.037, 0.026	-0.35	0.73	-0.5	-3.6, 2.6
ORG vs. CON	All*	1(6)	0.372 ± 0.065	0.245, 0.499	5.76	10^{-8}	46.1	35.4, 57.7
	Fungi	6 (8)	0.016 ± 0.022	-0.028, 0.059	0.70	0.48	1.6	-2.7, 6.1
	Prokaryotes	18 (23)	0.041 ± 0.014	0.013, 0.069	2.84	0.004	4.2	1.3, 7.1
ORG vs. NPK	All*	1(6)	0.419 ± 0.072	0.017, 0.072	5.79	10^{-8}	52.9	40.1, 66.8
	Fungi	6 (8)	-0.055 ± 0.021	-0.097, -0.013	2.61	0.009	-5.4	-9.2, -1.4
	Prokaryotes	23 (33)	0.045 ± 0.014	0.017, 0.073	3.20	0.001	4.6	1.8, 7.5

*A single study (Mahanta *et al*., 2017) contributed to this estimate.

FIGURES

Fig. 1. Comparison of H_{fun} (a-c) and H_{tax} (d-f) in (a,d) NPK vs. CON, (b,e) ORG vs. CON, (c,f) ORG

vs. NPK. Points show reported values with size proportional to log(1/variance). Results for Fungi are

in blue, Bacteria in red. Gel electrophoresis results are circles, sequencing results are triangles, TRFLP

are diamonds. The single soil dilution plate assay is shown as black squares.

Fig. 2. Functional soil diversity differences. a) NPK vs. CON, b) ORG vs. CON, c) ORG vs. NPK. Horizontal bars and ticks show 95% confidence intervals and means for effect sizes from the log-transformed ratio of means, back-transformed to give percentage differences. Summary estimates are given at the bottom of each plot. Random intercepts were fitted per Study.

Fig. 3. Taxonomic diversity effect sizes. a) NPK vs. CON, b) ORG vs. CON, c) ORG vs. NPK. Horizontal bars and ticks show 95% confidence intervals and means for effect sizes from the log-transformed ratio of means, back-transformed to give percentage differences. Summary estimates are given at the bottom of each plot, for All microbes (black), Fungi (blue) and Bacteria (red). Random intercepts were fitted per Study. "All" refers to results of a single study (Mahanta *et al*., 2017) which used soil dilution plate assays.

Fig. 4. Representation of processes influencing organic fertilizer affects on soil microbial diversity. See

Discussion for details.

SUPPLEMENTARY MATERIAL

SUPPLEMENTARY FIGURES

Fig. S1. Effect of fertilization on soil pH. a) Mean difference between NPK and CON vs. mean CON

pH. Blue solid and dashed lines show mean difference (-0.46 ± 0.20 SE, Z = -2.34, p = 0.019) and 95

% CI (-0.86, -0.08) of meta-analysis fit. b) Mean difference between ORG and CON vs. mean CON

pH. Blue solid and dashed lines show mean and 95% CI of meta-regression fit. Slope mean -0.174 ±

0.050, Z = -3.47, p = 0.0005, 95% CI = -0.272, -0.076. Points show means from individual studies with

size proportional to log(1/variance).

Fig. S2. Funnel plots for soil pH meta-analysis. Kendall's τ and p-values are given for rank correlation

tests for funnel plot asymmetry. a) NPK vs. CON (τ = -0.055, p = 0.88), b) ORG vs. CON (τ = -0.018,

p = 1.0).

Fig. S3. Funnel plots for functional diversity meta-analysis. Kendall's τ and p-values are given for rank

correlation tests for funnel plot asymmetry. a) NPK vs. CON (τ = -0.071, p = 0.90), b) ORG vs. CON

(τ = -0.167, p = 0.61), c) ORG vs. NPK (τ = -0.167, p = 0.61).

Fig. S4. Funnel plots for taxonomic diversity meta-analysis. Kendall's τ and p-values are given for rank

correlation tests for funnel plot asymmetry. a) NPK vs. CON (τ = -0.123, p = 0.34), b) ORG vs. CON

(τ = 0.156, p = 0.18), c) ORG vs. NPK (τ = 0.149, p = 0.14).

SUPPLEMENTARY TABLES

Table S1. Summary of studies included in the meta-analysis (alphabetical order).

Study	Location	Measure	Methodology
(Balachandar *et al.*, 2014)	Tamil Nadu, India	Taxonomic	16S seq
(Chaudhry *et al.*, 2012)	Maharashtra, India	Functional	Biolog Ecoplate
(Chen *et al.*, 2015)	Jiangxi, China	Functional	Biolog Ecoplate
(Chen *et al.*, 2016)	Jiangsu, China	Taxonomic	18S seq
(Cui *et al.*, 2018)	Hunan, China	Taxonomic	16S seq
(Ding *et al.*, 2016)	Heilongjiang, China	Taxonomic	16S seq
(Ding *et al.*, 2017)	Heilongjiang, China	Taxonomic	ITS seq
(Ge *et al.*, 2008)	Fengqiu, China	Taxonomic	16S seq
(Gu *et al.*, 2009)	Sichuan, China	Taxonomic	16S DGGE
(Guanghua *et al.*, 2008)	Jilin, China	Taxonomic	16S DGGE
		Functional	Biolog Ecoplate
(Hamm *et al.*, 2016)	Manitoba, Canada	Taxonomic	16S seq
(Han *et al.*, 2018)	Hunan, China	Taxonomic	nxr seq
(Hu *et al.*, 2018b)	Northeast China	Taxonomic	16S seq
(Ji *et al.*, 2018)	Fujian, China	Taxonomic	16S seq
(Kamaa *et al.*, 2011)	Kabete, Kenya	Taxonomic	16S and 28S DGGE
(Kumar *et al.*, 2017)	Odisha, India	Functional	Biolog Ecoplate
(Kumar *et al.*, 2018)	Odisha, India	Taxonomic	16S seq
(Li *et al.*, 2015)	Shandong, China	Taxonomic	16S DGGE
(Liu *et al.*, 2015)	Jiangxi, China	Taxonomic	16S seq
(Liu *et al.*, 2019)	Hebei, China	Taxonomic	TRFLP
(Mahanta *et al.*, 2017)	Uttarakhand, India	Taxonomic	Soil dilution plate
(Nautiyal *et al.*, 2010)	Maharashtra, India	Functional	Biolog Ecoplate
(Qiu *et al.*, 2014)	Fujian, China	Taxonomic	16S and 18S TGGE
(Ros *et al.*, 2006)	Linz, Austria	Taxonomic	16S PGGE
		Functional	Biolog Ecoplate
(Sun *et al.*, 2004)	Oklahoma, USA	Taxonomic	16S DGGE
(van Bruggen *et al.*, 2015)	Netherlands	Taxonomic	16S DGGE
(van der Bom *et al.*, 2018)	Denmark	Taxonomic	16S seq
(Wang *et al.*, 2018)	Heilongjiang, China	Taxonomic	16S seq
(Wei *et al.*, 2008)	Heilongjiang, China	Taxonomic	16S DGGE
		Functional	Biolog Ecoplate
(Yao *et al.*, 2018)	Inner Mongolia, China	Taxonomic	16S and 18S seq
(Zhong *et al.*, 2010)	Jiangxi, China	Functional	Biolog Ecoplate

628   Table S2. Meta-analysis diagnostics.

Test	I^2 (%)	Q	df	P
H_{fun} NPK vs. CON	75.8	20.6	7	0.0044
H_{fun} ORG vs. CON	77.8	23.2	8	0.0031
H_{fun} ORG vs. NPK	76.6	36.9	8	<0.0001
H_{tax} NPK vs. CON	97.0	406.6	33	<0.0001
H_{tax} ORG vs. CON	97.0	231.2	34	<0.0001
H_{tax} ORG vs. NPK	97.6	601.4	44	<0.0001
pH_{NPK} - pH_{CON} vs. pH_{CON}	98.5	1017.5	10	<0.0001
pH_{ORG} - pH_{CON} vs. pH_{CON}	69.6	23.1	9	0.0059