High-throughput phenotypic characterization of zebrafish CRISPR mutants of DNA repair genes

Unbeom Shin[1,†], Khriezhanuo Nakhro[2,†], Chang-Kyu Oh[2,†], Blake Carrington[3], Hayne Song[2], Gaurav Varshney[3], Youngjae Kim[1], Hyemin Song[2], Sangeun Jeon[2], Gabrielle Robbins[3], Sangin Kim[1], Suhyeon Yoon[2], Yongjun Choi[2], Suhyung Park[2], Yoo Jung Kim[3], Shawn Burgess[3], Sukhyun Kang[2], Raman Sood[3], Yoonsung Lee[1,2,*], Kyungjae Myung[1,2,*]

[1]School of Life Sciences, Ulsan National Institute for Science and Technology (UNIST), Ulsan 44919, Republic of Korea; [2]Center for Genomic Integrity, Institute for Basic Science (IBS), Ulsan 44919, Republic of Korea; [3]Translational and Functional Genomics Branch, National Human Genome Research Institute, National Institutes of Health, Bethesda, Maryland 20892, USA

[†] These authors contributed equally to this work

[*] To whom correspondence should be addressed

Email: yoonsunglee@ibs.re.kr

Email: kmyung@ibs.re.kr

ABSTRACT

A systematic knowledge of the roles of DNA repair genes at the level of the organism has been limited largely due to the lack of appropriate animal models and experimental techniques. Zebrafish has become a powerful vertebrate genetic model system with availability due to the ease of genome editing and large-scale phenotype screening. Here, we generated zebrafish loss-of-function mutants for 33 DNA repair and replication genes through multiplexed CRISPR/Cas9-mediated mutagenesis. High-throughput phenotypic characterization of our mutant collection revealed that four genes (*atad5a*, *ddb1*, *pcid2, pcna*) are essential for proper embryonic development and hematopoiesis; seven genes (*apex1*, *atrip*, *ino80*, *mre11a*, *shfm1*, *telo2*, *wrn*) are required for growth and development during juvenile stage and six genes (*blm*, *brca2*, *fanci , rad51*, *rad54l*, *rtel1*) play critical roles in sex development. Furthermore, mutation in seven genes (*atad5a*, *brca2*, *pcid2*, *polk*, *rad51*, *shfm1*, *xrcc1*) displayed hypersensitivity to DNA damage agents. Further characterization of *atad5a*$^{-/-}$ mutants demonstrated that Atad5a is required for normal brain development and that its loss induces ATM-p53 dependent apoptosis in the brain. Our zebrafish mutant collection provides a unique resource for understanding of the roles of DNA repair genes at the organismal level.

INTRODUCTION

DNA repair is critical for the maintenance of genomic integrity in living organisms. Our genomes continuously encounter various kinds of challenges such as oxidation, deamination, alkylation, and formation of bulky adducts, mismatches and DNA breaks. Those damages are derived from endogenous sources such as reactive oxygen species, alkylating agents and replication errors, and exogenous agents including ionizing radiation (IR), ultraviolet radiation and environmental chemicals (Chatterjee and Walker 2017). To protect our genome from those challenges, cellular systems employ several DNA repair pathways including base excision repair (BER), nucleotide excision repair (NER), mismatch repair (MMR), translesion synthesis (TLS), homologous recombination (HR) and non-homologous end-joining (NHEJ) (Friedberg 2003; Iyama and Wilson 2013). These mechanisms have been widely studied at molecular and cellular level, but a systematic investigation at the organismal level is not currently available.

Loss-of-function studies using knock-out (KO) mice have revealed numerous functional roles of DNA repair processes during vertebrate development (Specks et al. 2015). For example, normal neurogenesis requires distinct DNA damage and repair processes. Genetic defects in ATM, which activates DNA double-strand break (DSB) damage responses, cause abnormalities in the nerve system in adult mice (Barlow et al. 1996; Shiloh and Ziv 2013) and mutation of DNA-PKcs required for NHEJ leads to aberrant neurogenesis affecting neural progenitor cells and post-mitotic neurons (Enriquez-Rios et al. 2017). Moreover, certain DNA repair processes are involved in distinct mice developmental processes. Fanconi anemia (FA) genes critical for DNA interstrand cross-link (ICL) repair showed their diverse roles during embryogenesis. KO mice of *Fanca* and *Fancc* showed abnormal development of hematopoietic progenitors, while *Fanca* and *Fancg* mutant mice

demonstrated microcephaly with induced apoptosis of embryonic neural progenitors (Rathbun et al. 1997; Rio et al. 2002; Sii-Felice et al. 2008). These FA gene KO mice also showed defects of germ cell development with hypogonadism (Parmar et al. 2009). In addition, conditional KO studies revealed the function of genes related to DNA damage and repair processes on specific tissue or organ development. Mice deficient in the ubiquitin ligase adapter Ddb1 (DNA damage binding protein 1) in the brain causes abnormal brain growth and hemorrhages, while blood lineage-specific deletion of *Ddb1* leads to failure of hematopoietic stem cell development (Cang et al. 2006; Gao et al. 2015).

Although numerous KO mice of genes involved in DNA repair have been generated, uncovering how these genes affect developmental processes is hampered by the early embryonic lethality associated with mouse KO mutations in a large number of DNA repair genes (Lim and Hasty 1996; Suzuki et al. 1997; Ding et al. 2004; Cang et al. 2006; Buis et al. 2008; Qiu et al. 2016). For further understanding of gene function, tissue-specific or inducible genetic mutants in mice are critical for the study of DNA repair and replication (Cang et al. 2006; Lee et al. 2009). Teleost zebrafish has emerged as a powerful alternative genetic model system to mice, possessing similar DNA repair systems as mammals (Pei and Strauss 2013). A key advantage of zebrafish is their faster development bypassing possible early embryonic lethality of essential genes in zygotic KO mutants due to maternal mRNA deposition effects by early gastrulation stage, compared to mice system where maternal mRNA is mostly degraded before 2-cell stage (Tadros and Lipshitz 2009; Svoboda and Flemr 2010). Zebrafish therefore provide an optimal model to investigate the roles of genes critical during early embryogenesis. Moreover, recently reported multiplexed mutagenesis using CRISPR/Cas9 system allows for the generation of targeted gene mutation on a large scale (Varshney et al. 2015). Finally, mutant phenotypes can be

continuously examined throughout development in a high-throughput manner due to the visual transparency of embryos during embryogenesis and high fecundity from single zebrafish breeding (Fuentes et al. 2018).

Here, we generated 67 mutant alleles disrupting 33 genes involved in DNA repair and replication using multiplexed CRISPR/Cas9 mutagenesis and characterized their phenotypes in embryonic development, viability, growth, germ cell development, hematopoiesis and sensitivity to DNA damage. Our data revealed that deficiency of 19 of these 33 DNA repair genes lead to at least one phenotype in the homozygous mutant animals, thus revealing their critical roles in embryogenesis, hematopoiesis, growth during larval and juvenile stages, germ cell development and response to DNA damaging agents. We chose *atad5a* mutants, deficient in unloading the DNA clamp PCNA upon completion of DNA replication and repair, for more detailed phenotypic analysis and determined how *atad5a* deficiency leads to reduced brain size and hematopoietic defects in *atad5a* mutants. Our data showed that reduced brain structure and diminished number of hematopoietic stem and progenitor cells (HSPCs) are mainly caused by p53-induced apoptosis in the brain and caudal hematopoietic tissue (CHT), respectively and they can be rescued by inhibition of apoptosis by *p53* ablation or inhibition of ATM signaling. We therefore directly show that ATM-p53-dependent apoptosis is induced in the brain and blood population during embryogenesis without normal function of Atad5a and possible other crucial DNA repair genes.

RESULTS

Multiplexed CRISPR mutagenesis generated 33 loss-of-function zebrafish mutants of DNA repair genes

Aiming to mutate 57 genes related to DNA repair and replication processes, multiplexed injections of sgRNAs with *Cas9* mRNA in 16 injection groups (Supplemental Table S1) were performed as previously described (Varshney et al. 2015). Up to four gRNAs targeting distinct gene loci from different chromosomes were injected into fertilized zebrafish eggs at 1-cell stage with *Cas9* mRNA in each injection group. To identify whether potential in/del mutations generated by CRIPSR/Cas9 system are germline-transmitted, F1 progeny embryos from injected candidate founder fish were analyzed through genetic fragment analysis with fluorescence PCR for each gene in the injection group. We first identified in/del mutations in 50 genes, indicating that sgRNAs to 7 genes failed to induce mutations. Next, individual F1 animals carrying mutations were crossed with wild-type to generate F2 germline-transmitted mutant families, retrieving mutations in 35 genes, while mutant candidates with the other 15 genes failed to inherit potential in/del mutations. The siblings of F2 heterozygous mutants were then crossed to produce homozygous mutants for each allele, and genotyped again. Eventually, we recovered mutant animals containing 69 mutant alleles for 35 genes involved in DNA repair processes and replication (Supplemental Table S2). Among 35 genes, mutants of 2 genes (*polh* and *palb2*) only possessed in/del alleles with in-frame mutations, while other 33 genes were successfully mutated, generating multiple alleles with frameshift mutations that cause premature stop codons (Table 1).

Phenotypic evaluation of the mutants of 33 DNA repair genes

Previously, deletion of a number of DNA repair genes in mice KO system resulted in lethality in early embryogenesis (e.g. *Apex1*, *Atad5*, *Blm*, *Brca2*, *Ddb1*, *Ino80*, *Mre11a*, *Rad51*, *Rtel1*, *Telo2*, *Xrcc1*) (Lim and Hasty 1996; Chester et al. 1998; Ludwig et al. 1998; Tebbs et al. 1999; Ding et al. 2004; Yan et al. 2004; Cang et al. 2006; Takai et al. 2007; Buis et al. 2008;

Bell et al. 2011; Qiu et al. 2016). To confirm its critical functions during early development and further characterize the roles of DNA repair genes spatiotemporally, we screened phenotypic alterations of homozygous mutants generated from our multiplexed CRISPR mutagenesis in distinct categories, including 1) embryonic development, 2) adult survival and growth, 3) sex determination and germ cell development, 4) hematopoiesis and 5) sensitivity to genotoxic stress. For phenotype screening, we utilized mutants of 33 genes with 53 loss-of-function alleles causing frameshift mutations (Supplemental Table S3). All the mutants with defective phenotypes were only the ones with frameshift mutations except $pcna^{-/-}$ as described below.

Embryogenesis. Homozygous mutant embryos for four out of 33 genes showed morphological alterations followed by death before 10 dpf. $atad5a^{cu33/cu33}$ mutants showed aberrant head development with reduced size of the brain at 5 dpf, while $ddb1^{cu43/cu43}$ and $pcid2^{cu50/cu50}$ started to display anterior malformation including abnormalities of eye and jaw structure from 5 to 6 dpf compared to wild-type (WT) (Fig. 1A). Not surprisingly, in the absence of normal PCNA function, abnormal morphological appearance with reduced size of head and eyes as well as curved body trunk formation was observed at the earliest 50 hpf among all the mutant lines (Fig. 1B). Interestingly, the $pcna^{cu30/cu30}$ and $pcna^{cu31/cu31}$ mutants containing in-frame in/del mutations also displayed abnormal development (Fig. 1B and Supplemental Fig. S1A). Immunoprecipitation analysis of the $Pcna^{cu31/cu31}$ ($\Delta\,^{44}HVS^{46}$ mutation) cells showed that the protein was unstable to form the functional PCNA complex in those cells, while the $Pcna^{cu30/cu30}$ ($\Delta\,^{46}SL^{47}$ mutation) failed to form homo-trimer essential for clamp formation in DNA replication (Supplemental Fig. S1B and C). Due to lack of functional protein, $pcna^{cu31/cu31}$ replicates defective phenotypes of null $pcna^{cu28/cu28}$, while

pcna$^{cu30/cu30}$ showed distinctive morphology in the posterior part of the body, potentially suggesting multiple roles of PCNA during embryogenesis.

Survival and growth. In contrast to what was observed in mice, our embryonic screening demonstrated that zebrafish mutants of majority of the DNA repair and replication genes develop normally during early development. Morphological defects were only obvious more than 2 dpf, when the basic development is already completed (Fig. 1A and B). To further characterize potential phenotypic defects, we examined viability of homozygous mutant animals at 4-6 months of age when the adult zebrafish become fully mature to produce progeny. Excluding the four mutants with embryonic defects introduced above, progenies generated from inbred heterozygous parents for 29 genes were genotyped to determine whether the homozygous mutant animals survived at expected Mendelian ratio. We found that homozygous mutants in seven genes: *apex1$^{-/-}$, atrip$^{-/-}$, shfm1$^{-/-}$, ino80$^{-/-}$, mre11a$^{-/-}$, telo2$^{-/-}$, and wrn$^{-/-}$* failed to survive to adult stages (Fig. 1C). To closely understand phenotypes of these mutants, we examined a lethal time-point for each mutant by identifying the developmental stage at which the number of homozygous mutants was significantly reduced. Interestingly, each mutant showed reduced survival rate at distinctive developmental stage. Homozygous mutants of *telo2$^{cu41/cu41}$* and *shfm1$^{cu46/cu46}$* died by 2 weeks post fertilization (Fig. 2A). We further monitored possible morphological alterations and found that *telo2$^{cu41/cu41}$* embryos showed malformation of the head with a retrograded branchial arch as well as heart structure (Fig. 2B). Similarly, *shfm1$^{cu46/cu46}$* displayed defective head development with lens malformation. *apex1$^{cu2/cu2}$* and *atrip$^{cu37/cu37}$* mutants were not viable at 40 dpf, *mre11a$^{cu58/cu58}$* and *wrn$^{cu66/cu66}$* failed to survive to 60 dpf, while *ino80$^{cu36/cu36}$* scarcely survived to 90 dpf (Fig. 2C). These five mutants displayed consistent phenotypes with reduced body size compared

to their WT siblings at each designated developmental stage close to lethal time-point, suggesting potential function of each gene on either growth or physiology in vertebrates (Fig. 2D).

Sex reversal and germ cell development. Previous studies demonstrated that zebrafish mutants of HR and FA genes failed to generate female animals due to sex reversal defects without normal germ cell development (Shive et al. 2010; Botthof et al. 2017; Ramanagoudr-Bhojappa et al. 2018). To examine whether the DNA repair genes investigated here are involved in sex determination and germ cell development, we screened sexes of the 21 homozygous mutants that survive to 4- 6 month of age (Supplemental Fig. S2A). We found that all or most adult homozygous mutants were males in zebrafish lacking *blm, brca2*, *fanci, rad51*, *rad54l,* and *rtel1*, suggesting that sex reversal phenotypes developed in the absence of normal function of these genes (Fig. 3A and Supplemental Fig. S2B). Further fertility tests for these six mutants showed that male homozygotes of *brca2$^{-/-}$*, *rad51$^{-/-}$* and *blm$^{-/-}$* were infertile (fertilization success rate <6%), while the *rtel1$^{-/-}$*, *fanci$^{-/-}$* and *rad54l$^{-/-}$* male animals were found to be fertile (fertilization success rate >65%) based on outcrosses with WT female animals (Fig. 3B and Supplemental Table S4). Consistent with the finding of aberrant spermatogenesis due to meiotic arrest in *brca2$^{Q658X/Q658X}$* males (Shive et al. 2010), testes of our *brca2$^{cu53/cu53}$* males were void of spermatozoa, whereas WT siblings had normal spermatogenesis (Fig. 3C). Likewise, *blm$^{cu55/cu55}$* mutants exhibited the same skewed ratio with all homozygotes being males with possible sex reversal. Furthermore, although testis histology of *blm$^{cu55/cu55}$* homozygotes showed spermatogonia and an abundance of spermatocytes, the fraction of spermatozoa was severely diminished, indicating a failure of complete spermatogenesis similar to what was observed in *brca2$^{cu53/cu53}$* homozygotes (Fig.

3C).

Hematopoiesis. Although several studies have suggested that many DNA repair and
replication genes play a key role in hematopoiesis and human blood disorders (Xu et al.
2006; Alvarez et al. 2015; Gao et al. 2015; Botthof et al. 2017; Marsh et al. 2018), their
functions in hematopoietic progenitor development is not well understood. To understand
how HSPC development is affected in our DNA repair mutant library, we examined
expression of HSPC marker *cmyb* by whole-mount in situ hybridization (WISH).
Hypothesizing that normal function of HSPCs is critical for animal survival, we assessed
HSPC marker expression of 19 homozygous mutant embryos that show developmental and
viability defects (Fig. 4 and Supplemental Fig. S3A). WISH results showed that the *cmyb-*
positive HSPC population in the CHT where HSPCs expand to be functional (Tamplin et al.
2015) at 3-5 dpf was dramatically reduced in four mutants that showed morphological
defects before 6 dpf; *pcna$^{cu29/cu29}$*, *atad5a$^{cu33/cu33}$*, *ddb1$^{cu43/cu43}$* and *pcid2$^{cu51/cu51}$* (Fig. 4A).
cmyb expression in the dorsal aorta where HSPCs emerge at 36 hpf was normal in these
mutants, suggesting maintenance and expansion of HSPCs were affected, while HSPC
emergence and specification was normal (Supplemental Fig. S3B). We assessed
proliferation and apoptosis through EdU-incorporation assays and acridine orange (AO)
staining (Tucker and Lardelli 2007), respectively, in these animals. In the mutants of
ddb1$^{cu43/cu43}$ and *pcid2$^{cu51/cu51}$*, proliferation was significantly reduced in the CHT without
alteration of cell death events (Fig. 4B and C). In contrast, both *pcna$^{cu29/cu29}$* and
atad5a$^{cu33/cu33}$ mutants showed increased apoptosis as well as perturbed proliferation in the
CHT, suggesting that they may affect HSPC development in a distinct manner.

Sensitivity to genotoxic stress. In contrast to mice, zebrafish loss-of-function mutants of DNA repair genes were mostly viable through embryogenesis. Taking advantage of this unique property, we exposed mutant animals to genotoxic stresses to determine if hypersensitivity to specific types of DNA damage leads to distinct morphology in mutant zebrafish embryos. To induce DNA damage, embryos were treated with ionizing radiation (IR) to induce DNA DSBs or the alkylating agent methyl methanesulfonate (MMS) to stimulate base damage as well as DSBs (Beranek 1990; Mahaney et al. 2009). Mutant embryos at 6 hpf were exposed to 5 Gy IR or 0.001% MMS and their morphology was screened through 5 dpf (Supplemental Table S5). Among 31 genes, four mutants (*atad5a$^{-/-}$, pcid2$^{-/-}$, shfm1$^{-/-}$* and *rad51$^{-/-}$*) showed hypersensitivity to IR with abnormal morphology in the head and body trunk (Fig. 5A and Supplemental Fig. S4A). AO staining demonstrated that cell death was induced by IR in the head structures of the *atad5a$^{cu33/cu33}$*, *rad51$^{cu56/cu56}$* and *shfm1$^{cu46/cu46}$* mutants and in the posterior body trunk of *pcid2$^{cu50/cu50}$* (Fig. 5B and Supplemental Fig. S4B). MMS treatment also induced morphological malformation in the *atad5a$^{-/-}$, brca2$^{-/-}$, polk$^{-/-}$, rad51$^{-/-}$* and *xrcc1$^{-/-}$* mutants (Fig. 5C and Supplemental Fig. S4C). Apoptosis was robustly induced in the posterior trunk of *atad5a$^{cu33/cu33}$*, *brca2$^{cu53/cu53}$*, *rad51$^{cu56/cu56}$*, and *xrcc1$^{cu61/cu61}$* mutants, while *polk$^{cu13/cu13}$* mutant embryos showed no alteration of apoptotic events in the body trunk where morphological alteration was induced by MMS treatment (Fig. 5D and Supplemental Fig. S4D). Taken together, our data suggest that seven zebrafish mutants showed hypersensitivity to DNA damaging agents, resulting in induced morphological alterations, which were mainly caused by apoptosis.

Summary. Homozygous mutants of 19 out of 33 genes resulted in at least one defective in either embryogenesis, growth, sex development, hematopoiesis and hypersensitivity to

genotoxic stress (Table 2). In contrast to higher embryonic lethality in KO mice, only 4 out of 33 zebrafish loss-of-function mutants showed lethality before 10 dpf, successfully completing early development, including gastrulation, body patterning and initiation of organogenesis like hematopoiesis. Through further screening of HSPC formation, we found that the same four mutants failed to perform HSPC repopulation during embryogenesis. We discovered additional 13 homozygous mutant animals that displayed abnormal phenotypes at juvenile or adult stages, in an aspect of growth, viability and sex development. Finally, seven mutants showed hypersensitivity to treatment of IR and MMS. Since both damage stimuli can induce DSB, mutants of genes related to HR processes were mainly affected by IR or MMS.

Atad5a is required for normal brain development in zebrafish

We chose the *atad5a* mutants, which showed defects of embryonic morphogenesis and hematopoiesis and displayed hypersensitivity to DNA damage stress, to examine on KO animal in more detail. The ATPase family AAA domain-containing protein 5 (ATAD5) is known for regulating the function of PCNA by unloading PCNA from chromatin during DNA replication or deubiqutinating PCNA following TLS (Lee et al. 2010; Kubota et al. 2013). *Atad5* mutant mice did not survive past embryonic stage before E8.5, implying its crucial role for early development (Bell et al. 2011). Zebrafish possesses two paralogs of ATAD5; Atad5a and Atad5b (Supplemental Fig. S5A). In our study, we found that loss of *atad5a* leads to developmental aberration including abnormal structure of head and body trunk, and loss of blood stem cell precursors, while loss of *atad5b* displayed no discernable phenotype (Fig. 1C, Supplemental Fig. S5B). We further observed that *atad5a* is highly expressed during embryogenesis while the level of *atad5b* was present at much lower levels,

suggesting that Atad5a is the main functional paralog in zebrafish (Supplemental Fig. S5C).

To examine the roles of Atad5a in zebrafish embryogenesis, we analyzed the spatial distribution of *atad5a*. WISH demonstrated that *atad5a* started to express in the whole brain from 1 dpf and specifically localize in the ventricular zone of the midbrain region at 2 dpf, continuing its spatial expression in the brain up to 5 dpf (Fig. 6A, Supplemental Fig. S5D), suggesting Atad5a is potentially involved in brain development. To further examine the roles of Atad5a on brain development, we first analyzed the functional property of *atad5a*$^{cu33/cu33}$ mutants (referred as *atad5*$^{-/-}$ for brevity) and observed no functional Atad5a due to nonsense-mediated mRNA decay (NMD) (Tuladhar et al. 2019) (Supplemental Fig. S6). Moreover, western blot analysis of whole *atad5a*$^{-/-}$ embryo lysates showed that PCNA was accumulated in the chromatin, indicating that PCNA unloading was diminished, similar to what is observed in human cell lines (Kang et al. 2019b) (Fig. 6B). Finally, examination of brain morphology during embryogenesis revealed a clear reduction of brain size in the mutant at 5 dpf (Fig. 6C, Supplemental Fig.S7A). As no significant difference of whole-body length was found, this suggests a microcephaly-like phenotype of *atad5a*$^{-/-}$, indicative of a tissue-specific function (Fig. 6D, Supplemental Fig. S7B). Taken together, our results demonstrate that Atad5a is important for normal brain development during zebrafish embryogenesis.

The ventricular zone, where *atad5a* is expressed, is a highly proliferative region in the developing brain (Hibi and Shimizu 2012). Moreover, apoptosis was induced by DNA damage in the *atad5a*$^{-/-}$ embryos (Fig. 5B). We therefore hypothesized that altered proliferation and/or apoptosis in the brain of *atad5a*$^{-/-}$ would lead to malformation of the brain structures of the embryo. We assessed proliferation of brain cells by EdU staining and found EdU-positive cells mainly concentrated in the ventricular zone in WT animals at 2 dpf, while

the EdU signal was strongly reduced in the *atad5a*$^{-/-}$ mutants from 2 to 5 dpf (Fig. 6E, Supplemental Fig. S8A). In parallel, AO-positive apoptotic cells were significantly enriched in the mutant at 2 and 5 dpf, compared to WT (Fig. 6F, Supplemental Fig. S8B). These data suggest that diminished proliferation and induced cell death in the brain of *atad5a*$^{-/-}$ embryos leads to abnormal brain formation with microcephaly-like phenotype

ATM-Chk2 signaling induces p53-dependent apoptosis in the brain of the *atad5a*$^{-/-}$ mutants.

Along with DNA repair and cell cycle arrest, p53-dependent apoptosis is a well-known consequence of DNA damage induction (Chen 2016). Interestingly, *p53* transcription was robustly induced in the brain of *atad5a* mutants, compared to WT animals (Fig. 7A). To confirm whether p53-induced apoptosis is responsible for the malformation of the brain, we analyzed the phenotypes of the *atad5a*$^{-/-}$; *p53*$^{-/-}$ double mutant embryos. Depletion of p53 in the *atad5a* mutant zebrafish reduced apoptosis in the brain and recovered the size of brain without alteration of proliferation, suggesting that the p53 induction in the absence of Atad5a leads to activation of cell death and eventually failure of normal brain development (Fig. 7B, Supplemental Fig. S9A). To further understand molecular mechanism of p53-depedent apoptotic events, we examined expression of DNA damage markers. Western blot of lysates of whole embryos at 5 dpf showed that γH2AX as well as phospho-Chk2 levels were induced in the *atad5a*$^{-/-}$ mutants, suggesting DNA breaks and ATM-Chk2 signaling is activated to induce p53-depndent apoptosis in the absence of Atad5a (Taylor and Stark 2001) (Fig. 7C). To confirm if ATM activates p53 to induce cell death in the brain without functional Atad5a, we inhibited ATM signaling using the ATM inhibitor KU60019 (ATMi) (Danilova et al. 2014). Perturbation of ATM signaling significantly reduced apoptotic cell numbers in the *atad5a*$^{-/-}$

mutants (Fig. 7D). Taken together, our data showed that ATM/Chk2 signaling induces p53-dependent apoptosis in the developing zebrafish brain in the absence of Atad5a.

HSPC maturation is blocked by ATM/p53-dependent apoptosis in the *atad5a*[−/−] mutant

Our preliminary investigation of hematopoiesis showed that HSPC precursor cells failed to repopulate in the CHT of the *atad5a* mutants, although HSPC emergence in the dorsal aorta was normal (Fig. 4A, Supplemental Fig. S3B). We found that *atad5a* was expressed in the CHT at 3 dpf, when HSPCs start to repopulate and expand, suggesting its potential requirement for HSPC maintenance and maturation (Supplemental Fig. S9B). Similar to the brain development, the number of EdU-positive proliferating cells was reduced, while apoptotic events were induced in the CHT of the *atad5a*[−/−] at 3 dpf (Fig. 4B and C). To determine whether p53 activation by ATM/Chk2 signaling induces apoptosis in HSPC precursors in the absence of Atad5a, we performed *p53* WISH assays in the *atad5a*[−/−] and rescue experiment for HSPC marker expression using the *atad5a*[−/−];*p53*[−/−] double mutant and ATMi treatment in the *atad5a*[−/−] animals. *p53* expression was highly induced in the CHT of *atad5a*[−/−] mutants (Supplemental Fig. S9B). Furthermore, HSPC expansion was rescued in the CHT of *atad5a*[−/−]; *p53*[−/−] double mutants as well as ATMi-treated *atad5a*[−/−] embryos, compared to *atad5a*[−/−] single mutants or vehicle-treated *atad5a*[−/−] embryos (Fig. 7E and F). Taken together, our data suggest that *p53* activation by ATM/Chk2 triggered cell death in the HSPC population in the CHT as well as brain of *atad5a* homozygous mutants.

DISCUSSION

In this study, we generated 33 loss-of-function zebrafish mutants of genes related to DNA repair and replication process, including BER, NER, HR, TLS and MMR, via multiplexed

CRISPR mutagenesis followed by florescence PCR-based fragment analysis (Varshney et al. 2015). To fully understand functional annotations of genes *in vivo*, we screened the phenotypes of the mutants in embryogenesis, viability, growth, sex determination, hematopoiesis and sensitivity to DNA damaging agents. All the 33 homozygous mutants grew normally until 48 hpf with only four mutants showing morphological malformation and hematopoietic defects before 6 dpf, in line with observations that genomic homeostasis is linked to hematopoiesis (Farres et al. 2015; Botthof et al. 2017; Marsh et al. 2018). Mutants of *pcna* and *atad5a*, two critical genes for DNA replication (Kang et al. 2019a), showed reduced proliferation and induced apoptosis in the blood precursors at the CHT, while *ddb1* and *pcid2* homozygous mutants showed loss of HSPCs caused by reduced proliferation. PCNA and ATAD5 are critical for not only general DNA replication but also for the maintenance of replication fork stability (Park et al. 2019). Perturbation of genomic stability generally induces diverse DNA damage responses, such as cell-cycle arrest and cell death (Wang et al. 2016; Jiang et al. 2017; Gao et al. 2018). Activation of ATM pathway and p53-apoptotic process in the *atad5a* zebrafish mutants suggest that maintenance of genomic stability is critical for maintenance and repopulation of HSPCs.

Further monitoring viability and development of the mutants demonstrated that seven more KO zebrafish showed defects of survival during larval and juvenile stages; *shfm1*[-/-]*, telo2*[-/-]*, apex1*[-/-]*, atrip*[-/-]*, mre11a*[-/-]*, wrn*[-/-] *and ino80*[-/-] (Fig. 2). Unlike KO studies using murine system, all the mutant animals showed distinct defects of morphological appearance or growth at different developmental stages after embryogenesis. While KO mouse studies for the function of *Shfm1* is not yet reported, systematic analysis of KO mice in the other 6 genes have shown their critical roles in early embryogenesis resulting in embryonic lethality (Oshima et al. 1996; Xanthoudakis et al. 1996; Luo et al. 1999; Lombard et al. 2000; Takai

et al. 2007; Dickinson et al. 2016). Conditional KO (CKO) mouse approaches have shown the roles of Apex1 on neurological function with injury and functions of Ino80 on genomic stability in mammals (Min et al. 2013; Stetler et al. 2016). In addition, our six zebrafish mutants showed female-to-male sex reversal defects; $blm^{-/-}$, $brca2^{-/-}$, $fanci^{-/-}$, $rad51^{-/-}$, $rad54l^{-/-}$ and $rtel1^{-/-}$. While $rad54l^{-/-}$ and $rtel1^{-/-}$ mutants demonstrated a possible sex reversal phenotype with normal fertility like $fanci^{-/-}$ homozygotes (Ramanagoudr-Bhojappa et al. 2018), $blm^{-/-}$ male animals were infertile without normal functional germ cells in the testis like $brca2^{-/-}$ and $rad51^{-/-}$ where primordial germ cells failed to survive and proliferate in the testes (Shive et al. 2010; Botthof et al. 2017). Although Blm homozygous KO mouse embryos were developmentally delayed and died by embryonic day 13.5 (Chester et al. 1998), $blm^{cu55/cu55}$ zebrafish survived and we were able to observe aberrant spermatogenesis, potentially recapitulating a phenotype of human male infertility in patients with Bloom syndrome (de Renty and Ellis 2017). Taken together, our zebrafish KO mutants showed diverse developmental problems at various stages, implying distinct roles of each gene *in vivo*. Further phenotypic annotation of each zebrafish mutant generated in our study will provide valuable knowledge of biological roles of DNA repair genes excepting early embryogenesis without laborious CKO approaches.

Our CRISPR mutants showed phenotypes consistent with previously reported zebrafish loss-of-function studies, including the sex reversal defects of $brca2^{-/-}$, $rad51^{-/-}$ and $fanci^{-/-}$ mutants (Shive et al. 2010; Botthof et al. 2017; Ramanagoudr-Bhojappa et al. 2018) and embryonic malformation with aberrant head development in $ddb1^{-/-}$ zebrafish embryos (Hu et al. 2015). Interestingly, several zebrafish knock-down studies using morpholinos (MOs) yielded different results from our mutant studies. For example, knock-down of BER components, *apex1* or *unga* using MOs led to aberrant heart and brain development during

embryogenesis (Wang et al. 2006; Wu et al. 2014; Pei et al. 2019) and *hmgb1a* morphants showed abnormal morphology of the brain during early development (Zhao et al. 2011), whereas mutants of these three genes develop normally in embryogenesis. This discrepancy between studies of MOs and gene-edited mutants could be due to several reasons. First, early knock-down using translation blocking MOs might block the gene function earlier than KOs by blocking expression of target genes in maternal mRNA, removing maternal protection and resulting earlier phenotypes. To clarify this hypothesis using mutant animals, phenotyping maternal zygotic mutant embryos will be practical. The second possible cause is a potential genetic compensation in the zebrafish CRISPR mutants, which is rarely observed in the stable KO models by inducing genes related to the mutated one (Rossi et al. 2015). The presence of genetic compensation phenomenon can be identified by transcriptome analysis between the mutants and their sibling controls.

We further examined sensitivity of the mutants to two genotoxic agents. IR generally induces DNA DSBs, while MMS is known as an alkylation agent that triggers DNA repair processes including DSB and excision repair as it forms alkylated bases and DSBs (Beranek 1990; Lundin et al. 2005; Mahaney et al. 2009). As expected, mutants of HR-related genes like *rad51*, *pcid2* and *shfm1* showed hypersensitivity to DSBs induced by IR, while *xrcc1*[-/-] mutants with an BER defect were sensitive to MMS treatment. Additionally, MMS-induced hypersensitivity in the mutants of *polk* essential for TLS process is consistent with the study of *C. elegans polk* null mutants where mutation rate in the genome is robustly enhanced by MMS treatment (Volkova et al. 2020). Likewise, MMS-induced malformation of our zebrafish *polk*[-/-] mutants without induced apoptosis is possibly caused by highly induced mutagenesis in the somatic genome of the mutant. This is consistent with the role of Polk in bypassing minor groove DNA adducts such as 3-methyladenine (Plosky et al. 2008). In addition to

known function of each DNA repair gene, we uncovered potentially unknown functions of genes like *atad5a*. Beside 19 mutants with defects of development or genotoxic sensitivity (Table 2), the other 14 zebrafish mutants showed no altered phenotypes in our assays. To closely dissect these mutant animals without visible phenotypes in this study, it is important to prioritize more specific phenotyping strategies based on their general cellular functions. Therefore, further investigation of sensitivity to genotoxic stress using various DNA damage agents may lead to observation of new morphological alterations. In addition, further screening of non-lethal tissue-specific phenotypes along with analysis of genome expression will provide more information of functional roles of these DNA repair genes in a vertebrate.

To successfully complete highly proliferative neurogenesis, maintenance of genomic stability and DNA damage signaling are both critical (McKinnon 2009; McKinnon 2013). Loss of genomic stability by malfunction of DNA repair protein often leads to neurologic disease (Bender et al. 2006; Reynolds et al. 2009). Consistent with previous cellular studies, we found that *atad5a* mutant zebrafish showed sensitivity in response to both IR and MMS (Lee et al. 2013; Park et al. 2019). In addition, chromatin-bound PCNA is accumulated in the mutant background, showing that the cellular function of ATAD5, the removal of PCNA upon completion of replication, is conserved in zebrafish. *atad5a*[-/-] embryos showed reduced size of the brain where proliferation was diminished while cell death was induced by ATM/p53 activation. Based on our characterization of *atad5a*[-/-] phenotype, we speculate loss of Atad5a leads to the accumulation of PCNA on chromatin, causing replication stress and induction of cell death through ATM/Chk2 and p53. ATM is one of main DNA damage-activated kinases in the nervous system, responding to mainly DSB (Shiloh and Ziv 2013). Our data describing the function of *atad5a* using zebrafish

mutants demonstrated that Atad5 plays a pivotal role to maintain genomic integrity and regulate DNA damage signaling for normal neurogenesis and brain development.

Through high-throughput characterization of genetically functional materials, zebrafish KO lines can be utilized for functional studies of specific proteins with emphasis on the organ level or systemic phenotype on a large scale. Survival/lethality and phenotypic studies of DNA damage response genes will not only help unravel the importance of those genes at specific tissues, but also show the functional spectrum of influence on the disease phenotypes. These model systems can therefore also be vital platform technology for the study of genetic therapy and drug testing for human diseases.

MATERIALS AND METHODS

Zebrafish husbandry

All zebrafish embryos and adult animals were raised and maintained in a circulating aquarium system (Genomic Design or Aquaneering) at 28.5 °C in accordance with UNIST or NIH Institutional Animal Care and Use Committees (IACUC: UNISTIACUC-20-09; NIH G-05-5). For CRISPR mutant generation, wild-type TAB5 strain was used. For loss of function studies for p53, $tp53^{zdf1/zdf1}$ line was used (Berghmans et al. 2005).

Generation of zebrafish mutants through CRISPR/Cas9 mutagenesis

Single guide RNAs (sgRNAs) targeting 57 genes related to DNA repair processes were designed and generation of sgRNA template and preparation of sgRNA and *Cas9* mRNA was performed as previously described (Varshney et al. 2015; Varshney et al. 2016) (Supplemental Table S1). Up to four gRNAs (50pg per each gRNA) targeting different genes mixed with *Cas9* mRNA (300pg) were injected into wild-type TAB5 embryos at 1-cell stage

grown to adulthood. The multiplexing scheme and fluorescence PCR primer sequences for each gene are described in Supplemental Table S1. To confirm germline-transmitted in/del mutations, F1 progeny generated from injected founder fish were screened through fluorescence PCR as described previously (Varshney et al. 2016). Sanger sequencing was performed to determine the sequence of mutant alleles.

ACKNOWLEDGEMENT

This research was supported by the Institute for Basic Science (IBS-R022-D1) and Ulsan National Institute of Science and Technology grant (1.150054) to K Myung and Intramural Research Program of the National Human Genome Research Institute, National Institutes of Health to R Sood. We thank Hana Kim and Yun Ji Choi for technical assistance, Orlando Sharer for scientific guidance and members of Molecular Genetics Section in the Center for Genomic Integrity for providing helpful comments.

Author Contribution: Conceptualization and Methodology, Y.L. and K.M.; Validation, U.S., K.N., C.O. and Y.L.; Formal Analysis and Visualization, U.S., K.N., C.O., and Y.L; Investigation, U.S., K.N., C.O., B.C., H.S., G.V., Y.K., H.S., S.J., G.R., S.K., S.Y., Y.C., and S.P.; Writing-Original Draft, U.S., K.N., C.O. and Y.L.; Supervision, Y.L. and K.M.; Project Administration, Y.L. and K.M.

REFERENCES

Alvarez S, Diaz M, Flach J, Rodriguez-Acebes S, Lopez-Contreras AJ, Martinez D, Canamero M, Fernandez-Capetillo O, Isern J, Passegue E et al. 2015. Replication stress caused by low MCM expression limits fetal erythropoiesis and hematopoietic stem cell functionality. *Nat Commun* **6**: 8548.

Barlow C, Hirotsune S, Paylor R, Liyanage M, Eckhaus M, Collins F, Shiloh Y, Crawley JN, Ried T, Tagle D et al. 1996. Atm-deficient mice: a paradigm of ataxia telangiectasia. *Cell* **86**: 159-171.

Bell DW, Sikdar N, Lee KY, Price JC, Chatterjee R, Park HD, Fox J, Ishiai M, Rudd ML,

Pollock LM et al. 2011. Predisposition to cancer caused by genetic and functional defects of mammalian Atad5. *PLoS Genet* **7**: e1002245.

Bender A, Krishnan KJ, Morris CM, Taylor GA, Reeve AK, Perry RH, Jaros E, Hersheson JS, Betts J, Klopstock T et al. 2006. High levels of mitochondrial DNA deletions in substantia nigra neurons in aging and Parkinson disease. *Nat Genet* **38**: 515-517.

Beranek DT. 1990. Distribution of methyl and ethyl adducts following alkylation with monofunctional alkylating agents. *Mutat Res* **231**: 11-30.

Berghmans S, Murphey RD, Wienholds E, Neuberg D, Kutok JL, Fletcher CD, Morris JP, Liu TX, Schulte-Merker S, Kanki JP et al. 2005. tp53 mutant zebrafish develop malignant peripheral nerve sheath tumors. *Proc Natl Acad Sci U S A* **102**: 407-412.

Botthof JG, Bielczyk-Maczynska E, Ferreira L, Cvejic A. 2017. Loss of the homologous recombination gene rad51 leads to Fanconi anemia-like symptoms in zebrafish. *Proc Natl Acad Sci U S A* **114**: E4452-E4461.

Buis J, Wu Y, Deng Y, Leddon J, Westfield G, Eckersdorff M, Sekiguchi JM, Chang S, Ferguson DO. 2008. Mre11 nuclease activity has essential roles in DNA repair and genomic stability distinct from ATM activation. *Cell* **135**: 85-96.

Cang Y, Zhang J, Nicholas SA, Bastien J, Li B, Zhou P, Goff SP. 2006. Deletion of DDB1 in mouse brain and lens leads to p53-dependent elimination of proliferating cells. *Cell* **127**: 929-940.

Chatterjee N, Walker GC. 2017. Mechanisms of DNA damage, repair, and mutagenesis. *Environ Mol Mutagen* **58**: 235-263.

Chen J. 2016. The Cell-Cycle Arrest and Apoptotic Functions of p53 in Tumor Initiation and Progression. *Cold Spring Harb Perspect Med* **6**: a026104.

Chester N, Kuo F, Kozak C, O'Hara CD, Leder P. 1998. Stage-specific apoptosis, developmental delay, and embryonic lethality in mice homozygous for a targeted disruption in the murine Bloom's syndrome gene. *Genes Dev* **12**: 3382-3393.

Danilova N, Bibikova E, Covey TM, Nathanson D, Dimitrova E, Konto Y, Lindgren A, Glader B, Radu CG, Sakamoto KM et al. 2014. The role of the DNA damage response in zebrafish and cellular models of Diamond Blackfan anemia. *Dis Model Mech* **7**: 895-905.

de Renty C, Ellis NA. 2017. Bloom's syndrome: Why not premature aging?: A comparison of the BLM and WRN helicases. *Ageing Res Rev* **33**: 36-51.

Dickinson ME, Flenniken AM, Ji X, Teboul L, Wong MD, White JK, Meehan TF, Weninger WJ, Westerberg H, Adissu H et al. 2016. High-throughput discovery of novel developmental phenotypes. *Nature* **537**: 508-514.

Ding H, Schertzer M, Wu X, Gertsenstein M, Selig S, Kammori M, Pourvali R, Poon S, Vulto I, Chavez E et al. 2004. Regulation of murine telomere length by Rtel: an essential gene encoding a helicase-like protein. *Cell* **117**: 873-886.

Enriquez-Rios V, Dumitrache LC, Downing SM, Li Y, Brown EJ, Russell HR, McKinnon PJ. 2017. DNA-PKcs, ATM, and ATR Interplay Maintains Genome Integrity during Neurogenesis. *J Neurosci* **37**: 893-905.

Farres J, Llacuna L, Martin-Caballero J, Martinez C, Lozano JJ, Ampurdanes C, Lopez-Contreras AJ, Florensa L, Navarro J, Ottina E et al. 2015. PARP-2 sustains erythropoiesis in mice by limiting replicative stress in erythroid progenitors. *Cell Death Differ* **22**: 1144-1157.

Friedberg EC. 2003. DNA damage and repair. *Nature* **421**: 436-440.

Fuentes R, Letelier J, Tajer B, Valdivia LE, Mullins MC. 2018. Fishing forward and reverse: Advances in zebrafish phenomics. *Mech Dev* **154**: 296-308.

Gao J, Buckley SM, Cimmino L, Guillamot M, Strikoudis A, Cang Y, Goff SP, Aifantis I. 2015. The CUL4-DDB1 ubiquitin ligase complex controls adult and embryonic stem cell differentiation and homeostasis. *Elife* **4**.

Gao X, Han L, Ding N, Mu Y, Guan P, Hu C, Huang X. 2018. Bafilomycin C1 induces G0/G1 cell-cycle arrest and mitochondrial-mediated apoptosis in human hepatocellular cancer SMMC7721 cells. *J Antibiot (Tokyo)* **71**: 808-817.

Hibi M, Shimizu T. 2012. Development of the cerebellum and cerebellar neural circuits. *Dev Neurobiol* **72**: 282-301.

Hu Z, Holzschuh J, Driever W. 2015. Loss of DDB1 Leads to Transcriptional p53 Pathway Activation in Proliferating Cells, Cell Cycle Deregulation, and Apoptosis in Zebrafish Embryos. *PLoS One* **10**: e0134299.

Iyama T, Wilson DM, 3rd. 2013. DNA repair mechanisms in dividing and non-dividing cells. *DNA Repair (Amst)* **12**: 620-636.

Jiang Y, Wang X, Hu D. 2017. Furanodienone induces G0/G1 arrest and causes apoptosis via the ROS/MAPKs-mediated caspase-dependent pathway in human colorectal cancer cells: a study in vitro and in vivo. *Cell Death Dis* **8**: e2815.

Kang MS, Kim J, Ryu E, Ha NY, Hwang S, Kim BG, Ra JS, Kim YJ, Hwang JM, Myung K et al. 2019a. PCNA Unloading Is Negatively Regulated by BET Proteins. *Cell Rep* **29**: 4632-4645 e4635.

Kang MS, Ryu E, Lee SW, Park J, Ha NY, Ra JS, Kim YJ, Kim J, Abdel-Rahman M, Park SH et al. 2019b. Regulation of PCNA cycling on replicating DNA by RFC and RFC-like complexes. *Nat Commun* **10**: 2420.

Kubota T, Nishimura K, Kanemaki MT, Donaldson AD. 2013. The Elg1 replication factor C-like complex functions in PCNA unloading during DNA replication. *Mol Cell* **50**: 273-280.

Lee KY, Fu H, Aladjem MI, Myung K. 2013. ATAD5 regulates the lifespan of DNA replication factories by modulating PCNA level on the chromatin. *J Cell Biol* **200**: 31-44.

Lee KY, Yang K, Cohn MA, Sikdar N, D'Andrea AD, Myung K. 2010. Human ELG1 regulates the level of ubiquitinated proliferating cell nuclear antigen (PCNA) through Its interactions with PCNA and USP1. *J Biol Chem* **285**: 10362-10369.

Lee Y, Katyal S, Li Y, El-Khamisy SF, Russell HR, Caldecott KW, McKinnon PJ. 2009. The genesis of cerebellar interneurons and the prevention of neural DNA damage require XRCC1. *Nat Neurosci* **12**: 973-980.

Lim DS, Hasty P. 1996. A mutation in mouse rad51 results in an early embryonic lethal that is suppressed by a mutation in p53. *Mol Cell Biol* **16**: 7133-7143.

Lombard DB, Beard C, Johnson B, Marciniak RA, Dausman J, Bronson R, Buhlmann JE, Lipman R, Curry R, Sharpe A et al. 2000. Mutations in the WRN gene in mice accelerate mortality in a p53-null background. *Mol Cell Biol* **20**: 3286-3291.

Ludwig DL, MacInnes MA, Takiguchi Y, Purtymun PE, Henrie M, Flannery M, Meneses J, Pedersen RA, Chen DJ. 1998. A murine AP-endonuclease gene-targeted deficiency with post-implantation embryonic progression and ionizing radiation sensitivity. *Mutat Res-DNA Repair* **409**: 17-29.

Lundin C, North M, Erixon K, Walters K, Jenssen D, Goldman AS, Helleday T. 2005. Methyl methanesulfonate (MMS) produces heat-labile DNA damage but no detectable in vivo DNA double-strand breaks. *Nucleic Acids Res* **33**: 3799-3811.

Luo G, Yao MS, Bender CF, Mills M, Bladl AR, Bradley A, Petrini JH. 1999. Disruption of mRad50 causes embryonic stem cell lethality, abnormal embryonic development, and sensitivity to ionizing radiation. *Proc Natl Acad Sci U S A* **96**: 7376-7381.

Mahaney BL, Meek K, Lees-Miller SP. 2009. Repair of ionizing radiation-induced DNA double-strand breaks by non-homologous end-joining. *Biochem J* **417**: 639-650.

Marsh JCW, Gutierrez-Rodrigues F, Cooper J, Jiang J, Gandhi S, Kajigaya S, Feng X, Ibanez M, Donaires FS, Lopes da Silva JP et al. 2018. Heterozygous RTEL1 variants in bone marrow failure and myeloid neoplasms. *Blood Adv* **2**: 36-48.

McKinnon PJ. 2009. DNA repair deficiency and neurological disease. *Nat Rev Neurosci* **10**: 100-112.

-. 2013. Maintaining genome stability in the nervous system. *Nat Neurosci* **16**: 1523-1529.

Min JN, Tian Y, Xiao Y, Wu L, Li L, Chang S. 2013. The mINO80 chromatin remodeling complex is required for efficient telomere replication and maintenance of genome stability. *Cell Res* **23**: 1396-1413.

Oshima J, Yu CE, Piussan C, Klein G, Jabkowski J, Balci S, Miki T, Nakura J, Ogihara T, Ells J et al. 1996. Homozygous and compound heterozygous mutations at the Werner syndrome locus. *Hum Mol Genet* **5**: 1909-1913.

Park SH, Kang N, Song E, Wie M, Lee EA, Hwang S, Lee D, Ra JS, Park IB, Park J et al. 2019. ATAD5 promotes replication restart by regulating RAD51 and PCNA in response to replication stress. *Nat Commun* **10**: 5718.

Parmar K, D'Andrea A, Niedernhofer LJ. 2009. Mouse models of Fanconi anemia. *Mutat Res* **668**: 133-140.

Pei DS, Jia PP, Luo JJ, Liu W, Strauss PR. 2019. AP endonuclease 1 (Apex1) influences brain development linking oxidative stress and DNA repair. *Cell Death Dis* **10**: 348.

Pei DS, Strauss PR. 2013. Zebrafish as a model system to study DNA damage and repair. *Mutat Res* **743-744**: 151-159.

Plosky BS, Frank EG, Berry DA, Vennall GP, McDonald JP, Woodgate R. 2008. Eukaryotic Y-family polymerases bypass a 3-methyl-2 '-deoxyadenosine analog in vitro and methyl methanesulfonate-induced DNA damage in vivo. *Nucleic Acids Research* **36**: 2152-2162.

Qiu Z, Elsayed Z, Peterkin V, Alkatib S, Bennett D, Landry JW. 2016. Ino80 is essential for proximal-distal axis asymmetry in part by regulating Bmp4 expression. *BMC Biol* **14**: 18.

Ramanagoudr-Bhojappa R, Carrington B, Ramaswami M, Bishop K, Robbins GM, Jones M, Harper U, Frederickson SC, Kimble DC, Sood R et al. 2018. Multiplexed CRISPR/Cas9-mediated knockout of 19 Fanconi anemia pathway genes in zebrafish revealed their roles in growth, sexual development and fertility. *PLoS Genet* **14**: e1007821.

Rathbun RK, Faulkner GR, Ostroski MH, Christianson TA, Hughes G, Jones G, Cahn R, Maziarz R, Royle G, Keeble W et al. 1997. Inactivation of the Fanconi anemia group C gene augments interferon-gamma-induced apoptotic responses in hematopoietic cells. *Blood* **90**: 974-985.

Reynolds JJ, El-Khamisy SF, Katyal S, Clements P, McKinnon PJ, Caldecott KW. 2009. Defective DNA ligation during short-patch single-strand break repair in ataxia oculomotor apraxia 1. *Mol Cell Biol* **29**: 1354-1362.

Rio P, Segovia JC, Hanenberg H, Casado JA, Martinez J, Gottsche K, Cheng NC, Van de Vrugt HJ, Arwert F, Joenje H et al. 2002. In vitro phenotypic correction of hematopoietic progenitors from Fanconi anemia group A knockout mice. *Blood* **100**: 2032-2039.

Rossi A, Kontarakis Z, Gerri C, Nolte H, Holper S, Kruger M, Stainier DY. 2015. Genetic compensation induced by deleterious mutations but not gene knockdowns. *Nature*

524: 230-233.

Shiloh Y, Ziv Y. 2013. The ATM protein kinase: regulating the cellular response to genotoxic stress, and more. *Nat Rev Mol Cell Biol* **14**: 197-210.

Shive HR, West RR, Embree LJ, Azuma M, Sood R, Liu P, Hickstein DD. 2010. brca2 in zebrafish ovarian development, spermatogenesis, and tumorigenesis. *Proc Natl Acad Sci U S A* **107**: 19350-19355.

Sii-Felice K, Etienne O, Hoffschir F, Mathieu C, Riou L, Barroca V, Haton C, Arwert F, Fouchet P, Boussin FD et al. 2008. Fanconi DNA repair pathway is required for survival and long-term maintenance of neural progenitors. *EMBO J* **27**: 770-781.

Specks J, Nieto-Soler M, Lopez-Contreras AJ, Fernandez-Capetillo O. 2015. Modeling the study of DNA damage responses in mice. *Methods Mol Biol* **1267**: 413-437.

Stetler RA, Gao Y, Leak RK, Weng Z, Shi Y, Zhang L, Pu H, Zhang F, Hu X, Hassan S et al. 2016. APE1/Ref-1 facilitates recovery of gray and white matter and neurological function after mild stroke injury. *Proc Natl Acad Sci U S A* **113**: E3558-3567.

Suzuki A, de la Pompa JL, Hakem R, Elia A, Yoshida R, Mo R, Nishina H, Chuang T, Wakeham A, Itie A et al. 1997. Brca2 is required for embryonic cellular proliferation in the mouse. *Genes Dev* **11**: 1242-1252.

Svoboda P, Flemr M. 2010. The role of miRNAs and endogenous siRNAs in maternal-to-zygotic reprogramming and the establishment of pluripotency. *EMBO Rep* **11**: 590-597.

Tadros W, Lipshitz HD. 2009. The maternal-to-zygotic transition: a play in two acts. *Development* **136**: 3033-3042.

Takai H, Wang RC, Takai KK, Yang H, de Lange T. 2007. Tel2 regulates the stability of PI3K-related protein kinases. *Cell* **131**: 1248-1259.

Tamplin OJ, Durand EM, Carr LA, Childs SJ, Hagedorn EJ, Li P, Yzaguirre AD, Speck NA, Zon LI. 2015. Hematopoietic stem cell arrival triggers dynamic remodeling of the perivascular niche. *Cell* **160**: 241-252.

Taylor WR, Stark GR. 2001. Regulation of the G2/M transition by p53. *Oncogene* **20**: 1803-1815.

Tebbs RS, Flannery ML, Meneses JJ, Hartmann A, Tucker JD, Thompson LH, Cleaver JE, Pedersen RA. 1999. Requirement for the Xrcc1 DNA base excision repair gene during early mouse development. *Dev Biol* **208**: 513-529.

Tucker B, Lardelli M. 2007. A rapid apoptosis assay measuring relative acridine orange fluorescence in zebrafish embryos. *Zebrafish* **4**: 113-116.

Tuladhar R, Yeu Y, Tyler Piazza J, Tan Z, Rene Clemenceau J, Wu X, Barrett Q, Herbert J, Mathews DH, Kim J et al. 2019. CRISPR-Cas9-based mutagenesis frequently provokes on-target mRNA misregulation. *Nat Commun* **10**: 4056.

Varshney GK, Carrington B, Pei W, Bishop K, Chen Z, Fan C, Xu L, Jones M, LaFave MC, Ledin J et al. 2016. A high-throughput functional genomics workflow based on CRISPR/Cas9-mediated targeted mutagenesis in zebrafish. *Nat Protoc* **11**: 2357-2375.

Varshney GK, Pei W, LaFave MC, Idol J, Xu L, Gallardo V, Carrington B, Bishop K, Jones M, Li M et al. 2015. High-throughput gene targeting and phenotyping in zebrafish using CRISPR/Cas9. *Genome Res* **25**: 1030-1042.

Volkova NV, Meier B, Gonzalez-Huici V, Bertolini S, Gonzalez S, Vohringer H, Abascal F, Martincorena I, Campbell PJ, Gartner A et al. 2020. Mutational signatures are jointly shaped by DNA damage and repair. *Nat Commun* **11**: 2169.

Wang H, Zhang T, Sun W, Wang Z, Zuo D, Zhou Z, Li S, Xu J, Yin F, Hua Y et al. 2016.

Erianin induces G2/M-phase arrest, apoptosis, and autophagy via the ROS/JNK signaling pathway in human osteosarcoma cells in vitro and in vivo. *Cell Death Dis* **7**: e2247.

Wang Y, Shupenko CC, Melo LF, Strauss PR. 2006. DNA repair protein involved in heart and blood development. *Mol Cell Biol* **26**: 9083-9093.

Wu D, Chen L, Sun Q, Wu X, Jia S, Meng A. 2014. Uracil-DNA glycosylase is involved in DNA demethylation and required for embryonic development in the zebrafish embryo. *J Biol Chem* **289**: 15463-15473.

Xanthoudakis S, Smeyne RJ, Wallace JD, Curran T. 1996. The redox/DNA repair protein, Ref-1, is essential for early embryonic development in mice. *Proc Natl Acad Sci U S A* **93**: 8919-8923.

Xu Y, Leung CG, Lee DC, Kennedy BK, Crispino JD. 2006. MTB, the murine homolog of condensin II subunit CAP-G2, represses transcription and promotes erythroid cell differentiation. *Leukemia* **20**: 1261-1269.

Yan DH, Wen Y, Su LK, Xia WY, Wang SC, Zhang S, Gan L, Lee DF, Spohn B, Frey JA et al. 2004. A delayed chemically induced tumorigenesis in Brca2 mutant mice. *Oncogene* **23**: 1896-1901.

Zhao X, Kuja-Panula J, Rouhiainen A, Chen YC, Panula P, Rauvala H. 2011. High mobility group box-1 (HMGB1; amphoterin) is required for zebrafish brain development. *J Biol Chem* **286**: 23200-23213.

DNA repair process	Genes with frameshift mutant alleles
Base excision repair	*apex1, mutyh, neil1, unga, xrcc1*
Nucleotide excision repair	*ddb1, ercc1, xpc*
Homologous recombination	*brca2, blm, mre11a, pcid2, rad51, rad54l, shfm1, slx1b, wrn*
Translesion synthesis	*polk, rad18*
Template switching	*hltf, shprh*
Mismatch repair	*msh2*
Cross-link repair	*fanci, usp1*
DNA replication	*atad5a, atad5b, pcna*
Checkpoint	*atrip, hmgb1a, ino80, telo2*
Telomere related	*rtel1*
dNTP pool sanitation	*nudt1*

Table 1. List of genes related to DNA repair and replication with frameshift mutations obtained from CRISPR mutagenesis.

DNA repair process	Genes	Phenotyping				
		Embryogenesis	Hematopoiesis	Adult survival	Sex reversal	Sensitivity to genotoxic stress
BER	*apex1*			O		
	mutyh					
	neil1					
	unga					
	xrcc1					O
NER	*ddb1*	O	O			
	ercc1					
	xpc					
HR	*brca2*				O	O
	blm				O	
	mre11a			O		
	pcid2	O	O			O
	rad51				O	O
	rad54l				O	
	shfm1			O		O
	slx1b					
	wrn			O		
TLS	*polk*					O
	rad18					
Template switch	*hltf*					
	shprh					
MMR	*msh2*					
Cross-link Repair	*fanci*				O	
	usp1					
Replication	*atad5a*	O	O			O
	atad5b					
	pcna	O	O			
checkpoint	*atrip*			O		
	hmgb1a					
	ino80			O		
	telo2			O		
telomere	*rtel1*				O	
dNTP pool sanitation	*nudt1*					

Table 2. Summary of mutant phenotyping using 33 CRISPR mutants related to DNA repair processes. Letter "O" indicates defective phenotypes observed in designated mutants with each category of phenotyping.

FIGURE LEGENDS

Figure 1. Large-scale morphological characterization of zebrafish CRISPR mutants related to DNA repair genes. A. Lateral view of mutant embryos of $atad5a^{-/-}$($cu33/cu33$) at 5 dpf, $pcid2^{-/-}$($cu50/cu50$) at 5 dpf, and $ddb1^{-/-}$($cu43/cu43$) at 6dpf with their sibling WT controls. Red arrowheads indicate morphological alterations in the mutant embryos. **B.** Images of *pcna* mutants and WT control at 50 hpf. $pcna^{cu28/cu28}$ possessed 1 base pair (bp) deletion producing null mutants, while $pcna^{cu30/cu30}$ and $pcna^{cu31/cu31}$ contain in-frame deletion generating deleted protein of PCNA monomer of $\Delta^{46}SL^{47}$ and $\Delta^{44}HVS^{46}$, respectively. Red arrowheads indicate morphological alterations in the mutant embryos. **C.** A graph describes percentage of fish progeny with genotype (+/+, +/-, -/-) from each inbred heterozygous mutant line at 4 to 6 months of age. Numbers in each graph bar represent the number of fish with each genotype. Black asterisks indicate the mutant alleles in which homozygous mutants are lethal at adult stage. Scale bar = 500 μm.

Figure 2. Screening of viability and morphology of mutants during larval and juvenile stages. A. The number of genotyped zebrafish progeny from inbred heterozygous mutants of *telo2* and *shfm1* at designated developmental stages with three representative genotypes (+/+, +/-, -/-). Numbers in the bar of each graph represent the number of fish with each genotype. **B.** Images of anterior region in the $telo2^{-/-}$($cu41/cu41$) and $shfm1^{-/-}$($cu46/cu46$) mutants and their WT sibling controls. Lateral view of $telo2^{-/-}$($cu41/cu41$) at 9 dpf showed abnormal formation of jawbone and heart structure, while dorsal view of $shfm1^{-/-}$($cu46/cu46$) at 10 dpf demonstrated malformation of head part including mandible and lens. Red arrowheads indicate aberrant morphological structures in the mutants. **C.** Graphs describe genotyped progeny from inbred heterozygous mutants of *apex1*, *atrip*, *mre11a*, *wrn* and

ino80 at each designated developmental stage. Numbers in the bar of each graph represent the number of fish with each genotype. **D.** Representative phenotypic images of each mutant (*apex1*[-/-](*cu2/cu2*), *atrip*[-/-](*cu37/cu37*), *mre11a*[-/-](*cu58/cu58*), *wrn*[-/-](*cu66/cu66*) and *ino80*[-/-] (*cu36/cu36*)) and its WT sibling controls at the designated time and quantification of the body length (fork length) of each mutant and WT control. Scale bars in each image indicate 10 mm. Graphs represent mean ± S.E.M. with individual values. p-values were calculated by unpaired two-tailed Student's t-test. *** $p<0.001$, **$p<0.01$. Black scale bars = 200 μm.

Figure. 3. Mutants with sex reversal and abnormal germ cell development. A. The number of male and female animals genotyped from inbred heterozygous mutants of *blm, rtel1, brca2 and rad51*.   **B**. Fertilization success rate of progenies generated from crosses between male homozygous mutants (*blm*[-/-], *brca2*[-/-], *rad51*[-/-], *fanci*[-/-], *rad54l*[-/-] and *rtel1*[-/-]) and wild-type female zebrafish. **C.** Hematoxylin and eosin (H&E) staining of testis sections from the mutants of *blm*[cu55/cu55] and *brca2*[cu53/cu53] and their WT sibling controls. Green arrows indicate spermatogonia (sg), spermatocytes (sc) and mature spermatozoa (sz). Scale bar = 20 μm.

Figure 4. HSPC maintenance and repopulation defects in the *ddb1, pcid2, pcna* and *atad5a* mutant zebrafish. A. Representative images of *cmyb* WISH labeling HSPC population in the CHT at 3 or 4 dpf in *ddb1*[-/-](*cu43/cu43*), *pcid2*[-/-](*cu51/cu51*), *pcna*[-/-] (*cu29/cu29*) and *atad5a*[-/-](*cu33/cu33*), and their WT sibling controls. Black brackets indicate *cmyb*-positive HSPCs in CHT. Numbers at the bottom of images indicate the number of representative outcome observed out of the total number of embryos obtained. **B.** Merged images of bright-field and confocal microscopy to show EdU-positive proliferative cells in the

CHT between mutants and their WT sibling controls at 3 or 4 dpf and their quantification of the number of EdU-positive cells in the CHT. **C.** Merged images of bright-field and confocal microscopy to display apoptotic cells in the CHT using AO staining between mutants and WT controls at 3 or 4 dpf and their quantification of the apoptotic events in the CHT. Graphs represent mean ± S.E.M. with individual values. p-values were calculated by unpaired two-tailed Student's t-test. *** $p<0.001$, **$p<0.01$. n.s.; not significantly different. Scale bar = 200 μm.

Figure 5. Response of DNA replication/repair mutants to genotoxic stresses. A. Images of embryos generated from inbred heterozygous mutant of *atad5a*, *pcid2* and *shfm1*, in the presence or absence of 5Gy IR exposure at 6 hpf. Morphological alteration is indicated by red arrowheads in the homozygous mutants exposed to IR. **B.** Confocal microscope images of embryos stained with AO to show apoptotic cells in *atad5a* mutants and WT controls at 3 dpf in response to IR at 6 hpf and its quantification of the number of AO–positive apoptotic cells in the brain of *atad5a$^{-/-}$*(*cu33/cu33*) mutants and control with IR-treatment. Yellow dotted area indicates dorsal view of the brain area. **C.** Lateral views of *brca2, polk, and xrcc1* mutant embryos and their WT controls with treatment of 0.001% MMS from 6 hpf. Red arrowheads indicate altered body trunk of the mutants with MMS treatment. **D.** Confocal microscope images showing body trunk of AO-stained 2 dpf *xrcc1$^{-/-}$*(*cu61/cu61*) mutant embryos and WT siblings in the presence of 0.001% MMS from 6 hpf and its quantification of AO–positive apoptotic cells in the body trunk of *xrcc1* mutants and WT control with MMS treatment. White dotted box indicates neural tissues in the trunk of embryos. Quantification graphs represent mean ± S.E.M. with individual values. p-values were calculated by unpaired two-tailed Student's t-test. *** $p<0.001$, **$p<0.01$, *$p<0.05$. n.s.; not significantly

different. Black scale bar = 500 μm; white scale bar = 200 μm.

Figure 6. Atad5a is required for brain development during embryogenesis. A. Dorsal view of anterior part of zebrafish embryos showing expression of *atad5a* in the brain at 2 and 5 dpf using WISH. *atad5a*-expressing regions are indicated by abbreviated letters; Fb: forebrain, Mb: midbrain, Hb: hindbrain, Rt: retina. **B.** Western blot with Pcna antibody using chromatin fraction of whole embryo lysate from *atad5a*$^{-/-}$ and WT controls at 5 dpf. H3 is histone H3 antibody used to show equal loading of chromatin fraction for western blotting. **C.** Dorsal view of anterior part of *atad5a*$^{-/-}$ and WT controls at 5 dpf and its quantification of the brain size from dotted area. **D.** Quantification of the entire body length of *atad5a*$^{-/-}$ mutants and WT controls at 5 dpf. **E.** Confocal microscope images of EdU-positive cells and quantification of relative intensity of EdU signal in the brain (dotted area) between *atad5a*$^{-/-}$ and WT controls at 2 dpf. **F.** Confocal microscope images of AO-positive cells and quantification of AO-positive cell numbers in the brain (dotted area) between *atad5a*$^{-/-}$ mutants and WT controls at 2 dpf. Yellow-dotted area indicates the brain including forebrain to cerebellum. All graphs represent mean ± S.E.M. with individual values. p-values were calculated by unpaired two-tailed Student's t-test. *** p<0.001, **p<0.01, *p<0.05. n.s.; not significantly different. Scale bar = 200 μm.

Figure 7. p53-dependet apoptosis in the brain and CHT of *atda5a* mutant is induced by ATM/Chk2 activation. A. WISH staining shows that *p53* expression is highly induced in *atad5a* mutant brain compared to WT siblings. **B.** Upper panel shows confocal microscope images of AO-positive cells in the brain (dotted area) of *atad5a*$^{-/-}$ single, *atad5a*$^{-/-}$; *p53*$^{-/-}$

double mutants and WT controls at 2 dpf and their quantification of the number of apoptotic cells in the brain. Bottom panel displays dorsal views of zebrafish head area in the *atad5a*^{-/-} single, *atad5a*^{-/-}; *p53*^{-/-} double mutants and WT controls at 5 dpf and quantification of brain size around midbrain and forebrain (dotted area). Yellow-dotted area indicates the brain including forebrain to cerebellum. **C.** Western blot results with antibodies for phospho-Chk2 Thr68 (pChk2), Chk2, and phospho-histone H2AX Ser139 (pH2AX) using whole lysate of WT and *atad5a* mutant embryos at 5 dpf. H3 is histone H3 used to show equal loading of whole lysates for western blotting. **D.** Confocal images of AO staining using WT and *atad5a*^{-/-} embryos at 2 dpf with treatment of vehicle control (2% DMSO) and ATMi from 6 hpf and its quantification of the AO-positive cell number in the brain (dotted area). **E.** Images of *cmyb* WISH in the CHT of 5 dpf embryos of *atad5a*^{-/-} single, *atad5a*^{-/-}; *p53*^{-/-} double mutants and WT controls. **F.** *cmyb* WISH in the CHT of 4 dpf embryos of *atad5a*^{-/-} and WT controls with treatment of ATMi from 2 dpf. Numbers at the bottom of images indicate the number of representative outcome observed out of total number of embryos obtained. All graphs represent mean ± S.E.M. with individual values. p-values were calculated by unpaired two-tailed Student's t-test. *** p<0.001, *p<0.05. n.s.; not significantly different. Scale bar = 200 μm.

Figure 1. Large-scale morphological characterization of zebrafish CRISPR mutants related to DNA repair genes

Figure 2. Screening of viability and morphology of mutants during larval and juvenile stages

Figure 3. Mutants with sex reversal and abnormal germ cell development

A

blm · rtel1 · brca2 · rad51

■ Male   ■ Female

B

□ Fertilizaed eggs   ■ Unfertilized eggs

C

WT · blm^{cu55/cu55} · WT · brca2^{cu53/cu53}

Figure 4. . HSC maintenance and repopulation defects in the *ddb1*, *pcid2*, *pcna* and *atad5a* mutant zebrafish

Figure 5. Response of DNA replication/repair mutants to genotoxic stress

Figure 6. Atad5a is required for brain development during embryogenesis

Figure 7. p53-dependent apoptosis in the brain and CHT of *atad5a* mutants is induced by ATM/Chk2 activation.