CEP78 functions downstream of CEP350 to control biogenesis of primary cilia by

negatively regulating CP110 levels

André Brás Gonçalves[1], Beinta Biskopstø Joensen[1,5], Sarah Kirstine Hasselbalch[1,5], Pernille Mar-

tens[1], Signe Krogh Ohlsen[2], Mathieu Quinodoz[3,3.1,3.2], Konstantinos Nikopoulos[3], Magnus Per

Damsø Jeppesen[1], Catja Weiss[1], Søren Tvorup Christensen[1], Sebastian Patzke[4], Carlo Rivolta[3,3.1,3.2],

Jens S. Andersen[2], Pietro Farinelli[1*] and Lotte Bang Pedersen[1*]

[1]Department of Biology, Section for Cell Biology and Physiology, University of Copenhagen, Uni-

versitetsparken 13, DK-2100 Copenhagen Ø, Denmark

[2]Department of Biochemistry and Molecular Biology, University of Southern Denmark, Campusvej

55, 5230 Odense M, Denmark

[3]Institute of Molecular and Clinical Ophthalmology Basel (IOB), Basel, Switzerland

[3.1]Department of Ophthalmology, University of Basel, Basel, Switzerland

[3.2]Department of Genetics and Genome Biology, University of Leicester, Leicester, United King-

dom

[4]Department of Radiation Biology, Institute for Cancer Research, Norwegian Radium Hospital,

Oslo University Hospital, Oslo, Norway

[5]These authors contributed equally to this study

[*]For correspondance: Pietro.Farinelli@twelve.bio (PF); lbpedersen@bio.ku.dk (LBP)

Abstract

CEP78 is a centrosomal protein implicated in ciliogenesis and ciliary length control, and mutations in the *CEP78* gene cause retinal cone-rod dystrophy associated with hearing loss. However, the mechanism by which CEP78 affects cilia formation is unknown. Based on a recently-discovered disease-causing *CEP78* p.L150S mutation, we identified the disease-relevant interactome of CEP78. We confirmed that CEP78 interacts with the EDD1-DYRK2-DDB1^{VPRBP} E3 ubiquitin ligase complex, which is involved in CP110 ubiquitination and degradation, and identified a novel interaction between CEP78 and CEP350 that is weakened by the CEP78^{L150S} mutation. We show that CEP350 promotes centrosomal recruitment and stability of CEP78, which in turn leads to centrosomal recruitment of EDD1. Consistently, cells lacking CEP78 display significantly increased cellular and centrosomal levels of CP110, and depletion of CP110 in CEP78-deficient cells restored ciliation frequency to normal. We propose that CEP78 functions downstream of CEP350 to promote ciliogenesis by negatively regulating CP110 levels via an EDD1-dependent mechanism.

Introduction

Primary cilia are antenna-like sensory organelles that play pivotal roles in coordinating various signaling pathways important for human development and tissue homeostasis [1]. They comprise a microtubule-based axoneme core, which extends directly from the mother centriole-derived basal body and is surrounded by a bilayer membrane enriched for specific receptors, ion channels and lipids that endow the organelle with unique signaling properties [2]. Not surprisingly, mutations in genes that affect assembly, structure or function of primary cilia are causative for a growing number of pleiotropic diseases and syndromes called ciliopathies, which include cone-rod dystrophy in the retina and hearing loss (CRDHL; MIM# 617236) amongst others [3]. Ciliopathy genes include those coding for components of the centrosome, which contains the daughter and mother centriole and gives rise to the ciliary basal body. The mother centriole is distinguished from the daughter centriole by distal and subdistal appendages, which play critical roles in vesicle docking at the onset of ciliogenesis and in microtubule anchoring, respectively. In addition, the centrosome contains pericentriolar material and is associated with centriolar satellites that affect cilia biogenesis and function in various ways [4].

Assembly of primary cilia is a complex, multistep process that is tightly coordinated with the cell cycle. In actively-proliferating cells, centriolar coiled coil protein 110 (CP110; also known as CCP110) and centrosomal protein of 97 kDa (CEP97) cap the distal ends of both mother and daughter centrioles and suppress ciliogenesis. Furthermore, overexpression of CP110 in growth-arrested cells prevents ciliogenesis [5]. CP110 also regulates centriole duplication and length control during S phase and interacts with key regulators of centriole duplication, including PLK4 [6-9]. As cells enter G1/G0, ciliogenesis begins with recruitment of pre-ciliary vesicles to the distal end of the mother centriole. The vesicles subsequently fuse to form a larger ciliary vesicle underneath which the ciliary transition zone and axoneme are assembled. The axoneme is further extended by intraflagellar transport (IFT), and the ciliary vesicle expands and matures to form the ciliary membrane and a surrounding sheath that fuses with the plasma membrane upon completion of ciliogenesis [10-12].

Initiation of transition zone formation and axoneme extension during early stages of ciliogenesis requires removal of the CEP97-CP110 complex from the distal end of the mother centriole [5]. This process relies on M-Phase Phosphoprotein 9 (MPP9), which interacts directly with CEP97 and cooperates with kinesin KIF24 to recruit the CEP97-CP110 complex to the distal centriole end [13, 14]. During ciliogenesis, phosphorylation of MPP9 by Tau tubulin kinase 2 (TTBK2) leads to degradation of MPP9 by the ubiquitin proteasome system (UPS), which results in destabilization of

the CEP97-CP110 complex causing its removal from the distal end of the mother centriole [14].

TTBK2 is recruited to the mother centriole distal appendages by CEP164 [15, 16] where it also

phosphorylates CEP83 to promote CP110 removal [17]. Consequently, depletion of CEP164,

TTBK2, CEP83 or other centriole distal appendage proteins that regulate their localization and/or

function impairs ciliogenesis [18-23]. Mother centriole recruitment of TTBK2 and removal of the

distal CEP97-CP110 cap additionally requires the subdistal CEP350-FOP-CEP19 complex, which

furthermore interacts with RABL2 and IFT-B complex components to promote their axonemal en-

try [24-26].

Despite recent advances, the precise mechanisms by which CP110 regulates ciliogenesis and

is removed from the mother centriole at the onset of ciliogenesis remain incompletely understood.

For example, a recent study implicated the homologous to the E6AP carboxyl terminus (HECT)

type EDD1-DYRK2-DDB1^{VPRBP} E3 ligase complex in ubiquitination and degradation of CP110 via

a mechanism involving direct interaction of viral protein R binding protein (VPRBP; also known as

DCAF1) and centrosomal protein of 78 kDa (CEP78) [27]. Specifically, the authors reported that

CEP78 suppresses CP110 ubiquitination by EDD1 (also known as UBR5 and EDD) and it was pro-

posed that CP110 is phosphorylated by DYRK2 and thereby recognized and brought close to EDD1

by VPRBP. EDD1 then transfers ubiquitin to CP110 leading to its degradation. When CEP78 binds

VPRBP it induces a conformational change in the complex thereby preventing CP110 ubiquitina-

tion. Further, they demonstrated that depletion of CEP78 promoted centriole elongation, whereas

CEP78 over-expression inhibited primary cilia formation in hTERT-immortalized retinal pigment

epithelial (RPE1) cells [27]. On the other hand, knockout (KO) of *Dyrk2* in the mouse was reported

to result in elongation of primary cilia, but centrosomal CP110 levels appeared unaffected in *Dyrk2*

mouse KO cells [28]. Therefore, it remains uncertain how CEP78 and the EDD1-DYRK2-

DDB1^{VPRBP} complex affect CP110 homeostasis to control ciliogenesis.

CEP78 is composed of 16 exons and encodes a protein comprising 722 amino acids that

possesses five consecutive leucine rich repeats (LRR) at the N-terminus, and a coiled-coil (CC) do-

main at the C-terminus [29, 30]. Four independent studies have reported eight different mutations in

CEP78 in patients with CRDHL [29-32], whereas one study identified a homozygous *CEP78*-trun-

cating variant in a family with non-syndromic retinitis pigmentosa (MIM# 268003; [33]), another

form of retinal degeneration. Of these studies, two have investigated the functional consequences of

human *CEP78* mutations at the cellular level. In one study, whole-exome sequencing (WES) identi-

fied biallelic mutations in *CEP78* in two unrelated families from Greece and Sweden, respectively

[29]. The Greek subject had a homozygous base substitution at the splice donor site in intron 3 of *CEP78* (NM_032171.2:c.499+1G>T). Two subjects from a Swedish family carried heterozygous mutations, one base substitution in intron 3 (NM_032171.2:c.499+5G>A) and a single nucleotide deletion in exon 5 (NM_032171.2:c.633del; p.Trp212GlyfsTer18) causing a frameshift. These *CEP78* mutations lead to exon skipping and premature stop codons accompanied by almost unde-tectable levels of CEP78 protein in human skin fibroblasts (HSFs) of affected individuals. Further-more, it was found that HSFs from these patients have significantly longer primary cilia as com-pared to control cells [29]. More recently, a missense mutation in *CEP78* (NM_032171.2:c.449T>C; p.Leu150Ser) was identified in three unrelated families from Belgium and Germany diagnosed with CRDHL [30]. In the two Belgian families, affected individuals were homozygous for the p.Leu150Ser variant, whereas affected individuals from the German family dis-played compound heterozygosity for this variant and a novel splice site variant, NM_032171.2:c.1462-1G>T. In all cases, HSFs from patients harboring the p.Leu150Ser mutation (hereafter: L150S) displayed decreased cellular levels of CEP78 and significantly elongated cilia compared to control HSFs [30], as seen in patient-derived HSFs with CEP78 truncating mutations [29]. In addition, other ciliopathy features were reported in some of the affected individuals with the CEP78^{L150S} mutation, including obesity, respiratory problems, diabetes 2 and infertility [30]. In summary, available data derived from human patients indicates that depletion of CEP78 leads to elongation of primary cilia at the cellular level, which manifests in ciliopathy phenotypes at the or-ganism level. However, the mechanism by which CEP78 regulates ciliary length is not known.

In addition to the patient studies described above, some studies have addressed CEP78 func-tion in different cell culture models. First, a study showed that siRNA-mediated depletion of CEP78 in RPE1 cells reduces the frequency of cells forming primary cilia, possibly due to centriolar an-choring defects, but the length of the residual cilia was not assessed. Similarly, depletion of CEP78 in *Planarians* was shown to impair motile cilia formation due to defective docking of centrioles to the cell surface [34]. Another study identified CEP78 interaction with PLK4 and implicated CEP78 in PLK4-mediated centriole over duplication, whereas possible roles for CEP78 in relation to cilia were not addressed [35]. Finally, as mentioned above, a study reported that CEP78 directly interacts with VPRBP of the EDD1-DYRK2-DDB1^{VPRBP} E3 ligase complex to suppress CP110 ubiquitina-tion and thereby stabilize it [27]. How such CEP78-mediated stabilization of CP110 might lead to the long cilia phenotype observed in patient fibroblasts lacking CEP78 [29, 30] and/or the reduced ciliation frequencies seen in CEP78-depleted *Planarians* and RPE1 cells [34] is unclear.

Here were show that in RPE1 cells and patient-derived HSFs loss of CEP78 leads to reduced

ciliation frequency as well as increased length of the cilia that do form. By taking advantage of the

recently identified disease-causing CEP78^{L150S} mutation [30], we used a quantitative mass spec-

trometry-based approach to identify a disease-relevant interactome of CEP78. We confirmed that

CEP78 interacts with the EDD1-DYRK2-DDB1^{VPRBP} complex implicated in CP110 ubiquitination

and degradation [27], and furthermore identified a novel interaction between CEP78 and the N-ter-

minal region of CEP350. The interaction of CEP78 with both VPRBP and CEP350, as well as cen-

trosomal recruitment of CEP78, are dramatically reduced by the CEP78^{L150S} mutation. Lack of cen-

trosomal recruitment of CEP78^{L150S} is likely due to impaired interaction with CEP350 since centro-

somal and cellular levels of CEP78 were significantly decreased in *CEP350* KO cells. Conversely,

cells lacking CEP78 displayed significantly decreased centrosomal levels of EDD1 and unaltered or

slightly increased centrosomal levels of CEP350. In addition, CEP78-deficient cells showed signifi-

cantly increased cellular and centrosomal levels of CP110, presumably owing to the reduced centro-

somal levels of EDD1 observed in these cells, and depletion of CP110 in CEP78-deficient cells re-

stored ciliation frequency to normal. Collectively our results suggest that CEP78 functions down-

stream of CEP350 to promote cilia biogenesis by negatively regulating CP110 levels via an EDD1-

dependent mechanism.

Results

Depletion of CEP78 leads to fewer and longer primary cilia in cultured cells

Previous studies showed that patient-derived fibroblasts lacking CEP78 display significantly elon-

gated primary cilia compared to control cells [29, 30] whereas depletion of CEP78 in *Planarians*

and RPE1 cells was shown to significantly reduce cilia numbers [34]. To reconcile these seemingly

contradictory findings we analyzed cilia numbers and length in serum-deprived wild type (WT) and

CEP78 KO RPE1 cells [30] by immunofluorescence microscopy (IFM) with antibodies against cili-

ary (ARL13B) and centrosomal (CEP350) markers. We found that the frequency of ciliated cells is

significantly reduced in *CEP78* KO cells compared to WT cells (Figure 1A, B) whereas the length

of the remaining cilia is significantly increased in the *CEP78* KO cells (Figure 1A, C). Similar re-

sults were obtained using previously described patient-derived *CEP78*-deficient or control HSFs

(Figure 1D-F) [29, 30]. The low ciliation frequencies of serum-deprived *CEP78* mutant cells were

not secondary to cell cycle defects, as judged by western blot analysis with antibody against reti-

noblastoma protein phosphorylated at S807/811 (P-Rb; Figure 1G-L). However, in serum-fed cells,

P-Rb levels were significantly higher in *CEP78* mutant cells compared to controls (Figure 1G-L),

indicating that mutant cells may progress slower through S-phase [36]. This is in line with previous

reports indicating that CEP78 protein levels are upregulated in late S-G2 phase suggesting a role for

CEP78 at this cell cycle stage, e.g. during centriole duplication [27, 35]. We conclude that CEP78

deficient RPE1 and HSF cells display reduced ciliation frequencies as well as increased length of

remaining cilia, thus reconciling previous observations [29, 30, 34].

Analysis of RPE1 WT and *CEP78* KO cells expressing FLAG-CEP78 and FLAG-CEP78^{L150S}

To confirm the above results, we performed a rescue experiment by transiently expressing FLAG-

CEP78 in WT and *CEP78* KO RPE1 cells followed by serum-deprivation and IFM analysis using

FLAG and ARL13B antibodies. In parallel, we performed similar experiments with FLAG-

CEP78^{L150S}. We first analyzed the localization and expression levels of the two FLAG fusions in

serum-deprived RPE1 cells. The FLAG-CEP78 fusion protein localized to both centrioles at the

base of cilia as expected [27, 29, 35], whereas centriolar localization of FLAG-CEP78^{L150S} was se-

verely compromised although not completely abolished (Figure 2A, B). The latter result is in line

with previous reports showing that the LRR region in the N-terminus of CEP78, which encom-

passes residue L150, is important for its localization to centrioles [27, 35]. Both FLAG fusions were

expressed at similar levels in the cells (Figure 2-figure supplement 1A, B); this was somewhat sur-

prising given that endogenous CEP78^{L150S} is largely undetectable by western blot analysis of pa-

tient-derived HSFs suggesting its stability is compromised [30]. Therefore, we tested if the L150S

mutation might affect binding of the CEP78 antibody used in [30] to CEP78. However, FLAG im-

munoprecipitation (IP) and western blot analysis of 293T cells expressing FLAG-CEP78 or FLAG-

CEP78^{L150S} indicated that the CEP78 antibody binds equally well to WT CEP78 and the CEP78^{L150S}

mutant protein. This analysis also revealed that endogenous CEP78 co-IPs with FLAG-CEP78 indi-

cating that CEP78 can form homodimers/oligomers (Figure 2-figure supplement 2). These results

indicate that the L150S mutation compromises the recruitment of CEP78 to the centrosome, but not

its short-term stability, at least in RPE1 cells. We cannot rule out that the N-terminal FLAG tag may

stabilize FLAG-CEP78^{L150S} and prevent it from being degraded.

Next, we tested if FLAG-CEP78 or FLAG-CEP78^{L150S} could rescue the ciliary phenotypes

of *CEP78* KO cells. In WT cells, transient expression of FLAG-CEP78 lead to reduced ciliation fre-

quency compared to non-transfected cells, in agreement with a previous study [27], although this

result was not statistically significant (Figure 2C). Surprisingly, ciliated WT cells expressing

FLAG-CEP78 were significantly elongated compared to non-transfected controls, similar to un-transfected *CEP78* KO cells (Figure 1D). In contrast, expression of FLAG-CEP78 in the *CEP78* KO cells rescued the ciliary length phenotype of these cells (Figure 1D), indicating that precise control of cellular/centrosomal levels of CEP78 is needed for appropriate ciliary length control, with too much or too little CEP78 causing ciliary lengthening. Since transient expression of FLAG-CEP78 in the *CEP78* KO cells caused ciliary shortening (Figure 1D) it was not possible to assess whether this fusion protein could also rescue the reduced ciliation frequency phenotype of these cells. While the *CEP78* KO cells expressing FLAG-CEP78 still displayed significantly fewer cilia than WT cells (Figure 1C), we cannot exclude that the transfected *CEP78* KO cells initially had more cilia, which progressively became shorter and eventually disappeared during the course of the experiment (see also below and Figure 8 for rescue of the ciliation frequency phenotype of the *CEP78* KO cells). Finally, we obtained essentially identical results upon expression of FLAG-CEP78^{L150S} in both WT and *CEP78* KO cells (Figure 2C, D) indicating that despite its reduced localization to centrioles (Figure 1A, B), this mutant protein is still able to influence ciliary length in a manner paralleling WT FLAG-CEP78.

Taken together, our results indicate that 1) loss of CEP78 causes reduced ciliation frequency as well as increased length of remaining cilia in both RPE1 and HSF cells; 2) overexpression of CEP78 similarly leads to fewer but longer cilia; and 3) the L150S mutation compromises localization of CEP78 to centrioles, but not its short-term stability and ability to influence ciliary length.

CEP78 interacts with the EDD1-DYRK2-DDB1^{VPRBP} complex and CEP350

To explore how CEP78 is recruited to centrioles to regulate ciliary frequency and length we first used Stable Isotope Labeling by Amino acids in Cell culture (SILAC)-based quantitative mass spectrometry (MS) to identify the differential interactomes of FLAG-CEP78 and FLAG-CEP78^{L150S} expressed in 293T cells (see Materials and methods for details). We identified interaction between CEP78 and several components of the EDD1-DYRK2-DDB1^{VPRBP} complex (Figure 3A), in agreement with previous work showing direct binding of CEP78 to VPRBP within this complex [27]. Interestingly, the MS results indicated that the CEP78^{L150S} mutation dramatically reduces this interaction (Figure 3A), which we confirmed by IP and western blot analysis of 293T cells co-expressing Myc-VPRBP and FLAG-CEP78 or FLAG-CEP78^{L150S} (Figure 3B). In addition, our interactome analysis identified CEP350 as potential novel CEP78 binding partner (Figure 3A). Using IP and western blot analysis in 293T cells co-expressing FLAG-CEP78 or FLAG-CEP78^{L150S} with a Myc-

tagged N-terminal or C-terminal CEP350 truncation we found that CEP78 binds to the N-terminal

region of CEP350 (residues 1-983; Figure 3C, E); reciprocal co-IP with Myc-tagged CEP350 con-

structs confirmed this result (Figure 3-figure supplement 1). Furthermore, we found that the interac-

tion between CEP78 and CEP350 N-terminus is reduced by the CEP78^{L150S} mutation (Figure 3C).

Using a similar approach, we could not detect interaction between CEP78 and endogenous FOP

(Figure 3C), whereas under similar conditions endogenous FOP was readily co-IPed with the C-ter-

minus of CEP350 (Figure 3-figure supplement 1), in agreement with a previous report [37]. We also

mapped the CEP350 interaction site in CEP78 and found that CEP350 primarily binds to the C-ter-

minal region of CEP78 comprising residues 395-722 (Figure 3D, E), although a weak interaction

with a fragment comprising the entire LRR region (residues 2-403) was also observed. In contrast,

truncation of the latter fragment into two smaller fragments abolished binding to the CEP350 N-ter-

minus (Figure 3D, E). Binding of CEP78 to VPRBP similarly requires the entire LRR region of

CEP78 [27] and we therefore wondered whether interaction between CEP350 and CEP78 might be

mediated by VPRBP. However, IP experiments and western blot analysis failed to reveal physical

interaction between VPRBP and CEP350 N-terminus (Figure 3, figure supplement 1). We conclude

that CEP78 binds independently to VPRBP and the N-terminus of CEP350 and that both of these

interactions are reduced by the CEP78^{L150S} mutation.

CEP350 regulates the stability and centrosomal recruitment of CEP78

In non-ciliated HeLa and U2OS cells CEP78 localizes to both centrioles during all phases of the

cell cycle and appears to concentrate at the centriole wall [35]. In ciliated RPE1 cells CEP78 was

shown to localize to the distal end of both centrioles, with a preference for the mother centriole/ba-

sal body [27]. Similarly, CEP350 was reported to localize to the distal end of the mother centri-

ole/basal body, near the subdistal appendages, in ciliated RPE1 cells [24, 26, 38]. In agreement with

these reports and with our interactome and IP results (Figure 3), 3D Structured Illumination Micros-

copy (SIM) showed partial co-localization of CEP78 with CEP350 at the distal end of the basal

body of ciliated RPE1 cells, beneath the distal appendage marker CEP164 (Figure 4A). Further-

more, using a previously characterized RPE1 *CEP350* KO cell line [24], we found that depletion of

CEP350 not only reduces localization of CEP78 to the centrosome, but also its total cellular level

(Figure 4B-G). Similarly, we found significantly reduced levels of CEP78 in *FOP* KO cells (Figure

4 B, C), in line with the decreased centrosomal levels of CEP350 observed in these cells [24]. These

results indicate that CEP350 promotes the recruitment of CEP78 to the mother centriole/basal body
as well as its overall stability.

CEP78 mutant cells display reduced centrosomal levels of EDD1
Next, we asked if CEP78 might affect the centrosomal recruitment of components of the EDD1-
DYRK2-DDB1^{VPRBP} complex and/or CEP350, and analyzed the cellular or centrosomal levels of
relevant complex components in RPE1 WT and *CEP78* KO cells. The results showed that serum-
deprived, ciliated *CEP78* KO cells display significantly increased centrosomal levels of CEP350
(Figure 5A, B) as compared to controls. However, in serum-deprived, non -ciliated cells, the centro-
somal levels of CEP350 were not significantly different in *CEP78* KO cells compared to WT con-
trols (Figure 5C, D). Similar results were obtained for CEP350 in control and CEP78 deficient pa-
tient HSFs (Figure 5E-H). We also analyzed the cellular levels of VPRBP by western blotting and
the centrosomal levels of EDD1 by IFM in RPE1 WT and *CEP78* KO cells. Since the VPRBP anti-
body cross-reacted with an unknown protein of lower molecular mass in western blot analysis (data
not shown) we could not use this antibody to assess centrosomal levels of VPRBP in our cells, but
we note that a previous study concluded that CEP78 is not required for VPRBP recruitment to the
centrosome [27]. Similarly, we found that VPRBP cellular levels were not significantly increased in
serum-fed or serum-deprived *CEP78* KO cells compared to controls (Figure 6A-C). Thus, CEP78
seems to be dispensable for recruitment of VPRBP and CEP350 to the centrosome. In contrast, cen-
trosomal levels of EDD1 were significantly reduced in RPE1 *CEP78* KO cells compared to WT
controls, both in ciliated and non-ciliated cells (Figure 6D-G). Thus, CEP78 is required for efficient
recruitment of EDD1 to the centrosome. To corroborate this result, we measured the centrosomal
levels of EDD1 in RPE1 *CEP350* KO cells and found that EDD1 levels were also significantly re-
duced when compared to the controls (Figure 6-figure supplement 1). Together our data indicate
that CEP350, by recruiting CEP78 to the centrosome, also promotes the centrosomal recruitment of
EDD1.

Increased cellular and centrosomal levels of CP110 in CEP78 deficient cells
Since EDD1 is an E3 ubiquitin-protein ligase that negatively regulates CP110 levels [27], and
whose depletion impairs ciliogenesis [39, 40], we asked if the loss of CEP78 would lead to altered
levels of CP110 at the centrosome. Indeed, western blot and IFM analysis of RPE1 WT and *CEP78*
KO cells revealed that total cellular and centrosomal CP110 levels are significantly increased in

CEP78 KO cells compared to WT controls, both under serum-deprived (Figure 7A-D, F, G) and serum-fed conditions (Figure 7-figure supplement 1A-D). We also analyzed CP110 levels specifically at the mother centriole of ciliated RPE1 WT and *CEP78* KO cells and found no significant differences between them (Figure 7E). Similar results were obtained using *CEP78* mutant and control HSF cells (Figure 7G-H). Since a previous study reported that siRNA-mediated depletion of CEP78 in RPE1 and HeLa cells reduces the cellular levels of CP110 [27], a result contradicting our own observations (Figure 7; Figure 7-figure supplement 1), we tested if the elevated CP110 levels seen in our *CEP78* mutant cells might be due to compensatory upregulation of CP110 mRNA expression. However, RNA-Seq analysis indicated no significant changes in CP110 mRNA levels in patient-derived, *CEP78* deficient HSF cultures compared to controls; mRNA levels of CEP350, VPRBP and DDB1 in *CEP78* deficient HSF cultures were also similar to those of controls (Figure 7-figure supplement 2). We conclude that loss of CEP78 leads to elevated cellular and centrosomal levels of CP110 and that this is not due to upregulation of *CP110* transcription. Furthermore, our results indicate that in ciliated *CEP78* KO cells, the amount of CP110 present specifically at the mother centriole is not significantly higher than in control cells.

Partial depletion of CP110 normalizes ciliation frequency of *CEP78* KO cells

CP110 is a key negative regulator of ciliogenesis [5, 41] that was also shown to be required for ciliogenesis initiation as well as ciliary length control [42, 43]. The above results therefore prompted us to test if the reduced ciliation frequency and/or increased length of remaining cilia observed in *CEP78* deficient cells could result from elevated centrosomal CP110 levels in these cells. Indeed, when we used siRNA to partially deplete CP110 from RPE1 WT and *CEP78* KO cells and subjected the cells to serum deprivation, we found that the *CEP78* KO cells restored their ciliation frequency to WT level (Figure 8A, B), whereas partial CP110 depletion had little effect on cilia numbers in WT cells grown under similar conditions (Figure 8C, D). In addition, depletion of CP110 from *CEP78* KO cells also seemed to cause a reduction in length of residual cilia, although these cilia were still significantly longer than those of WT controls (Figure 8C, E). Thus, partial depletion of CP110 from *CEP78* KO cells can rescue their reduced ciliation frequency phenotype but likely not the increased length of remaining cilia.

Discussion

Several studies have shown that loss-of-function mutations in *CEP78* are causative of ciliopathy

characterized by CRDHL [29-32], and that cells lacking CEP78 display reduced ciliation frequency

[34] or abnormally long cilia [29, 30]. However, the molecular mechanisms by which CEP78 regu-

lates cilium biogenesis and length were so far unclear. In this study, we found that loss of CEP78

leads to decreased ciliation frequency as well as increased length of the residual cilia that do form,

both in RPE1 cells and patient-derived HSFs. In addition, overexpression of CEP78 in RPE1 cells

similarly causes reduced ciliation frequency, as reported previously [27]. The latter observation in-

dicates that overexpression of CEP78 has an inhibitory effect on ciliogenesis, possibly by sequester-

ing one or more ciliogenic binding partners, e.g. CEP350.

Interestingly, our study suggests that both ciliated and non-ciliated CEP78 deficient cells fail

to efficiently recruit EDD1 to the centrosome, whereas cellular and centrosomal levels of CP110 are

dramatically increased. The increased centrosomal levels of CP110 observed in serum-deprived

CEP78 mutant cells likely account, at least in part, for their reduced ciliation frequency, since par-

tial depletion of CP110 in these cells could restore ciliation frequency to normal. The removal of

CP110 from the distal end of the mother centriole constitutes a key step in the initiation of ciliogen-

esis and is regulated by a growing number of proteins that include TTBK2 [19] and components of

the UPS system such as Neurl-4 [44] and the cullin-3 (CUL3)–RBX1–KCTD10 complex [45]. The

EDD1-DYRK2-DDB1^{VPRBP} complex was similarly shown to mediate degradation of CP110 and to

interact with CEP78 [27], but the precise consequences of these activities for ciliogenesis were un-

clear. Our work strongly suggests that CEP78 functions together with the EDD1-DYRK2-

DDB1^{VPRBP} complex to negatively regulate centrosomal CP110 levels at the onset of ciliogenesis,

leading to its removal from the distal end of the mother centriole. This is in contrast to a previously

proposed model, which suggested that CEP78 inhibits degradation of CP110 by the EDD1-

DYRK2-DDB1^{VPRBP} complex [27]. The reason for this discrepancy is unclear, but may be due to

different experimental approaches used in our study and the paper by Hossain et al. [2]. In this latter

study, siRNA was used to partially deplete CEP78 from cultured cells, whereas we performed our

experiments on cells with endogenous CEP78 mutations. CP110 removal from the distal end of the

mother centriole occurs concomitantly with docking of the centriole to vesicles or the plasma mem-

brane at the onset of ciliogenesis [12]}. Therefore, a role for CEP78 in removal of CP110 from the

centrosome is compatible with ultrastructural analyses in *Planarians*, which indicated that basal

bodies fail to dock to the plasma membrane in CEP78 RNAi-depleted animals failure [34]. Alt-

hough not addressed in detail in the current study, the elevated centrosomal CP110 levels observed

in serum-fed, CEP78 deficient cells are also compatible with previous work implicating CEP78 in

regulation of PLK4-mediated centriole duplication [35].

We also uncovered a new physical interaction between CEP78 and the N-terminal region of

CEP350, and found that CEP350 is important for recruitment of CEP78 to the centrosome as well

as for its overall stability. The reduced interaction of the $CEP78^{L150S}$ mutant protein with CEP350

might explain its lack of detection by western blot analysis of patient HSFs [30], since reduced in-

teraction with CEP350 is likely to decrease the long-term stability of $CEP78^{L150S}$. Our IP and IFM

results furthermore suggests that CEP350 controls CP110 removal from the mother centriole not

only via recruitment of TTBK2 [24], but also via CEP78-dependent recruitment of EDD1. Indeed,

we observed reduced centrosomal levels of EDD1 in *CEP350* KO cells, consistent with this idea. It

remains to be determined whether these two pathways act in sequence or in parallel to control cen-

trosomal CP110 levels. Furthermore, as the recruitment of CEP350 to centrioles itself depends on

FOP [24, 26, 37] and on the distal centriole protein C2CD3 [14], loss of these proteins may simi-

larly affect the centrosomal recruitment of CEP78 and EDD1, and thereby negatively regulate

CP110 levels via this pathway. It will be important to investigate these possibilities in future stud-

ies.

While increased CP110 levels can account for the decreased ciliation frequency observed in

serum-deprived CEP78 mutant cells, it remains unclear why some of these cells form abnormally

long cilia despite elevated centrosomal and cellular levels of CP110. Our analysis indicated that the

ciliated mutant cells are able to locally displace CP110 from the mother centriole, as the amount of

CP110 present specifically at the mother centriole/basal body of these cells was the same as for

control cells. However, the underlying mechanism remains to be determined. Of note, the *CEP350*

KO cells used in our study also appear to be able to overcome increased centrosomal levels CP110

during ciliogenesis, since approximately 30% of these cells form cilia despite an inability to recruit

TTBK2 and remove CP110 from the mother centriole [24]. It should also be noted that some studies

have reported dual roles for CP110 during ciliogenesis, i.e. CP110 may not only function to inhibit

ciliogenesis but may also promote ciliary extension in some contexts [42, 43]. Although the mecha-

nism responsible for the elongated cilia phenotype seen in some of the CEP78 mutant cells remains

unclear, it is interesting that exogenously expressed WT CEP78 as well as $CEP78^{L150S}$ could rescue

this phenotype, even though the $CEP78^{L150S}$ mutant protein displayed reduced recruitment to the

centrosome. It is likely that in an over-expression context there are sufficient levels of $CEP78^{L150S}$

present at the centrosome to promote ciliary shortening in the mutant cells; alternatively, it might

sequester one or more factors in the cytosol needed for ciliary lengthening. One attractive candidate

for such a factor is CEP350, which we found to be present at abnormally high levels at the centro-

some of ciliated *CEP78* KO cells, specifically.

Interestingly, in addition to promoting ubiquitination and degradation of CP110 [27], EDD1

was also shown to mediate ubiquitination of CSPP-L in turn promoting its recruitment to centriolar

satellites and the centrosome [40]. CSPP-L is a well-known positive regulator of primary cilia for-

mation [46] that was recently shown to bind directly to CEP104 to control axoneme extension [47].

Furthermore, CEP104 is implicated in ciliogenesis through interaction with the CEP97-CP110 com-

plex and modulation of microtubule dynamics at the ciliary tip [47-49]. It will therefore be interest-

ing to investigate if and how CSPP-L and CEP104 localization is affected in CEP78 mutant cells.

Moreover, as CEP350 additionally binds to CYLD [50], a de-ubiquitinase that controls centriolar

satellite homeostasis and ciliogenesis through de-ubiquitination of the E3 ubiquitin ligase MIB1

[51, 52], and also regulates axonemal IFT particle injection through FOP-CEP19-RABL2 interac-

tions [24-26], it is tempting to speculate that CEP78 might impinge on these processes.

Materials and methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifier	Additional information
Cell line (*H. sapiens*)	hTERT-RPE1	ATCC	Cat# CRL-4000	Derived from human retinal pigment epithelium
Cell line (*H. sapiens*)	Puromycin-sensitive hTERT-RPE1	Francesc Garcia-Gonzalo, Autonomous University of Madrid, Madrid, Spain	N/A	
Cell line (*H. sapiens*)	hTERT-RPE1 *CEP78* KO	[30]	N/A	
Cell line (*H. sapiens*)	hTERT-RPE1-BFP-Cas9/pMCB306 (pool)	[24]	N/A	Control cell line for *CEP350* and *FOP* KO lines
Cell line (*H. sapiens*)	hTERT-RPE1-BFP-Cas9/ *CEP350* KO (pool)	[24]	N/A	
Cell line (*H. sapiens*)	hTERT-RPE1-BFP-Cas9/*FOP* KO (pool)	[24]	N/A	
Cell line (*H. sapiens*)	293T	ATCC	Cat# CRL-3216	Derived from human embryonic kidney
Primary cells (*H. sapiens*)	Control skin fibroblasts	[29]	N/A	
Primary cells (*H. sapiens*)	Compound heterozygous skin fibroblasts with two *CEP78* mutations: c.449T>C; p.Leu150Ser and c.1462-1G>T	[30]	F3: II:1	These cells do not express detectable levels of CEP78 [30]

Primary cells (*H. sapiens*)	Compound heterozygous skin fibroblasts with two *CEP78* mutations: c.633delC; p.Trp212Glyfs*18 and c.499+5G>A; IVS + 5G>A	[29]	2702r34, individual II-3	These cells do not express detectable levels of CEP78 [29]
Primary cells (*H. sapiens*)	Homozygous substitution in the first invariant base of intron 3 splice donor site in *CEP78*: c.499+1G>T (IVS3+1G>T)	[29]	KN10, individual II-1	These cells express almost undetectable levels of CEP78 [29]
Bacterial strain (*E. coli*)	DH10B	Lab stock	N/A	
Antibody	Anti-Acetylated-tubulin (mouse monoclonal)	Sigma-Aldrich	Cat# T7451	IFM (1:2000)
Antibody	Anti-α-tubulin (mouse monoclonal)	Sigma-Aldrich	Cat# T5168	WB (1:10000)
Antibody	Anti-ARL13B (rabbit polyclonal)	Proteintech	Cat# 17711-1-AP	IFM (1:500)
Antibody	Anti-CEP78 (rabbit polyclonal)	Bethyl Laboratories	Cat# A301-799A-M	IFM (1:200) WB (1:500)
Antibody	Anti-CEP350 (mouse monoclonal)	Abcam	Cat# Ab219831 (CL3423)	IFM (1:500)
Antibody	Anti-CEP350 (rabbit polyclonal)	Novus Biologicals	Cat# NB100-59811	IFM (1:200)
Antibody	Anti-CP110 (rabbit polyclonal)	Proteintech	Cat# 12780-1-AP	IFM (1:200) WB (1:1500)
Antibody	Anti-DCTN1/P-150 (mouse monoclonal)	BD Biosciences	Cat# 610474	IFM (1:500)
Antibody	Anti-EDD1 (rabbit polyclonal)	Bethyl Laboratories	Cat# A300-573A-M	IFM (1:100)

Antibody	Anti-FLAG (mouse monoclonal)	Sigma-Aldrich	Cat# F1804	IFM (1:500) WB (1:1000)
Antibody	Anti-FLAG (rabbit polyclonal)	Sigma-Aldrich	Cat# F7425	WB (1:1000)
Antibody	Anti-FGFR1OP/FOP (mouse monoclonal)	Novus Biologicals	Cat# H00011116-M01	WB (1:500)
Antibody	Anti-GAPDH (rabbit polyclonal)	Cell Signaling Technology	Cat# 2118	WB (1:1000)
Antibody	Anti-GFP (rabbit polyclonal)	Sigma-Aldrich	Cat# SAB4301138	WB (1:500)
Antibody	Anti-Myc (mouse monoclonal, 9B11)	Cell Signaling Technology	Cat# 2276S	WB (1:1000)
Antibody	Anti-phospho Rb (S807/811) (mouse monoclonal)	Cell Signaling Technology	Cat# 9308	WB (1:500)
Antibody	Anti-VPRBP (rabbit polyclonal)	Bethyl Laboratories	Cat# A301-877A-M	WB (1:200)
Antibody	Polyclonal Goat Anti-Mouse Immunoglobulins /Horseradish Peroxidase conjugated	Dako	Cat# P0447	WB (1:10000)
Antibody	Polyclonal Swine Anti-Rabbit Immunoglobulins /Horseradish Peroxidase conjugated	Dako	Cat# P0399	WB (1:10000)
Antibody	Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody/Alexa Fluor 488-conjugated	Thermo Fischer Scientific	Cat# A21202	IFM (1:500)

Antibody	Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody/Alexa Fluor 568-conjugated	Thermo Fischer Scientific	Cat # A10037	IFM (1:500)
Antibody	Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody/Alexa Fluor 488-conjugated	Thermo Fischer Scientific	Cat #A21206	IFM (1:500)
Antibody	Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody/Alexa Fluor 568-conjugated	Thermo Fischer Scientific	Cat# A10042	IFM (1:500)
Enzyme	USER® Enzyme	New England BioLabs	Cat# M5505L	
Enzyme	Phusion® High-Fidelity DNA polymerase	New England BioLabs	Cat# M0530L	
Recombinant DNA reagent	*H. sapiens* Angiomotin p80 (Ap80)/pCMV-FLAG	[53]	N/A	
Recombinant DNA reagent	*H. sapiens* CEP78 aa 2-217/pEGFP-C1	This study	CEP78 N-terminus (N)	PCR-amplified using primers h.CEP78.4f.kpn1 and hsEGFP-CEP78_Rv (651)
Recombinant DNA reagent	*H. sapiens* CEP78 aa 203-403/pEGFP-C1	This study	CEP78 Middle (M)	PCR-amplified using primers hsEGFP-CEP78_Fwd (607) and hsEGFP-CEP78_Rv (1209)

Recombinant DNA reagent	*H. sapiens* CEP78 aa 2-403/pEGFP-C1	This study	CEP78 N-terminus plus Middle (NM)	PCR-amplified using primers h.CEP78.4f.kpn1 and hsEGFP-CEP78_Rv (1209)
Recombinant DNA reagent	*H. sapiens* CEP78 aa 395-722/pEGFP-C1	This study	CEP78 C-terminus (C)	PCR-amplified using primers hsEGFP-CEP78_Fwd (1186) and hCEP78.rv_stop.bamh1
Recombinant DNA reagent	*H. sapiens* CEP78/pEGFP-C1	This study	N/A	PCR-amplified using primers h.CEP78.4f.kpn1 and hCEP78.rv_stop.bamh1
Recombinant DNA reagent	*H. sapiens* CEP78/pDEST306	This study	p3xFLAG-CEP78	
Recombinant DNA reagent	*H. sapiens* CEP78^{L150S}/pDEST306	This study	p3xFLAG-CEP78^{L150S}	
Recombinant DNA reagent	Plasmid pEGFP-C1 (empty vector)	TaKaRa Bio	Cat# 6084-1	
Recombinant DNA reagent	Gateway® pDONR^{TM}201	Invitrogen/Thermo Fischer Scientific	Cat# 11798-014	
Recombinant DNA reagent	Gateway® pDEST306	Invitrogen/Thermo Fischer Scientific	N/A (lab stock)	Cloning produces in-frame 3xFLAG tag in the N-terminus
Recombinant DNA reagent	*H. sapiens* CEP350 aa 1-983/pCS2$^+$ with a 6×myc tag	[50, 54]	CAP N	
Recombinant DNA reagent	*H. sapiens* CEP350 aa 2990-3117/pCS2$^+$ with a 6×myc tag	[50, 54]	CAP C	
Recombinant DNA reagent	*H. sapiens* VPRBP/pCMV-Myc	[55]	Myc-VPRBP	Cloned in Sal1 and Not1 sites

Sequence-based reagent	*H. sapiens CEP78* ORF/ GeneCopoiea pShuttle™Gateway® PLUS ORF	GeneCopoiea	NCBI reference Sequence NM_001098802.3	
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Sigma-Aldrich	attB forward	5'-GGGACAAGTTGTACAAAAAACAGGCTTCATCGACTCCGTGAAGCTGC-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Sigma-Aldrich	attB reverse	5-GGGGACCACTTTGTACAAGAAAGCTGGGTTTCAGGAATGCAGGTCCTTTCCAG-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Sigma-Aldrich	CEP78 mut_F	5'-AGAGACAGGTGCACCGAAGAAGCCGATTTATTCAATC-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Sigma-Aldrich	CEP78 mut_R	5'-GATTGAATAAATCGGCTTCTTCGGTGCACCTGTCTCT-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Eurofins Genomics	hCEP78.4f.kpn1	5'-AAGGTACCATCGACTCCGTGAAGCTGCG-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Eurofins Genomics	hsEGFP-CEP78_Rv (651)	5'-AAGGATCCTTAGCGAAGACTCTCAGCCCAG-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Eurofins Genomics	hsEGFP-CEP78_Fwd (607)	5'- GGGGTACCCAGACCATGAGAAGGCATGA-3'

Sequence-based reagent	*H. sapiens CEP78* PCR primer	Eurofins Genomics	hsEGFP-CEP78_Rv (1209)	5'-GGGGATCC TTAACCCCTG TGTCTTTTTGC AC-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Eurofins Genomics	hsEGFP-CEP78_Fwd (1186)	5'-GGGGTACC GCAGAACGTG CAAAAAGACA -3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Eurofins Genomics	hCEP78.rv_stop. bamh1	5'-GAGGATCC ACAGGAATGC AGGTCCTTTC-3'
Sequence-based reagent	Control siRNA	Eurofins Genomics	N/A	5'-UAAUGUAUU GGAAGGCA-3'
Sequence-based reagent	*H. sapiens CP110* siRNA	Eurofins Genomics /[5]	N/A	5'-GCAAAACCAG AAUACGAGAT T-3'
Chemical compound	DAPI (4', 6-diamidino-2-phenylindole)	Thermo Fischer Scientific	Cat# D1306	Stored at -20ºC as stock solution of 20 μg/ml in H_2O
Chemical compound	FuGENE®6	Promega	Cat# E2692	
Chemical compound	DharmaFECT™ Duo Transfection reagent	Dharmacon™	Cat# T-2010-03	
Software/ algorithm	cellSens Dimension	Olympus	Version 1.18	
Software/ algorithm	Zen Black 2012	Zeiss	Version 2012	
Software/ algorithm	Adobe Photoshop®	Adobe	Version 21.0.1	
Software/ algorithm	Adobe Illustrator®	Adobe	Version 24.2.1	
Software/ algorithm	ImageJ	NIH	Mac OS X or Windows	
Software/ algorithm	GraphPad Prism 6.0	GraphPad Software Inc.	Mac OS X or Windows	
Software/ algorithm	MaxQuant	[56]	Version 1.6.1.0	

Other	Anti-FLAG® M2 Affinity Gel	Sigma-Aldrich	Cat# A2220	
Other	Anti-c-Myc Agarose Affinity Gel antibody produced in rabbit	Sigma-Aldrich	Cat# A7470-1 ml	
Other	GFP-Trap® Agarose	ChromoTek GmbH	Cat# gta-20	
Other	QuickChange Lightning Site-Directed Mutagenesis Kit	Agilent	Cat# 210518	
Other	Direct-zol RNA Miniprep Kit	Zymo Research	Cat# R2051	
Other	Agilent RNA Nano 6000 Assay Kit	Agilent	Cat# 5067-1511	
Other	NEBNext® Ultra™ RNA Library Prep Kit for Illumina®	New England BioLabs	Cat# E7530L	
Other	Invitrogen™ Qubit® RNA Assay Kit	Thermo Fischer Scientific	Cat# 32852	
Other	AMPure XP system	Beckman Coulter Life Sciences	Cat# A63881	

PCR and cloning

For the establishment of the N-terminal FLAG-tagged CEP78 construct a cDNA clone coding for

human full-length CEP78 cDNA (GeneCopoiea pShuttle™Gateway® PLUS ORF, NCBI accession

number NM_001098802) was used as template for amplification with the Phusion High-Fidelity

DNA polymerase (New England BioLabs) under standard PCR conditions and attB forward and re-

verse primers (see Key Resources Table) compatible with the Gateway cloning technology (Invitro-

gen/Thermo Fisher Scientific) for inclusion of the insert into the pDonor201 Gateway vector (Invi-

trogen/Thermo Fischer Scientific). To produce human FLAG-CEP78^{L150S}, site directed mutagenesis

was performed with QuickChange Lightning Site-Directed Mutagenesis Kit (Agilent) and primers

CEP78 mut_F and CEP78 mut_R (see Key Resources Table). Accuracy of the *CEP78* ORF was

assessed by Sanger sequencing. Subsequently, the desired fragment was cloned into the pDEST306

vector (Invitrogen/Thermo Fisher Scientific.

Plasmids encoding full-length and truncated versions (N-terminus, middle, N-terminus plus

middle, C-terminus) of CEP78 tagged with GFP in the N-terminus were generated by PCR using

relevant primers (see Key Resources Table) and pFLAG-CEP78 as template; PCR products were

cloned into pEGFP-C1 by standard approaches, following digestion with BamHI (Roche, cat#

1056704001) and KpnI (Roche, cat#10899186001) and ligation with T4 DNA ligase (Applichem,

cat# A5188). Ligated plasmids were transformed into competent *E. coli* DH10B cells and cells har-

boring recombinant plasmids selected on Luria Bertani (LB; Sigma-Aldrich) agar plates with 50

μg/ml kanamycin. Plasmids were purified using Plasmid DNA Mini Kit I (Omega Biotech, cat#

D6943-02) or Nucleobond Xtra Mid kit (Macherey-Nagel, cat# 740410.50), according to protocols

supplied by the manufacturers. Sequences of plasmid inserts were verified by Eurofins Genomics.

Mammalian cell culture and transfection

Human skin fibroblasts were cultured as described previously [29, 30]. Human embryonic kidney-

derived 293T cells were cultured and transfected with plasmids as described previously [57]. RPE1

cells were grown in T-75 flasks in a 95% humidified incubator at 37 ºC with 5% CO2. Cells were

cultured in Dulbecco's Modified Eagle Medium (DMEM) (GIBCO®, cat# 41966-029) supple-

mented with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin (P/S). When cells were

about 80-90% confluent, they were passaged and set up for new experiments. Upon seeding, RPE1

cells were washed once with preheated 1 x phosphate buffered saline (PBS) and then incubated with

1% Trypsin-EDTA (ethylenediaminetetraacetic acid) (Sigma-Aldrich, cat# T4174) solution for 5

min at 37 ºC. The detached cells were aspired in an appropriate volume of new preheated DMEM

and seeded into petri dishes with or without sterile glass coverslips [58]. When the cells had reached

80-90% confluency they were serum starved for 24 hours to induce formation of primary cilia. A

small amount of cell solution was passaged on in a new T-75 flask containing new preheated

DMEM.

For transfection with plasmids, RPE1 cells were seeded in 60 mm petri dishes as described

above to about 50-70% confluency at the time of transfection. For rescue experiments, 2 μg

pFLAG-CEP78 or pFLAG-CEP78^{L150S} and 6 μl FuGENE®6 were mixed in 100 μl serum- and anti-

biotic free DMEM and incubated for 20 minutes at room temperature. New DMEM was added to

the cells prior to transfer of the transfection reagent, which was gently dripped onto the cells and

swirled to make sure it was dispersed well. After cells were incubated with transfection medium

for 4 hours, the medium was replaced with serum free DMEM for 24 hours to induce growth arrest.

Gene silencing in RPE1 cells by siRNA was performed using DharmaFect (see Key Resources Table). Briefly, RPE1 cells were seeded in DMEM medium with serum and antibiotics in 30 mm petri dishes at 20-25% confluency. Transfection was performed by combining 50 nM siRNA with 5 µl of DharmaFect in 200 µl of serum and antibiotic-free DMEM medium for 15 min at room temperature. After incubation, the complexes were added dropwise onto the cells and the media was swirled gently to ensure good dispersion prior to incubation at 37 °C. Cells were kept under these conditions for 48 hours and then serum starved for 24 hours before being processed for further experiments.

Immunofluorescence microscopy and image analysis

Standard epifluorescence IFM analysis was performed as described previously [58]. Briefly, glass coverslips containing cells of interest were washed in PBS and fixed with either 4% paraformaldehyde (PFA) solution for 15 min at room temperature or on ice, or with ice-cold methanol for 12-15 min at -20 ºC. After three washes in PBS the PFA-fixed cells were permeabilized 10 min in 0.2% Triton X-100 and 1% Bovine Serum Albumin (BSA) in PBS and blocked in 2% BSA in PBS for 1 hour at room temperature. Coverslips were incubated over night at 4 ºC in primary antibodies (see Key Resources Table) diluted in 2% BSA in PBS. The next day, coverslips were washed three times with PBS for 5 min and incubated for 1 hour at room temperature in secondary antibodies (see Key Resources Table) diluted in 2% BSA in PBS. Coverslips were washed three times in PBS for 5 min before staining with 2 µg/ml DAPI solution PBS for 30 sec. Finally, coverslips were washed once with PBS and before mounting on objective glass with 6% n-propyl gallate diluted in glycerol and 10x PBS and combined with Shandon™ Immuno-Mount™ (Thermo Scientific, cat# 9990402) in a 1:12 ratio. Cells were imaged on a motorized Olympus BX63 upright microscope equipped with a DP72 color, 12.8 megapixel, 4140 x 3096 resolution camera. cellSense Dimension software 1.18 from Olympus was used to measure cilia length; ImageJ version 2.0.0-rc-69/1.52i was used to measure the relative mean fluorescence intensity (MFI) of relevant antibody-labeled antigens at the centrosome/basal body. Images were first converted to 8 bit (grayscale) so that the intensity of the histogram always was constant between 0-255. A fixed circle was drawn around a centrosome in a cell. This same region of interest (ROI) was used to measure the MFI of a specific protein at the centrosome. A constant ROI was also drawn to measure the cell background signal which was subtracted from the MFI measured at the centrosome/basal body. Images were prepared for publication using Adobe Photoshop® and Adobe Illustrator®.

For super resolution imaging, RPE1 cells were photographed on a 3D-SIM Zeiss Elyra PS.1

microscope equipped with 488 and 561 nm diode lasers, two high-speed sCMOS pco-edge cameras

(scientific sCMOS 16-bit depth; 1280 x1280 pixel resolution; 0.04 μm pixel size, manufactured by

PCO), and a x100/ 1.46 NA alpha Plan-Apochromat oil immersion DIC M27 objective (Zeiss).

Multi-channel 3D-SIM image Z-stacks were acquired with three rotations of the grid and with 0.116

μm interval between each section. SIM images were reconstructed using Zeiss Zen Black 2012 soft-

ware. After acquisition, channel alignment was performed using calibrations based on GE reference

slide containing 100 nm diameter TetraSpeck Microspheres and the Z-stacks were three-dimension-

ally reconstructed using ImageJ (v 1.53c).

Statistical analysis

Statistical analyses were performed using GraphPad Prism 6.0. The background-corrected MFI

measured at the centrosome was normalized to relevant control cells. Mean and standard deviation

(SD) was calculated for all groups, and outliers were identified and removed by the ROUT method

before the statistical tests were conducted. The data was tested for normality using either D'Ago-

stino's K-squared test or Shapiro-Wilk test. Depending on the distribution of the data, two-tailed

and unpaired Student's t-test or Mann Whitney test were used when comparing two groups. One-

way ANOVA followed by Tukey's or Dunn's multiple comparison was used when comparing more

than two groups. Unless otherwise stated, the statistical analyses were performed on at least three

independent biological replicates (n=3). A p-value under 0.05 was considered statistically signifi-

cant and p-values are indicated in the figures with asterisks as follows; * $p<0.05$, ** $<p0.01$,

***<0.001, and **** $p<0.0001$.

Immunoprecipitation (IP), protein quantification, SDS-PAGE and western blot analysis

Immunoprecipitation of transfected 293T cell lysates was performed as described previously [57]

using relevant antibody-conjugated beads (see Key Resources Table). Protein concentrations were

measured using the DC^{TM} Protein Assay Kit I from Bio-Rad (cat # 5000111) by following the man-

ufacturer's protocol. For SDS-PAGE analysis of RPE1 and HSF cells, protein samples were pre-

pared by lysis of cells with 95 ºC SDS-lysis buffer (1% SDS, 10 mM Tris-HCl, pH 7.4) and cell ly-

sates transferred to Eppendorf tubes and heated shortly at 95 ºC. The samples were then sonicated

two times for 30 seconds to shear DNA followed by centrifugation at 20,000 x g for 15 min at room

temperature to pellet cell debris. Supernatants were transferred to new Eppendorf tubes and an ali-

quot of each sample used for determination of protein concentration. Samples with equal concentra-

tions of protein were prepared for SDS-PAGE analysis by addition of NuPAGE™ LDS Sample

Buffer (4X) from Thermo Fischer Scientific (cat# NP0007) and 50 mM DTT. Samples from IP

analysis in 293T cells, performed using modified EBC buffer [57], were prepared for SDS-PAGE in

a similar fashion. Protein samples were heated at 95 ºC for 5 minutes before loading them on a

Mini-PROTEAN® TGX™ Precast Gel 4-15% from Bio-Rab Laboratories, Inc. (cat# 456-1083 or

cat #456-1086). PageRuler™ Plus Prestained Protein Ladder (Thermo Fischer Scientific, cat#

26619) was used as molecular mass marker. SDS-PAGE was performed using the Mini-PRO-

TEAN® Tetra System from Bio-Rad. Gels were run at 100 V for 15 minutes and 200 V for 45

minutes. Using the Trans-BLOT® Turbo Transfer System from Bio-Rad Laboratories, Inc. the pro-

teins were transferred at 1.3 A, 25 V for 10 min from the gel to a Trans-Blot® Turbo™ Transfer

Pack, Mini format 0.2 μm Nitrocellulose membrane (cat# 1704158). Ponceau-S solution was added

to the membrane to visualize the proteins before blocking in 5% milk in Tris-buffered saline with

Tween-20 (TBS-T; 10 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.1% Tween-20) for 2 hours at room

temperature. Primary antibodies (see Key Resources Table) were diluted in 5% milk in TBS-T and

incubated with the membrane overnight at 4ºC. The membrane was washed three times 10 min at

room temperature in TBS-T on a shaker before incubation with secondary antibody (see Key Re-

sources Table), diluted in 5% milk in TBS-T, for 1 hour at room temperature. The membrane was

washed three times 10 min at room temperature in TBS-T on a shaker. Finally, SuperSignal™ west

Pico PLUS chemiluminescent Substrate (Thermo Scientific, cat # 34580) was mixed in a 1:1 ratio

and added to the membrane for 5 min before development on a FUSION FX SPECTRA machine

from Vilber.

Quantitative mass spectrometry (MS) and peptide identification

For SILAC-based mass spectrometry, three distinct SILAC culture media were used: light-light-

(Lys0 and Arg0); medium- (Lys4 and Arg6) and high-labeled media (Lys8 and Arg10). 293T cells

were seeded in 100 mm petri dishes and SILAC labeled for 1 week to ensure proper amino acid in-

corporation. After the incorporation phase, the cells were transfected with 2 μg of plasmid encoding

FLAG-Ap80 (control), FLAG-CEP78 or FLAG-CEP78^{L150S} using 6 μl FuGENE®6, as described

above, and then subjected to lysis using modified EBC buffer [56] and FLAG IP as described

above. Affinity enriched proteins were digested with trypsin and the resulting peptides desalted

prior to LCMS analysis using an Easy-nLC 1000 system (Thermo Scientific) connected to a Q Ex-active HF-X mass spectrometer (Thermo Scientific). Peptides were separated by a 70 minutes gradient using increased buffer B (95% ACN, 0.5% acetic acid). The instrument was running in positive ion mode with MS resolution set at 60,000 for a scan range of 300 to 1700 m/z. Precursors were fragmented by higher-energy collisional dissociation (HCD) with normalized collisional energy of 28 eV. For protein identification and quantitation, the obtained MS raw files were processed by MaxQuant software version 1.6.1.0 [56] and searched against a FASTA file from UniProt.

RNAseq analysis

RNA was extracted using the Direct-zol RNA Miniprep Kit (Zymo Research) following the manufacturer's instructions. RNA degradation and contamination were monitored on 1% agarose gels. RNA purity was checked using the NanoPhotometer® spectrophotometer (IMPLEN, CA, USA). RNA concentration was measured using Invitrogen™ Qubit® RNA Assay Kit in Qubit® 2.0 Fluorometer (Thermo Fischer Scientific). RNA integrity was assessed using the Agilent RNA Nano 6000 Assay Kit of the Bioanalyzer 2100 system (Agilent, CA, USA). A total amount of 3 µg RNA per sample was used as input material for the RNA sample preparations. Sequencing libraries were generated using NEBNext® Ultra™ RNA Library Prep Kit for Illumina® (New England BioLabs, USA) following manufacturer's recommendations and index codes were added to attribute sequences to each sample. Briefly, mRNA was purified from total RNA using poly-T oligo-attached magnetic beads. Fragmentation was carried out using divalent cations under elevated temperature in NEBNext First Strand Synthesis Reaction Buffer（5X). First strand cDNA was synthesized using random hexamer primer and M-MuLV Reverse Transcriptase（RNase H -）. Second strand cDNA synthesis was subsequently performed using DNA Polymerase I and RNase H. Remaining overhangs were converted into blunt ends via exonuclease/polymerase activities. After adenylation of 3' ends of DNA fragments, NEBNext Adaptor with hairpin loop structure were ligated to prepare for hybridization. In order to select cDNA fragments of preferentially 150~200 bp in length, the library fragments were purified with AMPure XP system (Beckman Coulter Life Sciences). Then 3 µl USER® Enzyme (New England BioLabs) was used with size-selected, adaptor-ligated cDNA at 37°C for 15 min followed by 5 min at 95 °C before PCR. Then PCR was performed with Phusion® High-Fidelity DNA polymerase, Universal PCR primers and Index (X) Primer (New England BioLabs). At last, PCR products were purified (AMPure XP system) and library quality was assessed on the Agilent Bioanalyzer 2100 system. The clustering of the index-coded samples was performed

on a cBot Cluster Generation System using HiSeq PE Cluster Kit cBot-HS (Illumina) according to the manufacturer's instructions. After cluster generation, the library preparations were sequenced on an Illumina Hiseq platform and 125 bp/150 bp paired-end reads were generated. Raw data (raw reads) of fastq format were firstly processed through in-house perl scripts. In this step, clean data (clean reads) were obtained by removing reads containing adapter, reads containing ploy-N and lowquality reads from raw data. At the same time, Q20, Q30 and GC content the clean data were calculated. All the downstream analyses were based on the clean data with high quality. Reference genome and gene model annotation files were downloaded from genome website directly. Index of the reference genome was built using Bowtie v2.2.3 [59] and paired-end clean reads were aligned to the reference genome using TopHat v2.0.12 [60]. We selected TopHat as the mapping tool for that TopHat can generate a database of splice junctions based on the gene model annotation file and thus a better mapping result than other non-splice mapping tools. HTSeq v0.6.1 [61] was used to count the reads numbers mapped to each gene and then Fragments Per Kilobase of transcript sequence per Millions base pairs sequenced (FPKM) of each gene was calculated based on the length of the gene and reads count mapped to this gene. FPKM considers the effect of sequencing depth and gene length for the reads count at the same time, and is currently the most commonly used method for estimating gene expression levels [62]. Prior to differential gene expression analysis, for each sequenced library, the read counts were adjusted by edgeR program package through one scaling normalized factor [63]. Differential expression analysis of two conditions was performed using the DEGSeq R package (1.20.0) [64].

Acknowledgements

We thank Søren Lek Johansen and Maria Schrøder Holm for expert technical assistance, Benedicte Schultz Kappel and Frida Roikjer Rasmussen for help with plasmid generation, and Nynne Christensen, Center for Advanced Bioimaging (CAB) Denmark, for help with 3D-SIM. We are grateful to Drs. Peter K. Jackson, Tomoharu Kanie, Francesc Garcia-Gonzalo, Laurence Pelletier, Brian David Dynlacht, Anne-Marie Tassin, Maria Gavilan, Rosa M. Rios, Éric A. Cohen and Peter ten Dijke for reagents. This study was supported grants from Independent Research Fund Denmark (#8020-00162B to PF and LBP, and #8021-00425A to JSA), from the Carlsberg Foundation (#CF18-0294 to LBP) and from the Swiss National Science Foundation (#176097 to CR).

Author contributions

ABG, Performed experiments, Analyzed the data, Prepared the figures; BBJ, Performed experi-

ments, Analyzed the data, Contributed to manuscript preparation; SKH, PM, SKO, MQ, KN,

MPDJ, CW, Performed experiments, Analyzed the data; STC, Contributed to manuscript prepara-

tion; SP, Analyzed data, Contributed to figure preparation; CR, JSA, Contributed to manuscript

preparation, Obtained funding; PF, Conception and design, Performed experiments, Contributed to

manuscript preparation, Obtained funding; LBP, Conception and design, Prepared the figures,

Wrote the manuscript, Obtained funding.

Additional files

Source data for Figure 3A

Source data for Figure 7-figure supplement 2

Figure 1. CEP78 mutant cells display fewer, but longer primary cilia. (**A**) Representative IFM im-

ages of serum-deprived RPE1 WT and *CEP78* KO cells stained with the indicated antibodies; DAPI

marks the nucleus. Scale bar, 5μm. (**B**, **C**) Quantification of the percentage of ciliated cells (**B**) and

the length of residual cilia (**C**) in WT and *CEP78* KO cells, based on images as shown in (A). Data

were normalized in relation to WT values. Student's t-test (two-tailed, unpaired) was used for the

statistical analysis. Graphs in (B) represent accumulated data from three individual experiments

(n=384 for WT and n=322 for *CEP78* KO cells). Graphs in (C) show data from three individual ex-

periments (n=338 for WT and n=198 for *CEP78* KO cells). (**D**) Representative IFM images of se-

rum-deprived control and *CEP78* mutant (Patient) HSFs stained with the indicated antibodies;

DAPI marks the nucleus. Scale bar, 5 μm. (**E**, **F**) Quantification of the percentage of ciliated cells

(**E**) and the length of residual cilia (**F**) in control and *CEP78* patient HSFs (data from HSFs derived

from patient 2702r34, individual II-3 [29]), based on images as shown in (D). Graphs in (E) repre-

sent accumulated data from seven individual experiments (n= 678 for control HSFs; n= 707 for pa-

tient-derived HSFs). Graphs in (F) show data from seven individual experiments (n= 237 for con-

trol HSFs; n= 216 for patient-derived HSFs). Data were normalized in relation to control values. A

Student's t-test (unpaired, two-tailed) was performed to address differences in the ciliary frequency

and length between control and patient HSFs. (**G**) Western blot analysis of Rb phosphorylated on

S807/811 (P-Rb) in RPE1 WT and *CEP78* KO cells grown with or without serum. (**H**, **I**) Quantifi-

cation of data shown in (G) from three independent experiments analyzed in duplicates. (**J**) P-Rb

blots from HSFs derived from control and patient HSFs grown with or without serum (data from

HSFs derived from patient F3: II:1; [30]). GAPDH was used as a loading control. (**K, L**) Quantification of data shown in (J) from three independent experiments analyzed in duplicates. Student's t-test (two-tailed, unpaired) was used for the statistical analysis in (H, I) and (K, L). Error bars of graphs represent SD and data are shown as mean ± SD. a.u., arbitrary units; (*) $p < 0.05$; (****) $p < 0.0001$; n.s., not statistically significant.

Figure 2. Analysis of RPE1 WT and *CEP78* KO cells expressing FLAG-CEP78 and FLAG-CEP78[L150S]. (**A**) Representative IFM images of serum-starved WT and *CEP78* KO cells transiently expressing FLAG-CEP78 and FLAG-CEP78[L150S] labeled with antibodies against FLAG (green) and ARL13B (magenta). Insets show enlarged views of the cilia-centrosome region. Scale bars: 5 μm in original images and 1 μm in magnified images. (**B**) Quantification of the relative MFI of FLAG-CEP78 and FLAG-CEP78[L150S] at the centrosome in serum-starved WT cells. Accumulated data from three individual experiments (n=145 and n=151 for cells transfected with FLAG-CEP78 and FLAG-CEP78[L150S], respectively). Student's t-test (two-tailed, unpaired) was used for the statistical analysis. Data is shown as mean ±SD. (**C**) Quantification of the percentage of ciliated cells in serum-deprived WT and *CEP78* KO cells expressing the indicated FLAG fusions shown in (A). (**D**) Quantification of the ciliary length in the same group of cells grown in the same conditions mentioned above. Analysis on cilia numbers and size was performed through accumulated data from three independent experiments (n=338 for untransfected WT; n=69 for WT+FLAG-CEP78; n=66 for WT+FLAG-CEP78[L150S]; n=198 for untransfected *CEP78* KO; n=35 for *CEP78* KO + FLAG-CEP78; n=16 for *CEP78* KO + FLAG-CEP78[L150S]). Ordinary one-way ANOVA with Dunnett's multiple comparison test was used for the statistical analysis of the ciliary frequency and length amongst all groups, in relation to untransfected WT cells mean (control group). Error bars in (C) indicate SD and data in (D) is presented as mean ± SD. Differences in cilia numbers and size between transfected and untransfected *CEP78* KO cells were assessed by Ordinary one-way ANOVA with Dunnett's multiple comparison test in which the untransfected *CEP78* KO cells were selected as the control group. a.u., arbitrary units; (*) $p < 0.05$; (**) $p < 0.01$; (***) $p < 0.001$; (****) $p < 0.0001$; n.s., not statistically significant.

Figure 3. CEP78 interacts with the EDD-DYRK2-DDB1[VPRBP] complex and CEP350. (**A**) Overview of results from CEP78 interactome analysis in 293T cells grown in SILAC medium. Cells expressing FLAG-CEP78 WT, FLAG-CEP78[L150S] or FLAG-Ap80 (negative control) were subjected to FLAG

IP and pellets analyzed by mass spectrometry. The upper panel displays affinity enrichment of Cep78

and interaction partners in FLAG-CEP78 WT cells relative to control cells. The lower panel shows

how the CEP78^{L150S} mutation weakens these interactions relative to CEP78 WT using ratios normal-

ized for equal affinity enrichment of FLAG-tagged CEP78. (**B-D**) 293T cells expressing the indicated

Myc-, FLAG-, or GFP-fusions were subjected to IP with anti-FLAG (**B**) or -GFP (**C**, **D**) beads and

input and pellet fractions analyzed by SDS-PAGE and western blotting with Myc, FLAG or GFP

antibodies as indicated. FLAG-Ap80-NB used in (B) is a negative control. Molecular mass markers

are indicated in kDa to the left of the blots. (**E**) Schematic of the CEP350 and CEP78 fusions used in

IP analysis. LRR, leucine-rich repeat; CC, coiled coil. The CEP350 constructs were described in [50,

54]; the CEP78 constructs were generated in this study based on published sequence information [27,

30, 35]. N, N-terminal region; M, middle region; NM, N-terminal plus middle region; C, C-terminal

region; FL, full length.

Figure 4. CEP350 regulates stability and centrosomal recruitment of CEP78. (**A**) 3D-SIM analysis

of CEP78 localization in serum-deprived WT RPE1 cells. Cells were fixed with PFA and stained

with the indicated antibodies. (**B**) Western blot analysis of CEP78 and α-tubulin (loading control)

in serum-deprived hTERT-RPE1-BFP-Cas9 control (pMCB306), *CEP350* KO or *FOP* KO cells

[24]. (**C**) Quantification of data in (B), based on three independent experiments analyzed in dupli-

cates. Error bars indicate SD. Statistical analysis was done using Student's t-test (unpaired, two-

tailed). (**D, F**) IFM analysis of serum-deprived control and *CEP350* KO ciliated (D) and non-cili-

ated cells (F) using antibodies as indicated. Dashed boxes indicate cropped images to highlight the

centrosomal/ciliary area. Scale bars: 5 μm in representative images and 1 μm in insets. (**E, G**)

Quantification of the relative MFI of CEP78 at the centrosome in serum-starved, ciliated (E) and

non-ciliated (G) control and *CEP350* KO cells, based on images as shown in (D) and (G), respec-

tively. Statistical analysis was done using a two-tailed and unpaired Mann-Whitney test for ciliated

cells and two-tailed and unpaired Student's t-test. Accumulated data from three individual experi-

ments (n=128 and 132 for control and *CEP350* KO ciliated cells, respectively; n= 101 and 107 for

control and *CEP350* KO non-ciliated cells, respectively). a.u., arbitrary units; (*) $p<0.05$; (**)

$p<0.01$; (****) $p<0.0001$.

Figure 5. Altered centrosomal levels of CEP350 in *CEP78* mutant cells. (**A, C, E, G**) Representa-

tive IFM images of ciliated (**A**) and non-ciliated (**C**) serum-deprived RPE1 WT and *CEP78* KO

cells, and ciliated (**E**) and non-ciliated (**G**) serum-deprived control and CEP78 deficient (Patient) HSFs (combined data from HSFs from patient F3: II:1 [30] and patient 2702r34, individual II-3 [29]) labeled with antibodies indicated in the figure. Insets show closeups of the centrosomal/ciliary area (dashed boxes). Scale bars: 5 μm in representative images and 1 μm in closeups. (**B, D**). Quantification of CEP350 relative MFI at the centrosome based on images as shown in (A, C). Data is shown as mean ± SD. A Student's t-test was used as statistical analysis between the two groups based on three individual experiments (n=157 or 158 for ciliated cells; n=144-152 for non-ciliated cells). (**F, H**) Quantification of CEP350 relative MFI at the centrosome in HSFs based on images as shown in (E, G). Data is shown as mean ± SD. Student's t-test was used as statistical analysis between the two cell groups based on seven individual experiments with a total of 154-156 analyzed ciliated cells (**F**), and three individual experiments with a total of 122-126 analyzed non-ciliated cells (**H**). a. u., arbitrary units; n.s, not statistically significant; (****) p<0.0001.

Figure 6. Analysis of EDD-DYRK2-DDB1^{VPRBP} complex components in CEP78 deficient cells. (**A**) Western blot analysis of VPRBP in lysates from serum-fed and serum-deprived RPE1 WT and *CEP78* KO cells. α-tubulin was used as a loading control (**B, C**) Quantification of the VPRBP relative levels in the different conditions depicted in (A). Statistical analysis was performed using a Student's t-test (unpaired, two-tailed) from five independent experiments analyzed in duplicates. Error bars indicate SD. (**D, F**) Representative IFM images of ciliated (**D**) and non-ciliated (**F**) serum-deprived RPE1 WT and *CEP78* KO cells labeled with antibodies against EDD1 (green) and DCTN1 plus acetylated tubulin (magenta). DAPI was used to mark the nucleus (blue). Insets show enlarged views of the cilium-centrosome region. Scale bars: 5 μm in original images and 1 μm in closeups. (**E, G**) Quantification of the EDD1 MFI at the centrosome based on images as shown in (D) and (F) using a two-tailed and unpaired Student's t-test. Data is shown as mean ± SD. Student's t-test was used from three independent experiments (n=194 and n=201 for ciliated WT and *CEP78* KO cells, respectively; n=194 and n=217 for non-ciliated WT and *CEP78* KO cells, respectively). a.u., arbitrary units; n.s., not statistically significant; (****) p<0.0001.

Figure 7. CEP78 deficient cells display elevated cellular and centrosomal levels of CP110. (**A**) Western blot analysis of lysates from serum-deprived RPE1 WT and *CEP78* KO cells using indicated antibodies. GAPDH was used as a loading control. (**B**) Quantification of the data shown in

(A), based on four independent experiments analyzed in duplicates. Error bars indicate SD. Statistical analysis was done using Student's t-test (unpaired, two-tailed). (**C, F**) Representative IFM images of serum-starved ciliated (**C**) and non-ciliated (**F**) RPE1 WT and *CEP78* KO cells labeled with antibodies against CP110 (green), combined DCTN1 and acetylated tubulin (Ac-tub; magenta) and DAPI to mark the nucleus (blue). Insets show enlarged views of the cilium-centrosome area. Scale bars: 5 μm in original images and 1 μm in closeups. (**D, G**) Quantification of the relative MFI of CP110 at the centrosome based on images as shown in (C) and (F), respectively. Student's t-test (unpaired, two-tailed) from three independent biological experiments (n= 90 and n=75 for ciliated RPE1 WT and *CEP78* KO cells, respectively; n=75 and n=82 for non-ciliated RPE1 WT and CEP78 KO cells, respectively) was used for statistical analysis. Data is presented as mean ± SD. (**E**) Quantification of the relative MFI of CP110 at the mother centriole based on images shown in (C) Mann Whitney test (two-tailed and unpaired) was used as statistical analysis based on two independent experiments (n=61 and n=62 for RPE1 WT and CEP78 KO cells, respectively). Data is presented as mean ± SD. (**H**) Representative IFM images of serum-deprived healthy and CEP78 deficient (Patient) HSFs (data from patient 2702r34, individual II-3 described in [29]) labeled with the indicated antibodies. DAPI was used as counterstaining to mark the nucleus. Dashed lines show closeup images of the centrosome region. Scale bars: 5 μm in original images and 1 μm in closeups. (**I**) Quantification based on observations of 151 and 155 cells of CEP78 control and patient cells, respectively, from three individual experiments. Student's t-test (unpaired and two-tailed) was used to assess differences between the two groups. a.u., arbitrary units; n.s., not statistically significant; (******) p<0.01; (*******) p<0.001; (********) p<0.0001.

Figure 8. Depletion of CP110 from *CEP78* KO cells rescues their ciliation frequency phenotype. (**A**) Representative western blots of RPE1 WT or *CEP78* KO cells treated with control (siMock) or CP110-specific siRNA (siCP110). Cells were deprived of serum and blots were probed with antibody against CP110 or α-tubulin (loading control). (**B**) Quantification of relative cellular CP110 levels under different conditions based on blots as shown in (A) (n=3 for WT and n=4 for *CEP78* KO cells; each sample was analyzed in duplicates). Data is represented as mean ±SD. A Student's t-test was used to infer CP110 knock down efficiency between WT cells treated with siMock and siCP110. The same statistical method was applied to the *CEP78* KO cells treated with the same siRNAs. (**C**) Representative IFM images of serum-deprived RPE1 WT and *CEP78* KO cells treated with control (siMock) or CP110-specific siRNA (siCP110) and stained for ARL13B (green). DAPI

marks the nucleus. Dashed boxes indicate cropped images to highlight the centrosomal/ciliary zone.

Scale bars, 5 μm in representative images and 1μm in insets. (**D**, **E**) Quantification of percentage of

ciliated cells (**D**) and length of remaining cilia (**E**) in RPE1 WT or *CEP78* KO cells treated with

control (siMock) or CP110-specific siRNA (siCP110). The quantification is based on images as

shown in (C) and represents data from three and four independent experiments, respectively, for

RPE1 WT (n=432-455 cells per condition) and *CEP78* KO cells (n=722-846 cells per condition).

Ordinary one-way ANOVA with Dunnett's multiple comparison test was used, where RPE1 WT

treated with siMock was selected as the control group. a.u., arbitrary units; (*) $p<0.005$; (***)

$p<0.001$; (****) $p<0.001$.

References

1. Anvarian, Z., K. Mykytyn, S. Mukhopadhyay, L.B. Pedersen, and S.T. Christensen, *Cellular signalling by primary cilia in development, organ function and disease.* Nat Rev Nephrol, 2019. **15**(4): p. 199-219.
2. Nachury, M.V. and D.U. Mick, *Establishing and regulating the composition of cilia for signal transduction.* Nat Rev Mol Cell Biol, 2019. **20**(7): p. 389-405.
3. Reiter, J.F. and M.R. Leroux, *Genes and molecular pathways underpinning ciliopathies.* Nat Rev Mol Cell Biol, 2017. **18**(9): p. 533-547.
4. Breslow, D.K. and A.J. Holland, *Mechanism and Regulation of Centriole and Cilium Biogenesis.* Annu Rev Biochem, 2019. **88**: p. 691-724.
5. Spektor, A., W.Y. Tsang, D. Khoo, and B.D. Dynlacht, *Cep97 and CP110 suppress a cilia assembly program.* Cell, 2007. **130**(4): p. 678-90.
6. Chen, Z., V.B. Indjeian, M. McManus, L. Wang, and B.D. Dynlacht, *CP110, a cell cycle-dependent CDK substrate, regulates centrosome duplication in human cells.* Dev Cell, 2002. **3**(3): p. 339-50.
7. Kleylein-Sohn, J., J. Westendorf, M. Le Clech, R. Habedanck, Y.D. Stierhof, and E.A. Nigg, *Plk4-induced centriole biogenesis in human cells.* Developmental cell, 2007. **13**(2): p. 190-202.
8. D'Angiolella, V., V. Donato, S. Vijayakumar, A. Saraf, L. Florens, M.P. Washburn, *et al.*, *SCF(Cyclin F) controls centrosome homeostasis and mitotic fidelity through CP110 degradation.* Nature, 2010. **466**(7302): p. 138-42.
9. Li, J., V. D'Angiolella, E.S. Seeley, S. Kim, T. Kobayashi, W. Fu, *et al.*, *USP33 regulates centrosome biogenesis via deubiquitination of the centriolar protein CP110.* Nature, 2013. **495**(7440): p. 255-9.
10. Sorokin, S., *Centrioles and the formation of rudimentary cilia by fibroblasts and smooth muscle cells.* J Cell Biol, 1962. **15**: p. 363-377.
11. Sorokin, S.P., *Reconstructions of centriole formation and ciliogenesis in mammalian lungs.* J Cell Sci, 1968. **3**(2): p. 207-30.
12. Shakya, S. and C.J. Westlake, *Recent advances in understanding assembly of the primary cilium* F1000 Research, In Press.
13. Kobayashi, T., W.Y. Tsang, J. Li, W. Lane, and B.D. Dynlacht, *Centriolar kinesin Kif24 interacts with CP110 to remodel microtubules and regulate ciliogenesis.* Cell, 2011. **145**(6): p. 914-25.
14. Huang, N., D. Zhang, F. Li, P. Chai, S. Wang, J. Teng, *et al.*, *M-Phase Phosphoprotein 9 regulates ciliogenesis by modulating CP110-CEP97 complex localization at the mother centriole.* Nat Commun, 2018. **9**(1): p. 4511.
15. Cajanek, L. and E.A. Nigg, *Cep164 triggers ciliogenesis by recruiting Tau tubulin kinase 2 to the mother centriole.* Proc Natl Acad Sci U S A, 2014. **111**(28): p. E2841-50.
16. Oda, T., S. Chiba, T. Nagai, and K. Mizuno, *Binding to Cep164, but not EB1, is essential for centriolar localization of TTBK2 and its function in ciliogenesis.* Genes Cells, 2014. **19**(12): p. 927-40.
17. Lo, C.H., I.H. Lin, T.T. Yang, Y.C. Huang, B.E. Tanos, P.C. Chou, *et al.*, *Phosphorylation of CEP83 by TTBK2 is necessary for cilia initiation.* J Cell Biol, 2019. **218**(10): p. 3489-3505.

18.   Graser, S., Y.D. Stierhof, S.B. Lavoie, O.S. Gassner, S. Lamla, M. Le Clech*, et al.*, *Cep164,*
a novel centriole appendage protein required for primary cilium formation. J Cell Biol,
2007. **179**(2): p. 321-30.

19.   Goetz, S.C., K.F. Liem, Jr., and K.V. Anderson, *The spinocerebellar ataxia-associated gene*
Tau tubulin kinase 2 controls the initiation of ciliogenesis. Cell, 2012. **151**(4): p. 847-58.

20.   Schmidt, K.N., S. Kuhns, A. Neuner, B. Hub, H. Zentgraf, and G. Pereira, *Cep164 mediates*
vesicular docking to the mother centriole during early steps of ciliogenesis. J Cell Biol,
2012. **199**(7): p. 1083-101.

21.   Tanos, B.E., H.J. Yang, R. Soni, W.J. Wang, F.P. Macaluso, J.M. Asara*, et al.*, *Centriole*
distal appendages promote membrane docking, leading to cilia initiation. Genes Dev, 2013.
27(2): p. 163-8.

22.   Ye, X., H. Zeng, G. Ning, J.F. Reiter, and A. Liu, *C2cd3 is critical for centriolar distal*
appendage assembly and ciliary vesicle docking in mammals. Proc Natl Acad Sci U S A,
2014. **111**(6): p. 2164-9.

23.   Kurtulmus, B., C. Yuan, J. Schuy, A. Neuner, S. Hata, G. Kalamakis*, et al.*, *LRRC45*
contributes to early steps of axoneme extension. J Cell Sci, 2018. **131**(18).

24.   Kanie, T., K.L. Abbott, N.A. Mooney, E.D. Plowey, J. Demeter, and P.K. Jackson, *The*
CEP19-RABL2 GTPase Complex Binds IFT-B to Initiate Intraflagellar Transport at the
Ciliary Base. Dev Cell, 2017. **42**(1): p. 22-36 e12.

25.   Nishijima, Y., Y. Hagiya, T. Kubo, R. Takei, Y. Katoh, and K. Nakayama, *RABL2 interacts*
with the intraflagellar transport-B complex and CEP19 and participates in ciliary assembly.
Mol Biol Cell, 2017. **28**(12): p. 1652-1666.

26.   Mojarad, B.A., G.D. Gupta, M. Hasegan, O. Goudiam, R. Basto, A.C. Gingras*, et al.*,
CEP19 cooperates with FOP and CEP350 to drive early steps in the ciliogenesis
programme. Open Biol, 2017. **7**(6).

27.   Hossain, D., Y. Javadi Esfehani, A. Das, and W.Y. Tsang, *Cep78 controls centrosome*
homeostasis by inhibiting EDD-DYRK2-DDB1 VprBP. EMBO Rep, 2017. **18**(4): p. 632-
644.

28.   Yoshida, S., K. Aoki, K. Fujiwara, T. Nakakura, A. Kawamura, K. Yamada*, et al.*, *The*
novel ciliogenesis regulator DYRK2 governs Hedgehog signaling during mouse
embryogenesis. Elife, 2020. **9**.

29.   Nikopoulos, K., P. Farinelli, B. Giangreco, C. Tsika, B. Royer-Bertrand, M.K. Mbefo*, et al.*,
Mutations in CEP78 Cause Cone-Rod Dystrophy and Hearing Loss Associated with
Primary-Cilia Defects. Am J Hum Genet, 2016. **99**(3): p. 770-776.

30.   Ascari, G., F. Peelman, P. Farinelli, T. Rosseel, N. Lambrechts, K.A. Wunderlich*, et al.*,
Functional characterization of the first missense variant in CEP78, a founder allele
associated with cone-rod dystrophy, hearing loss, and reduced male fertility. Hum Mutat,
2020. **41**(5): p. 998-1011.

31.   Namburi, P., R. Ratnapriya, S. Khateb, C.H. Lazar, Y. Kinarty, A. Obolensky*, et al.*, *Bi-*
allelic Truncating Mutations in CEP78, Encoding Centrosomal Protein 78, Cause Cone-
Rod Degeneration with Sensorineural Hearing Loss. Am J Hum Genet, 2016. **99**(5): p.
1222-1223.

32.   Fu, Q., M. Xu, X. Chen, X. Sheng, Z. Yuan, Y. Liu*, et al.*, *CEP78 is mutated in a distinct*
type of Usher syndrome. J Med Genet, 2017. **54**(3): p. 190-195.

33.   de Castro-Miro, M., R. Tonda, P. Escudero-Ferruz, R. Andres, A. Mayor-Lorenzo, J. Castro*,*
et al., *Novel Candidate Genes and a Wide Spectrum of Structural and Point Mutations*
Responsible for Inherited Retinal Dystrophies Revealed by Exome Sequencing. PLoS One,
2016. **11**(12): p. e0168966.

34. Azimzadeh, J., M.L. Wong, D.M. Downhour, A. Sanchez Alvarado, and W.F. Marshall,
Centrosome loss in the evolution of planarians. Science, 2012. **335**(6067): p. 461-3.

35. Brunk, K., M. Zhu, F. Barenz, A.S. Kratz, U. Haselmann-Weiss, C. Antony*, et al., Cep78 is*
a new centriolar protein involved in Plk4-induced centriole overduplication. J Cell Sci,
2016. **129**(14): p. 2713-8.

36. Knudsen, E.S. and J.Y. Wang, *Dual mechanisms for the inhibition of E2F binding to RB by*
cyclin-dependent kinase-mediated RB phosphorylation. Mol Cell Biol, 1997. **17**(10): p.
5771-5783.

37. Yan, X., R. Habedanck, and E.A. Nigg, *A complex of two centrosomal proteins, CAP350*
and FOP, cooperates with EB1 in microtubule anchoring. Mol Biol Cell, 2006. **17**(2): p.
634-644.

38. Wang, L., M. Failler, W. Fu, and B.D. Dynlacht, *A distal centriolar protein network*
controls organelle maturation and asymmetry. Nat Commun, 2018. **9**(1): p. 3938.

39. Kim, J., J.E. Lee, S. Heynen-Genel, E. Suyama, K. Ono, K. Lee*, et al., Functional genomic*
screen for modulators of ciliogenesis and cilium length. Nature, 2010. **464**(7291): p. 1048-
51.

40. Shearer, R.F., K.M. Frikstad, J. McKenna, R.A. McCloy, N. Deng, A. Burgess*, et al., The*
E3 ubiquitin ligase UBR5 regulates centriolar satellite stability and primary cilia. Mol Biol
Cell, 2018. **29**(13): p. 1542-1554.

41. Tsang, W.Y. and B.D. Dynlacht, *CP110 and its network of partners coordinately regulate*
cilia assembly. Cilia, 2013. **2**(1): p. 9.

42. Yadav, S.P., N.K. Sharma, C. Liu, L. Dong, T. Li, and A. Swaroop, *Centrosomal protein*
CP110 controls maturation of the mother centriole during cilia biogenesis. Development,
2016. **143**(9): p. 1491-501.

43. Walentek, P., I.K. Quigley, D.I. Sun, U.K. Sajjan, C. Kintner, and R.M. Harland, *Ciliary*
transcription factors and miRNAs precisely regulate Cp110 levels required for ciliary
adhesions and ciliogenesis. Elife, 2016. **5**.

44. Loukil, A., K. Tormanen, and C. Sutterlin, *The daughter centriole controls ciliogenesis by*
regulating Neurl-4 localization at the centrosome. J Cell Biol, 2017. **216**(5): p. 1287-1300.

45. Nagai, T., S. Mukoyama, H. Kagiwada, N. Goshima, and K. Mizuno, *Cullin-3-KCTD10-*
mediated CEP97 degradation promotes primary cilium formation. J Cell Sci, 2018. **131**(24).

46. Patzke, S., S. Redick, A. Warsame, C.A. Murga-Zamalloa, H. Khanna, S. Doxsey*, et al.,*
CSPP is a ciliary protein interacting with Nephrocystin 8 and required for cilia formation.
Molecular biology of the cell, 2010. **21**(15): p. 2555-67.

47. Frikstad, K.M., E. Molinari, M. Thoresen, S.A. Ramsbottom, F. Hughes, S.J.F. Letteboer*, et*
al., A CEP104-CSPP1 Complex Is Required for Formation of Primary Cilia Competent in
Hedgehog Signaling. Cell Rep, 2019. **28**(7): p. 1907-1922 e6.

48. Jiang, K., G. Toedt, S. Montenegro Gouveia, N.E. Davey, S. Hua, B. van der Vaart*, et al., A*
Proteome-wide screen for mammalian SxIP motif-containing microtubule plus-end tracking
proteins. Curr Biol, 2012. **22**(19): p. 1800-7.

49. Satish Tammana, T.V., D. Tammana, D.R. Diener, and J. Rosenbaum, *Centrosomal protein*
CEP104 (Chlamydomonas FAP256) moves to the ciliary tip during ciliary assembly. J Cell
Sci, 2013. **126**(Pt 21): p. 5018-29.

50. Eguether, T., M.A. Ermolaeva, Y. Zhao, M.C. Bonnet, A. Jain, M. Pasparakis*, et al., The*
deubiquitinating enzyme CYLD controls apical docking of basal bodies in ciliated epithelial
cells. Nat Commun, 2014. **5**: p. 4585.

51. Wang, L., K. Lee, R. Malonis, I. Sanchez, and B.D. Dynlacht, *Tethering of an E3 ligase by*
PCM1 regulates the abundance of centrosomal KIAA0586/Talpid3 and promotes
ciliogenesis. Elife, 2016. **5**.
52. Douanne, T., G. Andre-Gregoire, A. Thys, K. Trillet, J. Gavard, and N. Bidere, *CYLD*
Regulates Centriolar Satellites Proteostasis by Counteracting the E3 Ligase MIB1. Cell
Rep, 2019. **27**(6): p. 1657-1665 e4.
53. Yi, C., Z. Shen, A. Stemmer-Rachamimov, N. Dawany, S. Troutman, L.C. Showe*, et al.*,
The p130 isoform of angiomotin is required for Yap-mediated hepatic epithelial cell
proliferation and tumorigenesis. Sci Signal, 2013. **6**(291): p. ra77.
54. Hoppeler-Lebel, A., C. Celati, G. Bellett, M.M. Mogensen, L. Klein-Hitpass, M. Bornens*, et*
al., *Centrosomal CAP350 protein stabilises microtubules associated with the Golgi*
complex. J Cell Sci, 2007. **120**(Pt 18): p. 3299-308.
55. Belzile, J.P., G. Duisit, N. Rougeau, J. Mercier, A. Finzi, and E.A. Cohen, *HIV-1 Vpr-*
mediated G2 arrest involves the DDB1-CUL4AVPRBP E3 ubiquitin ligase. PLoS Pathog,
2007. **3**(7): p. e85.
56. Cox, J. and M. Mann, *MaxQuant enables high peptide identification rates, individualized*
p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol,
2008. **26**(12): p. 1367-72.
57. Schou, K.B., J.B. Mogensen, A. Aleliunaite, S.K. Morthorst, I.R. Veland, S.T. Christensen*,*
et al., *Kinesin-3 motor protein KIF13B localizes to centrosomes and primary cilia and*
regulates ciliary length and signaling. Cilia, 2014. **4**(Suppl 1): p. O3.
58. Verdier, P., S.K. Morthorst, and L.B. Pedersen, *Targeting of ASH Domain-Containing*
Proteins to the Centrosome. Methods Mol Biol, 2016. **1454**: p. 15-33.
59. Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nat Methods,
2012. **9**(4): p. 357-9.
60. Kim, D., G. Pertea, C. Trapnell, H. Pimentel, R. Kelley, and S.L. Salzberg, *TopHat2:*
accurate alignment of transcriptomes in the presence of insertions, deletions and gene
fusions. Genome Biol, 2013. **14**(4): p. R36.
61. Anders, S., P.T. Pyl, and W. Huber, *HTSeq--a Python framework to work with high-*
throughput sequencing data. Bioinformatics, 2015. **31**(2): p. 166-9.
62. Trapnell, C., B.A. Williams, G. Pertea, A. Mortazavi, G. Kwan, M.J. van Baren*, et al.*,
Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and
isoform switching during cell differentiation. Nat Biotechnol, 2010. **28**(5): p. 511-5.
63. Robinson, M.D., D.J. McCarthy, and G.K. Smyth, *edgeR: a Bioconductor package for*
differential expression analysis of digital gene expression data. Bioinformatics, 2010. **26**(1):
p. 139-40.
64. Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion*
for RNA-seq data with DESeq2. Genome Biol, 2014. **15**(12): p. 550.
966

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8