CEP78 functions downstream of CEP350 to control biogenesis of primary cilia by

negatively regulating CP110 levels

André Brás Gonçalves[1,6], Sarah Kirstine Hasselbalch[1,7], Beinta Biskopstø Joensen[1,7], Sebastian

Patzke[2], Pernille Martens[1], Signe Krogh Ohlsen[3], Mathieu Quinodoz[4,4.1,4.2], Konstantinos

Nikopoulos[5], Reem Suleiman[1], Magnus Per Damsø Jeppesen[1], Catja Weiss[1], Søren Tvorup Chris-

tensen[1], Carlo Rivolta[4,4.1,4.2], Jens S. Andersen[3], Pietro Farinelli[1*] and Lotte Bang Pedersen[1*]

[1]Department of Biology, Section for Cell Biology and Physiology, University of Copenhagen,

Universitetsparken 13, DK-2100 Copenhagen Ø, Denmark

[2]Department of Radiation Biology, Institute for Cancer Research, Norwegian Radium Hospital,

Oslo University Hospital, Oslo, Norway

[3]Department of Biochemistry and Molecular Biology, University of Southern Denmark, Campusvej

55, 5230 Odense M, Denmark

[4]Institute of Molecular and Clinical Ophthalmology Basel (IOB), Basel, Switzerland

[4.1]Department of Ophthalmology, University of Basel, Basel, Switzerland

[4.2]Department of Genetics and Genome Biology, University of Leicester, Leicester, United King-

dom

[5]Department of Computational Biology, University of Lausanne, Switzerland
25

26 [6] Current address: Instituto de Medicina Molecular João Lobo Antunes, Faculdade de Medicina,

27 Universidade de Lisboa, Lisbon, Portugal

30 [7]These authors contributed equally to this study

32 *For correspondance: Pietro.Farinelli@twelve.bio (PF); lbpedersen@bio.ku.dk (LBP)

Abstract

CEP78 is a centrosomal protein implicated in ciliogenesis and ciliary length control, and mutations in the *CEP78* gene cause retinal cone-rod dystrophy associated with hearing loss. However, the mechanism by which CEP78 affects cilia formation is unknown. Based on a recently discovered disease-causing *CEP78* p.L150S mutation, we identified the disease-relevant interactome of CEP78. We confirmed that CEP78 interacts with the EDD1-DYRK2-DDB1^{VPRBP} E3 ubiquitin ligase complex, which is involved in CP110 ubiquitination and degradation, and identified a novel interaction between CEP78 and CEP350 that is weakened by the CEP78^{L150S} mutation. We show that CEP350 promotes centrosomal recruitment and stability of CEP78, which in turn leads to centrosomal recruitment of EDD1. Consistently, cells lacking CEP78 display significantly increased cellular and centrosomal levels of CP110, and depletion of CP110 in CEP78-deficient cells restored ciliation frequency to normal. We propose that CEP78 functions downstream of CEP350 to promote ciliogenesis by negatively regulating CP110 levels via an EDD1-dependent mechanism.

Introduction

Primary cilia are antenna-like sensory organelles that play pivotal roles in coordinating various signaling pathways important for human development and tissue homeostasis [1]. They comprise a microtubule-based axoneme core, which extends directly from the mother centriole-derived basal body and is surrounded by a bilayer membrane enriched for specific receptors, ion channels and lipids that endow the organelle with unique signaling properties [2]. Not surprisingly, mutations in genes that affect assembly, structure or function of cilia are causative for a growing number of pleiotropic diseases and syndromes called ciliopathies, which include cone-rod dystrophy in the retina and hearing loss (CRDHL; MIM# 617236) amongst others [3]. Ciliopathy genes include those coding for components of the centrosome, which contains the daughter and mother centriole and gives rise to the ciliary basal body. The mother centriole is distinguished from the daughter centriole by distal and subdistal appendages, which play critical roles in vesicle docking at the onset of ciliogenesis and in microtubule anchoring, respectively. In addition, the centrosome contains pericentriolar material and is associated with centriolar satellites that affect cilia biogenesis and function in various ways [4].

Assembly of primary cilia is a complex, multistep process that is tightly coordinated with the cell cycle. In actively-proliferating cells, centriolar coiled coil protein 110 (CP110; also known as CCP110) and centrosomal protein of 97 kDa (CEP97) cap the distal ends of both mother and daughter centrioles and suppress ciliogenesis. Furthermore, overexpression of CP110 in growth-arrested cells prevents ciliogenesis [5]. CP110 also regulates centriole duplication and length control during S phase and interacts with key regulators of centriole duplication, including PLK4 [6-9]. As cells enter G1/G0, ciliogenesis begins with recruitment of pre-ciliary vesicles to the distal end of the mother centriole. The vesicles subsequently fuse to form a larger ciliary vesicle underneath which the ciliary transition zone and axoneme are assembled. The axoneme is further extended by intraflagellar transport (IFT), and the ciliary vesicle expands and matures to form the ciliary membrane and a surrounding sheath that fuses with the plasma membrane upon completion of ciliogenesis [10-12].

Initiation of transition zone formation and axoneme extension during early stages of ciliogenesis requires removal of the CEP97-CP110 complex from the distal end of the mother centriole [5]. This process relies on M-Phase Phosphoprotein 9 (MPP9), which interacts directly with CEP97 and cooperates with kinesin KIF24 to recruit the CEP97-CP110 complex to the distal centriole end [13, 14]. During ciliogenesis, phosphorylation of MPP9 by Tau tubulin kinase 2 (TTBK2)

leads to degradation of MPP9 by the ubiquitin proteasome system (UPS), which results in destabili-

zation of the CEP97-CP110 complex causing its removal from the distal end of the mother centriole

[14]. TTBK2 is recruited to the mother centriole distal appendages by CEP164 [15, 16] where it

also phosphorylates CEP83 to promote CP110 removal [17]. Consequently, depletion of CEP164,

TTBK2, CEP83 or other centriole distal appendage proteins that regulate their localization and/or

function impairs ciliogenesis [18-23]. Mother centriole recruitment of TTBK2 and removal of the

distal CEP97-CP110 cap additionally requires the subdistal CEP350-FOP-CEP19 complex, which

furthermore interacts with RABL2 and IFT-B complex components to promote their axonemal en-

try [24-26].

89   Despite recent advances, the precise mechanisms by which CP110 regulates ciliogenesis and

is removed from the mother centriole at the onset of ciliogenesis remain incompletely understood.

For example, a recent study implicated the homologous to the E6AP carboxyl terminus (HECT)

type EDD1-DYRK2-DDB1^{VPRBP} E3 ligase complex in ubiquitination and degradation of CP110 via

a mechanism involving direct interaction of viral protein R binding protein (VPRBP; also known as

DCAF1) and centrosomal protein of 78 kDa (CEP78) [27]. Specifically, the authors reported that

CEP78 suppresses CP110 ubiquitination by EDD1 (also known as UBR5 and EDD) and it was pro-

posed that CP110 is phosphorylated by DYRK2 and thereby recognized and brought close to EDD1

by VPRBP. EDD1 then transfers ubiquitin to CP110 leading to its degradation. When CEP78 binds

VPRBP it induces a conformational change in the complex thereby preventing CP110

ubiquitination. Further, they demonstrated that depletion of CEP78 promoted centriole elongation,

whereas *CEP78* over-expression inhibited primary cilia formation in hTERT-immortalized retinal

pigment epithelial (RPE1) cells [27]. On the other hand, knockout (KO) of *Dyrk2* in the mouse was

reported to result in elongation of primary cilia, but centrosomal CP110 levels appeared unaffected

in *Dyrk2* mouse KO cells [28]. Therefore, it remains uncertain how CEP78 and the EDD1-DYRK2-

DDB1^{VPRBP} complex affect CP110 homeostasis to control ciliogenesis.

CEP78 is composed of 16 exons and encodes a protein comprising 722 amino acids that

possesses five consecutive leucine rich repeats (LRR) at the N-terminus, and a coiled-coil (CC)

domain at the C-terminus [29, 30]. Four independent studies have reported eight different mutations

in *CEP78* in patients with CRDHL [29-32], whereas one study identified a homozygous *CEP78*-

truncating variant in a family with non-syndromic retinitis pigmentosa (MIM# 268003; [33]), an-

other form of retinal degeneration. Of these studies, two have investigated the functional conse-

quences of human *CEP78* mutations at the cellular level. In one study, whole-exome sequencing

(WES) identified biallelic mutations in *CEP78* in two unrelated families from Greece and Sweden,

respectively [29]. The Greek subject had a homozygous base substitution at the splice donor site in

intron 3 of *CEP78* (NM_032171.2:c.499+1G>T). Two subjects from a Swedish family carried het-

erozygous mutations, one base substitution in intron 3 (NM_032171.2:c.499+5G>A) and a single

nucleotide deletion in exon 5 (NM_032171.2:c.633del; p.Trp212GlyfsTer18) causing a frameshift.

These *CEP78* mutations lead to exon skipping and premature stop codons accompanied by almost

undetectable levels of CEP78 protein in human skin fibroblasts (HSFs) of affected individuals. Fur-

thermore, it was found that HSFs from these patients have significantly longer primary cilia as

compared to control cells [29]. More recently, a missense mutation in *CEP78*

(NM_032171.2:c.449T>C; p.Leu150Ser) was identified in three unrelated families from Belgium

and Germany diagnosed with CRDHL [30]. In the two Belgian families, affected individuals were

homozygous for the p.Leu150Ser variant, whereas affected individuals from the German family

displayed compound heterozygosity for this variant and a novel splice site variant,

NM_032171.2:c.1462-1G>T. In all cases, HSFs from patients harboring the p.Leu150Ser mutation

(hereafter: L150S) displayed decreased cellular levels of CEP78 and significantly elongated cilia

compared to control HSFs [30], as seen in patient-derived HSFs with CEP78 truncating mutations

[29]. In addition, other ciliopathy features were reported in some of the affected individuals with the

CEP78[L150S] mutation, including obesity, respiratory problems, diabetes 2 and infertility [30]. In

summary, available data derived from human patients indicates that depletion of CEP78 leads to

elongation of primary cilia at the cellular level, which manifests in ciliopathy phenotypes at the

organism level. However, the mechanism by which CEP78 regulates ciliary length is not known.

In addition to the patient studies described above, some studies have addressed CEP78 func-

tion in different cell culture models. First, a study showed that siRNA-mediated depletion of CEP78

in RPE1 cells reduces the frequency of cells forming primary cilia, possibly due to centriolar an-

choring defects, but the length of the residual cilia was not assessed. Similarly, depletion of CEP78

in *Planarians* was shown to impair motile cilia formation due to defective docking of centrioles to

the cell surface [34]. Another study identified CEP78 interaction with PLK4 and implicated CEP78

in PLK4-mediated centriole over duplication, whereas possible roles for CEP78 in relation to cilia

were not addressed [35]. Finally, as mentioned above, a study reported that CEP78 directly interacts

with VPRBP of the EDD1-DYRK2-DDB1[VPRBP] E3 ligase complex to suppress CP110

ubiquitination and thereby stabilize it [27]. How such CEP78-mediated stabilization of CP110

might lead to the long cilia phenotype observed in patient fibroblasts lacking CEP78 [29, 30] and/or

the reduced ciliation frequencies seen in CEP78-depleted *Planarians* and RPE1 cells [34] is un-

clear.

Here were show that in RPE1 cells and patient-derived HSFs loss of CEP78 leads to reduced

ciliation frequency as well as increased length of the cilia that do form. Further, we find that a mu-

tant line expressing a partially functional, truncated version of CEP78 displays reduced ciliation

frequency but normal length of residual cilia. By taking advantage of the recently identified disease-

causing CEP78^{L150S} mutation [30], we used a quantitative mass spectrometry-based approach to

identify a disease-relevant interactome of CEP78. We confirmed that CEP78 interacts with the

EDD1-DYRK2-DDB1^{VPRBP} complex implicated in CP110 ubiquitination and degradation [27], and

furthermore identified a novel interaction between CEP78 and the N-terminal region of CEP350.

The interaction of CEP78 with both VPRBP and CEP350, as well as centrosomal recruitment of

CEP78, are dramatically reduced by the CEP78^{L150S} mutation. Lack of centrosomal recruitment of

CEP78^{L150S} is likely due to impaired interaction with CEP350 since centrosomal and cellular levels

of CEP78 were significantly decreased in *CEP350* KO cells. Conversely, cells lacking CEP78 dis-

played significantly decreased centrosomal levels of EDD1 and unaltered or slightly increased

centrosomal levels of VPRBP and CEP350. In addition, CEP78-deficient cells showed significantly

increased cellular and centrosomal levels of CP110, presumably owing to the reduced centrosomal

levels of EDD1 observed in these cells. Depletion of CP110 in CEP78-deficient cells restored

ciliation frequency, but not the increased length of remaining cilia, to normal. Collectively our re-

sults suggest that CEP78 functions downstream of CEP350 to promote cilia biogenesis by negative-

ly regulating CP110 levels via an EDD1-dependent mechanism, but suppresses ciliary elongation

independently of CP110.

Results

Depletion of CEP78 leads to fewer and longer primary cilia in cultured cells

Previous studies showed that patient-derived fibroblasts lacking CEP78 display significantly elon-

gated primary cilia compared to control cells [29, 30] whereas depletion of CEP78 in *Planarians*

and RPE1 cells was shown to significantly reduce cilia numbers [34]. To reconcile these seemingly

contradictory findings we analyzed cilia numbers and length in serum-deprived wild type (WT)

RPE1 cells and four different *CEP78* KO clones generated by CRISPR/Cas9 methodology. We first

confirmed by western blot analysis with CEP78-specific antibody that endogenous CEP78 is lack-

ing in three of the mutant clones, designated #2, #52 and #73, whereas clone #44 expresses a short-

er version of CEP78 (Figure 1-figure supplement 1A). Sequencing of clone #44 indicated that it

contains an insertion of the px459-Cas9 plasmid in exon 1 of *CEP78*, suggestion that a shorter ver-

sion of CEP78 lacking the extreme N-terminus is expressed in this strain, possibly from an alterna-

tive promotor. Immunofluorescence microscopy (IFM) analysis with antibodies against ciliary

(ARL13B) and centrosomal (CEP350) markers showed that the frequency of ciliated cells is signif-

icantly reduced in all four *CEP78* KO clones compared to WT cells (Figure 1A, B; Figure 1-Figure

supplement 1B). In addition, the length of the remaining cilia was significantly increased in the

three *CEP78* KO clones that completely lack CEP78, but identical to the average WT cilia size in

clone #44 (Figure 1A, C; Figure 1-figure supplement 1C). The latter result indicates that the trun-

cated CEP78 protein expressed in clone #44 functions normally with respect to ciliary length con-

trol, in turn suggesting that CEP78 may affect ciliogenesis and ciliary length by distinct mecha-

nisms. For the rest of this manuscript RPE1 *CEP78* KO cells refer to clone #73 unless otherwise

indicated.

In agreement with our observations in RPE1 cells, similar results were obtained using previ-

ously described patient-derived *CEP78*-deficient or control HSFs (Figure 1D-F) [29, 30]. The low

ciliation frequencies of serum-deprived *CEP78* mutant cells were not secondary to cell cycle de-

fects, as judged by western blot analysis with antibody against retinoblastoma protein phosphory-

lated at S807/811 (P-Rb; Figure 1G-L). However, in serum-fed cells, P-Rb levels were significantly

higher in *CEP78* mutant cells compared to controls (Figure 1G-L), indicating that mutant cells may

progress slower through S-phase [36]. This is in line with previous reports indicating that CEP78

protein levels are upregulated in late S-G2 phase suggesting a role for CEP78 at this cell cycle

stage, e.g. during centriole duplication [27, 35]. We conclude that CEP78 deficient RPE1 and HSF

cells display reduced ciliation frequencies as well as increased length of remaining cilia, thus recon-

ciling previous observations [29, 30, 34].

Analysis of RPE1 WT and *CEP78* KO cells expressing FLAG- or mNG-tagged CEP78 or

CEP78^{L150S} fusions

To confirm the above results, we first set out to perform a rescue experiment by expressing FLAG-

CEP78 in WT and *CEP78* KO RPE1 cells followed by serum-deprivation and IFM analysis using

FLAG and ARL13B antibodies. In parallel, we performed similar experiments with FLAG-

CEP78^{L150S}. We first analyzed the localization and expression levels of the two FLAG fusions in

transiently transfected serum-deprived RPE1 cells. In 4% paraformaldehyde-fixed cells we ob-

served that the ARL13B antibody labeled the cilium itself as well as the basal body, but not the

daughter centriole. The basal body pool of ARL13B likely corresponds to that present in the mother

centriole-associated vesicle of mitotic centrosomes [37]. The transiently expressed FLAG-CEP78

fusion protein localized to both centrioles at the base of cilia as expected [27, 29, 35], whereas

centriolar localization of FLAG-CEP78^{L150S} was severely compromised although not completely

abolished (Figure 2A, B). The latter result is in line with previous reports showing that the LRR

region in the N-terminus of CEP78, which encompasses residue L150, is important for its localiza-

tion to centrioles [27, 35]. Both FLAG fusions were expressed at similar levels in the cells (Figure

2-figure supplement 1); this was somewhat surprising given that endogenous CEP78^{L150S} is largely

undetectable by western blot analysis of patient-derived HSFs suggesting its stability is compro-

mised [30]. Therefore, we tested if the L150S mutation might affect binding of the CEP78 antibody

used in [30] to CEP78. However, FLAG immunoprecipitation (IP) and western blot analysis of

293T cells expressing FLAG-CEP78 or FLAG-CEP78^{L150S} indicated that the CEP78 antibody binds

equally well to WT CEP78 and the CEP78^{L150S} mutant protein. This analysis also revealed that en-

dogenous CEP78 co-IPs with FLAG-CEP78 indicating that CEP78 can form

homodimers/oligomers (Figure 2-figure supplement 2). These results indicate that the L150S muta-

tion compromises the recruitment of CEP78 to the centrosome, but not its short-term stability, at

least in RPE1 cells. We cannot rule out that the N-terminal FLAG tag may stabilize FLAG-

CEP78^{L150S} and prevent it from being degraded.

Next, we tested if FLAG-CEP78 or FLAG-CEP78^{L150S} could rescue the ciliary phenotypes

of *CEP78* KO cells. However, upon transient expression of the FLAG-CEP78 fusions we could not

rescue the cilia frequency and length phenotype of the *CEP78* KO cells, and cilia frequency and

length were also affected in WT control cells transiently expressing these fusions (data not shown).

We therefore generated cell lines stably expressing mNeonGreen (mNG)-tagged CEP78 or

CEP78^{L150S} at ca. 3-4 times the endogenous CEP78 level by viral transduction of mNG-CEP78 con-

structs into WT and *CEP78* KO RPE1 cells, respectively (Figure 2-figure supplement 3A). Micro-

scopic examination of these lines indicated that mNG-CEP78 was concentrated at the centrosome as

expected, whereas mNG-CEP78^{L150S} was diffusely located in the cytosol (Figure 2-figure supple-

ment 3B), in agreement with results for transiently expressed FLAG-CEP78/CEP78^{L150S} fusions

(Figure 2A, B). Importantly, *CEP78* KO cells stably expressing mNG-CEP78 had higher ciliation

frequency than untransfected *CEP78* KO cells or WT cells expressing mNG-CEP78 or *CEP78* KO

cells expressing mNG-CEP78^{L150S} (Figure 2C). However, their ciliation frequency was still lower

than that of untransfected WT cells, which we attribute to the fact that mild overexpression of

mNG-CEP78 seems to inhibit ciliogenesis, as seen for WT cells stably expressing mNG-CEP78

(Figure 2C). Furthermore, WT or *CEP78* KO cells stably expressing mNG-CEP78 had significantly

shorter cilia than untransfected WT and *CEP78* KO cells, or *CEP78* KO cells expressing mNG-

CEP78^{L150S} (Figure 2D). Together these results suggest that mNG-CEP78 is able to at least partially

rescue the cilia frequency and length phenotype of CEP78 KO cells whereas mNG-CEP78^{L150S} is

not.). We conclude that the observed ciliary frequency and length phenotypes of the *CEP78* KO

cells are caused specifically by CEP78 loss, and that the L150S mutations impairs CEP78 centro-

some localization and function.

CEP78 interacts with the EDD1-DYRK2-DDB1^{VPRBP} complex and CEP350

To explore how CEP78 is recruited to centrioles to regulate ciliary frequency and length we first

used Stable Isotope Labeling by Amino acids in Cell culture (SILAC)-based quantitative mass spec-

trometry (MS) to identify the differential interactomes of FLAG-CEP78 and FLAG-CEP78^{L150S}

expressed in 293T cells (see Materials and methods for details). We identified interaction between

CEP78 and several components of the EDD1-DYRK2-DDB1^{VPRBP} complex (Figure 3A), in agree-

ment with previous work showing direct binding of CEP78 to VPRBP within this complex [27].

Interestingly, the MS results indicated that the CEP78^{L150S} mutation dramatically reduces this inter-

action (Figure 3A), which we confirmed by IP and western blot analysis of 293T cells co-

expressing Myc-VPRBP and FLAG-CEP78 or FLAG-CEP78^{L150S} (Figure 3B). In addition, our

interactome analysis identified CEP350 as potential novel CEP78 binding partner (Figure 3A). Us-

ing IP and western blot analysis in 293T cells co-expressing FLAG-CEP78 or FLAG-CEP78^{L150S}

with a Myc-tagged N-terminal or C-terminal CEP350 truncation we found that CEP78 binds to the

N-terminal region of CEP350 (residues 1-983; Figure 3C, E); reciprocal co-IP with Myc-tagged

CEP350 constructs confirmed this result (Figure 3-figure supplement 1A). Furthermore, we found

that the interaction between CEP78 and CEP350 N-terminus is reduced by the CEP78^{L150S} mutation

(Figure 3C). Using a similar approach, we could not detect interaction between CEP78 and endoge-

nous FOP (Figure 3C), whereas under similar conditions endogenous FOP was readily co-IPed with

the C-terminus of CEP350 (Figure 3-figure supplement 1A), in agreement with a previous report

[38]. We also mapped the CEP350 interaction site in CEP78 and found that CEP350 primarily

binds to the C-terminal region of CEP78 comprising residues 395-722 (Figure 3D, E), although a

weak interaction with a fragment comprising the entire LRR region (residues 2-403) was also ob-

served. In contrast, truncation of the latter fragment into two smaller fragments abolished binding to
the CEP350 N-terminus (Figure 3D, E). Binding of CEP78 to VPRBP similarly requires the entire
LRR region of CEP78 [27] and we therefore wondered whether interaction between CEP350 and
CEP78 might be mediated by VPRBP. However, Myc IP experiments and western blot analysis
failed to reveal physical interaction between VPRBP and Myc-CEP350 N-terminus (Figure 3, fig-
ure supplement 1A). Moreover, GFP IP of cells co-expressing GFP-CEP78 and Myc-CEP350 N- or
C-terminal fragments showed that GFP-CEP78 can bind to Myc-CEP350 N-terminus and endoge-
nous VPRBP at the same time, indicating that CEP350 does not compete with VPRBP for binding
to CEP78 (Figure 3, figure supplement 1B). We conclude that CEP78 binds independently to
VPRBP and the N-terminus of CEP350 and that both of these interactions are reduced by the
CEP78^{L150S} mutation.

CEP350 regulates the stability and centrosomal recruitment of CEP78
In non-ciliated HeLa and U2OS cells endogenous CEP78 was reported to localize to both centrioles
during all phases of the cell cycle and appeared to concentrate at the centriole wall [35]. In ciliated
RPE1 cells CEP78 was shown to localize to the distal end of both centrioles, with a preference for
the mother centriole/basal body [27]. Similarly, CEP350 was reported to localize to the distal end of
the mother centriole/basal body, near the subdistal appendages, in ciliated RPE1 cells [24, 26, 39].
In agreement with these reports and with our interactome and IP results (Figure 3), 3D Structured
Illumination Microscopy (SIM) showed partial co-localization of mNG-CEP78 with CEP350 at the
distal end of the centrioles in RPE1 cells, adjacent to the distal appendage marker CEP164 and the
centriole capping complex protein CP110 (Figure 4A). Superimposing fluorescence images on elec-
tron micrographs of the mother centriole [40] indicates that mNG-CEP78 localizes to a ring-
structure of centriole-sized diameter at the distal half of the centriole, encompassed by a wider un-
closed-ring structure of CEP350. mNG-CEP78 but not CEP350 depicted decreased levels at the
daughter centriole as compared to the mother. Furthermore, using a previously characterized RPE1
CEP350 KO cell line [24] that displays significantly reduced centrosomal levels of CEP350 (Figure
4-figure supplement 1), we found that depletion of CEP350 not only reduces localization of CEP78
to the centrosome, but also its total cellular level (Figure 4B-G). Similarly, we found significantly
reduced levels of CEP78 in *FOP* KO cells (Figure 4 B, C), in line with the decreased centrosomal
levels of CEP350 observed in these cells [24]. These results indicate that CEP350 promotes the
recruitment of CEP78 to the mother centriole/basal body as well as its overall stability.

CEP78 mutant cells display reduced centrosomal levels of EDD1

CEP78 mutant cells display reduced centrosomal levels of EDD1

Next, we asked if CEP78 might affect the centrosomal recruitment of components of the EDD1-

DYRK2-DDB1^{VPRBP} complex and/or CEP350, and analyzed the cellular or centrosomal levels of

relevant complex components in RPE1 WT and *CEP78* KO cells. The results showed that serum-

deprived, ciliated *CEP78* KO cells display significantly increased centrosomal levels of CEP350

(Figure 5A, B) as compared to controls. However, in serum-deprived, non -ciliated cells, the

centrosomal levels of CEP350 were not significantly different in *CEP78* KO cells compared to WT

controls (Figure 5C, D). Similar results were obtained for CEP350 in control and CEP78 deficient

patient HSFs (Figure 5E-H). We also analyzed the cellular levels of VPRBP by western blotting

and the centrosomal levels of VPRBP and EDD1 by IFM in RPE1 WT and *CEP78* KO cells. In

agreement with a previous study [27] we found that cellular and centrosomal levels of VPRBP were

not significantly reduced in serum-fed or serum-deprived *CEP78* KO cells compared to controls

(Figure 6A-C; Figure 6-figure supplement 1). Thus, CEP78 seems to be dispensable for recruitment

of VPRBP and CEP350 to the centrosome. In contrast, centrosomal levels of EDD1 were signifi-

cantly reduced in RPE1 *CEP78* KO cells compared to WT controls, both in ciliated and non-ciliated

cells (Figure 6D-G). Thus, CEP78 is required for efficient recruitment of EDD1 to the centrosome.

To corroborate this result, we measured the centrosomal levels of EDD1 in RPE1 *CEP350* KO cells

and found that EDD1 levels were also significantly reduced when compared to the controls (Figure

6-figure supplement 2). Together our data indicate that CEP350, by recruiting CEP78 to the centro-

some, also promotes the centrosomal recruitment of EDD1.

Increased cellular and centrosomal levels of CP110 in CEP78 deficient cells

Since EDD1 is an E3 ubiquitin-protein ligase that negatively regulates CP110 levels [27], and

whose depletion impairs ciliogenesis [41, 42], we asked if the loss of CEP78 would lead to altered

levels of CP110 at the centrosome. Indeed, western blot and IFM analysis of RPE1 WT and *CEP78*

KO cells revealed that total cellular and centrosomal CP110 levels are significantly increased in

CEP78 KO cells compared to WT controls, both under serum-deprived (Figure 7A-D, F, G; Figure

7-figure supplement 1A, B) and serum-fed conditions (Figure 7-figure supplement 2A-D), whereas

CP110 levels appeared normal upon stable expression of mNG-CEP78 in the *CEP78* KO back-

ground (Figure 7-figure supplement 1C). We also analyzed CP110 levels specifically at the mother

centriole of ciliated RPE1 WT and *CEP78* KO cells and found no significant differences between

them (Figure 7E). Similar results were obtained using *CEP78* mutant and control HSF cells (Figure

7G-H). Since a previous study reported that siRNA-mediated depletion of CEP78 in RPE1 and

HeLa cells reduces the cellular levels of CP110 [27], a result contradicting our own observations

(Figure 7; Figure 7-figure supplement 1 and 2), we tested if the elevated CP110 levels seen in our

CEP78 mutant cells might be due to compensatory upregulation of CP110 mRNA expression.

However, RNA-Seq analysis indicated no significant changes in CP110 mRNA levels in patient-

derived, *CEP78* deficient HSF cultures compared to controls; mRNA levels of CEP350, VPRBP

and DDB1 in *CEP78* deficient HSF cultures were also similar to those of controls (Figure 7-figure

supplement 3). Furthermore, a cycloheximide chase experiment indicated that CP110 is more stable

upon serum deprivation of *CEP78* KO cells compared to WT controls (Figure 7-figure supplement

4). Taken together, we conclude that loss of CEP78 leads to elevated cellular and centrosomal lev-

els of CP110 and that this is likely due to increased stability of CP110. Furthermore, our results

indicate that in ciliated *CEP78* KO cells, the amount of CP110 present specifically at the mother

centriole is not significantly higher than in control cells.

Partial depletion of CP110 normalizes ciliation frequency of *CEP78* KO cells

CP110 is a key negative regulator of ciliogenesis [5, 43] that was also shown to be required for

ciliogenesis initiation as well as ciliary length control [44, 45]. The above results therefore prompt-

ed us to test if the reduced ciliation frequency and/or increased length of remaining cilia observed in

CEP78 deficient cells could result from elevated centrosomal CP110 levels in these cells. When we

used siRNA to partially deplete cellular and centrosomal CP110 from RPE1 WT and *CEP78* KO

cells and subjected the cells to serum deprivation (Figure 8A, B; Figure 8-figure supplement 1), we

found that the *CEP78* KO cells restored their ciliation frequency to WT levels, whereas partial

CP110 depletion had little effect on cilia numbers in WT cells grown under similar conditions (Fig-

ure 8C, D). In contrast, depletion of CP110 from *CEP78* KO cells did not rescue the ciliary length

defect of residual cilia; these cilia were still significantly longer than those of WT controls (Figure

8C, E). Thus, partial depletion of CP110 from *CEP78* KO cells can rescue their reduced ciliation

frequency phenotype but not the increased length of remaining cilia. This result is in line with our

observation that a *CEP78* KO clone expressing a truncated version of CEP78 (clone #44) displays

significantly reduced ciliation frequency, but normal length of the cilia that do form (Figure 1-

figure supplement 1), implying that CEP78 regulates ciliogenesis and ciliary length by separate

mechanisms.

Discussion

Several studies have shown that loss-of-function mutations in *CEP78* are causative of ciliopathy characterized by CRDHL [29-32], and that cells lacking CEP78 display reduced ciliation frequency [34] or abnormally long cilia [29, 30]. However, the molecular mechanisms by which CEP78 regulates cilium biogenesis and length were so far unclear. In this study, we found that loss of CEP78 leads to decreased ciliation frequency as well as increased length of the residual cilia that do form, both in RPE1 cells and patient-derived HSFs. Interestingly, we found that both ciliated and non-ciliated CEP78 deficient cells fail to efficiently recruit EDD1 to the centrosome, whereas cellular and overall centrosomal levels of CP110 are dramatically increased. However, in the subpopulation of *CEP78* KO cells that are ciliated, CP110 levels specifically at the mother centriole were similar to those of ciliated control cells, implying that the elevated centrosomal CP110 level observed in these cells is caused by a pool of CP110 located at the daughter centriole and/or pericentriolar area. This result also indicates that the abnormally long cilia phenotype seen in the *CEP78* KO cells is not caused by increased CP110 levels at the mother centriole. Supportively, siRNA-mediated depletion of CP110 could restore cilia frequency, but not length, to normal in the *CEP78* KO cells. Furthermore, a *CEP78* mutant line, clone #44, expressing a shorter version of CEP78 likely lacking the N-terminus displayed reduced ciliation frequency but normal length of residual cilia, substantiating that CEP78 regulates ciliogenesis and cilia length by distinct mechanisms. Specifically, our results indicate that CEP78 promotes ciliogenesis by negatively regulating CP110 levels at the mother centriole via activation of VprBP, but negatively regulates ciliary lengthening independently of CP110 (Figure 8F).

The removal of CP110 from the distal end of the mother centriole constitutes a key step in the initiation of ciliogenesis and is regulated by a growing number of proteins that include TTBK2 [19] and components of the UPS system such as Neurl-4 [46] and the cullin-3 (CUL3)–RBX1–KCTD10 complex [47]. The EDD1-DYRK2-DDB1^{VPRBP} complex was similarly shown to mediate degradation of CP110 and to interact with CEP78 [27], but the precise consequences of these activities for ciliogenesis were unclear. Our work strongly suggests that CEP78 functions together with the EDD1-DYRK2-DDB1^{VPRBP} complex to negatively regulate centrosomal CP110 levels at the onset of ciliogenesis, leading to its removal from the distal end of the mother centriole (Figure 8F). This is in contrast to a previously proposed model, which suggested that CEP78 inhibits degradation of CP110 by the EDD1-DYRK2-DDB1^{VPRBP} complex [27]. The reason for this discrepancy is

unclear, but may be due to different experimental approaches used in our study and the paper by

Hossain et al. [27]. In this latter study, siRNA was used to partially deplete CEP78 from cultured

cells, whereas we performed our experiments on cells with endogenous CEP78 mutations. CP110

removal from the distal end of the mother centriole occurs concomitantly with docking of the cen-

triole to vesicles or the plasma membrane at the onset of ciliogenesis [12]. Therefore, a role for

CEP78 in removal of CP110 from the centrosome is compatible with ultrastructural analyses in

Planarians, which indicated that basal bodies fail to dock to the plasma membrane in CEP78

RNAi-depleted animals [34]. Although not addressed in detail in the current study, the elevated

centrosomal CP110 levels observed in serum-fed CEP78 deficient cells are also compatible with

previous work implicating CEP78 in regulation of PLK4-mediated centriole duplication [35].

We also uncovered a new physical interaction between CEP78 and the N-terminal region of

CEP350, and found that CEP350 is important for recruitment of CEP78 to the centrosome as well

as for its overall stability. The reduced interaction of the CEP78^{L150S} mutant protein with CEP350

might explain its lack of detection by western blot analysis of patient HSFs [30], since reduced in-

teraction with CEP350 is likely to decrease the long-term stability of CEP78^{L150S}. Our IP and IFM

results furthermore suggests that CEP350 controls CP110 removal from the mother centriole not

only via recruitment of TTBK2 [24], but also via CEP78-dependent recruitment of EDD1. Indeed,

we observed reduced centrosomal levels of EDD1 in *CEP350* KO cells, consistent with this idea. It

remains to be determined whether these two pathways act in sequence or in parallel to control

centrosomal CP110 levels. Furthermore, as the recruitment of CEP350 to centrioles itself depends

on FOP [24, 26, 38] and on the distal centriole protein C2CD3 [14], loss of these proteins may simi-

larly affect the centrosomal recruitment of CEP78 and EDD1, and thereby negatively regulate

CP110 levels via this pathway. It will be important to investigate these possibilities in future stud-

ies.

While increased CP110 levels can account for the decreased ciliation frequency observed in

serum-deprived CEP78 mutant cells, it remains unclear why some of these cells form abnormally

long cilia. Our analysis indicated that the ciliated mutant cells are able to locally displace or degrade

CP110 at the mother centriole, but the underlying mechanism remains to be determined. Of note,

the *CEP350* KO cells used in our study also seem to display a similar dual phenotype, since approx-

imately 30% of these cells form cilia despite an inability to recruit TTBK2 and remove CP110 from

the mother centriole [24]. Moreover, a recent genome-wide siRNA knockdown analysis showed

that depletion of DDB1 caused an elongated cilia phenotype whereas depletion of EDD1 (UBR5)

lead to fewer but normal length cilia [48], even though DDB1 and EDD1 are part of the same E3

ubiquitin ligase complex [49]. Thus, a slight imbalance in the regulation of this complex might de-

termine whether or not cilia are formed and how long they are. It should also be noted that some

studies have reported dual roles for CP110 during ciliogenesis, i.e. CP110 may not only function to

inhibit ciliogenesis but may also promote ciliary extension in some contexts [44, 45]. Although the

mechanism responsible for the elongated cilia phenotype seen in some of the CEP78 mutant cells

remains unclear, one attractive candidate to be involved is CEP350, which we found to be present at

abnormally high levels at the centrosome of ciliated *CEP78* KO cells, specifically. Consistent with

this idea, we have observed that the CEP78 truncation mutant (clone #44), which expresses fewer,

but normal length cilia, displays normal or slightly reduced levels of CEP350 at the centrosome in

ciliated as well as in non-ciliated cells (data not shown). Alternatively, one or more substrates of

EDD1 other than CP110 could also be affected in the *CEP78* mutant cells thereby affecting ciliary

length control. For example, EDD1 was also shown to mediate ubiquitination of CSPP-L in turn

promoting its recruitment to centriolar satellites and the centrosome [42]. CSPP-L is a well-known

positive regulator of primary cilia formation [50] that was recently shown to bind directly to

CEP104 to control axoneme extension [51]. Furthermore, CEP104 is implicated in ciliogenesis

through interaction with the CEP97-CP110 complex and modulation of microtubule dynamics at

the ciliary tip [51-53]. It will therefore be interesting to investigate if and how CSPP-L and CEP104

localization is affected in CEP78 mutant cells. Moreover, as CEP350 additionally binds to CYLD

[54], a deubiquitinase that controls centriolar satellite homeostasis and ciliogenesis through de-

ubiquitination of the E3 ubiquitin ligase MIB1 [55, 56], and was proposed to regulate axonemal IFT

particle injection through FOP-CEP19-RABL2 interactions [24-26], it is tempting to speculate that

CEP78 might impinge on these processes.

Materials and methods

PCR and cloning

For the establishment of the N-terminal FLAG-tagged CEP78 construct a cDNA clone coding for

human full-length CEP78 cDNA (GeneCopoiea pShuttle™Gateway® PLUS ORF, NCBI accession

number NM_001098802) was used as template for amplification with the Phusion High-Fidelity

DNA polymerase (New England BioLabs) under standard PCR conditions and attB forward and

reverse primers (see Key Resources Table) compatible with the Gateway cloning technology (Invi-

trogen/Thermo Fisher Scientific) for inclusion of the insert into the pDonor201 Gateway vector

(Invitrogen/Thermo Fisher Scientific). To produce human FLAG-CEP78^{L150S}, site directed muta-

genesis was performed with QuickChange Lightning Site-Directed Mutagenesis Kit (Agilent) and

primers CEP78 mut_F and CEP78 mut_R (see Key Resources Table). Accuracy of the *CEP78*

ORF was assessed by Sanger sequencing. Subsequently, the desired fragment was cloned into

p3xFLAG-CMV/DEST (gift from Dr. Ronald Roepman lab, Radboud University Medical School,

Nijmegen, NL; see Key Resources Table for details) to produce p3xFLAG-CEP78 or p3xFLAG-

CEP78^{L150S}.

Plasmids encoding full-length and truncated versions (N-terminus, middle, N-terminus plus

middle, C-terminus) of CEP78 tagged with GFP in the N-terminus were generated by PCR using

relevant primers (see Key Resources Table) and pFLAG-CEP78 as template; PCR products were

cloned into pEGFP-C1 by standard approaches, following digestion with BamHI (Roche, cat#

1056704001) and KpnI (Roche, cat#10899186001) and ligation with T4 DNA ligase (Applichem,

cat# A5188). Ligated plasmids were transformed into competent *E. coli* DH10B cells and cells har-

boring recombinant plasmids selected on Luria Bertani (LB; Sigma-Aldrich) agar plates with 50

µg/ml kanamycin. Plasmids were purified using Plasmid DNA Mini Kit I (Omega Biotech, cat#

D6943-02) or Nucleobond Xtra Mid kit (Macherey-Nagel, cat# 740410.50), according to protocols

supplied by the manufacturers. Sequences of plasmid inserts were verified by Eurofins Genomics.

Gateway system compatible pENTR vectors encoding for mNeonGreen-CEP78 or mNeon-

Green-CEP78L150S fusions were generated by subcloning of ORFs from pEGFP constructs de-

scribed above in frame with mNeonGreen using the BamH1/Kpn1 sites. For lentivirus particle pro-

duction these plasmids were recombined with pCDH-EF1a-Gateway-IRES-BLAST plasmids as

described in [51] using LR Clonase II (Invitrogen/Thermo Fisher Scientific).

Mammalian cell culture and transfection

Human skin fibroblasts were cultured as described previously [29, 30]. Human embryonic kidney-

derived 293T cells were cultured and transfected with plasmids as described previously [61]. RPE1

cells were grown in T-75 flasks in a 95% humidified incubator at 37 ºC with 5% CO2. Cells were

cultured in Dulbecco's Modified Eagle Medium (DMEM) (GIBCO®, cat# 41966-029) supplement-

ed with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin (P/S). When cells were

about 80-90% confluent, they were passaged and set up for new experiments. Upon seeding, RPE1

cells were washed once with preheated 1 x phosphate buffered saline (PBS) and then incubated with

1% Trypsin-EDTA (ethylenediaminetetraacetic acid) (Sigma-Aldrich, cat# T4174) solution for 5

min at 37 °C. The detached cells were aspired in an appropriate volume of new preheated DMEM

and seeded into petri dishes with or without sterile glass coverslips [62]. When the cells had reached

80-90% confluency they were serum starved for 24 hours to induce formation of primary cilia. A

small amount of cell solution was passaged on in a new T-75 flask containing new preheated

DMEM. All cell lines were tested negative for Mycoplasma and those not generated in this study

have been used in prior publications. New cell lines generated in this study were validated by

sanger sequencing.

For transfection with plasmids, RPE1 cells were seeded in 60 mm petri dishes as described

above to about 50-70% confluency at the time of transfection. For rescue experiments, 2 μg

pFLAG-CEP78 or pFLAG-CEP78^{L150S} and 6 μl FuGENE®6 were mixed in 100 μl serum- and an-

tibiotic free DMEM and incubated for 20 minutes at room temperature. New DMEM was added to

the cells prior to transfer of the transfection reagent, which was gently dripped onto the cells and

swirled to make sure it was dispersed well. After cells were incubated with transfection medium

for 4 hours, the medium was replaced with serum free DMEM for 24 hours to induce growth arrest.

Gene silencing in RPE1 cells by siRNA was performed using DharmaFect (see Key Resources Ta-

ble). Briefly, RPE1 cells were seeded in DMEM medium with serum and antibiotics in 30 mm petri

dishes at 20-25% confluency. Transfection was performed by combining 50 nM siRNA with 5 μl of

DharmaFect in 200 μl of serum and antibiotic-free DMEM medium for 15 min at room tempera-

ture. After incubation, the complexes were added dropwise onto the cells and the media was swirled

gently to ensure good dispersion prior to incubation at 37 °C. Cells were kept under these condi-

tions for 48 hours and then serum starved for 24 hours before being processed for further experi-

ments.

RPE1 cell transduction with lentivirus particles and Blasticidine selection (10μg/ml f.c.) was

conducted as described in [51].

Immunofluorescence microscopy and image analysis

Standard epifluorescence IFM analysis was performed as described previously [62]. Briefly, glass

coverslips containing cells of interest were washed in PBS and fixed with either 4% paraformalde-

hyde (PFA) solution for 15 min at room temperature or on ice, or with ice-cold methanol for 10-12

min at -20 °C. After three washes in PBS the PFA-fixed cells were permeabilized 10 min in 0.2%

Triton X-100 and 1% Bovine Serum Albumin (BSA) in PBS and blocked in 2% BSA in PBS for 1

hour at room temperature. Coverslips were incubated overnight at 4 ºC in primary antibodies (see

Key Resources Table) diluted in 2% BSA in PBS. The next day, coverslips were washed three

times with PBS for 5 min and incubated for 1 hour at room temperature in secondary antibodies

(see Key Resources Table) diluted in 2% BSA in PBS. Coverslips were washed three times in PBS

for 5 min before staining with 2 µg/ml DAPI solution PBS for 30 sec. Finally, coverslips were

washed once with PBS and before mounting on objective glass with 6% n-propyl gallate diluted in

glycerol and 10x PBS and combined with Shandon^{TM} Immuno-Mount^{TM} (Thermo Scientific, cat#

9990402) in a 1:12 ratio. Cells were imaged on a motorized Olympus BX63 upright microscope

equipped with a DP72 color, 12.8 megapixel, 4140 x 3096 resolution camera. cellSense Dimension

software 1.18 from Olympus was used to measure cilia length; ImageJ version 2.0.0-rc-69/1.52i

was used to measure the relative mean fluorescence intensity (MFI) of relevant antibody-labeled

antigens at the centrosome/basal body. A fixed circle was drawn around a centrosome in a cell. This

same region of interest (ROI) was used to measure the MFI of a specific protein at the centrosome.

A constant ROI was also drawn to measure the cell background signal which was subtracted from

the MFI measured at the centrosome/basal body. Images were prepared for publication using Adobe

Photoshop® and Adobe Illustrator®.

Super resolution imaging of RPE1 cells was conducted as described in [51] except that cells

were mounted in ProLong Diamond Antifade mountant. Settings for hardware-assisted axial and

software-assisted lateral channel alignment and image reconstruction were validated by imaging of

RPE1 cells stained for CEP164 and labeled with secondary antibodies conjugated to either

Alexa488, DyeLight550, or Alexa647. For superimposing 3D-SIM images on electron microscopy

micrographs of centrioles (kindly provided by Dr. Michel Bornens; [40]), 3D-SIM images were

sized to same digital pixel resolution as original EM images using the bicubic algorithm in ImageJ

and maximal intensity z-projections of single channel images overlaid on micrographs using centri-

ole centers / centriole distal ends as reference points.

Statistical analysis

Statistical analyses were performed using GraphPad Prism 6.0. The background-corrected MFI

measured at the centrosome was normalized to relevant control cells. Mean and standard deviation

(SD) was calculated for all groups, and outliers were identified and removed by the ROUT method

before the statistical tests were conducted. The data was tested for gaussian normality using either

D'Agostino's K-squared test or Shapiro-Wilk test. Depending on the distribution of the data, two-

tailed and unpaired Student's t-test or Mann Whitney test were used when comparing two groups.

Also, depending on the distribution of the data, One-way ANOVA followed by Tukey's, Dunnet´s

or Dunn's multiple comparison tests was used when comparing more than two groups. Unless oth-

erwise stated, the statistical analyses were performed on at least three independent biological repli-

cates (n=3). A p-value under 0.05 was considered statistically significant and p-values are indicated

in the figures with asterisks as follows; * p<0.05, ** <p0.01, ***<0.001, and **** p<0.0001.

Immunoprecipitation (IP), protein quantification, SDS-PAGE and western blot analysis

Immunoprecipitation of transfected 293T cell lysates was performed as described previously [61]

using relevant antibody-conjugated beads (see Key Resources Table). Protein concentrations were

measured using the DC^{TM} Protein Assay Kit I from Bio-Rad (cat # 5000111) by following the man-

ufacturer's protocol. For SDS-PAGE analysis of RPE1 and HSF cells, protein samples were pre-

pared by lysis of cells with 95 ºC SDS-lysis buffer (1% SDS, 10 mM Tris-HCl, pH 7.4) and cell

lysates transferred to Eppendorf tubes and heated shortly at 95 ºC. The samples were then sonicated

two times for 30 seconds to shear DNA followed by centrifugation at 20,000 x g for 15 min at room

temperature to pellet cell debris. Supernatants were transferred to new Eppendorf tubes and an ali-

quot of each sample used for determination of protein concentration. Samples with equal concentra-

tions of protein were prepared for SDS-PAGE analysis by addition of NuPAGE^{TM} LDS Sample

Buffer (4X) from Thermo Fisher Scientific (cat# NP0007) and 50 mM DTT. Samples from IP anal-

ysis in 293T cells, performed using modified EBC buffer [61], were prepared for SDS-PAGE in a

similar fashion. Protein samples were heated at 95 ºC for 5 minutes before loading them on a Mini-

PROTEAN® TGX^{TM} Precast Gel 4-15% from Bio-Rab Laboratories, Inc. (cat# 456-1083 or cat

#456-1086). PageRuler^{TM} Plus Prestained Protein Ladder (Thermo Fisher Scientific, cat# 26619)

was used as molecular mass marker. SDS-PAGE was performed using the Mini-PROTEAN® Tetra

System from Bio-Rad. Gels were run at 100 V for 15 minutes and 200 V for 45 minutes. Using the

Trans-BLOT® Turbo Transfer System from Bio-Rad Laboratories, Inc. the proteins were trans-

ferred at 1.3 A, 25 V for 10 min from the gel to a Trans-Blot® Turbo^{TM} Transfer Pack, Mini format

0.2 μm Nitrocellulose membrane (cat# 1704158). Ponceau-S solution was added to the membrane

to visualize the proteins before blocking in 5% milk in Tris-buffered saline with Tween-20 (TBS-T;

10 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.1% Tween-20) for 2 hours at room temperature. Prima-

ry antibodies (see Key Resources Table) were diluted in 5% milk in TBS-T and incubated with the

membrane overnight at 4ºC. The membrane was washed three times 10 min at room temperature in

TBS-T on a shaker before incubation with secondary antibody (see Key Resources Table), diluted

in 5% milk in TBS-T, for 1 hour at room temperature. The membrane was washed three times 10

min at room temperature in TBS-T on a shaker. Finally, SuperSignal[TM] west Pico PLUS

chemiluminescent Substrate (Thermo Scientific, cat # 34580) was mixed in a 1:1 ratio and added to

the membrane for 5 min before development on a FUSION FX SPECTRA machine from Vilber.

Quantitative mass spectrometry (MS) and peptide identification

For SILAC-based mass spectrometry, three distinct SILAC culture media were used: light- (Lys0

and Arg0); medium- (Lys4 and Arg6) and high-labeled media (Lys8 and Arg10). 293T cells were

seeded in 100 mm petri dishes and SILAC labeled for 1 week to ensure proper amino acid incorpo-

ration. After the incorporation phase, the cells were transfected with 2 µg of plasmid encoding

FLAG-Ap80 (control), FLAG-CEP78 or FLAG-CEP78[L150S] using 6 µl FuGENE®6, as described

above, and then subjected to lysis using modified EBC buffer [56] and FLAG IP as described

above. Affinity enriched proteins were digested with trypsin and the resulting peptides desalted

prior to LCMS analysis using an Easy-nLC 1000 system (Thermo Scientific) connected to a Q

Exactive HF-X mass spectrometer (Thermo Scientific). Peptides were separated by a 70 minutes

gradient using increased buffer B (95% ACN, 0.5% acetic acid). The instrument was running in

positive ion mode with MS resolution set at 60,000 for a scan range of 300 to 1700 m/z. Precursors

were fragmented by higher-energy collisional dissociation (HCD) with normalized collisional ener-

gy of 28 eV. For protein identification and quantitation, the obtained MS raw files were processed

by MaxQuant software version 1.6.1.0 [60] and searched against a FASTA file from UniProt.

RNAseq analysis

RNA was extracted using the Direct-zol RNA Miniprep Kit (Zymo Research) following the manu-

facturer's instructions. RNA degradation and contamination were monitored on 1% agarose gels.

RNA purity was checked using the NanoPhotometer® spectrophotometer (IMPLEN, CA, USA).

RNA concentration was measured using Invitrogen[TM] Qubit® RNA Assay Kit in Qubit® 2.0

Fluorometer (Thermo Fisher Scientific). RNA integrity was assessed using the Agilent RNA Nano

6000 Assay Kit of the Bioanalyzer 2100 system (Agilent, CA, USA). A total amount of 3 µg RNA

per sample was used as input material for the RNA sample preparations. Sequencing libraries were

generated using NEBNext® Ultra™ RNA Library Prep Kit for Illumina® (New England BioLabs,

USA) following manufacturer's recommendations and index codes were added to attribute se-

quences to each sample. Briefly, mRNA was purified from total RNA using poly-T oligo-attached

magnetic beads. Fragmentation was carried out using divalent cations under elevated temperature in

NEBNext First Strand Synthesis Reaction Buffer （5X）. First strand cDNA was synthesized using

random hexamer primer and M-MuLV Reverse Transcriptase （RNase H -）. Second strand cDNA

synthesis was subsequently performed using DNA Polymerase I and RNase H. Remaining over-

hangs were converted into blunt ends via exonuclease/polymerase activities. After adenylation of 3'

ends of DNA fragments, NEBNext Adaptor with hairpin loop structure were ligated to prepare for

hybridization. In order to select cDNA fragments of preferentially 150~200 bp in length, the library

fragments were purified with AMPure XP system (Beckman Coulter Life Sciences). Then 3 µl US-

ER® Enzyme (New England BioLabs) was used with size-selected, adaptor-ligated cDNA at 37°C

for 15 min followed by 5 min at 95 °C before PCR. Then PCR was performed with Phusion® High-

Fidelity DNA polymerase, Universal PCR primers and Index (X) Primer (New England BioLabs).

At last, PCR products were purified (AMPure XP system) and library quality was assessed on the

Agilent Bioanalyzer 2100 system. The clustering of the index-coded samples was performed on a

cBot Cluster Generation System using HiSeq PE Cluster Kit cBot-HS (Illumina) according to the

manufacturer's instructions. After cluster generation, the library preparations were sequenced on an

Illumina Hiseq platform and 125 bp/150 bp paired-end reads were generated. Raw data (raw reads)

of fastq format were firstly processed through in-house perl scripts. In this step, clean data (clean

reads) were obtained by removing reads containing adapter, reads containing poly-N and low quali-

ty reads from raw data. At the same time, Q20, Q30 and GC content the clean data were calculated.

All the downstream analyses were based on the clean data with high quality. Reference genome and

gene model annotation files were downloaded from genome website directly. Index of the reference

genome was built using Bowtie v2.2.3 [63] and paired-end clean reads were aligned to the reference

genome using TopHat v2.0.12 [64]. We selected TopHat as the mapping tool for that TopHat can

generate a database of splice junctions based on the gene model annotation file and thus a better

mapping result than other non-splice mapping tools. HTSeq v0.6.1 [65] was used to count the reads

numbers mapped to each gene and then Fragments Per Kilobase of transcript sequence per Millions

base pairs sequenced (FPKM) of each gene was calculated based on the length of the gene and

reads count mapped to this gene. FPKM considers the effect of sequencing depth and gene length

for the reads count at the same time, and is currently the most commonly used method for estimat-

ing gene expression levels [66]. Prior to differential gene expression analysis, for each sequenced

library, the read counts were adjusted by edgeR program package through one scaling normalized

factor [67]. Differential expression analysis of two conditions was performed using the DEGSeq R

package (1.20.0) [68].

Acknowledgements

We thank Søren Lek Johansen and Maria Schrøder Holm for expert technical assistance, Benedicte

Schultz Kappel and Frida Roikjer Rasmussen for help with plasmid generation, and Nynne Chris-

tensen, Center for Advanced Bioimaging (CAB) Denmark, for help with 3D-SIM acquisition. Dr.

Kay Schink kindly provided plasmids for lentivirus particle generation and assisted in 3D-SIM ac-

quisition at OUH Radiumhospitalet. We are grateful to Drs. Peter K. Jackson, Tomoharu Kanie,

Francesc Garcia-Gonzalo, Laurence Pelletier, Brian David Dynlacht, Anne-Marie Tassin, Maria

Gavilan, Rosa M. Rios, Éric A. Cohen and Peter ten Dijke for reagents, and Dr. Michel Bornens for

sharing original EM microcraphs. This study was supported grants from Independent Research

Fund Denmark (#8020-00162B to PF and LBP, and #8021-00425A to JSA), from the Carlsberg

Foundation (#CF18-0294 to LBP) and from the Swiss National Science Foundation (#176097 to

CR).

Figure legends

Figure 1. CEP78 mutant cells display fewer, but longer primary cilia. (**A**) Representative IFM im-

ages of serum-deprived RPE1 WT and *CEP78* KO cells stained with the indicated antibodies; DAPI

marks the nucleus. Scale bar, 5μm. (**B**, **C**) Quantification of the percentage of ciliated cells (**B**) and

the length of residual cilia (**C**) in WT and *CEP78* KO cells, based on images as shown in (A). Data

were normalized in relation to WT values. Student's t-test (two-tailed, unpaired) was used for the

statistical analysis. Graphs in (B) represent accumulated data from three individual experiments

(n=384 for WT and n=322 for *CEP78* KO cells). Graphs in (C) show data from three individual

experiments (n=338 for WT and n=198 for *CEP78* KO cells). (**D**) Representative IFM images of

serum-deprived control and *CEP78* mutant (Patient) HSFs stained with the indicated antibodies;

DAPI marks the nucleus. Scale bar, 5 μm. (**E**, **F**) Quantification of the percentage of ciliated cells

(**E**) and the length of residual cilia (**F**) in control and *CEP78* patient HSFs (data from HSFs derived

from patient 2702r34, individual II-3 [29]), based on images as shown in (D). Graphs in (E) repre-

sent accumulated data from seven individual experiments (n= 678 for control HSFs; n= 707 for

patient-derived HSFs). Graphs in (F) show data from seven individual experiments (n= 237 for con-

trol HSFs; n= 216 for patient-derived HSFs). Data were normalized in relation to control values. A

Student's t-test (unpaired, two-tailed) was performed to address differences in the ciliary frequency

and length between control and patient HSFs. (**G**) Western blot analysis of Rb phosphorylated on

S807/811 (P-Rb) in RPE1 WT and *CEP78* KO cells grown with or without serum. (**H, I**) Quantifi-

cation of data shown in (G) from three independent experiments analyzed in duplicates. (**J**) P-Rb

blots from HSFs derived from control and patient HSFs grown with or without serum (data from

HSFs derived from patient F3: II:1; [30]). GAPDH was used as a loading control. (**K, L**) Quantifi-

cation of data shown in (J) from three independent experiments analyzed in duplicates. Student's t-

test (two-tailed, unpaired) was used for the statistical analysis in (H, I) and (K, L). Error bars of

graphs represent SD and data are shown as mean ± SD. a.u., arbitrary units; (*) $p<0.05$; (****)

$p<0.0001$; n.s., not statistically significant.

Figure 2. Analysis of RPE1 WT and *CEP78* KO cells expressing FLAG- or mNG-tagged CEP78 or

CEP78^{L150S} fusions. (**A**) Representative IFM images of serum-starved WT and *CEP78* KO cells

transiently expressing FLAG-CEP78 and FLAG-CEP78^{L150S} labeled with antibodies against FLAG

(green) and ARL13B (magenta). Insets show enlarged views of the cilia-centrosome region. Scale

bars: 5 µm in original images and 1 µm in magnified images. (**B**) Quantification of the relative MFI

of FLAG-CEP78 and FLAG-CEP78^{L150S} at the centrosome in serum-starved WT cells. Accumulat-

ed data from three individual experiments (n=145 and n=151 for cells transfected with FLAG-

CEP78 and FLAG-CEP78^{L150S}, respectively). Student's t-test (two-tailed, unpaired) was used for

the statistical analysis. Data is shown as mean ±SD. (**C, D**) Quantification of the percentage of cili-

ated cells (**C**) or ciliary length (**D**) in serum-deprived WT and *CEP78* KO cells stably expressing

the mNG-CEP78 or mNG-CEP78^{L150S}. Analysis on cilia numbers and length was performed

through accumulated data from three independent experiments for WT, *CEP78* KO and WT +

mNG-CEP78 cells and five independent experiments for *CEP78* KO cells expressing either the

mNG-CEP78 or the mNG-CEP78^{L150S} (n=338 for untransfected WT; n=198 for untransfected

CEP78 KO; n=138 for WT + mNG-CEP78; n=280 for *CEP78* KO + mNG-CEP78 and n=141 for

CEP78 KO + mNG-CEP78^{L150S}). Ordinary one-way ANOVA with Dunnett's multiple comparison

test was used for the statistical analysis of the ciliary frequency and length amongst all groups, in

relation to the mean of untransfected WT cells, designated as the control group. Differences be-

tween WT + mNG-CEP78, *CEP78* KO + mNG-CEP78 and *CEP78* KO + mNG-CEP78^{L150S} cells

were addressed in a pairwise fashion using an unpaired and two-tailed Student´s t-test. Error bars in

(C) indicate SD and data in (D) is presented as mean ± SD. a.u., arbitrary units; (*) p<0.05; (**)

p<0.01; (****) p<0.0001.

Figure 3. CEP78 interacts with the EDD-DYRK2-DDB1^{VPRBP} complex and CEP350. (**A**) Over-

view of results from CEP78 interactome analysis in 293T cells grown in SILAC medium. Cells

expressing FLAG-CEP78 WT, FLAG-CEP78^{L150S} or FLAG-Ap80 (negative control) were subject-

ed to FLAG IP and pellets analyzed by mass spectrometry. The upper panel displays affinity en-

richment of Cep78 and interaction partners in FLAG-CEP78 WT cells relative to control cells. The

lower panel shows how the CEP78^{L150S} mutation weakens these interactions relative to CEP78 WT

using ratios normalized for equal affinity enrichment of FLAG-tagged CEP78. (**B-D**) 293T cells

expressing the indicated Myc-, FLAG-, or GFP-fusions were subjected to IP with anti-FLAG (**B**) or

-GFP (**C, D**) beads and input and pellet fractions analyzed by SDS-PAGE and western blotting with

Myc, FLAG or GFP antibodies as indicated. FLAG-Ap80-NB used in (B) is a negative control.

Molecular mass markers are indicated in kDa to the left of the blots. (**E**) Schematic of the CEP350

and CEP78 fusions used in IP analysis. LRR, leucine-rich repeat; CC, coiled coil. The CEP350 con-

structs were described in [54, 58]; the CEP78 constructs were generated in this study based on pub-

lished sequence information [27, 30, 35]. N, N-terminal region; M, middle region; NM, N-terminal

plus middle region; C, C-terminal region; FL, full length.

Figure 4. CEP350 regulates stability and centrosomal recruitment of CEP78. (**A**) 3D-SIM analysis

of mNG-CEP78 and CEP350 localization in WT RPE1 cells, relative to CP110 and CEP164 (max-

imal z-projections). Cells were fixed and stained with the indicated antibodies. Montage panels

show single channel images of the mother centriole region (left panels) and overlays on previously

published electron micrographs of isolated mother centrioles from KE37 cells, reproduced with

permission from [40] (top: cross-sections of proximal and subdistal-appendage region; bottom:

side-view). Scale bar, 200 nm. (**B**) Western blot analysis of CEP78 and α-tubulin (loading control)

in serum-deprived hTERT-RPE1-BFP-Cas9 control (pMCB306), *CEP350* KO or *FOP* KO cells

[24]. (**C**) Quantification of data in (B), based on three independent experiments analyzed in dupli-

cates. Error bars indicate SD. Statistical analysis was done using Student's t-test (unpaired, two-

tailed). (**D, F**) IFM analysis of serum-deprived control and *CEP350* KO ciliated (D) and non-

ciliated cells (F) using antibodies as indicated. Dashed boxes indicate cropped images to highlight

the centrosomal/ciliary area. Scale bars: 5 μm in representative images and 1 μm in insets. (**E, G**)

Quantification of the relative MFI of CEP78 at the centrosome in serum-starved, ciliated (E) and

non-ciliated (G) control and *CEP350* KO cells, based on images as shown in (D) and (F), respec-

tively. Statistical analysis was done using a two-tailed and unpaired Mann-Whitney test for ciliated

cells and two-tailed and unpaired Student's t-test for non-ciliated cells. Accumulated data from

three individual experiments (n=128 and 132 for control and *CEP350* KO ciliated cells, respective-

ly; n= 101 and 107 for control and *CEP350* KO non-ciliated cells, respectively). a.u., arbitrary

units; (*) $p<0.05$; (**) $p<0.01$; (****) $p<0.0001$.

Figure 5. Altered centrosomal levels of CEP350 in *CEP78* mutant cells. (**A, C, E, G**) Representa-

tive IFM images of ciliated (**A**) and non-ciliated (**C**) serum-deprived RPE1 WT and *CEP78* KO

cells, and ciliated (**E**) and non-ciliated (**G**) serum-deprived control and CEP78 deficient (Patient)

HSFs (combined data from HSFs from patient F3: II:1 [30] and patient 2702r34, individual II-3

[29]) labeled with antibodies indicated in the figure. Insets show closeups of the centrosomal/ciliary

area (dashed boxes). Scale bars: 5 µm in representative images and 1 µm in closeups. (**B, D**). Quan-

tification of CEP350 relative MFI at the centrosome based on images as shown in (A, C). Data is

shown as mean ± SD. A Student's t-test was used as statistical analysis between the two groups

based on three individual experiments (n=157 or 158 for ciliated cells and n=144-152 for non-

ciliated cells). (**F, H**) Quantification of CEP350 relative MFI at the centrosome in HSFs based on

images as shown in (E, G). Data is shown as mean ± SD. Student's t-test was used as statistical

analysis between the two cell groups based on seven individual experiments with a total of 154 and

156 analyzed ciliated cells (**F**), and three individual experiments with a total of 122 or 126 analyzed

non-ciliated cells (**H**). a. u., arbitrary units; n.s, not statistically significant; (****) $p<0.0001$.

Figure 6. Analysis of EDD-DYRK2-DDB1$^{\text{VPRBP}}$ complex components in CEP78 deficient cells.

(**A**) Western blot analysis of VPRBP in lysates from serum-fed and serum-deprived RPE1 WT and

CEP78 KO cells. α-tubulin was used as a loading control (**B, C**) Quantification of the VPRBP rela-

tive levels in the different conditions depicted in (A). Statistical analysis was performed using a

Student's t-test (unpaired, two-tailed) from five independent experiments analyzed in duplicates.

Error bars indicate SD. (**D, F**) Representative IFM images of ciliated (**D**) and non-ciliated (**F**) se-

rum-deprived RPE1 WT and *CEP78* KO cells labeled with antibodies against EDD1 (green) and

DCTN1 plus acetylated tubulin (magenta). DAPI was used to mark the nucleus (blue). Insets show

enlarged views of the cilium-centrosome region. Scale bars: 5 µm in original images and 1 µm in

closeups. (**E, G**) Quantification of the EDD1 MFI at the centrosome based on images as shown in

(D) and (F) using a two-tailed and unpaired Student's t-test. Data is shown as mean ± SD. Student's

t-test from three independent experiments (n=194 and n=201 for ciliated WT and *CEP78* KO cells,

respectively; n=194 and n=217 for non-ciliated WT and *CEP78* KO cells, respectively). Data is

shown as mean ± SD. a.u., arbitrary units; n.s., not statistically significant; (****) p<0.0001.

Figure 7. CEP78 deficient cells display elevated cellular and centrosomal levels of CP110. (**A**)

Western blot analysis of lysates from serum-deprived RPE1 WT and *CEP78* KO cells using indi-

cated antibodies. GAPDH was used as a loading control. (**B**) Quantification of the data shown in

(A), based on four independent experiments analyzed in duplicates. Error bars indicate SD. Statisti-

cal analysis was done using Student's t-test (unpaired, two-tailed). (**C, F**) Representative IFM im-

ages of serum-starved ciliated (**C**) and non-ciliated (**F**) RPE1 WT and *CEP78* KO cells labeled with

antibodies against CP110 (green), combined DCTN1 and acetylated tubulin (Ac-tub; magenta) and

DAPI to mark the nucleus (blue). Insets show enlarged views of the cilium-centrosome area. Scale

bars: 5 µm in original images and 1 µm in closeups. (**D, G**) Quantification of the relative MFI of

CP110 at the centrosome based on images as shown in (C) and (F), respectively. Student's t-test

(unpaired, two-tailed) from three independent biological experiments (n= 90 and n=75 for ciliated

RPE1 WT and *CEP78* KO cells, respectively; n=75 and n=82 for non-ciliated RPE1 WT and

CEP78 KO cells, respectively) was used for statistical analysis. Data is presented as mean ± SD. (**E**)

Quantification of the relative MFI of CP110 at the mother centriole based on images shown in (C)

Mann Whitney test (two-tailed and unpaired) was used as statistical analysis based on two inde-

pendent experiments (n=61 and n=62 for RPE1 WT and CEP78 KO cells, respectively). Data is

presented as mean ± SD. (**H**) Representative IFM images of serum-deprived healthy and CEP78

deficient (Patient) HSFs (data from patient 2702r34, individual II-3 described in [29]) labeled with

the indicated antibodies. DAPI was used as counterstaining to mark the nucleus. Dashed lines show

closeup images of the centrosome region. Scale bars: 5 µm in original images and 1 µm in closeups.

(**I**) Quantification based on observations of 151 and 155 cells of CEP78 control and patient cells,

respectively, from three individual experiments. Student's t-test (unpaired and two-tailed) was used

to assess differences between the two groups. a.u., arbitrary units; n.s., not statistically significant;

(**) p<0.01; (***) p<0.001; (****) p<0.0001.

Figure 8. Depletion of CP110 from *CEP78* KO cells rescues their ciliation frequency phenotype.

(**A**) Representative western blots of RPE1 WT or *CEP78* KO cells treated with control (siMock) or

CP110-specific siRNA (siCP110). Cells were deprived of serum and blots were probed with anti-

body against CP110 or α-tubulin (loading control). (**B**) Quantification of relative cellular CP110

levels under different conditions based on blots as shown in (A) (n=5 for WT and n=4 for *CEP78*

KO cells; each sample was analyzed in duplicates). Error bars indicate SD. Data was normalized in

relation to the WT cells transfected with Mock siRNA (WT siMock), designated as the control

group. Differences of CP110 relative levels between *CEP78* KO cells transfected with Mock or

CP110 siRNA and WT cells transfected with CP110 siRNA in relation to control were assessed by

performing an ordinary one-way ANOVA with a Dunnet´s multiple comparison test. Differences of

CP110 cellular levels between *CEP78* KO cells transfected with the same siRNAs was addressed

using an unpaired and two-tailed Student's t-test. (**C**) Representative IFM images of serum-

deprived RPE1 WT and *CEP78* KO cells treated with control (siMock) or CP110-specific siRNA

(siCP110) and stained for ARL13B (green). DAPI marks the nucleus. Dashed boxes indicate

cropped images to highlight the centrosomal/ciliary zone. Scale bars, 5 μm in representative images

and 1μm in insets. (**D**, **E**) Quantification of percentage of ciliated cells (**D**) and length of remaining

cilia (**E**) in RPE1 WT or *CEP78* KO cells treated with control (siMock) or CP110-specific siRNA

(siCP110). The quantification is based on images as shown in (C) and represents data from four and

five independent experiments, respectively, for RPE1 WT (n=554-617cells per condition) and

CEP78 KO cells (n=846-856 cells per condition). Ordinary one-way ANOVA with Dunnett's mul-

tiple comparison test was used, to compare the mean of the remaining groups in relation with the

mean of RPE1 WT cells treated with siMock, which was designated as the control group. Differ-

ences between *CEP78* KO cells transfected with the above-mentioned siRNA were discriminated

using an unpaired and two-tailed Student´s t-test. Bars in (D) represent SD and data in (E) is pre-

sented as mean ± SD. a.u., arbitrary units; (*) $p<0.05$; (**) $p<0.01$; (***) $p<0.001$; (****)

$p<0.0001$. (**F**) Proposed model for how CEP78 regulates ciliogenesis and ciliary length control. See

text for details.

Supplementary figure legends

Figure 1-figure supplement 1. Western blot analysis and ciliary frequency and length in RPE1 WT

and *CEP78* KO clones. (**A**) Western blot analysis of serum-deprived RPE1 WT and four different

CEP78 KO clones, generated by CRISPR/Cas9 methodology. Blots were probed with antibodies

against CEP78 and GAPDH (loading control), as indicated. The lower band in the CEP78 blot is

due to unspecific staining of the antibody. Molecular mass markers are indicated in kDa to the right.

Unless otherwise indicated, clone #73 was used for all experiments in RPE1 cells performed in this

manuscript, and is designated *CEP78* KO. (**B, C**) Quantification of ciliary frequency (B) and length

(C) in serum-deprived RPE1 WT and different *CEP78* KO clones. Differences in cilia numbers and

size between the WT and the several *CEP78* KO clones were addressed through accumulated data

from three independent experiments (for cilia frequency: n= 304 for WT; n=308 for clone #2;

n=321 for clone #44 and n=311 for clone #52; and for ciliary length: n=145 for WT; n=68 for clone

#2; n=98 for clone #44 and n=62 for clone #52) by performing an ordinary one-way ANOVA with

Dunnet´s multiple comparison test amongst all groups, in relation to WT cells mean (control

group). Error bars in (B, C) indicate mean ± SD. (*) $p<0.05$; (**) $p<0.01$; (***) $p<0.001$.

Figure 2-figure supplement 1. Expression of FLAG-CEP78 and FLAG-CEP78^{L150S} in RPE1 WT

and *CEP78 KO* cells. (**A**) FLAG blots from serum-deprived WT and *CEP78* KO lysates transfected

with indicated FLAG fusions. GAPH was used as a loading control. (**B**) Quantification of data de-

picted in (A). Graphs of the FLAG blots are presented as the mean + SD from three individual ex-

periments analyzed in duplicates. Ordinary one-way ANOVA with Tukey's multiple comparison

test was used for the statistical analysis, where no significant statistical differences were reported

between the groups.

Figure 2-figure supplement 2. CEP78 endogenous antibody binds equally well to FLAG-CEP78

WT and L150S mutant fusions. 293T cells transiently expressing FLAG-CEP78 or FLAG-

CEP78^{L150S} were subjected to IP with FLAG. Pellet fractions were analyzed by SDS-PAGE and

western blotting using antibodies against FLAG and endogenous CEP78. Molecular mass marker is

indicated in kDa to the left of the blots.

Figure 2-figure supplement 3. Stable expression of mNG-CEP78 and mNG-CEP78^{L150S} in RPE1

WT and *CEP78* KO cells. (**A**) Western blots of lysates of serum-deprived WT and *CEP78* KO

RPE1 cells lines stably expressing the indicated mNG-tagged fusions. The blots were probed with

antibodies against CEP78 (upper panel) and α-tubulin (loading control; lower panel). Molecular

mass markers are shown in kDa to the left. (**B**) Representative light (upper panels) and fluorescence

(lower panels) images of live cells expressing the indicated fusions. Note that mNG-CEP78 is con-

centrated at the centrosome whereas mNG-CEP78^{L150S} is not. Scale bar, 20 µm.

Figure 3-figure supplement 1. CEP78 associates independently with CEP350 and VPRBP. (**A**)

293T cells expressing the indicated Myc- or GFP-fusions were subjected to IP with Myc antibody-

coated beads and input and pellet (IP Myc) fractions analyzed by SDS-PAGE and western blotting

with VPRBP, Myc, FOP or GFP antibodies as indicated. (**B**) Similar analysis as in (A), but using

GFP antibody-coated beads for IP instead of Myc beads. Molecular mass markers are indicated in

kDa to the left of the blots. Arrowhead in (A) indicates IgG band observed in the lower part of the

GFP blot.

Figure 4-figure supplement 1. Validation of *CEP350* KO cells by IFM analysis. (**A**) RPE1 control

and *CEP350* KO cells were serum starved for 24 hours before fixation and staining with the indi-

cated antibodies and DAPI for nuclear visualization. Insets show closeups of the centrosomal/ciliary

area (dashed boxes). Scale bars: 5 µm in representative images and 1 µm in closeups. (**B**) Quantifi-

cation of CEP350 mean fluorescence intensity (MFI) at the centrosome, based on images as shown

in (A). A total of 38 and 26 cells were analyzed for control and *CEP350* KO cells, respectively. For

statistical analysis, the data was first normalized to the control group values and differences were

assessed using an unpaired and two-tailed Student´s t-test. Data is shown as the mean ± SD. a.u.,

arbitrary units; (**) $p < 0.01$.

Figure 6-figure supplement 1. VPRBP centrosomal levels are increased or unchanged in RPE1

CEP78 KO cells. (**A, C**) Representative IFM micrographs of ciliated (**A**) and non-ciliated (**C**) se-

rum-starved RPE1 WT and *CEP78* KO cells labeled with the antibodies indicated in the figure.

DAPI was used to mark the nucleus (blue). Insets show enlarged views of the cilium-centrosome

region. Scale bars: 5 µm in original images and 1 µm in closeups. (**B, D**) Quantification of the

VPRBP MFI at the centrosome based on images shown in (A) and (C) based on data from three

independent experiments (n= 156 and n=158 for ciliated WT and CEP78 KO cells, respectively; n=

162 and n= 166 for non-ciliated WT and CEP78 KO cells, respectively) using a two-tailed and un-

paired Student's t-test. Data is shown as mean ± SD and was first normalized in relation to WT val-

ues. a.u., arbitrary units; (**) $p < 0.01$; n.s., not statistically significant.

Figure 6-figure supplement 2. EDD1 centrosomal levels are reduced in RPE1 *CEP350* KO cells.

(**A, C**) Representative IFM micrographs of ciliated (**A**) and non-ciliated (**C**) serum-starved RPE1

control and *CEP350* KO cells labeled with the antibodies indicated in the figure. DAPI was used to

mark the nucleus (blue). Insets show enlarged views of the cilium-centrosome region. Scale bars: 5

μm in original images and 1 μm in closeups. (**B, D**) Quantification of the EDD1 MFI at the centro-

some based on images as shown in (A) and (C) using a two-tailed and unpaired Student´s t-test

from two independent experiments (n=105 and n= 111 for ciliated WT and *CEP78* KO cells, re-

spectively; n=118 for non-ciliated WT and CEP78 KO cells). Data is shown as mean ± SD and was

first normalized in relation to WT values. a.u., arbitrary units; (****) $p < 0.0001$.

Figure 7-figure supplement 1. Elevated cellular CP110 levels in RPE1 *CEP78* KO clones. (**A**)

Western blot analysis of serum deprived RPE1 WT and three different *CEP78* KO clones, as indi-

cated. Blots were probed with antibodies against CP110 and GAPDH (loading control). Molecular

mass markers are indicated in kDa to the right. (**B**) Quantification of relative cellular CP110 levels

based on blots as shown in (A) from two independent experiments. The means of the CEP78 KO

clones, were compared in relation to the WT mean, selected as the control group using an ordinary

one-way ANOVA followed by a Dunnet´s multiple comparison test. Error bars of graphs represent

SD and data are shown as mean ± SD. (**C**) Representative western blot of WT and *CEP78* KO cells

(KO; clone #73) stably expressing mNG-CEP78 or mNG-CEP78^{L150S} as indicated. Blots were

probed with indicated antibodies (α-tub=α-tubulin) and the molecular mass markers are indicated

in kDa to the right of the blot. Quantification of relative CP110 band intensities revealed no signifi-

cant changes between the three conditions; CP110 level relative to that of WT mNG-CEP78 was

95% +/-12% for the *CEP78* KO mNG-CEP78 line and 90% +/- 9% for the *CEP78* KO mNG-

CEP78^{L150S} line, respectively (n=3). Blots were probed with indicated antibodies.

Figure 7-figure supplement 2. Serum-fed CEP78 deficient cells display elevated cellular and

centrosomal levels of CP110. (**A**) CP110 western blot analysis of lysates from RPE1 WT and

CEP78 KO cells grown in the presence of serum; α-tubulin serves as loading control. (**B**) Quantifi-

cation of the CP110 blots depicted in (A) based on four individual experiments analyzed in dupli-

cates. Error bars indicate SD. Student's t-test (unpaired, two-tailed) was used to analyze the differ-

ences between both cellular groups. (**C**) Representative IFM images of cycling RPE1 WT and

CEP78 KO cells labeled with the indicated antibodies. DAPI was used to mark the nucleus. Insets

show closeup images of the centrosome region. Scale bars: 5 µm in original images and 1 µm in

closeups. (**D**) Data quantification of the relative MFI of CP110 based on images as shown in (C).

Data is present as mean ±SD. Statistical analysis was performed using Student's t-test (unpaired,

two-tailed) of three independent experiments (n=158 for WT and n= 147 for *CEP78* KO cells).

a.u., arbitrary units; (*) $p<0.05$; (****) $p<0.0001$.

Figure 7-figure supplement 3. Barplots showing expression of *CP110, CEP350,*

VRPBP and *EDD1* genes for patients (n=3) and controls (n=4), based on RNA-seq data of skin fi-

broblasts. Expression levels are shown in fpkm (Fragments Per Kilobase of transcript sequence per

Millions base pairs sequenced). The patient data are combined data from HSFs derived from three

different *CEP78* probands; F3 individual II:1 described in [30] and KN10 individual II-1 and

2716s15 individual II-2 described in [29]. Control data are combined data from HSFs from four

unaffected individuals described in the same studies. Nominal p-values derived from unpaired two-

tailed t-tests are shown on top of the bars. None of the comparisons were significantly different

($p<0.05$).

Figure 7-figure supplement 4. Relative CP110 levels in WT and *CEP78* KO cells treated with

cycloheximide. (**A**) Representative western blot of lysates from serum-deprived WT and *CEP78*

KO cells treated with 200 ng/ml cycloheximide (CHX) for the indicated times. The drug was added

at the onset of serum deprivation (time 0 h). Blots were probed with CP110 and GAPDH (loading

control) antibodies; molecular mass markers are shown in kDa to the left. (**B**) Quantification of av-

erage CP110 levels determined based on western blots as shown in (A). The analysis is based on

data from four independent experiments, normalized to the relative CP110 level for each cell type at

time zero. Note that the data indicates decline in CP110 levels with time in the WT cells but not in

the *CEP78* KO cells; however, due to fluctuations from experiment to experiment the changes were

not statistically significant (ns; p values denoted in parenthesis above the graphs). The data is pre-

sented as mean ±SD.

Figure 8-figure supplement 1. Validation of CP110 knockdown in RPE1 WT and *CEP78* KO

cells by IFM. (A, B) Quantification of CP110 centrosomal levels in ciliated and non-ciliated WT

and *CEP78* KO cells treated with control (siMock) or CP110-specific siRNA (siCP110). Both

graphs are based on one biological experiment with 35 observations of ciliated and 35 observations

of non-ciliated cells per condition. Statistical analysis was performed using and Ordinary One-way

ANOVA followed by a Dunnett´s multiple comparison test amongst all groups in relation to WT

siMock, designated as the control group. Differences between *CEP78* KO cells treated with siMock

or siCP110 were addressed by performing an unpaired and two-tailed Student´s t-test. Bars are rep-

resented as mean ±SD. (a.u.) arbitrary units; (*) $p < 0.05$; (**) $p < 0.01$.

Additional files

Figure 1-source data 1. Original western blots corresponding to Figure 1G. The upper blots are

labeled with pRb antibody and lower blots with GAPDH antibody.

Figure 1-source data 2. Original western blots corresponding to Figure 1J. The upper blots are

labeled with pRb antibody and lower blots with GAPDH antibody.

Figure 3-source data 1. Raw data from the CEP78 interactome analysis depicted in Figure 3A.

293T cells grown in SILAC medium and expressing FLAG-CEP78 WT, FLAG-CEP78^{L150S} or

FLAG-Ap80 (negative control) were subjected to FLAG IP and pellets analyzed by mass spectrom-

etry.

Figure 3-source data 2. Original western blots corresponding to Figure 3B. Upper left, input Myc

blot; upper middle, input 100 kDa FLAG blot; upper right, IP Myc and 100 kDa FLAG blots; bot-

tom left, input 35 kDa FLAG blot; bottom right, IP 35 kDa FLAG blot.

Figure 3-source data 3. Original western blots corresponding to Figure 3C. Upper left, input Myc

blot; upper middle, IP Myc blot; upper right, input GFP blot; bottom left, IP GFP blot; bottom mid-

dle, inpot FOP blot; bottom right, IP FOP blot.

Figure 3-source data 4. Original western blots corresponding to Figure 3D. Upper left, input Myc

blot; upper right, IP Myc blot; lower left, input GFP blot; lower right, IP GFP blot.

Figure 4-source data 1. Original western blots for Figure 4B. Left, CEP78 blot; right, α-tubulin

blot.

Figure 6-source data 1. Original western blots for Figure 6A. Left, VPRBP blot; right, α-tubulin

blot.

Figure 7-source data 1. Original western blots for Figure 7A. Left, CP110 blot; right, GAPDH

blot.

Figure 8-source data 1. Original bots for Figure 8A. Left, CP110 blot; right, α-tubulin blot.

Figure 1-figure supplement 1-source data 1. Original western blots for Figure 1-figure supple-

ment 1. Left, CEP78 blot; right, GAPDH blot.

Figure 2-figure supplement 1-source data 1. Original western blots for Figure 2-figure supple-

ment 1. Left, FLAG blot; right, GAPDH blot.

Figure 2-figure supplement 2-source data 1. Original western blots for Figure 2-figure supple-

ment 2. Left, FLAG blot; right, CEP78 blot.

Figure 2-figure supplement 3-source data 1. Original western blots for Figure 2-figure supple-

ment 3. Left, CEP78 blot; right, α-tubulin blot

Figure 3-figure supplement 1-source data 1. Original western blots for Figure 3-figure supple-

ment 1A. Top row from left to right: input VPRBP blot, IP VPRBP blot, input Myc blot. Second

row from left to right: IP Myc blot, input FOP blot, IP FOP blot. Third row from left to right: input

GFP blot (upper), IP GFP blot (upper). Fourth row left to right: input GFP blot (lower), IP GFP blot

(lower).

Figure 3-figure supplement 1-source data 2. Original western blots for Figure 3-figure supple-

ment 1B. Top row, input VPRBP blot; second row, IP VPRBP blot; third row left, input Myc blot;

third row right, IP Myc blot; fourth row left, input GFP blots; fourth row right, IP GFP blots.

Figure 7-figure supplement 1-source data 1. Original western blots for Figure 7-figure supple-

ment 1. Left, CP110 (upper) and GAPDH blots in panel A. Right, CP110 (upper) and α–tubulin

(lower) blots in panel C.

Figure 7-figure supplement 2-source data 1. Original western blots for Figure 7-figure supple-

ment 2A. Top, CP110 blot; bottom, α–tubulin blot.

Figure 7-figure supplement 3-source data 1. Raw RNA seq data for Figure 7-figure supplement

3.

Figure 7-figure supplement 4-source data 1. Original western blots for Figure 7-figure supple-

ment 4A. Upper left, WT CP110 blot; upper right, WT GAPDH blot. Lower left, *CEP78* KO

CP110 blot; lower right, *CEP78* KO GAPDH blot.

1031

References

1. Anvarian, Z., K. Mykytyn, S. Mukhopadhyay, L.B. Pedersen, and S.T. Christensen, *Cellular signalling by primary cilia in development, organ function and disease.* Nat Rev Nephrol, 2019. **15**(4): p. 199-219.
2. Nachury, M.V. and D.U. Mick, *Establishing and regulating the composition of cilia for signal transduction.* Nat Rev Mol Cell Biol, 2019. **20**(7): p. 389-405.
3. Reiter, J.F. and M.R. Leroux, *Genes and molecular pathways underpinning ciliopathies.* Nat Rev Mol Cell Biol, 2017. **18**(9): p. 533-547.
4. Breslow, D.K. and A.J. Holland, *Mechanism and Regulation of Centriole and Cilium Biogenesis.* Annu Rev Biochem, 2019. **88**: p. 691-724.
5. Spektor, A., W.Y. Tsang, D. Khoo, and B.D. Dynlacht, *Cep97 and CP110 suppress a cilia assembly program.* Cell, 2007. **130**(4): p. 678-90.
6. Chen, Z., V.B. Indjeian, M. McManus, L. Wang, and B.D. Dynlacht, *CP110, a cell cycle-dependent CDK substrate, regulates centrosome duplication in human cells.* Dev Cell, 2002. **3**(3): p. 339-50.
7. Kleylein-Sohn, J., J. Westendorf, M. Le Clech, R. Habedanck, Y.D. Stierhof, and E.A. Nigg, *Plk4-induced centriole biogenesis in human cells.* Developmental cell, 2007. **13**(2): p. 190-202.
8. D'Angiolella, V., V. Donato, S. Vijayakumar, A. Saraf, L. Florens, M.P. Washburn, *et al.*, *SCF(Cyclin F) controls centrosome homeostasis and mitotic fidelity through CP110 degradation.* Nature, 2010. **466**(7302): p. 138-42.
9. Li, J., V. D'Angiolella, E.S. Seeley, S. Kim, T. Kobayashi, W. Fu, *et al.*, *USP33 regulates centrosome biogenesis via deubiquitination of the centriolar protein CP110.* Nature, 2013. **495**(7440): p. 255-9.
10. Sorokin, S., *Centrioles and the formation of rudimentary cilia by fibroblasts and smooth muscle cells.* J Cell Biol, 1962. **15**: p. 363-377.
11. Sorokin, S.P., *Reconstructions of centriole formation and ciliogenesis in mammalian lungs.* J Cell Sci, 1968. **3**(2): p. 207-30.
12. Shakya, S. and C.J. Westlake, *Recent advances in understanding assembly of the primary cilium membrane.* Fac Rev, 2021. **10**: p. 16.
13. Kobayashi, T., W.Y. Tsang, J. Li, W. Lane, and B.D. Dynlacht, *Centriolar kinesin Kif24 interacts with CP110 to remodel microtubules and regulate ciliogenesis.* Cell, 2011. **145**(6): p. 914-25.
14. Huang, N., D. Zhang, F. Li, P. Chai, S. Wang, J. Teng, *et al.*, *M-Phase Phosphoprotein 9 regulates ciliogenesis by modulating CP110-CEP97 complex localization at the mother centriole.* Nat Commun, 2018. **9**(1): p. 4511.
15. Cajanek, L. and E.A. Nigg, *Cep164 triggers ciliogenesis by recruiting Tau tubulin kinase 2 to the mother centriole.* Proc Natl Acad Sci U S A, 2014. **111**(28): p. E2841-50.
16. Oda, T., S. Chiba, T. Nagai, and K. Mizuno, *Binding to Cep164, but not EB1, is essential for centriolar localization of TTBK2 and its function in ciliogenesis.* Genes Cells, 2014. **19**(12): p. 927-40.
17. Lo, C.H., I.H. Lin, T.T. Yang, Y.C. Huang, B.E. Tanos, P.C. Chou, *et al.*, *Phosphorylation of CEP83 by TTBK2 is necessary for cilia initiation.* J Cell Biol, 2019. **218**(10): p. 3489-3505.

18.   Graser, S., Y.D. Stierhof, S.B. Lavoie, O.S. Gassner, S. Lamla, M. Le Clech, *et al.*, *Cep164,*
a novel centriole appendage protein required for primary cilium formation. J Cell Biol,
2007. **179**(2): p. 321-30.
19.   Goetz, S.C., K.F. Liem, Jr., and K.V. Anderson, *The spinocerebellar ataxia-associated gene*
Tau tubulin kinase 2 controls the initiation of ciliogenesis. Cell, 2012. **151**(4): p. 847-58.
20.   Schmidt, K.N., S. Kuhns, A. Neuner, B. Hub, H. Zentgraf, and G. Pereira, *Cep164 mediates*
vesicular docking to the mother centriole during early steps of ciliogenesis. J Cell Biol,
2012. **199**(7): p. 1083-101.
21.   Tanos, B.E., H.J. Yang, R. Soni, W.J. Wang, F.P. Macaluso, J.M. Asara, *et al.*, *Centriole*
distal appendages promote membrane docking, leading to cilia initiation. Genes Dev, 2013.
27(2): p. 163-8.
22.   Ye, X., H. Zeng, G. Ning, J.F. Reiter, and A. Liu, *C2cd3 is critical for centriolar distal*
appendage assembly and ciliary vesicle docking in mammals. Proc Natl Acad Sci U S A,
2014. **111**(6): p. 2164-9.
23.   Kurtulmus, B., C. Yuan, J. Schuy, A. Neuner, S. Hata, G. Kalamakis, *et al.*, *LRRC45*
contributes to early steps of axoneme extension. J Cell Sci, 2018. **131**(18).
24.   Kanie, T., K.L. Abbott, N.A. Mooney, E.D. Plowey, J. Demeter, and P.K. Jackson, *The*
CEP19-RABL2 GTPase Complex Binds IFT-B to Initiate Intraflagellar Transport at the
Ciliary Base. Dev Cell, 2017. **42**(1): p. 22-36 e12.
25.   Nishijima, Y., Y. Hagiya, T. Kubo, R. Takei, Y. Katoh, and K. Nakayama, *RABL2 interacts*
with the intraflagellar transport-B complex and CEP19 and participates in ciliary assembly.
Mol Biol Cell, 2017. **28**(12): p. 1652-1666.
26.   Mojarad, B.A., G.D. Gupta, M. Hasegan, O. Goudiam, R. Basto, A.C. Gingras, *et al.*,
CEP19 cooperates with FOP and CEP350 to drive early steps in the ciliogenesis
programme. Open Biol, 2017. **7**(6).
27.   Hossain, D., Y. Javadi Esfehani, A. Das, and W.Y. Tsang, *Cep78 controls centrosome*
homeostasis by inhibiting EDD-DYRK2-DDB1 VprBP. EMBO Rep, 2017. **18**(4): p. 632-
644.
28.   Yoshida, S., K. Aoki, K. Fujiwara, T. Nakakura, A. Kawamura, K. Yamada, *et al.*, *The*
novel ciliogenesis regulator DYRK2 governs Hedgehog signaling during mouse
embryogenesis. Elife, 2020. **9**.
29.   Nikopoulos, K., P. Farinelli, B. Giangreco, C. Tsika, B. Royer-Bertrand, M.K. Mbefo, *et al.*,
Mutations in CEP78 Cause Cone-Rod Dystrophy and Hearing Loss Associated with
Primary-Cilia Defects. Am J Hum Genet, 2016. **99**(3): p. 770-776.
30.   Ascari, G., F. Peelman, P. Farinelli, T. Rosseel, N. Lambrechts, K.A. Wunderlich, *et al.*,
Functional characterization of the first missense variant in CEP78, a founder allele
associated with cone-rod dystrophy, hearing loss, and reduced male fertility. Hum Mutat,
2020. **41**(5): p. 998-1011.
31.   Namburi, P., R. Ratnapriya, S. Khateb, C.H. Lazar, Y. Kinarty, A. Obolensky, *et al.*, *Bi-*
allelic Truncating Mutations in CEP78, Encoding Centrosomal Protein 78, Cause Cone-
Rod Degeneration with Sensorineural Hearing Loss. Am J Hum Genet, 2016. **99**(5): p.
1222-1223.
32.   Fu, Q., M. Xu, X. Chen, X. Sheng, Z. Yuan, Y. Liu, *et al.*, *CEP78 is mutated in a distinct*
type of Usher syndrome. J Med Genet, 2017. **54**(3): p. 190-195.
33.   de Castro-Miro, M., R. Tonda, P. Escudero-Ferruz, R. Andres, A. Mayor-Lorenzo, J. Castro,
et al., *Novel Candidate Genes and a Wide Spectrum of Structural and Point Mutations*
Responsible for Inherited Retinal Dystrophies Revealed by Exome Sequencing. PLoS One,
2016. **11**(12): p. e0168966.

34.   Azimzadeh, J., M.L. Wong, D.M. Downhour, A. Sanchez Alvarado, and W.F. Marshall,
1126   *Centrosome loss in the evolution of planarians.* Science, 2012. **335**(6067): p. 461-3.
1127   35.   Brunk, K., M. Zhu, F. Barenz, A.S. Kratz, U. Haselmann-Weiss, C. Antony, *et al.*, *Cep78 is*
1128   *a new centriolar protein involved in Plk4-induced centriole overduplication.* J Cell Sci,
1129   2016. **129**(14): p. 2713-8.
1130   36.   Knudsen, E.S. and J.Y. Wang, *Dual mechanisms for the inhibition of E2F binding to RB by*
1131   *cyclin-dependent kinase-mediated RB phosphorylation.* Mol Cell Biol, 1997. **17**(10): p.
1132   5771-5783.
1133   37.   Paridaen, J.T., M. Wilsch-Brauninger, and W.B. Huttner, *Asymmetric inheritance of*
1134   *centrosome-associated primary cilium membrane directs ciliogenesis after cell division.*
1135   Cell, 2013. **155**(2): p. 333-44.
1136   38.   Yan, X., R. Habedanck, and E.A. Nigg, *A complex of two centrosomal proteins, CAP350*
1137   *and FOP, cooperates with EB1 in microtubule anchoring.* Mol Biol Cell, 2006. **17**(2): p.
1138   634-644.
1139   39.   Wang, L., M. Failler, W. Fu, and B.D. Dynlacht, *A distal centriolar protein network*
1140   *controls organelle maturation and asymmetry.* Nat Commun, 2018. **9**(1): p. 3938.
1141   40.   Paintrand, M., M. Moudjou, H. Delacroix, and M. Bornens, *Centrosome organization and*
1142   *centriole architecture: their sensitivity to divalent cations.* J Struct Biol, 1992. **108**(2): p.
1143   107-128.
1144   41.   Kim, J., J.E. Lee, S. Heynen-Genel, E. Suyama, K. Ono, K. Lee, *et al.*, *Functional genomic*
1145   *screen for modulators of ciliogenesis and cilium length.* Nature, 2010. **464**(7291): p. 1048-
1146   51.
1147   42.   Shearer, R.F., K.M. Frikstad, J. McKenna, R.A. McCloy, N. Deng, A. Burgess, *et al.*, *The*
1148   *E3 ubiquitin ligase UBR5 regulates centriolar satellite stability and primary cilia.* Mol Biol
1149   Cell, 2018. **29**(13): p. 1542-1554.
1150   43.   Tsang, W.Y. and B.D. Dynlacht, *CP110 and its network of partners coordinately regulate*
1151   *cilia assembly.* Cilia, 2013. **2**(1): p. 9.
1152   44.   Yadav, S.P., N.K. Sharma, C. Liu, L. Dong, T. Li, and A. Swaroop, *Centrosomal protein*
1153   *CP110 controls maturation of the mother centriole during cilia biogenesis.* Development,
1154   2016. **143**(9): p. 1491-501.
1155   45.   Walentek, P., I.K. Quigley, D.I. Sun, U.K. Sajjan, C. Kintner, and R.M. Harland, *Ciliary*
1156   *transcription factors and miRNAs precisely regulate Cp110 levels required for ciliary*
1157   *adhesions and ciliogenesis.* Elife, 2016. **5**.
1158   46.   Loukil, A., K. Tormanen, and C. Sutterlin, *The daughter centriole controls ciliogenesis by*
1159   *regulating Neurl-4 localization at the centrosome.* J Cell Biol, 2017. **216**(5): p. 1287-1300.
1160   47.   Nagai, T., S. Mukoyama, H. Kagiwada, N. Goshima, and K. Mizuno, *Cullin-3-KCTD10-*
1161   *mediated CEP97 degradation promotes primary cilium formation.* J Cell Sci, 2018. **131**(24).
1162   48.   Wheway, G., M. Schmidts, D.A. Mans, K. Szymanska, T.T. Nguyen, H. Racher, *et al.*, *An*
1163   *siRNA-based functional genomics screen for the identification of regulators of ciliogenesis*
1164   *and ciliopathy genes.* Nat Cell Biol, 2015. **17**(8): p. 1074-1087.
1165   49.   Nakagawa, T., K. Mondal, and P.C. Swanson, *VprBP (DCAF1): a promiscuous substrate*
1166   *recognition subunit that incorporates into both RING-family CRL4 and HECT-family*
1167   *EDD/UBR5 E3 ubiquitin ligases.* BMC Mol Biol, 2013. **14**: p. 22.
1168   50.   Patzke, S., S. Redick, A. Warsame, C.A. Murga-Zamalloa, H. Khanna, S. Doxsey, *et al.*,
1169   *CSPP is a ciliary protein interacting with Nephrocystin 8 and required for cilia formation.*
1170   Molecular biology of the cell, 2010. **21**(15): p. 2555-67.

51.   Frikstad, K.M., E. Molinari, M. Thoresen, S.A. Ramsbottom, F. Hughes, S.J.F. Letteboer, *et*
al., *A CEP104-CSPP1 Complex Is Required for Formation of Primary Cilia Competent in*
Hedgehog Signaling. Cell Rep, 2019. **28**(7): p. 1907-1922 e6.
52.   Jiang, K., G. Toedt, S. Montenegro Gouveia, N.E. Davey, S. Hua, B. van der Vaart, *et al.*, *A*
Proteome-wide screen for mammalian SxIP motif-containing microtubule plus-end tracking
proteins. Curr Biol, 2012. **22**(19): p. 1800-7.
53.   Satish Tammana, T.V., D. Tammana, D.R. Diener, and J. Rosenbaum, *Centrosomal protein*
CEP104 (Chlamydomonas FAP256) moves to the ciliary tip during ciliary assembly. J Cell
Sci, 2013. **126**(Pt 21): p. 5018-29.
54.   Eguether, T., M.A. Ermolaeva, Y. Zhao, M.C. Bonnet, A. Jain, M. Pasparakis, *et al.*, *The*
deubiquitinating enzyme CYLD controls apical docking of basal bodies in ciliated epithelial
cells. Nat Commun, 2014. **5**: p. 4585.
55.   Wang, L., K. Lee, R. Malonis, I. Sanchez, and B.D. Dynlacht, *Tethering of an E3 ligase by*
PCM1 regulates the abundance of centrosomal KIAA0586/Talpid3 and promotes
ciliogenesis. Elife, 2016. **5**.
56.   Douanne, T., G. Andre-Gregoire, A. Thys, K. Trillet, J. Gavard, and N. Bidere, *CYLD*
Regulates Centriolar Satellites Proteostasis by Counteracting the E3 Ligase MIB1. Cell
Rep, 2019. **27**(6): p. 1657-1665 e4.
57.   Yi, C., Z. Shen, A. Stemmer-Rachamimov, N. Dawany, S. Troutman, L.C. Showe, *et al.*,
The p130 isoform of angiomotin is required for Yap-mediated hepatic epithelial cell
proliferation and tumorigenesis. Sci Signal, 2013. **6**(291): p. ra77.
58.   Hoppeler-Lebel, A., C. Celati, G. Bellett, M.M. Mogensen, L. Klein-Hitpass, M. Bornens, *et*
al., *Centrosomal CAP350 protein stabilises microtubules associated with the Golgi*
complex. J Cell Sci, 2007. **120**(Pt 18): p. 3299-308.
59.   Belzile, J.P., G. Duisit, N. Rougeau, J. Mercier, A. Finzi, and E.A. Cohen, *HIV-1 Vpr-*
mediated G2 arrest involves the DDB1-CUL4AVPRBP E3 ubiquitin ligase. PLoS Pathog,
2007. **3**(7): p. e85.
60.   Cox, J. and M. Mann, *MaxQuant enables high peptide identification rates, individualized*
p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol,
2008. **26**(12): p. 1367-72.
61.   Schou, K.B., J.B. Mogensen, A. Aleliunaite, S.K. Morthorst, I.R. Veland, S.T. Christensen,
et al., *Kinesin-3 motor protein KIF13B localizes to centrosomes and primary cilia and*
regulates ciliary length and signaling. Cilia, 2014. **4**(Suppl 1): p. O3.
62.   Verdier, P., S.K. Morthorst, and L.B. Pedersen, *Targeting of ASH Domain-Containing*
Proteins to the Centrosome. Methods Mol Biol, 2016. **1454**: p. 15-33.
63.   Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nat Methods,
2012. **9**(4): p. 357-9.
64.   Kim, D., G. Pertea, C. Trapnell, H. Pimentel, R. Kelley, and S.L. Salzberg, *TopHat2:*
accurate alignment of transcriptomes in the presence of insertions, deletions and gene
fusions. Genome Biol, 2013. **14**(4): p. R36.
65.   Anders, S., P.T. Pyl, and W. Huber, *HTSeq--a Python framework to work with high-*
throughput sequencing data. Bioinformatics, 2015. **31**(2): p. 166-9.
66.   Trapnell, C., B.A. Williams, G. Pertea, A. Mortazavi, G. Kwan, M.J. van Baren, *et al.*,
Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and
isoform switching during cell differentiation. Nat Biotechnol, 2010. **28**(5): p. 511-5.
67.   Robinson, M.D., D.J. McCarthy, and G.K. Smyth, *edgeR: a Bioconductor package for*
differential expression analysis of digital gene expression data. Bioinformatics, 2010. **26**(1):
p. 139-40.

68.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion*
for RNA-seq data with DESeq2. Genome Biol, 2014. **15**(12): p. 550.
1221

1222   **Appendix**

1223   **Key Resources Table**

Reagent type (species) or resource	Designation	Source or reference	Identifier	Additional information
Cell line (*H. sapiens*)	hTERT-RPE1	ATCC	Cat# CRL-4000	Derived from human retinal pigment epithelium
Cell line (*H. sapiens*)	Puromycin-sensitive hTERT-RPE1	Francesc Garcia-Gonzalo, Autonomous University of Madrid, Madrid, Spain	N/A	
Cell line (*H. sapiens*)	hTERT-RPE1 *CEP78* KO	[30]	Clone #73	
Cell line (*H. sapiens*)	hTERT-RPE1 *CEP78* KO	This study	Clone #44	Generated as described in [30]; contains px459-Cas9 plasmid inserted in exon 1
Cell line (*H. sapiens*)	hTERT-RPE1 *CEP78* KO	This study	Clone #52	Generated as described in [30]; contains homozygous frameshift mutation in exon 1 of *CEP78* (single nucleotide insertion in codon 10; GAC>GAAC)

Cell line (*H. sapiens*)	hTERT-RPE1 *CEP78* KO	This study	Clone #2	Generated as described in [30]; homozygous frameshift mutation in exon 1 of *CEP78* (single nucleotide insertion in codon 10; GAC>GAAC)
Cell line (*H. sapiens*)	hTERT-RPE1 *CEP78* KO/mNG-CEP78	This study	N/A	Parental line is *CEP78* KO clone #73
Cell line (*H. sapiens*)	hTERT-RPE1 *CEP78* KO/mNG-CEP78^{L150S}	This study	N/A	Parental line is *CEP78* KO clone #73
Cell line (*H. sapiens*)	hTERT-RPE1 WT/mNG-CEP78	This study	N/A	Parental line is puromycin-sensitive hTERT-RPE listed above
Cell line (*H. sapiens*)	hTERT-RPE1-BFP-Cas9/pMCB306 (pool)	[24]	N/A	Control cell line for *CEP350* and *FOP* KO lines
Cell line (*H. sapiens*)	hTERT-RPE1-BFP-Cas9/ *CEP350* KO (pool)	[24]	N/A	
Cell line (*H. sapiens*)	hTERT-RPE1-BFP-Cas9/*FOP* KO (pool)	[24]	N/A	
Cell line (*H. sapiens*)	293T	ATCC	Cat# CRL-3216	Derived from human embryonic kidney
Cell line (*H. sapiens*)	Control skin fibroblasts	[29]	N/A	Primary cells
Cell line (*H. sapiens*)	Compound heterozygous skin fibroblasts with two *CEP78* mutations: c.449T>C; p.Leu150Ser and c.1462-1G>T	[30]	F3: II:1	Primary cells; these cells do not express detectable levels of CEP78 [30]

Cell line (*H. sapiens*)	Compound heterozygous skin fibroblasts with two *CEP78* mutations: c.633delC; p.Trp212Glyfs*18 and c.499+5G>A; IVS + 5G>A	[29]	2702r34, individual II-3	Primary cells; these cells do not express detectable levels of CEP78 [29]
Cell line (*H. sapiens*)	Homozygous substitution in the first invariant base of intron 3 splice donor site in *CEP78*: c.499+1G>T (IVS3+1G>T)	[29]	KN10, individual II-1	Primary cells; these cells express almost undetectable levels of CEP78 [29]
Strain, strain background (*E. coli*)	DH10B	Lab stock	N/A	
Antibody	Anti-Acetylated-tubulin (mouse monoclonal)	Sigma-Aldrich	Cat# T7451	IFM (1:2000)
Antibody	Anti-α-tubulin (mouse monoclonal)	Sigma-Aldrich	Cat# T5168	WB (1:10000)
Antibody	Anti-ARL13B (rabbit polyclonal)	Proteintech	Cat# 17711-1-AP	IFM (1:500)
Antibody	Anti-CEP78 (rabbit polyclonal)	Bethyl Laboratories	Cat# A301-799A-M	IFM (1:200) WB (1:500)
Antibody	Anti-CEP164 (rabbit polyclonal)	Sigma-Aldrich	Cat# HPA037606	IFM (1:500)
Antibody	Anti-CEP350 (mouse monoclonal)	Abcam	Cat# Ab219831 (CL3423)	IFM (1:500)
Antibody	Anti-CEP350 (rabbit polyclonal)	Novus Biologicals	Cat# NB100-59811	IFM (1:200)
Antibody	Anti-CP110 (rabbit polyclonal)	Proteintech	Cat# 12780-1-AP	IFM (1:200) WB (1:1500)

Antibody	Anti-DCTN1/P-150 (mouse monoclonal)	BD Biosciences	Cat# 610474	IFM (1:500)
Antibody	Anti-EDD1 (rabbit polyclonal)	Bethyl Laboratories	Cat# A300-573A-M	IFM (1:100)
Antibody	Anti-FLAG (mouse monoclonal)	Sigma-Aldrich	Cat# F1804	IFM (1:500) WB (1:1000)
Antibody	Anti-FLAG (rabbit polyclonal)	Sigma-Aldrich	Cat# F7425	WB (1:1000)
Antibody	Anti-FGFR1OP/FOP (mouse monoclonal)	Novus Biologicals	Cat# H00011116-M01	WB (1:500)
Antibody	Anti-GAPDH (rabbit polyclonal)	Cell Signaling Technology	Cat# 2118	WB (1:1000)
Antibody	Anti-GFP (rabbit polyclonal)	Sigma-Aldrich	Cat# SAB4301138	WB (1:500)
Antib0dy	Anti-GFP (rabbit polyclonal)	Santa Cruz Biotechnologies, Inc.	Cat# sc-8834	WB (1:500)
Antibody	Anti-Myc (mouse monoclonal, 9B11)	Cell Signaling Technology	Cat# 2276S	WB (1:1000)
Antibody	Anti-phospho Rb (S807/811) (mouse monoclonal)	Cell Signaling Technology	Cat# 9308	WB (1:500)
Antibody	Anti-VPRBP (rabbit polyclonal)	Bethyl Laboratories	Cat# A301-877A-M	WB (1:200)
Antibody	Polyclonal Goat Anti-Mouse Immunoglobulins /Horseradish Peroxidase conjugated	Dako	Cat# P0447	WB (1:10000)
Antibody	Polyclonal Swine Anti-Rabbit Immunoglobulins /Horseradish Peroxidase conjugated	Dako	Cat# P0399	WB (1:10000)

Antibody	Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody/Alexa Fluor 488-conjugated	Thermo Fisher Scientific	Cat# A21202	IFM (1:500)
Antibody	Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody/Alexa Fluor 568-conjugated	Thermo Fisher Scientific	Cat # A10037	IFM (1:500)
Antibody	Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody/Alexa Fluor 488-conjugated	Thermo Fisher Scientific	Cat #A21206	IFM (1:500)
Antibody	Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody/Alexa Fluor 568-conjugated	Thermo Fisher Scientific	Cat# A10042	IFM (1:500)
Antibody	AffiniPure Don-key Anti-Mouse IgG (H+L)/Alexa Fluor® 647-conjugated	Jackson ImmunoResearch	Cat# 715-605-150	IFM (1:1000)
Antibody	Donkey Anti-Rabbit IgG H&L Secondary Anti-bodies/ Dylight® 550-conjugated	Abcam	Cat# ab96892	IFM (1:1000)
Commercial assay, kit	USER® Enzyme	New England BioLabs	Cat# M5505L	
Commercial assay, kit	Phusion® High-Fidelity DNA polymerase	New England BioLabs	Cat# M0530L	

Commercial assay, kit	QuickChange Lightning Site-Directed Mutagenesis Kit	Agilent	Cat# 210518	
Commercial assay, kit	Direct-zol RNA Miniprep Kit	Zymo Research	Cat# R2051	
Commercial assay, kit	Agilent RNA Nano 6000 Assay Kit	Agilent	Cat# 5067-1511	
Commercial assay, kit	NEBNext® Ultra™ RNA Library Prep Kit for Illumina®	New England BioLabs	Cat# E7530L	
Commercial assay, kit	Invitrogen™ Qubit® RNA Assay Kit	Thermo Fisher Scientific	Cat# 32852	
Recombinant DNA reagent	*H. sapiens* CEP78 aa 2-217/pEGFP-C1	This study	CEP78 N-terminus (N)	PCR-amplified using primers h.CEP78.4f.kpn1 and hsEGFP-CEP78_Rv (651)
Recombinant DNA reagent	*H. sapiens* CEP78 aa 203-403/pEGFP-C1	This study	CEP78 Middle (M)	PCR-amplified using primers hsEGFP-CEP78_Fwd (607) and hsEGFP-CEP78_Rv (1209)
Recombinant DNA reagent	*H. sapiens* CEP78 aa 2-403/pEGFP-C1	This study	CEP78 N-terminus plus Middle (NM)	PCR-amplified using primers h.CEP78.4f.kpn1 and hsEGFP-CEP78_Rv (1209)
Recombinant DNA reagent	*H. sapiens* CEP78 aa 395-722/pEGFP-C1	This study	CEP78 C-terminus (C)	PCR-amplified using primers hsEGFP-CEP78_Fwd (1186) and hCEP78.rv_stop.bamh1

Recombinant DNA reagent	*H. sapiens* CEP78/pEGFP-C1	This study	N/A	PCR-amplified using primers h.CEP78.4f.kpn1 and hCEP78.rv_stop.bamh1
Recombinant DNA reagent	*H. sapiens* CEP78/p3xFLAG-CMV/DEST	This study	p3xFLAG-CEP78	
Recombinant DNA reagent	*H. sapiens* CEP78^{L150S}/p p3xFLAG-CMV/DEST	This study	p3xFLAG-CEP78^{L150S}	
Recombinant DNA reagent	Plasmid pEGFP-C1 (empty vector)	TaKaRa Bio	Cat# 6084-1	
Recombinant DNA reagent	Gateway® pDONR^{TM}201	Invitrogen/Thermo Fisher Scientific	Cat# 11798-014	
Recombinant DNA reagent	p3xFLAG-CMV-10	Sigma-Aldrich	Cat# E7658	Cloning produces in-frame 3xFLAG tag in the N-terminus
Recombinant DNA reagent	p3xFLAG-CMV/DEST	Gift from Ronald Roepman lab, Radboud University Medical Center, Nijmegen, NL	pDEST306	p3xFLAG-CMV-10 with Gateway Rf-B cassette inserted into the blunted HindIII/BamHI site
Recombinant DNA reagent	*H. sapiens* CEP350 aa 1-983/pCS2^{+} with a 6×myc tag	[54, 58]	CAP N	
Recombinant DNA reagent	*H. sapiens* CEP350 aa 2990-3117/pCS2^{+} with a 6×myc tag	[54, 58]	CAP C	
Recombinant DNA reagent	*H. sapiens* VPRBP/pCMV-Myc	[59]	Myc-VPRBP	Cloned in Sal1 and Not1 sites
Sequence-based reagent	*H. sapiens* *CEP78* ORF/ GeneCopoiea pShuttle™Gateway® PLUS ORF	GeneCopoiea	NCBI reference Sequence NM_001098802.3	

Sequence-based reagent	*H. sapiens CEP78* PCR primer	Sigma-Aldrich	attB forward	5'-GGGACAAGTTGTACAAAAAACAGGCTTCATCGACTCCGTGAAGCTGC-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Sigma-Aldrich	attB reverse	5-GGGGACCACTTTGTACAAGAAAGCTGGGTTTCAGGAATGCAGGTCCTTTCCAG-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Sigma-Aldrich	CEP78 mut_F	5'-AGAGACAGGTGCACCGAAGAAGCCGATTTATTCAATC-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Sigma-Aldrich	CEP78 mut_R	5'-GATTGAATAAATCGGCTTCTTCGGTGCACCTGTCTCT-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Eurofins Genomics	hCEP78.4f.kpn1	5'-AAGGTACCATCGACTCCGTGAAGCTGCG-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Eurofins Genomics	hsEGFP-CEP78_Rv (651)	5'-AAGGATCCTTAGCGAAGACTCTCAGCCCAG-3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Eurofins Genomics	hsEGFP-CEP78_Fwd (607)	5'- GGGGTACCCAGACCATGAGAAGGCATGA -3'
Sequence-based reagent	*H. sapiens CEP78* PCR primer	Eurofins Genomics	hsEGFP-CEP78_Rv (1209)	5'-GGGGATCCTTAACCCCTGTGTCTTTTTGCAC-3'

Sequence-based reagent	*H. sapiens* CEP78 PCR primer	Eurofins Genomics	hsEGFP-CEP78_Fwd (1186)	5'-GGGGTACC GCAGAACGTG CAAAAAGACA -3'
Sequence-based reagent	*H. sapiens* CEP78 PCR primer	Eurofins Genomics	hCEP78.rv_stop. bamh1	5'-GAGGATCC ACAGGAATGC AGGTCCTTTC-3'
Sequence-based reagent	Control siRNA	Eurofins Genomics	N/A	5'-UAAUGUAUU GGAAGGCA-3'
Sequence-based reagent	*H. sapiens* CP110 siRNA	Eurofins Genomics /[5]	N/A	5'-GCAAAACCAG AAUACGAGAT T-3'
Chemical compound, drug	DAPI (4′, 6-diamidino-2-phenylindole)	Thermo Fisher Scientific	Cat# D1306	Stored at -20^{o}C as stock solution of 20 µg/ml in H$_2$O
Chemical compound drug	FuGENE®6	Promega	Cat# E2692	
Chemical compound, drug	DharmaFECT^{TM} Duo Transfection reagent	Dharmacon^{TM}	Cat# T-2010-03	
Software, algorithm	cellSens Dimension	Olympus	Version 1.18	
Software, algorithm	Zen Black 2012	Zeiss	Version 2012	
Software, algorithm	Adobe Photoshop®	Adobe	Version 21.0.1	
Software, algorithm	Adobe Illustrator®	Adobe	Version 24.2.1	
Software, algorithm	ImageJ	NIH	Mac OS X or Windows	
Software, algorithm	GraphPad Prism 6.0	GraphPad Software Inc.	Mac OS X or Windows	
Software, algorithm	MaxQuant	[60]	Version 1.6.1.0	
Other	Anti-FLAG® M2 Affinity Gel	Sigma-Aldrich	Cat# A2220	
Other	Anti-c-Myc Agarose Affinity Gel antibody produced in rabbit	Sigma-Aldrich	Cat# A7470-1 ml	

Other	GFP-Trap® Agarose	ChromoTek GmbH	Cat# gta-20	
Other	AMPure XP system	Beckman Coulter Life Sciences	Cat# A63881	

1224