Lexical parafoveal previewing predicts reading speed

Yali Pan[1,2], Steven Frisson[1,2] and Ole Jensen[1,2*]

[1] Centre for Human Brain Health, [2] School of Psychology, University of Birmingham, Birmingham B15 2TT, United Kingdom

*Correspondence: o.jensen@bham.ac.uk

Abstract

While reading is an essential human skill, many of the neuronal mechanisms supporting reading are not well understood. In spite of the reduced visual acuity, parafoveal information plays a critical role in natural reading[1]; however, it is strongly debated whether words are previewed parafoveally at the lexical level[2–9]. This is a key dispute for competing models on reading and of great significance since lexical retrieval is important for guiding eye movements. We found evidence for parafoveal previewing at the lexical level by combining a novel *rapid invisible frequency tagging* (RIFT)[10,11] approach with magnetoencephalography (MEG) and eye-tracking. In a silent reading task, target words of either low or high lexical frequency were tagged (flickered) subliminally at 60 Hz. The tagging response measured when fixating on the pre-target word reflected parafoveal previewing of the target word. We observed a stronger tagging response as early as 100 ms following fixations on pre-target words that were followed by target words of low compared with high lexical frequency. Moreover, the difference in tagging response with respect to target lexicality predicted individual reading speed. Our findings demonstrate that reading unfolds in the fovea and parafovea simultaneously to support fluent reading.

Main

Humans have developed the remarkable skill of reading, allowing for efficient acquisition of information from busy pages or screens of text. Given the importance of written text for communication, individuals with reading disabilities are highly disadvantaged in modern society. Yet, we know little about the neuronal mechanism underlying natural reading. Our study aimed at answering if lexical information is retrieved for upcoming words in the parafovea (i.e., 2 to 5 degrees from the current fixation) during natural reading. We used a novel technique combining *rapid invisible frequency tagging* (RIFT) with magnetoencephalography (MEG) to provide neural insights on this question. In the present study, participants read 228 sentences in total (composed of two sets of sentences). All sentences were plausible and contained unpredictable target words of either low or high lexical frequency (see Supplementary information for sentence details). Word length for both pre-target and target words were matched with respect to low and high lexical frequency of the target words (Supplementary Table 1). Target words were flickered at 60 Hz throughout the reading of each sentence while the neuronal activity was measured by MEG (Fig. 1). When participants fixated on the pre-target word, the flickering target induced a reliable tagging response at 60 Hz, reflecting the neuronal resources associated with parafoveal previewing. Thus, this experimental design allowed us to investigate neural activity associated with lexical parafoveal previewing without interfering with natural reading.

Figure 1 | The reading task. Participants (n = 39) read sentences silently at their own pace, while eye-movements and brain activity were recorded. Each sentence was embedded with either a low or high lexical frequency target word (see dashed rectangle; not shown in the experiment). A Gaussian smoothed patch below the target word was flickered at 60 Hz continuously while the sentence was on the screen. This allowed us to measure the associated neuronal response using MEG: rapid invisible frequency tagging (RIFT). One-quarter of the sentences were followed by a simple yes-or-not comprehension question to ensure that participants read the sentences carefully.

In line with previous studies[2,4,5,8,9], we did not find an effect of target word lexical frequency

on pre-target first fixation durations (i.e. the duration of the *first* fixation on a word) ($t_{(38)}$ =

0.17, p = 0.86, d = 0.03, two-tailed pairwise t-test, Fig. 2a). However, the target fixation

durations were longer for low compared with high lexical frequency targets ($t_{(38)}$ = 6.94, p =

3×10^{-8}, d = 1.11, two-tailed pairwise t-test). This classic word frequency effect indicates the

successful manipulation of target lexical frequency. We observed the same pattern using gaze

durations (i.e. *total* duration of first-pass fixations on a word; Supplementary Fig. 1). These

findings demonstrate that parafoveal lexical processing is not reflected in the eye movement

data, at least in the well-controlled experimental studies.

Next, we analyzed MEG data to uncover the brain activity associated with lexical processing

before saccading to the target. In previous researches on spatial attention, we demonstrated

that RIFT captured covert attention, reflected by stronger tagging responses for attended

compared with unattended stimuli[10,11]. Here we hypothesized that lexical parafoveal

previewing would be observed as a modulation in the tagging response with respect to target

lexical frequency during pre-target fixation. A measure of time-resolved coherence (r^2)

between the 60 Hz visual flicker and the brain activity was used to quantify the tagging

responses (see Methods for details).

We found a strong tagging response over the left visual cortex sensors (Fig. 2b), reflecting

the neuronal resources associated with parafoveal previewing. This was observed as a robust

60 Hz visual flicker-to-MEG coherence during pre-target fixations (caused by the target

flickering in the parafovea) as compared with a baseline period (cross-fixation presented

before sentence onset). In 26 out of the 39 participants, one or more sensors had a significant

tagging response (5.4 ± 4.0 sensors per participant, mean ± SD). The subsequent analyses

were based on these participants and sensors. A source modelling approach revealed that the

generators of the coherence were localized in the early visual cortex (Brodmann area 17, 18;

Fig 2c). The time course of the 60 Hz coherence is shown in Fig. 2d. Note the robust 60 Hz

coherence from the target word when fixating on the pre-target word. These results

demonstrate that RIFT is a sensitive tool for measuring brain activity associated with

parafoveal previewing during natural reading.

Figure 2 | Eye-movement metrics and tagging responses. (**a**) The first fixation duration
difference for pre-target and target words when comparing low- versus high lexical frequency
target words. There was no difference in fixation times on pre-target words with respect to
target lexicality (left). The fixation times were however longer for low than high lexical
frequency target words (*** $p < 0.001$, n = 39, two-tailed paired t-test). The horizontal bar in
violin plot indicates the mean value; each dot represents one participant. (**b**) Topography for
sensors from all participants that showed stronger tagging responses during the pre-target
period (flicker) compared with the baseline period (no-flicker). Twenty-six participants
showed a robust RIFT response. The tagging response was quantified as the coherence
between the tagging signal and the MEG signal. (**c**) Source modelling based on a beam-
forming approach identified the neural sources of tagging response to the early visual cortex
(Brodmann area 17, 18). (**d**) The time-course of neuronal tagging response (coherence)
during the pre-target (left panel), target (middle panel), and post-target fixation periods (right
panel, n = 26). The analysis was time-locked to fixations onsets (white vertical lines; the grey
vertical lines indicated the average fixation offsets). The strong coherence in the pre-target
fixation interval captures brain activity associated with parafoveal target processing.

Next, we addressed if the RIFT was modulated by the lexical frequency of the target word.

Our key finding was that the coherence at 60 Hz in the pre-target fixation was significantly

stronger when followed by a low compared with a high lexical frequency target word (Fig.

3a; $t_{(25)} = 2.20$, $p = 0.037$, $d = 0.43$, two-tailed pairwise t-test). To ensure that the coherence in

pre-target fixation was not contaminated by temporal smoothing from target fixation, the time

window for averaging coherence was adjusted individually according to the shortest pre-

target fixation duration (88.3 ± 8.9 ms across participants, mean $\pm$ SD). Furthermore, as the

number of trials biases coherence magnitude, we subsampled the same number of trials for

both conditions in each participant (by randomly selecting trials from the condition with more

trials). In sum, we found evidence for a stronger neuronal response for low compared with

high lexical frequency target words in the interval when participants fixated on the pre-target

word. This finding supports lexical extraction in the parafovea.

Figure 3 | Neuronal activity associated with parafoveal processing. (**a**) Time-frequency
representation of coherence differences during pre-target word fixations (low- minus high
lexical frequency target words; time-locked to fixation onsets; n = 26). (**b**) The pre-target
coherence at 60 Hz during pre-target word fixations for low- (blue) and high lexical
frequency (orange) target words (n = 26, violin horizontal bar indicates mean value). The
coherence was significantly stronger for low- compared with high lexical frequency target
words (* $p = 0.037$; two-tailed paired t-test). (**c,d**) During target word fixations, the
coherence was similar for low and high lexical frequency target words.

Did the lexical frequency of the target word affect onset latency for parafoveal previewing?

The pre-target coherence onset latency was defined by the time to reach its half maximum

(see Supplementary information). We then used a jackknife procedure to statistically evaluate

if the coherence onset was significantly different for low compared with high lexical

frequency target words. While no statistical difference was found for the combined sets of

sentences (Supplementary Fig. 2a left panel, $t_{(25)}$ = -1.69, p > 0.1, two-tailed pairwise t-test),

we did however find a significant difference for the set with shorter pre-target and target

words (2nd sentences set; see Supplementary Fig. 2c left panel, $t_{(25)}$ = -2.85, p < 0.01, two-

tailed pairwise t-test). Importantly, this effect was already visible around the first 100ms of

the pre-target fixation (72ms for low lexical frequency target; 116ms for high lexical

frequency target). The null finding for the combined sentence set might be explained by the

longer and more varying length of the pre-target and target words (Supplementary Fig. 2b).

The faster onset for the low lexical frequency target indicates that parafoveal neuronal

processing is modulated in time by upcoming lexical information as well, but this effect is

only visible when pre-target and target words are relatively short.

Did lexical parafoveal previewing facilitate reading? To address this, we correlated the

coherence difference with respect to lexical frequency of the target word when fixating the

pre-target word with individual reading speed. Reading speed was quantified as the number

of words read per second (calculated as the total number of words in a sentence divided by

the respective sentence reading time). We found a positive correlation (Fig. 4; $r_{(25)}$ = 0.45, p =

0.022, Spearman correlation), indicating that participants who captured more lexical

information in the parafovea were also faster readers.

Figure 4 | Relation between lexical parafoveal processing difference and individual
reading speed. The coherence-difference with respect to target lexical frequency during
fixations on pre-target words were derived per participant (see **Fig. 3b**). The reading speed
was quantified as the number of words read per second. Each dot represents a participant. A
Spearman correlation demonstrated a significant relation (n = 26). The shaded area indicates

a 95% confidence interval. Histograms of individual reading speed and pre-target coherence
differences are shown on the top and to the right.
159

In addition to the main analyses above, we also carried out two further analyses to rule out
alternative explanations for our effects. First, we asked if the lexical parafoveal effect on the
pre-target was coming from the target word but was smeared to the pre-target word (Fig. 3c).
Therefore, we tested whether the lexical frequency effect could also be observed during target
fixations. As before, the averaging time window for the coherence was the minimum target
fixation duration for each participant (87.1 ± 9.4 ms, mean $\pm$ SD). No significant coherence
difference was found when fixating on low versus high lexical frequency target words (Fig.
3d; $t_{(25)} = 0.01$, $p = 0.992$, $d = 0.002$, two-tailed pairwise t-test). Presumably, the finding that
the flickering did not affect target word viewing might be related to the fact that flicker
sensitive photoreceptors (rod cells) mainly exist in the parafovea of the retina[12]. As a
consequence, there would be reduced sensitivity to flickering when the target word is in
foveal vision.

Next, we performed an initial landing position analysis to exclude the possibility that the
lexical parafoveal effect was due to mislocated fixations or oculomotor error. If one fixation
was intended for the target word, but the saccade undershot and landed on the pre-target
word, then the lexical frequency effect of the target word should have been observed for the
eye movements to the pre-target word. This was not the case in our data as no target lexical
frequency effect was found for pre-target fixation durations (Fig. 2a). Furthermore, we
showed that saccade landings on the pre-target words were highly accurate (-1.1 ± 1.9 letters
with respect to the centre of the pre-target word, mean $\pm$ SD) and were similarly distributed
as saccades to all words (Supplementary Fig. 3). The lexical frequency of the target word also
did not affect the pre-target landing position ($t_{(25)} = 0.35$, $p = 0.727$, $d = 0.07$, two-tailed
pairwise t-test). In addition, the coherence during the pre-target word fixations did not
correlate with the landing position on the pre-target word ($r_{(25)} = 0.16$, $p = 0.420$; Spearman
correlation).

Our findings shed new light on the long-standing debate between models arguing for either
serial attention shift[13,14] or parallel graded processing [15,16]. Both models regard spatial
attention as important for reading but they differ on whether more than one word can be
accessed simultaneously at the lexical level. Parallel graded models predict that the lexical
frequency of both foveal and parafoveal words can be processed simultaneously. This idea is

challenged by the fact that only a few correlation-based corpus studies found lexical

parafoveal effects in eye movement data[6,7], while more well-controlled experimental studies

did not[2,4,5] (and neither did our eye movement results, Fig. 2a). But these conclusions are

based on eye movement studies, we found neural support for parallel graded models: lexical

parafoveal effects were observed within 100ms from pre-target onset (Fig. 3b). This finding

does not seem to be compatible even with the most temporally compressed version of a serial

model, in which the attentional shift and a significant amount of lexical parafoveal

previewing occur during saccadic programming[17]. According to such an account, the

neuronal computations that need to be completed before parafoveal previewing can even

begin – retina-brain lag, visual encoding, foveal lexical processing, and shift of attention –

take far too long (around 150ms) to also accommodate the extraction of lexical information

from the next parafoveal word. Moreover, evoked response studies based on EEG[18] and

MEG[19] provide evidence for lexical frequency effects for foveally fixated words no earlier

than 100 ms. Our results showed a lexical frequency effect for the parafoveal word at around

100 ms, which supports the idea of parallel processing of foveal and parafoveal information.

In particular, we provide evidence that parafoveal lexical frequency modulates neural

excitability in support of a parallel model of reading.

One might ask why lexical parafoveal previewing is reflected in the neuronal responses (Fig.

3b) but not in fixation durations (Fig. 2a)? This discrepancy indicates that two systems are at

play simultaneously: covert and overt attention. The fixation duration on a foveal word (pre-

target) captures overt attention, while the RIFT, which reflects the steady-state response to

the parafoveal word (target), captures covert attention. However, these two parallel processes

might not get equal attentional resources. Because the parafoveal word attracts less attention

than the foveal word, its activation level might be insufficient to affect the current foveal

saccade planning and fixation duration. Other studies using a rapid simultaneous visual

presentation paradigm also show that words can be identified in parallel without rising to a

high level of awareness[20]. Nevertheless, even though readers might not be consciously aware

of their parafoveal previewing, it does accelerate individual reading as shown in our

correlation analysis (Fig. 4). Taken together, our study shows that natural reading involves

the simultaneous processing of several words, providing novel neural evidence for the idea

that "readers are parallel processors"[21].

The neuronal response reflecting previewing was observed in the early visual cortex. This

might be a surprise, as functional Magnetic Resonance Imaging studies have localized lexical

frequency to e.g. the visual word area[22]. According to interactive processing theories,

higher-level lexical information interacts with lower-level visual information during word

recognition[18,23]. Thus, lexical frequency information extracted in the parafovea could direct

visual attention covertly. Increased spatial attention will boost tagging responses[10,11],

resulting in stronger coherence for the pre-target word followed by a low compared with a

high lexical frequency target word.

Our results show that RIFT is a new powerful technique to investigate parafoveal reading. A

classic paradigm in this field is the gaze-contingent boundary task developed by Keith

Rayner in 1975[24]. In this task, parafoveal information is manipulated by changing the target

word while saccading to it[4,25]. This approach allows for manipulating parafoveal previewing

and has made great contributions to studies on parafoveal previewing[26]. However, the

approach is limited, as changing the target word inevitably interferes with natural reading.

Fixation related potentials based on EEG is another method used in reading paradigms[27].

While this approach has provided important insights by demonstrating a lexical frequency

effect for foveal word recognition on the N1 component[18,28], it fails to provide conclusive

results with regard to parafoveal lexical processing [4,5]. The novel approach we present in this

study based on subliminal frequency tagging allows for capturing brain activity associated

with parafoveal previewing during natural reading.

It will be important for future studies to use the RIFT to investigate other factors of parallel

information capture during reading. Can parafoveal information be extracted at the semantic

level? Another direction is investigating the primary determinants of reading proficiency in

relation to parafoveal previewing. For instance, previewing at the phonological level

identified using gaze-contingent paradigms has been found to reflect reading proficiency[29].

We found that the neuronal signature of lexical previewing predicted reading speed, which

could be used as a potential indicator to diagnose reading disorders such as dyslexia. Besides,

some researchers argue that dyslexia is due to spatial processing problems in the

magnocellular visual pathway[30]. Our frequency tagging approach could be helpful to

understand the underlying neural mechanism of dyslexia in relation to the allocation of

spatial attention and various factors of previewing including parafoveal previewing at the

phonological level. In sum, the present study demonstrates that RIFT is a powerful tool for

investigating natural reading, and provides novel neural evidence for lexical parafoveal

previewing in support of parallel graded models of reading.

Methods

Participants

Our study recruited forty-three participants (28 females), aged 22 ± 2.6 (mean ± SD), right-handed, with normal or corrected-to-normal vision, and without a neurological history or language disorder diagnosis. Four of them were excluded from analysis due to poor eye tracking or falling asleep during the recordings, which left thirty-nine participants (25 females). The University of Birmingham Ethics Committee approved the study. The participants provided written informed consent and received £15 per hour or course credits as compensation for their participation.

Reading task

Each trial started with a central fixation cross on a grey screen centre presented for 1.2 – 1.6 s. Then followed by a square (1 degree wide) presented 2 degrees to the right of the middle of the screen left edge. A gaze of at least 0.2 s on this square triggered sentence onset. The square was replaced by the first word of the sentence. The text was presented in the equal spaced Courier New font, and each letter occupied 0.35 visual degrees (Fig. 1). After reading the sentence, participants were instructed to fixate on a square below the screen centre for 0.1 s to trigger the sentence offset. One-quarter of the trials were followed by a simple yes-or-no comprehension question to ensure careful reading. Two sets of sentences, consisting of respectively 142 and 86 sentences, were divided into 7 blocks. Each block took approximately 7 minutes to read and was followed by a rest for at least 1 minute (see Supplementary Information for sentence sets details).

Eye-tracking

Each session began with a nine-point calibration and validation test. After every three trials, we performed a one-point drift checking test. If a participant failed to pass drift checking or was unable to trigger sentence onset through gazing, the nine-point calibration and validation test was conducted again. The eye-tracking error was limited to below 1 visual degree both horizontally and vertically.

Visual projection and frequency tagging

All visual stimuli were presented by a PROPixx DLP LED projector (VPixx Technologies

Inc., Canada) at a 1440 Hz refresh rate that provided a stable 60 Hz tagging signal (for a

detailed description about how the projector works, please see Supplementary information).

A Gaussian smoothed patch underlying the target word was flickered at 60 Hz during the

whole sentence presentation. A custom made photodiode (Aalto NeuroImaging Centre,

Finland) was attached to the right-below corner of the screen to record tagging signal from a

square whose luminance was kept the same as the flickering patch.

Coherence calculation

To measure the tagging response associated with target word processing, coherence was

estimated between the MEG sensors and the tagging response of the photodiode (for MEG

sensor selection see Supplementary information). First, 1s segments were filtered using a

phase preserving, two-pass, Butterworth bandpass filters (4th order) with a hamming taper.

The centre filter frequencies were from 40 to 80 Hz in steps of 2 Hz with a 10 Hz frequency

smoothing. For each frequency step, the analytic signals were determined by the Hilbert

transform which then was used as the input for coherence at time point t:

$$299 \quad coh(t) = \frac{\left| n^{-1} \sum_{j=1}^{n} m_{x_j}(t) m_{y_j}(t) e^{i\emptyset_{xy_j}(t)} \right|^2}{n^{-1} \sum_{j=1}^{n} m_{x_j}(t)^2 \, m_{y_j}(t)^2}$$

where j is the trial, n is the number of trials, $m_x(t)$ and $m_y(t)$ are the time-varying

magnitude of the analytic signals from respectively a MEG sensor and a photodiode, $\emptyset_{xy}(t)$

is the phase difference as a function of time between them. A time-frequency coherence

representation was obtained as applied in Fig. 2d and Fig. 3a and 3c (for details of the source

modelling of the MEG data see Supplementary information).

Data and code availability

The data in this study, as well as the codes to generate the associated figures, will be

available upon request from the corresponding author.

References

1.   Rayner, K. & Bertera, J. H. Reading without a fovea. *Science.* **206**, 468–469 (1979).

2.   White, S. J. Eye movement control during reading: Effects of word frequency and

orthographic familiarity. *J. Exp. Psychol. Hum. Percept. Perform.* **34**, 205–223 (2008).

3. Brothers, T., Hoversten, L. J. & Traxler, M. J. Looking back on reading ahead: No
evidence for lexical parafoveal-on-foveal effects. *J. Mem. Lang.* **96**, 9–22 (2017).

4. Degno, F. *et al.* Parafoveal previews and lexical frequency in natural reading :
Evidence from eye movements and Fixation-related potentials. *J. Exp. Psychol. Gen.*
148, 453–474 (2019).

5. Kretzschmar, F., Schlesewsky, M. & Staub, A. Dissociating word frequency and
predictability effects in reading: Evidence from coregistration of eye movements and
EEG. *J. Exp. Psychol. Learn. Mem. Cogn.* **41**, 1648–1662 (2015).

6. Kennedy, A. & Pynte, J. Parafoveal-on-foveal effects in normal reading. *Vision Res.*
45, 153–168 (2005).

7. Kliegl, R., Nuthmann, A. & Engbert, R. Tracking the mind during reading: The
influence of past, present, and future words on fixation durations. *J. Exp. Psychol.*
Gen. **135**, 12–35 (2006).

8. Niefind, F. & Dimigen, O. Dissociating parafoveal preview benefit and parafovea-on-
fovea effects during reading: A combined eye tracking and EEG study.
Psychophysiology **53**, 1784–1798 (2016).

9. Risse, S. & Kliegl, R. Dissociating preview validity and preview difficulty in
parafoveal processing of word n + 1 during reading. *J. Exp. Psychol. Hum. Percept.*
Perform. **40**, 653–668 (2014).

10. Zhigalov, A., Herring, J. D., Herpers, J., Bergmann, T. O. & Jensen, O. Probing
cortical excitability using rapid frequency tagging. *Neuroimage* **195**, 59–66 (2019).

11. Zhigalov, A. & Jensen, O. Alpha oscillations do not implement gain control in early
visual cortex but rather gating in parieto-occipital regions. *Hum. Brain Mapp.*
hbm.25183 (2020) doi:10.1002/hbm.25183.

12. Conner, J. D. & Macleod, D. I. A. Rod photoreceptors detect rapid flicker. *Science.*
195, 698–699 (1977).

13. Engbert, R., Longtin, A. & Kliegl, R. A dynamical model of saccade generation in
reading based on spatially distributed lexical processing. *Vision Res.* **42**, 621–636

(2002).

14.   Engbert, R., Nuthmann, A., Richter, E. M. & Kliegl, R. Swift: A dynamical model of
343   saccade generation during reading. *Psychol. Rev.* **112**, 777–813 (2005).

344   15.   Reichle, E. D., Pollatsek, A., Fisher, D. L. & Rayner, K. Toward a model of eye
345   movement control in reading. *Psychol. Rev.* **105**, 125–157 (1998).

346   16.   Reichle, E. D., Rayner, K. & Pollatsek, A. The E-Z reader model of eye movement
347   control in reading: Comparisons to other models. *Behav. Brain Sci.* **26**, 445–476
348   (2003).

349   17.   Reichle, E. D. & Reingold, E. M. Neurophysiological constraints on the eye-mind link.
350   *Front. Hum. Neurosci.* **7**, 361 (2013).

351   18.   Hauk, O., Davis, M. H., Ford, M., Pulvermüller, F. & Marslen-Wilson, W. D. The time
352   course of visual word recognition as revealed by linear regression analysis of ERP
353   data. *Neuroimage* **30**, 1383–1400 (2006).

354   19.   Assadollahi, R. & Pulvermüller, F. Neuromagnetic evidence for early access to
355   cognitive representations. *Neuroreport* **12**, 207–213 (2001).

356   20.   Snell, J. & Grainger, J. Consciousness Is Not Key in the Serial-versus-Parallel Debate.
357   *Trends Cogn. Sci.* **23**, 814–815 (2019).

358   21.   Snell, J. & Grainger, J. Readers are parallel processors. *Trends Cogn. Sci.* **23**, 537–546
359   (2019).

360   22.   Dehaene, S. & Cohen, L. The unique role of the visual word form area in reading.
361   *Trends Cogn. Sci.* **15**, 254–262 (2011).

362   23.   Carreiras, M., Armstrong, B. C., Perea, M. & Frost, R. The what, when, where, and
363   how of visual word recognition. *Trends Cogn. Sci.* **18**, 90–98 (2014).

364   24.   Rayner, K. The perceptual span and peripheral cues in reading. *Cogn. Psychol.* **7**, 65–
365   81 (1975).

366   25.   Dimigen, O., Kliegl, R. & Sommer, W. Trans-saccadic parafoveal preview benefits in
367   fluent reading: A study with fixation-related brain potentials. *Neuroimage* **62**, 381–393
368   (2012).

26.   Schotter, E. R., Angele, B. & Rayner, K. Parafoveal processing in reading. *Attention,*

Perception, Psychophys. **74**, 5–35 (2012).

27.   Dimigen, O., Sommer, W., Hohlfeld, A., Jacobs, A. M. & Kliegl, R. Coregistration of

eye movements and EEG in natural reading: Analyses and review. *J. Exp. Psychol.*

Gen. **140**, 552–572 (2011).

28.   Sereno, S. C., Rayner, K. & Posner, M. I. Establishing a time-line of word recognition:

Evidence from eye movements and event-related potentials. *Neuroreport* **9**, 2195–

2200 (1998).

29.   Pollatsek, A., Lesch, M., Morris, R. K. & Rayner, K. Phonological codes are used in

integrating information across saccades in word identification and reading. *J. Exp.*

Psychol. Hum. Percept. Perform. **18**, 148–162 (1992).

30.   Vidyasagar, T. R. & Pammer, K. Dyslexia: a deficit in visuo-spatial attention, not in

phonological processing. *Trends Cogn. Sci.* **14**, 57–63 (2010).

Acknowledgements

We thank F. Degno and S. Liversedge for sharing the second sentence set, Geoffrey
Brookshire and Yang Cao for feedbacks on the manuscript. This work was supported by the
following grants to O.J.: the James S. McDonnell Foundation Understanding Human
Cognition Collaborative Award (grant number 220020448), Wellcome Trust Investigator
Award in Science (grant number 207550), and the BBSRC grant (BB/R018723/1) as well as
the Royal Society Wolfson Research Merit Award.

Author Contributions

Y.P. and O.J. devised and designed the experiments with assistance from S.F., Y.P. made the
sentences with assistance from S.F, Y.P. programmed and conducted the experiments, Y.P.
carried out the analyses with assistance from O.J., Y.P., O.J. and S.F. wrote the paper
together.

Supplementary table and figures

Supplementary Table 1 | Characteristics of words in the experimental materials (1st set)

Measure	Pre-target	Target				Post-target
	Length	Low frequency	High frequency	Length	Position	Length
Mean	6.1	5.3	95.3	5.8	6.7	6.7
SD	1.5	4.5	135.5	0.8	2.3	1.7

Low (<10) and high lexical frequency (>30) target words are reported in terms of the total CELEX frequency per million[1]. Length is the number of letters in a word. Position refers to the location in a sentence where the target word is presented. The sentences were 11.2 ± 2.1 words long. For details about words in the 2nd sentence set, please see Degno et al., 2019[2].

Supplementary Figure 1 | Gaze-duration difference for pre-target and target words.
Gaze duration refers to the sum of all first-pass fixations for a given word. The differences for low- minus high- lexical frequency of the target words. A significant lexical frequency effect was only observed for target words, with longer gaze durations for low- compared with high- lexical frequency (*** $p < 0.001$, n = 39, two-tailed paired t-test). We observed no significant gaze duration difference for pre-target words ($p = 0.977$, n = 39, two-tailed paired t-test). The horizontal bar in the violin plot indicates mean value; each dot represents one participant. This finding supports the conclusion of Fig. 2a.

Supplementary Figure 2 | Pre-target coherence onset latency was modulated by parafoveal target lexical frequency in the short-words sentence set. A jackknife-based method was used to identify the onset latency for 60 Hz coherence on the group level. Here onset latency refers to the time when coherence reached half maximum, denoted by the dotted lines. (**a**) Each participant read two sets of sentences; for combined sentence set, no significant difference was found in neither pre-target fixation (left panel, $p > 0.1$, n = 26, two-tailed paired t-test) nor target fixation (right panel, $p > 0.4$, n = 26, two-tailed paired t-test). (**b**) The histogram of word length showed that in the 2nd sentence set (in red), both pre-target

and target words were shorter compared with the 1st sentence set (in blue). (**c**) For the 2nd

sentence set, a significant coherence onset difference was found in pre-target fixation (left

panel, $p < 0.01$, n = 26, two-tailed paired t-test), but not in target fixation (right panel, $p > 0.4$,

n = 26, two-tailed paired t-test). (All analyses in the study were based on the combined

sentence set except this one).

Supplementary Figure 3 | Initial landing position for pre-target words and all words.

Landing positions for the first fixation on words were estimated in the unit of 'letter' aligned

according to the word centre. Distribution in dark grey was the landing position error for pre-

target words (-1.1 ± 1.9, mean ± SD); in light grey, was the landing position error for all

words (-0.8 ± 1.8, mean ± SD). This demonstrates that the landing position of fixations was

around ~1 letter to the left of the centre of the words fixated on.

Supragraph Supplementary Information

Sentence material

1ˢᵗ sentence set

We constructed 142 sentences embedded with 71 target word pairs (low/high lexical frequency). For each sentence, the pre-target, target, and post-target words were in the same structure as adjective + noun + verb. For each target pair, two different sentence frames (versions) were made, so each participant can read both target words embedded in different frames. For example, for target pair **waltz/music** (low/high lexical frequency), one participant read version A, another one read version B (see below, targets were bold for illustration, but not in the real experiment).

```
A. Mike thought this difficult waltz received lots of criticism.
   It was obvious that the beautiful music captured her attention.
B. Mike thought this difficult music received lots of criticism.
   It was obvious that the beautiful waltz captured her attention.
```
The sentences in version B were made from version A by circular shifting the first and second half of the sentences. For both versions, no more than 3 successive sentences were from the same target lexical frequency condition.

2ⁿᵈ sentence set

This sentence set was adapted from Degno et al., 2019[2]. After removing sentences that contained the same pre-target or target words as in the 1ˢᵗ sentence set, left 86 sentences. Each sentence was embedded with two target words from the same lexical frequency condition (see below, version A contained two low lexical frequency targets, while version B contained two high lexical frequency targets).

```
A. I felt quite bleak after discussing that really risky subject with Paul.
B. I felt quite weird after discussing that really nasty subject with Paul.
```
Each participant read either version A or B. The same control for sentence sequence as in the 1ˢᵗ set.

Pre-tests for target predictability and sentence plausibility

473 We only did pre-test for the 1st sentence set, for pre-test details about the 2nd set please see

474 Degno et al., 2019[2]. All participants in the pre-test were different from participants in the

475 MEG session.

Predictability

A cloze test was performed with a partial sentence just before the target word, and

participants were asked to read the sentence then write down the first word/s that came to

mind ssto complete the sentence. See below as an example for target word **waltz/music**:

`Mike thought this difficult ________________`

If a target word was predicted by less than 10% of the participants, this target was considered

unpredictable. First, 22 participants (1 male) took part in this pre-test, and 6 target words

turned out to be highly predictable. These 6 target words were changed to other words and

the predictability test for the whole sentence set was conducted again with another 22

participants (3 males). And none of the target words was predictable.

Plausibility

Participants were instructed to rate how plausible (or acceptable) each sentence was. In a 7-

points rating scale, from 1 to 7 indicated implausible to plausible for each sentence (see

below for examples). To occupy the full range of plausibility scale, we made 142 filler

sentences, half of which were middle plausible (e.g., sentence 1) and another half were

implausible (e.g., sentence 3). Here in this example, the second sentence was from the

experiment.

	Implausible Plausible
	1 2 3 4 5 6 7
1. Kate said that she saw lots of stars twinkling in the sky at noon.	1 2 3 4 5 6 7
2. It was obvious that the beautiful music captured her attention.	1 2 3 4 5 6 7
3. Sarah used an adhesive to glue the traffic and just made it to the wedding.	1 2 3 4 5 6 7
……	1 2 3 4 5 6 7

For version A (n = 22, 2 males, 2 invalid), the rating for experimental sentence plausibility

was 5.7 ± 0.5 (mean $\pm$ SD), which was significantly higher than filler sentences (two-tailed

pair-wised $t_{(19)} = 23.98$, $p < 0.001$). For version B (n = 23, 4 males), the rating was 5.5 ± 0.8

(mean ± SD), which was also significantly higher than filler sentences (two-tailed pair-wised

$t_{(22)} = 16.68$, $p < 0.001$). These results showed that all the sentences in both versions were

plausible.

Procedure

Participants were seated 145 cm away from the projection screen in a dimly-lit magnetically

shielded room. One-line sentence was presented on a grey screen using Psychophysics

Toolbox -3[3]. All sentences started at the same position: two degrees to the right of the middle

of the screen left edge. Words were displayed in black using an equal-space Courier New font

(font size as 22). Each letter occupied 0.35 visual degrees, the same for the space between

words. In total, each sentence was no longer than 27 visual degrees horizontally. Participants

were instructed to read each sentence silently at their own pace and to keep the head and

body stable during the MEG session. Eye-movements were also acquired . One-quarter of the

sentences were followed by a yes-or-no comprehension question to ensure careful reading.

Participants put their left and right index fingers of the right hand on a two buttons response

box to respond to the question. The mapping between yes-or-no responses and button presses

was balanced over participants.

Rapid Invisible frequency tagging

Flickering target word

To flicker the target word, we added a rectangular patch underneath the target. The length of

the patch was the width of the target word plus the spaces on both sides. The height of the

patch was 1.5 times of the word height. The target word was placed in the centre of this

patch. All pixels within the patch were flickered at 60 Hz by multiplying the luminance of the

pixels with a 60 Hz sinusoid (the modulation depth was 100%). Typically, the patch was

perceived as indistinguishable from the grey background, making it invisible. To reduce the

visibility of the patch edges during saccades, a Gaussian smoothed transparent mask was

applied on top of the flickering patch. The mask was created by a two-dimensional Gaussian

function:

$$mask = \exp\left(-\frac{x^2}{2\sigma^2} - \frac{y^2}{2\sigma^2}\right)$$

Where, x and y are the mesh grid coordinates for the flickering patch, and σ is the x and y

spared of the blob with σ = 0.02 degrees. By applying a Gaussian smoothed mask, the

flickering patch was hardly noticed. Only 3 out of 39 participants noticed the flickering patch

according to a questionnaire after the MEG session.

Projector

To achieve a stable rapid invisible frequency tagging, sentence stimuli were presented with a

refresh rate up to 1440 Hz using a PROPixx DLP LED projector (VPixx Technologies Inc.,

Canada). This was done by presenting the sentence stimuli repeatedly in four quadrants on

the stimulus computer screen (1920 × 1200 pixels resolution) with a refresh rate of 120 Hz.

For each quadrant, the stimuli were coded in RGB three colour channels. Then the projector

interpreted these 12 colour channels (3 channels × 4 quadrants) as 12 individual grayscale

frames and projected them onto the projector screen in rapid succession. This allowed for a

1440 Hz (120 Hz × 12) refresh rate.

Data acquisition

MEG

MEG data were acquired with a 306-sensor TRIUX Elekta Neuromag system with 204

orthogonal planar gradiometers and 102 magnetometers (Elekta, Finland). The data were

band-pass filtered online using anti-aliasing filters from 0.1 to 330 Hz and sampled at 1,000

Hz. To 3D-digitize the head shape of each participant, we first used the Polhemus Fastrack

electromagnetic digitizer system (Polhemus Inc, USA) to digitalize location of four head-

position indicator coils (HPI coils) as well as three bony fiducial points: the nasion, left and

right preauricular points. Two coils were attached on the left/right mastoid bone and another

two on the forehead at least 3 cm in between. Furthermore, at least 200 extra points on the

scalp were acquired for each participant. After preparations, participants were seated upright

under the MEG dewar at a 60° orientation position.

Eye-movements

553 The eye-tracker (EyeLink 1000 Plus, SR Research Ltd, Canada) was placed on a wooden

554 table in front of the bottom edge of the projector screen. The distance between the eye-tracker

555 camera and participant's left-eye was 90 cm. It was used throughout the experiment to

556 acquire horizontal and vertical positions of the left eye as well as pupil size sampled at 1,000

557 Hz. We also placed one pair of electrodes above and below the right eye (vertical

558 electrooculogram; EOG) and one pair to the left and right of the eyes (horizontal EOG) to

559 provide additional measures for ocular and eye-blink artefacts.

560 *MRI*

561 After MEG data acquisition, we acquired the T1-weighted structural MRI using a 3-Tesla

562 Siemens PRISMA scanner (TR = 2000 ms, TE = 2.01 ms, TI = 880 ms, flip angle = 8

563 degrees, FOV = 256×256×208 mm, 1 mm isotropic voxel). Out of all 39 participants, 3

564 dropped out for the MRI acquisition, and the standardized template brain was used in later

565 source modelling analysis.

567 **MEG data analyses**

568 The data analyses were performed in MATLAB R2019b (Mathworks Inc, USA) by using the

569 FieldTrip[4] toolbox (version 20200220) and custom-made scripts.

570 *Pre-processing*

571 The MEG data were band-pass filtered from 0.5 to 100 Hz using phase preserving two-pass

572 Butterworth filters. First, the MEG segments were extracted from the -0.5 – 0.5 s interval

573 aligned with the first fixation onset to pre-target, target, and post-target words separately.

574 Only segments with fixation duration ranging from 0.08 to 1 s entered further analyses.

575 Baseline segments (1 s long) were aligned with the fixation cross onset before sentence

576 presentation. Next, the data segments were demeaned by removing the linear trend and the

577 mean. After removing malfunctioning sensors (0 to 2 sensors per participant), all these

578 segments entered an independent component analysis (ICA[5]). Before the ICA, a PCA

579 approach was used to reduce the rank of the data to 30 components after which components

580 related to eye blinks, eye movements, and heartbeat were rejected. Finally, we manually

581 inspected all these segments to identify and remove segments with excessive noise, like

582 remaining ocular, muscle and movement artefacts.

 Tagging response sensor selection

To identify MEG sensors that showed a reliable 60 Hz coherence with the photodiode during

the pre-target segments compared with the baseline segments, a permutation test was

conducted. Only sensors in the visual cortex (52 planar sensors) entered the permutation test.

For each MEG sensor and photodiode channel combination, coherence at 60Hz was

computed for the 1 s pre-target and 1 s baseline segments over trials and transformed into a z-

value. Next, the segments for the 'pre-target' and 'baseline' were randomly relabelled after

which coherence was re-computed and one z-value was obtained. After 10000 shuffles, a null

distribution of z-values was established. If the empirical z-value was larger than 99% of the

null distribution, this sensor was considered to have a robust tagging response (at a $p = 0.01$

'significance level'). After this permutation procedure, 26 out of all 39 participants showed a

robust tagging response in one or more sensors (5.4 ± 4.0 sensors per participant, mean $\pm$

SD).

Source modelling

In order to estimate the neural source of the tagging response, a beamforming approach was

performed using dynamic imaging coherent sources [9] (DICS) implemented in FieldTrip.

Participants with robust tagging responses entered this source analysis (n = 26; see Fig. 2b).

The DICS approach computes a spatial filter from the lead-field matrix and a cross-spectral

matrix (CSD).

To calculate the individual lead-field matrices, we first constructed a forward model by

aligning the individual structural MRI image with the head-position in the MEG system

according to the three fiducial anatomical markers from the head shape digitization (the

nasion and the two preauricular points). Next, a realistically shaped single-shell head model

based on harmonic functions[10] was constructed based on the segmented anatomical MRI for

each participant. Then the brain volumes were divided into a 5 mm spaced grid and warped to

an MNI template brain (Montreal, Quebec, Canada).

The CSD matrix was calculated as follows. For each participant, the time window was the

shortest pre-target fixation interval (same time window as in the coherence t-test in Fig. 3b

left panel). The centre frequency was 60 Hz. For the lead-field matrix and CSD matrix, a

common spatial filter was computed based on all pre-target fixation intervals. The coherence

value at each grid point was then calculated by applying this spatial filter. Finally, we

obtained the neural sources for the coherence as shown in Fig. 2c by the grand average across

all participants (n = 26).

Jackknife-based method for onset latency

We used a jackknife-based method[11] to assess the difference in onset latency for pre-target

and target coherence with respect to lexical frequency (Supplementary Fig. 2). The time-point

of the onset was estimated from when the averaged coherence reached its half-maximum

$(coh_{min} + (coh_{max} - coh_{min})/2)$. Using a leave-one-subject-out jackknife approach, the

coherence onset difference (high versus low lexical frequency) was re-computed across the

remaining participants. The standard error (S_D) was estimated from these latency differences

when looping over subjects. The latency difference observed in the overall sample (without

leaving out a participant) was also computed and then divided by the S_D to obtained the *t*-

value. The statistical significance for onset latency difference between the two conditions

(high versus low lexical frequency) was then estimated using a standard *t* table (for details

see Miller et al., 1998[11]).

Statistical information

For all the t-tests conducted in this study, we used a two-sided pairwise student's t-test using

R[12].

Method references

1.   Davis, C. J. N-watch: A program for deriving neighborhood size and other

psycholinguistic statistics. *Behav. Res. Methods* **37**, 65–70 (2005).

2.   Degno, F. *et al.* Parafoveal previews and lexical frequency in natural reading :

Evidence from eye movements and Fixation-related potentials. *J. Exp. Psychol. Gen.*

148, 453–474 (2019).

3.   Kleiner, M., Brainard, D. & Pelli, D. What's new in Psychtoolbox-3? *Percept. 36*

ECVP Abstr. Suppl. (2007).

4.   Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: Open Source

Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological

Data. *Comput. Intell. Neurosci.* **2011**, (2011).

5.   Ikeda, S. & Toyama, K. Independent component analysis for noisy data - MEG data
analysis. *Neural Networks* **13**, 1063–1074 (2000).

6.   Zhigalov, A. & Jensen, O. Alpha oscillations do not implement gain control in early
visual cortex but rather gating in parieto-occipital regions. *Hum. Brain Mapp.*
hbm.25183 (2020) doi:10.1002/hbm.25183.

7.   Zhigalov, A., Herring, J. D., Herpers, J., Bergmann, T. O. & Jensen, O. Probing
cortical excitability using rapid frequency tagging. *Neuroimage* **195**, 59–66 (2019).

8.   Duecker, K., Gutteling, T. P., Herrmann, C. S. & Jensen, O. No evidence for
entrainment: endogenous gamma oscillations and 1 responses to rhythmic visual
stimulation coexist in visual cortex. *bioRxiv* 2020.09.02.279497 (2020)
doi:10.1101/2020.09.02.279497.

9.   Gross, J. *et al.* Dynamic imaging of coherent sources: Studying neural interactions in
the human brain. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 694–699 (2001).

10.   Nolte, G. The magnetic lead field theorem in the quasi-static approximation and its use
for magnetoenchephalography forward calculation in realistic volume conductors.
Phys. Med. Biol. **48**, 3637–3652 (2003).

11.   Miller, J., Patterson, T. & Ulrich, R. Jackknife-based method for measuring LRP onset
latency differences. *Psychophysiology* **35**, 99–115 (1998).

12.   Team, R. C. R: A language and environment for statistical computing. (2013)
doi:10.1016/j.dendro.2008.01.002.