1 **Title:** A novel subset of follicular helper-like MAIT cells has capacity for B cell help and antibody
2 production in the mucosa.
3

4 **Author List:** Owen Jensen[1,2], Shubhanshi Trivedi[1], Jeremy D. Meier[3,4], Keke Fairfax[2], J. Scott Hale[2],
5 Daniel T. Leung[1,2*]
6
[1] Division of Infectious Diseases, Department of Internal Medicine, University of Utah School of
Medicine, Salt Lake City, USA
[2] Division of Microbiology & Immunology, Department of Pathology, University of Utah School of
Medicine, Salt Lake City, USA
[3] Division of Otolaryngology-Head and Neck Surgery, University of Utah School of Medicine, Salt Lake
City, USA
[4] Primary Children's Hospital, Salt Lake City, USA
14
15 * Corresponding author: daniel.leung@utah.edu

16
17 **One Sentence Summary:** We identified and characterized a novel subset of T follicular helper-like
18 MAIT (MAITfh) cells that has the capacity to provide B cell help, and show the sufficiency of MAIT
19 cells to promote production of pathogen-specific IgA antibodies and B cell differentiation in mucosal
20 challenge.
21
22
Abstract

Mucosal-associated invariant T (MAIT) cells are innate-like T lymphocytes that aid in protection against

bacterial pathogens at mucosal surfaces via release of inflammatory cytokines and cytotoxic molecules.

Recent evidence suggests MAIT cells are capable of providing B cell help. In this study, we describe a

previously unreported population of CXCR5[+] T follicular helper (Tfh)-like MAIT cells, MAITfh, that

have the capacity to provide B cell help within mucosal lymphoid organs. MAITfh cells are preferentially

located near germinal centers in human tonsils and express the classical Tfh-associated transcription

factor, B-cell lymphoma 6 (BCL-6), co-stimulatory markers, inducible T cell costimulatory (ICOS) and

programmed death receptor 1 (PD-1), and cytokines, interleukin (IL)-21. Furthermore, we demonstrate

the ability of MAIT cells to provide B cell help *in vivo* following mucosal challenge with *Vibrio cholerae*.

Specifically, we show that adoptive transfer of MAIT cells into αβ T cell-deficient mice promoted B cell

differentiation and increased serum *V. cholerae*-specific IgA responses. Our data demonstrate the capacity

of MAIT cells to participate in adaptive immune responses, and suggest that MAIT cells may be potential

targets for mucosal vaccines.

Introduction

Mucosal surfaces are in constant contact with both commensal and pathogenic microbes. One of the primary means of protection against invading pathogens is the production of secretory IgA and IgM by plasmablasts (PB) and plasma cells (PC) (*1*). In the lamina propria of the gut and lung, T independent low affinity IgA is produced against non-protein antigens by short-lived PBs. Conversely, T dependent high affinity antibodies against protein antigens are typically generated during germinal center reactions between T follicular helper (Tfh) and follicular B cells within local lymph nodes (LNs) or mucosal associated lymphoid tissues (MALT) (*2*). T follicular helper (Tfh) cells migrate to B cell follicles via upregulation of CXCR5 and downregulation of CCR7 (*3*). Within GCs, Tfh cells highly express the lineage-defining transcription factor, BCL6, and co-stimulatory molecules, PD1, ICOS, and CD40L (*4*). PD1 is important for Tfh GC positioning and function (*5*), while ICOS (*6*) and CD40L (*7*) engagement with B cells are essential for Tfh activation and GC formation (*4*). Tfh also produce cytokines, such as IL-21, that promote GC B cell responses (*8*).

Although Tfh cells are the primary drivers of T dependent GC responses, innate T cells, including invariant Natural Killer T (iNKT) cells and γδ T cells, also have Tfh like subsets capable of B cell help. In particular, iNKT cells have been well established in mice to provide both cognate and non-cognate help to B cells (*9–18*). Murine iNKTfh cells engage in cognate help leading to germinal center formation, and antibody class switching and production (*12, 14, 18*). iNKTfh cells have also been shown to promote non-cognate B cell help by licensing dendritic cells to recruit and activate Tfh cells (*10, 11*). CXCR5$^+$ γδ T cells can promote antibody production *in vitro* (*19, 20*), and promote Tfh differentiation *in vivo* (*21*). More recent evidence suggests that another type of innate-like lymphocyte, Mucosal-associated invariant T (MAIT) cells, are capable of B cell help (*22–26*).

MAIT cells are innate-like $\alpha\beta$ T cells defined by the expression of an invariant α chain, generally Vα7.2

linked to Jα33, 12 or 20 in humans and Vα19 linked to Jα33 in mice, and a limited array of TCRβ chains

(27–29). MAIT cells are highly enriched in human blood, liver and mucosa and are appreciated for their

rapid ability to respond to microbial Vitamin B metabolites presented on the MHC class I related protein,

MR1, or cytokine stimulation (30–33). Upon stimulation, MAIT cells produce pro-inflammatory

cytokines including IFNγ, TNFα, and IL-17A and cytotoxic molecules including Granzyme B and

Perforin (34–36). Furthermore, using MAIT deficient mice (MR1$^{-/-}$), several studies have exemplified the

importance of MAIT cells in immunity against mucosal bacterial pathogens (37–40).

Recent evidence suggests MAIT cells play a role in adaptive immune responses through B cell help.

Analysis of human peripheral blood MAIT cells and serum antibody responses following *Vibrio cholerae*

infection (22) and *Shigella* vaccination (35), revealed strong associations between MAIT frequency and

activation with polysaccharide-specific IgA and IgG responses, but not with protein antibody responses

(22). We have recently shown that human blood MAIT cells have the capacity to induce antibody

production and B cell differentiation *in vitro*, and can secrete the B cell help cytokines following

stimulation (23). Analysis of pleural effusions from tuberculosis patients revealed a population of PD1^{High}

MAIT cells secreting key B cell help cytokines (24). Furthermore, two recent animal studies demonstrate

the importance of MAIT cells in B cell help in murine autoimmunity (25) and mucosal vaccine immunity

in non-human primates (26).

Our aims for this study were two-fold. We first wanted to determine whether a specific subset of MAIT

cells are responsible for the B cell help phenotype. Our second aim was to determine *in vivo* if MAIT

cells were sufficient to induce antibody production and humoral immune protection in the context of

mucosal challenge. We found that like other innate-like T cells, MAIT cells have a Tfh like subset

enriched within mucosal lymphoid organs. This MAITfh population expresses classical Tfh co-

stimulatory markers, transcription factors, and cytokines, and is localized near B cell follicles. We further

show, in a murine model, that adoptively transferred MAIT cells are capable of generating microbe-

specific antibody responses against a mucosal bacterial pathogen in the absence of other αβ T cells.

Additionally, we find that in the context of mucosal challenge, MAIT cells promote increased production

of microbe-specific IgA antibodies and mucosal B cell differentiation. These results suggest that MAIT

cells have the capacity to enhance mucosal antibody mediated immunity, and thus may be a target in

future mucosal vaccine development.

Results

CXCR5$^+$ MAIT cells are increased in tonsils and express higher levels of Tfh co-stimulatory markers

compared to peripheral blood.

With recent evidence highlighting the ability of MAIT cells to provide B cell help, and studies showing a

modest population of MAIT cells within human and mouse mucosal lymphoid tissues (*39*, *41*, *42*), we

aimed to determine if MAIT cells had a Tfh-like subset capable of B cell help. To investigate this, we

used flow cytometry to analyze MAIT cell expression of classical Tfh markers in human peripheral blood

mononuclear cells (PBMCs) and tonsils. We obtained PBMC's from adult blood donors, and tonsils from

children, ages 2-16, undergoing tonsillectomy for recurrent tonsillitis or tonsillar hyperplasia. MAIT cells

were defined as CD3$^+$ Vα7.2$^+$ MR1-5-OPRU Tetramer$^+$ cells and gated based on a PE conjugated MR1-

6FP Tetramer negative control (fig S1A, S1B and Fig. 1A). Median MAIT frequency among total CD3$^+$

cells in tonsils was 0.23% (interquartile range (IQR) = 0.14%, 0.29%) compared to 1.03% (IQR = 0.62%,

1.35%) among PBMCs (Fig. 1B). We found that a higher percentage (Median=18.1%, IQR=9.2%,

28.6%) of tonsil MAIT cells were CXCR5$^+$ compared to PBMC MAIT cells (Median=1.74%,

IQR=0.37%, 2.16%, p<0.0001), although there was considerable variability among tonsil MAIT cells

(Fig. 1C and fig. S1C). To further confirm the MAIT cell phenotype, we measured expression of two C

type lectin receptors, CD161, a common delineator of MAIT cells (*30*), and CD69, a marker of T cell

Fig. 1. Increased expression of Tfh co-stimulatory molecules in tonsil MAIT cells. (A) Representative FACS plots of unstimulated CD3$^+$ PBMC (top) and tonsil (bottom) cells with MAIT cell (gated in left panel) co-expression of CXCR5 with ICOS (middle) and PD1 (right). **(B)** Quantification of MAIT cell frequency as percentage of CD3$^+$. **(C)** Frequency of CXCR5$^+$ MAIT cells. Frequency of **(D)** ICOS$^+$ and **(E)** PD1$^+$ MAIT (CXCR5$^-$), MAITfh (CXCR5$^+$) and Tfh cells. Data are represented as Median from 2 pooled independent experiments. n ≥ 13. *p < 0.05, **p<0.01, ***p<0.001, ****p<0.0001 by two-tailed Mann-Whitney U test

tissue residency (*43*). We found that significantly fewer CXCR5+ MAIT cells expressed CD161 in both

PBMC (p<0.01) and tonsils (p<0.001) than did CXCR5- MAIT cells (fig. S2D and F). The loss of CD161

expression has been demonstrated in vitro following TCR stimulation, suggesting CXCR5+ MAIT cells

may be previously activated (*44*). Despite being MR1-Tetramer+ and TRAV1-2+, we wanted to confirm

that the CD161- fraction were indeed MAIT cells and therefore utilized paired TCRαβ single-cell

Illumina sequencing as previously described (*45*). We sorted and single-cell sequenced CD161+ and

CD161- MAIT cells (identified as live CD3$^+$ Vα7.2$^+$ MR1-Tetramer$^+$ cells) from four tonsil samples and

analyzed *TRAJ* expression in *TRAV1-2* expressing wells. We found no statistically significant differences

in *TRAJ* expression frequency between CD161+ and CD161- MAIT cells, (fig. S3A) with the majority

expressing *TRAJ-33* with average frequencies of 66.3% ± 17.5 and 65.5% ± 7.9, respectively. In

concurrence with previous studies (*29, 46–48*), *TRAJ-34* and *TRAJ-12* were also highly expressed in both

groups highlighting that these are bona fide MAIT cells. Furthermore, we found no statistically

significant differences in the percentages of CXCR5+ and CXCR5- MAIT cells that expressed CD69 in

PBMC and tonsils (fig. S2C and E), though both CXCR5+ and CXCR5- MAIT cells in the tonsils had

higher cell surface levels CD69 expression, indicating a tissue resident memory (Trm) phenotype. We

next wanted to determine if there were differences in Tfh co-stimulatory marker expression between

CXCR5+ and CXCR5- MAIT populations. Significantly more CXCR5+ MAIT cells in tonsils expressed

ICOS (p<0.0001) (Fig. 1D and fig S2B) than did CXCR5- MAIT cells; moreover, significantly more

CXCR5+ MAIT cells in both tonsils and PMBC's expressed PD1 (Fig. 1E and fig S2A) than did CXCR5-

MAIT cells in the same tissue. Additionally, significantly more CXCR5+ MAIT cells in tonsils expressed

PD1 (p<0.0001) and ICOS (p<0.0001) than did PBMC CXCR5+ MAIT cells. Notably, ICOS and PD1

frequency and fluorescence intensity among tonsil CXCR5$^+$ MAIT cells are similar to that seen in tonsil

CD4$^+$ CXCR5$^+$ (Tfh) cells (Fig.1D and 1E, and fig. S2A and S2B). In order to confirm that the

differences between PBMC and tonsil MAIT cells are not age dependent, we analyzed MAIT cell

CXCR5, ICOS and PD1 expression in 5 paired PBMC and tonsil samples from pediatric donors ages 3-15

(fig. S4). Similar to adult PBMC samples we found increased total MAIT frequency and decreased

CXCR5 expression in PMBC compared to tonsil samples (fig. S4B-C). Similar trends in ICOS and PD1

expression were found in the pediatric PBMC MAIT and MAITfh groups compared to the adult PBMC

donors supporting the use of adult PBMC samples in the following figures where low pediatric blood

sample volumes would limit analysis (Fig. 1D-E and fig. S4D-E). Taken together, a high proportion of

MAIT cells in tonsils have a CXCR5$^+$ phenotype with Tfh-like co-stimulatory markers.

Transcriptional analyses of CXCR5$^+$ MAIT cells reveals expression of Tfh associated cytokines and

transcription factors

We have previously demonstrated that peripheral blood MAIT cells can secrete B cell help cytokines IL-

6, IL-10 and low levels of IL-21 *in vitro* (*23*). MAIT cells are also potent producers of IFNγ (*49*), which is

known to have both inhibitory and activating roles in B cell development, proliferation, and antibody

responses (*50*). In order to investigate the transcription factor and cytokine expression of CXCR5$^+$ and

CXCR5$^-$ MAIT cells, we examined transcript levels of fluorescent associated cell sorted (FACS) MAIT

and non-MAIT CD3$^+$ T cell populations from tonsil and blood. We refer to the sorted populations, as

CD8$^+$ (CD8$^+$ CXCR5$^-$), Tfh (CD4$^+$ CXCR5$^+$), MAIT (MR1-Tet$^+$, Vα7.2$^+$, CXCR5$^-$), and MAITfh (MR1-

Tet$^+$, Vα7.2$^+$, CXCR5$^+$) (fig. S5A). To confirm our flow analysis, we assayed for *CXCR5* expression. We

found significant increases in *CXCR5* relative transcript levels in the MAITfh compared to the MAIT

groups in both PBMC's ($p<0.01$) and tonsils ($p<0.001$) (Fig. 2A). Though notably, the MAITfh group had

lower CXCR5 expression than the control Tfh group in both tissues. We next assayed for the Tfh lineage-

defining transcription factor (TF), *BCL6*. The MAITfh population within PBMCs had higher *BCL6*

expression compared to MAIT ($p<0.05$) and Tfh ($p<0.01$) cells. Within tonsils, MAITfh cells had higher

BCL6 expression compared to Tfh ($p<0.01$) and CXCR5- MAIT ($p<0.08$) populations (Fig. 2B). In

comparison, analysis of canonical T cell associated TFs *RORC* (Th17) ($p<0.05$), and *TBX21* (Th1)

($p<0.05$)*,* revealed higher expression in the tonsil MAIT over the MAITfh population (Fig. 2C and D).

MAITfh groups had higher expression of *RORC* and *TBX21* compared to Tfh groups in PBMCs ($p<0.01$)

and tonsils ($p<0.01$) (Fig. 2C and D). Given that conventional Tfh cells are known to co-express TFs

associated with other helper T cell subsets based on lineage and environment (*51–53*), our data suggest

that co-expression of non-follicular helper-associated TFs in the MAITfh population may suggest

potential for plasticity in phenotype or differentiation from a Th1 or Th17-like subset.

Fig. 2. CXCR5$^+$ MAIT cells highly express B cell help cytokine IL-21. (**A**) *CXCR5*, (**B**) *BCL6*, (**C**) *RORC*, (**D**) *TBX21*, (**E**) *IL-21*, (**F**) *IL-6*, (**G**) *IL-10*, and (**H**) *TNFSF13B* qPCR data of unstimulated FAC sorted PBMC and tonsil T cell populations. Data are represented as ΔΔCt relative to an internal housekeeping gene, β-Actin, and the PBMC-CD8$^+$ CXCR5$^-$ population. Data are Median with IQR from 2 independent experiments. n=6-8. (**I**) Representative flow cytometry from PBMC and tonsil T cell populations of IL-21 and IFNγ co-expression following 6 h stimulation with PMA/Ionomycin and Brefeldin A for final 4h. (**J**) Frequency of T cell populations expressing IL-21, IFNγ or both. Data are Median with IQR from 2 pooled independent experiments. n≥11. (**K**) IL-21 ELISA following 42 h PMA/Ionomycin stimulation of FAC sorted mono-cultured T cell populations from tonsils described below. Data are Median with IQR, n=5. T cell populations described gated on live CD3+ cells are as follows: Black =CD8$^+$ CXCR5$^-$, Blue = Tfh (CD4$^+$ CXCR5$^+$), Red= MAIT (MR1-Tetramer$^+$ Va7.2$^+$) CXCR5$^-$, Green= MAIT CXCR5$^+$. Representative gating for populations found in Sup. Fig 3a. *p < 0.05, **p<0.01, ***p<0.001, ****p<0.0001 by two-tailed Mann-Whitney U test.

We next assayed for cytokines known to play a role in B cell differentiation or antibody production. *IL-21*

expression, although below the limit of detection (<LD) in PBMC MAIT and MAITfh groups, had a

trending increase in the tonsil MAITfh compared to the MAIT groups (p<0.07) (Fig. 2E). Notably,

although 2-5 fold lower than the tonsil Tfh group, both tonsil MAIT (median=361.3) and MAITfh

(Median=977.9) groups highly expressed *IL-21* relative to the tonsil CD8$^+$ CXCR5$^-$ group (Median=6.0)

(Fig. 2E). We found similar increases in *IL-6* expression in both PBMC and tonsil MAITfh compared to

MAIT and Tfh groups (Fig. 2F). Interestingly, both *TNFSF13B (BAFF)* and *IL-10* expression were

significantly higher in both MAIT and MAITfh groups within tonsils and PBMCs compared to Tfh cells

in their respective compartments (Fig. 2G and H). Taken together, CXCR5$^+$ MAIT (MAITfh) cells

express Tfh-associated cytokines and transcription factors at levels similar or higher than Tfh cells in their

respective compartments

CXCR5$^+$ MAIT cells highly express IL-21 compared to CXCR5$^-$ MAIT cells

We next wanted to investigate the relationship between IL-21 and IFNγ expression among MAIT subsets

as both cytokines are highly expressed and generally associated with a Tfh vs Th1 like phenotype. When

we stimulated PBMCs with PMA/Ionomycin, we saw a clear dichotomy between IFNγ and IL-21 positive

cells in all populations with very low frequency of double positive cells, and the vast majority being

IFNγ$^+$ (Fig. 2 I and J). MAITfh cells seemed to have a substantially lower frequency of IFNγ$^+$ cells

compared to conventional MAIT cells, though these data were limited by low PBMC-MAITfh cell

numbers, potentially biasing results. This effect was also notable in tonsils (p<0.001), where IFNγ

production in MAITfh cells was significantly reduced compared to that of MAIT cells (p<0.05), and IL21

production was significantly increased (p<0.001). Very few double-positive cells were recorded in either

population. Furthermore, the tonsil MAITfh population had a similar frequency of IL-21$^+$ cells compared

to conventional Tfh cells following stimulation (Fig. 2I and J). IFNγ$^+$ and IL21$^+$ frequency were low in all

unstimulated populations (fig. S6A-B). Conversely, when stimulated with a combination of 5-amino-6-D-

ribitylaminouracil (5-A-RU) and methylglyoxal to simulate MR1-dependent MAIT activation through

formation of the MAIT ligand, 5-OP-RU (*54*), we found minimal IL-21$^+$ MAIT or MAITfh cells and

statistically similar IFNγ$^+$ frequencies (fig. S7A-B). This suggests that IL-21 production by MAIT cells

may be MR1-independent, or dependent on alternative stimulation conditions. To confirm the IL-21

production capacity of MAIT cells and MAITfh cells within tonsils, we sorted MAIT (CD3$^+$ Vα7.2$^+$

MR1-Tetramer$^+$ CXCR5$^-$), MAITfh (CD3$^+$ Vα7.2$^+$ MR1-Tetramer$^+$ CXCR5$^+$), CD8$^+$ CXCR5$^-$ and GC

Tfh (CD4$^+$ CXCR5$^+$ PD1^{high}) populations, and stimulated mono-cultures of 17,000-40,000 cells with

PMA/Ionomycin (*55*). To account for differences in cell number per well we normalized all data to pg IL-

21 per cell. We found a trending increase in IL-21 production in the MAITfh over MAIT groups

(p<0.056) and no statistical difference in IL-21 production between MAITfh and GC Tfh populations

(Fig. 2K). Taken together, these data demonstrate the potent ability of the tonsil MAITfh population to

produce B-cell help cytokines and dichotomy in phenotype between classical MAIT cells and MAITfh

cells in both peripheral blood and tonsil tissue.

PD1$^+$ MAIT cells localize near germinal centers within tonsils

Having established that CXCR5$^+$ MAIT cells exhibit a Tfh-like phenotype *in vitro*, we next sought to

determine the *in vivo* spatial differences between CXCR5$^-$ and CXCR5$^+$ MAIT cells in relation to

germinal centers (GCs) in tonsil cryosections. MAIT cells were defined by co-staining of TRAV1-2 and

CD161 as described in Leng et al. 2019 (*56*). Anti-IgD was used to stain naïve B cells making up the

follicular mantle, and potential IgD$^+$ memory B cells in the marginal or superficial zone (*57*). Due to

difficulties with co-staining MAIT cells and CXCR5, we used PD1 to mark MAITfh and non-MAIT GC

Tfh cells (Fig. 3A). To confirm PD1 expression on Tfh cells, we stained tonsil cyrosections with anti-

CD4, anti-CD3, anti-IgD and anti-PD1. We found high co-expression of CD3$^+$CD4$^+$ cells and PD1 within

tonsil GCs, while minimum PD1 expression was found in the CD3$^+$CD4$^+$ cells outside of the follicular

mantle (fig. S8A). These data suggest that GC Tfh cells can be marked by high PD1 expression. PD1 has

also been used to mark GC light zone Tfh cells in human adenoid samples (*58*) and non-human primates

(*59*). Furthermore, flow cytometry analysis revealed significantly higher expression of PD1 among

CXCR5$^+$ (Median=44.1%, IQR=37.1%, 49.2%) vs CXCR5$^-$ (Median=10.5%, IQR=4.88%, 18.2%) MAIT

cells (Fig. 1E). ImageJ was used to quantify and measure PD1$^{+/-}$ MAIT cell distance to the edge of the

nearest GC, outlined in Fig. 3A (dashed white line). A total of 349 MAIT cells from n=5 tonsils were

analyzed with 32% ± 15.9 being PD1$^+$. In comparison, flow results of unstimulated tonsil cells showed

38% ± 14.9 of MAIT cells to be PD1$^+$, thus confirming the ability of the immunohistochemistry imaging

and quantification to assess this cell population. PD1$^+$ MAIT cells were generally located within or

closely surrounding GC's with a median distance of 66.8 μm from a GC edge compared to PD1$^-$MAIT

cells with a median distance of 220.2 μm (Fig. 3B). 28.4% of PD1$^+$ MAIT cells were located within GC's

and thus were recorded as 0 μm. PD1^{High} expressing MAIT cells tended to localize with other PD1^{High}

cells within GC light zones. In contrast, PD1^{low} or PD1$^-$MAIT cells were often found outside of the

IgD^{high} follicular mantle zone (Fig. 3A-yellow dashed line) and frequently in contact with IgD^{low} cells.

Overall, combined with phenotypic analysis above, these data reveal PD1$^+$ (and likely CXCR5$^+$) MAIT

Fig. 3. PD1^{high} MAIT cells preferentially locate near B cell follicles. (**A**) Location of MAIT cells within tonsils. Representative immunofluorescence images from tonsil cryosections stained with anti-IgD (blue), anti-PD1 (white), anti-CD161 (green), and anti-TRAV1-2 (red) and imaged on a Leica SP8 confocal microscope with 20X objective. Enlarged composite image below with MAIT cells (TRAV1-2$^+$CD161$^+$) denoted with arrows. White arrows indicate PD1$^+$MAIT cells. Yellow arrows indicate PD1$^-$MAIT cells. White dashed line outlines germinal center (GC) highlighted by PD1^{high} cells. Yellow dashed line outlines follicular mantle (FM) highlighted by IgD$^+$naïve B cells. All scale bars represent 100 μm. (**B**) Quantification of MAIT cell distance to edge of nearest GC. Tiled images from tonsil donors were visually assessed for co-staining of CD161, TRAV1-2 and then PD1. Distance to nearest GC was then measured using ImageJ. n=359 MAIT cells were quantified (PD1$^+$= 103; PD1$^-$ = 246) from n=5 tonsil donors. Data are represented as medians with dots representing single MAIT cells. *p < 0.05, **p<0.01, ***p<0.001, ****p<0.0001 by two-tailed Mann-Whitney U test.

MAIT cells repopulate mucosal and lymphoid organs following adoptive transfer in TCRα⁻/⁻ mice

To study the sufficiency of MAIT cells to provide B cell help in the context of mucosal immune

responses *in vivo*, we utilized an adoptive MAIT transfer model into T*cra^{tm1Mom}*/J (TCRα⁻/⁻) mice followed

by a prime-boost model of mucosal bacterial infection. Expansion of lung MAIT cells from WT C57Bl/6

(B6) mice using an intranasal *Salmonella* Typhimurium challenge have shown to result in MAIT cells

that retain function even after adoptive transfer (*40*). TCRα⁻/⁻ mice have been successfully used in T cell

transfer experiments to study the B cell help potential of non-Tfh CD4 T cells without the potential

masking effects of classic Tfh cells (*60*). We thus adoptively transferred expanded lung MAIT cells or

PBS as an injection control (Sham groups) from B6 into TCRα⁻/⁻ mice and then utilized an intranasal

prime-boost challenge model with live *Vibrio cholerae* O1 Inaba (*V.c*) that induces systemic and mucosal

LPS IgA and IgG responses (*61*) (Fig. 4A). MAIT cells have been shown to be activated during cholera

infection *in vivo* (*22*), and *V. cholerae* can biosynthesize riboflavin and thus can activate MAIT cells via

MR1 presentation of riboflavin intermediates (*62*). CXCR5 expression was present but low on D7 *S.*

Typhimurium expanded MAIT cells prior to adoptive transfer (fig. S10A-B). To confirm MAIT cells

were successfully transferred into recipient mice, we measured MAIT frequencies and total cell numbers

by flow cytometry of digested lung tissue and spleen (Fig. 4B and fig. S9A-C). The frequency and total

cell count of lung MAIT cells at Day 42 in MAIT-*V.c* and MAIT-PBS groups were statistically similar to

the WT- *V.c* group (Fig 4C and fig. S9B). Importantly, no MAIT cells were recorded in either Sham-*V.c*

or Sham-PBS group. Similar MAIT cell counts were also observed in the spleen (fig. S9C), thus

demonstrating WT physiologic MAIT frequencies and repopulation of various organs following adoptive

transfer of lung MAIT cells. Furthermore, we found no differences between all TCRα⁻/⁻ groups in total

non-MAIT TCRβ⁺ T cells, thus confirming no significant expansion of contaminating non-MAIT T cells

following adoptive transfer (fig. S9D). We next analyzed the expression of CXCR5 in CD8 T cells, CD4

T cells and MAIT cells from the WT-*V.c* and MAIT cells from the MAIT-*V.c* and MAIT-PBS groups

(Fig. 4D and E). MAIT cells from WT-*V.c*, MAIT-*V.c* and MAIT-PBS groups, though they had decreased

CXCR5 MFI compared to WT CD4 T cell populations, had significantly increased CXCR5 expression

compared to the control CD8 T cell group in lungs, suggesting a potential mouse MAITfh population

(Fig. 4D and E). Taken together, we found that adoptive transfer of expanded MAIT cells into TCRα$^{-/-}$

mice repopulates MAIT cells, including CXCR5$^+$ MAIT cells in lymphoid and mucosal organs, to levels

similar to that of WT mice.

MAIT cell adoptive transfer drives increase in plasmablast differentiation and memory B cell

development

We next sought to determine the impact of MAIT adoptive transfer on mucosal B cell differentiation. Day

42 digested lung suspensions were analyzed by flow cytometry with B cells being defined as B220$^+$

lymphocytes (fig S11A). Interestingly, total B220$^+$ cell frequency (Fig. 4F) and total B cell count per lung

(fig. S11B) were decreased in both the MAIT-*V.c* and MAIT-PBS groups compared to Sham-*V.c*, Sham-

PBS and WT-*V.c* groups. Similar decreases in IgD$^+$ naïve B cell frequency were observed in MAIT

transfer groups (Fig. 4G), although no statistical differences were observed in total naïve B cells (fig.

S11C). In contrast, we found that both MAIT-*V.c* and MAIT-PBS groups had higher Memory B cell

(B220$^+$ IgD$^-$ CD27^{mid} CD38$^+$ CD138$^-$) and Plasmablast (PB)/Plasma cell (PC) (B220^{low} IgD$^-$ CD27$^+$

CD38$^+$ CD138^{++}) frequency as a percentage of total B cells compared to Sham-*V.c,* Sham-PBS and WT-

V.c groups (Fig. 4H and I). Total Memory B cell count per lung was not statistically different between

groups with significant variability observed in the MAIT-*V.c* group (fig. S11D), though we saw a

trending increase in total PB/PC cell count per lung in MAIT-*V.c* compared to Sham-*V.c* (p=0.097) and

WT-*V.c* (p=0.179) groups (fig. S11E). Overall, these data suggest that in the absence of αβ T cells,

transfer of activated MAIT cells help induce B cell differentiation from naïve B cells to PB/PCs or

memory B cells.

Fig. 4. Adoptively transferred MAIT cells expand in lungs of TCRα$^{-/-}$ mice driving B cell differentiation. (**A**) MAIT adoptive transfer and *V.c* intranasal challenge time-line. (**B**) Representative FACS plots of MAIT cells (TCRβ^{+} MR1-Tetramer^{+}) from lung tissue of WT-*V.c* (left), MAIT-*V.c* (middle), and Sham-*V.c* (right) mice at Day 42 post initial *V. cholerae* challenge. (**C**) MAIT cell frequency as percentage of CD3^{+}. Representative flow cytometry histograms (**D**) and Mean fluorescence intensity quantification (**E**) of CXCR5 in WT-*V.c* CD8 T cells (grey), CD4 T cells (green) and MAIT cells (orange), MAIT-*V.c* MAIT cells (red), and MAIT-PBS MAIT cells (blue). (**F**) Frequency of B220^{+} B cells among live lymphocytes. Frequency of IgD^{+} naïve B cells (**G**), IgD^{-} CD27^{+} CD38^{+} CD138^{-} Memory B cells (**H**) and IgD^{-} CD27^{+} CD38^{+} CD138^{++} PB/PCs (**I**) as percent of B220$^{low-high}$ lymphocytes. Data are represented as Median with IQR from 4 independent experiments. n= 4-15 mice per group. *$p < 0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by two-tailed Mann-Whitney *U* test. Legend in black box denotes experimental groups: MAIT-*V.c* = TCRα$^{-/-}$ plus MAIT Transfer-*V.c* challenge, MAIT-PBS = TCRα$^{-/-}$ plus MAIT Transfer-PBS challenge, Sham-*V.c* = TCRα$^{-/-}$ plus Sham Transfer-*V.c* challenge, Sham-PBS = TCRα$^{-/-}$ plus Sham Transfer-PBS challenge, WT-*V.c* = WT C57Bl/6J plus Sham Transfer-*V.c* challenge.

Increase in V. cholerae specific IgA responses following MAIT transfer

V.c-specific antibodies in blood is a surrogate for protection against cholera (*63, 64*). Specifically, the

vibriocidal assay, a measurement of complement-fixing bactericidal antibody activity, has been in use

since the mid twentieth century (*65*) and shown to be the best marker of protection (*66, 67*). In addition to

the vibriocidal titer, cholera toxin (CT) and lipopolysaccharide (LPS) serum ELISAs have been widely

used to estimate cholera incidence (*67–70*). The O-specific polysaccharide (OSP) component of LPS and

CT are the immunodominant antigens following cholera infection (*71*). Therefore, in order to measure the

impact of MAIT transfer on pathogen-specific antibody kinetics we measured serum IgG, IgA, and IgM

antibody responses against whole *V.c*-lysate, OSP, and CT. We found comparable *V.c*-lysate IgA

responses in the MAIT-*V.c* and WT-*V.c* groups particularly following *V.c* re-challenge (D28), suggesting

development of a memory response (Fig. 5A). Day 42 endpoint ELISAs reveal a significantly higher *V.c*-

lysate IgA responses in the MAIT-*V.c* group compared to Sham-*V.c* (p<.0001), and MAIT-PBS (p<0.05)

groups, and similar responses compared to the WT-*V.c* group (p=0.58) (Fig. 5C). Importantly, minimal

V.c-lysate IgA responses were observed in the Sham-PBS group. MAIT-*V.c* and MAIT-PBS groups had

no significant differences in *V.c*-lysate IgG (p=0.66, Fig. 5B and D) and IgM (p=0.08, fig. S12A and B)

responses, despite higher compared to Sham transfer groups (p<0.01), thus indicating a non-specific

response induced by MAIT transfer. We saw a non-significant (p=0.054) higher OSP IgA response in the

MAIT-*V.c.* group compared to the Sham-*V.c* group, which can largely be attributed to high values in one

mouse (Fig. 5E and G). No differences were observed in OSP-IgG responses between MAIT-*V.c* and

Sham-*V.c* groups (p=0.64), Fig. 5F and H). Notably, even Sham-*V.c* TCRα-/- mice were able to mount

(albeit lower than WT) OSP IgG responses, indicating that a lack of T cells blunted, but did not eliminate,

the ability to mount an IgG response to a polysaccharide antigen, which is classically T-independent. In

contrast, OSP-IgM levels were similar between the MAIT-*V.c* and Sham-*V.c* (p=0.33), and MAIT-*V.c* and

WT-*V.c* groups (p=0.11), though were lower in the Sham-*V.c* compared to the WT-*V.c* (p=0.03) groups.

All *V.c* infected groups had higher OSP-IgM responses compared to MAIT-PBS (p<0.05) and Sham-PBS

(p<0.01) groups, indicating a specific but primarily T-independent response (fig. S12E and F). Overall,

very low levels of CT IgA and IgG were observed in all TCRα$^{-/-}$ groups (Fig. 5I-L) compared to the WT-

V.c group, supporting the role of classical helper T cells in protein antigen B cell help, a response that

adoptive transfer of MAIT cells did not rescue. Analysis of total serum IgA revealed a highly variable but

substantial increase in the MAIT-*V.c* group over both Sham-*V.c* (p<0.0001)) and WT-*V.c* groups

(p=0.01), fig. S12H). No statistically significant changes were observed in total IgM and IgG between

MAIT-*V.c* and Sham-*V.c* groups (fig. S12G and I). To confirm the phenotype observed in the serum

corresponded to the mucosa, Bronchoalveolar lavage (BAL) fluid from a subset of MAIT-*V.c* and Sham-

V.c mice were assayed for *V.c* lysate IgA, IgG, and IgM responses. We found increased anti-lysate

responses in all isotypes in the MAIT-*V.c* compared to the Sham-*V.c* group which corresponded with

findings from serum responses (fig. S12J-L).

Fig. 5. MAIT cells promote *V. cholerae*-specific IgA antibody responses in TCRα[-/-] mice. Serum ELISAs against *V. cholerae* whole lysate, OSP and CT. (**A-D**) IgA and IgG *V. cholerae* lysate ELISAs from weekly cheek bleed (A & B) and Day 42 endpoint serum (C & D). (**E-H**) IgA and IgG OSP ELISAs from weekly cheek bleed (E & F) and Day 42 endpoint serum (G & H). (**I-L**) IgA and IgG CT ELISAs from weekly cheek bleed (I & J) and Day 42 endpoint serum (K & L). Data are represented as ELISA units normalized to a pooled serum positive control from WT B6 mice challenged with *V. cholerae*. (**M**) Vibriocidal titer of Day 42 endpoint serum. Data are represented as Median with IQR from 5 independent experiments. n= 6-22 mice per group. *$p < 0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by two-tailed Mann-Whitney *U* test. Legend in black box denotes experimental groups.

We further utilized the vibriocidal assay to measure complement-mediated bactericidal activity, the best

studied and most commonly used correlate of protection from cholera in humans and a proxy for

protection in murine cholera challenge models (*66*, *67*, *72*). We found a trending increase in vibriocidal

titer in the MAIT-*V.c* group compared to the Sham-*V.c* group (p=0.069), and no difference when

compared to the WT-*V.c* group (Fig. 6M). Unlike the ELISAs, no background vibriocidal responses were

observed in the MAIT-PBS group, demonstrating the specificity of this assay. Taken together, adoptive

transfer of MAIT cells into T-cell deficient mice rescued deficiencies in *V. cholerae*-specific IgA and

promoted slight increases in vibriocidal antibody responses.

Discussion

MAIT cells are widely appreciated for their ability to respond rapidly to microbial antigens and cytokines

through potent production of Th1 and Th17-associated inflammatory cytokines and cytotoxic molecules.

Recent studies have highlighted the capacity of human MAIT cells to provide B cell help *in vitro*, and

promote B cell differentiation and antibody production in animal models (*23*, *25*, *26*). Despite these

findings, whether there remains a pre-defined MAIT subset capable of B cell help remains uncertain.

Additionally, many studies have detected MAIT cells within mucosal lymphoid tissues but have yet to

understand their role in these environments (*33*, *41*, *73*). In this study we begin to define the role of MAIT

cells in human mucosal lymphoid organs and uncover a novel T follicular helper-like MAIT population

defined by CXCR5. This MAITfh subset highly expresses PD1, ICOS, *BCL6*, and IL-21, and

preferentially locates near germinal centers. We further add to the body of literature proving the ability of

MAIT cells to provide B cell help *in vivo* through sufficiency experiments following mucosal challenge.

Here we show that adoptively transferred MAIT cells can rescue microbe specific IgA responses, and

promote B cell differentiation in the absence of other αβ T lymphocytes.

We report a subset of CXCR5$^+$ MAIT (MAITfh) cells that are enriched in mucosal lymphoid organs and

represent a small percentage of peripheral blood MAIT cells (Fig. 1C). We further report that MAITfh

cells highly express the B cell help co-stimulatory molecules, PD1 and ICOS, compared to CXCR5$^-$

MAIT cells (Fig 1D and E). In addition, we show high relative expression ($\approx$30-45 fold) of *CD40L* in

both MAIT and MAITfh groups compared to the control CD8+ population (fig. S5B). PD1, ICOS and

CD40L expression are a hallmark of GC Tfh cells (*74*). PD1, although long considered a marker of T cell

exhaustion, has an important role in Tfh GC positioning and function (*5*). ICOS and CD40L engagement

with ICOS Ligand and CD40 expressed on B cells is essential for GC formation and Tfh activation (*6, 7*).

CD40L is also implicated in MAIT induced maturation of dendritic cells (*75*). Based on the Tfh

costimulatory molecule profile, we hypothesized that MAITfh cells may share further transcriptional

similarities with Tfh cells. In support, we found that MAITfh cells in PBMCs have higher *BCL6*

expression than MAIT and Tfh cells and a trending increase in *BCL6* expression in tonsils (Fig. 2E).

BCL6 is considered the master regulator of the Tfh lineage (*76*) and is similarly required for iNKTfh B

cell help responses (*18*). Although further research into the transcriptional regulation of MAITfh cells is

necessary to understand the requirements for differentiation, plasticity and function, these data support a

B cell help transcriptional phenotype among the MAITfh subset.

We report MAITfh expression of the B cell help cytokines *IL-21*, *IL-10*, *IL-6* and *BAFF* (Fig. 2 and fig.

S3). Specifically, we found trending increases in *IL-21* and *IL-6* expression in MAITfh compared to

MAIT cells in tonsils, while *BAFF* and *IL-10* were highly expressed in both PBMC and tonsil MAIT cells

compared to Tfh controls (Fig. 2). IL-21 plays an important role in T dependent B cell differentiation,

class switching and antibody production (*77*) and together with IL-6 promotes Tfh differentiation (*78*).

IL-10 promotes antibody production and class switching (*79*). IL-21 expression in MAIT cells has been

demonstrated in low concentrations in PBMCs (*23*), and in a subset of PD1^{High} CXCR5$^-$ MAIT cells in

pleural effusions from Tuberculosis patients (*24*). This study also presents a small subset of CXCR5$^+$

PD1^{high} MAIT cells, though they do not investigate whether this population expressed IL-21 as well (*24*).

In contrast, we show that tonsil CXCR5$^+$ MAIT cells have a higher frequency of IL-21$^+$ IFNγ^- cells (Fig.

2J) and produce more IL-21 than CXCR5$^-$ MAIT cells (Fig. 2K). It is unclear whether PD1^{High} CXCR5$^-$

MAIT cells in pleural effusions are transcriptionally or developmentally related as potential precursors to

CXCR5$^+$ MAIT cells found in lymphoid tissues. Furthermore, the B cell survival cytokine, BAFF, is

integral in T independent class switching and differentiation (*80*), and is similarly produced by iNKTfh

cells to promote B cell responses (*18*). Thus, it is possible MAITfh cells can aid in B cell help through

cognate interactions via direct MR1-TCR, co-stimulatory molecule engagement and cytokine expression,

or through non-cognate interactions such as promoting Tfh differentiation or licensing of dendritic cells.

We have observed that PD1^{high} MAIT cells preferentially locate within or around GCs in human tonsils

(Fig. 3A and B). This is noteworthy since spatial positioning of T and B cells within lymphoid tissues is

critical in understanding their phenotype. For instance, CXCR5$^+$ PD1^{High} GC Tfh cells reside primarily in

the light zone of GCs where they promote affinity maturation and differentiation of GC B cells leading to

high affinity antibody responses (*74*). Thus, by combining imaging results with *in vitro* MAITfh

expression of B cell help co-stimulatory markers and cytokines (Fig. 1 and 2), we can conclude that

MAITfh cells share both phenotype and positioning with traditional GC Tfh cells. Thus we hypothesize

that this population may be involved in cognate or contact-dependent B cell help as is seen in other

innate-like Tfh subsets, such as iNKTfh (*12*). Though as this population is relatively small in tonsils,

further research will need to address the *in vivo* relevance to overall mucosal antibody production, the role

of MAITfh-B cell interactions in affinity maturation, and development of B cell memory.

We report that adoptive transfer of murine MAIT cells boosts pathogen-specific antibody responses in the

absence of other αβ T lymphocytes. We further previous work on MAIT-B cell help responses in animal

models (*25*, *26*), by using an adoptive transfer model to show that MAIT cells are sufficient to increase

functional anti-bacterial antibodies in the context of a mucosal challenge. Most notably, we show that

adoptive transfer of MAIT cells rescues the deficiency in *V.c*-specific IgA of infected TCRα$^{-/-}$ mice when

compared to infected WT mice. Examination of antibody kinetics of *V.c* specific IgA in mice receiving

MAIT transfer suggest a memory phenotype, based on substantial increases on Day 35 (one-week post re-

challenge) (Fig. 5A), and is associated with higher lung PB/PC and memory B cell frequencies (Fig. 4H

and I). Interestingly, we also report MAIT-*V.c* mice had significant systemic *V.c* specific IgA and total

IgA responses compared to Sham-*V.c* mice (Fig. 5A and fig. S12H). These data may be a result of the

mucosal nature of the intranasal challenge, which has shown to induce both mucosal and systemic

immunity in other bacterial infections (*81*). Notably, increases in total IgA in MAIT-PBS mice suggests

MAIT cells may specifically promote IgA class switching. In support, pulmonary MAIT cells in mice

have recently been shown to produce both IL-10 (*82*) and TGF1-β (*83*), key cytokines in IgA class

switching (*84*). Although CXCR5 staining of lung MAIT cells suggests a similar Tfh-like MAIT

phenotype in mice (Fig. 4D and E), limitations of available reagents for identifying murine MAIT cells by

IHC precludes our ability to confirm a MAITfh subset in mice. Future analysis of lung and lymph node

MAIT cells is necessary to confirm a MAITfh subset in mice and its potential role in IgA class switching

and antibody production. Given that systemic *V. cholerae* specific IgA and vibriocidal responses strongly

correlate with protection against subsequent cholera re-infection (*64*, *66*, *67*), these are promising data in

the context of mucosal vaccines, where IgA is often the primary means of protection, and development of

long-lasting protective memory responses is limiting (*85*). These data support further research into the

potential adjuvant activity of MAIT ligand administration during mucosal infection or vaccination to

enhance humoral immune responses.

In addition to promoting microbe-specific IgA responses, we report amplification of non-specific

antibodies following MAIT transfer. We report similar *V. cholerae* lysate IgG (Fig. 5B and D) and IgM

(fig. S11A and B), OSP (Fig. 5E-H), CT (Fig. 5I-L), total IgA (fig. S11H), and similar changes in B cell

subset frequencies in our MAIT-PBS and MAIT-*V.c* groups (Fig. 4F-I). Thus, regardless of *V. cholerae*

infection, adoptively transferred MAIT cells are promoting B cell differentiation and antibody production.

Similar MAIT amplification of non-specific responses were reported in non-human primates (*26*) and

FcγRIIB$^{-/-}$ mice (*25*). Specifically, Rahman et al. reported increased total IgM and IgG antibody

production and increased IgD$^-$ B cells (indicating class switching) when non-human primate MAIT cells

were co-cultured with B cells. Murayama et. al 2019 show increases in total IgG and anti-dsDNA IgG and

IgA when MAIT cells were co-cultured with B cells from FcγRIIB$^{-/-}$ mice. They also report reduction in

anti-dsDNA IgG in MR1$^{-/-}$ (MAIT deficient) FcγRIIB$^{-/-}$ mice, though total IgG was not published. It

should be noted that in all studies, MAIT cells were either activated *in vitro* in direct co-culture, or *in vivo*

in the absence of regulatory T cells (TCR$\alpha^{-/-}$ mice). Therefore, such responses may be a result of lack of

either MAIT and/or B cell regulation leading to aberrant antibody responses. This has interesting

consequences in the setting of mucosal immunity and in autoimmunity. In a setting of proper T and B cell

regulation, MAIT cells may help to promote anti-microbial responses and therefore could potentially be

targeted as a vaccine adjuvant. Conversely, in a dysregulated environment, such as in FcγRIIB$^{-/-}$ or TCRα^-

$^{/-}$ mice, MAIT cells may enhance aberrant antibody production against self or commensal microbes thus

enhancing the pro-inflammatory autoimmune state. These aberrant responses open many questions into

cell extrinsic MAIT regulation which remains relatively unstudied.

There are a number of limitations that need to be addressed when considering the results of this study. In

our human *in vitro* experiments, we primarily compare PBMCs from healthy adults ages 18+ to tonsils

from pediatric patients, ages 2-16, undergoing tonsillectomy. MAIT cells expand immediately after birth

and reach adult frequencies by approximately 10 years of age, and most reach maturity (CD45RO$^+$) by 6

months of age (*86*). Therefore, it is possible that age may explain low MAIT percentage in tonsil samples,

though analysis of 5 paired PBMC and tonsil samples pediatric donors revealed little differences in MAIT

frequency when compared to the healthy donor adult PBMCs (fig. S4). Limited MAIT frequency in

tonsils also limited our abilities to perform mechanistic analysis of our MAITfh populations. Future co-

culture studies with B cells and expanded MAITfh cells may be useful to understand the MAIT-B cell

help mechanism. Regarding our animal studies, a limitation of this study is the under sampled analysis of

mucosal antibody measurement, as we focused primarily on systemic responses in serum. While mucosal

secretory IgA (sIgA) and circulating IgA responses may differ, measurement of systemic antibodies is

standard practice for mucosal vaccine response testing in humans, and *V.c* specific serum IgA response

and vibriocidal titer strongly correlate with protection in humans (*64*, *66*, *67*). Additionally, our limited

analysis of BAL antibody responses correlated well with serum results. Thus, these serum responses are

likely relevant to mucosal protection, though lack of conclusive protective immunity via survival curves

or colony counts limits this finding. Secondly, our study lacks insight into the mechanism and location of

MAIT B cell help in our murine model. Future studies, potentially using congenic markers, are necessary

to address whether MAIT cells provide help within local lymph nodes, MALT or within the mucosal

lamina propria. Additionally, the nature of *the V. cholerae* antigens targeted by the IgA antibodies

remains unknown. Despite previous evidence showing association of human MAIT cells with

polysaccharide specific antibody responses (*22*, *35*), we found only moderate changes in OSP antibody

responses in our MAIT transferred mice, suggesting an alternate antigen target. Future studies are

necessary to address the whether MAIT cells, like other innate-like T cells, promote antibody responses to

specific antigen targets.

In conclusion, we identify a Tfh like MAIT population enriched near germinal centers of mucosal

lymphoid tissues that express cytokine and transcriptional profiles consistent with B cell help. In addition,

we show in an animal model that MAIT cells are sufficient to rescue pathogen-specific IgA antibody

responses following mucosal challenge in the absence of other classical T cells. Our work strengthens a

growing body of evidence supporting the capacity of MAIT cells to aid in B cell help and antibody

mediated immunity, and support the expanding phenotypic heterogeneity of MAIT cells (*87*). As MAIT

cells are enriched at the mucosa and can be activated with specific ligands, we hypothesize that they may

be a promising target to enhance mucosal vaccines.

Materials and Methods

Peripheral blood and tonsil collection

Healthy adult blood was collected from residual leukocyte packs following blood donation (ARUP, Sandy, UT), and PBMC's were isolated by density gradient centrifugation (Lymphoprep, StemCell Technologies). Residual palatine tonsil samples were acquired from pediatric patients, ages 2-16, undergoing routine tonsillectomies for tonsillar hyperplasia or recurrent tonsillitis. All tonsil samples were stored on ice in complete media (RPMI 1640 with 10% FBS, 1% penicillin/streptomycin, and 15 mM HEPES) (R10) and processed within 2 hours of surgery. 1 cm x 1 cm pieces were cut and frozen in O.C.T. Compound (Fisher) at -80°C for later use. The remaining tonsil tissue was minced in media and dissociated using a 100 μM cell strainer. Tonsil mononuclear cells were isolated by density gradient centrifugation (Lymphoprep, StemCell Technologies). PBMC's from pediatric donors were obtained during surgery with informed consent granted by a parent/guardian. PBMC and tonsil mononuclear cells were frozen in media containing 25% FBS and 10% DMSO at -80°C for later use. All human samples were de-identified prior to receipt, and the research protocol was deemed exempt by the University of Utah Institutional Review Board (Protocol 100683).

Flow cytometry and cell sorting

Prior to surface staining, human and mouse single cell suspensions were labeled with Fixable Viability Dye eFluor 780 (eBioscience) according to manufacturer's protocol to delineate live cells. Cells were washed and incubated with fluorochrome conjugated antibodies for 20 min at room temperature (RT). Mouse cells were incubated in anti-mouse CD16/CD132 (Fc block) for 15 minutes prior to surface staining. Mouse and human MR1-Tetramers (NIH Tetramer Core) diluted 1:400 were added to surface stain. Prior to mouse MR1-Tetramer staining, cells were incubated for 15 minutes with unlabeled i6FP-Tetramer (NIH Tetramer Core) to minimize non-specific binding. For intracellular cytokine staining, cells were fixed and permeabilized using Foxp3/Transcription Factor Staining Buffer set (eBioscience)

according to manufacturer's protocol and incubated with fluorochrome conjugated antibodies for 40 min

at RT. Cells were analyzed using BD LSR Fortessa or Cytek Aurora for phenotypic analyses, or Aria II

(BD Biosciences) for cell sorting. The following monoclonal antibodies to human were used : CD3-

AF700 (clone OKT3), CD8-FITC/BV605 (cloneSK1, TCR Vα7.2-BV711/PE-Cy7 (clone 3C10), PD1-

BV605/AF700 (clone EH12-2H7), ICOS-PerCP-Cy5.5/BV605 (clone C398.4A), CXCR5-PE-Cy7/APC

(clone J252D4), IL-21-APC (clone 3A3-N2), IFNγ-BV421 (clone B27), IL-6-FitC (clone MQ2-13A5),

IL-10-PerCP-Cy5.5 (clone JES2-9D7), CD161-AF488 (clone HP-3610) (Biolegend), CD4-BUV496

(clone SK3), CD3-BUV395 (clone SK7) (BD Biosciences). The following monoclonal antibodies to

mouse were used: CD3-PE-Dazzle594 (clone 17A2), TCRβ-BV421 (clone H57-597), CXCR5-BV605

(clone L138D7), CD44-BV650 (clone IM7), TCR$\gamma\delta$-PE-Cy7 (clone GL3), CD8-APC (clone 53-6.7),

CD45R-PE-Cy5 (clone RA3-6B2), IgD-BV510 (clone 11-26c.2a), CD138-BV605 (clone 281-2), CD38-

PE-Cy7 (clone 90, CD27-APC (clone LG-3A10) (Biolegend), CD4-FITC (clone GK1.5) (Tonbo

Biosciences).

In vitro stimulation assays

For flow cytometry intracellular cytokine staining, 1-2 x 10^6 mononuclear cells isolated from blood or

tonsil samples were cultured in a round bottom 96 well plate at 37°C 5% CO_2 in R10 media for 6 hours

with or without 200 ng/ml phorbol myristate acetate (PMA) and 1.0 μg/ml Ionomycin (Sigma-Aldrich) or

0.5 μM/ml 5-amino-6-D-ribitylaminouracil (5-A-RU) (courtesy of Jeffrey Aubé, UNC) and 50 μM/ml

Methylglyoxal (Sigma). Brefeldin A Solution (Biolegend) was added at 1X for the final 4 hours. For

measurement of IL-21 production, tonsil mononuclear cells were sorted into CD8$^+$ CXCR5$^-$, CD4$^+$

CXCR5$^+$PD1^{High}, MR1-Tet$^+$, Vα7.2$^+$, CXCR5$^-$, MR1-Tet$^+$, Vα7.2$^+$, CXCR5$^+$ populations and 4x10^4 cells

in R10 media were cultured in a round bottom 96 well plate for 48 h with 20 ng/ml PMA and 1.0 μg/ml

Ionomycin as described in Shen et al (*55*). Plates were centrifuged and cell supernatant was isolated for

ELISAs.

qRT-PCR

PBMC and tonsil CD8$^+$ CXCR5$^-$, CD4$^+$ CXCR5$^+$, MR1-Tet$^+$ Vα7.2$^+$ CXCR5$^-$, MR1-Tet$^+$ Vα7.2$^+$

CXCR5$^+$ cells were sorted into TRIzol Reagent (Qiagen). Total RNA was isolated using the RNeasy

Micro kit and cDNA synthesized a High-Capacity cDNA Reverse Transcription Kit (Applied

Biosystems). Targeted qPCR was performed using the QuantStudio 6 Real-Time PCR system using the

following Taqman Primers (Applied Biosystems): BCL6 (Hs00153368_m1), Tnfsf13b

(Hs00198106_m1), IL6 (Hs00174131_m1), IL10 (Hs00961622_m1), IL21 (Hs00222327_m1), TNFα

(Hs00174128_m1), TGFB1 (HS00998133_m1), Tcf7 (Hs01556515_m1), Tbx21 (Hs00894392_m1),

Runx3 (Hs01091094_m1), Rorc (Hs01076112_m1), Perforin (Hs00169473_m1), Pdcd1

(Hs01550088_m1), Il17a (Hs00174383_m1), Ifnγ (Hs00989291_m1), Icos (Hs00359999_m1), Gzmb

(Hs00188051_m1, Cxcr5 (Hs00173527_m1), Cd40l (Hs00163934_m1), Cd38 (Hs01120071_m1),

Blimp1 (Hs00153357_m1), Ascl2 (Hs00270888_s1), ActB (Hs99999903_m1).

Single cell TCR sequencing

MAITs from PBMC's and tonsils were single cell sorted using the Aria II cell sorter (BD Biosciences)

directly into One Step RT-PCR reaction mix (NEB) loaded in MicroAmp Optical 96-well reaction plates

(Applied Biosystems). MAITs were defined as CD3$^+$ Vα7.2$^+$ MR1-Tetramer$^+$ cells and sorted based on

CD161 expression. Following reverse transcription and preamplification reaction, a series of three nested

PCR's were run using primers for TCR sequence and gene expression as described (*45*). Subsequent

sequencing data analysis was performed as described (*45*).

ELISAs and Vibriocidal assay

IL-21 concentration from tonsil mono-cultures was measured using Human IL-21 ELISA MAX Deluxe

kit (Biolegend) according to manufacturer's protocol. Serum *V.c* Lysate, OSP:BSA and CT ELISAs were

performed as follows. Nunc Maxisorp flat bottom 96 well plates (Invitrogen) were coated with *V.c* Lysate

or OSP:BSA (gift from Dr. Edward Ryan (Massachusetts General Hospital, Boston, MA) (1.0 μg/ml) in

PBS and incubated overnight at 4°C. For CT ELISAs, Nunc plates were first coated with

Monosialoganglioside G_{M1} (Sigma-Aldrich) (1.0 μg/ml) in 50mM carbonate buffer overnight at 4°C.

Plates were subsequently blocked with 1% bovine serum albumin (BSA) (Sigma-Aldrich) and incubated

overnight with CT (Sigma-Aldrich) (2.5 μg/ml) in carbonate buffer. All plates were then blocked with 1%

BSA and 25 μl of mouse serum diluted 1:10 in 0.1% BSA-0.5% Tween 20 in PBS (BSA-PBST) was

added for 2 hours. Plates were incubated with anti-mouse IgG (Invitrogen), IgA (Invitrogen), or IgM (Life

Technologies) HRP conjugate diluted 1:1000 in BSA-PBST for 2 hours. 100 μl of TMB substrate

(Thermo Fisher) was added and plates were immediately read kinetically at 405 nm (7 min x 1 min

intervals) using plate reader (Biotek). ELISA measurements were recorded as max interval slope and

normalized to pooled positive mouse sera included on each plate. Total Ig ELISAs were performed as

described above except Nunc plates were pre-coated with anti-mouse IgA (clone RMA1), anti-mouse IgG

(clone Poly 4053), or anti-mouse IgM (clone RMM-1) (Biolegend) and standard curves were generated

using 2-fold dilutions of purified Mouse IgA, IgM and IgG (Southern Biotech). Plates were incubated

with corresponding anti-mouse HRP antibodies and TMB substrate was neutralized after 10 min with 0.2

N HCl. Plates were read at 605 nm and concentrations were calculated based on the standard curve.

Vibriocidal titer was measured as described (*88*). *Vibrio cholerae* serotype 01 Strain El Tor Inaba N16961

grown in LB media was used as target organism for assay, and serum samples were diluted to a starting

concentration of 1:5.

Immunofluorescence staining

Unfixed pediatric tonsil samples frozen in OCT medium were cut into 9μm thick sections using a cryostat

(Leica) and stored at -80°C. Tonsil sections were brought to room temperature and then fixed at -20°C for

10 min in pre-cooled 50:50 Acetone:Methanol solution. Sections were dried and then rehydrated in PBS

for 5 min with gentle shaking. The sections were then blocked in 2% normal mouse serum (Invitrogen)

for 1 hour and subsequently stained overnight at 4°C with mouse anti-human TCR Vα7.2-PE (Clone

3C10, Biolegend), mouse anti-human CD161-AF488 (Clone NKR-P1A, Biolegend), mouse anti-human

IgD-BV421 (Clone IA6-2, Biolegend) and mouse anti-human PD1-APC (Clone EH1202H7, Biolegend)

all diluted 1:100. Sections were washed 3x in PBS and mounted using ProLong Gold Antifade Mounting

media (Thermo Fisher). Full tonsil cross section images were acquired using the Leica Sp8 Confocal

microscope and then processed and analyzed using ImageJ.

Bacterial culture and V. cholerae lysate preparation

The attenuated *Salmonella* Typhimurium strain BMM50 was a gift from Dr. Stephen McSorely (UC

Davis, Davis, CA). BMM50, like previously published *S.* Typhimurium BRD509 (*39*), has a deleted *aroA*

gene and intact riboflavin pathway. *Vibrio cholerae* serotype 01 Strain El Tor Inaba N16961 was a gift

from Dr. Edward Ryan (Massachusetts General Hospital, Boston, MA). All strains were inoculated from

single colonies into Luria Bertani (LB) broth shaking at 220 RPM 37°C overnight. They were then

subcultured 1:10 in fresh LB and incubated for 3-4 hours shaking at 37°C until log phase was reached.

Cultures were washed 2x in sterile PBS and normalized to 0.1 OD600 using a microplate absorbance

spectrophotometer (BioRad). *V. cholerae* at 0.1 OD600 was used as inoculum for intranasal challenge

model. *S.* Typhimurium BMM50 was diluted 1:10 with PBS to an OD600 of 0.01 for adoptive transfer

inoculum. Inoculums ranged from 1×10^6-1×10^7 colony forming units (CFU) confirmed by plate counts. *V.*

cholerae lysate for ELISAs was prepared as follows. *V. cholerae* was cultured from a single colony in LB

broth shaking at 220 RPM 37°C overnight. *V. c* culture was then pelleted and washed 3x with cold sterile

PBS. Following resuspension in PBS, sample was sonicated at 40% power 4 x 1 min intervals, keeping on

ice for 1 minute between runs. Sonicated sample was then centrifuged at 15000 rpm x 10 min at 4°C, and

the supernatant removed and sterilized using 0.2 μm syringe filter (Thermo Fisher). Protein content was

measured using 260/280 absorbance ratio with Take3 Micro-Volume Plate reader (BioTek)

Mice

T*cra*[tm1Mom]/J mice (*89*) (on a B6 background) were obtained from The Jackson Laboratory (Bar Harbor, ME) and bred at the University of Utah Comparative Medicine Center mouse facility. WT C57BL/6J mice were obtained directly from The Jackson Laboratory prior to the start of the experiment. 6-10-week-old female and male mice were used in adoptive transfer studies. All mice were housed under specific pathogen-free conditions. Experiments were performed in strict accordance with the NIH guide for the Care and Use of Laboratory animals and institutional guidelines for animal care at the University of Utah under approved protocol #17-01011.

Adoptive transfer

Adoptive transfer of pulmonary MAIT cells was adapted from protocol described (*40*). C57BL/6 mice were infected i.n. with 10^6 CFU *S.* Typhimurium BMM50 suspended in 50 µl sterile PBS (25 µl per nares). Mice were rested for 7 days to allow for MAIT expansion, in which mice were weighed daily and evaluated for signs of clinical pneumonia symptoms (respiratory distress, inactivity, ruffled fur, hunched posture). Mice with loss of >20% of body weight and/or severe clinical symptoms were euthanized. On day 7, mice were euthanized and the lungs were perfused with 10 mls of PBS through the heart and then removed. Single cells suspensions of lungs were prepared using the gentleMacs Lung Dissociation kit, mouse (Miltenyi Biotech) according to the manufacturer's protocol. Red blood cells were lysed using ACK Lysis buffer (Thermo Fisher), and single cell suspensions were stained and sorted using the BD FACS Aria II. MAIT cells were defined as live $CD3^+$ $B220^-$ $TCR\gamma\delta^-$ $CD44^{high}$ $TCR\beta^+$ MR1-Tetramer$^+$ lymphocytes. Approximately 5×10^4 - 1×10^5 MAIT cells were sorted per mouse. 10^5 MAIT cells suspended in 100 µl was transferred via retroorbital injections into lightly anesthetized mice. Transferred mice were monitored to confirm recovery following injections, and rested for 2 weeks to allow for MAIT expansion before use in intranasal challenge model.

Intranasal challenge model

The following protocol was adapted from Nygren et al (*61*) to induce *V. cholerae* specific antibody

responses through a prime-boost live bacterial model. In brief, on Day 0 mice were anesthetized using

isoflurane and infected i.n. with 10^6-10^7 CFU *V. cholerae* 01 Biotype El Tor Serotype Inaba suspended in

50 µl sterile PBS (25 µl per nares). Mice were monitored for 7 days post infection for weight loss and

clinical signs of pneumoniae (outlined above). Mice with loss of >20% of body weight and/or severe

clinical symptoms were euthanized. Mice were subsequently re-challenged on Day 28 using the above

protocol. Blood samples were collected weekly, prior to infections on Day 0 and 28, from submandibular

bleeds on lightly isoflurane anesthetized mice. Blood serum was isolated using BD Microtainer SST tubes

(BD Biosciences) according to the manufacturer's protocol. On Day 42 (2 weeks post re-challenge), mice

were euthanized using a bell jar and isoflurane. Blood was collected immediately via cardiac puncture and

serum isolated. Lungs were perfused and single cell suspensions were processed as described above.

Spleens were excised, ground through 70 µm filters in cold R10 media. Red blood cells were lysed using

ACK Lysis Buffer (Thermo Fisher). 1-2x10^6 cells from lung and spleen cell suspensions were stained

with fluorochrome conjugated antibodies for flow cytometric analyses.

Statistical analysis

All statistical tests were performed using Prism version 8.4.2 (GraphPad Software, La Jolla, CA, USA).

Differences were compared using two-tailed Mann Whitney *U* tests, or Multiple t tests comparisons

accounting for False Discoveries using the Benjamini, Krieger and Yekutieli correction as indicated.

Graphs were created using Prism or R (Vienna, Austria) and the graphing package ggplot2 (H. Wickham,

Springer-Verlag, NY, USA).

References

1.   C. Gutzeit, G. Magri, A. Cerutti, Intestinal IgA production and its role in hot-microbe interaction.

Immunol. Rev. **260**, 76–85 (2014).

2.   O. Pabst, E. Slack, IgA and the intestinal microbiota: the importance of being specific. *Mucosal*

Immunol. **13**, 12–21 (2020).

3.   S. Hardtke, L. Ohl, R. Förster, Balanced expression of CXCR5 and CCR7 on follicular T helper

cells determines their transient positioning to lymph node follicles and is essential for efficient B-

cell help. *Blood.* **106**, 1924–1931 (2005).

4.   H. Qi, T follicular helper cells in space-time. *Nat. Rev. Immunol.* **16**, 612–625 (2016).

5.   J. Shi, S. Hou, Q. Fang, X. Liu, X. Liu, H. Qi, PD-1 Controls Follicular T Helper Cell Positioning

and Function. *Immunity.* **49**, 264-274.e4 (2018).

6.   A. Tafuri, A. Shahinian, F. Bladt, S. K. Yoshinaga, M. Jordana, A. Wakeham, L. M. Boucher, D.

Bouchard, V. S. F. Chan, G. Duncan, B. Odermatt, A. Ho, A. Itie, T. Horan, J. S. Whoriskey, T.

Pawson, J. M. Penninger, P. S. Ohashi, T. W. Mak, ICOS is essential for effective T-helper-cell

responses. *Nature.* **409**, 105–109 (2001).

7.   G. Van Kooten, J. Banchereau, CD40-CD40 ligand. *J. Leukoc. Biol.* **67**, 2–17 (2000).

8.   K. Ozaki, R. Spolski, C. G. Feng, C. F. Qi, J. Cheng, A. Sher, H. C. Morse, C. Liu, P. L.

Schwartzberg, W. J. Leonard, A critical role for IL-21 in regulating immunoglobulin production.

Science (80-.). **298**, 1630–1634 (2002).

9.   G. Galli, P. Pittoni, E. Tonti, C. Malzone, Y. Uematsu, M. Tortoli, D. Maione, G. Volpini, O.

Finco, S. Nuti, S. Tavarini, P. Dellabona, R. Rappuoli, G. Casorati, S. Abrignani, Invariant NKT

cells sustain specific B cell responses and memory. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 3984–9

(2007).

10. E. Tonti, G. Galli, C. Malzone, S. Abrignani, G. Casorati, P. Dellabona, NKT-cell help to B

lymphocytes can occur independently of cognate interaction. *Blood.* **113**, 370–376 (2009).

11.   S. T. Scanlon, S. Y. Thomas, C. M. Ferreira, L. Bai, T. Krausz, P. B. Savage, A. Bendelac,

672   Airborne lipid antigens mobilize resident intravascular NKT cells to induce allergic airway

673   inflammation. *J. Exp. Med.* **208**, 2113–2124 (2011).

674   12.   E. A. Leadbetter, M. Brigl, P. Illarionov, N. Cohen, M. C. Luteran, S. Pillai, G. S. Besra, M. B.

675   Brenner, NK T cells provide lipid antigen-specific cognate help for B cells. *Proc. Natl. Acad. Sci.*

676   **105**, 8339–8344 (2008).

677   13.   P. Barral, J. Eckl-Dorna, N. E. Harwood, C. De Santo, M. Salio, P. Illarionov, G. S. Besra, V.

678   Cerundolo, F. D. Batista, B cell receptor-mediated uptake of CD1d-restricted antigen augments

679   antibody responses by recruiting invariant NKT cell help in vivo. *Proc. Natl. Acad. Sci.* **105**,

680   8345–8350 (2008).

681   14.   I. L. King, A. Fortier, M. Tighe, J. Dibble, G. F. M. Watts, N. Veerapen, A. M. Haberman, G. S.

682   Besra, M. Mohrs, M. B. Brenner, E. A. Leadbetter, Invariant natural killer T cells direct B cell

683   responses to cognate lipid antigen in an IL-21-dependent manner. *Nat. Immunol.* **13**, 44–50

684   (2012).

685   15.   E. Tonti, M. Fedeli, A. Napolitano, M. Iannacone, U. H. von Andrian, L. G. Guidotti, S.

686   Abrignani, G. Casorati, P. Dellabona, Follicular Helper NKT Cells Induce Limited B Cell

687   Responses and Germinal Center Formation in the Absence of CD4 + T Cell Help . *J. Immunol.*

688   **188**, 3217–3222 (2012).

689   16.   H. B. Shah, S. K. Joshi, M. L. Lang, CD40L-null NKT cells provide B cell help for specific

690   antibody responses. *Vaccine*. **29**, 9132–9136 (2011).

691   17.   P. Rampuria, G. A. Lang, T. S. Devera, C. Gilmore, J. D. Ballard, M. L. Lang, Coordination

692   between T helper cells, iNKT cells, and their follicular helper subsets in the humoral immune

693   response against *Clostridium difficile* toxin B. *J. Leukoc. Biol.* **101**, 567–576 (2017).

694   18.   P. P. Chang, P. Barral, J. Fitch, A. Pratama, C. S. Ma, A. Kallies, J. J. Hogan, V. Cerundolo, S. G.

695   Tangye, R. Bittman, S. L. Nutt, R. Brink, D. I. Godfrey, F. D. Batista, C. G. Vinuesa,

Identification of Bcl-6-dependent follicular helper NKT cells that provide cognate help for B cell

responses. *Nat. Immunol.* **13**, 35–43 (2012).

19. N. Caccamo, L. Battistini, M. Bonneville, F. Poccia, J. J. Fournie, S. Meraviglia, G. Borsellino, R.

A. Kroczek, C. La Mendola, E. Scotet, F. Dieli, A. Salerno, CXCR5 Identifies a Subset of V 9V 2

T Cells which Secrete IL-4 and IL-10 and Help B Cells for Antibody Production. *J. Immunol.* **177**,

5290–5295 (2006).

20. R. R. Bansal, C. R. Mackay, B. Moser, M. Eberl, IL-21 enhances the potential of human γδ T cells

to provide B-cell help. *Eur. J. Immunol.* **42**, 110–119 (2012).

21. R. M. Rezende, A. J. Lanser, S. Rubino, C. Kuhn, N. Skillin, T. G. Moreira, S. Liu, G. Gabriely,

B. A. David, G. B. Menezes, H. L. Weiner, γδ T cells control humoral immune response by

inducing T follicular helper cell differentiation. *Nat. Commun.* **9** (2018), doi:10.1038/s41467-018-

05487-9.

22. D. T. Leung, T. R. Bhuiyan, N. S. Nishat, M. R. Hoq, A. Aktar, M. A. Rahman, T. Uddin, A. I.

Khan, F. Chowdhury, R. C. Charles, J. B. Harris, S. B. Calderwood, F. Qadri, E. T. Ryan,

Circulating mucosal associated invariant T cells are activated in Vibrio cholerae O1 infection and

associated with lipopolysaccharide antibody responses. *PLoS Negl. Trop. Dis.* **8**, e3076 (2014).

23. M. S. Bennett, S. Trivedi, A. S. Iyer, J. S. Hale, D. T. Leung, Human mucosal-associated invariant

T (MAIT) cells possess capacity for B-cell help. *J. Leukoc. Biol.* **102**, 1261–1269 (2017).

24. J. Jiang, Z. Cao, PD-1-expressing MAIT cells from patients with tuberculosis exhibit elevated

production of CXCL13, 1–15 (2020).

25. G. Murayama, A. Chiba, H. Suzuki, A. Nomura, T. Mizuno, T. Kuga, S. Nakamura, H. Amano, S.

Hirose, K. Yamaji, Y. Suzuki, N. Tamura, S. Miyake, A Critical Role for Mucosal-Associated

Invariant T Cells as Regulators and Therapeutic Targets in Systemic Lupus Erythematosus. *Front.*

Immunol. **10**, 2681 (2019).

26. M. A. Rahman, E. Ko, F. Bhuyan, G. Enyindah-asonye, R. Hunegnaw, S. H. Hait, C. J. Hogge, D.

J. Venzon, T. Hoang, M. Robert-guroff, Mucosal-associated invariant T (MAIT) cells provide B-cell help in vaccinated and subsequently SIV- infected Rhesus Macaques. *Sci. Rep.*, 1–17 (2020).

27. F. Tilloy, E. Treiner, S. H. Park, C. Garcia, F. Lemonnier, H. de la Salle, A. Bendelac, M. Bonneville, O. Lantz, An invariant T cell receptor alpha chain defines a novel TAP-independent major histocompatibility complex class Ib-restricted alpha/beta T cell subpopulation in mammals. *J. Exp. Med.* **189**, 1907–21 (1999).

28. R. Reantragoon, A. J. Corbett, I. G. Sakala, N. A. Gherardin, J. B. Furness, Z. Chen, S. B. G. Eckle, A. P. Uldrich, R. W. Birkinshaw, O. Patel, L. Kostenko, B. Meehan, K. Kedzierska, L. Liu, D. P. Fairlie, T. H. Hansen, D. I. Godfrey, J. Rossjohn, J. McCluskey, L. Kjer-Nielsen, Antigen-loaded MR1 tetramers define T cell receptor heterogeneity in mucosal-associated invariant T cells. *J. Exp. Med.* **210**, 2305–20 (2013).

29. M. Lepore, A. Kalinicenko, A. Colone, B. Paleja, A. Singhal, A. Tschumi, B. Lee, M. Poidinger, F. Zolezzi, L. Quagliata, P. Sander, E. Newell, A. Bertoletti, L. Terracciano, G. De Libero, L. Mori, Parallel T-cell cloning and deep sequencing of human MAIT cells reveal stable oligoclonal TCRβ 2 repertoire. *Nat. Commun.* **5** (2014), doi:10.1038/ncomms4866.

30. E. Martin, E. Treiner, L. Duban, L. Guerri, H. Laude, C. Toly, V. Premel, A. Devys, I. C. Moura, F. Tilloy, S. Cherif, G. Vera, S. Latour, C. Soudais, O. Lantz, Stepwise Development of MAIT Cells in Mouse and Human. *PLoS Biol.* **7**, e1000054 (2009).

31. L. Le Bourhis, E. Martin, I. Péguillet, A. Guihot, N. Froux, M. Coré, E. Lévy, M. Dusseaux, V. Meyssonnier, V. Premel, C. Ngo, B. Riteau, L. Duban, D. Robert, M. Rottman, C. Soudais, O. Lantz, Antimicrobial activity of mucosal-associated invariant T cells. *Nat. Immunol.* **11**, 701–708 (2010).

32. M. Dusseaux, E. Martin, N. Serriari, I. Péguillet, V. Premel, D. Louis, M. Milder, L. Le Bourhis, C. Soudais, E. Treiner, O. Lantz, Human MAIT cells are xenobiotic-resistant, tissue-targeted, CD161 hi IL-17-secreting T cells. *Blood*. **117**, 1250–1259 (2011).

33.   A. Rahimpour, H. F. Koay, A. Enders, R. Clanchy, S. B. G. Eckle, B. Meehan, Z. Chen, B.

Whittle, L. Liu, D. P. Fairlie, C. C. Goodnow, J. McCluskey, J. Rossjohn, A. P. Uldrich, D. G.

Pellicci, D. I. Godfrey, Identification of phenotypically and functionally heterogeneous mouse

mucosal-associated invariant T cells using MR1 tetramers. *J. Exp. Med.* **212**, 1095–1108 (2015).

34.   A. Kurioka, J. E. Ussher, C. Cosgrove, C. Clough, J. R. Fergusson, K. Smith, Y. H. Kang, L. J.

Walker, T. H. Hansen, C. B. Willberg, P. Klenerman, MAIT cells are licensed through granzyme

exchange to kill bacterially sensitized targets. *Mucosal Immunol.* **8**, 429–440 (2015).

35.   L. Le Bourhis, M. Dusseaux, A. Bohineust, S. Bessoles, E. Martin, V. Premel, M. Coré, D. Sleurs,

N. E. Serriari, E. Treiner, C. Hivroz, P. Sansonetti, M. L. Gougeon, C. Soudais, O. Lantz, MAIT

Cells Detect and Efficiently Lyse Bacterially-Infected Epithelial Cells. *PLoS Pathog.* **9** (2013),

doi:10.1371/journal.ppat.1003681.

36.   J. Dias, E. Leeansyah, J. K. Sandberg, Multiple layers of heterogeneity and subset diversity in

human MAIT cell responses to distinct microorganisms and to innate cytokines. *Proc. Natl. Acad.*

Sci. **114**, E5434–E5443 (2017).

37.   A. Meierovics, W.-J. C. Yankelevich, S. C. Cowley, MAIT cells are critical for optimal mucosal

immune responses during in vivo pulmonary bacterial infection. *Proc. Natl. Acad. Sci.* **110**,

E3119–E3128 (2013).

38.   P. Georgel, M. Radosavljevic, C. Macquin, S. Bahram, The non-conventional MHC class I MR1

molecule controls infection by Klebsiella pneumoniae in mice. *Mol. Immunol.* **48**, 769–775

(2011).

39.   Z. Chen, H. Wang, C. D'Souza, S. Sun, L. Kostenko, S. B. G. Eckle, B. S. Meehan, D. C. Jackson,

R. A. Strugnell, H. Cao, N. Wang, D. P. Fairlie, L. Liu, D. I. Godfrey, J. Rossjohn, J. McCluskey,

A. J. Corbett, Mucosal-associated invariant T-cell activation and accumulation after in vivo

infection depends on microbial riboflavin synthesis and co-stimulatory signals. *Mucosal Immunol.*

10, 58–68 (2017).

40.   H. Wang, C. D'Souza, X. Y. Lim, L. Kostenko, T. J. Pediongco, S. B. G. Eckle, B. S. Meehan, M.

Shi, N. Wang, S. Li, L. Liu, J. Y. W. Mak, D. P. Fairlie, Y. Iwakura, J. M. Gunnersen, A. W.

Stent, D. I. Godfrey, J. Rossjohn, G. P. Westall, L. Kjer-Nielsen, R. A. Strugnell, J. McCluskey,

A. J. Corbett, T. S. C. Hinks, Z. Chen, MAIT cells protect against pulmonary Legionella

longbeachae infection. *Nat. Commun.* **9**, 3350 (2018).

41.   F. J. Radcliff, S. Waldvogel-thurlow, F. Clow, M. Mahadevan, J. Johnston, G. Li, T. Proft, R. G.

Douglas, J. D. Fraser, Impact of Superantigen-Producing Bacteria on T Cells from Tonsillar

Hyperplasia. *Pathogens.* **8**, 1–17 (2019).

42.   E. Leeansyah, L. Loh, D. F. Nixon, J. K. Sandberg, Acquisition of innate-like microbial reactivity

in mucosal tissues during human fetal MAIT-cell development. *Nat. Commun.* **5**, 1–10 (2014).

43.   P. A. Szabo, M. Miron, D. L. Farber, Location, location, location: Tissue resident memory T cells

in mice and humans. *Sci. Immunol.* **4**, 1–12 (2019).

44.   M. L. Freeman, S. R. Morris, M. M. Lederman, CD161 Expression on Mucosa-Associated

Invariant T Cells is Reduced in HIV-Infected Subjects Undergoing Antiretroviral Therapy Who

Do Not Recover CD4+ T Cells. *Pathog. Immun.* **2**, 335 (2017).

45.   A. Han, J. Glanville, L. Hansmann, M. M. Davis, Linking T-cell receptor sequence to functional

phenotype at the single-cell level. *Nat. Biotechnol.* **32**, 684–692 (2014).

46.   V. Voillet, M. Buggert, C. K. Slichter, J. D. Berkson, F. Mair, M. M. Addison, Y. Dori, G.

Nadolski, M. G. Itkin, R. Gottardo, M. R. Betts, M. Prlic, Human MAIT cells exit peripheral

tissues and recirculate via lymph in steady state conditions. *JCI insight.* **3**, 1–10 (2018).

47.   L. Loh, N. A. Gherardin, S. Sant, J. C. Crawford, N. L. Bird, C. E. Van De Sandt, M. L. Moreira,

E. K. Allen, J. Crowe, T. Loudovaris, K. L. Flanagan, K. M. Quinn, J. Rossjohn, G. Paul, S. B. G.

Eckle, J. Mccluskey, I. Dale, K. Kedzierska, J. C. Crawford, N. L. Bird, H. Koay, C. E. Van De

Sandt, M. L. Moreira, M. Lappas, E. K. Allen, J. Crowe, T. Loudovaris, K. L. Flanagan, K. M.

Quinn, J. Rossjohn, P. G. Thomas, Human Mucosal-Associated Invariant T Cells in Older

Individuals Display Expanded TCR α β Clonotypes with Potent Antimicrobial Responses (2020),

doi:10.4049/jimmunol.1900774.

48.   M. C. Gold, J. E. McLaren, J. A. Reistetter, S. Smyk-Pearson, K. Ladell, G. M. Swarbrick, Y. Y.

L. Yu, T. H. Hansen, O. Lund, M. Nielsen, B. Gerritsen, C. Kesmir, J. J. Miles, D. A. Lewinsohn,

D. A. Price, D. M. Lewinsohn, MR1-restricted MAIT cells display ligand discrimination and

pathogen selectivity through distinct T cell receptor usage. *J. Exp. Med.* **211**, 1601–1610 (2014).

49.   K. Franciszkiewicz, M. Salou, F. Legoux, Q. Zhou, Y. Cui, S. Bessoles, O. Lantz, MHC class I-

related molecule, MR1, and mucosal-associated invariant T cells. *Immunol. Rev.* **272**, 120–138

(2016).

50.   M. I. Vazquez, J. Catalan-Dibene, A. Zlotnik, B cells responses and cytokine production are

regulated by their immune microenvironment. *Cytokine*. **74**, 318–326 (2015).

51.   A. G. Zaretsky, J. J. Taylor, I. L. King, F. A. Marshall, M. Mohrs, E. J. Pearce, T follicular helper

cells differentiate from Th2 cells in response to helminth antigens. *J. Exp. Med.* **206**, 991–999

(2009).

52.   D. Fang, K. Cui, K. Mao, G. Hu, R. Li, M. Zheng, N. Riteau, S. L. Reiner, A. Sher, K. Zhao, J.

Zhu, Transient T-bet expression functionally specifies a distinct T follicular helper subset. *J. Exp.*

Med. **215**, 2705–2714 (2018).

53.   P. Wang, Y. Wang, L. Xie, M. Xiao, J. Wu, L. Xu, Q. Bai, Y. Hao, Q. Huang, X. Chen, R. He, B.

Li, S. Yang, Y. Chen, Y. Wu, L. Ye, The transcription factor T-bet is required for optimal type I

follicular helper T cell maintenance during acute viral infection. *Front. Immunol.* **10**, 1–17 (2019).

54.   K. Li, C. K. Vorkas, A. Chaudhry, D. L. Bell, R. A. Willis, A. Rudensky, J. D. Altman, M. S.

Glickman, J. Aubé, Synthesis, stabilization, and characterization of the MR1 ligand precursor 5-

amino-6-D-ribitylaminouracil (5-A-RU). *PLoS One*. **13**, 20–22 (2018).

55.   J. Shen, X. Luo, Q. Wu, J. Huang, G. Xiao, L. Wang, B. Yang, H. Li, C. Wu, A subset of

CXCR5+CD8+ T cells in the germinal centers from human tonsils and lymph nodes help B cells

821   produce immunoglobulins. *Front. Immunol.* **9**, 1–12 (2018).

822 56. T. Leng, H. D. Akther, C. P. Hackstein, K. Powell, T. King, M. Friedrich, Z. Christoforidou, S.

823   McCuaig, M. Neyazi, C. V. Arancibia-Cárcamo, J. Hagel, F. Powrie, R. S. Peres, V. Millar, D.

824   Ebner, R. Lamichhane, J. Ussher, T. S. C. Hinks, E. Marchi, C. Willberg, P. Klenerman, TCR and

825   Inflammatory Signals Tune Human MAIT Cells to Exert Specific Tissue Repair and Effector

826   Functions. *Cell Rep.* **28**, 3077-3091.e5 (2019).

827 57. M. Lettau, A. Wiedemann, E. V. Schrezenmeier, C. Giesecke-Thiel, T. Dörner, Human CD27+

828   memory B cells colonize a superficial follicular zone in the palatine tonsils with similarities to the

829   spleen. A multicolor immunofluorescence study of lymphoid tissue. *PLoS One*. **15**, 1–17 (2020).

830 58. E. Donnadieu, K. B. Reisinger, S. Scharf, Y. Michel, J. Bein, S. Hansen, A. G. Loth, N. Flinner, S.

831   Hartmann, M.-L. Hansmann, Landscape of T Follicular Helper Cell Dynamics in Human

832   Germinal Centers. *J. Immunol.*, ji1901475 (2020).

833 59. J. J. Hong, P. K. Amancha, K. Rogers, A. A. Ansari, F. Villinger, Spatial Alterations between

834   CD4 + T Follicular Helper, B, and CD8 + T Cells during Simian Immunodeficiency Virus

835   Infection: T/B Cell Homeostasis, Activation, and Potential Mechanism for Viral Escape . *J.*

836   *Immunol.* **188**, 3247–3256 (2012).

837 60. J. A. Kotov, M. K. Jenkins, Cutting Edge: T Cell–Dependent Plasmablasts Form in the Absence of

838   Single Differentiated CD4 [+] T Cell Subsets. *J. Immunol.*, ji1801349 (2018).

839 61. E. Nygren, J. Holmgren, S. R. Attridge, Murine antibody responses following systemic or mucosal

840   immunization with viable or inactivated Vibrio cholerae. *Vaccine*. **26**, 6784–6790 (2008).

841 62. I. S. Cisternas, A. Torres, A. F. Flores, V. A. G. Angulo, Differential regulation of riboflavin

842   supply genes in Vibrio cholerae. *Gut Pathog.* **9**, 1–7 (2017).

843 63. D. J. Haney, M. D. Lock, J. K. Simon, J. Harris, M. Gurwith, Antibody-based correlates of

844   protection against cholera: Analysis of a challenge study of a cholera-naive population. *Clin.*

845   *Vaccine Immunol.* **24**, 1–11 (2017).

846   64.   J. B. Harris, R. C. LaRocque, F. Chowdhury, A. I. Khan, T. Logvinenko, A. S. G. Faruque, E. T.

847   Ryan, F. Qadri, S. B. Calderwood, Susceptibility to Vibrio cholerae infection in a cohort of

848   household contacts of patients with cholera in Bangladesh. *PLoS Negl. Trop. Dis.* **2** (2008),

849   doi:10.1371/journal.pntd.0000221.

850   65.   R. Finkelstein, Vibriocidal Antibody Inhibition (VAI) Analysis: A Technique for the Identification

851   of the Predominant Vibriocidal Antibodies in Serum and for the Detection and Identification of

852   Vibrio Cholerae Antigens. *J. Immunol.* **89**, 264–271 (1962).

853   66.   R. I. Glass, A. M. Svennerholm, M. R. Khan, S. Huda, M. Imdadul Huq, J. Holmgren,

854   Seroepidemiological studies of EI tor cholera in bangladesh: Association of serum antibody levels

855   with protection. *J. Infect. Dis.* **151**, 236–242 (1985).

856   67.   A. S. Azman, J. Lessler, F. J. Luquero, T. R. Bhuiyan, A. I. Khan, F. Chowdhury, A. Kabir, M.

857   Gurwith, A. A. Weil, J. B. Harris, S. B. Calderwood, E. T. Ryan, F. Qadri, D. T. Leung,

858   Estimating cholera incidence with cross-sectional serology. *Sci. Transl. Med.* **11**, 1–11 (2019).

859   68.   D. T. Leung, T. Uddin, P. Xu, A. Aktar, R. A. Johnson, M. A. Rahman, M. M. Alam, M. K.

860   Bufano, G. Eckhoff, Y. Wu-Freeman, Y. Yu, T. Sultana, F. Khanam, A. Saha, F. Chowdhury, A. I.

861   Khan, R. C. Charles, R. C. LaRocque, J. B. Harris, S. B. Calderwood, P. Kováč, F. Qadri, E. T.

862   Ryan, Immune responses to the O-specific polysaccharide antigen in children who received a

863   killed oral cholera vaccine compared to responses following natural cholera infection in

864   Bangladesh. *Clin. Vaccine Immunol.* **20**, 780–788 (2013).

865   69.   S. M. Patel, M. A. Rahman, M. Mohasin, M. A. Riyadh, D. T. Leung, M. M. Alam, F.

866   Chowdhury, A. I. Khan, A. A. Weil, A. Aktar, M. Nazim, R. C. LaRocque, E. T. Ryan, S. B.

867   Calderwood, F. Qadri, J. B. Harris, Memory B Cell Responses to Vibrio cholerae O1

868   Lipopolysaccharide Are Associated with Protection against Infection from Household Contacts of

869   Patients with Cholera in Bangladesh. *Clin. Vaccine Immunol.* **19**, 842–848 (2012).

70.   A. S. Iyer, A. S. Azman, M. Bouhenia, L. O. Deng, C. P. Anderson, M. Graves, P. Kováč, P. Xu,

E. T. Ryan, J. B. Harris, D. A. Sack, F. J. Luquero, D. T. Leung, Dried Blood Spots for Measuring

Vibrio cholerae-specific Immune Responses. *PLoS Negl. Trop. Dis.* **12** (2018),

doi:10.1371/journal.pntd.0006196.

71. R. C. Kauffman, T. R. Bhuiyan, R. Nakajima, L. M. Mayo-Smith, R. Rashu, M. R. Hoq, F.

Chowdhury, A. I. Khan, A. Rahman, S. K. Bhaumik, L. Harris, J. T. O'Neal, J. F. Trost, N. H.

Alam, A. Jasinskas, E. Dotsey, M. Kelly, R. C. Charles, P. Xu, P. Kováč, S. B. Calderwood, E. T.

Ryan, P. L. Felgner, F. Qadri, J. Wrammert, J. B. Harris, Single-cell analysis of the plasmablast

response to Vibrio cholerae demonstrates expansion of cross-reactive memory B cells. *MBio.* **7**, 1–

11 (2016).

72. B. Sit, T. Zhang, B. Fakoya, A. Akter, R. Biswas, E. T. Ryan, M. K. Waldor, Oral immunization

with a probiotic cholera vaccine induces broad protective immunity against Vibrio cholerae

colonization and disease in mice. *PLoS Negl. Trop. Dis.*, 1–20 (2019).

73. N. Rout, Enhanced Th1/Th17 Functions of CD161+ CD8+ T Cells in Mucosal Tissues of Rhesus

Macaques. *PLoS One.* **11**, e0157407 (2016).

74. S. Crotty, T Follicular Helper Cell Differentiation, Function, and Roles in Disease. *Immunity.* **41**,

529–542 (2014).

75. M. Salio, O. Gasser, C. Gonzalez-Lopez, A. Martens, N. Veerapen, U. Gileadi, J. G. Verter, G.

Napolitani, R. Anderson, G. Painter, G. S. Besra, I. F. Hermans, V. Cerundolo, Activation of

Human Mucosal-Associated Invariant T Cells Induces CD40L-Dependent Maturation of

Monocyte-Derived and Primary Dendritic Cells. *J. Immunol.*, ji1700615 (2017).

76. R. J. Johnston, A. C. Poholek, D. DiToro, I. Yusuf, D. Eto, B. Barnett, A. L. Dent, J. Craft, S.

Crotty, Bcl6 and Blimp-1 are reciprocal and antagonistic regulators of T follicular helper cell

differentiation. *Science (80-.).* **325**, 1006–1010 (2009).

77. R. Ettinger, S. Kuchen, P. E. Lipsky, The role of IL-21 in regulating B-cell function in health and

disease. *Immunol. Rev.* **223**, 60–86 (2008).

78.   D. Eto, C. Lao, D. DiToro, B. Barnett, T. C. Escobar, R. Kageyama, I. Yusuf, S. Crotty, IL-21 and

IL-6 are critical for different aspects of B cell immunity and redundantly induce optimal follicular

helper CD4 T cell (Tfh) differentiation. *PLoS One*. **6**, e17739 (2011).

79.   L. Moens, S. G. Tangye, Cytokine-mediated regulation of plasma cell generation: IL-21 takes

center stage. *Front. Immunol.* **5**, 1–13 (2014).

80.   A. Cerutti, M. Cols, I. Puga, Activation of B cells by non-canonical helper signals. *EMBO Rep.* **13**,

798–810 (2012).

81.   X. Shen, T. Lagergard, Y. Yang, M. Lindblad, M. Fredriksson, J. Holmgren, Systemic and

mucosal immune responses in mice after mucosal immunization with group B streptococcus type

III capsular polysaccharide-cholera toxin B subunit conjugate vaccine. *Infect. Immun.* **68**, 5749–

5755 (2000).

82.   H. Yu, A. Yang, S. Derrick, J. Y. W. Mak, L. Liu, D. P. Fairlie, S. Cowley, Artificially induced

MAIT cells inhibit M . bovis BCG but not M . tuberculosis during in vivo pulmonary infection.

Sci. Rep., 1–12 (2020).

83.   S. Sakai, K. D. Kauffman, S. Oh, C. E. Nelson, C. E. B. Iii, D. L. Barber, MAIT cell-directed

therapy of Mycobacterium tuberculosis infection. *Mucosal Immunol.*, 1–10 (2020).

84.   A. Cerutti, The regulation of IgA class switching. *Nat. Rev. Immunol.* **8**, 421–434 (2008).

85.   J. Holmgren, A. M. Svennerholm, Vaccines against mucosal infections. *Curr. Opin. Immunol.* **24**,

343–353 (2012).

86.   G. Ben Youssef, M. Tourret, M. Salou, L. Ghazarian, V. Houdouin, S. Mondot, Y. Mburu, M.

Lambert, S. Azarnoush, J.-S. Diana, A.-L. Virlouvet, M. Peuchmaur, T. Schmitz, J.-H. Dalle, O.

Lantz, V. Biran, S. Caillat-Zucman, Ontogeny of human mucosal-associated invariant T cells and

related T cell subsets. *J. Exp. Med.* **215**, 459–479 (2018).

87.   F. Legoux, M. Salou, O. Lantz, MAIT Cell Development and Functions: the Microbial

Connection. *Immunity*. **53**, 710–723 (2020).

88.   F. Qadri, G. Mohi, J. Hossain, T. Azim, A. M. Khan, M. A. Salam, R. B. Sack, M. J. Albert, A. M.

Svennerholm, Comparison of the vibriocidal antibody response in cholera due to Vibrio cholerae

O139 Bengal with the response in cholera due to Vibrio cholerae O1. *Clin. Diagn. Lab. Immunol.*

2, 685–8 (1995).

89.   P. Mombaerts, A. R. Clarke, M. A. Rudnicki, J. Iacomini, S. Itohara, J. J. Lafaille, L. Wang, Y.

Ichikawa, R. Jaenisch, M. L. Hooper, S. Tonegawa, Mutations in T-cell antigen receptor genes α

and β block thymocyte development at different stages. *Nature.* **360**, 225–231 (1992).

929
930

Acknowledgements

Funding: This research was supported by the National Institutes of Health (R01 AI130378 to D.T.L. and

T32 AI138945 to O.J.). This work was also supported by the University of Utah Flow Cytometry Facility

in addition to the National Cancer Institute through Award Number P30CA042014. We would also like to

thank the staff of the Office of Comparative Medicine, and University of Utah Cell Imaging Core, and

Kelin Li and Jeffrey Aubé (UNC Chapel Hill) for providing 5-amino-6-D-ribitylaminouracil (5-A-RU)

for our study.

Author Contributions: O.J. and D.T.L. designed and directed the project. O.J and S.T. performed the

experiments. K.F. and J.S.H. contributed to experimental methodology. J.D.M. provided samples and

resources. D.T.L acquired funding. O.J. and D.T.L wrote the paper, and all authors discussed results,

provided critical feedback, and contributed to the final manuscript.

Declaration of interests: None.

Supplementary Materials

Supplementary Fig. 1. Representative flow cytometry gating strategy for Fig. 1. Representative flow cytometry gating strategy for MAIT and non-MAIT T cells in (**A**) PBMC and (**B**) tonsils. PE conjugated 6-FP MR1-Tetramer gating controls for PBMC and tonsils are located below 5-OP-RU MR1-Tetramer staining gates. (**C**) Representative CXCR5 staining for PBMC and tonsil CD4[+] T and MAIT cells quantified in Fig. 1C. Gating was based on CXCR5 fluorescence minus one (FMO) control.

946
947

Supplementary Fig. 2. Flow gating strategy for T cell co-stimulatory molecule expression analysis. Representative flow cytometry gating strategy for PBMC and tonsil Tfh, MAIT and MAITfh cell (**A**) PD1, (**B**) ICOS, (**C**) CD69, and (**D**) CD161 expression. FMO's were not included for CD69 and CD161 due to clarity of antibody staining. (**E**) Frequency of Tfh, MAIT, and MAITfh cell populations expressing CD69. (**F**) Frequency of CD161$^+$ MAIT cells broken down by CXCR5 expression. Data are represented as median from 1 independent experiment. n = 5-7. *p < 0.05, **p<0.01, ***p<0.001, ****p<0.0001 by two-tailed Mann-Whitney U test.

949
950
951

952
953

Supplementary Fig. 3. Similar TRAJ expression in CD161+ and CD161- tonsil MAIT cells. Heat map of TRAJ usage percent frequencies based on Illumina single cell TCR-Phenotype sequencing in tonsil MAIT cells broken down by CD161 negative (top) and positive (bottom) populations. Each column represents an individual donor (n=4). MAIT cells were defined as MR1-5-OP-RU-tetramer positive, and TRAV1-2 antibody positive. No statistical significance was found between CD161- and CD161+ groups using Multiple t tests comparison accounting for False Discoveries using the Benjamini, Krieger and Yekutieli method.

Supplementary Fig. 4. Tfh co-stimulatory marker expression in paired pediatric tonsil and PBMC samples. (**A**) Representative FACS plots of unstimulated paired pediatric CD3[+] PBMC (top) and tonsil (bottom) donor cells with MAIT cell (gated in left panel) co-expression of CXCR5 with ICOS (middle) and PD1 (right). (**B**) Quantification of MAIT cell frequency as percentage of CD3[+]. (**C**) Frequency of CXCR5[+] MAIT cells. Frequency of (**D**) ICOS[+] and (**E**) PD1[+] MAIT, MAITfh and Tfh cells. Data are represented as Median from 2 pooled independent experiments. $n \geq 13$. *$p < 0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by two-tailed Mann-Whitney U test

Supplementary Fig. 5. qPCR and flow gating strategy (Fig 2) and additional FAC sorted qPCR targets. Representative flow cytometry gating strategy for qPCR FAC sorted and flow populations in Fig.2 from (**A**) PBMC and (**B**) tonsils. (**C**) Heat map of additional qPCR targets described in Fig 2. As described, data are represented as ΔΔCt relative to β-Actin and then the PBMC-CD8$^+$CXCR5$^-$ population. Rows denote gene ID and columns denote sample group (P=PBMC, T=tonsil). Heat map data are represented by mean of n=6-8 samples per group.

955
956
957

Supplementary Fig. 6. IFNγ and IL-21 expression in unstimulated PMBC and tonsil T cell populations. (**A**) Representative flow cytometry from 6 hour no stimulation controls with BFA added for final 4 hours in PBMC and tonsil T cell populations showing IL-21 and IFNγ co-expression. (**B**) Frequency of T cell populations expressing IL-21, IFNγ or both. Data are represented as median from two pooled independent experiments. n≥5. *$p < 0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by two-tailed Mann-Whitney *U* test. Legend in black box denotes experimental groups.

Supplementary Fig. 7. IFNγ and IL-21 expression in 5-A-RU stimulated PMBC and tonsil T cell populations. (**A**) Representative flow cytometry of 6 hr 5-A-RU (0.5 μM/ml) and Methylglyoxal (50 μg/ml) stimulated cultures with BFA in PBMC and tonsil T cell populations showing IL-21 and IFNγ co-expression. (**B**) Frequency of T cell populations expressing IL-21, IFNγ or both. Data are represented as median from one independent experiment. n=5. *$p < 0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by two-tailed Mann-Whitney *U* test. Legend in black box denotes experimental groups.

Supplementary Fig. 8. GC Tfh cells are marked by high PD1 expression in tonsils. (A) Representative immunofluorescence images from tonsil cryosections stained with anti-IgD (blue), anti-PD1 (white), anti-CD4 (green), and anti-CD3 (red) and imaged on a Leica SP8 confocal microscope with 20X objective. Enlarged composite image on bottom with a germinal center (GC) highlighted by PD1^{High} GC Tfh cells outlined with white dashed line, and the follicular mantle (FM) made up of IgD^{+} naïve B cells outlined with a yellow dashed line. All scale bars represent 100 μm.

Supplementary Fig. 9. Flow gating for murine lung MAIT and T lymphocyte counts. (**A**) Representative flow gating of murine MAIT cells from digested lung tissue highlighted in Fig. 4a. MAIT cells were defined as live B220$^-$ CD3$^+$ TCRγδ$^-$ CD44^{High} TCRβ$^+$ MR1-Tetramer$^+$. Only WT-*V.c.*, MAIT-*V.c.*, and Sham-*V.c.* groups are portrayed. (**B**) Total MAIT cell count per lung. (**C**) Total MAIT cell count per spleen. (D) Total non-MAIT CD3$^+$ TCRβ$^+$ T cells per lung. Data are represented as Median with IQR from 4 independent experiments. n= 4-15 mice per group. *$p < 0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ by two-tailed Mann-Whitney U test. Legend in black box denotes experimental groups.

Supplementary Fig. 10. CXCR5 expression on *S. enterica* Typhimurium expanded MAIT cells prior to adoptive transfer. Representative flow cytometry histograms (**A**) and mean fluorescence intensity (**B**) of CXCR5 expression in T cell subsets of Day 7 WT C57Bl/6 mice given intranasal PBS (Left) or 10^6 CFU *S.* Typhimurium (Left). Data are represented as Median from 1 experiment. n= 2-4 mice per group.

Supplementary Fig. 11. Flow gating for murine lung B lymphocyte counts. (A) Representative
flow cytometry gating strategy of murine B cell subsets from digested lung tissue. **(B)** Total B220$^+$ B
cells per lung. **(C)** Total B220$^+$ IgD$^+$ Naïve B cells per lung. **(D)** Total B220$^+$ IgD$^-$
CD38$^+$CD27$^+$CD138$^-$ Memory B cells per lung. **(E)** Total B220^{low} IgD$^-$ CD38$^+$CD27$^+$CD138$^+$ PB/PCs
per lung. Data are represented as Median with IQR from 4 independent experiments. n= 4-15 mice per
group. *p < 0.05, **p<0.01, ***p<0.001, ****p<0.0001 by two-tailed Mann-Whitney U test. Legend
in black box denotes experimental groups.

Supplementary Fig. 12. Serum IgM, total IgM, IgG and IgA and BAL fluid ELISAs. (A-F)
Serum *V. cholerae* lysate, OSP, and CT IgM ELISAs representing weekly antibody kinetics (A, C &
E) and Day 42 endpoint responses (B, D & F). Data are represented as ELISA units normalized to a
pooled serum positive control from WT B6 mice challenged with V. cholerae. Total IgM (G), IgA (H)
and IgG (I) ELISAs from Day 42 endpoint serum. Total Ig was quantitated using a standard curve
based on known Ig quantities. Data are represented as Median with IQR from 5 independent
experiments. n= 6-22 mice per group. *p < 0.05, **p<0.01, ***p<0.001, ****p<0.0001 by two-tailed
Mann-Whitney U test. *V. cholerae* lysate IgA (J), IgG (K) and IgM (L) ELISAs from D42 BAL fluid.
Legend in black box denotes experimental groups.