1   Corresponding Author:

2
Dr. Megan L. Fritz
Department of Entomology
University of Maryland
4291 Field House Dr.
Plant Sciences Bldg. Rm. 4112
College Park, MD 20742
mfritz13@umd.edu

Genome evolution in an agricultural pest following adoption of transgenic crops

Megan L. Fritz[1*], Kelly A. Hamby[1], Katherine Taylor[1], Alexandra M. DeYonke[2], and Fred Gould[2]

[1]Department of Entomology, University of Maryland, College Park, MD 20742 USA

[2]Department of Entomology and Plant Pathology, North Carolina State University, Raleigh, NC 27607

USA

* Corresponding author

Running Title: Resistance Evolution in Agricultural Ecosystems

Keywords

Helicoverpa zea, Bt resistance, temporal genomic change, polygenic selection

Abstract

Replacement of synthetic insecticides with transgenic crops for pest management has been both

economically and environmentally beneficial. These benefits have often eroded as pests evolved

resistance to the transgenic crops, but a broad understanding of the timing and complexity of the

adaptive changes which lead to field-evolved resistance in pest species is lacking. Wild populations of

Helicoverpa zea, a major lepidopteran crop pest and the target of transgenic Cry toxin-expressing

cotton and corn, have recently evolved widespread, damaging levels of resistance. Here, we quantified

patterns of genomic change in wild *H. zea* collected between 2002 and 2017 when adoption rates of

Cry-expressing crops expanded in North America. Using a combination of genomic and genotypic

approaches, we identified significant temporal changes in allele frequency throughout the genomes of

field-collected *H. zea*. Many of these changes occurred concurrently with increasingly damaging levels

of resistance to Cry toxins between 2012 and 2016, in a pattern consistent with polygenic selection.

Surprisingly, none of the eleven previously described Cry resistance genes showed signatures of

selection in wild *H. zea*. Furthermore, we observed evidence of a very strong selective sweep in one

region of the *H. zea* genome, yet this strongest change was not additively associated with Cry

resistance. This first, whole genome analysis of field-collected specimens to study evolution of Cry

resistance demonstrates the potential and need for a more holistic approach to examining pest

adaptation to changing agricultural practices.

Introduction

The study of rapid evolutionary responses to natural selection has gained momentum in the past decade, and along with it recognition of the relevance of pest adaptation to control measures (Reznick et al. 2019). Melander published the first observation of insecticide resistance evolution in 1914 (Melander 1914), and since then, pests and weeds have adapted to myriad chemical, biological, and cultural control measures (Gould 1991, Palumbi 2001, Délye et al. 2013, Gould et al. 2018). While there has been an emphasis on finding single genetic loci that confer resistance to pesticides, recognition that surviving insecticide or herbicide applications could involve a suite of genomic changes has increased (Baucom 2019). Yet a clear understanding of the timing and complexity of changes associated with adaptation to agricultural ecosystems is lacking (Leon et al. 2020).

The bacterium, *Bacillus thuringiensis* (Bt), produces one or more crystalline (Cry) proteins that are each toxic to a narrow range of insect pests when ingested. Genes encoding Cry toxins were engineered into cultivated crops to reduce pest damage, producing so-called Bt crops, a major innovation in agricultural pest management that has been adopted around the globe (NASEM 2016). The economic and environmental benefits of this technology are significant: they can replace heavily used synthetic insecticides, which have damaging off target effects on pollinators and natural enemies, and have contributed to area wide pest suppression in non-Bt crops (Hutchison et al. 2010, NASEM 2016, Dively et al. 2018). Yet a number of pest species have evolved resistance to one or more Cry toxins and there is concern about the durability of these transgenic crops (Tabashnik and Carrière 2017). Resistance management strategies have been implemented effectively for a few major pest species with high susceptibility to the toxins (NASEM 2016). Unfortunately, these measures have been inadequate for species where Cry toxins cause only moderate levels of mortality, providing opportunities for the evolution of resistance (Tabashnik and Carrière 2017).

Discovery of resistance-conferring genes has been of long-standing interest, in part due to expectations that this knowledge could lead to molecular diagnostic approaches that facilitate detection

of early resistance evolution in wild pest populations (US-EPA 1998, US-EPA 2018). Molecular

signals of resistance could then serve as triggers for regulatory actions that would minimize resistance

risk and preserve environmentally sustainable technologies, like Cry toxins (US-EPA 1998). Genes

discovered to have major impacts on Cry tolerance are being used to a limited extent for molecular

resistance monitoring (Morin et al. 2004, Gahan et al. 2007, Zhang et al. 2013, Jin et al. 2018). To

date, there are at least 11 such genes in the Lepidoptera (Wu 2014, Lin 2019), many of which were

discovered in laboratory-selected populations and from screening wild populations for single genes

with major effects on resistance (Morin et al. 2003, Gahan et al. 2001, Zhang et al. 2012, Jin et al.

2018). The question of whether molecular monitoring of these genes would effectively track resistance

evolution in wild populations remains unresolved.

Helicoverpa zea is a migratory pest species (Gould et al. 2002) which comprises a single

panmictic population in North America (Seymour et al. 2016). It has been controlled with Bt crops

since 1996, when corn and cotton expressing Cry1A became commercially available. In 2007, corn

and cotton cultivars producing both Cry1A and Cry2A were introduced. Since their

commercialization, North American adoption rates of Cry-expressing corn and cotton grew from 8 and

15% of corn and cotton acreage in 1997 to *ca.* 80% by 2012. This rapid expansion of Cry-expressing

crops placed significant selection pressure on their target pests, and widespread, damaging levels of

resistance to Cry-expressing crops by *H. zea* were documented by 2016 (Dively et al. 2016, Reisig et

al. 2018, Kaur et al. 2019, Yang et al. 2019). Existing Bt corn and cotton cultivars do not produce a

high dose of Cry protein relative to what is required to kill heterozygote *H. zea* larvae with resistance

alleles (Luttrell and Jackson 2012). This likely contributed to their recent resistant status (Tabashnik

and Carrière 2017). Under these conditions, resistance to Cry proteins was predicted to be polygenic

(Gould 1998, Tabashnik et al. 2004) and to arise from standing genetic variation (Roush and McKenzie

1987).

Between 2002 and 2017, we saved *ca.* 1000 adult moths in ethanol each year, and we used

samples from these collections to track genomic changes in *H. zea* over time. Instead of simply

monitoring for changes in previously identified candidate genes, we took a gene agnostic approach and

scanned entire genomes for signatures of selection. Using two sequencing approaches, double-digest

Restriction-site Associated DNA sequencing (ddRAD-seq) and whole genome resequencing, we

identified multiple genomic regions with significant changes in allele frequency during a 15 year time

period. Shifts in allele frequency were used to calculate the strength of selection imposed on wild *H.*

zea in agricultural ecosystems. We also examined whether any of the 11 previously identified

candidate Cry resistance genes showed evidence of strong shifts in allele frequency over time.

Surprisingly, none of these genes could be implicated in Cry-toxin resistance evolution for *H. zea*.

Instead, we found a number of novel candidate genes within regions of greater than expected allele

frequency change. Due to changes in synthetic insecticide use following the introduction of Bt crops,

we also examined whether the frequency of alleles of *H. zea* genes associated with detoxification of

synthetic insecticides or even plant defensive compounds had changed, as was observed in another

species where Bt crops replaced insecticidal control (Fritz et al. 2018).

This first whole genome analysis of field-collected specimens to study impacts of Bt crops

demonstrates the potential and need for a more holistic approach to examining pest adaptation to

changing agricultural practices. While monitoring previously identified candidate genes may prove

useful in some cases, our results indicate that such an approach is likely to miss critical evolutionary

events. Analysis of patterns of genomic changes in pest populations following the introduction of a

new control tactic or tool could prove useful, but requires follow-up experimental confirmation to

demonstrate clear, causal relationships among genes and adaptive phenotypes.

Results

Strong temporal ddRAD-seq allele frequency shifts

H. zea males were collected from five pheromone baited traps (Hartstack 1979) in Bossier Parish, LA, and pooled by collection date in the months of May through October, 2002 through 2017. Randomly sampled moths from a subset of years were selected for genetic analysis; 2002, 2007, 2012, as well as in 2016, when strong phenotypic resistance to Cry1Ab and increasing resistance to Cry2Ab was being documented (Dively et al. 2016, Kaur et al. 2019). Two hundred and sixty-five specimens were prepared into ddRAD-seq libraries and sequenced according to Fritz *et al.* (2016, 2018). Following filtering and alignment to the 341 Mb *H. zea* genome, we identified 14,398 single nucleotide polymorphisms (SNPs) from 259 individuals with an average read count of 404,003 (Figure S1). Nucleotide diversity (π), and heterozygosity (F) did not change between years (Table S2), and overall genomic divergence was low according to Weir and Cockerham's F_{ST} ($F_{ST} < 0.004$; Table S3). This low divergence was also supported by a sequential k-means analysis, which showed that a single genetic cluster or population (k = 1) best described our ddRAD-seq genotyped *H. zea* (Figure S2). Interestingly, genetic divergence that occurred between the 2012-2016 collection periods was at least 3 times higher ($F_{ST} = 0.0025$) than for any other sequentially sampled periods. Pairwise F_{ST} values were 0.0009 and 0.0002 for the 2002-2007 and 2007-2012 collection periods, respectively (Table S3).

We used a discriminant analysis of principal components (DAPC; R package adegenet v. 2.1.1, Jombart et al. 2010, Jombart and Ahmed 2011) to further examine temporal changes in the genomic structure of wild *H. zea*. Initially, we identified ddRAD SNP combinations that clustered individuals into groups (k = 2-4) by maximizing between group variation and minimizing within group variation. Using these SNP combinations, we then estimated posterior membership probabilities for each individual from our dataset from each collection year. When k was equal to 2, samples within each of the collection years 2002, 2007, and 2012 were assigned to both genetic clusters. In contrast, samples collected in 2016 were strongly biased toward a single genetic cluster (Figure 1), and this trend

persisted regardless of the k-value used (Figure S3). Interestingly, the genotypic shift between 2012

and 2016 corresponded to the emergence of widespread field-evolved phenotypic resistance to Cry

proteins in *H. zea* (Dively et al. 2016, Kaur et al. 2019). We sequentially removed the top 1%, 2.5%

and 5% of ddRAD SNPs that had the greatest influence on genetic clustering, and re-evaluated

population membership probability (assuming k=2). From this we determined that up to 5% (n = 720)

of ddRAD SNPs were responsible for this observed genotypic shift (Figure S4).

To identify which of the 14,398 ddRAD SNPs experienced greater than expected shifts in allele

frequency over time, we conducted a genomic outlier analysis (R package OutFLANK v. 0.2, Whitlock

and Lotterhos 2015). When we calculated genetic divergence (F_{ST}) for each SNP across all 4

collection years, 53 had higher than expected divergence (Figure 2, File S1). Twenty-two of those 53

were among the top variance contributing SNPs that strongly influenced genetic clustering in our

DAPC analysis. The 53 ddRAD SNP outliers were spread across 47 (of 2,975) unique scaffolds, and

their among-year F_{ST} values ranged from 0.068 to 0.278.

No significant allelic changes within candidate Bt resistance genes

We found ddRAD-seq markers within or near (<56kb) 10 of the 11 previously identified

candidate genes for Bt resistance, but none corresponded to outliers found over the time course of the

collections (Table 1). To verify this result we conducted Illumina whole genome sequencing (WGS) of

35 *H. zea* males collected from the same location in LA in 2002 (n = 13), 2012 (n = 11), and 2017 (n =

11) (Table S4). After genome alignment and filtering, this approach yielded 4,999,128 SNPs, and

2,622 fell within the 11 candidate Cry resistance genes. We used a 10kb sliding window-averaged

Weir and Cockerham's F_{ST} analysis and observed that F_{ST} values calculated for each gene ranged from

-0.018 to 0.056 (Table 1; Figure S5). The empirical threshold for statistically significant genetic

divergence, generated according to Rubin et al. (2010), was 6 standard deviations ($ZF_{ST} > 6$) from the

mean pairwise F_{ST} value in a window. None of the eleven candidate genes contained windows that

reached our threshold for statistical significance of $F_{ST} = 0.079$ ($ZF_{ST} > 6$) per 10kb window. Taken

together, these results indicated that the Cry resistance in *H. zea* was not caused by changes in the

coding regions of any of these eleven candidate genes or their flanking cis-regulatory sequences.

Selection led to significant temporal divergence in novel genomic regions

Using our full WGS SNP dataset, we conducted pairwise analyses of genomic divergence

between populations collected from 2002, 2012, and 2017 using 5, 10, 20 and 40 kb sliding window-

averaged Weir and Cockerham's F_{ST} along the length of the genome (Figure S6). We report on the 40

kb sliding window analysis because it captured the major genomic changes that occurred for all sliding

window sizes, but reduced the numbers of windows that appeared highly divergent due to few SNPs in

the window. Thirty-six genomic windows on 11 different scaffolds had F_{ST} values that exceeded our

empirical threshold for statistical significance of $F_{ST} = 0.047$ per 40 kb when 2002 and 2017

populations were compared (Figure 3). For paired comparisons of populations collected in 2002 and

2012, and 2012 and 2017 there were 55 and 30 genomic windows on 16 and 10 scaffolds with

statistically significant genomic divergence, respectively (Figure 3; File S2).

Both genetic drift and selection influence the landscape of genomic divergence between

populations. Therefore, we quantified the frequency with which our empirical threshold of $F_{ST} =$

0.047, calculated from the pairwise comparison of samples from 2002 and 2017, could be reached in a

40kb genomic window due to drift alone (Hudson 2002). Genotypic simulations were conducted for a

single population sampled at two timepoints, but in the absence of selection. Allele frequency changes

within 40kb windows were examined for multiple population sizes (N = 21,000 - 140,000;

Supplemental Methods). The proportion of genotypic simulations (of 20,000) that produced $F_{ST} >$

0.047 was the probability that our empirical threshold of $F_{ST} = 0.047$ could be exceeded due to genetic

drift. Results from our simulations demonstrated this probability was very low (p = 0.0189 - 0) and

negatively correlated with present day population size (Figure S7; Table S5).

Because the population sizes used in the simulations are likely underestimates of the effective population size of this migratory pest (Jones et al. 2019), it is highly improbable that our empirical F_{ST} threshold could have been exceeded by the action of drift. As a demonstration of its large effective population size, we collected fifth and sixth instar *H. zea* from ears of Obsession I (n = 8) and Cry-expressing Obsession II (n = 8) sweet corn grown in Central Maryland Research and Education Center (CMREC) demonstration plots. The genomes of these individuals, collected in September of 2017, were sequenced with Illumina short reads, and their genotypes compared to samples collected in Bossier Parish, LA, in that same year. After genome alignment and filtering for quality and linkage disequilibrium, we examined 14,697 SNPs in the dataset, which yielded a genome-wide F_{ST} value of 0.0004. The bootstrapped 95% confidence intervals were (-0.0002, 0.0008), and their overlap with zero indicated that allele frequencies in the LA and MD populations did not significantly differ from one another.

Identification of novel genes under selection

We examined the genes near regions of greatest temporal genomic divergence to determine whether any were associated with detoxification of plant-produced or human-applied toxicants. Our list of candidates included those from families of genes associated with Cry resistance, but also resistance to synthetic insecticides and xenobiotics. Eleven genes, including 3 peptidases, 4 choline/carboxylesterases, 2 cytochrome p450s, 1 ABC transporter, and 1 tetraspanin, were near or within regions of significant temporal genetic divergence. No proteases, proteinases, cadherins, voltage-gated channels, alkaline phosphatases, or glutathione-S-transferases were found within 50 kb of regions with elevated divergence. Evidence from a recent study suggests that increasing adoption of Bt crops led to a reduction in the frequency of synthetic insecticide resistance-conferring alleles, likely due to a reduction in synthetic insecticide applications (Fritz et al. 2018). Therefore, we specifically examined two genes known to be involved in pyrethroid resistance. Neither the voltage-gated sodium

channel gene (*vgsc*), nor a cytochrome p450 (*cyp337b3*) which underlie pyrethroid resistance in the

close *H. zea* relatives, *Chloridea virescens* (previously *Heliothis virescens*; Taylor et al. 1993) and

Helicoverpa armigera (Joußen et al. 2012), respectively, were near regions of elevated genomic

divergence. All genes found within 50kb of genomic windows with higher than expected divergence

can be found in File S3.

We reasoned that increasing adoption of Bt crops over time would result in strong, sustained

selection over time, leading to significant allele frequency changes across multiple collection years.

Therefore, we identified scaffolds with elevated temporal genomic divergence in more than one

pairwise comparison. Five scaffolds showed signs of significant allele frequency change in at least two

pairwise comparisons. These included scaffold 173, which contained a *venom dipeptidyl peptidase 4 -*

like gene (HzOG204148), and scaffold 569, which contained two *carboxypeptidase Q (cpq)* -like genes

(HzOG204151 and HzOG204153). Scaffold 1964 contained an RNA-directed DNA polymerase

(HzOG206500), and scaffold 65 had one genomic window of elevated divergence in multiple pairwise

comparisons which was near 7 genes, including *tetraspanin (tspan) 68C* (HzOG211510) and a

krueppel-like gene (HzOG211515). Finally, scaffold 1612 contained 2 genomic windows with

elevated divergence across all three by-year comparisons, which overlapped with *cyp333b3*

(HzOG200024). Of these five scaffolds, 569 and 1612, had levels of genetic divergence that greatly

exceeded all others between 2002 and 2017 ($F_{ST} > 0.089$), and such divergence was never observed for

genetic simulations in the absence of selection (Table S5).

The genome assembly of *H. zea* is fragmented into 2,975 genomic scaffolds, which makes

detecting the extent of a selective sweep challenging. Under conditions of strong selection, the

footprint in the genome should be broad, and we reasoned there was potential for some of these

temporally divergent scaffolds to be physically linked. We used tblastn to align the genes from the 5

scaffolds showing strong temporal divergence in multiple by-year comparisons to the *Bombyx mori*

genome (ASM15162v1), and identified their relative positions assuming conserved macrosynteny

among Lepidoptera (d'Alençon et al. 2010). Four of these five scaffolds were adjacent to one another

and syntenic to *B. mori* chromosome 13, indicating that the breadth of the selective sweep extended for

ca. 3 Mb (Fig S8). Linkage disequilibrium within the putative sweep region decayed more slowly than

both genome-wide average rates of decay, as well as decay along single, similarly-sized scaffolds (>

1Mb) without evidence of genomic divergence (Fig S9). This region of the *B. mori* genome contained

134 unique genes, none of which were previously known to be involved in Cry resistance (File S4).

Interestingly, we re-examined our ddRAD-seq dataset and detected two ddRAD-seq markers found

within this 3 Mb genomic region (on scaffold 569), which were outliers according to OutFLANK. This

further confirmed the significance of changes occurring in this genomic region, but with a much larger

sample size (n > 45 per year).

The 3 Mb putative sweep region experienced the greatest temporal change in frequency

according to our whole genome scan. Yet our DAPC analysis identified 720 ddRAD SNPs on 505

unique scaffolds which caused a major architectural shift in the genomes of *H. zea* collected between

2012 and 2016 (Figure 1, Figure S3, Figure S4). Of those 720 ddRAD SNPs, only 5 aligned to this 3

Mb region. This pattern, consistent with polygenic selection on standing variation, indicated that allele

frequency changes at many loci are involved in *H. zea* adaptation over the study period. Our further

analyses in this study focused on the 3 Mb putative sweep region due to the strong changes in allele

frequency and the breadth of the putative selective sweep.

Temporal shifts in the frequency of non-synonymous mutations

Given the significant levels of genomic divergence near *cyp333b3* and the two

*carboxypeptidase Q-like (*hereafter *cpq*) genes detected by ddRAD-seq and WGS datasets, as well as

the putative functional roles of these genes in insect metabolism and detoxification, we further

examined protein-coding changes at these genes over time. Based upon our temporal analysis,

genomic divergence was strongest at the 3' end of all three genes (Figure S10). While we detected 50

and 40 SNPs within *cpq1* and *cpq2*, respectively, only 3 were associated with putative changes in

amino acid sequence (Table 3).   One transition from cytosine to thymine caused a putative alanine to

valine substitution in *cpq1*, and over time, the cytosine declined significantly from 0.73 in 2002, to 0.32

and 0.14 in 2012 and 2017, respectively (Freeman-Halton Fisher's exact p = 0.0001; Table 2). Two

SNPs in *cpq2* showed modest changes in allele frequency over time, although not statistically

significant when a Bonferroni-corrected α value was used to account for multiple comparisons (Table

2). One cytosine to thymine transition encoded an alanine to valine amino acid substitution at 152,595

bp, and a second occurred at 152,183 bp, encoding a threonine to isoleucine substitution. Finally, there

were 28 SNPs present in the protein-coding sequence of *cyp333b3*, 10 of which produced putative

amino acid changes. Five of these non-synonymous mutations underwent statistically significant

changes over time (p < 0.003; Table 2).

We also compared the frequencies of these non-synonymous mutations in the LA and MD

populations collected in 2017, and for all 8 SNPs, there was no statistically significant difference in

allele frequency based on collection location (Table S7). These results and the low overall F_{ST} value

for the two-population comparison (reported above) support previous findings that *H. zea* is highly

migratory with little population structure.

Strength of selection experienced by wild H. zea

Using the temporal changes in allele frequency, we calculated the strength of selection required

to produce these statistically significant protein-coding changes over time. The selection coefficient(*s*)

gave the strength of selection against *q*, the declining allele, examined under 3 different dominance

scenarios of *p:* complete dominance, no dominance, and recessiveness. For 5 of the 8 non-synonymous

mutations (Table 2), *q* was the allele present in the reference genome, and in the remaining cases, *q* was

the alternate allele. We calculated mean values of *s* from allele frequency changes which occurred

during the 2002-2012 collection period, as well as the 2012-2017 collection period. Values of s varied

according to selection period and dominance scenario (Figure 4), but most fell within the range of 0.03-

0.17. When p was assumed to be recessive, stronger selection was required to produce the observed

changes in allele frequency in the 2002-2012 collection period due to its low frequency (*ca.* 0.25 - 0.3).

Yet when p was dominant, stronger selection was required to produce the observed changes in allele

frequency between 2012-2017, due to its relatively high starting frequency (*ca.* 0.65-0.75). In the case

of the SNP at 36,959 bp on scaffold 1612, it's frequency (p) was already 0.95 in 2012. Assuming

dominance of p, the strength of selection that would have been required to increase its frequency to 1

by 2017 was very high ($s = 0.70$), which increased the variance around the mean value of s (Figure 4).

Frequency of non-synonymous substitutions prior to 2002

The intermediate frequency of the non-synonymous polymorphisms in 2002 compelled us to

examine the frequencies of those polymorphisms in *H. zea* collected from our LA trapping sites in

1998, shortly after the release of Cry-expressing crops in 1996. Moths from 1998 had DNA quality

that was too low for use in ddRAD-seq and WGS analyses, but short fragments could be PCR

amplified. We therefore amplified and Sanger sequenced *ca.* 200bp DNA fragments spanning one

non-synonymous polymorphism in each of *cpq1* and *cyp333b3* from 20-26 *H. zea* collected in 1998.

The respective frequencies of the reference alleles for *cpq1* and *cyp333b3* were 0.93 and 0.90 in 1998

compared with 0.73 and 0.65 in 2002, and 0.14 and 0.18 in 2017 (Table 2) showing further evidence of

genomic divergence at these sites. The change in allele frequency between 1998 and 2002 occurred

prior to introduction of varieties with Cry2 genes.

Strongest selective sweep is not linked to Cry resistance

To test whether this 3 Mb region with significant allele frequency divergence was linked to Cry

resistance, we bioassayed the progeny of two F_2 genetic crosses between recently field-collected Cry

resistant grandparents and Cry susceptible grandparents (Benzon Research Inc., Carlisle, PA). Cry

resistant larval growth was 7 and 10 times greater than that of Benzon susceptibles when fed a

diagnostic dose of diet containing Cry1 and Cry1 + Cry2 sweetcorn leaf tissue, made according to

Dively et al. (2016). Following 7 days of exposure to Cry1 or Cry1 + Cry2 treated diet, F_2 larval

growth phenotypes from each family were measured and individuals genotyped by PCR at non-

synonymous mutations in the *cyp333b3* and *cpq1* genes. A Kruskal Wallis test found no significant

difference in larval growth according to genotype on either diet treatment, indicating these non-

synonymous mutations were not additively associated with resistance to Cry toxins (Figure 5; Table

S9).   For the *H. zea* system, the question of what environmental factor could have selected for such

rapid change in allele frequency is fascinating but unsolved.

Discussion

In this study, we compared the genomes of *H. zea* collected between 2002 and 2017 to quantify

changes that occurred during a period of rapid adoption of a major agricultural innovation, Cry toxin-

expressing crops, in the United States. We found substantial changes in the frequencies of SNPs

throughout the *H. zea* genome that occurred in concert with the evolution of resistance to Cry toxins in

field populations of *H. zea*. A major finding of our work is that none of the previously identified

candidate Cry resistance genes in Lepidoptera showed signatures of selection in the field-collected *H.*

zea, nor were genes in the strongest selective sweep region associated with resistance in an additive

way. Instead, the timing and complexity of the genomic changes that occurred in *H. zea* were

consistent with the predicted polygenic nature of Cry resistance in this species, and provide insight into

how, when and whether genomic approaches could be used for resistance risk monitoring.

Over the past four decades there has been an emphasis on the search for single genes that confer

resistance to control measures in insect pests and weeds, and we now have a catalogue of such genes

(Ffrench-Constant 2013, Feyereisen et al. 2015, Délye et al. 2013, Heckel 2020). There are cases

where a single gene diagnostic (Donnelly et al. 2016) or a genomic diagnostic targeted at multiple

previously established candidate genes (Weetman et al. 2018) has been useful in monitoring for

resistance, but recently there has been concern that these focused approaches could lead to

complacency, as resistance evolves based on unexpected genes (Heckel 2020) or based on multiple

genes (Délye et al. 2013, Van Leeuwen et al. 2020, Leon et al. 2020) that are already at low

frequencies in pest or weed populations. Our work demonstrates that these concerns are appropriate. If

any or all of the previously identified candidate genes had been used in a diagnostic for Cry resistance

in *H. zea*, the diagnostic would have missed the coming problem.

While functional roles of specific genes that we identified have not yet been demonstrated,

multiple *H. zea* genomic regions have changed dramatically since the introduction of Cry toxin-

producing cotton and corn. The most compelling changes were found within a broad selective sweep

on the *H. zea* homolog of *B. mori* chromosome 13. WGS and ddRAD-seq datasets both showed

stronger than expected temporal genomic changes in a roughly 3 Mb region over the 15 year sampling

period. When the results of the genome analyses are combined with PCR analysis of SNPs within

candidate genes from 1998 samples from the same location, allele frequency changes of up to 0.79

occurred in these genes with metabolic functions or that are associated with xenobiotic resistance

(Ward 1976, Guzov et al. 1998, Bown and Gatehouse 2004). Analysis of genetic crosses of resistant

and susceptible strains found that this genomic region was not linked to Cry resistance in an additive

way (Figure 5). This was surprising due to rapid allele frequency changes, and the fact that two of the

candidates (*cpq1* and *cpq2*) were genes from a family capable of binding Cry toxins in *H. armigera*

(Da Silva et al. 2018). While this does not eliminate the possibility that genes in this region have

epistatic interactions that confer resistance, it highlights that monitoring solely for strong selective

sweeps could potentially mislead when it comes to resistance gene identification and resistance risk monitoring.

We emphasize, however, that we were able to detect that multiple regions of the genome changed during the course of our sampling in a pattern consistent with polygenic adaptation from standing genetic variation. Our findings are consistent with a recent study of crosses of Cry2 resistant *H. zea* to a susceptible strain that found phenotypic variance fitting a polygenic model (Yang et al. 2020). In our study, strong polygenic change was most apparent between the years 2012 and 2016 (Figure 1) when widespread and damaging levels of resistance to Cry toxins were documented (Dively et al. 2016). Ongoing, experimental, quantitative genomic studies will shed light on which of these genomic regions contribute most substantially to Cry tolerance in field-collected larvae.

The use of genomic approaches to study rapid evolution in natural populations has gained momentum (*e.g.* Crossley et al. 2017, Fritz et al. 2018, Nosil et al. 2018, Weetman et al. 2018, Bi et al. 2019, Dayan et al. 2019, Laurentino et al. 2020). Yet their use for studying pest adaptation to agricultural systems has been limited to a few cases (reviewed in Pélissié et al. 2018). As we attempt to apply these approaches to a variety of agricultural systems, system- and method-specific issues will need to be addressed to ensure accuracy. Results from our experiments indicate that genomic approaches may hold promise for monitoring resistance risk in pest populations exposed to novel management practices, with several caveats for how we interpret our results.

One concern is that major genomic changes identified through genomic monitoring might not be due to the control tool being studied. Deployment of pest management innovations do not take place in the absence of other existing selection pressures in the environment. When transgenic cotton and corn were introduced into North American agricultural systems, the use of synthetic insecticides were reduced, though certainly not eliminated. In a previous study using ddRAD-seq, we found that

another pest of cotton in Louisiana, *Chloridea virescens*, did not experience changes in the frequency

of Bt resistance candidate genes. Rather, changes to the frequency of an allele that conferred resistance

to pyrethroid insecticides were observed over a 15 year period (Fritz et al. 2018). Indeed, our data

suggest that the greatest allele frequency changes in the *H. zea* genome were not directly associated

with Cry resistance. Therefore, confirming the strength of association between any major genomic

change with a resistance phenotype is critical.

This confirmation testing can be conducted using the types of simple crosses and PCR-based

assays that we carried out in our study. Such assays can falsify the importance of rapidly changing loci

to resistance phenotypes and alleviate concerns regarding resistance risk. Conversely, knowledge of

allele frequency changes and confirmation of their impact on field resistance could then be used to

make predictions for future changes in resistance in field populations of pests. In cases similar to *H.*

zea on Cry-expressing crops where the control tactic does not kill all pest individuals or has sublethal

effects, collections from paired sentinel plots could also point to genomic regions associated with

resistance. Such paired sentinel plots are already being used for determining the current extent of

resistance, but typically monitor damage to the crop rather than the genetics of the pest (Dively et al.

2016). Focusing on damage and the economic impact of resistance does little to predict risk of

resistance and then mitigate it. Once widespread damage has occurred, resistance alleles have likely

already spread widely in the landscape. Instead, sampling and sequencing early instar larvae found in

treated and untreated sentinel plots would allow researchers to respectively quantify genomic change

associated with the pest management tool compared to the population at large.

A second concern is that population demographic processes, including migration patterns and

population expansion can result in the appearance of genome-wide shifts in allele frequency, which are

not due to selection. When this occurs, the risk for genomic monitoring would be to falsely implicate a

region of the genome in evolving resistance. We were fortunate to be studying a highly migratory

species (Hartstack et al. 1982, Gould et al. 2002, Seymour et al. 2016) with little population structure

over large geographic areas (Perera and Blanco 2011, Seymour et al. 2016). Our finding of negligible

genomic differences between samples collected in 2017 from Louisiana and Maryland are consistent

with results from these previous studies. Additionally, *H. zea* feeds on a number of host plants, and this

seems to have resulted in the population densities remaining similar to those prior to the use of

transgenic cotton and corn (Micinski et al. 2008). It is not surprising that with this species, measured

overall genomic diversity remained constant. For a species that interbreeds over such a large area,

such population demographic processes are unlikely to be an issue. Had we been studying an insect or

weed with strong local population structure, or one in which the control measure caused severe

reduction and resurgence of the pest, our experimental approaches would have required additional

careful consideration.

Finally, many so-called "mega-pest" lineages are thought to have genomes that already contain

variants, which prime them for the evolution of resistance to novel pest control measures (Crossley et

al. 2017, Pélissié et al. 2018). When selection is applied to these populations, they experience soft

selective sweeps at genomic regions that are typically spread throughout the genome. Such signatures

of polygenic selection in field-collected samples can be difficult to bioinformatically tease apart from

the impacts of drift. Fortunately, new computational approaches which can better differentiate soft

selective sweeps from background selection or other population demographic processes are becoming

available (Schrider and Kern 2016), and some even detect the proportion of genome-wide changes in

populations due to selection using temporal datasets (Buffalo and Coop 2019). Further development of

bioinformatic approaches specifically aimed at detecting temporal genomic changes at both single loci

and those spread throughout the genome will provide critical tools for resistance risk monitoring in

agricultural systems.

The benefits of genomic monitoring of pest species go beyond identifying the mechanisms of

resistance and applications for resistance risk mitigation. Adoption of genomic approaches for

resistance monitoring will be crucial for addressing the decades-long debate on exactly how pest

species rapidly adapt to management practices (Crow 1957). Even today, the extent to which a species

response to selection is mono- or polygenic, from standing genetic variation or novel mutation remains

unclear. Here we present evidence on the nature of genomic change during expansion of a major

agricultural innovation, Cry-expressing crops, in one important pest species. Extension of genomic

monitoring programs, accompanied by targeted experimental confirmation, to additional pest species

on broad spatial and temporal scales could provide critical insight into the nature of adaptation by these

species to human-imposed selection.

Data Availability

Data for this study are available at: to be completed after manuscript is accepted for publication.

445
All scripts and barcodes files used to perform these analyses can be found at

https://github.com/mcadamme/FieldHz_Pop_Genomics.

Competing Interests

The authors declare they have no competing interests.

Acknowledgements

This work was funded by USDA NIFA Biotechnology Risk Assessment Grants 2016-33522-25640 and

2019-33522-29992.

Methods

Insect collections for ddRAD-seq and WGS analyses

H. zea adults were collected by pheromone-baited trap bi-weekly from May through September in Bossier Parish, Louisiana (Table S1). Collections were made in the years 2002, 2007, 2012, 2016, and 2017, and specimens were stored in 100% ethanol at -20° C until use. Late instar (5-6th) *H. zea* were also collected in August and September of 2017 at the University of Maryland CMREC farms in Prince George's county, MD, from the ears of Bt-expressing and non-Bt sweet corn isolines according to Dively *et al.* (2016) and reared to pupation under laboratory conditions of 16:8 LD at 25° C and 50% RH. Newly emerged adults were sacrificed by freezing and held at -80° C until use. Genomic DNA was isolated from half of an adult thorax with a Qiagen DNeasy kit (Qiagen, Inc., Valencia, CA, USA) using a modified mouse tail protocol according to Fritz et al. (2020).

DdRAD-seq library preparation

Specimens from the years 2002, 2007, 2012, and 2016 were prepared into double-digest RAD-seq (ddRAD-seq) libraries for population genomic analysis according to Fritz *et al*. (2016 2018; further details in SI). Barcoded and indexed libraries were combined in equimolar amounts (4 nM) for sequencing on 8 runs of an Illumina MiSeq. Our final data set contained 259 individuals after quality control and filtering to remove individuals with low read counts (Figure S1).

DdRAD-seq Bioinformatic Analysis

After adapter trimming (cutadapt v. 1.18; Martin, 2011), paired end reads were merged with FLASh (v. 1.2.11; Magoc and Salzberg, 2011), demultiplexed and filtered using the process_radtags script from Stacks (v. 2.2; Catchen et al. 2011), and aligned to the *H. zea* genome assembly (v. 1.0, NCBI Bioproject PRJNA378438; Pearce et al., 2017) with Bowtie2 (v. 2.2.6-2; Langmead and Salzberg, 2012). We called genotypes with BCFtools mpileup and call functions (Danecek et al., 2016)

using standard options. Our final SNP dataset was filtered with vcftools (Danecek et al., 2011) to

include SNPs that i) had a depth of coverage > 3, ii) were present in at least 75% of the 259 individuals

in our filtered dataset, iii) had a minor allele frequency of 0.05, iv) had a maximum of 2 alleles, and v)

were thinned to 1 per 200 bp to reduce the degree of linkage disequilibrium between SNPs on the same

ddRAD-seq marker.

We calculated average nucleotide diversity (π) and heterozygosity (the inbreeding coefficient

F) values and their corresponding 95% bootstrapped confidence intervals (N = 5000) for each year

using vcftools. Genome wide allele frequency divergence between years was examined by pairwise

F_{ST} (Weir and Cockerham 1984) using the StAMPP package (v. 1.5.1, Pembleton *et al.,* 2013) in R (v.

3.4.4, R Development Core Team, 2008). A k-means analysis, implemented with adegenet (v. 2.1.1,

Jombart and Ahmed, 2011), modeled the number of putative populations or clusters (k = 1:10) into

which our *H. zea* fit according to genotype. Using the three best-fitting numbers of clusters (k = 2:4)

as priors, we applied a Discriminant Analysis of Principal Components (DAPC; adegenet) to identify

groups of genetically similar *H. zea* individuals.   The R package OutFLANK (v. 0.2, Whitlock and

Lotterhos 2015) calculated Weir and Cockerham's F_{ST} for each SNP to quantify genetic divergence

across years and identify those with higher than expected divergence over time.

A number of genes have already been implicated in Bt resistance in other Lepidopteran pest

species (Table 1). Blastn and tblastn searches (https://blast.ncbi.nlm.nih.gov) determined the locations

of these putative Bt resistance genes in the *H. zea* genome, and we used custom scripts to identify the

nearest ddRAD-seq marker to each (Table 1).

WGS library preparation

To confirm our ddRAD-seq results, and extend our analysis to regions not covered by ddRAD-

seq markers, we conducted whole genome sequencing of additional *H. zea* collected from this same

location in LA. Genomic DNA from additional adults collected in 2002 (n= 13), 2012 (n = 11), and

507 2017 (n = 11) was prepared according to an Illumina TruSeq LT library preparation (Illumina, Inc. San

508 Diego, CA), and indexed libraries were sequenced on two Illumina NextSeq500 runs using 150 bp

509 paired-end reads.

511 *WGS bioinformatic analysis*

512  Following filter-trimming (Trimmomatic v. 0.38; Bolger et al., 2014), and reference genome

513 alignment with Bowtie2, we filtered poor quality and duplicate (PCR and optical) reads from our

514 dataset using GATK (v. 4.0.12.0; McKenna et al. 2010) and SAMtools (v. 0.1.19; Li et al., 2009).

515 Genotypes were called with BCFtools mpileup and call functions, and indels were removed with

516 vcftools. Further filtering with vcftools removed SNPs i) with a coverage depth < 3, ii) absent from 8

517 or more individuals sequenced, iii) with minor allele frequency of < 0.05, and iv) with > 2 alleles per

518 locus. Genomic change at the 11 Bt resistance candidate genes was measured by pairwise F_{ST}

519 averaged across SNPs found within each gene for samples collected in 2002 and 2017. According to

520 Rubin et al. (2010), we standardized F_{ST} estimates with a z-transformation, and excluded windows with

521 fewer than 10 SNPs. Windows with an average F_{ST} value greater than 6 standard deviations from the

522 genome-wide averaged F_{ST} value ($ZF_{ST} > 6$) were considered to have undergone statistically significant

523 genetic divergence. For each scaffold containing a Bt resistance candidate, 10kb averaged F_{ST} values

524 with a 1kb step size were examined for the length of the scaffold and compared against our empirical

525 threshold determined for each genome-wide by-year comparison.

526  Sliding window averaged F_{ST} values were used to identify signatures of strong selection across

527 the entire genome. For each pair of years we calculated averaged F_{ST} across 40kb windows with a

528 10kb step size. Additional sliding window and step sizes were also examined for the comparison of

529 samples collected in 2002 and 2017, and are reported in Figure S2. Genotype simulations under

530 neutral expectations using realistic population demographic conditions generated a distribution of F_{ST}

531 values, which were compared against empirical F_{ST} values from our genome scan results as a

secondary approach to examining statistical significance (Hudson 2002; Supplemental Methods).

Results of our whole genome scan were compared to results from our ddRAD-enabled outlier analysis

to determine which, if any, regions were detected in both datasets.

We examined the genes within 50kb of each genomic window showing signs of statistically

significant genetic divergence between each pair of years, and queried the gene list to identify those

from families generally known to be associated with detoxification of plant-produced or human-applied

toxicants. These included peptidases/proteases (Gill et al. 1995, Banks et al. 2001, Liu et al. 2015),

carboxyl/cholinesterases (Newcomb et al. 1997, Russell et al. 2011), proteinases (Ahmed et al. 1998),

cytochrome p450s (reviewed in Russell et al. 2011, Howe and Herde 2015), cadherins (Gahan et al.

2001, Morin et al. 2003, Fabrick et al. 2014; Zhang et al. 2017), ABC transporters (Gahan et al. 2010),

alkaline phosphatases (Jurat-Fuentes and Adang 2004), glutathione-S-transferases (reviewed in Russell

et al. 2011, Howe and Herde 2015), voltage-gated sodium channels (reviewed in Dong et al. 2014), and

tetraspanins (Jin et al. 2018).

The genome assembly of *H. zea* is fragmented into 2,975 genomic scaffolds, which makes

detecting the extent of a selective sweep challenging. A tblastn alignment of genes from the 5 most

divergent scaffolds to the *Bombyx mori* genome (NCBI ASM15162v1) identified their relative

positions to one another using conserved macrosynteny among Lepidoptera (d'Alençon et al. 2010).

Genome similarity between H. zea *collected from MD and LA in 2017*

A global F_{ST} analysis was implemented using the R package hierfstat (v. 0.04-22; Goudet

2005), and compared *H. zea* collected at CMREC in Prince George's county, MD, in 2017 to those

collected in Bossier Parish, LA, in 2017. Bootstrapped 95% confidence intervals around the global F_{ST}

value were generated using the R package bootstrap (v. 2019.6; Tibshirani and Efron 1993).   Overlap

of the 95% confidence intervals with zero indicated a lack of statistical significance.

Gene-specific analyses

Putative protein-coding changes were examined for three genes found within regions of highest

genetic divergence across years, *cpq1*, *cpq2* and *cyp333b3*. We extracted the nucleotide sequence for

the coding region of each gene using a custom bioinformatic pipeline, and ExPASy

(https://web.expasy.org/translate/) was used to produce the putative protein sequences from individuals

collected in 2002 and 2017. Protein sequences were aligned to one another using MUSCLE (Edgar

2004) to identify putative amino acid substitutions. Allele frequencies for SNPs which caused these

substitutions were calculated for each collection year. A two-sided 3x2 Freeman-Halton extension of

the Fisher's exact tests with a Bonferroni-corrected alpha value ($\alpha = 0.006$) was used to examine

whether allele frequencies differed significantly among years. Frequencies of these 8 non-synonymous

SNPs were also compared in 2017 collections of *H. zea* from MD and LA using this same approach to

quantify the extent of similarity over geographical distance.

Strength of selection

The strength of selection (*s*) imposed upon ancestral codons in the protein-coding regions of

cpq1, *cpq2*, and *cyp333b3* were examined over time (Falconer and Mackay 1996). We calculated *s* for

two periods of selection: 2002 - 2012, as well as 2012 - 2017. We assumed 5 generations per year

(Reay-Jones 2019) for each selection period, and calculations were performed assuming dominance of

the allele that was increasing in frequency, *p,* no dominance of *p*, and recessiveness of *p*.

Frequency of non-synonymous substitutions prior to 2002

Genomic DNA was isolated from *H. zea* collected from our LA traps in 1998 using a Qiagen

DNeasy isolation kit as indicated above. DNA from each individual was PCR amplified using primers

flanking each non-synonymous mutation (Table S8), which allowed us to determine the frequency of

these mutations prior to 2002. Amplicons were sequenced on an ABI3730xl (Applied Biosystems,

Foster City, CA, USA), and chromatograms were visually inspected for quality filtering and sample

genotyping.

Test for linkage

In 2019, we crossed a population of Cry resistant *H. zea* initiated from late instars collected out

of Cry1 + Cry2 expressing sweet corn ears to Cry susceptible *H. zea* (Benzon Research Inc., Carlisle,

PA). Following 7 days of exposure to Cry1 or Cry1 + Cry2 treated diet (Supplemental Methods),

larval growth phenotypes for F2s from each family were scored and individuals genotyped at non-

synonymous mutations in the *cyp333b3* and *cpq1* genes. For genotyping, DNA from each individual

was PCR amplified using primers in Table S8 and genotyped using a Sanger approach, as above.

Associations between larval genotype and Cry resistance phenotype were analyzed using ANOVA.

References

Ahmed S, Wilkins RM, Mantle D (1998) Comparison of proteolytic enzyme activities in adults of insecticide resistant and susceptible strains of the housefly *M. domestica* L. *Insect Biochem Mol Biol* 28(9): 629-639.

Bagwell RD, Graves JB, Leonard BR, Burris E, Micinski S, Fife JH, Mascarenhas V (1996) Status of insecticide resistance in tobacco budworm and bollworm in Louisiana during 1995. In Proceedings Beltwide Cotton Prod. Conf., National Cotton Council, Memphis, TN. pp. 1059-1067

Banks DJ, Jurat-Fuentes JL, Dean DH, Adang MJ (2001) *Bacillus thuringiensis* Cry1Ac and Cry1Fa delta-endotoxin binding to a novel 110kDa aminopeptidase in *Heliothis virescens* is not N-acetylgalactosamine mediated. *Insect Biochem Mol Biol* 31: 909-918.

Baucom RS (2019) Evolutionary and ecological insights from herbicide-resistant weeds: what have we learned about plant adaptation, and what is left to uncover? *New Phytol* 223:68-82.

Bi K, Linderoth T, Singhal S, Vanderpool D, Patton JL, Nielsen R, et al. (2019) Temporal genomic contrasts reveal rapid evolutionary responses in an alpine mammal during recent climate change. *PLoS Genet* 15(5): e1008119. https://doi.org/10.1371/journal.pgen.1008119

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: A flexible trimmer for Illumina Sequence Data. *Bioinformatics*, btu170.

Bown DP and Gatehouse JA (2004) Characterization of a digestive carboxypeptidase from the insect pest corn earworm (*Helicoverpa armigera*) with novel specificity towards C-terminal glutamate residues. *EJB* 271(10): 2000-2011.

Catchen J, Amores A, Hohenlohe P, Cresko W, Postlethwait J (2011). *Stacks: building and genotyping loci de novo from short-read sequences. G3* 1:171-182.

Crossley MS, Chen YH, Groves RL, Schoville SD. Landscape genomics of Colorado potato beetle provides evidence of polygenic adaptation to insecticides (2017). Mol Ecol 26(22):6284-6300. doi: 10.1111/mec.14339.

Crow JF (1957) Genetics of insect resistance to chemicals. *Ann Rev Entomol* 2: 227-246.

d'Alençon E, Sezutsu G, Legeai F, Permal E, Bernard-Samain S, Gimenez S, Gagneur C, Cousserans F, Shimomura M, Brun-Barale A, Flutre T, Couloux A, East P, Gordon K, Mita K, Quesneville H, Fournier P, Feyereisen R (2010) Extensive synteny conservation of holocentric chromosomes in Lepidoptera despite high rates of local genome rearrangements. *PNAS* 107(17): 7680-7685; DOI: 10.1073/pnas.0910413107

Da Silva IHS, Goméz I, Sánchez J, Martínez de Castro DL, Valicente FH, Soberón M, et al. (2018) Identification of midgut membrane proteins from different instars of *Helicoverpa armigera* (Lepidoptera: Noctuidae) that bind to Cry1Ac toxin. *PLoS ONE* 13(12): e0207789. https://doi.org/10.1371/journal.pone.0207789

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker R, LUnter G, Marth G, Sherry ST, McVean G, Durbin R and 1000 Genomes Project Analysis Group (2011) The Variant Call Format and VCFtools. *Bioinformatics* 27(15): 2156–2158.

Danecek P, Schiffels S, Durbin R (2016) Multiallelic calling model in bcftools (-m),
samtools.github.io/bcftools/call-m.pdf

Délye C, Jasieniuk M, Le Corre V (2013) Deciphering the evolution of herbicide resistance in weeds.
Trends Genet 29:649-658.

Dively GP, Venugopal PD, Finkenbinder C (2016) Field-Evolved Resistance in Corn Earworm to Cry
Proteins Expressed by Transgenic Sweet Corn. *PLoS ONE* 11(12): e0169115.

Dively G.P., P. D. Venugopal, D. Bean, J. Whalen, K. Holmstrom, T.P. Kuhar. H.B. Doughty, T.
Patton, B. Cissel, and W.D. Hutchison. (2018) Regional pest suppression associated with
widespread Bt maize adoption benefits vegetable growers. *Proc Natl Acad Sci* 115:3320-3325.

Dong K, Du Y, Rinkevich F, Nomura Y, Xu P, Wang L, Silver K, and Zhorov BS (2014) Molecular
biology of insect sodium channel and pyrethroid resistance. *Insect Biochem Mol Biol* 50: 1-17.

Donnelly MJ, Isaacs A, Weetman D (2016) Identification, validation, and application of molecular
diagnostics for insecticide resistance in malaria vectors. *Trends Parasitol* 32:197-206.

Edgar RC (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nuc Acids Res 32(5):1792–1797. https://doi.org/10.1093/nar/gkh340

Fabrick JA, Ponnuraj J, Singh A, Tanwar RK, Unnithan GC, Yelich AJ, Li X, Carrière Y, Tabashnik
BE (2014) Alternative splicing and highly variable cadherin transcripts associated with field-
evolved resistance of pink bollworm to Bt cotton in India. *PLoS One* 9(5): e97900.

Falconer DS, Mackay TFC (1996) Introduction to Quantitative Genetics. 4th Edition, Prentice Hall,
Essex.

Feyereisen R, Dermauw W, Van Leeuwen T (2015)Genotype to phenotype, the molecular and
physiological dimensions of resistance in arthropods. *Pestic Biochem Physiol* 121:61-77.

Ffrench-Constant RH (2013) The molecular genetics of insecticide resistance. *Genet* 194(4):807-815.

Fritz ML, Paa S, Baltzegar J, Gould F (2016) Application of a dense genetic map for assessment of
genomic responses to selection and inbreeding in *Heliothis virescens*. *Insect Mol Biol* 25(4): 385-
400.

Fritz ML, DeYonke AM, Papanicolaou A, Micinski S, Westbrook J, Gould F (2018) Contemporary
evolution of a Lepidopteran species, *Heliothis virescens*, in response to modern agricultural
practices. *Mol Ecol* 27(1): 167-181.

Fritz ML, Nunziata SO, Guo R, Tabashnik BE, Carrière Y (2020) Mutations in a novel cadherin
gene associated with Bt resistance in *Helicoverpa zea*. *G3* 10(5):1563-1574.

Gahan LJ, Gould F, Heckel DG (2001) Identification of a gene associated with Bt resistance in
Heliothis virescens. *Sci* 293(5531): 857-60.

Gahan LJ, Gould F, Lopez JD, Micinski S, and Heckel DG (2007) A polymerase chain reaction screen
of field populations of *Heliothis virescens* for a retrotransposon insertion conferring resistance to
Bacillus thuringiensis toxin. *J Econ Entomol* 100: 187-194.

Gahan LJ, Pauchet Y, Vogel H, Heckel DG (2010) An ABC transporter mutation is correlated with insect resistance to *Bacillus thuringiensis* Cry1Ac toxin. *PLoS Genet* 6: e1001248.

Gill S, Cowles EA, Francis V (1995) Identification, isolation, and cloning of a *Bacillus thuringiensis* Cry1Ac toxin-binding protein form the midgut of the lepidopteran insect, *Heliothis virescens*. *J Biol Chem* 270: 27277-27282.

Goudet J (2005) hierfstat, a package for r to compute and test hierarchical *F*-statistics. *Mol Ecol Notes* 5(1):184-186.

Gould F (1991) The evolutionary potential of crop pests. *Am Sc* 79: 496-507.

Gould F (1998) Sustainability of transgenic insecticidal cultivars: integrating pest genetics and ecology. *Annu Rev Entomol* 43: 701–726.

Gould F, Blair N, Reid M, *et al*. (2002) *Bacillus thuringiensis*-toxin resistance management: stable isotope assessment of alternate host use by *Helicoverpa zea*. *Proc Natl Acad Sci USA* 99:16581–16586.

Gould F, Brown ZS and Kuzma J (2018) Wicked evolution: Can we address the sociobiological dilemma of pesticide resistance? *Sci* 360(6390): 728-32.

Guo Z, Kang S, Zhu Z, Zia J, Wu Q, Wang S, Xie W, Zhang Y (2015a) Down-regulation of a novel ABC transporter gene (Pxwhite) is associated with Cry1Ac resistance in the diamondback moth, *Plutella xylostella* (L.) *Insect Biochem Mol Biol* 59: 30-40.

Guo Z, Kang S, Chen D, Wu Q, Wang S, Xie W, et al. (2015b) MAPK Signaling Pathway Alters Expression of Midgut ALP and ABCC Genes and Causes Resistance to *Bacillus thuringiensis* Cry1Ac Toxin in Diamondback Moth. *PLoS Genet* 11(4): e1005124. https://doi.org/10.1371/journal.pgen.1005124

Guzov VM, Unnithan GC, Chernogolov AA, Feyereisen R (1998) CYP12A1, a Mitochondrial Cytochrome P450 from the House Fly. *Arc Biochem Biophys* 359(2):231-240.

Hartstack AW, Witz JA, Buck DR (1979) Moth traps for the tobacco budworm. *J Econ Entomol* 72: 519-522.

Hartstack AW, Lopez JD, Muller RA, Sterling WL, King EG (1982) Evidence of long range migration of *Heliothis zea* (Boddie) into Texas and Arkansas. *Southwestern Entomol* 7:188–201.

Heckel DG (2020) How do toxins from Bacillus thuringiensis kill insects? An evolutionary perspective. *Arc Insect Biochem Physiol* 104:e21673

Howe GA, Herde M (2015) Interaction of plant defense compounds with the insect gut: new insights from genomic and molecular analyses. *Curr Opin Insect Sci* 9: 62-68.

Hudson RR (2002) Generating samples under a Wright–Fisher neutral model of genetic variation. *Bioinformatics* 18(2): 337–338.

Hutchison, W. D., E. C. Burkness, P. D. Mitchell, R. D. Moon, T. W. Leslie, S. J. Fleischer, M. Abrahamson, K. L. Hamilton, K. L. Steffey, M. E. Gray, et al. (2010) Areawide suppression of European corn borer with Bt maize reaps savings to non-Bt maize growers. *Science* 330: 222–225.

Jin L, Wang J, Guan F, Zhang J, Yu S, Liu S, Xue Y, Li L, Wu S, Wang X, Yang Y, Abdelgaffar H, Jurat-Fuentes JL, Tabashnik B, Wu Y (2018) Dominant point mutation in a tetraspanin gene associated with field-evolved resistance of cotton bollworm to transgenic Bt cotton. *Proc Natl Acad Sci USA* 115(46): 11760-11765.

Jurat-Fuentes JL, Adang MJ (2004) Characterization of a Cry1Ac-receptor alkaline phosphatase in susceptible and resistant *Heliothis virescens* larvae. *Eur J Biochem* 271(15): 3127-3135.

Jombart T, Devillard S, Balloux F (2010) Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genet* 11:94. doi:10.1186/1471-2156-11-94.

Jombart T, Ahmed I (2011) adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. *Bioinformatics* 27(21): 3070-3071. https://doi:10.1093/bioinformatics/btr521

Jones CM, Parry H, Tay WT, Reynolds DR, Chapman JW (2019) Movement ecology of pest *Helicoverpa*: implications for ongoing spread. *Annu Rev Entomol* 64: 277–295.

Joußen N, Agnolet S, Lorenz S, et al. (2012) Resistance of Australian *Helicoverpa armigera* to fenvalerate is due to the chimeric P450 enzyme CYP337B3. *Proc Natl Acad Sci U S A* 109(38):15206-15211. doi:10.1073/pnas.1202047109

Kaur G, Guo J, Brown S, Head GP, Price PA, *et al.* (2019) Field-evolved resistance of *Helicoverpa zea* (Boddie) to transgenic maize expressing pyramided Cry1A.105/Cry2Ab2 proteins in northeast Louisiana. *J Inv Pathol* 163: 11-20.

Langmead B, Salzberg S (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357-359.

Laurentino TG, Moser D, Roesti M, Ammann M, Frey A, Ronco F, Kueng B, Berner D (2020) Genomic release-recapture experiment in the wild reveals within-generation polygenic selection in stickleback fish. *Nat Comm Biol* 11(1) DOI: 10.1038/s41467-020-15657-3

Leon RG, Dunnea JC, Gould F (2020) The role of population and quantitative genetics in modern sequencing technologies to understand evolved herbicide resistance and weed fitness. *Pest Manag Sci* DOI 10.1002/ps.5988

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R; 1000 Genome Project Data Processing Subgroup (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25(16):2078-9. doi: 10.1093/bioinformatics/btp352. Epub 2009 Jun 8.

Liu C, Xiao Y, Li X, Oppert B, Tabashnik BE, Wu K (2015) *Cis*-mediated down-regulation of a trypsin gene associated with Bt resistance in cotton bollworm. *Sci Rep* 4: 7219.

Luttrell RG, Jackson RE (2012) *Helicoverpa zea* and Bt cotton in the United States. *GM Crops Food* 3: 213.

Magoc T, Salzberg S (2011). FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27(21): 2957-63.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet. journal*, *17*(1), 10-12.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, *et al.* (2010). The Genome
Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.
Genome Res 20(9): 1297–1303. doi:10.1101/gr.107524.110

Melander AL (1914). Can insects become resistant to sprays? *J Econ Entomol* 7:167-173.

Micinski S, Blouin DC, Waltman WF, Cookson C (2008) Abundance of *Helicoverpa zea* and *Heliothis*
virescens in pheromone traps during the past twenty years in Northwestern Louisiana. *Southwest*
Entomol 33:139 - 49. http://dx.doi.org/10.3958/0147-1724-33.2.139

Morin S, Henderson S, Fabrick JA, *et al.* (2004) DNA-based detection of Bt resistance alleles in pink
bollworm. *Insect Biochem Mol Biol* 34(11):1225-1233. doi:10.1016/j.ibmb.2004.08.003

Morin S, Biggs RW, Sisterson MS, Shriver L, Ellers-Kirk C, Higginson D, Holley D, Gahan LJ,
Heckel DG, Carrière Y, Dennehy TJ (2003) Three cadherin alleles associated with resistance to
Bacillus thuringiensis in pink bollworm. *Proc Nat Acad Sci U S A* 100(9): 5004-9.

National Academies of Sciences, Engineering, and Medicine [NASEM] (2016) Genetically Engineered
Crops: Experiences and Prospects. Washington, DC: The National Academies Press. doi:
10.17226/23395.

Newcomb RD, Campbell PM, Ollis DL, Heah EC, Russell RJ, and Oakeshott JG (1997) A single
amino acid substitution converts a carboxylesterase to an organophosphorus hydrolase and confers
insecticide resistance on a blowfly. *Proc Natl Acad Sci USA* 94:7464 –7468.

Nosil P, Villoutreix R, de Carvalho CF, Farkas TE, Soria-Carrasco V, Feder JL, Crespi BJ, Gompert Z
(2018) Natural selection and the predictability of evolution in Timema stick insects. *Sci* 359
(6377):765-770.

Palumbi SR (2001) Humans as the world's greatest evolutionary force. *Sci* 293(5536):1786-1790.

Pearce SL, Clarke DF, East PD *et al.* (2017) Genomic innovations, transcriptional plasticity and gene
loss underlying the evolution and divergence of two highly polyphagous and invasive *Helicoverpa*
pest species. *BMC Biol* 15: 63. doi:10.1186/s12915-017-0402-6

Pélissié B, Crossley MS, Cohen ZP, and Schoville SD (2018) Rapid evolution in insect pests: the
importance of space and time in population genomics studies. *Curr Opin Insect Sci* 26: 8-16.

Pembleton LW, Cogan NOI, Forster JW (2013) StAMPP: an R package for calculation of genetic
differentiation and structure of mixed-ploidy level populations. *Mol Ecol Res* 13(5): 946-952.

Perera OP and Blanco CA (2011). Microsatellite variation in *Helicoverpa zea* (Boddie) populations in
the Southern United States. *Southwestern Entomol* 36(3):271-286.

R Development Core Team (2008). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, http://www.R-
project.org.

Reay-Jones FPF (2019) Pest Status and Management of Corn Earworm (Lepidoptera: Noctuidae) in
Field Corn in the United States. *J Int Pest Manag* 10(1):19 https://doi.org/10.1093/jipm/pmz017

Reisig DD, Huseth AS, Bacheler JS, Aghaee MA, Braswell L, *et al*. (2018) Long-term empirical and observational evidence of practical *Helicoverpa zea* resistance to cotton with pyramided Bt toxins. *J Econ Entomol* 111(4): 1824-33.

Reznick DN, Losos JB, Travis J (2019) From low to high gear: There has been a paradigm shift in our understanding of evolution. *Ecol Lett* 22:233–44.

Roush RT, McKenzie JA (1987) Ecological genetics of insecticide and acaricide resistance. *Ann Rev Entomol* 32:361-80.

Rubin CJ, Zody MC, Eriksson J, Meadows JR, Sherwood E, Webster MT, Jiang L, Ingman M, Sharpe T, Ka S, Hallböök F (2010) Whole-genome resequencing reveals loci under selection during chicken domestication. *Nature* 464(7288): 587.

Russell RJ, Scott C, Jackson CJ, et al. (2011) The evolution of new enzyme function: lessons from xenobiotic metabolizing bacteria versus insecticide-resistant insects. *Evol App* 4(2):225-248. DOI: 10.1111/j.1752-4571.2010.00175.x.

Seymour M, Perera OP, Fescemyer HW, Jackson RE, Fleischer SJ, Abel CA (2016) Peripheral genetic structure of *Helicoverpa zea* indicates asymmetrical panmixia. *Ecol Evol* 6(10):3198-207. https://doi:10.1002/ece3.2106

Tabashnik BE, Gould F, Carrière Y (2004) Delaying evolution of insect resistance to transgenic crops by decreasing dominance and heritability. *J Evol Biol* 17:904-912.

Tabashnik BE, Carrière Y (2017) Surge in insect resistance to transgenic crops and prospects for sustainability. *Nature Biotechnol* 35(10):926.

Tay WT, Mahon RJ, Heckel DG, Walsh TK, Downes S, James WJ, Lee SF, Reineke A, Williams AK, Gordon KHJ (2015) Insect resistance to *Bacillus thuringiensis* toxin Cry2Ab is conferred by mutations in an ABC transporter subfamily A protein. *PLoS Genet* 11: e1005534.

Taylor MF, Heckel DG, Brown TM, Kreitman ME, Black B (1993) Linkage of pyrethroid insecticide resistance to a sodium channel locus in the tobacco budworm. *Insect Biochem Mol Biol* 23(7): 763-775.

Tibshirani RJ, Efron B (1993). An introduction to the bootstrap. *Mono Statist Applied Prob* 57: 1-436.

U.S. Environmental Protection Agency [US-EPA] (1998) The environmental protection agency's white paper on Bt plant-pesticide resistance management. Washington: http://www.epa.gov/scipoly/sap/meetings/1998/february/finalfeb.pdf

U.S. Environmental Protection Agency [US-EPA]. (2018) Resistance of Lepidopteran Pests to Bacillus thuringiensis (Bt) Plant Incorporated Protectants (PIPs) in The United States. SAP Minutes No. 2018-06. FIFRA Scientific Advisory Panel Meeting, 17-19 July 2018 Arlington, VA.

Van Leeuwen T, Dermauw W, Mavridis K, Vontas J (2020) Significance and interpretation of molecular diagnostics for insecticide resistance management of agricultural pests. *Curr Opin Insect Sci* 39:69-76.

Ward CW (1976) Properties of the major carboxypeptidase in the larvae of the webbing clothes moth *Tineolla bisselliella*. *Biochim Biophys Acta* 429: 564–572.

Weetman D, Wilding CS, Neaffsey DE, Muller P, Ochomo E, Isaacs AT, *et al.* (2018) Candidate-gene
based GWAS identifies reproducible DNA markers for metabolic pyrethroid resistance from
standing genetic variation in East African *Anopheles gambiae*. *Sci Rep* 8:2920.

Weir BS, Cockerham CC (1984) Estimating F-Statistics for the Analysis of Population-Structure.
Evolution 38: 1358-1370.

Whitlock MC, Lotterhos KE (2015) Reliable detection of loci responsible for local adaptation:
Inference of a neutral model through trimming the distribution of FST. *Am Nat* 186(S1): S24-S36.

WuY (2014) Detection and mechanisms of resistance evolved in insects to Cry toxins from *Bacillus*
thuringiensis. *Adv Insect Physiol* 47, 297–342.

Yang F, González JC, Williams J, Cook DC, Gilreath RT (2019) Occurrence and ear damage of
Helicoverpa zea on transgenic *Bacillus thuringiensis* maize in the field in Texas, US and its
susceptibility to Vip3A protein. *Toxins* 11(2): 102.

Yang F, Head GP, Price PA, Gonzáleza JCS, Kerns DL (2020) Inheritance of *Bacillus thuringiensis*
Cry2Ab2 protein resistance in *Helicoverpa zea* (Lepidoptera: Noctuidae). *Pest Manag Sci* DOI
10.1002/ps.5916

Zhang H, Wu S, Yang Y, Tabashnik BE, Wu Y (2012) Non-Recessive Bt Toxin Resistance Conferred
by an Intracellular Cadherin Mutation in Field-Selected Populations of Cotton Bollworm. *PLoS*
ONE 7(12): e53418. https://doi.org/10.1371/journal.pone.0053418

Zhang H, Tang M, Yang F, Yang Y, Wu Y (2013) DNA-based screening for an intracellular cadherin
mutation conferring non-recessive Cry1Ac resistance in field populations of *Helicoverpa armigera.*
Pesticide Biochem Physiol 107(1):148-152.

Zhang Z, Teng X, Li F (2017) Knockdown of two Cadherin genes confers resistance to Cry2A and
Cry1C in *Chilo suppressalis*. *Sci Rep* 7: 5992.

Zhang M, Wei J, Ni X, Zhang J, Jurat-Fuentes JL *et al*. (2019) Decreased Cry1Ac activation by midgut
proteases associated with Cry1Ac resistance in *Helicoverpa zea*. *Pest Manag Sci* 75: 1099–1106.
https://doi.org/10.1002/ps.5224

Tables

Table 1 - Candidate genes previously documented as being involved in Bt resistance in lab or field-

collected Lepidopteran species, their best alignments to the *H. zea* genome. We calculated the average

Weir and Cockerham's F_{ST} value for SNPs within the gene-containing region of each scaffold. The

number of SNPs used to calculate the average F_{ST} value can be found in parentheses. Whole genome

sequencing samples from the years 2002 and 2017 were used for the comparison. We also report

where the nearest SNP marker produced by ddRAD sequencing was found relative to each candidate

gene and its F_{ST} value. When ddRAD SNPs were found within the gene, the average F_{ST} value across

SNPs within the gene are reported. When the ddRAD SNP was found outside of the gene, the F_{ST}

value for the nearest SNP is reported.

Candidate Gene	Uniprot/ Genbank ID	Citation	*H. zea* scaffold	Gene Position (bp)	Avg F_{ST} (Num SNPs)	Nearest ddRAD SNP (bp)	ddRAD Avg F_{ST} (Num SNPs)
alp	ACP39715.1	Jurat-Fuentes and Adang 2004	KZ117832.1	1,946 - 16,347	-0.011 (171)	NA	NA
apn1	Q11000.1	Gill *et al.* 1995	KZ118301.1	140,429 - 161,526	-0.005 (353)	in gene	-0.003 (3)
apn4	AF378666.1	Banks *et al.* 2001	KZ118301.1	168,017 - 174,495	0.056 (170)	22,238	-0.005 (1)
abcA2	KP259911.1	Tay *et al.* 2015	KZ118207.1	171,103 - 192,554	0.002 (217)	in gene	0.006 (2)

abcC2	GQ332571.1	Gahan *et al.* 2010	KZ118297.1	98,882 - 114,139	-0.004 (100)	3,178	-0.004 (1)
abcG1 (white)	KM667971.1	Guo *et al.* 2015a	KZ118424.1	2,680 - 24,508	0.015 (58)	12,964	0.001 (1)
calp	AAK85198.1	Gahan *et al.* 2001	KZ117563.1	271,234 - 288,320	0.015 (201)	56,926	-0.007 (1)
cad2	AGG36450.2	Zhang *et al.* 2017	KZ118195.1	87,164 - 95,177	-0.003 (44)	20,535	-0.003 (1)
cad-86C	*H. zea* assembly v. 1.0 ID 537580	Fritz *et al.* 2020	KZ117463.1	550,935- 603,715	-0.008 (1079)	in gene	-0.004 (2)
map4K4	AKC96184.1	Guo *et al.* 2015b	KZ118817.1	92,551 - 104,845	-0.018 (65)	35,125	0.005 (1)
tspan1	AYM54687.1	Jin *et al.* 2018	KZ118424.1	106,132 - 117,025	-0.008 (164)	27,211	0.003 (1)

Table 2 - Putative amino acid substitutions (AA Sub) for two *carboxypeptidase Q* genes (*cpq1* and
cpq2) and *cyp333b3*. We quantified frequency changes in the codon present in the reference genome
(Ref Codon) for *H. zea* populations collected in 1998 (n chr = 40-52), 2002 (n chr = 26), 2012 (n chr =
22), and 2017 (n chr = 22). A Freeman-Halton extension of a Fisher's exact test with a bonferroni
correction ($\alpha = 0.006$) determined statistical significance (*). *Cpq1* extended from 8583 to 10,070 bp
on scaffold 569, and *cpq2* extended from 151,736 to 160,706 bp, while cyp333b3 extended from
32,819 to 39,355 bp on scaffold 1612. All three genes are transcribed from the reverse strand.
861

Scaffold	Gene	SNP Pos (bp)	AA Sub	Ref Codon	2002 freq	2012 freq	2017 freq	p-val
KZ118395.1	*cpq1*	9,409	A → V	GCT	0.73	0.32	0.14	0.0001*
KZ118395.1	*cpq2*	152,183	T → I	ATA	0.83	0.73	1	0.022
KZ118395.1	*cpq2*	152,595	A → V	GTA	0.84	0.77	1	0.049
KZ117131.1	*cyp333b3*	35,685	S → T	AGC	0.65	0.32	0.18	0.0001*
KZ117131.1	*cyp333b3*	36,339	K → N	TTT	0.69	0.27	0.09	0.0003*
KZ117131.1	*cyp333b3*	36,579	G → V	GGA	0.69	0.27	0.09	< 0.0001*
KZ117131.1	*cyp333b3*	36,959	T → A	GCC	0.62	0.95	1	< 0.0001*
KZ117131.1	*cyp333b3*	37,490	S → N	AGC	0.73	0.32	0.09	0.003*

Figures

Figure 1 - Posterior membership probabilities for *H. zea* individuals collected in Bossier Parish, LA, in 2002, 2007, 2012, and 2016.
Genotypic clustering used ddRAD-seq generated SNP data and assumed a prior number of clusters (k) equal to two.

871
Figure 2 - F_{ST} outlier analysis of *H. zea* collected from Bossier Parish, LA in the years 2002, 2007, 2012, and 2016. Each grey point
represents one of the 14,398 SNPs examined in our total dataset. Those circled in black represent SNPs that were identified as outliers and
indicate they were under directional selection.

877
878 **Figure 3 -** Manhattan plots of genomic divergence for individuals collected from pheromone-baited
879 traps in Bossier Parish, LA in A) 2002 and 2017, B) 2002 and 2012, and C) 2012 and 2017.
880 Alternating light and dark grey points demarcate averaged F_{ST} values for 40kb windows with a 10kb
881 step for each of the 2,975 genomic scaffolds. Red points represent statistically significant genetic
882 divergence.
883
884

885
886 **Figure 4** - Values of *s*, the selection coefficient, required to produce the observed allele frequency
887 changes at SNPs within protein-coding regions in field-collected *H. zea* during two selection periods:
888 2002-2012 and 2012-2017. Points represent the mean values (n = 6 SNPs), and error bars represent
889 standard error of the mean.

Figure 5 – Association of F$_2$ larval growth phenotypes on Cry-treated diet with nonsynonymous substitutions that cause protein coding
changes in *cpq1* and *cyp333b3*. After one week of growth with **A.** Cry1 or **B.** Cry1 + Cry2 containing diet the susceptible control strain (light
gray) grew 7 and 10 times less on the two diets respectively than the resistant Maryland population (dark gray). No significant association
was detected between Cry toxin resistance phenotype and non-synonymous substitutions in the sweep region for F$_2$ progeny of these lines
according to a Kruskall Wallis test. Box plots compare weight after one week of growth on Cry1 (**C, E**) or Cry1 + Cry2 containing diet (**D,
F**) for F$_2$ progeny across genotypes in two genes, *cpq1* (**C, D**), *cyp333b3* (**E, F**).

898