A novel juxtamembrane basolateral targeting motif regulates TGF-β receptor signaling in *Drosophila*

Aidan J. Peterson [1], Stephen J. Murphy [2], Melinda G. Mundt [1], Maryjane Shimell [1], Edward B. Leof [3], and Michael B. O'Connor [1#]

1) Department of Genetics, Cell Biology & Development and the Developmental Biology Center. University of Minnesota, Minneapolis, MN 55455, USA

2) Center for Individualized Medicine, Mayo Clinic College of Medicine, Rochester, MN 55905

3) Thoracic Diseases Research Unit, Department of Pulmonary and Critical Care Medicine, Mayo Clinic College of Medicine, Rochester, MN 55905

#Correspondence
Michael B. O'Connor
5-235 Moos Tower
515 Delaware St SE
Minneapolis, MN 55455
Email: moconnor@umn.edu

Abstract

In polarized epithelial cells, receptor-ligand interactions can be restricted by different spatial distributions of the two interacting components giving rise to an underappreciated layer of regulatory complexity. We explored whether such regulation occurs in the Drosophila wing disc, an epithelial tissue that requires the TGF-β family member Dpp for growth and patterning. Dpp protein has been observed in a gradient within the columnar cells of the disc, but also uniformly in the disc lumen leading to the question of how graded signaling is achieved in the face of two distinctly localized pools. We find the Dpp type II receptor Punt, but not the type I receptor Tkv, is enriched at the basolateral membrane, and depleted at the junctions and apical surface. Wit, a second type II receptor, shows a markedly different behavior, with the protein detected on all membrane regions but enriched at the apical side. Mutational studies enabled us to identify the BLT, a novel, short juxtamenbrane sequence that confers basolateral targeting of Punt in both wing discs and mammalian MDCK cells. Rescue of *punt* mutants with transgenes altered in the targeting motif showed that flies expressing apicalized Punt due to the lack of a functional BLT displayed developmental defects, female sterility and significant lethality. We also show that apicalized Punt does not produce an ectopic signal, indicating that the apical pool of Dpp is not a significant signaling source even when presented with Punt. Finally, we present evidence that the BLT acts through polarized sorting machinery that differs between types of epithelia. This suggests a code whereby each epithelial cell type may differentially traffic common receptors to enable distinctive responses to spatially localized pools of extracellular ligands.

Introduction

Polarization of cells underlies core behaviors of cells and tissues by imparting directionality to important processes including adhesion, uptake, and secretion. A common structural motif in metazoans is the epithelium, a contiguous sheet of cells connected by cell-cell junctions. Epithelial cells are inherently polarized, typically possessing a basement membrane at the basal surface and an apical surface exposed to a lumen. The junctions provide compartmentalization by partitioning the basolateral and apical membrane domains *within* a cell, and collectively form a barrier separating the basolateral and apical fluids pools *surrounding* the cell (Bryant & Mostov, 2008). Beyond these common features, epithelia have diverse morphologies and molecular characteristics befitting their tissue type.

In the context of cell-cell signal transduction, polarized epithelia have access to regulatory schemes beyond the simple ligand-receptor paradigm. Because the ligand pools and the membrane domains are each divided into two independent partitions, there are conceptually four combinations of ligand and receptor availability in which effective signaling only occurs when exposed receptors can bind the ligand. Some signaling cascades originate at the apical surface, such as the Notch pathway wherein apical receptors bind to ligands in the lumen (Sasaki et al 2007). TGF-β receptors are restricted to basolateral membrane domains in MDCK epithelial cell culture, and addition of exogenous ligand produces robust signaling only when applied to the basolateral fluid (Murphy et al 2004). In other cases, ligand and receptor isolation conditionally limits signaling. For example, the heregulin-α ligand and its receptor are physically separated in intact airway epithelium, but are exposed to each other upon wounding and thence signal to promote tissue repair (Vermeer et al 2003).

In a case featuring two pools of ligand, the localization of receptors could control the response to multiple potentially overlapping signals in a tissue. Gibson and Schubiger (2002) described such a configuration in the *Drosophila* wing disc, where the Dpp ligand is detectable in two spatial pools: a graded stripe centered around the A/P border of the wing pouch, and in the lumenal space between the wing pouch and the squamous peripodial membrane. This raised the question of how the graded domain can act as a morphogen if all cells are exposed to the lumenal Dpp pool. Given the reported basolateral restriction of mammalian TGF-β receptors, we hypothesized that specific receptor localization in the wing disc plays a role in signal transduction and tissue patterning.

We thus undertook a study of TGF-β receptors in the *Drosophila* model system to determine if restricted basolateral localization occurs and how it impacts signaling. Three questions were experimentally addressed: Do receptors exhibit polarized localization? What is the *cis*-acting determinant for basolateral restriction? How does localization impact signal transduction? We find that membrane localization is qualitatively different for Type II receptors, with Punt exhibiting tight basolateral localization in the wing disc. A *cis*-acting basolateral determinant from Punt that functions in insect and mammalian epithelial cells was mapped to the cytoplasmic juxtamembrane region. We show that membrane targeting controls signal output in flies to influence patterning and viability.

Results

<u>TGF-β receptors exhibit diverse membrane localization in wing disc epithelium</u>

TGF-β superfamily signaling is crucial for the growth and patterning of the wing disc. Most prominently, a Dpp gradient patterns the tissue during larval development while promoting proliferation (Affolter & Basler, 2007). The main receptors required for this signaling are Punt (Type II) and Thickveins (Type I). The Gbb ligand and Saxophone receptor also contribute to BMP signaling in the disc (Upadhyay et al 2017). The Activin branch has been reported to promote proliferation (Hevia et al 2017) and influence a subset of Dpp target genes (Peterson et al 2013, Hevia et al 2013). As a first step to build a spatial model of signaling that incorporates epithelial polarity, we assessed the membrane location of TGF-β receptors in wing disc epithelial cells. The disc is composed of epithelial cells that form a sac with a lumenal space. We detected receptors in fixed tissues and compared the membrane distribution to endogenous protein markers for junctions and the subapical membrane domain (Figure 1, Fig1-S1). Tkv was found at both basolateral and apical positions, with some enrichment near the junctions (Fig 1-S1). Because the main Type I receptor does not exhibit restricted membrane positioning, we turned to the critical Type II receptors.

The Type II receptors Punt (encoded by *put*) and Wishful Thinking (Wit, encoded by *wit*) are both present in the wing imaginal disc (Wrana et al 1994, Marques et al 2002), where Punt function is critical for wing development (Nellen et al, 1996). Overexpressed Punt protein exhibited basolateral restriction, with nearly all of the protein staying basal to the septate junction, which is marked by a concentrated stripe of FasIII or Dlg staining (Figure 1, A). By contrast, Wit was enriched in the subapical region overlapping aPKC, with weaker staining along the basolateral membrane (Figure 1, B). Because of the clear difference in membrane localization between these two functionally and structurally similar Type II receptors, we conducted further studies to explore the *cis*-acting determinants of their membrane targeting.

The cytoplasmic domain of Punt directs basolateral restriction in evolutionarily distant epithelia

To coarsely map the targeting activity, we assessed the localization of chimeric Type II proteins (Figure 1, C). Upon expression in the wing discs of transgenic larvae, the Punt:Wit protein had similar distribution to the control Wit:Wit protein, with a clear apical enrichment over the aPKC stripe (Figure 1, D-E). However, the Wit:Punt protein was basolaterally restricted, indicating that the cytoplasmic domain of Punt is responsible for localization (Figure 1, F).

The mammalian TGF-β Type II receptor (TβRII) utilizes a basolateral targeting motif in the cytoplasmic kinase domain (Murphy et al 2007, Yin et al 2013). Given the parallel membrane localization between fly and mammalian receptors, we expressed Punt in MDCK epithelial cells to test the conservation of protein targeting. Remarkably, Punt was targeted to the basolateral membrane domain (Figure 1, H), revealing the action of a cell biology pathway operating on TGF-β receptors separated by over 500 million years of evolution (Huminiecki et al 2009). To dissect the protein regions driving this localization, a series of progressive truncations was analyzed to detect a presumptive targeting motif (Figure 1, G and Figure 1-S2). Surprisingly, removal of the sequence region corresponding to the mammalian "LTA" targeting motif (Murphy et al 2007) did not disrupt basolateral targeting (Figure 1, I). Indeed, removal of nearly the entire cytoplasmic portion of Punt produced a protein with basolateral localization (Figure 1, K and Figure 1-S1). These data reveal that basolateral targeting of TGF-β receptors in epithelia is phenomenologically conserved from flies to mammals, but that different portions of the protein and amino acid sequences direct the sorting of fly Punt and mammalian TβRII.

Identification of the BLT basolateral targeting motif in the juxtamembrane region of Punt

Combining the results from the cytoplasmic truncations in MDCK cells with the chimeric receptors in wing discs, we predicted that the cytoplasmic juxtamembrane portion of Punt is a basolateral determinant. We examined the 19 amino acids between the transmembrane domain and the end of the smallest truncation for predicted targeting motifs and evolutionary conservation. No canonical basolateral targeting motif is present in this sequence (Harada, 2010). There is no detectable amino acid conservation in this region amongst the complete set of *Drosopohila* TGF-β receptors (3 Type I and 2 Type II) (Peterson et al, 2014). However, there is strong conservation of several positions within insect Punt homologs (Figure 2, A). These amino acids fall between the transmembrane motif and the kinase domain, a region not linked to the core ligand binding and signal transduction functions of the receptors.

To directly test the requirement of this region in basolateral localization, we deleted 10 or 19 amino acids and determined the membrane targeting of the altered proteins. The 10 amino acid deletion removes the insect-conserved residues, and the 19 amino acid deletion additionally removes the remaining residues present in the shortest truncation (Figure 2, A and Figure 2-S1). These internal deletions did not destroy signal transduction activity of Punt as judged by accumulation of P-Mad in wing discs (Figure 2-S1). Compared to the Punt-*PstI* pseudo-wt control, the deletion proteins display apical staining, with the 19 amino acid deletion causing more severe apicalization than the 10 amino acid deletion (Figure 2, B-D and Figure2-S2). Strikingly, the same behavior was observed in MDCK cells: the removal of 10 amino acids led to clear apical staining, and Δ19 had somewhat stronger apical localization (Figure 2, E-H). The prediction that a targeting determinant resides in the cytoplasmic juxtamembrane portion of Punt was borne out by these results, and we coin this motif of Punt the BLT (BasoLateral Targeting).

To confirm this result in a context preserving the spacing of the intracellular portion of Punt relative to the transmembrane domain, we tested the behavior of two variants of fruit fly Punt harboring juxtamembrane sequences from related proteins. We used the corresponding portion of honeybee Punt protein as one donor to test the functional conservation of this sequence in insects (Figure 2, A). Transgenic flies expressing the Punt[Apis] protein showed predominantly basolateral staining in wing discs, indicating functional conservation among insect Punt proteins (Figure 2, I-J). Because Wit is apically enriched, we used the *Drosophila* Wit sequence to test if the juxtamembrane amino acid sequence correlates with membrane location. Punt[Wit] showed apical enrichment, much like full length Wit, confirming that the Punt BLT is required for basolateral targeting (Figure 2, K).

To further define the structure-function aspects of the BLT at the level of amino acid sequence, we characterized a series of point mutations in the BLT targeting the amino acids conserved in insects. Alanine substitutions were made for three positions at a time and the localization of the resulting protein, named for conserved mutated residues, was determined in wing discs. The FNE, EIP, and PTE variants were detected primarily basolateral (Figure 2-S3 and Figure 2, L), similar to the WT starting protein. To specifically address a potential role for charged side chains, an EEE variant was tested, which was also basolateral (Figure 2-S3). More extensive mutation of seven conserved residues (FNEIPTE) led to mixed apical and basolateral distribution, and a more drastic alteration converting the entire 19 AA stretch comprising the minimal BLT mapped by the truncations series led to strong apical localization (Figure 2, M-N), which resembled the Δ19 and Punt[Wit] proteins. Considering these results together, we conclude that the BLT amino acid sequence is robust in the face of mutation, and that the conserved amino acids and the surrounding sequence contribute to the basolateral targeting of Punt.

The Punt BLT motif is a dominant, transferable targeting signal

Having established the requirement of the Punt juxtamembrane region in basolateral restriction, we conducted swap experiments to address sufficiency by asking if the BLT acts as a dominant basolateral targeting motif. Wit shares many signaling properties with Punt, but as described above showed a starkly different localization in wing disc epithelium. We thus replaced the corresponding portion of Wit with the Punt BLT. A pseudo-wildtype version of Wit (with a single amino acid change from a cloning scar) and Wit[Apis] displayed strong apical staining that overlaps with the aPKC marker (Figure 2, O-P). Wit[Punt] behaved like intact Punt, with nearly all staining excluded from the junction and apical membranes (Figure 2, Q). The Punt BLT is thus a dominant moveable basolateral targeting motif that confers basolateral restriction in wing disc epithelia on an otherwise apically enriched protein. We queried the positional requirement of the BLT by appending it to the C-terminus of an otherwise apical protein and assaying epithelial distribution. Punt-Δ19-CBLT retains the apical staining of Punt-Δ19 (Figure 2-S3), indicating that the motif does not function at the C-terminus. Since it functions in a juxtamembrane position but not at the C-terminus, we conclude that the function of the BLT motif as a dominant *cis*-acting determinant is contingent on topology.

Punt activity is regulated by polarized membrane localization

All variants of Punt used for our localization tests are active when overexpressed. This may mask functional differences since any overexpressed Type II receptor can bypass normal controls and generate ectopic signaling. To correlate the localization of Type II receptors with function under controlled expression conditions, we generated transgenic lines with variant UAS constructs recombined into fixed genomic attP sites. For Wit, we tested three constructs for their ability to rescue *wit* mutants when expressed in neurons with elav-GAL4 (Marques et al 2002). The control protein, Wit-FLAG, provided full rescue to viability in this assay (Figure 3, A). Wit[Apis] and Wit[Punt] also provided full or substantial rescue (Figure 3, A), indicating that the juxtamembrane domain of Wit is not critical for its function in this assay, and that the Punt BLT does not grossly interfere with Wit function in neurons. We also tested the ability of Punt variants to rescue *wit* mutants. Any form of Punt overexpressed pan-neuronally supported rescue of otherwise lethal *wit* mutants (Figure 3, A). This indicates that some Type II activity is sufficient in neurons to support viability, but yields no evidence that the BLT region is critical to regulate the activity.

A parallel attempt to rescue *put* mutants with UAS-controlled proteins was hampered by toxicity of over-expressed Punt. Widespread ectopic expression of Punt was lethal at 25 degrees in an otherwise wildtype background and in a *put* mutant background (Figure 3-S1). The *put* alleles we used for the rescue tests are temperature sensitive, showing variable survival at lower temperatures (Letsou et al 1995), thus precluding assessment of relative rescue activities with reduced UAS expression levels at lower temperatures. However, we did leverage the temperature sensitivity of the GAL4 system to uncover differential activity of Punt variants with altered BLT regions. Punt[Wit] produced a lower degree of lethality than Punt[Punt] or Punt[Apis] over a range of temperatures with the spatially restricted *en*-GAL4 driver (Figure 3, B). Similar results were obtained for gender-biased lethality with the X-linked A9-GAL4 driver, which is more active in males than females (Figure 3, C).

In vivo rescue

To bypass the toxicity of over-expressed Punt, we generated transgenic lines encoding Punt variants expressed from endogenous regulatory elements. A BAC-based construct expressing GFP-tagged Punt

(Punt-GFP) provided full rescue of *punt*^{135/P1} animals at 25 degrees. A smaller construct including the *punt* genomic region was also able to provide genetic function, with equivalent activity for Punt and Punt-GFP versions. Survival, fertility, and overt behavior were fully rescued by this construct. Rescued animals displayed variable eye defects, however this phenotype tracked with the attP docking site rather than BLT status or basolateral localization (Figure 3-S2). To assess the role of the BLT motif, we compared the WT rescue construct to a BLT point mutant (EG) and a deletion of the BLT (Δ19). As shown by anti-GFP IHC, the WT construct was basolaterally restricted in the wing disc epithelia (Figure 3, D), in line with the distribution of over-expressed Punt. The EG variant showed a limited degree of increased apical staining, and the Δ19 protein was more strongly enriched apically (Figure 3, E-F). In this context, the EG construct provided rescue activity equivalent to the WT construct. Deletion of the BLT domain, however, severely reduced the rescue activity of Punt as assayed by survival to eclosed adults (Figure 3, G). Lethal phase analysis showed that animals with Punt-Δ19 as the sole source of Punt frequently failed during pupation (Figure 3, H). Additionally, eclosed adults displayed reduced movement and died within several days. Finally, the surviving females were completely sterile (Figure 3, H). Taken together, these results reveal that the BLT portion of Punt is crucial for full viability and fertility.

Having established the requirement of the Punt BLT for overall fitness, we returned to the wing to probe the signaling consequence of altering protein localization. We considered two general outcomes: That apical Punt would create a gain-of-function situation due to exposure to and activation by the lumenal Dpp pool, or that a re-direction of Punt from basolateral to apical membranes would create a loss-of-function situation for Dpp signaling. To distinguish between these outcomes, we analyzed molecular and morphological features associated with the Dpp morphogen gradient. In an over-expression context, the lethality results described above for UAS-driven Punt variants are consistent with basolateral Punt having more toxicity, which could suggest this position supports more signaling activity. We linked this to signal transduction by examining p-Mad levels upon expression of Punt variants with the *en*-GAL4 driver. Based on stronger p-Mad staining in the posterior compartment, the basolateral proteins Punt[Punt] and Punt[Apis] produce more ectopic signaling than the apicalized Punt[wit] version (Figure 3, J-N).

To address the same question under physiological conditions, we stained rescued animals for p-Mad as a direct readout of BMP signaling downstream of endogenous Dpp. Larvae with a single copy of Punt-WT had a stereotypical p-Mad pattern and overtly normal wing discs (Figure 3, O). The corresponding EG point mutant genotype also had normal p-Mad detection within normally shaped discs (Figure 3, P). The Δ19 rescue animals, however, frequently displayed a reduced p-Mad staining intensity, and a fraction of the discs presented an abnormal split morphology (Figure 3, Q-R). Both of these phenotypes are consistent with reduced Dpp signaling (Affolter & Basler 2007, Gibson & Schubiger 2002). These results indicate that a Dpp morphogen gradient forms without the membrane targeting function of Punt's BLT, but the output is weakened and exposes the tissue to developmental error. Wings from the subset of surviving adults correspondingly displayed variable shape and veination defects compared to the fully rescued genotypes (Figure 3, S-V).

Tissue specific utilization of the BLT and *trans*-acting sorting factors

Taken together, the overexpression and endogenous level activity assays indicate that altering the localization of Punt changes its activity in at least one tissue, but it is not strictly required for signaling. We were intrigued that the activity of Punt in the wing tracked with overall animal survival. To define the generality of the localization program, we determined the localization of Punt and Wit in two other epithelial tissues. In larval salivary glands, both Punt and Wit were only detected in basolateral

membrane regions, and were excluded from junctions (Fig 4, A-B). In follicle cells of the egg chamber, both Punt and Wit were distributed basolaterally and apically (Figure 4, C-D). Thus, in three different fly epithelia, we see three distinct distribution profiles, indicating that for a given receptor, membrane targeting schemes vary with tissue.

The conclusion that the Punt BLT confers basolateral restriction in the wing disc epithelium is based on observation of steady state protein distribution. Generically, this preferential localization can be achieved by specific delivery to basolateral surfaces or by selective removal from apical surfaces, or by a combination of these actions. As a broad test of these mechanisms, we disrupted a group of trafficking factors to ascertain their requirement for Punt basolateral restriction. We induced RNAi in clones to knock down candidate genes in animals expressing Punt-GFP from a rescuing transgene. This approach is advantageous because it precludes any over-expression or ectopic signaling effects of Punt, avoids secondary consequences of loss of cell biology functions in the entire disc, and permits direct comparison of neighboring cells within the same tissue, thus avoiding potential changes from developmental stage or environmental conditions.

Adaptor protein (AP) complexes are key mediators of vesicular sorting and membrane trafficking. The *Drosophila* genome encodes subunits for 3 AP complexes (Boem and Bonifacino 2001). AP-1 has been implicated in sorting mechanisms that direct cargo to restricted membrane domains, whereas AP-2 has been linked to endocytosis and early endosome movement (Gravotta et al 2012, Bonaficino 2014). We obtained and tested RNAi lines for genes encoding AP-1 and AP-2 components in Drosophila. RNAi constructs for AP-1/2β, AP-1γ, AP-2α, AP-2σ and AP-2μ were active as judged by lethality upon expression with a constitutive GAL driver (see Methods). Wing disc clones expressing each construct were generated by hsFlp and the localization of PuntGFP was compared in clones and nearby normal cells. In nearly all AP-1/2β knock-down clones, PuntGFP was detected at the apical membrane at levels higher than surrounding cells (Figure 4, E). Because in *Drosophila* the β1/2 subunit (Bap) is thought to assemble into both AP-1 and -2 complexes, we tested AP-1γ to clarify the requirement of AP-1. AP-1γ knock-down clones also have apical Punt (Figure 4, F). We thus conclude that the AP-1 complex is required to produce or maintain the basolateral distribution of Punt (Figure 4-S1).

To test the requirement of endocytosis, we knocked down several AP-2-specific subunit genes and Rab5, which is required for early endosome formation in *Drosophila*. In each case, Punt-GFP distribution was not altered in the clones and there was no apparent change in level (Figure 4, H and Figure 4-S1). Even in cases where the morphology of the wing disc was abnormal, Punt-GFP remained in membrane domains basal to the junctions (Figure 4, G). We conclude that endocytosis does not play a significant role in restricting Punt's membrane distribution. Another potential mechanism to remove Punt from the apical membrane is proteolytic cleavage, as has been observed for other receptors including mammalian TGF-β receptors (Liu et al 2009). We analyzed HA-Punt[Punt] and HA-Punt[Wit] expressed in the wing disc by Western blot. We observed no change in protein level and did not detect any shorter peptide containing either the N-terminal epitope or the C-terminal Punt antigen that differed between Punt[Punt] and Punt[Wit] (Figure 4, I-J). Combining the AP-1 data with the negative results for AP-2 and Rab5, we conclude that selective delivery is the most likely mechanism generating the basolateral distribution of Punt via its BLT motif.

Discussion

Punt displays restricted basolateral localization in the wing disc

The goal of this study was to determine if polarized membrane localization of TGF-β receptors impacts growth and patterning of the wing disc. Two observations motivated this investigation. First, Dpp is detected in two distinct pools in the wing disc: A graded population centered in the disc proper, and a more uniform pool in the lumen of the epithelial sac structure (Entchev et al 2000, and Gibson & Schubiger, 2004). The second finding is that TGF-β receptors are basolateral restricted in mammalian epithelia (Murphy et al, 2004). If *Drosophila* receptors display similar restricted membrane presentation, this could spatially restrict signaling by the Dpp ligand.

The first step in probing this question was to determine the membrane distribution of TGF-β receptors. We initially deployed constructs to over-express tagged proteins in the wing disc. Tkv, the primary BMP pathway Type I receptor, was detected in both basolateral and apical membrane domains, with apparent enrichment at the cellular junctions, consistent with studies showing Tkv protein in apical cytoneme projections (Roy et al 2014). This observation supports the general statement that basolateral restriction is not a universal property of TGF-β receptors. We also studied Type II receptors to determine if their distribution suggested a role in limiting Dpp signaling. Indeed, Punt was detected in the basolateral membranes in the wing disc epithelium, but excluded from the apical membrane and cell-cell junctions. In contrast, Wit was enriched apically but detected in basolateral domains. As a coarse *cis*-determinant mapping test, we expressed chimeric receptors swapping the Wit and Punt ecto- and cytodomains. The localization patterns of the chimeras indicated that the cytoplasmic portion of Punt directs basolateral localization. Overall, we observed diverse membrane distribution patterns among receptors expected to mediate Dpp signaling in the wing disc. We focused on Punt because of its obvious restricted localization and implications for interpretation of Dpp signaling, and on the cytodomain in particular because basolateral sorting signals are typically cytoplasmic (Bonifacino 2014) and the targeting activity mapped to this portion of Punt.

The BLT motif is an active basolateral determinant that functions in evolutionarily diverse epithelia

To what extent is membrane targeting conserved between insect and mammalian epithelia? The basolateral restriction of TβRII and Punt prompted us to assess evolutionary conservation. Notably, Punt exhibited basolateral restriction in MDCK cells, both in the context of a chimera with an unrelated ectodomain and as a full length protein. A series of progressive C-terminally truncated Punt proteins were assayed in MDCK cells to identify regions important for targeting. Deletion of the cytoplasmic region that corresponds to the LTA motif, which is required for TβRII restriction (Murphy et al 2007), did not alter Punt's localization, suggesting that the same behavior of these homologous proteins is mediated by different regions of the proteins. Further deletion variants, including one that retained just 19 amino acids of the Punt cytodomain, also preserved basolateral restriction. This result, combined with the chimera data, suggested that the cytoplasmic juxtamembrane region of Punt harbors a basolateral targeting motif. Deletion of the BLT abolished basolateral restriction in both MDCK cells and wing disc epithelial cells, establishing the necessity of this portion of the receptor. Sufficiency is suggested by the basolateral detection of severely truncated Punt in MDCK cells. To prove that the BLT is a dominant, moveable element we placed it in the cytoplasmic juxtamembrane position of Wit and found it bestowed basolateral restriction on the otherwise apically enriched protein.

None of the several previously described flavors of basolateral targeting motifs (Bonifacino 2014) is present in the short active region of Punt. However, sequence alignments revealed a core KxxxFNEφPTxE sequence within the BLT region among Punt insect homologs, indicative of a sequence-specific element. Simultaneous mutation of several sets of conserved residues did not completely abolish the basolateral preference of overexpressed Punt, indicating robustness of the BLT

motif function. In fact, the only two mutations that completely erased the basolateral preference were 19AA and Punt[wit]. On the other hand, a single E-to-G mutation within the conserved sequence led to partial apicalization of Punt-GFP expressed at the endogenous level. There may be a biochemical or structural difference between the E-to-G and alanine substitution included in the PTE variant, or there may be differences in the trafficking flux of endogenous and over-expressed Punt. In addition to the mixed behavior of Punt[conserved], we uncovered other evidence that non-conserved residues contribute to BLT function. The Δ19 protein displayed more apicalization than the Δ10 version. The robust basolateral sorting of the Punt[Apis] protein is interesting in this context. If viewed as a mutant of all the non-conserved residues, this would mean that the non-conserved positions are not important. However, since the 19 AA stretch is from a native protein, it is likely the BLT from honeybee Punt has been under functional selection for both the strictly conserved residues and the neighboring side chains that support BLT activity. Further work, including identifying binding proteins as discussed below, is required to explain the mechanism of the novel basolateral sorting signal embedded in the BLT.

Mapping studies of three TGF-β receptors with basolateral restriction have uncovered three different targeting motifs, none of which corresponds to canonical motifs (this work, Murphy et al 2007, Yin et al 2018). In the context of myriad families of transmembrane proteins, all of which have the potential for regulated apicobasal presentation, we can speculate that there is a substantial pool of heretofore unrecognized sorting motifs.

<u>Relevance to Dpp signaling in the wing disc</u>

Armed with the identification of the *cis*-acting BLT motif of Punt, we assessed its role in regulating signaling by blocking its function and monitoring signaling *in vivo*. This approach requires direct study of each protein of interest to identify and modify the *cis*-acing sorting determinants. An alternative approach has been developed to assess the outcome of forcing proteins to different apicobasal membrane domains (Harmansa et al 2017). That approach may offer a higher throughput platform for exploring targeting-function relationships in a variety of membrane proteins, but will likely not shed light on the *cis*- and *trans*-acting machinery that execute the normal sorting. For the case of Punt and the wing disc, we wished to define the consequence of mis-localization and the mechanism of specific basolateral targeting.

At the organismal level, removal of the BLT severely harms fitness. Wildtype Punt rescue constructs restored full viability and fertility, but a construct lacking the BLT provided only partial viability and rendered the females sterile. Multiple functional domains of TGF-β receptors have been characterized (Heldin & Moustakas, 2016), including the ectoplasmic ligand-binding domain, cytoplasmic GS box (found in Type I receptors), and cytoplasmic kinase domain. However, function has traditionally not been assigned to the cytoplasmic juxtamembrane portion. Notably, one exception is the VxxEED juxtamembrane motif that acts as a basolateral sorting motif for TβRI (Yin et al 2018). This portion of the receptor thus appears to offer a target for natural selection that can regulate receptor function outside the core ligand-binding and kinase domains. Our data suggests that for Punt, this regulatory function is to control polarized receptor presentation in epithelia.   Wing disc experiments with basolateral and apicalized Punt revealed that apicalized Punt is less active. This holds for the ectopic activity of over-expressed proteins, and for patterning activities of proteins expressed at the endogenous level.

Gibson and Schubiger (2002) speculated about the role of the lumenal Dpp pool in the wing disc, particularly how it relates to the observed morphogen gradient within the plane of the disc proper epithelium. Recent work using a morphotrap system confirmed the presence of Dpp in the lumen, but a

variant technique with apicobasal specificity revealed that the Dpp pool within the tissue is required for patterning and growth of the disc (Harmansa et al 2015, Harmansa et al 2017). This is consistent with our finding that apicalized Punt is less active, in the sense that Dpp and Punt need to be present in the basolateral cell region for effective signaling. On its face, this behavior is somewhat surprising as the exposure of Punt to the lumenal pool might be predicted to generate ectopic signaling. We note that even strongly apicalized Punt has significant basolateral distribution, so it is not technically possible to assess the function of purely apical Punt. Thus, our results can not formally differentiate between complete inactivity of apical Punt, or significantly less activity than basolaterally situated receptor. Tkv should be available in all membrane domains, so is not likely to block apical Dpp signaling. Independent trafficking of Type I and Type II receptors is observed in multiple species (Gleason et al 2014, Norman et al 2016). Further molecular studies are needed to define *in vivo* the spatial and kinetic details of when the Type I and Type II receptors are bound in active signaling complexes. Presently it is not known why apical signaling is tepid, but the overall effect of basolateral targeting of Punt in the wing disc is that it is where it needs to be in order to properly interpret the Dpp morphogen gradient.

A tissue-specific sorting mechanism "reads" the BLT in the wing disc

Returning to the mechanism of how basolateral restriction of Punt is achieved, we screened for *trans*-acting factors that caused relaxed targeting of a functional Punt-GFP protein. With the goal of differentiating between the broad categories of polarized *delivery* and/or polarized *removal*, we perturbed the function of AP1 and AP2 Adapter Protein complexes involved with vesicular movement. The AP1 complex controls delivery of vesicles to the plasma membrane (Bonifacino 2014). Clones with knock-down of AP1 subunits exhibited apical localization of Punt-GFP, indicating that the AP1 complex is required for effective basolateral restriction, and suggests that polarized sorting is a key mechanism. In contrast, loss of AP2-specific subunits or the early endosome regulator Rab5 did not perturb the steady-state basolateral localization of Punt, suggesting that removal of apical Punt is not a significant factor.

The conserved localization behavior directed by the Punt BLT in *Drosophila* wing disc epithelial and mammalian MDCK cells indicates that the machinery to process the BLT instructions are deeply conserved in the insect and mammalian genomes. Thus, we were somewhat surprised to find that polarized localization schemes vary between different *Drosophila* epithelia. In salivary glands, both Punt and Wit were detected at basolateral membranes, but excluded from the junctions and the apical surface. In follicle cells, the same proteins are distributed to basolateral and apical surfaces. This clearly indicates that polarized delivery is cell-type dependent. The BLT motif of Punt is therefore best described as an imaginal disc basolateral targeting motif. Many sorted proteins bind directly to AP complex subunits (Boehm & Bonifacino 2001), but the tissue-specific nature of the sorting renders this unlikely for the BLT motif. We thus propose that the readout of the *cis*-acting motif depends on the *trans*-acting sorting factors expressed in each epithelial tissue. Extrapolating, we postulate that there is a sorting code whereby each epithelial tissue expresses a defined set of sorting factors that read the *cis*-acting determinants in the set of expressed membrane proteins and sorts them to the membrane domains resulting in a unique signaling potential profile for each cell type.

Materials and Methods

<u>Receptor expression constructs and Transgenic lines</u>

UAS-Wit-GFP is described in Smith et al (2012). UAS-Punt-GFP DNA was provided by G. Marques, and random transposition insertions were recovered after injection. Type II chimera constructs were generated by PCR amplification and ligation of the ectoplasmic and cytoplasmic portions of Wit-Flag and Punt-Flag constructs (Gesualdi & Haerry 2007). The WWF construct is a psuedo-WT transgene with a cloning scar that localizes similarly to Wit-GFP. WWF, PW, and WP coding regions were cloned into pUAST and random transposition lines were recovered after injection.

Internal deletion of Punt sequences was achieved by PCR amplification of desired cytoplasmic portion (Δ10 or Δ19) and cloning into a vector modified by Quikchange with a *PstI* sequence at the transmembrane boundary. The pseudo-wt HA-Punt-PstI protein localizes the same as HA-Punt. Intact coding regions were cloned into pUAST and transgenic lines with random transposition insertions were recovered after injection.

Punt constructs with wholesale substitution of the BLT motif were made by dropping oligos into a construct engineered to have restriction sites at the beginning (*PstI*, K185L change) and end (*HindIII*, codon silent) of the 19 AA BLT motif. UAS-attB constructs were injected for recovery of transgenic lines with recombination at the VK20 attP landing site. Similarly, Wit[BLT] variants were made by oligo drop-in to a Wit construct with restriction sites (*NgoMIV* and *HindIII*). UAS-attB constructs were injected and transgenic lines with recombination at the VK31and attP1 docking sites were recovered.

Smaller Punt Point Mutants (3 amino acids converted to Alanine) were generated by QuickChange site-directed mutagenesis, and longer substitutions were generated by oligo drop-in as above. UAS-attB constructs were used to generate transgenic lines at the same site as UAS-Punt[BLT]. C-BLT was generated by appending the 19AA BLT sequence to Punt-Δ10 or Punt-Δ19 with 3' PCR primer. Lines with these variants used the VK20 attP site.

Two versions of Punt rescue constructs were used. A larger version was made by recombineering GFP-encoding sequence at the C-terminus of Punt in BAC CH322-93G2. A smaller rescue construct was generated by PCR amplification of two genomic regions totaling 7 kb that encompass the *put* gene, with addition of a C-terminal GFP tag. Sequencing of the entire coding region revealed a serendipitous substitution converting a Glu to Gly within the BLT (EG variant). The WT Punt-GFP construct was made by deploying Quikchange mutagenesis to revert to the expected sequence. The Punt BLT coding region spans two exons. The Punt-GFP-Δ19 construct was thus made by successive deletions mediated by QuikChange mutagenesis. Constructs were injected to recover transgenic lines recombined into the attP1 or VK15 landing sites.

<u>MDCK cell culture and protein detection</u>

MDCK cell culture, transfection, protein detection, and imaging were performed as described (Murphy et al 2004, Murphy et al 2007). Briefly, the extracellular pool of receptors was detected by antibody staining in fully polarized MDCK cells after transient transfection. Multiple regions containing positive cells were imaged by confocal microscopy, and xz projections were generated to visualize the apicobasal distribution of the proteins of interest. For Punt detection an HA epitope tag (YDVPDYALE) was inserted into the extracellular domain after Pro27. Truncations were generated by PCR amplification of portions of HA-Punt and cloned into pCMV5 or a GM-CSF shuttle vector for expression.

<u>Protein detection</u>

Wing discs and salivary glands from wandering third instar larvae were fixed and subjected to IHC staining as described (Peterson et al 2013). Overexpressed Receptor-GFP constructs were detected

using GFP fluorescence. Other UAS constructs were detected by antibody staining against Punt (rabbit polyclonal, Fabgennix Punt-112-AP), HA (rat monoclonal 3F10), Flag (mouse monoclonal M2, Sigma F3165), Tkv-Pan (rabbit polyclonal), Tkv2 (rabbit polyclonal). Endogenous proteins were detected with anti-aPKC (goat polyclonal, Santa Cruz sc-216), FasIII (mouse monoclonal, DHSB 7G10), Dlg (mouse monoclonal, DHSB 4F3), phospho-Mad (Peterson et al 2013). Punt-GFP rescue constructs are expressed at low levels, so anti-GFP (Abcam, ab6556, pre-adsorbed against fixed larvae prior to use) was used to image these proteins. Fluorescent secondary antibodies were Alexafluor-488, 568, or 647 (Invitrogen). DAPI was used a routine stain for nuclei. For follicle cell imaging, ovaries from yeast-fed females were dissected and fixed as for wing discs. Punt-GFP expression caused severe ovariole dysgenesis with a reduced number of egg chambers.

Most imaging was carried out on a Zeiss LSM710 unit using a 20X objective. Several images were collected using a CARV spinning disc attachment on a Zeiss Axio microscope. Single confocal images are shown for all *Drosophila* images.

For Western blot analysis, rabbit anti-Punt (as above) and anti-HA (rabbit monoclonal, CST C29F4) were used to probed blots, with anti-rabbit-HRP as a secondary antibody.

Experimental Genotypes
Wing disc expression was achieved with A9-Gal4 (BDSC 8761), nub-Gal4 (Calleja et al 1996), or vg-Gal4 (BDSC 8229). en-Gal4 (BDSC 30564) was used to drive expression in posterior compartments, A9-Gal4 was used for salivary gland expression, and C768-Gal was used for follicle cell expression.

Rescue tests of *wit* by UAS-Wit constructs were conducted by crossing elav-Gal4; *wit^{B11}*/Balancer females to UAS-Wit[BLT]; *wit^{412}*/Balancer males. Progeny were counted by genotype from crosses incubated at the indicated temperature. Rescue tests of wit by UAS-Punt constructs used UAS-Punt[BLT], *wit^{412}* recombinants. Rescue tests for Punt used an armGAL4, *put^{P1}* recombinant crossed to PH[BLT], *put^{135}* or WF[BLT], *put^{135}* recombinants, or the UAS elements alone for toxicity tests.

RNAi hs-flp clones were generated and analyzed as described (Peterson et al 2013), but with Punt-GFP (attP1) in the background. UAS-RNAi lines used include AP1/2-beta (BDSC 31707), AP1-gamma (BDSC 27533), AP-2mu (BDSC 28040), AP2-alpha (BDSC 32866), AP2-sigma (BDSC 27322) and Rab5 (BDSC 51847). Activity of these lines was confirmed by production of lethality when crossed to Tub-Gal4.

Acknowledgements

We thank Guillermo Marques for the Punt-GFP construct and for providing Wit-GFP flies prior to publication, Ela Serpe for initial characterization of Punt-GFP flies, Melissa Ritter for cloning and analysis of Punt & Wit, and Autumn Pace for preliminary studies of AP-1 components in the wing. MBO was funded by grant R35GM118029.

References

AFFOLTER, M.; BASLER, K. The Decapentaplegic morphogen gradient: from pattern formation to growth regulation. **Nat Rev Genet**, 8, n. 9, p. 663-674, Sep 2007.

APODACA, G.; GALLO, L. I.; BRYANT, D. M. Role of membrane traffic in the generation of epithelial cell asymmetry. **Nat Cell Biol**, 14, n. 12, p. 1235-1243, Dec 2012.

BOEHM, M.; BONIFACINO, J. S. Adaptins: the final recount. **Mol Biol Cell**, 12, n. 10, p. 2907-2920, Oct 2001.

BONIFACINO, J. S. Adaptor proteins involved in polarized sorting. **J Cell Biol**, 204, n. 1, p. 7-17, Jan 2014.

BRUMMEL, T. J.; TWOMBLY, V.; MARQUES, G.; WRANA, J. L. *et al.* Characterization and relationship of Dpp receptors encoded by the saxophone and thick veins genes in Drosophila. **Cell**, 78, n. 2, p. 251-261, Jul 29 1994.

BRYANT, D. M.; MOSTOV, K. E. From cells to organs: building polarized tissue. **Nat Rev Mol Cell Biol**, 9, n. 11, p. 887-901, Nov 2008.

BURKE, R.; BASLER, K. Dpp receptors are autonomously required for cell proliferation in the entire developing Drosophila wing. **Development**, 122, n. 7, p. 2261-2269, Jul 1996.

CALLEJA, M.; MORENO, E.; PELAZ, S.; MORATA, G. Visualization of gene expression in living adult Drosophila. **Science**, 274, n. 5285, p. 252-255, Oct 1996.

ENTCHEV, E. V.; SCHWABEDISSEN, A.; GONZALEZ-GAITAN, M. Gradient formation of the TGF-beta homolog Dpp. **Cell**, 103, n. 6, p. 981-991, Dec 8 2000.

GESUALDI, S. C.; HAERRY, T. E. Distinct signaling of Drosophila Activin/TGF-beta family members. **Fly (Austin)**, 1, n. 4, p. 212-221, 2007 Jul-Aug 2007.

GIBSON, M. C.; LEHMAN, D. A.; SCHUBIGER, G. Lumenal transmission of decapentaplegic in Drosophila imaginal discs. **Dev Cell**, 3, n. 3, p. 451-460, Sep 2002.

GLEASON, R. J.; AKINTOBI, A. M.; GRANT, B. D.; PADGETT, R. W. BMP signaling requires retromer-dependent recycling of the type I receptor. **Proc Natl Acad Sci U S A**, 111, n. 7, p. 2578-2583, Feb 2014.

GRAVOTTA, D.; CARVAJAL-GONZALEZ, J. M.; MATTERA, R.; DEBORDE, S. *et al.* The clathrin adaptor AP-1A mediates basolateral polarity. **Dev Cell**, 22, n. 4, p. 811-823, Apr 2012.

HARADA, A. Molecular mechanism of polarized transport. **J Biochem**, 147, n. 5, p. 619-624, May 2010.

HARMANSA, S.; ALBORELLI, I.; BIELI, D.; CAUSSINUS, E. *et al.* A nanobody-based toolset to investigate the role of protein localization and dispersal in. **Elife**, 6, 04 2017.

HARMANSA, S.; HAMARATOGLU, F.; AFFOLTER, M.; CAUSSINUS, E. Dpp spreading is required for medial but not for lateral wing disc growth. **Nature**, 527, n. 7578, p. 317-322, Nov 2015.

HELDIN, C. H.; MOUSTAKAS, A. Signaling Receptors for TGF-β Family Members. **Cold Spring Harb Perspect Biol**, 8, n. 8, 08 2016.

HEVIA, C. F.; DE CELIS, J. F. Activation and function of TGFbeta signalling during Drosophila wing development and its interactions with the BMP pathway. **Dev Biol**, 377, n. 1, p. 138-153, May 1 2013.

HEVIA, C. F.; LÓPEZ-VAREA, A.; ESTEBAN, N.; DE CELIS, J. F. A Search for Genes Mediating the Growth-Promoting Function of TGFβ in the. **Genetics**, 206, n. 1, p. 231-249, 05 2017.

HUMINIECKI, L.; GOLDOVSKY, L.; FREILICH, S.; MOUSTAKAS, A. *et al.* Emergence, development and diversification of the TGF-beta signalling pathway within the animal kingdom. **BMC Evol Biol**, 9, p. 28, 2009.

LETSOU, A.; ARORA, K.; WRANA, J. L.; SIMIN, K. *et al.* Drosophila Dpp signaling is mediated by the punt gene product: a dual ligand-binding type II receptor of the TGF beta receptor family. **Cell**, 80, n. 6, p. 899-908, Mar 1995.

LIU, C.; XU, P.; LAMOUILLE, S.; XU, J. *et al.* TACE-mediated ectodomain shedding of the type I TGF-beta receptor downregulates TGF-beta signaling. **Mol Cell**, 35, n. 1, p. 26-36, Jul 10 2009.

MARQUÉS, G.; BAO, H.; HAERRY, T. E.; SHIMELL, M. J. *et al.* The Drosophila BMP type II receptor Wishful Thinking regulates neuromuscular synapse morphology and function. **Neuron**, 33, n. 4, p. 529-543, Feb 2002.

MURPHY, S. J.; DORÉ, J. J.; EDENS, M.; COFFEY, R. J. *et al.* Differential trafficking of transforming growth factor-beta receptors and ligand in polarized epithelial cells. **Mol Biol Cell**, 15, n. 6, p. 2853-2862, Jun 2004.

MURPHY, S. J.; SHAPIRA, K. E.; HENIS, Y. I.; LEOF, E. B. A unique element in the cytoplasmic tail of the type II transforming growth factor-beta receptor controls basolateral delivery. **Mol Biol Cell**, 18, n. 10, p. 3788-3799, Oct 2007.

NELLEN, D.; BURKE, R.; STRUHL, G.; BASLER, K. Direct and long-range action of a DPP morphogen gradient. **Cell**, 85, n. 3, p. 357-368, May 1996.

NORMAN, M.; VUILLEUMIER, R.; SPRINGHORN, A.; GAWLIK, J. *et al.* Pentagone internalises glypicans to fine-tune multiple signalling pathways. **Elife**, 5, 06 2016.

PETERSON, A. J.; JENSEN, P. A.; SHIMELL, M.; STEFANCSIK, R. *et al.* R-smad competition controls activin receptor output in Drosophila. **PLoS One**, 7, n. 5, p. e36548, 2012.

PETERSON, A. J.; O'CONNOR, M. B. Activin receptor inhibition by Smad2 regulates Drosophila wing disc patterning through BMP-response elements. **Development**, 140, n. 3, p. 649-659, Feb 1 2013.

PETERSON, A. J.; O'CONNOR, M. B. Strategies for exploring TGF-beta signaling in Drosophila. **Methods**, 68, n. 1, p. 183-193, Jun 15 2014.

ROY, S.; HUANG, H.; LIU, S.; KORNBERG, T. B. Cytoneme-mediated contact-dependent transport of the Drosophila decapentaplegic signaling protein. **Science**, 343, n. 6173, p. 1244624, Feb 21 2014.

SASAKI, N.; SASAMURA, T.; ISHIKAWA, H. O.; KANAI, M. *et al.* Polarized exocytosis and transcytosis of Notch during its apical localization in Drosophila epithelial cells. **Genes Cells**, 12, n. 1, p. 89-103, Jan 2007.

SMITH, R. B.; MACHAMER, J. B.; KIM, N. C.; HAYS, T. S. *et al.* Relay of retrograde synaptogenic signals through axonal transport of BMP receptors. **J Cell Sci**, 125, n. Pt 16, p. 3752-3764, Aug 2012.

UPADHYAY, A.; MOSS-TAYLOR, L.; KIM, M. J.; GHOSH, A. C. *et al.* TGF-β Family Signaling in. **Cold Spring Harb Perspect Biol**, 9, n. 9, Sep 2017.

VERMEER, P. D.; EINWALTER, L. A.; MONINGER, T. O.; ROKHLINA, T. *et al.* Segregation of receptor and ligand regulates activation of epithelial growth factor receptor. **Nature**, 422, n. 6929, p. 322-326, Mar 2003.

WRANA, J. L.; TRAN, H.; ATTISANO, L.; ARORA, K. *et al.* Two distinct transmembrane serine/threonine kinases from Drosophila melanogaster form an activin receptor complex. **Mol Cell Biol**, 14, n. 2, p. 944-950, Feb 1994.

YIN, X.; KANG, J. H.; ANDRIANIFAHANANA, M.; WANG, Y. *et al.* Basolateral delivery of the type I transforming growth factor beta receptor is mediated by a dominant-acting cytoplasmic motif. **Mol Biol Cell**, 28, n. 20, p. 2701-2711, Oct 2017.

YIN, X.; MURPHY, S. J.; WILKES, M. C.; JI, Y. *et al.* Retromer maintains basolateral distribution of the type II TGF-β receptor via the recycling endosome. **Mol Biol Cell**, 24, n. 14, p. 2285-2298, Jul 2013.

Figure Legends.

Figure 1. **Punt exhibits basolateral membrane targeting in wing disc and MDCK cell epithelia**.
(A,B) GFP signal reveals the membrane distribution of Type II receptors relative to junctions. Punt-
GFP is largely excluded from the septate junction and sub-apical regions but detected at basolateral
surfaces (A' and A"). Wit-GFP is enriched at the sub-apical membrane but detected basolaterally (B'
and B") Single prime (') images are magnified views where two portions of the epithelium lean
towards each other. Double prime (") images are intensity profiles perpendicular to the line demarcated
by the junction staining. (C-F) Chimeric proteins of Punt and Wit map the basolateral targeting activity
to the cytoplasmic domain of Punt. Tagged Wit (WWF) staining overlapped with the apical marker
(D), as did the Punt:Wit PW chimera (E). The Wit:Punt WP chimera displayed basolateral distribution
(F).   (G-K) Fly Punt expressed in polarized mammalian MDCK cells displays basolateral localization
but does not require the same region as TβRII. A schematic showing the position of Punt relative to the
plasma membrane (G). The N-terminal portion is ectoplamic and the C-terminal portion is
cytoplasmic. The corresponding position of the LTA motif is shown in red. (H-K) xz projections from
confocal images of intact epithelium transiently expressing HA-Punt constructs, with apical to the top
of each image. Nuclei are stained with DAPI (blue) and surface receptors were stained with anti-HA
(red). Full-length Punt and variants harboring deletions of extended segments of the cytoplasmic
domain are all detected at the basolateral membrane but excluded from the apical membrane. See
Figure 1-S2 for additional deletion constructs that retain basolateral distribution.

Figure 2. **The cytoplasmic juxtamembrane region of Punt harbors the BLT basolateral targeting
motif**. (A) Amino acid sequence alignment of several insect Punt homologs reveals a core conserved
sequence. The corresponding region of Wit does not resemble the Punt sequence. (B-D) Deletion of the
BLT abolishes the basolateral restriction of Punt proteins expressed in wing disc epithelia. The red
arrow in B points to the furthest apical reach of HA-Punt, which is basolateral to the septate junction
(B). Yellow arrows indicate ectopic staining near (C) or apical to (D) the septate junction. (E-H)
Deletion of the BLT also abolishes basolateral restriction in MDCK cells. HA-Punt-PstI was detected
in mainly in the lateral membrane region (F), whereas Punt-Δ10 and Punt-Δ19 displayed robust apical
staining (level marked by red arrows) in addition to basolateral staining (F, G). (I-K) BLT function is
conserved in insect Punt proteins. A9-Gal4 driving the control UAS-Punt[Punt] protein revealed the
absence of staining at or apical to the junctions (I). The corresponding sequence from honeybee also
supported basolateral restriction of the Punt[Apis] proteing (J). The corresponding Wit sequence
instead of Punt failed to provide basolateral targeting activity, with the Punt[Wit] protein detected in a
pattern resembling Wit-GFP (K). Single prime (') images are magnified views of the rectangles
depicted in the whole wing disc image. (L-N) Mutation of the BLT leads to varied apicalization
depending on the severity of the mutation. Representative staining and signal profiles for mutant
versions of Punt with increasing number of changed amino acids expressed with A9-Gal4 are shown.
The Punt-PTE variant showed a lack of staining in the apical domain, similar to WT Punt (L and L').
Mutation of the insect-conserved residues to Alanines led to moderate and varied overlap with the
aPKC apical marker (M and M'). Mutation of the entire 19 AA of the BLT abolished basolateral
enrichment as shown by the overlap of Punt staining with Dlg and aPKC (N and N'). (O-Q) Moving
the BLT to the juxtamembrane region of Wit causes basolateral restriction. Flag IHC gives background
signal enriched at the apical membrane (O). The control Wit[Wit] protein driven by A9-Gal4 shows
enrichment at the apical membrane (P). Wit[Punt] displays primarily basolateral distribution with
many apical regions showing staining at background levels (Q). Double prime (") images are Fire
false-color displays of the anti-Flag channel.

Figure 3. **Basolateral targeting of Punt is required for animal fitness and normal Dpp signaling**.
(A) *wit* mutants are rescued to viability by all variants of Wit and Punt proteins. Column labels
indicate the rescue construct being driven by elav-Gal4. N=none, WW = Wit[Wit], WA=Wit[Apis],
WP=With[Punt], PP =Punt[Punt] PA=Punt[Apis] and PW= Punt[Wit], with temperature indicated by
21 or 25, and color coded. (B-C) BLT-containing Punt proteins caused more ectopic lethality than
apicalized Punt. *en*-GAL4 driving UAS-Punt[BLT] proteins revealed differential temperature
sensitivity for lethality (B). With A9-GAL4 gender differences in survival at different temperatures
indicate differential activity of over-expressed Punt[BLT] variants (C). (D-I) Genomic rescue
constructs link viability to presence of the BLT. Punt-GFP staining shows basolateral restriction (D).
The EG point mutant displayed mild apicalization (E) whereas the Δ19 construct presented a stronger
apical signal (F).   PuntWT and EG constructs provide sufficient activity to support survival above
Normalized Viability predicted ratios, but the Δ19 construct supported only limited survival (G). Lethal
phase observations showed that Punt-GFPΔ19 animals exhibited increased pupal lethality (H). WT and
EG rescue constructs supported female fertility, but the Δ19 construct did not (I). (J-N) Ectopic activity
of over-expressed Punt is diminished upon apicalization. p-Mad pattern in the wing disc for control (J)
and *en*>Punt[BLT] proteins (K-M). The ratio of p-Mad staining from the posterior compartment (en-
positive) versus anterior compartment increased for each Punt[BLT] protein, but less so for Punt[Wit]
(N). (Q-V) Apicalized Punt did not support proper Dpp signaling or wing development.
Representative wing disc P-Mad staining from larvae rescued by the indicated construct (O-R). Δ19
animals showed a variable morphology, with 4 of 11 discs having an abnormal furrow along the A/P
bounday (R). Wings from adults from rescued genotypes (S-V). Δ19 wings (U-V) showed variability
in vein patterns and overall morphology.

Figure 4. **Membrane distribution varies by tissue, and requires sorting machinery but not
endocytosis in the wing disc**. (A-B) Punt-GFP and Wit-GFP displayed basolateral restriction in the
larval salivary gland epithelium. Red arrows in point to gap in GFP signal at the septate junction (A',
B'). (C-D) Punt-GFP and Wit-GFP showed general membrane presentation in the follicle cell
epithelium in the ovary. Green arrows point to clear GFP signal at the apical surface (C', D'). (E-H)
Screening for apicalization of Punt-GFP in RNAi clones for trafficking factors. Clones are marked in
red in whole disc confocal image, and double-prime insets. GFP signal is shown in Fire false-color
scale in single-prime insets. Green arrows in the red channel point to apical membrane regions near
RNAi clones whereas white arrows point to control no-RNAi regions. Green arrows in the GFP
channel indicate ectopic apical signal and white asterisks show the background level of apical signal.
Punt-GFP is observed at the apical membrane near marked clones with RNAi knockdown of AP-1/2β
(E) or AP1-γ (F). No such apical staining was observed in clones with RNAi knockdown for Rab5 (G)
or AP-2 components such as AP-2μ (H). (I)-J Western blot detection of Punt[Punt] and Punt[Wit].
Samples were collected from nub-GAL4 driving the indicated Punt construct. 'ant' is anterior half of
the larvae, including discs, and 'WD' is specific dissection of Wing Discs. The HA epitope near the N-
terminus did not detect significant specific degradation products (I). Detection of the Punt antibody
epitope near C-terminus also did not reveal any significant specific bands that would suggest clipping
(J).   Mobility markers are indicated by kDa labels, arrows indicate expected position of Punt, and
asterisks mark background bands not attributed to specific Punt detection. Punt[Wit] level was not
increased with either antibody (I,J).

A

```
Fruit fly Punt:     [ECD][TM]RRRKQAHFNEIPTHEAITNS...[KD]
Mosquito Punt:              ...KQAMFNEIPTVE...
Honey bee Punt:             ...KSGYFNEVPTLE...
Beetle Punt:                ...KF-YFNELPTSE...
Punt consensus:             KxxxFNEφPTxE

fruit fly Wit:      [ECD][TM]RTAKEKPEPEESPLAPSGPGYSS...[KD]
```

vg-GAL4>   punt(PstI pseudo)   punt-Δ10   punt-Δ19

anti-HA (ECD)

B — FasIII / HA-punt

C — FasIII / HA-punt

D — FasIII / HA-punt

E

FL-Punt PstI — 1, 145, 175, 516, HA, PstI

Punt Δ10aa — 1, 145, 185, 516, HA, Δ10aa

Punt Δ19aa — 1, 145, 194, 516, HA, Δ19aa

F

G

H

I — Punt[Punt] — Dlg / aPKC   I'

J — Punt[apis] — Dlg / aPKC   J'

K — Punt[Wit] — Dlg / aPKC   K'

L — Punt-PTE — Anti-Punt   L'

M — Punt-Conserved-Ala — Anti-Punt   M'

N — Punt-19-Ala — Anti-Punt   N'

O — Background — Apical (aPKC) / Anti-Flag   O' — O'' Anti-Flag

P — Wit[Wit] — Apical (aPKC) / Anti-Flag   P' — P''

Q — Wit[Punt] — Apical (aPKC) / Anti-Flag   Q' — Q''

Larval Salivary Gland

A Punt-GFP — DAPI / GFP / FASIII / aPKC
B Wit-GFP
A' Punt-GFP — DAPI / GFP / aPKC
B' Wit-GFP

Follicle Cells of Egg Chamber

C Punt-GFP — aPKC
D Wit-GFP — aPKC
C' Punt-GFP
D' Wit-GFP

E AP-1/2 β RNAi — Punt-GFP
E' GFP/Fire
E''
F AP-1γ RNAi — Punt-GFP
F'
F''

G Rab5 RNAi — Punt-GFP
G'
G''
H AP-2μ RNAi — Punt-GFP
H'
H''

I anti-HA (N-terminal)
1 2 3 4 kDa
100 75 50 37 25
ant — none
ant — Punt[Punt]
ant — Punt[Wit]
WD — Punt[Punt]

J anti-Punt (C-terminal)
1 2 3 4
ant — none
ant — Punt[Punt]
ant — Punt[Wit]
WD — Punt[Punt]