Taxonomic hypotheses and the biogeography of speciation in the Tiger Whiptail complex (*Aspidoscelis tigris*: Squamata, Teiidae)

Tyler K. Chafin[1], Marlis R. Douglas[1], Whitney J.B. Anthonysamy[1,2], Brian K. Sullivan[3], James M. Walker[1], James E. Cordes[4], and Michael E. Douglas[1]

[1] Department of Biological Sciences, University of Arkansas, Fayetteville, Arkansas 72701, USA; Email: (TKC) tkchafin@uark.edu, send reprint requests to this address; (MRD) mrd1@uark.edu; (JMW) jmwalker@uark.edu; and (MED) med1@uark.edu.

[2] Present address: St. Louis College of Pharmacy, 4588 Parkview Place, St. Louis, Missouri 63110, USA; Email: whitney.anthonysamy@stlcop.edu.

[3] School of Mathematical and Natural Sciences, P. O. Box 37100, Arizona State University, Phoenix, Arizona 85069, USA; Email: bsullivan@asu.edu.

[4] Division of Mathematics and Sciences, Louisiana State University, Eunice, Louisiana 70535, USA; Email: jcordes1@live.com.

Running head: *Aspidoscelis tigris* phylogeography

Keywords: whiptail lizards, climate change; mtDNA; southwestern United States; species concept; subspecies

(225)Biodiversity in southwestern North America has a complex biogeographic history involving

2   tectonism interspersed with climatic fluctuations. This yields a contemporary pattern replete with

historic idiosyncrasies often difficult to interpret when viewed from through the lens of modern

4   ecology. The *Aspidoscelis tigris* (Tiger Whiptail) complex (Squamata: Teiidae) is one such

group in which potential taxonomic boundaries have been confounded by a series of complex

6   biogeographic processes that have defined the evolution of the clade. To clarify this situation, we

first generated multiple taxonomic hypotheses, which were subsequently tested using

8   mitochondrial DNA sequences (ATPase 8 and 6) evaluated across 239 individuals representing

five continental members of this complex. To do so, we evaluated the manner by which our

10   models parsed phylogenetic and biogeographic patterns. We found considerable variation among

species 'hypotheses', which we suggest in part reflects inflated levels of inter-population genetic

12   divergence caused by historical demographic expansion and contraction cycles. Inter-specific

boundaries with *A. marmoratus* juxtaposed topographically with the Cochise Filter Barrier that

14   separates Sonoran and Chihuahuan deserts (interpreted herein as case of 'soft' allopatry).

Patterns of genetic divergence were consistent across the Cochise Filter Barrier, regardless of

16   sample proximity. Surprisingly, this also held true for intraspecific comparisons that spanned the

Colorado River. These in turn suggest geomorphic processes as a driver of speciation in the *tigris*

18   complex, with intraspecific units governed locally by demographic processes.

HIGHLIGHTS(185)

1. Phylogeographies of vertebrates within the southwestern deserts of North America have been shaped by climatic fluctuations imbedded within broad geomorphic processes.

2. The resulting synergism drives evolutionary processes, such as an expansion of within-species genetic divergence over time. Taxonomic inflation often results (i.e., an increase in recognized taxa due to arbitrary delineations), such as when morphological divergences fail to juxtapose with biogeographic hypotheses.

3. However, isolated groups can be discriminated within-species by mapping genetic variability onto geographic distances. This approach can often diagnose 'hard' barriers to dispersal, or alternatively, strong selection acting against hybridization. On the other hand, elevated genetic divergences among groups less-isolated would underscore isolation-by-distance (i.e., an increase in genetic differentiation concomitant with geographic distance).

4. The biogeographic patterns we identified in Tiger Whiptail are largely synonymous with those found in other regional species, particularly given the geomorphic separation of Mohave and Sonoran deserts by the Colorado River, and Sonoran/ Chihuahuan deserts by the Cochise Filter Barrier.

5. Our results for the Tiger Whiptail complex broaden and extend the context within which polytypic species are conserved and managed, particularly those that reflect an incongruence among molecular and morphological standards.

INTRODUCTION

Whiptail lizards (*Aspidoscelis*; Teiidae) comprise one of the most conspicuous groups of lizards in western North America. Their complex history of hybridization/ introgression coupled

with multiple transitions to parthenogenic life histories fuels their extraordinary capacity as an evolutionary model (Barley et al. 2019). However, intraspecific morphological variation coupled

with high rates of hybridization has made species-delineation difficult, which in turn has muddied their taxonomy. The occurrence of paraphyly in the group was partially resolved by

subsequently partitioning species between a resurrected *Aspidoscelis* and the nominate *Cnemidophorus* (Reeder et al., 2002). While a positive step, it also necessitated reallocation of

several groups, such as: *Aspidoscelis cozumelus* (parthenogenetic species), *A. deppi* (gonochoristic species), *A. sexlineatus* (parthenogenetic and gonochoristic species), *A.*

tesselatus (parthenogenetic species), and *A. tigris* (gonochoristic species). As a result, many systematic issues inherent within *Cnemidophorus* were simply transferred to *Aspidoscelis*.

The *A. tigris* species-group, for example, represents a perplexing systematic challenge as it comprises a widely distributed complex of gonochoristic populations located within desert

regions of southwestern North America, northern México (to include Baja California), as well as numerous islands east and west of the peninsula. Although the group *per se* is characterized by a

readily assessed set of morphological characters (Walker and Maslin, 1981) and a distinctive karyotype (Lowe et al., 1970), its diversity (i.e., ranging from a single species to more than a

dozen) has been a contentious issue in North American Herpetology. Much of its diversity is represented by distinct populations located on relatively small, vulnerable islands in the Gulf of

California. Several of these were designated as distinct species (Walker et al., 1966a, b; Walker and Taylor, 1968; Walker and Maslin, 1981), based snout-vent length, color pattern, and/or

morphology (e.g., *A. bacatus* of Isla San Pedro Nolasco; *A. catalinensis* of Isla Santa Catalina; *A.*

martyris of Isla San Pedro Martír; and *A. celeripes* of Isla San Jose). However, some authors

68   (e.g. Lowe et al., 1970; Wright, 1993) have considered them instead as subspecific variants.

Continental populations have also proven taxonomically contentious. For example, a long-

70   standing disagreement (Hendricks and Dixon, 1986; Dessauer et al., 2000) centers on whether *A.*

tigris and *A. marmoratus* in small areas in Hidalgo County NM represent intergrading subspecies

72   or hybridizing species.

Taxonomic difficulties are thus quite apparent within the *A. tigris* complex, and primarily

74   stem from complex patterns of intergradation and hybridization confounded by non-

discriminating morphological variation. To provide clarification, we evaluated a complex of four

76   continental subspecies (*A. t. mundus*, *A. t. punctilinealis*, *A. t. septentrionalis*, and *A. t. tigris*;

Fig. 1), plus a recently elevated putative sister species, *A. marmoratus* (Hendricks and Dickson,

78   1986; Reeder et al., 2002; Tucker et al., 2016). Our goal was to examine how mitochondrial

(mt)DNA variation in *A. tigris* subspecies compares with morphological discordance [e.g.,

80   subspecies seemingly supported by color pattern yet weakly differentiated by scalation and

morphometrics (Taylor et al., 1994; Walker et al., 2015; Fig. 2)]. Furthermore, morphological

82   intergrades and overlapping genetic polymorphisms suggest admixture among several groups: *A.*

t. tigris and *A. t. septentrionalis* (Taylor 1988), *A. t. tigris* and *A. t. punctilinealis* (Taylor 1990),

84   and *A. t. punctilinealis* and *A. t. septentrionalis* (Dessauer et al. 2000).

The contact zone separating *A. t. punctilinealis* and *A. marmoratus* (i.e., *A. t. marmoratus*

86   of the following authors) coincides with a genetic cline at multiple loci (Dessauer and Cole 1991,

Dessauer et al. 2000), suggesting secondary contact following a period of isolation (Barton and

88   Hewitt, 1985). A steep cline involving six color-pattern characteristics is also apparent in the

hybrid zone between *A. t. punctilinealis* and *A. marmoratus* (Dessauer et al., 2000), arguing for

90 the utility of these characters for delineation of *A. tigris* lineages.

In the present study, we focused on defining phylogeographic patterns and interpreting

92 phylogenetic delineations primarily within the range of *A. t. punctilinealis* in Arizona. To the

east, it forms steep morphological/ genetic clines at a hybrid zone with *A. marmoratus* (Dessauer

94 et al., 2000), whereas to the west it exhibits a relatively gradual phenotypic gradient with *A. t.*

tigris (Taylor 1990). Interestingly, both *A. t. septentrionalis* and *A. t. punctilinealis* also form

96 exceptionally gradual morphoclines with *A. t. tigris* that lack apparent ecological underpinnings

(Taylor et al. 1994, Taylor 1990). This stands in contrast to that found between *A. t.*

98 *punctilinealis* and *A. marmoratus* (<2km; Dessauer et al., 2000; Zweifel 1962). Given this

discontinuity, we also examined the association between morphological gradation in the complex

100 and its genetic discontinuity. To do so, we extended the phylogenetic resolution of lineages

within the delineated *A. tigris* complex, then contrasted our phylogenetic patterns with

102 morphological species identifications. We generated taxonomic hypotheses by employing

'species delimitation' algorithms, then employed subsequent downstream analyses to compare

104 the manner by which each model partitioned genetic variation.

106 *The Colorado Plateau as a biogeographic entity*

The Colorado Plateau is a tectonically intact landform (~2000m elevation) with scant

108 internal deformation, thus providing a relatively linear configuration of layered sedimentary rock

for the Plateau interior that contrasts with an eroding margin since Late Miocene (Levander et

110 al., 2011). It also stands in sharp contrast with the more fractured geomorphic features of

neighboring regions: The Rocky Mountains (immediately north and east), Rio Grande Uplift

112   (southeast), and the Basin and Range Province (immediately west) (Fig. 1 of Flowers, 2010). The

uplift of the Plateau and the integration of the Colorado River were defining geomorphic events

114   (Minckley et al., 1986; Spencer et al., 2008).

Prior to integration, the Colorado River emptied into an endorheic basin, most likely

116   following the early path of a major modern tributary (Little Colorado River; House et al., 2008).

In late Miocene-early Pliocene, headwater erosion and basin 'spillover' forged a connection

118   between the river and the Gulf of California via the Salton Trough (Dorsey et al., 2018;

Nicholson et al., 2019). The Gila River system, which had previously drained directly into the

120   Gulf (Eberly & Stanley, 1978) was similarly incorporated. Other drainages that terminated into

closed basins during the Miocene (i.e. Salt and Verde rivers) were also integrated with the Gila

122   River system by late Pliocene (Blakely & Ranney, 2008).

Thus, an historic perspective is necessary context for deciphering phylogeographic

124   patterns of biodiversity on the Colorado Plateau and surrounding regions. This is because the

North American southwest was essentially defined by tectonism and subsequent climatic

126   oscillations, with phylogeographic/ phylogenetic reverberations manifest within the endemic

biodiversity (Douglas et al., 2016; Holycross & Douglas, 2007; Mussmann et al., 2020; Sullivan

128   et al., 2014). Ancillary genetic footprints of hybridization, diversification, and rapid demographic

change have also become increasingly apparent in modern evaluations (Chafin et al., 2020;

130   Bangs et al., 2020a; Bangs et al., 2020b; Douglas et al., 2006).

Opportunities for biotic exchange (or alternatively, vicariant events) hinged largely upon

132   drainage evolution (e.g., Chafin et al., 2020; Douglas et al., 1999). Small-bodied terrestrial

vertebrates were also involved in these biogeograpnic opportunities, despite the fact that the

Colorado River had long been hypothesized as a vicariant barrier to dispersal (e.g. Lamb et al., 1992; Pounds and Jackson, 1981).

Geomorphic events actively modulated geographic boundaries and were thus pivotal with regard to biological diversification on the Plateau (Bangs et al., 2020a, 2020b; Douglas et al.,

1999; Douglas et al., 2016). This process acted in synergy with widespread physiographic and climatic alterations to drive high levels of endemism. The Miocene saw contraction of the Basin

and Range province, as well as climatic oscillation creating 'boom-and-bust' demographic cycles, as Plateau habitats expanded and contracted (Douglas et al., 2006). The Early and Middle

Pliocene brought decreased seasonality as well as elevated temperatures and humidity that were subsequently reversed by Plio-Pleistocene glaciation (Thompson, 1991).

Quaternary climate effects such as glaciation were limited primarily to higher elevations (Douglas et al., 2006), whereas average temperatures within the continental interior declined

substantially due to suppression of the western North American monsoon (Bhattacharya et al., 2018). Desert vegetation, initially restricted to lower elevations (<300m), soon expanded with

Holocene drought, such that demographic signatures were recorded as genetic bookmarks within desert herpetofauna (Douglas et al., 2006; Sullivan et al., 2014). The extent of desert habitat was

modulated by fluctuating interglacials (e.g., Cole, 1986; Douglas et al., 2009) that also re-organized stream diversions as potential dispersal barriers. The genetic legacy of these climatic

fluctuations is recorded at both continental and global scales (e.g., Brunner et al., 2001; Bryson et al., 2011; Hewitt, 2000; Erickson et al., 2019; Kohli et al., 2015; Migliore et al., 2019;

Morales-Barbero et al., 2018).

MATERIALS AND METHODS

158   *Sampling, amplifying and sequencing of mtDNA*

We evaluated a total of 239 samples representing five members of the *A. tigris* complex

160   (*A. marmoratus*, *A. tigris mundus*, *A. t. punctilinealis*, *A. t. septentrionalis*, and *A. t. tigris*), their

putative secondary intergrades (e.g., hybrids), plus two out-groups (*Tupinambis rufus* and

162   *Aspidoscelis gularis*) (Fig. 1). Genomic DNA was isolated from tail clippings or liver tissue

using the Chelex Instagene-matrix extraction protocol (Bio-Rad).

164   We utilized published primers and PCR protocols (Douglas et al., 2002) to amplify two

overlapping mtDNA genes (ATP8 and ATP6). Poor amplification in *A. marmoratus* also forced

166   the design of species-specific primers (PRIMER3; Untergasser et al., 2012). These were: ATP_F

(3-CGCACCCTGATTTATTGTTT-5) and ATP_R (3-GGGTGAAAACGTATGCTTGA-5).

168   Each 20 µl PCR reaction consisted of 1.5 µl of 1/10x DNA template, 10 µl of HotStarTaq Master

Mix (QIAGEN), 0.08 µM of each primer, with the reaction volume finalized with distilled water.

170   After verification via agarose gel electrophoresis, amplicons were purified using an Exonuclease

I/Shrimp Alkaline Phosphatase. Forward and reverse sequencing reactions were performed using

172   BigDye v3.1 Dye Terminator chemistry (Applied Biosystems, Inc.) and sequenced at the

University of Illinois Roy J. Carver Biotechnology Center. Raw sequence chromatograms were

174   manually curated (SEQUENCHER ver. 5.0, Gene Codes, Ann Arbor, MI) and aligned (MEGA_5;

Tamura et al., 2011).

Phylogenetic inference

178   To develop initial hypotheses, we first inferred phylogenetic relationships using both

Bayesian and distance-based methods. We tested models of nucleotide substitution (N=88-

180　JMODELTEST; Darriba et al., 2012), using Akaike and Bayesian information criteria (AIC and

BIC, respectively). We generated an ultrametric tree using the HKY-model (selected by both

182　AIC and BIC) with gamma distributed rates and invariant sites (BEAST2; Bouckaert et al.,

2014). We applied a Yule model with default parameters and a relaxed log-normal clock process.

184　MCMC chains were sampled every 500th of 10 million MCMC iterations, following a burn-in

period of 2 million. Chain convergence and mixing were visually assessed, and MCMC samples

186　thinned so as to maintain effective sample sizes >200 while reducing autocorrelation of samples

(TRACER; Rambaut and Drummond, 2007). The posterior sampled trees were summarized as a

188　maximum clade credibility tree with branch lengths representing median heights. For contrast,

we also generated a neighbor-joining (NJ) tree using model-corrected distances (R-package APE;

190　Paradis et al., 2004) in R (R Core Team, 2020), with nodal support assessed via 100 bootstrap

pseudo-replicates, as well as a maximum-likelihood tree in PHYML (Guindon et al. 2010). Basic

192　population summary statistics for major phylogenetic clades (as diagnosed via nodal support and

statistical tests of monophyly) were also computed in PEGAS (Paradis, 2010).

Delimiting groups within A. tigris

196　　　We first evaluated the currently accepted taxonomic designations for subspecies within *A.*

tigris (e.g., Crother et al., 2017; hereby abbreviated as 'TAX'), then employed an array of

198　analytical approaches to generate six additional delimitation hypotheses. Distance-based

clustering methods, used in the context of species delimitation, assume that biological units (i.e.,

200　species) form non-overlapping genetic clusters. We first employed a recursive method for

'Automated Barcode Gap Discovery' (ABGD; Puillandre et al., 2012) that statistically infers a

202　threshold distance from the data (i.e., the genetic distance at which groups are considered to

represent different clusters). We also generated clusters using corrected genetic distances as input

204 (CD-HIT; Fu et al., 2012), with manually specified distance thresholds of 2% and 5% (hereafter

referred to as CD2 and CD5). These hard thresholds were chosen because they represent

206 approximate 'breakpoints' at which changes occur in the ABGD analysis regarding the number

of sampled groups.

208   We defined phylogenetic species using several approaches: The first, a 'naïve' method,

defined groups as the shallowest divergences statistically supported and reciprocally

210 monophyletic (Rosenberg, 2007), hereafter abbreviated as 'ROS' (SPIDER R package; Brown et

al., 2012). A second approach derived groups from tree characteristics under an assumed

212 speciation process. The first of these methods ['Generalized Mixed Yule Coalescent' (GMYC;

Pons et al., 2006)] represents the species boundary as a distinguishable change in branching

214 patterns within the phylogeny, with independently evolving clusters diagnosed as those at which

branching patterns transition from expected under two different models: An assumed

216 interspecific (Yule process; Nee et al., 1994) and an intraspecific (neutral coalescent; Hudson,

1990). In the second ['Poisson Tree Process' (PTP; Zhang et al., 2013)] the probability of

218 speciation is considered in terms of mutational distance between cladogenesis events (as

assumed with a Poisson distribution). PTP requires a tree with branch lengths scaled per

220 substitutions, and given this, our input represented a tree topologically consistent with that

inferred by BEAST but derived instead with PHYML (Guindon et al. 2010). GMYC was run

222 using the Splits R package (Fujisawa et al., 2013); PTP using the Bayesian implementation

provided by the bPTP server (Zhang et al., 2013).

Comparingf delimitations

226   We used 2-way and 3-way Analyses of Molecular Variance (AMOVA; Excoffier et al.

1992) with group assignments defined by individual membership under each hypothesis. We

228 then used isolation-by-distance (IBD) as a null model under the assumption that the entire

complex represented a single species. Our justification stemmed from the pervasive nature of

230 IBD with regard to dispersal-limited species (Sexton et al. 2014). In this sense, deviations from

the expected IBD pattern are hypothesized as diagnostic of some alternative non-spatial process

232 (e.g. reproductive isolation). Partial Mantel tests (Smouse et al. 1986), employing pairwise

genetic distances with geographic distances as a covariate (Douglas and Endler, 1982; Meirmans,

234 2012), tested the fit of individual assignments under each delimitation hypothesis. Simple and

partial Mantel tests (R-package VEGAN; Oksanen et al., 2007) utilized corrected genetic distance

236 (APE; Paradis et al., 2004) with confidence assessed using 9,999 permutations. We also employed

a stratified Mantel test (per Meirmans, 2012) with permutations constrained within clusters, to

238 test if significance of the IBD model is impacted by sampling across structured populations, as

delimited by each of our hypotheses.

240   We also used a spatially explicit partitioning analysis with genetic variance [measured as

a fixation index (F_{CT})] maximized among groups by iteratively rearranging samples into K-

242 clusters [Spatial Analysis of Molecular Variance (SAMOVA); Dupanloup et al., 2008]. These

were executed for K-values 2-20, using 100 bootstrapped alignments and replicated without

244 explicit geographic constraints. We also identified spatial transitions in genetic distances using

an interpolation approach [ALLELES-IN-SPACE (AIS); Miller, 2005].

RESULTS

248 *Phylogenetic and spatial relationships within* Aspidoscelis tigris

The final 777bp alignment resulted in 120 haplotypes. Bayesian analysis (BEAST2)

250   yielded strong support for *A. t. tigris* + *A. t. punctilinealis* (ROS clades A–F, Fig. 3; also see

Supplementary Figure S1) as being sister to *A. t. mundus* + *A. t. septentrionalis* (ROS clade G).

252   All clades were significantly supported as being reciprocally monophyletic [Rosenberg (2007)

test], and all reflected similarly high bootstrap support in the neighbor-joining analysis. The

254   maximum-likelihood analysis resulted in a tree topologically identical, and was thus not

displayed.

256   All four subspecies formed a monophyletic group sister to a well-defined *A. marmoratus*

(clade H). *Aspidoscelis t. septentrionalis* (clade G, excluding *mundus* samples) demonstrates

258   significantly negative values for both Tajima's and Fu and Li's *D* statistics, with exceptionally

low nucleotide diversity and haplotype diversity much less than that recorded for *A. t.*

260   *punctilinealis* and *A. t. tigris* (Table 1). These suggest a population expansion following a

bottleneck (Douglas et al., 2010). *Aspidoscelis marmoratus* showed similar signatures, although

262   with a much greater differential between haplotype and nucleotide diversities, and a reduced

number of segregating sites relative to sample size (Table 1). Together, *A. t. tigris* + *A. t.*

264   *punctilinealis* were subdivided into five major clades with strong nodal support and reciprocal

monophyly (clades A–F, Fig. 3).

266   Morphologically-diagnosed *A. t. tigris* was polyphyletic, with all samples but one

allocated to either clade A or F. The single exception was a morphologically diagnosed *A. t.*

268   *tigris* (Nevada) that fell within the *A. t. septentrionalis*/*mundus* haplogroup (clade G). Clade F,

which was >2% divergent from the remaining *A. t. tigris*/*punctilinealis* clades (A–E), contained

270   both morphologically consistent *A. t. tigris* (Clark County, southern Nevada), as well as a small

subset of *A. t. punctilinealis* (Mohave County, northwest Arizona). All other morphologically

272   identified *A. t. tigris* from California had mitochondrial haplotypes aligning with clade A of *A. t.*

punctilinealis. This clade formed a monophyletic grouping in the BA analysis with samples from

274   southeastern Arizona (clades D and E). Spatially, clades D+E are separated by the mainstem Gila

River and defined to the east by the Cochise Filter Barrier that separates Sonoran and

276   Chihuahuan deserts (Castoe et al. 2007).

        All clades within *A. t. tigris/punctilinealis* generally showed higher numbers of

278   segregating sites relative to nucleotide diversity (negative values for Tajima's *D*; clades A–E

Table 1). Clades A–D also showed a high rate of singleton haplotypes. Although non-significant

280   (Table 1), they suggest past population expansions. Within the remainder of the *A. t.*

punctiliealis, clade A (southwestern Arizona) is sister to a monophyletic group composed of

282   clade B (Grand Canyon and Colorado River mainstem at juncture between Arizona, Nevada,

California), and clade C (central Basin and Range physiographic region of Arizona). The latter is

284   bounded on the north by the Mogollon Rim and is separated from clade B to the south by the

Gila River.

286   The appearance of clade C haplotypes south of the Gila River mainstem indicates limited

gene flow between the two groups, possibly reflecting trans-riverine dispersal as promoted by

288   low seasonal flows. Similarly, clade B haplotypes are found east and west of the Colorado River

mainstem, again suggesting a riverine barrier historically permeable to dispersal, or with

290   mitochondrial exchange facilitated by shifts in flow. However, no apparent geographic features

support the phylogenetic separation of clades A and C. Their sister relationship is also sustained

292   in the NJ analysis, although with a bootstrap support of 56% (compared to 0.87 posterior

probability in the Bayesian phylogeny).

Species delimitation models

296   We found considerable variation in the number of proposed clades generated by species

delimitation models. Fig. 3B depicts the manner by which each model maps onto the phylogeny,

298   using an identification scheme consistent with Fig. 4 and Supplementary Figure S1.

Phylogenetically defined clades ('ROS') and a taxonomic hypothesis ('TAX') both collapsed *A.*

300   *t. tigris* and *A. t. punctilinealis* on the basis of extensive intergradation (Taylor et al. 1994).

        The ABGD analysis was by far the most conservative, collapsing both *A. t. punctilinealis*

302   and *A. t. tigris* with *A. t. septentrionalis* + *A. t. mundus*, while only segregating *A. marmoratus*

(Fig. 3B). CD-HIT, with an *a priori* sequence divergence threshold of 5%, produced a 4-species

304   model (CD5) that separated *A. t. septentrionalis* + *A. t. mundus*, *A. marmoratus*, and *A. t.*

punctilinealis + *A. t. tigris*. The latter was also partitioned into two groups (i.e., separating clade

306   F). One consisted of haplotypes from northwestern Arizona and southern Nevada (clade F; a

subset of most morphologically consistent *A. t. tigris*), while the second comprised *A. t.*

308   *punctilinealis* (clades A–E) and remaining *A. t. tigris*, to include those with ranges overlapping

clade F. Decreasing the threshold to 2% (i.e., model CD2) then divided *A. t. punctilinealis* into

310   eight groups, and five representing *A. t. septentrionalis* (i.e., two from California and three

occupying the Colorado Plateau; Fig. 4). GMYC and PTP produced large numbers of groups,

312   with 13 in the former and 16 in the latter, with many consisting of but a single haplotype.

314   *Partitioning of genetic variance*

        The spatial analysis supported models with fewer species. In the bootstrapped datasets,

316   F_{CT} overlapped among *K* values (Fig. 5), with the greatest increase in among-group variance

(ΔF_{CT}) occurring at *K*=3. Individual assignment at *K*=2 corresponded to the ABGD model (two

species: *A. marmoratus* and *A. tigris*), with three groupings in *K*=3: *A. marmoratus*; *A. t. septentrionalis* + *A. t. mundus*; and *A. t. punctilinealis* + *A. t. tigris*.

A significant correlation was found between matrices of pairwise genetic and geographic distances (Mantel test: r=0.576, P=0.0001; Table 2; Fig. 6). Significant associations only

occurred with the ROS, TAX, ABGD and CD5 models when permutations were limited to those strata defined by delimitation models (i.e., stratified Mantel). The partial Mantel tests with the

ABGD model produced the strongest association (r=0.777), with values generally declining as the number of species increased (Table 2). Correlation of group membership with genetic

distance under all models was greater than in the simple Mantel test for all cases, save for the PTP model (r=0.5754), suggesting that all delimitations better encapsulate spatial genetic

variation than one in which all examined samples are collapsed.

        Overall, those models partitioning *A. tigris* subspecies were supported by AMOVA

330   analyses (Supplementary Table S1, S2), with high genetic variance among groups rejecting panmixia. Among-group genetic variance (analogous to F_{CT}) varied between 72–90% and was

332   maximized in models that recognizing greater numbers of partitions (i.e., CD2, GMYC, and PTP), with 89-90% of variance explained. These values generally declined as models became

334   more conservative.

        Interpolating the AIS results with a diffusion model (Fig. 7) produced apparent peaks in

336   relative pairwise genetic diversity at contact zones between *A. t. punctilinealis* (in this case clade D) and *A. marmoratus* (clade H). These occurred in southeastern Arizona near the shared border

338   with New Mexico, and in northwestern Arizona, near the contact zone with *A. t. tigris* clade F.

340   **DISCUSSION**

We used mtDNA sequence data to: Infer phylogenetic and phylogeographic patterns

342   within the *Aspidoscelis tigris* complex; Generate species delimitation hypotheses therein; and

compare model fit using several approaches. We judged how well each model partitioned genetic

344   variance, both hierarchically and spatially. Our results supported *A. marmoratus* as a distinct

species. However, our weak phylogenetic recognition of *A. t. tigris* and *A. t. punctilinealis*, as

346   well as *A. t. mundus* and *A. t. septentrionalis* questions their subspecific designations, and

suggests the need for additional work utilizing larger number of molecular markers to assess

348   potential intergradation. However, pairwise genetic divergences among the *A. t.*

tigris/punctilinealis and *A. t. septentrionalis/mundus* groups, regardless of their geographic

350   proximity (e.g., <10km versus >1000km) may indicate reproductive isolation, again prompting

further study that incorporates nuclear markers.

Biogeography of speciation in Aspidoscelis tigris

354   Deep relationships within the *A. tigris* complex generally supported the current taxonomy

and reflected divisions among biotic regions. Here, we specifically refer to *A. t. septentrionalis*

356   on the Colorado Plateau, *A. marmoratus* within the Chihuahan Desert, *A. t. punctilinealis* in the

Sonoran Desert, and *A. t. tigris* within Mohave and Great Basin deserts. Haplotypes of *A. t.*

358   *mundus,* restricted to the California Central Valley, were inferred as monophyletic but nested

within a larger *A. t. septentrionalis* clade. This highlights the need for increased sampling,

360   particularly for *A. t. mundus* and *A. t. tigris,* whose distributions were sampled less thoroughly.

Topographically, the separation of *A. t. punctilinealis* and *A. marmoratus* aligns with the

362   Cochise Filter Barrier (CFB), a biogeographic entity that has similarly been implicated as

influential in defining phylogeographic relationships among other taxa (e.g., Castoe et al., 2007;

364   O'Connell et al., 2017; Morafka, 1977). Pyron et al. (2015) invoked this barrier as a potential

driver of so-called 'soft allopatry' (Hickerson and Meyer 2008), in which niche differentiation

366   drives speciation *in lieu* of a hard barrier to dispersal. Such barriers can often lead to

permeability at contact zones, given that divergence is maintained by adaptation rather than

368   physical separation. As might be expected, a well-characterized hybrid zone separates *A. t.*

punctilinealis and *A. marmoratus,* replete with steep clines in color polymorphism (Dessauer et

370   al., 2000). We also identified a distinct east-west divide in mitochondrial haplotypes that is

concordant with an hypothesis of soft allopatry.

The potential for hybridization

374   We found phylogenetic subdivision less clear among subspecies of *A. tigris* and surmise

this likely reflects a cumulative effect of contemporary physiography coupled with historic

376   processes. *Aspidoscelis t. tigris* and *A. t. punctilinealis* were not distinct. Both transverse the

Colorado River, and neither is strictly defined by the Mohave-Sonoran transition. This is perhaps

378   not surprising, given that morphological intergradation between *A. t. tigris* and *A. t.*

punctilinealis does not follow a strict ecological gradient, but instead manifests itself as an

380   exceptionally wide and smooth morphocline (Taylor et al. 1994).

Aspidoscelis t. tigris from the most northerly part of the range had mitochondrial

382   sequences aligning instead with *A. t. septentrionalis*, potentially suggesting exchange between

the two (Taylor, 1988). Here, we emphasize that increased genetic sampling is necessary so as to

384   understand the nature of this putative exchange. In contrast, *A. t. punctilinealis* and *A. t.*

septentrionalis were demarcated according to recognized physiographic provinces, with *A. t.*

386   *septentrionalis* occupying the Colorado Plateau and *A. t. punctilinealis* the Sonoran Desert (Fig.

4, S1). This was expected, particularly given our understanding of their respective ranges (e.g.

388   Fig. 1) and associations with the disparate thorn-scrub communities therein (e.g., Zweifel, 1962;

Pianka, 1970). We suggest this ecological gradient represents a 'soft boundary,' similar to those

390   that mediate the transition between *A. t. punctilinealis* and *A. marmoratus*. Unique communities

characterize both the Colorado Plateau and Sonoran Desert (e.g., Douglas et al., 2002, 2010), and

392   this suggests a common, underlying process driving diversification therein (Riddle and Hafner,

2006; Pianka, 1967).

The validity of taxonomic designations

396   Genetic delineation of *A. t. tigris* was inconsistent with traditional taxonomy, in that

specimens allocated morphologically to *A. t. tigris* were not monophyletic within the

398   phylogenetic analyses, nor were those groupings consistent with the historically accepted range

boundaries for *A. t. tigris* and *A. t. punctilinealis*. Our expectations were that haplotypes sampled

400   from the Great Basin and Mohave deserts (i.e., those regions encompassing *A. t. tigris*) would be

monophyletic. Instead, we found a closer alignment with *A. t. septentrionalis* to the northeast and

402   *A. t. punctilinealis* to the southeast. This is likely a result of mitochondrial exchange among these

subspecies, again highlighting the need for a genomic evaluation.

404   Taylor et al. (1994) demonstrated the existence of a wide, smooth morphocline >180 km

in width that traversed the Colorado River and separated *A. t. tigris* from *A. t. punctilinealis*

406   along the southern extent of their contact (Fig. 1A). This stands in contrast to a 16 km-wide

morphocline found at the northern extent of their contact, and that discrepancy begs an obvious

408   ecological explanation (Taylor 1990). The northern morphocline coincides with the Arizona-

California-Nevada nexus, a hotspot for mitochondrial diversity as well, where sub-clades overlap

410   without clear physical boundaries (Fig. 4, S1). In addition, multiple clades (A and F) traverse the

Colorado River, with the highly divergent clade F (found primarily within morphologically

412   defined *A. t. tigris*) possibly representing a more widespread Mohave clade.

One explanation for the above variation in clinal width evokes an historic perspective

414   wherein the northern contact between *A. t. tigris* and *A. t. punctilinealis* is secondary, whereas

range overlap to the south was maintained. This would generate morphoclines of varying ages

416   and would offer one potential explanation for the variance in mitochondrial divergence (i.e.,

elevated to the north, nonexistent to the south), as well as morphocline width. Here, we posit that

418   the duration and extent of contact may be a driving mechanism. A related hypothesis would be

that the width of the cline is in response to an underlying environmental gradient, such as the

420   transition between Sonoran and Mohave habitats. However, neither hypothesis can be tested with

the data presented herein.

422   An additional possibility is the fluctuating capacity of the Colorado River as a barrier to

dispersal. We offer scant evidence for it serving as a vicariant structure, a result corroborated by

424   studies of other reptiles (e.g., desert tortoise: Lamb et al., 1989; iguanids: Lamb et al., 1992).

Although the Colorado River does seemingly constrain dispersal (Pounds and Jackson 1981),

426   gene flow could still be facilitated by the periodic fluctuations in its trajectory and flow. In fact,

the shallow divergences found among clades transecting the river provide evidence for how

428   recent these events were.

430   *Climate driven population oscillations*

One unexpected finding was the exceptional genetic divergences among *A. t.*

432   *punctilinealis* clades (A–E). One possible explanation evokes fluctuating habitat availability over

time, rather than contemporary features of the landscape. Concordance in relative crown ages

434 among the majority of *A. t. punctilinealis* clades (A–E; Fig. 1B) could suggest that intraspecific

divergence of these groups shares a common cause. Douglas et al. (2006) offered an applicable

436 model that explained genetic structuring and population expansion in four desert rattlesnake

species as a 'boom-and-bust' demographic cycle, caused by historical climate fluctuations.

438 Under this model, a 'refugium phase' (='bust') condensed continuous distributions into smaller,

allopatric populations such that restricted gene flow promoted genetic drift as manifested by

440 reduced effective population size (N_e). As climate shifted, desert habitat expanded such that

ecological barriers separating refugia were removed, allowing distributions to expand (='boom').

442 These oscillating contractions and expansions placed characteristic molecular signatures

within populations, most prominently as a loss of genetic diversity and an exaggerated

444 differentiation among sub-populations post-expansion (Hewitt 2000). This is reflected in an

increased number of low-frequency haplotypes, and low nucleotide diversity among

446 contemporary haplotypes (Table 1). We also noted similar patterns in *A. t. septentrionalis* and *A.

marmoratus*, suggesting bottleneck-expansion events within these taxa as well.

448 The effects of the 'refugium phase' can be seen when examining modern short-range

endemics [i.e., those with distributions easily contained within a 100x100 km grid (i.e., <10,000

450 km^2]. An example would be the isolated New Mexico Ridge-nosed Rattlesnake (*Crotalus

willardi obscurus*) that reflects signals of relatively recent divergence and population reduction

452 (Holycross and Douglas 2007). Its historic connectivity in the Sky-islands has been truncated by

contemporary climate change, whereas its refugium phase, as gauged by demographic and niche

454 modeling, underscored a continued loss of habitat (Davis et al. 2015). We suspect a similar

refugium-expansion model would explain the extant genetic differentiation found in the present

456   study.

458   *Comparing species delimitation approaches*

We found considerable variance in the number of groups delimited using the various

460   algorithmic methods, especially when contrasted with the current taxonomy (Fig. 3). This could

reflect the capacity of the various models to diagnose cryptic lineages, or alternatively, suggest

462   the data violate implicit assumptions (Carstens et al. 2013). Phylogenetic models (i.e., ROS,

PTP, GMYC) generally produced the largest numbers of groups, but did not explain the

464   geographic patterns of genetic variation as well as did other models (Table 2; Figure 3). All

models, however, captured greater genetic variance among groups (i.e., 72-90% in AMOVA)

466   than within-group (10-28%), indicating the recognition of an authentic biological signal (Table

S1, S2).

468   Species delimitation employing the coalescent, such as GMYC, have previously been

criticized for a tendency to over-split when applied to structured populations (Lohse 2009). This,

470   in turn, suggests that a single-threshold delimitation model may be inappropriate for use with

hierarchically structured data (Talavera et al. 2013). Strong intraspecific structure in

472   *Aspidoscelis*, similar to that found in other desert reptiles (Holycross and Douglas 2007, Douglas

et al. 2002), may explain the over-fitting of GMYC. A more comprehensive test using additional

474   scenarios is warranted, as this will more appropriately characterize those biases that can emerge

when the method is applied to empirical data.

476   We also employed a second phylogenetic method (PTP) that essentially modeled

speciation as a clock-like process, with its probability of occurrence represented as a function of

478   time (i.e., branch length). Given this, any aspect that would promote rate heterogeneity could be

sufficient to violate the model, such as the extreme demographic fluctuations suspected for

480   *Aspidoscelis* populations.

Clustering-based methods, on the other hand, may be biased by elevated genetic distances

482   or incomplete sampling, with group delimitation being driven entirely by the arbitrary nature of

the (user-selected) threshold. The ABGD model attempts to mitigate this by automatically

484   detecting distance-based partitions, as bounded by priors that account for maximum allowable

intraspecific divergence (Puillandre et al. 2012). ABGD and clustering methods appeared to

486   delimit groups that more appropriately represented the genetic/ geographic distances paradigm.

When examining genetic distances defined by these methods within- versus among-group, we

488   found *A. marmoratus* approximately equal to *A. tigris* with regard to genetic distances, regardless

of whether *A. tigris* was intact or partitioned (Fig. 6). Among-group genetic distances were equal

490   regardless of how proximate were the samples (i.e., genetic x geographic distance slope

approaches zero). This suggests that divergence is maintained not by physical distance but

492   instead by some intrinsic qualities possessed by those groups, such as reproductive isolation or

mutually exclusive habitat requirements (Fig. 6). In this case, near-zero slopes in the among-

494   group comparisons lend some validity to a species-level designation for *A. marmoratus* and

suggested the potential for reproductive isolation within the *tigris* complex (i.e., between *A. t.*

496   *tigris*/*punctilinealis* and *A. t. septentrionalis*; Fig. 6B). However, reduced sample sizes for *A. t.*

mundus and *A. t. tigris* remain an issue. Within-group comparisons resulted in a positive

498   relationship with geographic distance, as was expected (Fig. 6).

500   *Species concepts and taxonomic conclusions for* Aspidoscelis

Species concepts present a philosophical quandary (Coyne and Orr, 2004). For example,

502 many modern interpretations [i.e., biological (BSC; Mayr, 1942); evolutionary (ESC; Frost and

Hillis, 1990); phylogenetic (PSC; Cracraft, 1983)] offer a mechanistic or philosophical

504 interpretation for the existence of species, yet do not themselves present an explicit operational

criterion that can universally define species. The definitions also have fundamental conflicts with

506 regards to the secondary characteristics each relies upon (e.g., reproductive isolation,

morphological diagnosability; de Queiroz, 2007).

508 Given that inferring a direct relationship between genotypic (or 'haplotypic') clusters is a

questionable approach, we instead relied upon a 'theory-independent' definition (the 'genotypic

510 cluster' concept: Douglas et al., 2007; Sullivan et al., 2014) wherein clusters are defined as '…

distinguishable groups of individuals that have few or no intermediates when in contact' (Mallet,

512 1995). We used this basis to compare among delimitation hypotheses for *Aspidoscelis tigris*, and

how they may (or may not) extend to theoretical species definitions such as the BSC.

514 An expectation of the '… little to no intermediates in contact' criterion is that individuals

from separate genotypic clusters should not be any more genetically similar when sampled in

516 proximity than when sampled from more geographically disparate locations. Such a relationship

would suggest continuous gene flow limited only by the ability of individuals to disperse (i.e.,

518 'isolation-by-distance' *sensu* Wright, 1940, 1943).

We tested the applicability of this criterion to our species hypotheses using spatial and

520 non-spatial analyses of molecular variance (AMOVA/ SAMOVA), by testing the manner by

which each partitioning scheme explained spatial signatures of genetic differentiation (IBD/

522 Mantel), and with qualitative comparisons of landscape-level genetic differentiation (AIS

interpolation). We found the ABGD model (i.e., separation of *A. marmoratus* from all *A. tigris*

524   subspecies) did well with partitioning spatial haplotypic variation, with *A. marmoratus* being

highly differentiated (>10% sequence divergence) even from localities sampled <2km from *A. t.*

526   *punctilinealis* (Fig. 6A). These patterns are consistent with many commonly applied species

concepts (e.g. reproductive isolation; per BSC), and given this, the elevated status of *A.*

528   *marmoratus* is sustained.

We also found limited (albeit, existing) exchange between two major clades within the

530   remaining *A. tigris*: *A. t. tigris/punctilinealis* versus *A. t. septentrionalis/mundus* (i.e., TAX

model). Here, genetic differentiation was again uncorrelated with spatial distances (Fig. 6B),

532   possibly implicating intrinsic or extrinsic factors that may satisfy species-level criteria. Previous

estimates place divergence of these groups to be no greater than 5-10 kya, based upon

534   availability of suitable habitat following the Wisconsin glacial period (Zweifel, 1962). If so, then

our observed mean corrected genetic distance of 0.058 equates to a mutation rate of 5.8

536   substitutions per million years, a rate considerably greater than the expected rate of ~0.0045 for

ATPase 6, the more rapidly evolving of our two mitochondrial genes (Kumar, 1996; Mueller,

538   2006).

Although we recognize that numerous factors contribute to mtDNA rate heterogeneity,

540   our rudimentary calculation suggests that subspecies within *A. tigris* may have diverged much

earlier than previously thought. Although our results support the continued recognition of *A. t.*

542   *septentrionalis* as distinct, we deem them insufficient to propose elevation beyond a subspecific

level. Instead, we argue for greater sampling at both geographical and genomic scales to properly

544   adjudicate taxonomies within *A. t. tigris*. The latter, while enabling tests of hybridization/

introgression within a more statistically robust framework (e.g., Chafin et al., 2019), would also

546   open avenues for novel species delimitation algorithms to be formulated using unsupervised

machine learning approaches (Derkarabetian et al., 2019; Martin et al., 2020; Mussmann et al.,

548   2020).

The CD5 model identified potential cryptic lineages within a clade comprising *A. t. tigris*

550   and *A. t. punctilinealis*, although sample sizes representing this group were limited. Of note, *A. t.*

tigris and *A. t. punctilinealis* were not reciprocally monophyletic, however assigning causality of

552   this to hybridization/ intergradation, stochastic lineage sorting, or a lack of phylogenetic signal

requires additional data (Chafin et al., 2020). Again, we suggest a more in-depth examination

554   using a larger panel of molecular markers so as to confirm or reject continued recognition of *A. t.*

tigris as distinct from *A. t. punctilinealis*, and likewise to adjudicate recognition of *A. t. mundus*

556   from *A. t. septentrionalis*.

558   *Conclusion*

We found that algorithmic species delimitation methods vary substantially in the manner

560   by which they partition hierarchical phylogenetic diversity, particularly when contrasted against

the respected overlays of historical biogeography, morphology, ecology, and prior genetic

562   evaluations. By this we mean that some algorithms (e.g., ABGD) yielded groupings which

support external evidence of divergence at the 'species' level (e.g., according to a BSC-like

564   interpretation), while others (e.g., CD5) yield groupings that could potentially fit various

operational species concepts, yet require further evaluation using genomic data. Others (GMYC,

566   PTP) seemingly delimit populations, a conclusion in line with prior evaluations of simulated and

empirical datasets. The latter was hypothesized to be impacted by inflated levels of inter-

568   population divergence, a phenomenon which has been observed in other co-occuring desert

fauna subjected to the same historical climatic fluctuation. These observations prompt us to

570 support the elevation of *A. marmoratus*, but additional groups withinthe *A. tigris* complex will

requiring additional data to properly adjudicate.

574 **ACKNOWLEDGMENTS**

Samples were collected by BKS and JEC under authority of permits provided by Arizona and New

576 Mexico Game and Fish departments. Collecting methods were approved as part of IACUC protocols for

surveying and vouchering lizards (BKS and JEC). We are also indebted to students, post-doctorals, and

578 faculty who have promoted this research: M. R. Bangs, L. Graham, B. A. Levine, B. T. Martin, S. M.

Mussmann, Z. D. Zbinden, and especially K.E. Harrell. This research was made possible through

580 generous endowments to the University of Arkansas: A Doctoral Dissertation Fellowship (TKC), the

Bruker Professorship in Life Sciences (MRD), and the Twenty-First Century Chair in Global Change

582 Biology (MED), all of which provided salaries and/or research funds for this work. Additional analytical

resources were provided by the Arkansas Economic Development Commission (Arkansas Settlement

584 Proceeds Act of 2000) and the Arkansas High Performance Computing Center (AHPCC), and from an

NSF-XSEDE Research Allocation (TG-BIO160065) to access the Jetstream cloud. Summer support of

586 field work for JEC was derived from consecutive Endowed Professorships in Science by Opelousas

General Hospital of Opelousas, Louisiana.

LITERATURE CITED

590 Bangs, M.R., Douglas, M.R., Brunner, P.C., & Douglas, M.E. (2020a) Reticulate evolution as a
management challenge: Patterns of admixture with phylogenetic distance in endemic fishes
592 of western North America. Evolutionary Applications, 13, 1400-1419.

Bangs, M.R., Douglas, M.R., Chafin, T.K., & Douglas, M.E. (2020b) Gene flow and species
594 delimitation in fishes of Western North America: Flannelmouth (*Catostomus latipinnis*) and

Bluehead sucker (*C. Pantosteus discobolus*). Ecology and Evolution, 10 6477-6493.

596   Barley, A.J., Nieto-Montes de Oca, A., Reeder, T.W., Manríquez-Morán, N.L., Arenas Monroy,
       J.C., Hernández-Gallegos, O. & Thomson, R.C. (2019) Complex patterns of hybridization
598   and introgression across evolutionary timescales in Mexican whiptail lizards (Aspidoscelis).
       Molecular Phylogenetics and Evolution, 132, 284–295.

600   Barton, N.H. & Hewitt, G.M. (1985) Analysis of hybrid zones. Annual Review of Ecology and
       Systematics, 16, 113–148.

602   Bhattarachya, T., Tierney, J.E., Addison, J.A., & Murray, J.W. (2018) Ice-sheet modulation of
       deglacial North American monsoon intensification. Nature Geoscience, 11, 848-852.

604   Blakely, R., & Ranney, W. (2008) Ancient Landscapes of the Colorado Plateau. Grand Canyon
       Association, Grand Canyon, AZ.

606   Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., Suchard, M.A.,
       Rambaut, A. & Drummond, A.J. (2014) BEAST 2: A software platform for Bayesian
608   evolutionary analysis. PLoS Computational Biology, 10, e1003537.

       Brown, S.D.J, Collins, R.A., Boyer, S., Lefort, M., Malumbres-Olarte, J., Vink, C.J., &
610   Cruickshank, R.H. (2012) Spider: An R package for the analysis of species identity and
       evolution, with particular reference to DNA barcoding. Molecular Ecology Resources,
612   12, 562-565.

614   Brunner, P.C., Douglas, M.R., Osinov, A., Wilson, C.C., & Bernatchez, L. (2001) Holarctic
       phylogeography of Arctic char (*Salvelinus alpinus* L.) inferred from mitochondrial DNA
616   sequences. Evolution, 55, 573-586.

       Bryson, R.W., Murphy, R.W., Lathrop, A., & Lazcano-Villareal, D. (2011) Evolutionary drivers
618   of phylogeographical diversity in the highlands of Mexico: A case study of the *Crotalus
       triseriatus* species group of montane rattlesnakes. Journal of Biogeography, 38, 697-710.

620   Carstens, B.C., Pelletier, T. a., Reid, N.M. & Satler, J.D. (2013) How to fail at species
       delimitation. Molecular Ecology, 22, 4369–4383.

622   Castoe, T.A., Spencer, C.L. & Parkinson, C.L. (2007) Phylogeographic structure and historical
       demography of the western diamondback rattlesnake (*Crotalus atrox*): A perspective on
624   North American desert biogeography. Molecular Phylogenetics and Evolution, 42, 193–
       212.

626   Chafin, T.K., Douglas, M.R., Bangs, M.R., Martin, B.T., Mussmann, S.M., & Douglas, M.E.
       (2020) Taxonomic uncertainty and the anomaly zone: Phylogenomics disentangle a rapid

628   radiation to resolve contentious species (*Gila robusta* complex) in the Colorado River. bioRxiv, 692509.

630   Cole, K.L. (1986) The Lower Colorado River Valley: A Pleistocene desert. Quaternary Research 25, 392-400.

632   Cracraft, J. (1983) Species concepts and speciation analysis. In: Current ornithology (ed. by Ketterson, E., & Nolan, V.), pp. 159-187, Springer, New York, NY.

634   Crother, B.I. (ed) (2017) Scientific and standard english names of amphibians and reptiles of North America north of Mexico, with comments regarding confidence in our understanding
636   8th edition. Herpetological Circular, 43, 1–102.

Darriba, D., Taboada, G.L., Doallo, R. & Posada, D. (2012) jModelTest 2: more models, new
638   heuristics and parallel computing. Nature Methods, 9, 772–772.

Davis, M.A., Douglas, M.R., Webb, C.T., Collyer, M.L., Holycross, A.T., Painter, C.W.,
640   Kamees, L.K. & Douglas, M.E. (2015) Nowhere to go but up: Impacts of climate change on demographics of a short-range endemic (*Crotalus willardi obscurus*) in the sky-islands of
642   southwestern North America. PLoS ONE, 10, 1–20.

Dessauer, H.C. & Cole, C.J. (1991) Genetics of whiptail lizards (Reptilia: Teiidae:
644   *Cnemidophorus*) in a hybrid zone in southwestern New Mexico. Copeia, 1991, 622–637.

Dessauer, H.C., Cole, C.J. & Townsend (2000) Hybridization among western whiptail lizards
646   (*Cnemidophorus tigris*) in southwestern New Mexico: Population genetics, morphology, and ecology in three contact zones. Bulletin of the American Museum of Natural History,
648   246, 1–148.

Dorsey, R.J., Fluette, A., McDougall, K., House, B.A., Janecke, S.U., Axen, G.J., & Shirvell,
650   C.R. (2007) Chronology of Miocene-Pliocene deposits at Split Mountain Gorge, Southern California: A record of regional tectonic and Colorado River evolution. Geology, 35, 57-60.

652   Douglas, M.E., Douglas, M.R., Schuett, G.W., Beck, D.D. & Sullivan, B.K. (2010) Conservation phylogenetics of helodermatid lizards using multiple molecular markers and a supertree
654   approach. Molecular Phylogenetics and Evolution, 55, 153–167.

Douglas M.E., Douglas, M.R., Schuett, G.W., & Porras, L.W. (2009) Climate change and the
656   evolution of the New World pitviper genus *Agkistrodon* (Viperidae). Journal of Biogeography, 36, 1164-1180.

658   Douglas, M.E., Douglas, M.R., Schuett, G.W. & Porras, L.W. (2006) Evolution of rattlesnakes (Viperidae; *Crotalus*) in the warm deserts of western North America shaped by Neogene

660   vicariance and Quaternary climate change. Molecular Ecology, 15, 3353–3374.

Douglas, M.E., Douglas, M.R., Schuett, G.W., Porras, L.W. & Holycross, A.T. (2002)
662   Phylogeography of the western rattlesnake (*Crotalus viridis*) complex, with emphasis on the
       Colorado Plateau. In: Biology of the Vipers (ed. by Schuett, G.W., Höggren, M., Douglas,
664   M.E., & Greene, H.W.), pp. 11-50, EEagle Mountain Publishing, Eagle Mountain, UT.

Douglas, M.E., Minckley, W.L., & DeMarais, B.D. (1999) Did vicariance mold phenotypes of
666   Western North American fishes? Evidence from Gila River cyprinids. Evolution, 53, 238-
       246.

668   Douglas, M.R., Davis, M.A., Amarello, M., Smith, J.J., Schuett, G.W., Herrmann, H.W.,
       Holycross, A.T., & Douglas, M.E. (2016) Anthropogenic impacts drive niche and
670   conservation metrics of a cryptic rattlesnake on the Colorado Plateau of western North
       America. Royal Society Open Science, 3, 160047.

672   Drummond, C.S. & Hamilton, M.B. (2007) Hierarchical components of genetic variation at a
       species boundary: population structure in two sympatric varieties of *Lupinus microcarpus*
674   (Leguminosae). Molecular Ecology, 16, 753–769.

Dupanloup, I., Schneider, S. & Excoffier, L. (2008) A simulated annealing approach to define
676   the genetic structure of populations. Molecular Ecology, 11, 2571–2581.

Eberly, L.D., & Stanley, T.B. (1978) Cenozoic stratigraphy and geologic history of southwestern
678   Arizona. Geological Society of America Bulletin, 89, 921-940.

Ericson, P.G.P., Qu, Y., Rasmussen, P.C., Blom, M.P.K., Rheindt, F.E., & Irestedt, M. (2019)
680   Genomic differentiation tracks earth historic isolation in an Indo-Australasian archipelagic
       pitta (Pittidae; Aves) complex. BMC Evolutionary Biology, 19, 151.

682   Excoffier, L., Smouse, P.E. & Quattro, J.M. (1992) Analysis of molecular variance inferred from
       metric distances among DNA haplotypes: application to human mitochondrial DNA
684   restriction data. Genetics, 131, 479–491.

Frost, D.R. & Hillis, D.M. (1990) Species in concept and practice : Herpetological applications.
686   Herpetologica, 46, 86–104.

Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. (2012) CD-HIT: Accelerated for clustering the next-
688   generation sequencing data. Bioinformatics, 28, 3150–3152.

Fujisawa, T. & Barraclough, T.G. (2013) Delimiting species using single-locus data and the
690   Generalized Mixed Yule Coalescent Approach: A revised method and evaluation on
       simulated data sets. Systematic Biology, 62, 707-724.

Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W. & Gascuel, O. (2010) New
694   algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the
        performance of PhyML 3.0. Systematic Biology, 59, 307–321.

696   Hewitt, G.M. (2000) The genetic legacy of the Quaternary ice ages. Nature, 405, 907–913.

Hickerson, M.J. & Meyer, C.P. (2008) Testing comparative phylogeographic models of marine
698   vicariance and dispersal using a hierarchical Bayesian approach. BMC Evolutionary
        Biology, 8, 1–18.

700   Holycross, A.T. & Douglas, M.E. (2007) Geographic isolation, genetic divergence, and
        ecological non-exchangeability define ESUs in a threatened sky-island rattlesnake.
702   Biological Conservation, 134, 142–154.

House, P.K., Pearthree, P.A., & Perkins, M.E. (2008) Stratigraphic evidence for the role of lake
704   spillover in the inception of the lower Colorado River in southern Nevada and western
        Arizona. Special Paper of the Geological Society of America, 439, 335-353.

706   Kohli, B.A., Fedorov, V.B., Waltari, E., & Cook, J.A. (2015) Phylogeography of the Holarctic
        rodent (*Myodes rutilus*): Testing high-latitude biogeographical hypotheses of range shifts.
708   Journal of Biogeography, 42, 377-389.

Kumar, S. (1996) Patterns of nucleotide substitution in mitochondrial protein coding genes of
710   vertebrates. Genetics, 143, 537–548.

Lamb, T., Avise, J.C. & Gibbons, J.W. (1989) Phylogeographic patterns in mitochondrial-DNA
712   of the desert tortoise (*Xerobates agassizi*), and evolutionary relationships among the north-
        american gopher tortoises. Evolution, 43, 76–87.

714   Lamb, T., Jones, T.R. & Avise, J.C. (1992) Phylogeographic histories of representative
        herpetofauna of the southwestern U.S.: mitochondrial DNA variation in the desert iguana
716   (*Dipsosaurus dorsalis*) and the chuckwalla (*Sauromalus obesus*). Journal of Evolutionary
        Biology, 5, 465–480.

718   Lohse, K. (2009) Can mtDNA barcodes be used to delimit species? A response to Pons et al.
        (2006). Systematic Biology 58, 439–442.

720   Mayr, E. (1942) Systematics and the Origin of Species, Columbia University Press, New York,
        NY.

722   Meirmans, P.G. (2012) The trouble with isolation by distance. Molecular Ecology, 21, 2839–
        2846.

724   Miller, M.P. (2005) Alleles In Space (AIS): Computer software for the joint analysis of
            interindividual spatial and genetic information. Journal of Heredity, 96, 722–724.

726   Mueller, R.L. (2006) Evolutionary rates, divergence dates, and the performance of mitochondrial
            genes in Bayesian phylogenetic analysis. Systematic Biology, 55, 289–300.

728   O'Connell, K.A., Streicher, J.W., Smith, E.N. & Fujita, M.K. (2017) Geographical features are
            the predominant driver of molecular diversification in widely distributed North American
730   whipsnakes. Molecular Ecology, 26, 5729–5751.

        Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J. & Suggests,
732   M. (2007) The vegan package. Community ecology package, 10, 631–637.

        Martin, B.T., Chafin, T.K., Douglas, M.R., Placyk Jr., J.S., Birkhead, R.D., Phillips, C.A., &
734   Douglas, M.E. (2020) Species delimitation using machine learning recovers a
            phylogenomically consistent classification for North American box turtles (*Terrapene* spp.)
736   bioRxiv, 205039.

        Migliore, J., Kaymak, E., Mariac, C., Couvreur, T.L.P., Lissambo, B.-J., Piñeiro, R., & Hardy,
738   O.J. (2019) Pre-PLeistocene origin of phylogeographical breaks in African rain forest trees:
            New insights from *Greenwayodendron* (Annonaceae). Journal of Biogeography, 46, 212-
740   223.

        Minckley, W.L., Hendrickson, D.A., & Bond, C.E. (1986) Geography of western North
742   American freshwater fishes: Description and relations to intracontinental tectonism. In:
            Zoogeography of North American Freshwater Fishes (ed. by Hocutt, C.H., & Wiley, E.O).
744   pp. 519-613. John Wiley & Sons, New York, NY.

        Morales-Barbero, J., Martinez, P.A., Ferrer-Castán, D., & Olalla-Tárraga, M.A. (2018)
746   Quaternary refugia are associated with higher speciation rates in mammalian faunas of the
            western Palearctic. Ecography, 41, 607-621.

748   Mussmann, S.M., Douglas, M.R., Oakey, D.D., & Douglas, M.E. (2020) Defining relictual
            biodiversity: Conservation units in speckled dace (Leuciscidae: *Rhinichthys osculus*) of the
750   Greater Death Valley ecosystem. Ecology and Evolution, Early View.

        Paradis, E., Claude, J. & Strimmer, K. (2004) APE: Analyses of phylogenetics and evolution in
752   R language. Bioinformatics, 20, 289–290.

        Pianka, E.R. (1970) Comparative autecology of the lizard *Cnedimophorus tigris* in different parts
754   of its geographic range. Ecology, 51, 703–720.

        Pianka, E.R. (1967) On lizard species diversity: North American flatland deserts. Ecology, 48,

756   333–351.

Pons, J., Barraclough, T.G., Gomez-Zurita, J., Cardoso, A., Duran, D.P., Hazell, S., Kamoun, S.,
758   Sumlin, W.D. & Vogler, A.P. (2006) Sequence-based species delimitation for the DNA
      taxonomy of undescribed insects. Systematic Biology, 55, 595–609.

760   Pounds, J.A. & Jackson, J.F. (1981) Riverine barriers to gene flow and the differentiation of
      fence lizard populations. Evolution, 35, 516–528.

762   Puillandre, N., Lambert, A., Brouillet, S. & Achaz, G. (2012) ABGD, automatic barcode gap
      discovery for primary species delimitation. Molecular Ecology, 21, 1864–1877.

764   Pyron, R.A., Costa, G.C., Patten, M.A. & Burbrink, F.T. (2015) Phylogenetic niche conservatism
      and the evolutionary basis of ecological speciation. Biological Reviews, 90, 1248–1262.

766   de Queiroz, K. (2005) A unified species concept and its consequences for the future of
      taxonomy. Proceedings of the California Academy of Sciences, 56, 196–215.

768   R Core Team (2020) R: A language and environment for statistical computing. R Foundation for
      Statistical Computing, Vienna, Austria.

770   Rambaut, A. & Drummond, A.J. (2007) Tracer v1.4. Digital resource available at
      http://tree.bio.ed.ac.uk/software/tracer/

772   Riddle, B.R. & Hafner, D.J. (2006) A step-wise approach to integrating phylogeographic and
      phylogenetic biogeographic perspectives on the history of a core North American warm
774   deserts biota. Journal of Arid Environments, 66, 435–461.

Rosenberg, N.A. (2007) Statistical tests for taxonomic distinctiveness from observations of
776   monophyly. Evolution, 61, 317–323.

Sarna-Wojcicki, A.M., Deino, A.L., Fleck, R.J., McLaughlin, R.J., Wagner, D., Wan, E., Wahl,
778   D., Hillhouse, J.W., & Perkins, M. (2011) Age, composition, and areal distribution of the
      Pliocene Lawlor Tuff, and thee younger Pliocene tuffs, California and Nevada. Geosphere,
780   7, 599-628.

Sexton, J.P., Hangartner, S.B. & Hoffmann, A.A. (2014) Genetic isolation by environment or
782   distance: Which pattern of gene flow is most common? Evolution, 68, 1–15.

Smouse, P.E., Long, J.C. & Sokal, R.R. (1986) Multiple regression and correlation extensions of
784   the Mantel test of matrix correspondence. Systematic Zoology, 35, 627.

Spencer, J.E., Smith, G.R., & Dowling, T.E. (2008) Middle to Late Cenozoic geology,

786   hydrography, and fish evolution in the American Southwest. Geolocial Society of America, 80301.

788   Talavera, G., Dincă, V. & Vila, R. (2013) Factors affecting species delimitations with the GMYC model: Insights from a butterfly survey. Methods in Ecology and Evolution, 4,
790   1101–1110.

        Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011) MEGA5:
Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. Molecular Biology and Evolution, 28, 2731–2739.

Taylor, H.L. (1988) A morphological analysis of intergredation between the teiid lizards *Cnemidophorus tigris tigris* and *C. tigris septentrionalis*. Herpetologica, 44, 176–185.

Taylor, H.L. (1990) A morphological analysis of the teiid lizards *Cnemidophorus tigris tigris* and *C. tigris gracilis* from a contact zone in northwestern Arizona. Herpetologica, 46, 447–456.

Taylor, H.L., Beyer, C., Harris, L. & Pham, H. (1994) Subspecific relationships in the Teiid lizard *Cnemidophorus tigris* in southwestern Arizona. Journal of Herpetology, 28, 247–253.

Thompson, R.S. (1991) Pliocene environments and climates in the western United States. Quaternary Science Review, 10, 115-132.

Tucker, D.B., Colli, G.R., Giugliano, L.G., Hedges, S.B., Hendry, C.R., Lemmon, E.M., Lemmon, A.R., Sites Jr., J.W., & Pyron, R.A. (2016) Methodological congruence in
phylogenomic analyses with morphological support for teiid lizards (Sauria: Teiidae). Molecular Phylogenetics and Evolution, 103, 75-84.

Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B.C., Remm, M. & Rozen, S.G. (2012) Primer3-new capabilities and interfaces. Nucleic Acids Research, 40, 1–12.

Wright, S. (1940) Breeding structure of populations in relation to speciation. The American Naturalist, 74, 232–248.

Wright, S. (1943) Isolation by distance. Genetics, 28, 114–138.

        Zhang, J., Kapli, P., Pavlidis, P. & Stamatakis, A. (2013) A general species delimitation method
812   with applications to phylogenetic placements. Bioinformatics, 29, 2869–2876.

        Zweifel, R.G. (1962) Analysis of hybridization between two subspecies of the desert whiptail
814   lizard, *Cnemidophorus tigris*. Copeia, 1962, 749–766.

816   Dorsey, R.J., O'Connell, B., McDougall, K., Homan, M.B. (2018) Punctuated sediment
          discharge during Early Pliocene birth of the Colorado River: Evidence from regional
818   stratigraphy, sedimentology, and paleontology. Sedimentary Geology, 363, 1-33.
          Doi:/10.1016/j.sedgeo.2017.09.018

    Flowers, R.M. (2010) The enigmatic rise of the Colorado Plateau. Geology, 38, 671-672.
822   Doi:/10.1130/focus072010.1

824   Levander, A., Schmandt, B., Miller, M.S., Liu, K., Karlstrom, K.E., Crow, R.S., Lee, C-T.A.,
          Humphreys, E.D. (2011) Continuing Colorado Plateau uplift by delamination-style
826   convective lithospheric downwelling. Nature, 472, 461-466. Doi:/10.1038/nature10001

828   Nicholson, U., Carter, A., Robinson, P., Macdonald, D.I.M. (2019) Eocene–Recent drainage
          evolution of the Colorado River and its precursor: An integrated provenance perspective
830   from SW California. Geological Society Special Publications, 488, 47-72.
          Doi:/10.1144/SP488-2019-272

Figures and Tables

836 Table 1: Population summary statistics for clades within the *Aspidoscelis tigris* complex as determined by sequence analysis of 777bp of the ATPase 6 and 8 genes across 239 samples.

838 Abbreviations are as follows: H = haplotype diversity; π = nucleotide diversity; θ = population mutation rate; θ_s/N = number of segregating sites / clade sample size; H/N = number of

840 haplotypes / clade sample size. *A. t. mundus* was excluded due to low sample size.

CLADE	TAJIMA D	TAJIMA D P-VALUE	H	π	θ	FU & LI F	FU & LI D	θ_s/N	H/N
A	-1.522	0.128	0.993	0.013	15.409	-1.680	-1.595	2.889	0.944
B	-0.938	0.348	0.969	0.011	11.258	-0.442	-0.356	2.833	0.916
C	-1.628	0.103	0.997	0.012	17.033	-2.051	-1.935	2.461	0.923
D	-1.889	0.059	0.966	0.018	26.468	-2.968	-3.008	3.885	0.654
E	-0.524	0.600	1.000	0.013	10.363	0.050	0.524	4.75	1.000
F	0.616	0.538	1.000	0.021	14.890	0.660	0.374	5.667	1.000
G	-2.114	0.017	0.478	0.006	15.600	-2.609	-2.123	1.123	0.292
H	-1.778	0.075	0.828	0.002	3.762	0.457	-2.933	0.4	0.425

Table 2: Comparison of alternative species delimitation models according to the fit of the data under an isolation-by-distance model. Results are shown for 2-way, stratified, and partial Mantel

tests, reporting a correlation coefficient (r) between distance matrices composed of geographic and genetic distances. Correlation coefficients in the partial Mantel tests reflect association

between clade membership and genetic distance, with geographic distances as a covariate. Values indicated by (**) are significant at p>0.0001. Species delimitation models are abbreviated

as: ABGD=Automated Barcode-Gap Discovery; TAX=qualitative hypothesis derived from current taxonomy; CD5=CD-Hit with 5% divergence threshold for clustering; CD2= CD-Hit

with 2% divergence threshold for clustering; GMYC=Generalized mixed Yule coalescent; PTP=Poisson tree process; ROS=Clade delimitation based on Rosenberg test of reciprocal

monophyly.

MODEL	2-WAY MANTEL TEST	STRATIFIED 2-WAY MANTEL	PARTIAL MANTEL
	Mantel r	Mantel r	Mantel r
ALL	0.5762**	-	-
ROS	-	0.5762**	0.6317**
ABGD	-	0.5762**	**0.7770****
TAX	-	0.5762**	0.7118**
CD5	-	0.5762**	0.7025**
CD2	-	0.5762	0.5816**
GMYC	-	0.5762	0.5818**
PTP	-	0.5762	0.5754**

**Significant at α=0.0001

Figure 1: Range map and sampling sites for *Aspidoscelis marmoratus* and subspecies of *A. tigris* (subspecific units within *A. marmoratus* and Mexican subspecies of *A. tigris* not shown). Three

identified contact zones are depicted between: (A) *A. t. tigris* and *A. t. punctilinealis*, roughly transected by the Colorado River with morphoclines varying from 16km in the north to ~180km

in the south; (B) *A. t. tigris* x *A. t. septentrionalis*, where color-pattern intergradation corresponds to a ~50km transition from Mohave (inset; M) to Great Basin (GB) desert-scrub communities to

the Colorado Plateau ecoregion (CP); and (C) *A. t. punctilinealis* x *A. marmoratus*, a hybrid zone of <2km in width traversing the Cochise Filter Barrier transition zone between the Sonoran

(inset; S) and Chihuahuan (Ch) deserts. Note that both sample points and ranges are colored by subspecies, as defined in the legend. Map based on observations and prior work by BKS.

Figure 2: Pattern polymorphism in the *Aspidoscelis tigris* group: (A–B) *Aspidoscelis tigris mundus*; (C) *A. t. septentrionalis* from northern Arizona; (D) *A. t. tigris* from California; (E–F) *A.*
t. punctilinealis from central Arizona; (G–H) *A. marmoratus* from New Mexico; (I) Dorsal and (J) ventral view of suspected *A. t. tigris* x *A. t. punctilinealis* hybrids. Pictures provided by BKS.
876

878   Figure 3: Neighbor-joining (A) and Bayesian (B) phylogenetic trees based on sequence analysis of 777bp of the ATPase 6 and 8 genes
       across 239 samples, contrasted against results of *de novo* species delimitation models represented by colored bars. Branches are
880   labelled according to the morphological diagnoses of the tissue samples from which mtDNA sequences originated and are abbreviated
       as follows: *tigris*=*A. tigris tigris*; *punct.*=*A. t. punctilinealis*; *sept.*=*A. t. septentrionalis*; *mundus*=*A. t. mundus*; *marm.*=*A. marmoratus*;
882   *shared*=found in *A. t. tigris* and *A. t. punctilinealis*. Model abbreviations are: ABGD=Automated Barcode-Gap Discovery;
       TAX=qualitative hypothesis derived from current taxonomy; CD5=CD-Hit with 5% divergence threshold for clustering; CD2= CD-
884   Hit with 2% divergence threshold for clustering; GMYC=Generalized mixed Yule coalescent; PTP=Poisson tree process; ROS=Clade
       delimitation based on statistical tests of monophyly. Node labels represent posterior probabilities from Bayesian inference (B), and
886   neighbor-joining bootstrap support (A; omitted for shallow nodes). Nodes passing the statistical test of reciprocal monophyly are
       indicated with a red dot. Clades in both trees are colored according to the ROS model.

Figure 4: Geo-referenced haplotype tree generated via Bayesian analysis, representing 777bp of the ATPase 8 and 6 genes across 239 samples. Clade coloring follows the ROS model (see Fig. 1)

Figure 5: Among-group genetic variance (F_{CT}) for genetic clusters within the *Aspidoscelis tigris*
898   complex, as derived by SAMOVA clustering runs. Cluster optimization constrained by (A)
geographic proximity of samples and (B) without a geographic constraint. The x-axis (*K*) reports
900   the number of genetic clusters sampled.

904   Figure 6: Within- and among-group linear regression of geographic x genetic distances for clades delimited by (A) the ABGD (Automated barcode-gap discovery) model and (B) TAX (taxonomic) model. Coefficients of correlation (r^2) ranged from 0.01-0.449.
906   Note that "*tigris*" refers to both *A. t. tigris* and *A. t. punctilinealis*, as these subspecies are not differentiated in the models. *A. t. mundus* was excluded due to the geographically restricted sampling presented herein.
908

910   Figure 7: Map depicting results of an 'Alleles in Space' analysis for members of the *Aspidoscelis*
tigris complex. Relative genetic differentiation is interpolated across the landscape, with
912   individuals (as dots) colored by the CD5 model (see Fig. 1B). Heat map colors depict relative
genetic distances per unit of geographic distance, with red corresponding to a greater genetic
914   distance, and blue being lesser genetic distance.