Taxonomic hypotheses and the biogeography of speciation in the Tiger Whiptail complex (*Aspidoscelis tigris*: Squamata, Teiidae)

Tyler K. Chafin[1], Marlis R. Douglas[1], Whitney J.B. Anthonysamy[1,2], Brian K. Sullivan[3], James M. Walker[1], James E. Cordes[4], and Michael E. Douglas[1]

[1] Department of Biological Sciences, University of Arkansas, Fayetteville, Arkansas 72701, USA; Email: (TKC) tkchafin@uark.edu, send reprint requests to this address; (MRD) mrd1@uark.edu; (JMW) jmwalker@uark.edu; and (MED) med1@uark.edu.

[2] Present address: St. Louis College of Pharmacy, 4588 Parkview Place, St. Louis, Missouri 63110, USA; Email: whitney.anthonysamy@stlcop.edu.

[3] School of Mathematical and Natural Sciences, P. O. Box 37100, Arizona State University, Phoenix, Arizona 85069, USA; Email: bsullivan@asu.edu.

[4] Division of Mathematics and Sciences, Louisiana State University, Eunice, Louisiana 70535, USA; Email: jcordes1@live.com.

Running head: *Aspidoscelis tigris* phylogeography

Keywords: whiptail lizards, climate change; mtDNA; southwestern United States; species concept; subspecies

(225)Biodiversity in southwestern North America has a complex biogeographic history involving

2   tectonism interspersed with climatic fluctuations. This yields a contemporary pattern replete with

historic idiosyncrasies often difficult to interpret when viewed from through the lens of modern

4   ecology. The *Aspidoscelis tigris* (Tiger Whiptail) complex (Squamata: Teiidae) is one such

group in which potential taxonomic boundaries have been confounded by a series of complex

6   biogeographic processes that have defined the evolution of the clade. To clarify this situation, we

first generated multiple taxonomic hypotheses, which were subsequently tested using

8   mitochondrial DNA sequences (ATPase 8 and 6) evaluated across 239 individuals representing

five continental members of this complex. To do so, we evaluated the manner by which our

10   models parsed phylogenetic and biogeographic patterns. We found considerable variation among

species 'hypotheses', which we suggest in part reflects inflated levels of inter-population genetic

12   divergence caused by historical demographic expansion and contraction cycles. Inter-specific

boundaries with *A. marmoratus* juxtaposed topographically with the Cochise Filter Barrier that

14   separates Sonoran and Chihuahuan deserts (interpreted herein as case of 'soft' allopatry).

Patterns of genetic divergence were consistent across the Cochise Filter Barrier, regardless of

16   sample proximity. Surprisingly, this also held true for intraspecific comparisons that spanned the

Colorado River. These in turn suggest geomorphic processes as a driver of speciation in the *tigris*

18   complex, with intraspecific units governed locally by demographic processes.

HIGHLIGHTS(185)

1. Phylogeographies of vertebrates within the southwestern deserts of North America have been shaped by climatic fluctuations imbedded within broad geomorphic processes.

2. The resulting synergism drives evolutionary processes, such as an expansion of within-species genetic divergence over time. Taxonomic inflation often results (i.e., an increase in recognized taxa due to arbitrary delineations), such as when morphological divergences fail to juxtapose with biogeographic hypotheses.

3. However, isolated groups can be discriminated within-species by mapping genetic variability onto geographic distances. This approach can often diagnose 'hard' barriers to dispersal, or alternatively, strong selection acting against hybridization. On the other hand, elevated genetic divergences among groups less-isolated would underscore isolation-by-distance (i.e., an increase in genetic differentiation concomitant with geographic distance).

4. The biogeographic patterns we identified in Tiger Whiptail are largely synonymous with those found in other regional species, particularly given the geomorphic separation of Mohave and Sonoran deserts by the Colorado River, and Sonoran/ Chihuahuan deserts by the Cochise Filter Barrier.

5. Our results for the Tiger Whiptail complex broaden and extend the context within which polytypic species are conserved and managed, particularly those that reflect an incongruence among molecular and morphological standards.

INTRODUCTION

Whiptail lizards (*Aspidoscelis*; Teiidae) comprise one of the most conspicuous groups of lizards in western North America. Their complex history of hybridization/ introgression coupled

with multiple transitions to parthenogenic life histories fuels their extraordinary capacity as an evolutionary model (Barley et al. 2019). However, intraspecific morphological variation coupled

with high rates of hybridization has made species-delineation difficult, which in turn has muddied their taxonomy. The occurrence of paraphyly in the group was partially resolved by

subsequently partitioning species between a resurrected *Aspidoscelis* and the nominate *Cnemidophorus* (Reeder et al., 2002). While a positive step, it also necessitated reallocation of

several groups, such as: *Aspidoscelis cozumelus* (parthenogenetic species), *A. deppi* (gonochoristic species), *A. sexlineatus* (parthenogenetic and gonochoristic species), *A.*

tesselatus (parthenogenetic species), and *A. tigris* (gonochoristic species). As a result, many systematic issues inherent within *Cnemidophorus* were simply transferred to *Aspidoscelis*.

The *A. tigris* species-group, for example, represents a perplexing systematic challenge as it comprises a widely distributed complex of gonochoristic populations located within desert

regions of southwestern North America, northern México (to include Baja California), as well as numerous islands east and west of the peninsula. Although the group *per se* is characterized by a

readily assessed set of morphological characters (Walker and Maslin, 1981) and a distinctive karyotype (Lowe et al., 1970), its diversity (i.e., ranging from a single species to more than a

dozen) has been a contentious issue in North American Herpetology. Much of its diversity is represented by distinct populations located on relatively small, vulnerable islands in the Gulf of

California. Several of these were designated as distinct species (Walker et al., 1966a, b; Walker and Taylor, 1968; Walker and Maslin, 1981), based snout-vent length, color pattern, and/or

morphology (e.g., *A. bacatus* of Isla San Pedro Nolasco; *A. catalinensis* of Isla Santa Catalina; *A. martyris* of Isla San Pedro Martír; and *A. celeripes* of Isla San Jose). However, some authors

(e.g. Lowe et al., 1970; Wright, 1993) have considered them instead as subspecific variants. Continental populations have also proven taxonomically contentious. For example, a long-

standing disagreement (Hendricks and Dixon, 1986; Dessauer et al., 2000) centers on whether *A. tigris* and *A. marmoratus* in small areas in Hidalgo County NM represent intergrading subspecies

or hybridizing species.

Taxonomic difficulties are thus quite apparent within the *A. tigris* complex, and primarily

stem from complex patterns of intergradation and hybridization confounded by non-discriminating morphological variation. To provide clarification, we evaluated a complex of four

continental subspecies (*A. t. mundus*, *A. t. punctilinealis*, *A. t. septentrionalis*, and *A. t. tigris*; Fig. 1), plus a recently elevated putative sister species, *A. marmoratus* (Hendricks and Dickson,

1986; Reeder et al., 2002; Tucker et al., 2016). Our goal was to examine how mitochondrial (mt)DNA variation in *A. tigris* subspecies compares with morphological discordance [e.g.,

subspecies seemingly supported by color pattern yet weakly differentiated by scalation and morphometrics (Taylor et al., 1994; Walker et al., 2015; Fig. 2)]. Furthermore, morphological

intergrades and overlapping genetic polymorphisms suggest admixture among several groups: *A. t. tigris* and *A. t. septentrionalis* (Taylor 1988), *A. t. tigris* and *A. t. punctilinealis* (Taylor 1990),

and *A. t. punctilinealis* and *A. t. septentrionalis* (Dessauer et al. 2000).

The contact zone separating *A. t. punctilinealis* and *A. marmoratus* (i.e., *A. t. marmoratus*

86 of the following authors) coincides with a genetic cline at multiple loci (Dessauer and Cole 1991, Dessauer et al. 2000), suggesting secondary contact following a period of isolation (Barton and

88 Hewitt, 1985). A steep cline involving six color-pattern characteristics is also apparent in the

hybrid zone between *A. t. punctilinealis* and *A. marmoratus* (Dessauer et al., 2000), arguing for

90   the utility of these characters for delineation of *A. tigris* lineages.

In the present study, we focused on defining phylogeographic patterns and interpreting

92   phylogenetic delineations primarily within the range of *A. t. punctilinealis* in Arizona. To the

east, it forms steep morphological/ genetic clines at a hybrid zone with *A. marmoratus* (Dessauer

94   et al., 2000), whereas to the west it exhibits a relatively gradual phenotypic gradient with *A. t.*

tigris (Taylor 1990). Interestingly, both *A. t. septentrionalis* and *A. t. punctilinealis* also form

96   exceptionally gradual morphoclines with *A. t. tigris* that lack apparent ecological underpinnings

(Taylor et al. 1994, Taylor 1990). This stands in contrast to that found between *A. t.*

98   *punctilinealis* and *A. marmoratus* (<2km; Dessauer et al., 2000; Zweifel 1962). Given this

discontinuity, we also examined the association between morphological gradation in the complex

100 and its genetic discontinuity. To do so, we extended the phylogenetic resolution of lineages

within the delineated *A. tigris* complex, then contrasted our phylogenetic patterns with

102 morphological species identifications. We generated taxonomic hypotheses by employing

'species delimitation' algorithms, then employed subsequent downstream analyses to compare

104 the manner by which each model partitioned genetic variation.

106 *The Colorado Plateau as a biogeographic entity*

The Colorado Plateau is a tectonically intact landform (~2000m elevation) with scant

108 internal deformation, thus providing a relatively linear configuration of layered sedimentary rock

for the Plateau interior that contrasts with an eroding margin since Late Miocene (Levander et

110 al., 2011). It also stands in sharp contrast with the more fractured geomorphic features of

neighboring regions: The Rocky Mountains (immediately north and east), Rio Grande Uplift

112   (southeast), and the Basin and Range Province (immediately west) (Fig. 1 of Flowers, 2010). The

uplift of the Plateau and the integration of the Colorado River were defining geomorphic events

114   (Minckley et al., 1986; Spencer et al., 2008).

Prior to integration, the Colorado River emptied into an endorheic basin, most likely

116   following the early path of a major modern tributary (Little Colorado River; House et al., 2008).

In late Miocene-early Pliocene, headwater erosion and basin 'spillover' forged a connection

118   between the river and the Gulf of California via the Salton Trough (Dorsey et al., 2018;

Nicholson et al., 2019; Sarna-Wojcicki et al., 2011). The Gila River system, which had

120   previously drained directly into the Gulf (Eberly & Stanley, 1978) was similarly incorporated.

Other drainages that terminated into closed basins during the Miocene (i.e. Salt and Verde rivers)

122   were also integrated with the Gila River system by late Pliocene (Blakely & Ranney, 2008). The

Pliocene also brought decreased seasonality as well as elevated temperatures and humidity, a

124   climatic regime subsequently reversed by Quaternary glaciation (Thompson, 1991).

Geomorphic events were not only pivotal with regard to biological diversification on the

126   Plateau (Bangs et al., 2020a, 2020b; Douglas et al., 1999; Douglas et al., 2016) but also

provoked endemism as well, in synergy with widespread physiographic and climatic oscillations.

128   These created 'boom-and-bust' demographic cycles as Plateau habitats expanded and contracted

(Douglas et al., 2006), with bookmarks subsequently deposited within the genomes of desert

130   herpetofauna (Douglas et al., 2006; Douglas et al., 2016; Mussmann et al., 2020; Sullivan et al.,

2014). Ancillary genetic footprints of hybridization are also increasingly apparent in modern

132   evaluations (Chafin et al., 2020; Bangs et al., 2020a; Bangs et al., 2020b). Opportunities for

genetic exchange (or alternatively, vicariant events) hinged largely upon drainage evolution (e.g.,

134   Chafin et al., 2020; Douglas et al., 1999). Small-bodied terrestrial vertebrates were also involved

in these biogeographic opportunities, despite the fact that the Colorado River had long been

136   hypothesized as a vicariant barrier to dispersal (Lamb et al., 1992; Pounds and Jackson, 1981).

138   **MATERIALS AND METHODS**

Sampling, amplifying and sequencing of mtDNA

140   We evaluated a total of 239 samples representing five members of the *A. tigris* complex

(*A. marmoratus*, *A. tigris mundus*, *A. t. punctilinealis*, *A. t. septentrionalis*, and *A. t. tigris*), their

142   putative secondary intergrades (e.g., hybrids), plus two out-groups (*Tupinambis rufus* and

Aspidoscelis gularis) (Fig. 1). Genomic DNA was isolated from tail clippings or liver tissue

144   using the Chelex Instagene-matrix extraction protocol (Bio-Rad).

We utilized published primers and PCR protocols (Douglas et al., 2002) to amplify two

146   overlapping mtDNA genes (ATP8 and ATP6). Poor amplification in *A. marmoratus* also forced

the design of species-specific primers (PRIMER3; Untergasser et al., 2012). These were: ATP_F

148   (3-CGCACCCTGATTTATTGTTT-5) and ATP_R (3-GGGTGAAAACGTATGCTTGA-5).

Each 20 µl PCR reaction consisted of 1.5 µl of 1/10x DNA template, 10 µl of HotStarTaq Master

150   Mix (QIAGEN), 0.08 µM of each primer, with the reaction volume finalized with distilled water.

After verification via agarose gel electrophoresis, amplicons were purified using an Exonuclease

152   I/Shrimp Alkaline Phosphatase. Forward and reverse sequencing reactions were performed using

BigDye v3.1 Dye Terminator chemistry (Applied Biosystems, Inc.) and sequenced at the

154   University of Illinois Roy J. Carver Biotechnology Center. Raw sequence chromatograms were

manually curated (SEQUENCHER ver. 5.0, Gene Codes, Ann Arbor, MI) and aligned (MEGA_5;

156   Tamura et al., 2011).

 Phylogenetic inference

To develop initial hypotheses, we first inferred phylogenetic relationships using both

160   Bayesian and distance-based methods. We tested models of nucleotide substitution (N=88-

JMODELTEST; Darriba et al., 2012), using Akaike and Bayesian information criteria (AIC and

162   BIC, respectively). We generated an ultrametric tree using the HKY-model (selected by both

AIC and BIC) with gamma distributed rates and invariant sites (BEAST2; Bouckaert et al.,

164   2014). We applied a Yule model with default parameters and a relaxed log-normal clock process.

MCMC chains were sampled every 500^{th} of 10 million MCMC iterations, following a burn-in

166   period of 2 million. Chain convergence and mixing were visually assessed, and MCMC samples

thinned so as to maintain effective sample sizes >200 while reducing autocorrelation of samples

168   (TRACER; Rambaut and Drummond, 2007). The posterior sampled trees were summarized as a

maximum clade credibility tree with branch lengths representing median heights. For contrast,

170   we also generated a neighbor-joining (NJ) tree using model-corrected distances (R-package APE;

Paradis et al., 2004) in R (R Core Team, 2020), with nodal support assessed via 100 bootstrap

172   pseudo-replicates, as well as a maximum-likelihood tree in PHYML (Guindon et al. 2010). Basic

population summary statistics for major phylogenetic clades (as diagnosed via nodal support and

174   statistical tests of monophyly) were also computed in PEGAS (Paradis, 2010).

176   *Delimiting groups within A. tigris*

We first evaluated the currently accepted taxonomic designations for subspecies within *A.*

178   *tigris* (e.g., Crother et al., 2017; hereby abbreviated as 'TAX'), then employed an array of

analytical approaches to generate six additional delimitation hypotheses. Distance-based

180   clustering methods, used in the context of species delimitation, assume that biological units (i.e.,

species) form non-overlapping genetic clusters. We first employed a recursive method for

182 'Automated Barcode Gap Discovery' (ABGD; Puillandre et al., 2012) that statistically infers a

threshold distance from the data (i.e., the genetic distance at which groups are considered to

184 represent different clusters). We also generated clusters using corrected genetic distances as input

(CD-HIT; Fu et al., 2012), with manually specified distance thresholds of 2% and 5% (hereafter

186 referred to as CD2 and CD5). These hard thresholds were chosen because they represent

approximate 'breakpoints' at which changes occur in the ABGD analysis regarding the number

188 of sampled groups.

We defined phylogenetic species using several approaches: The first, a 'naïve' method,

190 defined groups as the shallowest divergences statistically supported and reciprocally

monophyletic (Rosenberg, 2007), hereafter abbreviated as 'ROS' (SPIDER R package; Brown et

192 al., 2012). A second approach derived groups from tree characteristics under an assumed

speciation process. The first of these methods ['Generalized Mixed Yule Coalescent' (GMYC;

194 Pons et al., 2006)] represents the species boundary as a distinguishable change in branching

patterns within the phylogeny, with independently evolving clusters diagnosed as those at which

196 branching patterns transition from expected under two different models: An assumed

interspecific (Yule process; Nee et al., 1994) and an intraspecific (neutral coalescent; Hudson,

198 1990). In the second ['Poisson Tree Process' (PTP; Zhang et al., 2013)] the probability of

speciation is considered in terms of mutational distance between cladogenesis events (as

200 assumed with a Poisson distribution). PTP requires a tree with branch lengths scaled per

substitutions, and given this, our input represented a tree topologically consistent with that

202 inferred by BEAST but derived instead with PHYML (Guindon et al. 2010). GMYC was run

using the Splits R package (Fujisawa et al., 2013); PTP using the Bayesian implementation

204   provided by the bPTP server (Zhang et al., 2013).

206   *Comparingf delimitations*

        We used 2-way and 3-way Analyses of Molecular Variance (AMOVA; Excoffier et al.

208   1992) with group assignments defined by individual membership under each hypothesis. We

        then used isolation-by-distance (IBD) as a null model under the assumption that the entire

210   complex represented a single species. Our justification stemmed from the pervasive nature of

        IBD with regard to dispersal-limited species (Sexton et al. 2014). In this sense, deviations from

212   the expected IBD pattern are hypothesized as diagnostic of some alternative non-spatial process

        (e.g. reproductive isolation). Partial Mantel tests (Smouse et al. 1986), employing pairwise

214   genetic distances with geographic distances as a covariate (Douglas and Endler, 1982; Meirmans,

        2012), tested the fit of individual assignments under each delimitation hypothesis. Simple and

216   partial Mantel tests (R-package VEGAN; Oksanen et al., 2007) utilized corrected genetic distance

        (APE; Paradis et al., 2004) with confidence assessed using 9,999 permutations. We also employed

218   a stratified Mantel test (per Meirmans, 2012) with permutations constrained within clusters, to

        test if significance of the IBD model is impacted by sampling across structured populations, as

220   delimited by each of our hypotheses.

        We also used a spatially explicit partitioning analysis with genetic variance [measured as

222   a fixation index (F_{CT})] maximized among groups by iteratively rearranging samples into *K*-

        clusters [Spatial Analysis of Molecular Variance (SAMOVA); Dupanloup et al., 2008]. These

224   were executed for *K*-values 2-20, using 100 bootstrapped alignments and replicated without

explicit geographic constraints. We also identified spatial transitions in genetic distances using

226   an interpolation approach [ALLELES-IN-SPACE (AIS); Miller, 2005].

228   **RESULTS**

Phylogenetic and spatial relationships within Aspidoscelis tigris

230   The final 777bp alignment resulted in 120 haplotypes. Bayesian analysis (BEAST2)

yielded strong support for *A. t. tigris + A. t. punctilinealis* (ROS clades A–F, Fig. 3; also see

232   Supplementary Figure S1) as being sister to *A. t. mundus + A. t. septentrionalis* (ROS clade G).

All clades were significantly supported as being reciprocally monophyletic [Rosenberg (2007)

234   test], and all reflected similarly high bootstrap support in the neighbor-joining analysis. The

maximum-likelihood analysis resulted in a tree topologically identical, and was thus not

236   displayed.

All four subspecies formed a monophyletic group sister to a well-defined *A. marmoratus*

238   (clade H). *Aspidoscelis t. septentrionalis* (clade G, excluding *mundus* samples) demonstrates

significantly negative values for both Tajima's and Fu and Li's *D* statistics, with exceptionally

240   low nucleotide diversity and haplotype diversity much less than that recorded for *A. t.*

punctilinealis and *A. t. tigris* (Table 1). These suggest a population expansion following a

242   bottleneck (Douglas et al., 2010). *Aspidoscelis marmoratus* showed similar signatures, although

with a much greater differential between haplotype and nucleotide diversities, and a reduced

244   number of segregating sites relative to sample size (Table 1). Together, *A. t. tigris + A. t.*

punctilinealis were subdivided into five major clades with strong nodal support and reciprocal

246   monophyly (clades A–F, Fig. 3).

Morphologically-diagnosed *A. t. tigris* was polyphyletic, with all samples but one

248   allocated to either clade A or F. The single exception was a morphologically diagnosed *A. t.*

tigris (Nevada) that fell within the *A. t. septentrionalis/mundus* haplogroup (clade G). Clade F,

250   which was >2% divergent from the remaining *A. t. tigris/punctilinealis* clades (A–E), contained

both morphologically consistent *A. t. tigris* (Clark County, southern Nevada), as well as a small

252   subset of *A. t. punctilinealis* (Mohave County, northwest Arizona). All other morphologically

identified *A. t. tigris* from California had mitochondrial haplotypes aligning with clade A of *A. t.*

254   *punctilinealis*. This clade formed a monophyletic grouping in the BA analysis with samples from

southeastern Arizona (clades D and E). Spatially, clades D+E are separated by the mainstem Gila

256   River and defined to the east by the Cochise Filter Barrier that separates Sonoran and

Chihuahuan deserts (Castoe et al. 2007).

258   All clades within *A. t. tigris/punctilinealis* generally showed higher numbers of

segregating sites relative to nucleotide diversity (negative values for Tajima's *D*; clades A–E

260   Table 1). Clades A–D also showed a high rate of singleton haplotypes. Although non-significant

(Table 1), they suggest past population expansions. Within the remainder of the *A. t.*

262   *punctiliealis*, clade A (southwestern Arizona) is sister to a monophyletic group composed of

clade B (Grand Canyon and Colorado River mainstem at juncture between Arizona, Nevada,

264   California), and clade C (central Basin and Range physiographic region of Arizona). The latter is

bounded on the north by the Mogollon Rim and is separated from clade B to the south by the

266   Gila River.

The appearance of clade C haplotypes south of the Gila River mainstem indicates limited

268   gene flow between the two groups, possibly reflecting trans-riverine dispersal as promoted by

low seasonal flows. Similarly, clade B haplotypes are found east and west of the Colorado River

270   mainstem, again suggesting a riverine barrier historically permeable to dispersal, or with

       mitochondrial exchange facilitated by shifts in flow. However, no apparent geographic features

272   support the phylogenetic separation of clades A and C. Their sister relationship is also sustained

       in the NJ analysis, although with a bootstrap support of 56% (compared to 0.87 posterior

274   probability in the Bayesian phylogeny).

276   *Species delimitation models*

               We found considerable variation in the number of proposed clades generated by species

278   delimitation models. Fig. 3B depicts the manner by which each model maps onto the phylogeny,

       using an identification scheme consistent with Fig. 4 and Supplementary Figure S1.

280   Phylogenetically defined clades ('ROS') and a taxonomic hypothesis ('TAX') both collapsed *A.

       t. tigris* and *A. t. punctilinealis* on the basis of extensive intergradation (Taylor et al. 1994).

282   The ABGD analysis was by far the most conservative, collapsing both *A. t. punctilinealis*

       and *A. t. tigris* with *A. t. septentrionalis* + *A. t. mundus*, while only segregating *A. marmoratus*

284   (Fig. 3B). CD-HIT, with an *a priori* sequence divergence threshold of 5%, produced a 4-species

       model (CD5) that separated *A. t. septentrionalis* + *A. t. mundus*, *A. marmoratus*, and *A. t.*

286   *punctilinealis* + *A. t. tigris*. The latter was also partitioned into two groups (i.e., separating clade

       F). One consisted of haplotypes from northwestern Arizona and southern Nevada (clade F; a

288   subset of most morphologically consistent *A. t. tigris*), while the second comprised *A. t.*

       punctilinealis (clades A–E) and remaining *A. t. tigris*, to include those with ranges overlapping

290   clade F. Decreasing the threshold to 2% (i.e., model CD2) then divided *A. t. punctilinealis* into

       eight groups, and five representing *A. t. septentrionalis* (i.e., two from California and three

292   occupying the Colorado Plateau; Fig. 4). GMYC and PTP produced large numbers of groups,

with 13 in the former and 16 in the latter, with many consisting of but a single haplotype.

Partitioning of genetic variance

296   The spatial analysis supported models with fewer species. In the bootstrapped datasets,

F_{CT} overlapped among *K* values (Fig. 5), with the greatest increase in among-group variance

298   (ΔF_{CT}) occurring at *K*=3. Individual assignment at *K*=2 corresponded to the ABGD model (two

species: *A. marmoratus* and *A. tigris*), with three groupings in *K*=3: *A. marmoratus*; *A. t.*

300   *septentrionalis* + *A. t. mundus*; and *A. t. punctilinealis* + *A. t. tigris*.

        A significant correlation was found between matrices of pairwise genetic and geographic

302   distances (Mantel test: r=0.576, P=0.0001; Table 2; Fig. 6). Significant associations only

occurred with the ROS, TAX, ABGD and CD5 models when permutations were limited to those

304   strata defined by delimitation models (i.e., stratified Mantel). The partial Mantel tests with the

ABGD model produced the strongest association (r=0.777), with values generally declining as

306   the number of species increased (Table 2). Correlation of group membership with genetic

distance under all models was greater than in the simple Mantel test for all cases, save for the

308   PTP model (r=0.5754), suggesting that all delimitations better encapsulate spatial genetic

variation than one in which all examined samples are collapsed.

310   Overall, those models partitioning *A. tigris* subspecies were supported by AMOVA

analyses (Supplementary Table S1, S2), with high genetic variance among groups rejecting

312   panmixia. Among-group genetic variance (analogous to F_{CT}) varied between 72–90% and was

maximized in models that recognizing greater numbers of partitions (i.e., CD2, GMYC, and

314 PTP), with 89-90% of variance explained. These values generally declined as models became

more conservative.

316 Interpolating the AIS results with a diffusion model (Fig. 7) produced apparent peaks in

relative pairwise genetic diversity at contact zones between *A. t. punctilinealis* (in this case clade

318 D) and *A. marmoratus* (clade H). These occurred in southeastern Arizona near the shared border

with New Mexico, and in northwestern Arizona, near the contact zone with *A. t. tigris* clade F.

DISCUSSION

322 We used mtDNA sequence data to: Infer phylogenetic and phylogeographic patterns

within the *Aspidoscelis tigris* complex; Generate species delimitation hypotheses therein; and

324 compare model fit using several approaches. We judged how well each model partitioned genetic

variance, both hierarchically and spatially. Our results supported *A. marmoratus* as a distinct

326 species. However, our weak phylogenetic recognition of *A. t. tigris* and *A. t. punctilinealis*, as

well as *A. t. mundus* and *A. t. septentrionalis* questions their subspecific designations, and

328 suggests the need for additional work utilizing larger number of molecular markers to assess

potential intergradation. However, pairwise genetic divergences among the *A. t.*

330 *tigris/punctilinealis* and *A. t. septentrionalis/mundus* groups, regardless of their geographic

proximity (e.g., <10km versus >1000km) may indicate reproductive isolation, again prompting

332 further study that incorporates nuclear markers.

334 *Biogeography of speciation in* Aspidoscelis tigris

Deep relationships within the *A. tigris* complex generally supported the current taxonomy

336 and reflected divisions among biotic regions. Here, we specifically refer to *A. t. septentrionalis*

on the Colorado Plateau, *A. marmoratus* within the Chihuahan Desert, *A. t. punctilinealis* in the

338   Sonoran Desert, and *A. t. tigris* within Mohave and Great Basin deserts. Haplotypes of *A. t.*

mundus, restricted to the California Central Valley, were inferred as monophyletic but nested

340   within a larger *A. t. septentrionalis* clade. This highlights the need for increased sampling,

particularly for *A. t. mundus* and *A. t. tigris,* whose distributions were sampled less thoroughly.

342   Topographically, the separation of *A. t. punctilinealis* and *A. marmoratus* aligns with the

Cochise Filter Barrier (CFB), a biogeographic entity that has similarly been implicated as

344   influential in defining phylogeographic relationships among other taxa (e.g., Castoe et al., 2007;

O'Connell et al., 2017; Morafka, 1977). Pyron et al. (2015) invoked this barrier as a potential

346   driver of so-called 'soft allopatry' (Hickerson and Meyer 2008), in which niche differentiation

drives speciation *in lieu* of a hard barrier to dispersal. Such barriers can often lead to

348   permeability at contact zones, given that divergence is maintained by adaptation rather than

physical separation. As might be expected, a well-characterized hybrid zone separates *A. t.*

350   *punctilinealis* and *A. marmoratus,* replete with steep clines in color polymorphism (Dessauer et

al., 2000). We also identified a distinct east-west divide in mitochondrial haplotypes that is

352   concordant with an hypothesis of soft allopatry.

354   *The potential for hybridization*

We found phylogenetic subdivision less clear among subspecies of *A. tigris* and surmise

356   this likely reflects a cumulative effect of contemporary physiography coupled with historic

processes. *Aspidoscelis t. tigris* and *A. t. punctilinealis* were not distinct. Both transverse the

358   Colorado River, and neither is strictly defined by the Mohave-Sonoran transition. This is perhaps

not surprising, given that morphological intergradation between *A. t. tigris* and *A. t.*

360   *punctilinealis* does not follow a strict ecological gradient, but instead manifests itself as an

exceptionally wide and smooth morphocline (Taylor et al. 1994).

362   *Aspidoscelis t. tigris* from the most northerly part of the range had mitochondrial

sequences aligning instead with *A. t. septentrionalis*, potentially suggesting exchange between

364   the two (Taylor, 1988). Here, we emphasize that increased genetic sampling is necessary so as to

understand the nature of this putative exchange. In contrast, *A. t. punctilinealis* and *A. t.*

366   *septentrionalis* were demarcated according to recognized physiographic provinces, with *A. t.*

septentrionalis occupying the Colorado Plateau and *A. t. punctilinealis* the Sonoran Desert (Fig.

368   4, S1). This was expected, particularly given our understanding of their respective ranges (e.g.

Fig. 1) and associations with the disparate thorn-scrub communities therein (e.g., Zweifel, 1962;

370   Pianka, 1970). We suggest this ecological gradient represents a 'soft boundary,' similar to those

that mediate the transition between *A. t. punctilinealis* and *A. marmoratus*. Unique communities

372   characterize both the Colorado Plateau and Sonoran Desert (e.g., Douglas et al., 2002, 2010), and

this suggests a common, underlying process driving diversification therein (Riddle and Hafner,

374   2006; Pianka, 1967).

376   *The validity of taxonomic designations*

Genetic delineation of *A. t. tigris* was inconsistent with traditional taxonomy, in that

378   specimens allocated morphologically to *A. t. tigris* were not monophyletic within the

phylogenetic analyses, nor were those groupings consistent with the historically accepted range

380   boundaries for *A. t. tigris* and *A. t. punctilinealis*. Our expectations were that haplotypes sampled

from the Great Basin and Mohave deserts (i.e., those regions encompassing *A. t. tigris*) would be

382   monophyletic. Instead, we found a closer alignment with *A. t. septentrionalis* to the northeast and

A. t. punctilinealis to the southeast. This is likely a result of mitochondrial exchange among these

384   subspecies, again highlighting the need for a genomic evaluation.

Taylor et al. (1994) demonstrated the existence of a wide, smooth morphocline >180 km

386   in width that traversed the Colorado River and separated *A. t. tigris* from *A. t. punctilinealis*

along the southern extent of their contact (Fig. 1A). This stands in contrast to a 16 km-wide

388   morphocline found at the northern extent of their contact, and that discrepancy begs an obvious

ecological explanation (Taylor 1990). The northern morphocline coincides with the Arizona-

390   California-Nevada nexus, a hotspot for mitochondrial diversity as well, where sub-clades overlap

without clear physical boundaries (Fig. 4, S1). In addition, multiple clades (A and F) traverse the

392   Colorado River, with the highly divergent clade F (found primarily within morphologically

defined *A. t. tigris*) possibly representing a more widespread Mohave clade.

394   One explanation for the above variation in clinal width evokes an historic perspective

wherein the northern contact between *A. t. tigris* and *A. t. punctilinealis* is secondary, whereas

396   range overlap to the south was maintained. This would generate morphoclines of varying ages

and would offer one potential explanation for the variance in mitochondrial divergence (i.e.,

398   elevated to the north, nonexistent to the south), as well as morphocline width. Here, we posit that

the duration and extent of contact may be a driving mechanism. A related hypothesis would be

400   that the width of the cline is in response to an underlying environmental gradient, such as the

transition between Sonoran and Mohave habitats. However, neither hypothesis can be tested with

402   the data presented herein.

An additional possibility is the fluctuating capacity of the Colorado River as a barrier to

404   dispersal. We offer scant evidence for it serving as a vicariant structure, a result corroborated by

studies of other reptiles (e.g., desert tortoise: Lamb et al., 1989; iguanids: Lamb et al., 1992).

406   Although the Colorado River does seemingly constrain dispersal (Pounds and Jackson 1981),

gene flow could still be facilitated by the periodic fluctuations in its trajectory and flow. In fact,

408   the shallow divergences found among clades transecting the river provide evidence for how

recent these events were.

Climate driven population oscillations

412   One unexpected finding was the exceptional genetic divergences among *A. t.*

punctilinealis clades (A–E). One possible explanation evokes fluctuating habitat availability over

414   time, rather than contemporary features of the landscape. Concordance in relative crown ages

among the majority of *A. t. punctilinealis* clades (A–E; Fig. 1B) could suggest that intraspecific

416   divergence of these groups shares a common cause. Douglas et al. (2006) offered an applicable

model that explained genetic structuring and population expansion in four desert rattlesnake

418   species as a 'boom-and-bust' demographic cycle, caused by historical climate fluctuations.

Under this model, a 'refugium phase' (='bust') condensed continuous distributions into smaller,

420   allopatric populations such that restricted gene flow promoted genetic drift as manifested by

reduced effective population size (N_e). As climate shifted, desert habitat expanded such that

422   ecological barriers separating refugia were removed, allowing distributions to expand (='boom').

These oscillating contractions and expansions placed characteristic molecular signatures

424   within populations, most prominently as a loss of genetic diversity and an exaggerated

differentiation among sub-populations post-expansion (Hewitt 2000). This is reflected in an

426   increased number of low-frequency haplotypes, and low nucleotide diversity among

contemporary haplotypes (Table 1). We also noted similar patterns in *A. t. septentrionalis* and *A.*

428   *marmoratus*, suggesting bottleneck-expansion events within these taxa as well.

The effects of the 'refugium phase' can be seen when examining modern short-range

430   endemics [i.e., those with distributions easily contained within a 100x100 km grid (i.e., <10,000

km^2]. An example would be the isolated New Mexico Ridge-nosed Rattlesnake (*Crotalus*

432   *willardi obscurus*) that reflects signals of relatively recent divergence and population reduction

(Holycross and Douglas 2007). Its historic connectivity in the Sky-islands has been truncated by

434   contemporary climate change, whereas its refugium phase, as gauged by demographic and niche

modeling, underscored a continued loss of habitat (Davis et al. 2015). We suspect a similar

436   refugium-expansion model would explain the extant genetic differentiation found in the present

study.

Comparing species delimitation approaches

440   We found considerable variance in the number of groups delimited using the various

algorithmic methods, especially when contrasted with the current taxonomy (Fig. 3). This could

442   reflect the capacity of the various models to diagnose cryptic lineages, or alternatively, suggest

the data violate implicit assumptions (Carstens et al. 2013). Phylogenetic models (i.e., ROS,

444   PTP, GMYC) generally produced the largest numbers of groups, but did not explain the

geographic patterns of genetic variation as well as did other models (Table 2; Figure 3). All

446   models, however, captured greater genetic variance among groups (i.e., 72-90% in AMOVA)

than within-group (10-28%), indicating the recognition of an authentic biological signal (Table

448   S1, S2).

Species delimitation employing the coalescent, such as GMYC, have previously been

450   criticized for a tendency to over-split when applied to structured populations (Lohse 2009). This,

in turn, suggests that a single-threshold delimitation model may be inappropriate for use with

452   hierarchically structured data (Talavera et al. 2013). Strong intraspecific structure in

Aspidoscelis, similar to that found in other desert reptiles (Holycross and Douglas 2007, Douglas

454   et al. 2002), may explain the over-fitting of GMYC. A more comprehensive test using additional

scenarios is warranted, as this will more appropriately characterize those biases that can emerge

456   when the method is applied to empirical data.

        We also employed a second phylogenetic method (PTP) that essentially modeled

458   speciation as a clock-like process, with its probability of occurrence represented as a function of

time (i.e., branch length). Given this, any aspect that would promote rate heterogeneity could be

460   sufficient to violate the model, such as the extreme demographic fluctuations suspected for

Aspidoscelis populations.

462   Clustering-based methods, on the other hand, may be biased by elevated genetic distances

or incomplete sampling, with group delimitation being driven entirely by the arbitrary nature of

464   the (user-selected) threshold. The ABGD model attempts to mitigate this by automatically

detecting distance-based partitions, as bounded by priors that account for maximum allowable

466   intraspecific divergence (Puillandre et al. 2012). ABGD and clustering methods appeared to

delimit groups that more appropriately represented the genetic/ geographic distances paradigm.

468   When examining genetic distances defined by these methods within- versus among-group, we

found *A. marmoratus* approximately equal to *A. tigris* with regard to genetic distances, regardless

470   of whether *A. tigris* was intact or partitioned (Fig. 6). Among-group genetic distances were equal

regardless of how proximate were the samples (i.e., genetic x geographic distance slope

472   approaches zero). This suggests that divergence is maintained not by physical distance but

instead by some intrinsic qualities possessed by those groups, such as reproductive isolation or

474   mutually exclusive habitat requirements (Fig. 6). In this case, near-zero slopes in the among-

group comparisons lend some validity to a species-level designation for *A. marmoratus* and

476   suggested the potential for reproductive isolation within the *tigris* complex (i.e., between *A. t.*

tigris/*punctilinealis* and *A. t. septentrionalis*; Fig. 6B). However, reduced sample sizes for *A. t.*

478   *mundus* and *A. t. tigris* remain an issue. Within-group comparisons resulted in a positive

relationship with geographic distance, as was expected (Fig. 6).

Species concepts and taxonomic conclusions for Aspidoscelis

482   Species concepts present a philosophical quandary (Coyne and Orr, 2004). For example,

many modern interpretations [i.e., biological (BSC; Mayr, 1942); evolutionary (ESC; Frost and

484   Hillis, 1990); phylogenetic (PSC; Cracraft, 1983)] offer a mechanistic or philosophical

interpretation for the existence of species, yet do not themselves present an explicit operational

486   criterion that can universally define species. The definitions also have fundamental conflicts with

regards to the secondary characteristics each relies upon (e.g., reproductive isolation,

488   morphological diagnosability; de Queiroz, 2007).

Given that inferring a direct relationship between genotypic (or 'haplotypic') clusters is a

490   questionable approach, we instead relied upon a 'theory-independent' definition (the 'genotypic

cluster' concept: Douglas et al., 2007; Sullivan et al., 2014) wherein clusters are defined as '…

492   distinguishable groups of individuals that have few or no intermediates when in contact' (Mallet,

1995). We used this basis to compare among delimitation hypotheses for *Aspidoscelis tigris*, and

494   how they may (or may not) extend to theoretical species definitions such as the BSC.

An expectation of the '… little to no intermediates in contact' criterion is that individuals

496   from separate genotypic clusters should not be any more genetically similar when sampled in

proximity than when sampled from more geographically disparate locations. Such a relationship

498   would suggest continuous gene flow limited only by the ability of individuals to disperse (i.e.,

'isolation-by-distance' *sensu* Wright, 1940, 1943).

500   We tested the applicability of this criterion to our species hypotheses using spatial and

non-spatial analyses of molecular variance (AMOVA/ SAMOVA), by testing the manner by

502   which each partitioning scheme explained spatial signatures of genetic differentiation (IBD/

Mantel), and with qualitative comparisons of landscape-level genetic differentiation (AIS

504   interpolation). We found the ABGD model (i.e., separation of *A. marmoratus* from all *A. tigris*

subspecies) did well with partitioning spatial haplotypic variation, with *A. marmoratus* being

506   highly differentiated (>10% sequence divergence) even from localities sampled <2km from *A. t.*

punctilinealis (Fig. 6A). These patterns are consistent with many commonly applied species

508   concepts (e.g. reproductive isolation; per BSC), and given this, the elevated status of *A.*

marmoratus is sustained.

510   We also found limited (albeit, existing) exchange between two major clades within the

remaining *A. tigris*: *A. t. tigris/punctilinealis* versus *A. t. septentrionalis/mundus* (i.e., TAX

512   model). Here, genetic differentiation was again uncorrelated with spatial distances (Fig. 6B),

possibly implicating intrinsic or extrinsic factors that may satisfy species-level criteria. Previous

514   estimates place divergence of these groups to be no greater than 5-10 kya, based upon

availability of suitable habitat following the Wisconsin glacial period (Zweifel, 1962). If so, then

516   our observed mean corrected genetic distance of 0.058 equates to a mutation rate of 5.8

substitutions per million years, a rate considerably greater than the expected rate of ~0.0045 for

518   ATPase 6, the more rapidly evolving of our two mitochondrial genes (Kumar, 1996; Mueller,

2006).

Although we recognize that numerous factors contribute to mtDNA rate heterogeneity, our rudimentary calculation suggests that subspecies within *A. tigris* may have diverged much

earlier than previously thought. Although our results support the continued recognition of *A. t. septentrionalis* as distinct, we deem them insufficient to propose elevation beyond a subspecific

level. Instead, we argue for greater sampling at both geographical and genomic scales to properly adjudicate taxonomies within *A. t. tigris*. The latter, while enabling tests of hybridization/

introgression within a more statistically robust framework (e.g., Chafin et al., 2019), would also open avenues for novel species delimitation algorithms to be formulated using unsupervised

machine learning approaches (Derkarabetian et al., 2019; Martin et al., 2020; Mussmann et al., 2020).

The CD5 model identified potential cryptic lineages within a clade comprising *A. t. tigris* and *A. t. punctilinealis*, although sample sizes representing this group were limited. Of note, *A. t.*

tigris and *A. t. punctilinealis* were not reciprocally monophyletic, however assigning causality of this to hybridization/ intergradation, stochastic lineage sorting, or a lack of phylogenetic signal

requires additional data (Chafin et al., 2020). Again, we suggest a more in-depth examination using a larger panel of molecular markers so as to confirm or reject continued recognition of *A. t.*

tigris as distinct from *A. t. punctilinealis*, and likewise to adjudicate recognition of *A. t. mundus* from *A. t. septentrionalis*.

Conclusion

540   We found that algorithmic species delimitation methods vary substantially in the manner by which they partition hierarchical phylogenetic diversity, particularly when contrasted against

542   the respected overlays of historical biogeography, morphology, ecology, and prior genetic

evaluations. By this we mean that some algorithms (e.g., ABGD) yielded groupings which

544   support external evidence of divergence at the 'species' level (e.g., according to a BSC-like

interpretation), while others (e.g., CD5) yield groupings that could potentially fit various

546   operational species concepts, yet require further evaluation using genomic data. Others (GMYC,

PTP) seemingly delimit populations, a conclusion in line with prior evaluations of simulated and

548   empirical datasets. The latter was hypothesized to be impacted by inflated levels of inter-

population divergence, a phenomenon which has been observed in other co-occuring desert

550   fauna subjected to the same historical climatic fluctuation. These observations prompt us to

support the elevation of *A. marmoratus*, but additional groups withinthe *A. tigris* complex will

552   requiring additional data to properly adjudicate.

ACKNOWLEDGMENTS

556   Samples were collected by BKS and JEC under authority of permits provided by Arizona and New

Mexico Game and Fish departments. Collecting methods were approved as part of IACUC protocols for

558   surveying and vouchering lizards (BKS and JEC). We are also indebted to students, post-doctorals, and

faculty who have promoted this research: M. R. Bangs, L. Graham, B. A. Levine, B. T. Martin, S. M.

560   Mussmann, Z. D. Zbinden, and especially K.E. Harrell. This research was made possible through

generous endowments to the University of Arkansas: A Doctoral Dissertation Fellowship (TKC), the

562   Bruker Professorship in Life Sciences (MRD), and the Twenty-First Century Chair in Global Change

Biology (MED), all of which provided salaries and/or research funds for this work. Additional analytical

564   resources were provided by the Arkansas Economic Development Commission (Arkansas Settlement

Proceeds Act of 2000) and the Arkansas High Performance Computing Center (AHPCC), and from an

566   NSF-XSEDE Research Allocation (TG-BIO160065) to access the Jetstream cloud. Summer support of

field work for JEC was derived from consecutive Endowed Professorships in Science by Opelousas

568   General Hospital of Opelousas, Louisiana.

570   **LITERATURE CITED**

Bangs, M.R., Douglas, M.R., Brunner, P.C., & Douglas, M.E. (2020a) Reticulate evolution as a
572   management challenge: Patterns of admixture with phylogenetic distance in endemic fishes
         of western North America. Evolutionary Applications, 13, 1400-1419.

574   Bangs, M.R., Douglas, M.R., Chafin, T.K., & Douglas, M.E. (2020b) Gene flow and species
         delimitation in fishes of Western North America: Flannelmouth (*Catostomus latipinnis*) and
576   Bluehead sucker (*C. Pantosteus discobolus*). Ecology and Evolution, 10 6477-6493.

Barley, A.J., Nieto-Montes de Oca, A., Reeder, T.W., Manríquez-Morán, N.L., Arenas Monroy,
578   J.C., Hernández-Gallegos, O. & Thomson, R.C. (2019) Complex patterns of hybridization
         and introgression across evolutionary timescales in Mexican whiptail lizards (Aspidoscelis).
580   Molecular Phylogenetics and Evolution, 132, 284–295.

Barton, N.H. & Hewitt, G.M. (1985) Analysis of hybrid zones. Annual Review of Ecology and
582   Systematics, 16, 113–148.

Bhattarachya, T., Tierney, J.E., Addison, J.A., & Murray, J.W. (2018) Ice-sheet modulation of
584   deglacial North American monsoon intensification. Nature Geoscience, 11, 848-852.

Blakely, R., & Ranney, W. (2008) Ancient Landscapes of the Colorado Plateau. Grand Canyon
586   Association, Grand Canyon, AZ.

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., Suchard, M.A.,
588   Rambaut, A. & Drummond, A.J. (2014) BEAST 2: A software platform for Bayesian
         evolutionary analysis. PLoS Computational Biology, 10, e1003537.

590   Brown, S.D.J, Collins, R.A., Boyer, S., Lefort, M., Malumbres-Olarte, J., Vink, C.J., &
         Cruickshank, R.H. (2012) Spider: An R package for the analysis of species identity and
592   evolution, with particular reference to DNA barcoding. Molecular Ecology Resources,
         12, 562-565.

Carstens, B.C., Pelletier, T. a., Reid, N.M. & Satler, J.D. (2013) How to fail at species
596   delimitation. Molecular Ecology, 22, 4369–4383.

Castoe, T.A., Spencer, C.L. & Parkinson, C.L. (2007) Phylogeographic structure and historical

598   demography of the western diamondback rattlesnake (*Crotalus atrox*): A perspective on North American desert biogeography. Molecular Phylogenetics and Evolution, 42, 193–
600   212.

Chafin, T.K., Douglas, M.R., Bangs, M.R., Martin, B.T., Mussmann, S.M., & Douglas, M.E.
602   (2020) Taxonomic uncertainty and the anomaly zone: Phylogenomics disentangle a rapid radiation to resolve contentious species (*Gila robusta* complex) in the Colorado River.
604   bioRxiv, 692509.

Cracraft, J. (1983) Species concepts and speciation analysis. In: Current ornithology (ed. by
606   Ketterson, E., & Nolan, V.), pp. 159-187, Springer, New York, NY.

Crother, B.I. (ed) (2017) Scientific and standard english names of amphibians and reptiles of
North America north of Mexico, with comments regarding confidence in our understanding 8th edition. Herpetological Circular, 43, 1–102.

Darriba, D., Taboada, G.L., Doallo, R. & Posada, D. (2012) jModelTest 2: more models, new heuristics and parallel computing. Nature Methods, 9, 772–772.

Davis, M.A., Douglas, M.R., Webb, C.T., Collyer, M.L., Holycross, A.T., Painter, C.W., Kamees, L.K. & Douglas, M.E. (2015) Nowhere to go but up: Impacts of climate change on
demographics of a short-range endemic (*Crotalus willardi obscurus*) in the sky-islands of southwestern North America. PLoS ONE, 10, 1–20.

Dessauer, H.C. & Cole, C.J. (1991) Genetics of whiptail lizards (Reptilia: Teiidae: *Cnemidophorus*) in a hybrid zone in southwestern New Mexico. Copeia, 1991, 622–637.

Dessauer, H.C., Cole, C.J. & Townsend (2000) Hybridization among western whiptail lizards (*Cnemidophorus tigris*) in southwestern New Mexico: Population genetics, morphology,
and ecology in three contact zones. Bulletin of the American Museum of Natural History, 246, 1–148.

Dorsey, R.J., O'Connell, B., McDougall, K., Homan, M.B. (2018) Punctuated sediment discharge during Early Pliocene birth of the Colorado River: Evidence from regional
stratigraphy, sedimentology, and paleontology. Sedimentary Geology, 363, 1-33.

Douglas, M.E., Douglas, M.R., Schuett, G.W., Beck, D.D. & Sullivan, B.K. (2010) Conservation phylogenetics of helodermatid lizards using multiple molecular markers and a supertree
approach. Molecular Phylogenetics and Evolution, 55, 153–167.

Douglas, M.E., Douglas, M.R., Schuett, G.W. & Porras, L.W. (2006) Evolution of rattlesnakes
630   (Viperidae; *Crotalus*) in the warm deserts of western North America shaped by Neogene vicariance and Quaternary climate change. Molecular Ecology, 15, 3353–3374.

632   Douglas, M.E., Douglas, M.R., Schuett, G.W., Porras, L.W. & Holycross, A.T. (2002)
Phylogeography of the western rattlesnake (*Crotalus viridis*) complex, with emphasis on the
634   Colorado Plateau. In: Biology of the Vipers (ed. by Schuett, G.W., Höggren, M., Douglas,
M.E., & Greene, H.W.), pp. 11-50, EEagle Mountain Publishing, Eagle Mountain, UT.

636   Douglas, M.E., Minckley, W.L., & DeMarais, B.D. (1999) Did vicariance mold phenotypes of
Western North American fishes? Evidence from Gila River cyprinids. Evolution, 53, 238-
638   246.

Douglas, M.R., Davis, M.A., Amarello, M., Smith, J.J., Schuett, G.W., Herrmann, H.W.,
640   Holycross, A.T., & Douglas, M.E. (2016) Anthropogenic impacts drive niche and
conservation metrics of a cryptic rattlesnake on the Colorado Plateau of western North
642   America. Royal Society Open Science, 3, 160047.

Drummond, C.S. & Hamilton, M.B. (2007) Hierarchical components of genetic variation at a
644   species boundary: population structure in two sympatric varieties of *Lupinus microcarpus*
(Leguminosae). Molecular Ecology, 16, 753–769.

646   Dupanloup, I., Schneider, S. & Excoffier, L. (2008) A simulated annealing approach to define
the genetic structure of populations. Molecular Ecology, 11, 2571–2581.

648   Eberly, L.D., & Stanley, T.B. (1978) Cenozoic stratigraphy and geologic history of southwestern
Arizona. Geological Society of America Bulletin, 89, 921-940.

650   Excoffier, L., Smouse, P.E. & Quattro, J.M. (1992) Analysis of molecular variance inferred from
metric distances among DNA haplotypes: application to human mitochondrial DNA
652   restriction data. Genetics, 131, 479–491.

Flowers, R.M. (2010) The enigmatic rise of the Colorado Plateau. Geology, 38, 671-672.
654

Frost, D.R. & Hillis, D.M. (1990) Species in concept and practice : Herpetological applications.
656   Herpetologica, 46, 86–104.

Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. (2012) CD-HIT: Accelerated for clustering the next-
658   generation sequencing data. Bioinformatics, 28, 3150–3152.

Fujisawa, T. & Barraclough, T.G. (2013) Delimiting species using single-locus data and the
660   Generalized Mixed Yule Coalescent Approach: A revised method and evaluation on
simulated data sets. Systematic Biology, 62, 707-724.
662

Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W. & Gascuel, O. (2010) New
664   algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the
performance of PhyML 3.0. Systematic Biology, 59, 307–321.

666   Hewitt, G.M. (2000) The genetic legacy of the Quaternary ice ages. Nature, 405, 907–913.

Hickerson, M.J. & Meyer, C.P. (2008) Testing comparative phylogeographic models of marine
668   vicariance and dispersal using a hierarchical Bayesian approach. BMC Evolutionary
        Biology, 8, 1–18.

670   Holycross, A.T. & Douglas, M.E. (2007) Geographic isolation, genetic divergence, and
        ecological non-exchangeability define ESUs in a threatened sky-island rattlesnake.
672   Biological Conservation, 134, 142–154.

House, P.K., Pearthree, P.A., & Perkins, M.E. (2008) Stratigraphic evidence for the role of lake
674   spillover in the inception of the lower Colorado River in southern Nevada and western
        Arizona. Special Paper of the Geological Society of America, 439, 335-353.

676   Kumar, S. (1996) Patterns of nucleotide substitution in mitochondrial protein coding genes of
        vertebrates. Genetics, 143, 537–548.

678   Lamb, T., Avise, J.C. & Gibbons, J.W. (1989) Phylogeographic patterns in mitochondrial-DNA
        of the desert tortoise (*Xerobates agassizi*), and evolutionary relationships among the north-
680   american gopher tortoises. Evolution, 43, 76–87.

Lamb, T., Jones, T.R. & Avise, J.C. (1992) Phylogeographic histories of representative
682   herpetofauna of the southwestern U.S.: mitochondrial DNA variation in the desert iguana
        (*Dipsosaurus dorsalis*) and the chuckwalla (*Sauromalus obesus*). Journal of Evolutionary
684   Biology, 5, 465–480.

Levander, A., Schmandt, B., Miller, M.S., Liu, K., Karlstrom, K.E., Crow, R.S., Lee, C-T.A.,
686   Humphreys, E.D. (2011) Continuing Colorado Plateau uplift by delamination-style
        convective lithospheric downwelling. Nature, 472, 461-466.
688

Lohse, K. (2009) Can mtDNA barcodes be used to delimit species? A response to Pons et al.
690   (2006). Systematic Biology 58, 439–442.

Martin, B.T., Chafin, T.K., Douglas, M.R., Placyk Jr., J.S., Birkhead, R.D., Phillips, C.A., &
692   Douglas, M.E. (2020) Species delimitation using machine learning recovers a
        phylogenomically consistent classification for North American box turtles (*Terrapene* spp.)
694   bioRxiv, 205039.

Mayr, E. (1942) Systematics and the Origin of Species, Columbia University Press, New York,
696   NY.

Meirmans, P.G. (2012) The trouble with isolation by distance. Molecular Ecology, 21, 2839–
698   2846.

Miller, M.P. (2005) Alleles In Space (AIS): Computer software for the joint analysis of
700   interindividual spatial and genetic information. Journal of Heredity, 96, 722–724.

Minckley, W.L., Hendrickson, D.A., & Bond, C.E. (1986) Geography of western North
702   American freshwater fishes: Description and relations to intracontinental tectonism. In:
         Zoogeography of North American Freshwater Fishes (ed. by Hocutt, C.H., & Wiley, E.O).
704   pp. 519-613. John Wiley & Sons, New York, NY.

Mueller, R.L. (2006) Evolutionary rates, divergence dates, and the performance of mitochondrial
706   genes in Bayesian phylogenetic analysis. Systematic Biology, 55, 289–300.

Mussmann, S.M., Douglas, M.R., Oakey, D.D., & Douglas, M.E. (2020) Defining relictual
708   biodiversity: Conservation units in speckled dace (Leuciscidae: *Rhinichthys osculus*) of the
         Greater Death Valley ecosystem. Ecology and Evolution, Early View.

710 Nicholson, U., Carter, A., Robinson, P., Macdonald, D.I.M. (2019) Eocene–Recent drainage
         evolution of the Colorado River and its precursor: An integrated provenance perspective
712   from SW California. Geological Society Special Publications, 488, 47-72.

714 O'Connell, K.A., Streicher, J.W., Smith, E.N. & Fujita, M.K. (2017) Geographical features are
         the predominant driver of molecular diversification in widely distributed North American
716   whipsnakes. Molecular Ecology, 26, 5729–5751.

Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J. & Suggests,
718   M. (2007) The vegan package. Community ecology package, 10, 631–637.

Paradis, E., Claude, J. & Strimmer, K. (2004) APE: Analyses of phylogenetics and evolution in
720   R language. Bioinformatics, 20, 289–290.

Pianka, E.R. (1970) Comparative autecology of the lizard *Cnedimophorus tigris* in different parts
722   of its geographic range. Ecology, 51, 703–720.

Pianka, E.R. (1967) On lizard species diversity: North American flatland deserts. Ecology, 48,
724   333–351.

Pons, J., Barraclough, T.G., Gomez-Zurita, J., Cardoso, A., Duran, D.P., Hazell, S., Kamoun, S.,
726   Sumlin, W.D. & Vogler, A.P. (2006) Sequence-based species delimitation for the DNA
         taxonomy of undescribed insects. Systematic Biology, 55, 595–609.

728 Pounds, J.A. & Jackson, J.F. (1981) Riverine barriers to gene flow and the differentiation of
         fence lizard populations. Evolution, 35, 516–528.

730 Puillandre, N., Lambert, A., Brouillet, S. & Achaz, G. (2012) ABGD, automatic barcode gap

discovery for primary species delimitation. Molecular Ecology, 21, 1864–1877.

Pyron, R.A., Costa, G.C., Patten, M.A. & Burbrink, F.T. (2015) Phylogenetic niche conservatism and the evolutionary basis of ecological speciation. Biological Reviews, 90, 1248–1262.

de Queiroz, K. (2005) A unified species concept and its consequences for the future of taxonomy. Proceedings of the California Academy of Sciences, 56, 196–215.

R Core Team (2020) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Rambaut, A. & Drummond, A.J. (2007) Tracer v1.4. Digital resource available at http://tree.bio.ed.ac.uk/software/tracer/

Riddle, B.R. & Hafner, D.J. (2006) A step-wise approach to integrating phylogeographic and phylogenetic biogeographic perspectives on the history of a core North American warm
deserts biota. Journal of Arid Environments, 66, 435–461.

Rosenberg, N.A. (2007) Statistical tests for taxonomic distinctiveness from observations of
744 monophyly. Evolution, 61, 317–323.

Sarna-Wojcicki, A.M., Deino, A.L., Fleck, R.J., McLaughlin, R.J., Wagner, D., Wan, E., Wahl,
746 D., Hillhouse, J.W., & Perkins, M. (2011) Age, composition, and areal distribution of the Pliocene Lawlor Tuff, and thee younger Pliocene tuffs, California and Nevada. Geosphere,
748 7, 599-628.

Sexton, J.P., Hangartner, S.B. & Hoffmann, A.A. (2014) Genetic isolation by environment or
750 distance: Which pattern of gene flow is most common? Evolution, 68, 1–15.

Smouse, P.E., Long, J.C. & Sokal, R.R. (1986) Multiple regression and correlation extensions of
752 the Mantel test of matrix correspondence. Systematic Zoology, 35, 627.

Spencer, J.E., Smith, G.R., & Dowling, T.E. (2008) Middle to Late Cenozoic geology,
754 hydrography, and fish evolution in the American Southwest. Geolocial Society of America, 80301.

756 Talavera, G., Dincă, V. & Vila, R. (2013) Factors affecting species delimitations with the GMYC model: Insights from a butterfly survey. Methods in Ecology and Evolution, 4,
758 1101–1110.

Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011) MEGA5:
Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. Molecular Biology and Evolution, 28, 2731–2739.

Taylor, H.L. (1988) A morphological analysis of intergredation between the teiid lizards *Cnemidophorus tigris tigris* and *C. tigris septentrionalis*. Herpetologica, 44, 176–185.

Taylor, H.L. (1990) A morphological analysis of the teiid lizards *Cnemidophorus tigris tigris* and *C. tigris gracilis* from a contact zone in northwestern Arizona. Herpetologica, 46, 447–456.

Taylor, H.L., Beyer, C., Harris, L. & Pham, H. (1994) Subspecific relationships in the Teiid lizard *Cnemidophorus tigris* in southwestern Arizona. Journal of Herpetology, 28, 247–253.

Thompson, R.S. (1991) Pliocene environments and climates in the western United States. Quaternary Science Review, 10, 115-132.

Tucker, D.B., Colli, G.R., Giugliano, L.G., Hedges, S.B., Hendry, C.R., Lemmon, E.M., Lemmon, A.R., Sites Jr., J.W., & Pyron, R.A. (2016) Methodological congruence in
phylogenomic analyses with morphological support for teiid lizards (Sauria: Teiidae). Molecular Phylogenetics and Evolution, 103, 75-84.

Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B.C., Remm, M. & Rozen, S.G. (2012) Primer3-new capabilities and interfaces. Nucleic Acids Research, 40, 1–12.

Wright, S. (1940) Breeding structure of populations in relation to speciation. The American Naturalist, 74, 232–248.

Wright, S. (1943) Isolation by distance. Genetics, 28, 114–138.

Zhang, J., Kapli, P., Pavlidis, P. & Stamatakis, A. (2013) A general species delimitation method
780   with applications to phylogenetic placements. Bioinformatics, 29, 2869–2876.

Zweifel, R.G. (1962) Analysis of hybridization between two subspecies of the desert whiptail
782   lizard, *Cnemidophorus tigris*. Copeia, 1962, 749–766.

Figures and Tables

Table 1: Population summary statistics for clades within the *Aspidoscelis tigris* complex as determined by sequence analysis of 777bp of the ATPase 6 and 8 genes across 239 samples. Abbreviations are as follows: H = haplotype diversity; π = nucleotide diversity; θ = population mutation rate; θ_s/N = number of segregating sites / clade sample size; H/N = number of haplotypes / clade sample size. *A. t. mundus* was excluded due to low sample size.

CLADE	TAJIMA D	TAJIMA D P-VALUE	H	π	θ	FU & LI F	FU & LI D	θ_s/N	H/N
A	-1.522	0.128	0.993	0.013	15.409	-1.680	-1.595	2.889	0.944
B	-0.938	0.348	0.969	0.011	11.258	-0.442	-0.356	2.833	0.916
C	-1.628	0.103	0.997	0.012	17.033	-2.051	-1.935	2.461	0.923
D	-1.889	0.059	0.966	0.018	26.468	-2.968	-3.008	3.885	0.654
E	-0.524	0.600	1.000	0.013	10.363	0.050	0.524	4.75	1.000
F	0.616	0.538	1.000	0.021	14.890	0.660	0.374	5.667	1.000
G	-2.114	0.017	0.478	0.006	15.600	-2.609	-2.123	1.123	0.292
H	-1.778	0.075	0.828	0.002	3.762	0.457	-2.933	0.4	0.425

Table 2: Comparison of alternative species delimitation models according to the fit of the data under an isolation-by-distance model. Results are shown for 2-way, stratified, and partial Mantel tests, reporting a correlation coefficient (*r*) between distance matrices composed of geographic and genetic distances. Correlation coefficients in the partial Mantel tests reflect association between clade membership and genetic distance, with geographic distances as a covariate. Values indicated by (**) are significant at p>0.0001. Species delimitation models are abbreviated as: ABGD=Automated Barcode-Gap Discovery; TAX=qualitative hypothesis derived from current taxonomy; CD5=CD-Hit with 5% divergence threshold for clustering; CD2= CD-Hit with 2% divergence threshold for clustering; GMYC=Generalized mixed Yule coalescent; PTP=Poisson tree process; ROS=Clade delimitation based on Rosenberg test of reciprocal monophyly.

MODEL	2-WAY MANTEL TEST	STRATIFIED 2-WAY MANTEL	PARTIAL MANTEL
	Mantel *r*	Mantel *r*	Mantel *r*
ALL	0.5762**	-	-
ROS	-	0.5762**	0.6317**
ABGD	-	0.5762**	**0.7770****
TAX	-	0.5762**	0.7118**
CD5	-	0.5762**	0.7025**
CD2	-	0.5762	0.5816**
GMYC	-	0.5762	0.5818**
PTP	-	0.5762	0.5754**

**Significant at α=0.0001

Figure 1: Range map and sampling sites for *Aspidoscelis marmoratus* and subspecies of *A. tigris* (subspecific units within *A. marmoratus* and Mexican subspecies of *A. tigris* not shown). Three identified contact zones are depicted between: (A) *A. t. tigris* and *A. t. punctilinealis*, roughly transected by the Colorado River with morphoclines varying from 16km in the north to ~180km in the south; (B) *A. t. tigris* x *A. t. septentrionalis*, where color-pattern intergradation corresponds to a ~50km transition from Mohave (inset; M) to Great Basin (GB) desert-scrub communities to the Colorado Plateau ecoregion (CP); and (C) *A. t. punctilinealis* x *A. marmoratus*, a hybrid zone of <2km in width traversing the Cochise Filter Barrier transition zone between the Sonoran (inset; S) and Chihuahuan (Ch) deserts. Note that both sample points and ranges are colored by subspecies, as defined in the legend. Map based on observations and prior work by BKS.

Figure 2: Pattern polymorphism in the *Aspidoscelis tigris* group: (A–B) *Aspidoscelis tigris mundus*; (C) *A. t. septentrionalis* from northern Arizona; (D) *A. t. tigris* from California; (E–F) *A. t. punctilinealis* from central Arizona; (G–H) *A. marmoratus* from New Mexico; (I) Dorsal and (J) ventral view of suspected *A. t. tigris* x *A. t. punctilinealis* hybrids. Pictures provided by BKS.

Figure 3: Neighbor-joining (A) and Bayesian (B) phylogenetic trees based on sequence analysis of 777bp of the ATPase 6 and 8 genes
830   across 239 samples, contrasted against results of *de novo* species delimitation models represented by colored bars. Branches are
labelled according to the morphological diagnoses of the tissue samples from which mtDNA sequences originated and are abbreviated
832   as follows: *tigris=A. tigris tigris*; *punct.=A. t. punctilinealis*; *sept.=A. t. septentrionalis*; *mundus=A. t. mundus*; *marm.=A. marmoratus*;
shared=found in *A. t. tigris* and *A. t. punctilinealis*. Model abbreviations are: ABGD=Automated Barcode-Gap Discovery;
834   TAX=qualitative hypothesis derived from current taxonomy; CD5=CD-Hit with 5% divergence threshold for clustering; CD2= CD-
Hit with 2% divergence threshold for clustering; GMYC=Generalized mixed Yule coalescent; PTP=Poisson tree process; ROS=Clade
836   delimitation based on statistical tests of monophyly. Node labels represent posterior probabilities from Bayesian inference (B), and
neighbor-joining bootstrap support (A; omitted for shallow nodes). Nodes passing the statistical test of reciprocal monophyly are
838   indicated with a red dot. Clades in both trees are colored according to the ROS model.

840   Figure 4: Geo-referenced haplotype tree generated via Bayesian analysis, representing 777bp of
the ATPase 8 and 6 genes across 239 samples. Clade coloring follows the ROS model (see Fig.
842   1)

848   Figure 5: Among-group genetic variance (F_{CT}) for genetic clusters within the *Aspidoscelis tigris* complex, as derived by SAMOVA clustering runs. Cluster optimization constrained by (A)
850   geographic proximity of samples and (B) without a geographic constraint. The x-axis (K) reports the number of genetic clusters sampled.

Figure 6: Within- and among-group linear regression of geographic x genetic distances for clades delimited by (A) the ABGD
856 (Automated barcode-gap discovery) model and (B) TAX (taxonomic) model. Coefficients of correlation (r^2) ranged from 0.01-0.449.
Note that "*tigris*" refers to both *A. t. tigris* and *A. t. punctilinealis*, as these subspecies are not differentiated in the models. *A. t. mundus*
858 was excluded due to the geographically restricted sampling presented herein.

Figure 7: Map depicting results of an 'Alleles in Space' analysis for members of the *Aspidoscelis tigris* complex. Relative genetic differentiation is interpolated across the landscape, with individuals (as dots) colored by the CD5 model (see Fig. 1B). Heat map colors depict relative genetic distances per unit of geographic distance, with red corresponding to a greater genetic distance, and blue being lesser genetic distance.