Stabilizing role of seed banks and the maintenance of bacterial diversity

Nathan I. Wisnoski (wisnoski@indiana.edu) and Jay T. Lennon (lennonj@indiana.edu)

Department of Biology, Indiana University, Bloomington, Indiana, USA 47405

Statement of authorship: JTL designed the study and coordinated data collection. NIW

performed statistical analyses and wrote the first draft of the manuscript. Both authors

contributed substantially to edits and revisions.

Running head: Stabilization of freshwater bacterioplankton

Keywords: bacterioplankton, community dynamics, dormancy, negative frequency dependence,

seasonality

Type of Article: Letter

Abstract: 150 words; **Main text:** 4120 words

References: 60

Figures: 5; **Tables:** 1; **Text boxes:** 0

Correspondence: Jay T. Lennon, 1001 E. Third St., Bloomington, IN, 47405. Tel: (812) 856-

7235. Email: lennonj@indiana.edu.

ABSTRACT

Coexisting species often exhibit negative frequency dependence due to mechanisms that promote

population growth and persistence when rare. These stabilizing mechanisms maintain diversity

through interspecific niche differences, but also through life-history strategies like dormancy that

buffer dynamics in fluctuating environments. However, there are few tests demonstrating how

seed banks contribute to long-term community dynamics and the maintenance of diversity. Using

multi-year, high-frequency bacterial community data from a north temperate lake, we

documented patterns consistent with stabilizing coexistence, including differential responses of

bacterial taxa to environmental conditions and the presence of a seed bank. By maintaining rare

taxa over winter periods, the seed bank contributed to recurrent seasonal dynamics. Last, strong

negative frequency dependence in rare, but metabolically active, taxa suggested biotic

interactions promoted coexistence. Together, our results provide field-based evidence that niche

differences and seed banks stabilize bacterial community dynamics and contribute to the

maintenance of microbial diversity in nature.

INTRODUCTION

The maintenance of biodiversity is important for regulating species interactions, stabilizing ecosystem functions, and promoting resilience in response to perturbations. Diversity is maintained by a range of processes including niche differentiation in resource use (Tilman 1982; Gudelj *et al.* 2010; Johnson *et al.* 2012), defensive abilities (Leibold 1996; Thingstad *et al.* 2014; Cadier *et al.* 2019), and abiotic constraints (Holt 2009). These stabilizing niche differences among species contribute to the maintenance of diversity not only by preventing competitive exclusion, but also by allowing populations to recover from low abundances (Chesson 2000; Adler *et al.* 2007). Stabilizing coexistence mechanisms that rely on environmental fluctuations can further increase the number of species in a community (Chesson 1994; Chesson & Huntly 1997; Descamps-Julien & Gonzalez 2005). For example, in seasonal environments, species that are favored at different times of the year may be able to coexist in the community if they can survive through periods of unfavorable environmental conditions (Pake & Venable 1996).

Many species invest in persistence strategies that increase survival and facilitate recovery from rarity. Such strategies can also contribute to stabilization, especially in fluctuating environments where diversity can be maintained by the storage effect (Warner & Chesson 1985; Chesson 2000). The storage effect requires that taxa differ in their responses to environmental conditions, that intraspecific limitation peaks during favorable conditions, and that population growth is buffered in suboptimal environments by a long-lived life stage (Pake & Venable 1996; Cáceres 1997; Angert *et al.* 2009). The storage effect may be particularly important in communities where species experience periods of extremely slow growth (Gray *et al.* 2019) or engage in various forms of dormancy, which are common among plants and animals, but also microorganisms (Lennon & Jones 2011). For example, the accumulation of dormant individuals

into a "seed bank" not only reduces the probability of extinction associated with rarity, but it can

also facilitate the re-establishment of populations when favorable conditions return via the

process of resuscitation. While coexistence mediated by seed banks has been a key hypothesis

for the maintenance of microbial diversity (Jones & Lennon 2010; Lennon & Jones 2011;

Sorensen & Shade 2020), empirical evidence from temporally fluctuating environments is

lacking.

Stabilization from niche differences and seed banks should generate negative frequency-

dependence (NFD) in population growth. The implication of NFD is that rare populations grow

faster than common populations (Chesson 2000; Adler *et al.* 2007). Support for stabilizing

coexistence has largely come from plant and animal communities (Cáceres 1997; Angert *et al.*

2009; Yenni *et al.* 2017), while evidence for its role in complex microbial systems is less

common (Zhang *et al.* 2010). In microbial communities, which contain a disproportionately large

number of extinction-prone rare taxa (Sogin *et al.* 2006; Lynch & Neufeld 2015; Shade *et al.*

2018), taxa may vary widely in their stability and long-term contributions to diversity. Although

some low-abundance taxa reflect transient dynamics, other rare taxa persist for longer periods of

time (Alonso-Sáez *et al.* 2015; Lynch & Neufeld 2015; Newton & Shade 2016). While it is

unclear whether the persistence of these rare taxa is aided by niche differences and seed banks,

their potential coexistence suggests that NFD may be important for maintaining high levels of

biodiversity that are commonly observed in microbial communities.

In this study, we tracked bacterioplankton dynamics over time in a north temperate lake

using high-resolution molecular data to infer ecological processes that maintain microbial

diversity. Bacterial communities in fluctuating aquatic environments often exhibit recurrent,

seasonal community patterns (Shade *et al.* 2007; Gilbert *et al.* 2012; Fuhrman *et al.* 2015; Ward

et al. 2017), but the potential mechanisms that contribute to cyclical dynamics and maintain

diversity in nature are poorly resolved. We characterized how persistent (and putatively

coexisting) taxa respond to environmental fluctuations and used null models to assess whether

stabilizing biotic interactions (e.g., self-limitation, as evidenced by strong NFD) help maintain

rare, but metabolically active, taxa in the community (Yenni *et al.* 2017; Rovere & Fox 2019).

Specifically, we compared patterns of NFD and population dynamics in the active and total

portions of the community (inferred by 16S rRNA transcripts and genes, respectively) to

quantify the importance of slow growth or dormancy strategies for the maintenance of diversity.

Consistent with predictions from the storage effect, our results provide empirical evidence that

stabilizing biotic interactions and seed bank dynamics underlie seasonal community dynamics

and play key roles in maintaining bacterial diversity in natural ecosystems.

97   **METHODS**

Study site and sampling: University Lake is a 3.2 ha meso-eutrophic reservoir located in

the Indiana University Research and Teaching Preserve, Bloomington, Indiana, USA (39°11' N,

86°30' W). The surrounding watershed is dominated by oak, beech, and maple forests. Three

streams drain into University Lake, which has an estimated volume of 150,000 m^3 and a

maximum depth of 10 m. From April 2013 to September 2015, we took weekly water samples

from the surface (0-0.5 m) of the epilimnion for microbial biomass, and measured environmental

variables commonly associated with aquatic microbial community dynamics: total phosphorus

(TP), total nitrogen (TN), and dissolved organic carbon (DOC). Microbial biomass was filtered

on 0.2 µm Supor filters (Pall, Port Washington, NY, USA) and frozen at -80 °C. We quantified

TP using the ammonium molybdate method (Wetzel & Likens 2000) and TN with the second

derivative method after persulfate digestion (Crumpton *et al.* 1992). DOC was quantified on 0.7

μm filtrates using nondispersive infrared (NDIR) detection on a Shimadzu TOC-V (Kyoto,

Japan). We also quantified water transparency with a Secchi disk and used a Quanta Hydrolab

(OTT, Kempton, Germany) water sonde to measure temperature, conductivity, dissolved oxygen,

salinity, and pH of the samples.

Bacterial community structure: We characterized the structure of the bacterial

community using high-throughput 16S rRNA sequencing. We extracted total nucleic acids from

biomass retained on 0.2 μm filters using the MoBio PowerWater RNA extraction kit and the

DNA elution accessory kit. Because sequences obtained from DNA can come from

metabolically active or inactive (e.g., slow growing or dormant) individuals, this sample

represents the "total" community. In contrast, RNA is a more ephemeral molecule that is

essential for synthesizing proteins; therefore, it is often used to characterize the metabolically

"active" subset of the community (Molin & Givskov 1999; Steiner *et al.* 2019; Locey *et al.*

2020). After extracting nucleic acids, we used DNase (Invitrogen) to remove DNA from the

RNA extractions and then synthesized cDNA with SuperScript III First Strand Synthesis kit and

random hexamer primers (Invitrogen). To amplify the 16S rRNA gene (DNA) and transcripts

(cDNA), we used barcoded V4 primers (515F and 806R) designed for the Illumina MiSeq

platform (Caporaso *et al.* 2012). We then purified the PCR products with AMPure XP,

quantified DNA concentrations using PicoGreen, and pooled samples at 10 ng per sample. The

resulting libraries were sequenced on an Illumina MiSeq at the Indiana University Center for

Genomic and Bioinformatics Sequencing Facility using 250×250 bp paired end reads (Reagent

Kit v2). Paired-end sequences were subsequently processed using the software package mothur

(version 1.41.1) (Schloss *et al.* 2009). We assembled contigs, removed low quality sequences

(minimum score of 35), aligned sequences to the SILVA Database (version 132) (Quast *et al.*

2013), removed chimeras using the VSEARCH algorithm (Rognes *et al.* 2016), and created 97%

similar operational taxonomic units (OTUs) using the OptiClust algorithm (Westcott & Schloss

2017), and classified sequences with the RDP taxonomy (Cole *et al.* 2009). To account for

variation in sequencing depth, subsequent analyses were performed on rarefied abundance data

subsampled to the fewest number of reads in the time series ($N = 5979$ per sample) using R

(version 3.6.0) (R Core Team 2018).

Differential responses to environment: To evaluate the first criterion of the storage effect,

we looked for signatures of niche partitioning along a suite of environmental variables. First, we

performed a principal component analysis (PCA) on Hellinger-transformed abundances to

visualize seasonal patterns of compositional trajectories. Next, we identified environmental

drivers of community dynamics using redundancy analysis (RDA). We then looked more closely

at the persistent, and potentially coexisting, subset of taxa in the community (OTUs present in

≥80% of the DNA-based samples, Table S1). To determine whether environmental fluctuations

facilitated temporal niche partitioning, we identified the week of the year when each persistent

OTU (n = 82) experienced its maximum average growth rate (see calculations in *Stabilizing*

biotic interactions below). For each of the persistent OTUs, we compared the environmental

conditions at the time of the year when they experienced maximum growth (as an index of

optimal conditions along an annual environmental cycle). Specifically, we performed a principal

component analysis (PCA) on the environmental variables including temperature, specific

conductivity, Secchi depth, pH, TP, TN, and DOC (all standardized to mean = 0, standard

deviation = 1). We plotted each time point along the first two PC axes, along with the PC

loadings of the environmental variables, and labeled each point with the name of the OTU that

exhibited maximum growth at that particular time point (Fig. S1).

Seed bank dynamics: To assess the importance of the next criterion of the storage effect,

buffered population growth, we analyzed the temporal role of the seed bank for maintaining

bacterial diversity. First, we examined whether the seed bank served as a reservoir of taxonomic

diversity by comparing the ratio of total richness to active richness at each time point in the time

series, where larger ratios indicate that the total community had higher α-diversity than the active

subset of the community. Second, we sought to determine whether seed bank dynamics were

more important for the maintenance of rare or common taxa in the community. To do so, we

developed a reactivation metric to quantify each OTUs frequency of reactivation from the seed

bank. For each OTU, its reactivation score is the number of times an OTU was present (i.e.,

detected in the DNA pool) but likely in an inactive (i.e., absent from the RNA pool) state at time

point t, yet active (present in the RNA pool) at the subsequent time point $t+1$. This represents a

transition from the inactive to active state mediated by slow growth or dormancy based on

recovery of sequences in the DNA and RNA pools. Thus, OTUs with higher reactivation scores

may be more reliant on the seed bank for long-term persistence in the community. We then

analyzed the relationship between the average relative abundance of active OTUs (excluding

zeroes) and their reactivation score to determine whether seed banking was more important for

maintaining rare taxa than common taxa in the bacterioplankton community.

Stabilizing biotic interactions: In relation to the last criterion of the storage effect, that

taxa experience the strongest self-limitation during favorable environments, we examined

patterns of negative frequency dependence in the community. In particular, we examined (1)

whether growth exhibited negative frequency dependence overall, which is indicative of

stabilization in the community, and (2) whether rare taxa experienced stronger negative

frequency dependence than common taxa (Yenni *et al.* 2012, 2017; Rovere & Fox 2019). Such

patterns of NFD shed light on the storage effect for three reasons. First, taxa are more likely to be

common in the community when they are environmentally favored relative to other taxa in the

community. Second, taxa that experience strong intraspecific limitation during environmentally

favorable periods would grow faster when rare than when common, thereby generating NFD.

Third, differences among taxa in the strength of self-limitation during favored growth periods

would lead to different average relative abundances in the community, implying that rare yet

persistent taxa may by strongly stabilized (Yenni *et al.* 2012, 2017; Rovere & Fox 2019).

However, metabolically inactive individuals are unlikely to engage in the biotic interactions that

generate NFD. Therefore, we focused on NFD in the metabolically active (i.e., RNA-based)

portion of the community comprised of the persistent taxa (Table S1).

We then calculated NFD for each OTU by comparing rates of change in relative

abundance between weekly samples. We inferred the strength of NFD for a given OTU as the

magnitude of the negative slope of the relationship between an OTU's relative abundance and its

per capita growth rate at each time step (t) across the time series. We calculated the relative

abundance ($x_{t,s}$), of each OTU (s) as its abundance ($N_{t,s}$) in the community of s OTUs relative

to the total abundance of all s OTUs ($\sum_s N_{t,s}$) at a given time step (t), such that $x_{t,s} = \frac{N_{t,s}}{\sum_s N_{t,s}}$.

From this, we then calculated the natural log of the per capita growth rate of each OTU as $y_{t,s} =$

$\log_e \left(\frac{N_{t+1,s}}{N_{t,s}} \right)$. To estimate the strength of negative frequency dependence for each OTU, we fit

simple linear regressions ($y_s = \beta_{0,s} + \beta_{1,s} x_s + \epsilon_s$), where the equilibrium frequency of an OTU

(f) is the x-intercept, $f = -\frac{\beta_{0,s}}{\beta_{1,s}}$, and the degree of frequency dependence (NFD) is the slope

$NFD = \beta_{1,s}$. In the end, f describes whether an OTU is common or rare, and negative slopes

with greater magnitudes indicate stronger negative frequency dependence.

Stabilization of rare bacterial taxa: We used null models to infer whether rare taxa

exhibited stronger NFD than expected by chance. Stabilization of rare taxa would be supported if

OTUs with lower equilibrium frequencies (smaller values of f) had more negative slopes (larger

$|\beta_{1,s}|$) indicating stronger self-limitation. We inferred the overall relationship between rarity and

strength of NFD from the covariance between the $\log(NFD)$ and $\log(f)$: a more negative

covariance would indicate that rarer taxa were more strongly stabilized than common taxa. To

account for the fact that the expectation of this covariance is already negative, and to control for

spurious statistical correlations in the temporal data due to other factors, we implemented a null

model approach (Yenni *et al.* 2017; Rovere & Fox 2019). We shuffled the abundances of each

OTU independently, recalculated relative abundances and per capita growth rates, estimated

equilibrium frequencies (fs) and negative frequency dependences (NFDs), and calculated the

covariance, repeating this procedure 5000 times to generate a null distribution of covariance

values ($COV[\log(f), \log(NFD)]$) (Yenni *et al.* 2017). We then compared our observed

covariance with the null distribution to infer the strength of asymmetry in NFD (i.e., the degree

to which rare OTUs experience disproportionately stronger self-limitation than common OTUs).

We quantified divergence from null distributions using standardized effect sizes (SES = mean

observed covariance / standard deviation of covariances in the null distribution) and the ratio of

observed covariance to the average covariance of the null distribution (Yenni *et al.* 2017). More

negative SES values and larger ratios would indicate greater deviations from the null expectation

of equal NFD across taxa. We inferred the degree of statistical significance by calculating a p-

value as the proportion of null covariance values less than or equal to our observed covariance.

227   **RESULTS**

Differential responses to environment: Bacterial community dynamics were related to

environmental variability, with different taxa favored at different times of the year. During the

summer months, the community followed a recurrent successional trajectory (Fig. 1A). This

trajectory was strongly aligned with seasonal trends in temperature (Fig. 1B). Across longer time

scales, inter-annual variation in dissolved oxygen and pH was associated with compositional

differences in the active bacterial community during winter months. Within an annual cycle, the

persistent OTUs (n = 82) demonstrated temporal partitioning in their maximal growth rates in the

active portion of the community (Fig. 2, Table S1), corresponding to different environmental

conditions (Fig. S1).

Seed bank dynamics: Our data suggest seed banks of dormant or slow growing

individuals contribute to the maintenance of diversity. Over the course of our study, total

richness ranged from 1.2–2.0 times higher than the richness of the active portion of the

community (Fig. 3). Furthermore, this discrepancy between total and active richness exhibited

seasonality, demonstrating a time-varying role for the bacterial seed bank. In particular, the seed

bank played a weaker role (i.e., active and total richness were more similar in magnitude) during

the summer, while proportionally higher diversity was found in the seed bank over winter, when

growing conditions may be less optimal (Fig. 3). In addition, the taxa that exhibited more

reactivations from the seed bank were also the taxa that were, on average, consistently rare when

active in the community (Fig. 4). In contrast, common taxa exhibited fewer transitions between

active and inactive states in the community. Thus, our findings suggest that rare taxa in the

community rely on life-history strategies, such as slow growth or dormancy, to avoid local

extinction.

Stabilizing biotic interactions: Persistent taxa exhibited stabilizing NFD, which varied in

strength depending on each taxon's mean relative abundance in the community (Fig. S2). In

particular, NFD was significantly stronger for rare taxa than common taxa, but only in the active

portion of the community ($p = 0.0002$; SES = -4.03, covariance ratio = 1.08), not in the total

community ($p = 0.221$; SES = -0.777, covariance ratio = 1.01) (Fig. 5). The p-values reflect the

rank of observed NFD compared with null simulations, while SES values take into account the

variance in the null distribution. In other words, the total community showed nearly the same

degree of stabilization (|SES| < 2, covariance ratio ~ 1) as the null communities.

262   **DISCUSSION**

Our findings from a multi-year survey support the view that biodiversity was maintained

by stabilizing mechanisms, including niche differentiation and seed bank dynamics, that

generated negative frequency dependence (NFD) in a natural bacterioplankton community. High

resolution sampling revealed recurrent seasonality in community dynamics (Fig. 1), driven by

taxon-specific responses to annual environmental fluctuations (Fig. 2, Fig. S2). Our results also

showed that the maintenance of diversity may be enhanced by life-history strategies, such as

slow growth or dormancy, that buffered rare taxa from local extinction during environmentally

unfavorable periods (e.g., winter) and facilitated reestablishment when conditions improved (Fig.

3-4). These apparent niche differences and seed bank dynamics contributed to stabilizing biotic

interactions (e.g., stronger intra- than interspecific limitation) among rare, but metabolically

active, taxa in the community (Fig. 5).

Negative frequency dependence in microbial communities: We found evidence for

stabilization through negative frequency dependence (NFD). While documented in some plant

and animal assemblages (Harpole & Suding 2007; Yenni *et al.* 2017; Rovere & Fox 2019),

observations of NFD in complex microbial communities are uncommon. In particular, our study

revealed disproportionately strong NFD for rare taxa, offering an explanation for why some taxa

appear to stably persist at low relative abundances in nature (Alonso-Sáez *et al.* 2015; Lindh *et*

al. 2015), potentially as members of the "rare biosphere" (Sogin *et al.* 2006; Lynch & Neufeld

2015; Shade *et al.* 2018). Our approach also allowed us to identify stabilizing mechanisms

operating among metabolically active rare taxa that were not detectable from the dynamics of the

total bacterial community (Fig. 5). Ignoring this metabolic heterogeneity can obscure inferences

of underlying ecological processes (Wisnoski *et al.* 2020), and would have gone otherwise

undetected in this study as well.

Asymmetric NFD is also important for the coexistence of rare plant and animal species,

but the magnitude of this effect varies across taxonomic groups (Yenni *et al.* 2017; Rovere &

Fox 2019). For example, NFD is less asymmetric for herpetofauna than plant or mammal

communities (Yenni *et al.* 2017), possibly due to higher evenness (Rovere & Fox 2019).

Compared to macro-organismal systems, the degree of NFD asymmetry in our highly uneven

bacterial community was moderate (SES = -4.03, covariance ratio = 1.08), suggesting that

coexistence among rare taxa may be weak, or that additional factors not captured by this metric

are important for maintaining diversity in this system. However, our comparison between the

metabolically active and total portions of the bacterioplankton community provides critical

evidence that active bacteria mediate the biotic interactions that generate asymmetry in NFD and

allow rare taxa to persist. Consistent with prior work showing that rare taxa may be

disproportionately active in freshwater bacterial communities (Jones & Lennon 2010), our study

demonstrates that rare, metabolically active bacteria may also be critical for the long-term

maintenance of bacterial diversity.

Storage effect in microbial communities: Temporal niche partitioning mediated by seed

bank dynamics may explain the maintenance of diversity in fluctuating environments. Evidence

for coexistence via the storage effect comes largely from communities of desert annuals (Pake &

Venable 1996; Angert *et al.* 2009), grasslands (Adler *et al.* 2006), tropical trees (Usinowicz *et al.*

2012), zooplankton (Cáceres 1997), and marine fish (Secor 2007). In most bacterial studies, the

role of seed banks for coexistence has been inferred from short-term observations, but here we

provide temporal evidence that bacterial seed banks may be important for community dynamics

and the maintenance of diversity. In the temperate climate of our study lake, different taxa

showed maximum growth rates at different times of the year, coinciding with seasonal transitions

in environmental conditions (Fig. 2), and contrasting active and total community dynamics

suggested buffered population dynamics (Figs. 3-5). In addition, species showed NFD in their

growth such that they experienced greater self-limitation (consistent with stronger intraspecific

than interspecific competition) when they were more common in the active community (and

were thus more likely favored by the environment) (Fig. 5, Fig. S2). While these conditions are

not definitive proof of a storage effect acting in our study system, they are consistent with the

criteria needed for a storage effect to contribute to the long-term maintenance of bacterial

diversity in the lake.

Seed banks may also have more general implications for bacterial community dynamics.

The persistence of taxa with temporal niche differences may contribute to the repeatability of

summer community dynamics in the active portion of the community (Hellweger *et al.* 2008) by

favoring overwinter survival (Fig. 3). For example, the seasonality of the seed bank indicates that

diversity stored in the seed bank was maximized when environmental conditions (e.g., water

temperature, resource/consumer densities) were least favorable for bacterial growth

(Neuenschwander *et al.* 2015). This pattern is consistent with the presence of a dormant seed

bank that buffers individuals from harsh conditions. In addition, transitions from inactive to

active metabolic states were more frequently detected among taxa that were, on average, rare

when active in the community. Analogous to the methodological challenges of finding and

identifying dormant individuals in non-microbial seed banks (e.g., plants), RNA detection limits

may affect the classification of metabolically active bacteria. Nevertheless, our reactivation

metric should capture rapid shifts in the metabolically active portion of the community. Overall,

these patterns suggest that recurrent environmental cues regulate active community dynamics by

favoring different taxa at different times of the year, and that seed banks are important for

maintaining these seasonal community trajectories at multi-annual timescales.

Future directions and conclusions: Our study provides empirical evidence supporting the

theory that biotic mechanisms stabilize bacterial communities and maintain diversity in nature.

While our results indicate differences in the diversity, dynamics, and stabilization between active

and total subsets of the community, an ultimate goal is to tighten the mechanistic links between

rates of ribosomal RNA transcription and *in situ* growth rates for individual taxa (Newton &

Shade 2016; Papp *et al.* 2018) or through other techniques that involve the physical sorting of

cells based on metabolic activity prior to sequencing (Couradeau *et al.* 2019; Reichart *et al.*

2020). It may also be important to consider the dispersal of terrestrial bacteria into aquatic

ecosystems (Crump *et al.* 2012). However, previous work in our study system revealed that most

terrestrial bacteria were metabolically inactive, most likely reflecting the abrupt environmental

transitions that accompany cross-ecosystem dispersal (Wisnoski *et al.* 2020). Thus, the lack of

asymmetric NFD we observed in the total community could also arise in part from allochthonous

inputs of inactive bacteria that decouple population growth of each OTU from its relative

abundance in the community. However, by focusing on the dynamics of metabolically active

bacteria, our approach was able to uncover the presence of stabilizing biotic interactions that

may have otherwise been obscured by metabolic heterogeneity in the total community.

In conclusion, we show that stabilizing biotic interactions and the ability to engage in

dormancy or slow growth strategies play important roles in maintaining microbial diversity in a

natural ecosystem over a multi-year time scale. Our results demonstrate the mechanisms at the

community scale that preserve Earth's vast microbial diversity, building on other explanations

that emphasize the importance of metabolic diversity (Sala *et al.* 2008), capacity for rapid growth

(Shade *et al.* 2014), and spatial scale (Vos *et al.* 2013). In particular, strong NFD offers a new

explanation for why the majority of bacterial taxa persist at low average relative abundances in

nature (Lynch & Neufeld 2015). Furthermore, our work builds on inferences about the roles of

microbial dormancy (and other persistence strategies) obtained from shorter time scales, and

provides temporal evidence that dormancy is an important buffer against local extinction over

longer time scales. More generally, our work demonstrates the importance of stabilization in

microbial systems, offering new insight into the long-term maintenance of microbial diversity.

ACKNOWLEDGEMENTS

We acknowledge technical support from Brent Lehmkuhl and Drew Meyer. This work

was supported by the National Science Foundation (DEB-1442246 to JTL), US Army Research

Office Grant (W911NF-14-1-0411 to JTL), National Aeronautics and Space Administration

(80NSSC20K0618 to JTL), and the Department of Biology at Indiana University (Louise

Constable Hoover Fellowship to NIW). Data and code for the project can be found at NCBI

(BioProject PRJNA664410) and a Zenodo archive of the GitHub repository

(https://github.com/LennonLab/ul-seedbank).

REFERENCES

Adler, P.B., HilleRisLambers, J., Kyriakidis, P.C., Guan, Q. & Levine, J.M. (2006). Climate
variability has a stabilizing effect on the coexistence of prairie grasses. *Proc. Natl. Acad.*
Sci., 103, 12793–12798.
Adler, P.B., HilleRisLambers, J. & Levine, J.M. (2007). A niche for neutrality. *Ecol. Lett.*, 10,
95–104.
Alonso-Sáez, L., Díaz-Pérez, L. & Morán, X.A.G. (2015). The hidden seasonality of the rare
biosphere in coastal marine bacterioplankton. *Environ. Microbiol.*, 17, 3766–3780.
Angert, A.L., Huxman, T.E., Chesson, P. & Venable, D.L. (2009). Functional tradeoffs
determine species coexistence via the storage effect. *Proc. Natl. Acad. Sci.*, 106, 11641–
11645.
Cáceres, C.E. (1997). Temporal variation, dormancy, and coexistence: a field test of the storage
effect. *Proc. Natl. Acad. Sci.*, 94, 9171–9175.

Cadier, M., Andersen, K.H., Visser, A.W. & Kiørboe, T. (2019). Competition–defense tradeoff
388   increases the diversity of microbial plankton communities and dampens trophic cascades.
389   *Oikos*, 128, 1027–1040.
390   Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., *et al.*
391   (2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and
392   MiSeq platforms. *ISME J.*, 6, 1621–1624.
393   Chesson, P. (1994). Multispecies competition in variable environments. *Theor. Popul. Biol.*, 45,
394   227–276.
395   Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.*, 31,
396   343–366.
397   Chesson, P. & Huntly, N. (1997). The roles of harsh and fluctuating conditions in the dynamics
398   of ecological communities. *Am. Nat.*, 150, 519–553.
399   Cole, J.R., Wang, Q., Cardenas, E., Fish, J., Chai, B., Farris, R.J., *et al.* (2009). The Ribosomal
400   Database Project: Improved alignments and new tools for rRNA analysis. *Nucleic Acids*
401   *Res.*, 37, 141–145.
402   Couradeau, E., Sasse, J., Goudeau, D., Nath, N., Hazen, T.C., Bowen, B.P., *et al.* (2019).
403   Probing the active fraction of soil microbiomes using BONCAT-FACS. *Nat. Commun.*,
404   10, 2770.
405   Crump, B.C., Amaral-Zettler, L.A. & Kling, G.W. (2012). Microbial diversity in arctic
406   freshwaters is structured by inoculation of microbes from soils. *ISME J.*, 6, 1629–1639.
407   Crumpton, W.G., Isenhart, T.M. & Mitchell, P.D. (1992). Nitrate and organic N analyses with
408   second-derivative spectroscopy. *Limnol. Oceanogr.*, 37, 907–913.
409   Descamps-Julien, B. & Gonzalez, A. (2005). Stable coexistence in a fluctuating environment: an
410   experimental demonstration. *Ecology*, 86, 2815–2824.
411   Fuhrman, J.A., Cram, J.A. & Needham, D.M. (2015). Marine microbial community dynamics
412   and their ecological interpretation. *Nat. Rev. Microbiol.*, 13, 133–146.
413   Gilbert, J.A., Steele, J.A., Caporaso, J.G., Steinbrück, L., Reeder, J., Temperton, B., *et al.*
414   (2012). Defining seasonal marine microbial community dynamics. *ISME J.*, 6, 298–308.
415   Gray, D.A., Dugar, G., Gamba, P., Strahl, H., Jonker, M.J. & Hamoen, L.W. (2019). Extreme
416   slow growth as alternative strategy to survive deep starvation in bacteria. *Nat. Commun.*,
417   10, 890.
418   Gudelj, I., Weitz, J.S., Ferenci, T., Claire Horner-Devine, M., Marx, C.J., Meyer, J.R., *et al.*
419   (2010). An integrative approach to understanding microbial diversity: from intracellular
420   mechanisms to community structure. *Ecol. Lett.*, 13, 1073–1084.
421   Harpole, W.S. & Suding, K.N. (2007). Frequency-dependence stabilizes competitive interactions
422   among four annual plants. *Ecol. Lett.*, 10, 1164–1169.
423   Hellweger, F.L., Kravchuk, E.S., Novotny, V. & Gladyshev, M.I. (2008). Agent-based modeling
424   of the complex life cycle of a cyanobacterium (Anabaena) in a shallow reservoir. *Limnol.*
425   *Oceanogr.*, 53, 1227–1241.
426   Holt, R.D. (2009). Bringing the Hutchinsonian niche into the 21st century: ecological and
427   evolutionary perspectives. *Proc. Natl. Acad. Sci.*, 106, 19659–19665.
428   Johnson, D.R., Goldschmidt, F., Lilja, E.E. & Ackermann, M. (2012). Metabolic specialization
429   and the assembly of microbial communities. *ISME J.*, 6, 1985–1991.
430   Jones, S.E. & Lennon, J.T. (2010). Dormancy contributes to the maintenance of microbial
431   diversity. *Proc. Natl. Acad. Sci.*, 107, 5881–5886.

Leibold, M.A. (1996). A graphical model of keystone predators in food webs: trophic regulation
of abundance, incidence, and diversity patterns in communities. *Am. Nat.*, 147, 784–784.
Lennon, J.T. & Jones, S.E. (2011). Microbial seed banks: the ecological and evolutionary
implications of dormancy. *Nat. Rev. Microbiol.*, 9, 119–130.
Lindh, M.V., Sjöstedt, J., Andersson, A.F., Baltar, F., Hugerth, L.W., Lundin, D., *et al.* (2015).
Disentangling seasonal bacterioplankton population dynamics by high-frequency
sampling. *Environ. Microbiol.*, 17, 2459–2476.
Locey, K.J., Muscarella, M.E., Larsen, M.L., Bray, S.R., Jones, S.E. & Lennon, J.T. (2020).
Dormancy dampens the microbial distance–decay relationship. *Philos. Trans. R. Soc. B
Biol. Sci.*, 375, 20190243.
Lynch, M.D.J. & Neufeld, J.D. (2015). Ecology and exploration of the rare biosphere. *Nat. Rev.
Microbiol.*, 13, 217–229.
Molin, S. & Givskov, M. (1999). Application of molecular tools for in situ monitoring of
bacterial growth activity. *Environ. Microbiol.*, 1, 383–391.
Neuenschwander, S.M., Pernthaler, J., Posch, T. & Salcher, M.M. (2015). Seasonal growth
potential of rare lake water bacteria suggest their disproportional contribution to carbon
fluxes. *Environ. Microbiol.*, 17, 781–795.
Newton, R.J. & Shade, A. (2016). Lifestyles of rarity: understanding heterotrophic strategies to
inform the ecology of the microbial rare biosphere. *Aquat. Microb. Ecol.*, 78, 51–63.
Pake, C.E. & Venable, D.L. (1996). Seed banks in desert annuals: implications for persistence
and coexistence in variable environments. *Ecology*, 77, 1427–1435.
Papp, K., Hungate, B.A. & Schwartz, E. (2018). Microbial rRNA synthesis and growth
compared through quantitative stable isotope probing with $H_2^{18}O$. *Appl. Environ.
Microbiol.*, 84, e02441-17.
Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., *et al.* (2013). The SILVA
ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic Acids Res., 41, 590–596.
R Core Team. (2018). *R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.*
Reichart, N.J., Jay, Z.J., Krukenberg, V., Parker, A.E., Spietz, R.L. & Hatzenpichler, R. (2020).
Activity-based cell sorting reveals responses of uncultured archaea and bacteria to
substrate amendment. *ISME J.*, 1–11.
Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. (2016). VSEARCH: a versatile open
source tool for metagenomics. *PeerJ*, 4, e2584–e2584.
Rovere, J. & Fox, J.W. (2019). Persistently rare species experience stronger negative frequency
dependence than common species: A statistical attractor that is hard to avoid. *Glob. Ecol.
Biogeogr.*, 28, 508–520.
Sala, M.M., Terrado, R., Lovejoy, C., Unrein, F. & Pedrós-Alió, C. (2008). Metabolic diversity
of heterotrophic bacterioplankton over winter and spring in the coastal Arctic Ocean.
Environ. Microbiol., 10, 942–949.
Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., *et al.* (2009).
Introducing mothur: open-source, platform-independent, community-supported software
for describing and comparing microbial communities. *Appl. Environ. Microbiol.*, 75,
7537–7541.

Secor, D.H. (2007). The year-class phenomenon and the storage effect in marine fishes. *J. Sea*
Res., Proceedings of the Sixth International Symposium on Flatfish Ecology, Part I, 57,
91–103.
Shade, A., Dunn, R.R., Blowes, S.A., Keil, P., Bohannan, B.J.M., Herrmann, M., *et al.* (2018).
Macroecology to unite all life, large and small. *Trends Ecol. Evol.*, 33, 731–744.
Shade, A., Jones, S.E., Gregory Caporaso, J., Handelsman, J., Knight, R., Fierer, N., *et al.*
(2014). Conditionally rare taxa disproportionately contribute to temporal changes in
microbial diversity. *mBio*, 5, 1–9.
Shade, A., Kent, A.D., Jones, S.E., Newton, R.J., Triplett, E.W. & McMahon, K.D. (2007).
Interannual dynamics and phenology of bacterial communities in a eutrophic lake.
Limnol. Oceanogr., 52, 487–494.
Sogin, M.L., Morrison, H.G., Huber, J.A., Welch, D.M., Huse, S.M., Neal, P.R., *et al.* (2006).
Microbial diversity in the deep sea and the underexplored "rare biosphere." *Proc. Natl.*
Acad. Sci., 103, 12115–12120.
Sorensen, J.W. & Shade, A. (2020). Dormancy dynamics and dispersal contribute to soil
microbiome resilience. *Philos. Trans. R. Soc. B Biol. Sci.*, 375, 20190255.
Steiner, P.A., Corte, D.D., Geijo, J., Mena, C., Yokokawa, T., Rattei, T., *et al.* (2019). Highly
variable mRNA half-life time within marine bacterial taxa and functional genes. *Environ.*
Microbiol.
Thingstad, T.F., Våge, S., Storesund, J.E., Sandaa, R.-A. & Giske, J. (2014). A theoretical
analysis of how strain-specific viruses can control microbial species diversity. *Proc. Natl.*
Acad. Sci., 111, 7813–7818.
Tilman, D. (1982). *Resource competition and community structure*. Princeton University Press,
Princeton, NJ.
Usinowicz, J., Wright, S.J. & Ives, A.R. (2012). Coexistence in tropical forests through
asynchronous variation in annual seed production. *Ecology*, 93, 2073–2084.
Vos, M., Wolf, A.B., Jennings, S.J. & Kowalchuk, G.A. (2013). Micro-scale determinants of
bacterial diversity in soil. *FEMS Microbiol. Rev.*, 37, 936–954.
Ward, C.S., Yung, C.-M., Davis, K.M., Blinebry, S.K., Williams, T.C., Johnson, Z.I., *et al.*
(2017). Annual community patterns are driven by seasonal switching between closely
related marine bacteria. *ISME J.*, 11, 1412–1422.
Warner, R.R. & Chesson, P.L. (1985). Coexistence mediated by recruitment fluctuations: a field
guide to the storage effect. *Am. Nat.*, 125, 769–787.
Westcott, S.L. & Schloss, P.D. (2017). OptiClust, an improved method for assigning amplicon-
based sequence data to operational taxonomic units. *mSphere*, 2, e00073-17.
Wetzel, R.G. & Likens, G.E. (2000). *Limnological analyses*. 3rd ed. Springer, New York, NY.
Wisnoski, N.I., Muscarella, M.E., Larsen, M.L., Peralta, A.L. & Lennon, J.T. (2020). Metabolic
insight into bacterial community assembly across ecosystem boundaries. *Ecology*, 101,
e02968.
Yenni, G., Adler, P.B. & Ernest, S.K.M. (2012). Strong self-limitation promotes the persistence
of rare species. *Ecology*, 93, 456–461.
Yenni, G., Adler, P.B. & Ernest, S.K.M. (2017). Do persistent rare species experience stronger
negative frequency dependence than common species? *Glob. Ecol. Biogeogr.*, 26, 513–
523.
Zhang, Q.-G., Buckling, A. & Godfray, H.C.J. (2010). Quantifying the relative importance of
niches and neutrality for coexistence in a model microbial system. *J. Ecol.*, 1139–1147.

 FIGURE LEGENDS

Figure 1. Seasonal dynamics of the active bacterial community in University Lake. (A) The

compositional trajectory of the active community (determined by high-throughput sequencing of

16S rRNA transcripts) shows strong seasonality, but the community remains relatively static

over winter. The first two axes of the principal component analysis (PCA) depict summer/winter

differences (PC1) along the major axis, and slight inter-annual differences in winter composition

(cool colors) along the minor axis (PC2). The summer successional trajectory (warmer colors) is

highly repeatable across years. (B) Constrained ordination using redundancy analysis (RDA)

shows the environmental drivers of community structure, along with strong correlates of

individual taxa in the community. This analysis reveals that differences in pH and dissolved

oxygen (DO) explain variation in winter composition among years.

Figure 2. Temporal partitioning of maximum growth rate among persistent bacterial taxa in

University Lake. Lines represent the mean daily growth rate for each taxon over the time series.

Points indicate the maximum growth observed for each bacterial taxon (OTU). Overall, the 82

persistent OTUs have maximum growth rates at different seasons of the year. Points are color-

coded such that warmer colors correspond to spring and summer months and cooler colors

correspond to winter months. Colored lined trace out the growth dynamics of three individual

taxa with different environmental responses (blue = OTU 1, Betaproteobacteria; yellow = OTU

17, Actinobacteria; mauve = OTU 18, Gammaproteobacteria). More taxonomic details can be

found in Table S1.

Figure 3. Seasonal importance of the seed bank for bacterial diversity in University Lake.

Richness was much higher in the total community, relative to the active community, during the

fall and winter months. The active and total communities converged over the summer, indicated

by values on the y-axis closer to 1. Warmer colors correspond to spring and summer months,

while cooler colors correspond to winter months. Shaded regions correspond to the fall and

winter months (October through March).

Figure 4. Rare taxa showed more seed bank transitions than common taxa. For the 82 persistent

taxa identified over the time series, OTUs that were (on average) rare in the active portion of the

community had a higher number of reactivations from the seed bank, while more common taxa

had fewer reactivations. The fit was performed via locally estimated scatterplot smoothing

(LOESS).

Figure 5. Asymmetric negative frequency dependence (NFD) for persistent bacterial taxa (n =

82) was only detected in the active portion of the community. The degree of asymmetry in NFD

is determined by the covariance between the equilibrium frequency of each OTU and its strength

of NFD; negative covariance indicates that rarer taxa exhibit stronger NFD. Compared with

expected covariances from a null distribution, the standardized effect size (SES) of the observed

covariance in the active portion of the community was -4.03, while the SES of the total

community was -0.77. The overall strength of NFD (observed NFD / mean NFD) was 1.08 in the

active portion and 1.01 in the total community. The metabolic state on the y-axis indicates

whether the NFD comparison is for the active portion of the community (inferred from 16S

rRNA transcripts) or the total portion (inferred from 16S rRNA genes, i.e., DNA).

FIGURES

569   Figure 1

573   Figure 2.

577   Figure 3.

581   Figure 4.

585   Figure 5.

Covariance between Equilibrium Frequency
and Negative Frequency Dependence

590   **Table S1**. Operational taxonomic units (OTUs) that were classified as persistent in the

591   bacterioplankton community based on being detected in ≥80% of the total (i.e., DNA)

592   community samples. The table is sorted by Julian date of max growth.

OTU	Class	Max growth rate (d^{-1})	Date of max growth
Otu00034	Alphaproteobacteria	0.3426	2014-01-17
Otu00045	Betaproteobacteria	0.8527	2014-01-03
Otu00196	Actinobacteria	0.5873	2015-01-09
Otu00019	Cytophagia	0.7346	2014-02-14
Otu00039	Betaproteobacteria	0.7167	2014-02-14
Otu00102	Betaproteobacteria	0.6729	2015-02-28
Otu00105	Alphaproteobacteria	0.8666	2014-02-28
Otu00065	Sphingobacteriia	0.8563	2014-03-21
Otu00292	Alphaproteobacteria	0.6851	2014-03-07
Otu00006	Sphingobacteriia	0.3643	2013-04-25
Otu00012	Betaproteobacteria	0.8784	2014-04-18
Otu00014	Actinobacteria	0.4906	2015-04-26
Otu00016	Actinobacteria	0.7836	2014-04-18
Otu00017	Actinobacteria	1.0200	2015-04-04
Otu00021	Gammaproteobacteria	1.0449	2014-04-18
Otu00033	Alphaproteobacteria	0.5306	2014-04-25
Otu00048	Verrucomicrobiae	0.7426	2015-04-11

Otu00049	Actinobacteria	0.6090	2014-04-04
Otu00055	Flavobacteriia	0.7950	2014-04-18
Otu00058	Armatimonadia	0.8153	2015-04-11
Otu00148	Bacteria sp.	0.7646	2013-04-25
Otu00172	Gammaproteobacteria	0.6729	2015-04-11
Otu00219	Betaproteobacteria	0.7503	2014-04-18
Otu00002	Actinobacteria	0.3190	2015-05-03
Otu00008	Actinobacteria	0.3637	2013-05-09
Otu00022	Opitutae	0.8003	2015-05-03
Otu00031	Cytophagia	0.6593	2014-05-09
Otu00051	Flavobacteriia	0.9187	2013-05-09
Otu00062	Flavobacteriia	0.7836	2015-05-23
Otu00064	Alphaproteobacteria	0.6964	2013-05-29
Otu00113	Bacteroidetes sp.	0.7950	2013-05-09
Otu00116	Betaproteobacteria	0.8373	2014-05-09
Otu00151	Betaproteobacteria	0.7259	2013-05-17
Otu00183	Bacteria sp.	0.6090	2015-05-03
Otu00200	Bacteria sp.	0.6851	2015-05-23
Otu00083	Flavobacteriia	1.0031	2015-06-06
Otu00087	Betaproteobacteria	0.5873	2014-06-05
Otu00098	Betaproteobacteria	0.8104	2013-06-14
Otu00123	Sphingobacteriia	0.7259	2014-06-20
Otu00194	Deltaproteobacteria	0.9277	2014-06-13

Otu00294	Alphaproteobacteria	0.6964	2013-06-21
Otu00004	Actinobacteria	0.3500	2015-07-11
Otu00009	Gammaproteobacteria	1.3329	2013-07-26
Otu00011	Betaproteobacteria	0.9841	2015-07-18
Otu00192	Bacteria sp.	0.4350	2013-07-26
Otu00195	Actinobacteria	0.6277	2014-07-18
Otu00020	Betaproteobacteria	0.4150	2013-08-01
Otu00026	Betaproteobacteria	0.6277	2013-08-01
Otu00029	Actinobacteria	0.6593	2013-08-23
Otu00036	Alphaproteobacteria	0.7576	2013-08-16
Otu00037	Actinobacteria	0.7646	2014-08-29
Otu00052	Alphaproteobacteria	0.6444	2013-08-09
Otu00076	Actinobacteria	0.6444	2014-08-08
Otu00112	Alphaproteobacteria	0.5873	2014-08-23
Otu00226	Opitutae	0.6729	2015-08-02
Otu00250	Actinobacteria	0.5306	2013-08-23
Otu00010	Proteobacteria sp.	0.6277	2014-09-26
Otu00024	Bacteroidetes sp.	0.7259	2014-09-19
Otu00056	Cytophagia	0.7711	2013-09-06
Otu00067	Betaproteobacteria	0.5306	2015-09-02
Otu00177	Proteobacteria sp.	0.5306	2013-09-20
Otu00005	Sphingobacteriia	0.5294	2014-10-04
Otu00015	Actinobacteria	0.6277	2014-10-17

Otu00018	Gammaproteobacteria	0.9516	2014-10-11
Otu00060	Betaproteobacteria	1.0031	2013-10-25
Otu00066	Betaproteobacteria	0.8153	2013-10-25
Otu00082	Bacteroidetes sp.	0.8003	2014-10-04
Otu00154	Alphaproteobacteria	0.7070	2014-10-04
Otu00158	Gammaproteobacteria	0.9141	2013-10-04
Otu00001	Betaproteobacteria	0.2860	2013-11-15
Otu00007	Betaproteobacteria	0.4577	2013-11-15
Otu00038	Actinobacteria	0.7346	2014-11-29
Otu00047	Betaproteobacteria	0.6729	2013-11-15
Otu00073	Betaproteobacteria	0.4906	2013-11-22
Otu00077	Flavobacteriia	0.7774	2013-11-15
Otu00109	Actinobacteria	0.6729	2013-11-15
Otu00118	Actinobacteria	0.6090	2013-11-22
Otu00129	Alphaproteobacteria	0.4906	2013-11-15
Otu00198	Betaproteobacteria	0.9210	2013-11-15
Otu00208	Betaproteobacteria	0.6964	2013-11-22
Otu00217	Proteobacteria sp.	0.7167	2013-11-22

Figure S1. Temporal partitioning of maximum growth rates corresponds to environmental

conditions. Here, we depict a principal component analysis of the environmental conditions at

each date. OTU labels indicate the individual taxa that exhibit maximum growth at the

corresponding time of the year. Vectors indicate the loadings of environmental variables along

the two PC axes. Warmer colors correspond to spring and summer months, cooler colors

correspond to winter months.

Figure S2. Negative frequency dependence (NFD) in the active portion of the community for the

82 persistent bacterial taxa. (A) Relationship between the rate of change of an OTU and its

relative abundance. Depicted in this graph are simple linear regression fits for the 82 taxa

individually, data points not shown to reduce clutter. Negative relationships indicate NFD

growth and variation in slopes indicates variation in the strength of NFD. (B) Rare taxa (lower

equilibrium frequencies) exhibit stronger NFD, while common taxa (higher equilibrium

frequency) have weaker NFD.