The landscape of metabolic pathway dependencies in cancer cell lines

James H. Joly[1][†], Brandon T.L. Chew[1][#], and Nicholas A. Graham[1,2,3]

[1]Mork Family Department of Chemical Engineering and Materials Science,

[2]Norris Comprehensive Cancer Center, University of Southern California

[3]Leonard Davis School of Gerontology,

University of Southern California, Los Angeles, CA 90089

[†]Present address: Nautilus Biotechnology, San Carlos, CA

[#]Present address: Department of Pharmaceutical Sciences and Pharmacogenomics, University of California, San Francisco

Running title:

Metabolic pathway dependencies in cancer

To whom correspondence should be addressed: Nicholas A. Graham, University of Southern California, Los Angeles, 3710 McClintock Ave., RTH 509, Los Angeles, CA 90089. Phone: 213-240-0449; E-mail: nagraham@usc.edu

Abstract

The metabolic reprogramming of cancer cells creates metabolic vulnerabilities that can be therapeutically targeted. However, our understanding of metabolic dependencies and the pathway crosstalk that creates these vulnerabilities in cancer cells remains incomplete. Here, by integrating gene expression data with genetic loss-of-function and pharmacological screening data from hundreds of cancer cell lines, we identified metabolic vulnerabilities at the level of pathways rather than individual genes. This approach revealed that metabolic pathway dependencies are highly context-specific such that cancer cells are vulnerable to inhibition of one metabolic pathway only when activity of another metabolic pathway is altered. Notably, we also found that the no single metabolic pathway was universally essential, suggesting that cancer cells are not invariably dependent on any metabolic pathway. In addition, we confirmed that cell culture medium is a major confounding factor for the analysis of metabolic pathway vulnerabilities. Nevertheless, we found robust associations between metabolic pathway activity and sensitivity to clinically approved drugs that were independent of cell culture medium. Lastly, we used parallel integration of pharmacological and genetic dependency data to confidently identify metabolic pathway vulnerabilities. Taken together, this study serves as a comprehensive characterization of the landscape of metabolic pathway vulnerabilities in cancer cell lines.

Author Summary

Cancer cells rewire their metabolism, which creates targetable metabolic vulnerabilities.

Previous analyses of metabolic vulnerabilities in cancer cells have been limited to the analysis of

individual genes or metabolites. However, metabolic *pathways* exhibit significant cross talk and

compensation for one another. We developed a computational method to answer the question:

when a metabolic pathway's activity is high, which other metabolic pathways become more

essential or less essential? By integrating genetic screen data with drug response data from FDA

approved drugs, we identified cancer cell line dependence on metabolic *pathways* as opposed to

individual *genes*. For example, we found that identifying key regulators of metabolic pathways,

such as the Pentose Phosphate Pathway, may serve as a biomarker to identify which patients

may benefit from antifolate chemotherapies (e.g. methotrexate, 5-fluorouracil). The efforts

outlined here serve as the first characterization of the landscape of metabolic pathway

vulnerabilities in cancer cell lines. Our results demonstrate the benefit of analyzing dependencies

on metabolic pathways as opposed to metabolic genes.

Introduction

The reprogramming of cellular metabolism was one of the earliest discovered hallmarks

of cancer [1]. Cancer cells rewire their metabolism to satisfy the bioenergetic, biosynthetic, and

redox demands of tumors. In turn, these metabolic adaptations create cancer-specific

vulnerabilities that can be therapeutically targeted [2]. Much research has focused on how

individual mutations or DNA copy number alterations reprogram tumor metabolism and create

therapeutic opportunities [3–7]. For example, leukemias and gliomas with mutations in isocitrate

dehydrogenase (IDH) are sensitive to inhibitors specific to mutant IDH [8,9]. In addition, some

tumors are sensitive to depletion or restriction of amino acids, including cysteine in tumors with

deleted methylthioadenosine phosphorylase (*MTAP*) [10], serine in tumors with deleted p53

(*TP53*) [11], asparagine in leukemias with low expression of asparagine synthetase (*ASNS*) [12],

asparagine in tumors with inhibition of the electron transport chain [13], and methionine in RAS-

driven tumors [14]. Other efforts have identified metabolic vulnerabilities related to redox balance

including in cancer cells with dysregulated PI3K/AKT signaling [15] or overexpression of the L-

glutamate/L-cystine antiporter SLC7A11 [16,17].

While these studies have proved fruitful for advancing the therapeutic targeting of cancer

metabolism [2], they have traditionally been limited to the study of individual genes. However,

metabolic pathways consist of multiple enzymes which collectively regulate metabolic flux.

Furthermore, metabolic pathways often have similar byproducts (e.g. ATP, NAD(P)H), suggesting

that pathways may compensate for one another under different contexts. For example, NADPH

can be produced by three different pathways: the oxidative pentose phosphate pathway, serine-

driven one-carbon metabolism, and malic enzyme [18]. These three pathways can presumably

be up- and down- regulated to respond to changes in activity of the other pathways. Since these

compensatory mechanisms exist across human metabolism, studying the effects of individual

genes may not reflect cancer cell metabolic vulnerabilities at the pathway level. As a result, our

understanding of cancer cell dependency on metabolic pathways remains incomplete.

Recent developments in large scale CRISPR-based genetic [19,20] and pharmacologic

screening [21] along with large panels of comprehensively characterized cancer cell lines [22]

have proved powerful tools for identification of genes essential for cancer cell survival [23],

elucidation of drug mechanism-of-action [19,24,25], and discovery of novel candidate drug targets

[26,27]. Furthermore, parallel integration of both pharmacologic and gene loss-of-function data

has been used to identify drug mechanism(s) of action [25,28–30]. While these databases have

served as a rich resource to explore individual gene vulnerabilities and drug sensitivities, there

exists a need to probe these datasets on the pathway level.

Here, we aimed to identify cancer cell dependencies on metabolic pathways rather than

individual metabolic genes. To do so, we used gene expression data from the Cancer Cell Line

Encyclopedia (CCLE) to infer metabolic pathway activity and then integrated these pathway

activities with data from genetic and pharmacologic screens across hundreds of cell lines. We

show that this approach provides a comprehensive characterization of the landscape of metabolic

pathway dependencies in cancer cell lines. In addition to demonstrating the context-specific

nature of metabolic pathway dependence, we identified both known and novel metabolic

vulnerabilities, discovered robust associations between drug response and metabolic pathway

activity, and independently found metabolic pathway essentialities in both genetic and

pharmacological screens. Collectively, we present an approach to integrate gene expression,

gene dependency, and drug response data to identify cancer cell dependencies on metabolic

pathways.

Results

Genetic Pathway Dependency Enrichment Analysis Identifies Metabolic Pathway Dependencies

in Genetic Screens

To identify metabolic pathway dependencies, we analyzed gene expression data and

CRISPR-Cas9 loss-of-function screens from 689 cancer cell lines overlapping between the

Cancer Cell Line Encyclopedia (CCLE) [22,26] and the Cancer Dependency Map [20]. First, we

inferred metabolic pathway activity for each cell line using single-sample gene set enrichment

analysis (ssGSEA) of the RNAseq data from each cell line [31]. To focus on metabolism, we

queried 69 metabolic pathways from the Kyoto Encyclopedia of Genes and Genomes (KEGG)

[32]. We chose to focus on pathways that are expressed in human metabolism and had more

than five genes in each set. Because metabolism is influenced by culture type [33] and culture

medium [23], we first divided cancer cell lines by culture type (e.g., adherent v. suspension

culture) and media (e.g., RPMI v. DMEM) (**Fig. 1A**). Cell lines without annotations for either of

these features were removed, leaving 300 adherent cell lines cultured in RPMI, 153 adherent cell

lines cultured in DMEM, 66 suspension cell lines cultured in RPMI, and 2 suspension cell lines

cultured in DMEM. Since the number of suspension cell lines was small, we focused our analysis

on adherent cell lines. The resulting ssGSEA normalized enrichment scores (ssNESs) represent

the metabolic pathway activity relative to all other cell lines within the respective cell culture

medium. Next, we correlated the cell line-specific NESs for each metabolic pathway with cell

fitness effects from CRISPR-Cas9 loss-of-function screens (16,643 gene knockouts). Here, each

correlation coefficient represents the association between metabolic pathway activity and gene

essentiality, with positive values representing increased gene dependency in cell lines with

increased metabolic pathway activity. Conversely, a negative correlation indicates increased

gene dependency in cell lines with decreased metabolic pathway activity. Finally, to measure the

130
131
Figure 1. Integration of gene expression and CRISPR gene dependencies to identify
metabolic pathway dependencies. A) Schematic outlining the approach for Genetic Pathway
Dependency Enrichment Analysis (Genetic PDEA). Cancer cell lines from the CCLE were first
stratified by culture type (adherent, suspension) and culture medium (RPMI, DMEM), and then
their metabolic pathway activity was inferred using single-sample GSEA (ssGSEA). The resulting
pathway activities were integrated with gene dependency to assess association with metabolic
pathway activity. **B-C)** Simulated data (see methods) was used to assess the sensitivity of the
Genetic PDEA approach. The heatmaps represent the percentage of significant results at each
gradient added. Values added to the expression gradient resulted in slightly stronger correlation
coefficients and Genetic PDEA results compared to values added to dependency gradient.
142

essentiality of the entire metabolic pathway, as opposed to individual genes, we then ranked the

resulting 16,643 correlation coefficients and analyzed the rank list using GSEA again querying

the KEGG metabolic pathways. Here, positive NES values represent increased pathway

dependency upon increased pathway activity, whereas negative NES values represent increased

pathway dependency upon decreased pathway activity. Because this approach integrates the

essentiality of all genes across a metabolic pathway into a single metric of pathway dependency,

we termed this approach genetic pathway dependency enrichment analysis (Genetic PDEA).

To analyze the sensitivity of our Genetic PDEA approach, we analyzed simulated gene

expression and gene dependency data using the pipeline outlined in Fig. 1A. Gene expression

data (16,643 genes) was simulated for 300 cell lines using a normal distribution for each cell line

($\mu = 0$, $\sigma = 0.5$) which reflects the shape of CCLE gene expression data (**Supp. Fig 1**). Then, a

synthetic gene set of 25 genes was perturbed using a normal distribution gradient. In cell line 1,

the 25 genes were replaced with values from a normal distribution with $\mu = -X$, $\sigma = 0.5$, and in cell

line 300, the 25 genes were replaced with values from a normal distribution with $\mu = +X$, $\sigma = 0.5$.

For cell lines 2-299, the 25 genes were replaced with values from normal distributions with μ

sequentially increasing from -X to X. ssGSEA NESs were then calculated for the synthetic gene

set for all 300 cell lines. Next, gene dependency data was simulated for the same 300 cell lines

using a similar normal distribution gradient method. For both gene expression and gene

dependency data, values for the perturbation X were varied from 0 to 0.5. Then, Spearman

correlation coefficients between synthetic gene set activity (ssGSEA NESs) and gene

dependency were calculated for all 16,643 genes. Finally, GSEA was run to calculate the

simulated Genetic PDEA values as outlined in Fig. 1A. Analyzing 50 replicates of this simulation

pipeline, we found that both the correlation coefficients and Genetic PDEA NES results were more

strongly influenced by expression gradients added than dependency gradients added (**Fig. 1B,C**).

This suggests that our Genetic PDEA approach to identify metabolic pathway vulnerabilities in

cancer cells is more sensitive to changes in metabolic pathway activity than changes in gene

dependency. However, when the perturbation X was large for the dependency gradient and small

for the expression gradient (or vice versa), significant Genetic PDEA NES values were still

obtained.

Having validated the sensitivity of our Genetic PDEA approach, we next investigated

cancer cell line dependency across all KEGG metabolic pathways. We tested a total of 4,692

metabolic pathway combinations and identified 190 significant enrichments (FDR < 0.05) between

metabolic pathway activity and pathway dependency in the adherent RPMI data set and an

additional 190 significant enrichments in the adherent DMEM data set (**Supp. Table 1**). Leading

edge genes for each significant pathway dependency are listed in **Supp. Table 1**. We next

clustered Genetic PDEA NES values across all pathway activities (columns) and pathway

dependencies (rows) (**Fig. 2A & Supp. Fig. 2A**) and found that related pathways clustered

together on the x-axis (pathway activity) but not on the y-axis (pathway dependency). This

suggests that pathway activity but not pathway dependency is similar within a group of related

pathways. For example, glycan biosynthesis pathways exhibit similar correlations with activity of

other glycan pathways (columns) but differ in their own dependency (rows). We also found that

the no single metabolic pathway had universally positive or negative NES values, suggesting that

cancer cells are not universally dependent on any metabolic pathway. Interestingly, the strongest

metabolic dependency of adherent RPMI cancer cell lines was Folate Biosynthesis (hsa00790)

when One-Carbon Pool by Folate (hsa00670) pathway activity was high (**Fig. 2B**). This

enrichment was driven by the genes *QDPR*, *ALPI*, *ALPP*, and *GCH1*. Notably, Folate

Biosynthesis is directly upstream of One-Carbon Pool by Folate. Furthermore, one of the

strongest Genetic PDEA results in both the adherent DMEM and adherent RPMI analyses

recapitulated a link between increased dependency on One-Carbon Pool by Folate in cells with

increased TCA cycle activity [34] (**Fig. 2C & Supp. Fig. 2B**). Additionally, some metabolic

Figure 2. Global analysis of metabolic dependency data reveals context-specific pathway essentialities. A) Metabolic pathway activity was inferred using ssGSEA for 300 adherent cell lines cultured in RPMI and correlated to gene dependency data from The Cancer Dependency

Map (DepMap). Correlation coefficients were then ranked and Genetic Pathway Dependency
Enrichment Analysis (Genetic PDEA) was run using the KEGG metabolic pathways (see Fig. 1).
Hierarchical clustering was performed on the Genetic PDEA normalized enrichment scores
(NES). Results for pathways with FDR < 0.25 are plotted. Dots are colored according to their NES
and sized according to the -$\log_{10}$ of the false discovery rate (FDR). Numerical values for each
pathway can be found in Supp. Table 1. Results shown in B and C are highlighted with a black
outline. **B)** Cancer cell dependency on Folate Biosynthesis (hsa00790) was increased when One-
Carbon Pool by Folate (hsa00670) pathway activity was high. The scatter plots of pathway activity
NES and gene dependency (-CERES) for leading-edge genes *QDPR* and *ALPI* are shown**. C)**
Dependency on One-Carbon Pool by Folate metabolism (hsa00670) is increased when TCA cycle
(hsa00020) activity is increased. The scatter plots of pathway activity NES and gene dependency
(-CERES) for leading-edge genes *MTR* and *MTHFD1* are shown.

pathways exhibited context-specific dependencies. For example, the dependency on the TCA

Cycle (hsa00020) was increased in adherent RPMI cell lines with decreased Glycolysis-

Gluconeogenesis (hsa00010) activity, whereas dependency on the TCA cycle was increased in

adherent RPMI cell lines with increased Pentose Phosphate Pathway (hsa00030) activity (**Supp.**

Fig. 3). This new finding suggests that the diversion of glucose from glycolysis to the pentose

phosphate pathway may confer increased dependency on the TCA Cycle. We next asked whether

there existed a general relationship between a metabolic pathway's activity and its own

essentiality. Of the 69 metabolic pathways queried, 36 had a negative NES and 33 had a positive

NES for adherent RPMI cell lines (**Supp. Fig. 4A, Supp. Table 2**). A similar distribution was

observed in DMEM, although the pathways with positive and negative NES values were not the

same as in RPMI (**Supp. Fig. 4B**). While there was not general agreement between the self-

dependencies, one gene set did exhibit significant Genetic PDEA NES for both Adherent RPMI

and Adherent DMEM cells. Specifically, Riboflavin Metabolism (hsa00740) exhibited significant

negative Genetic PDEA NES, indicating that when Riboflavin Metabolism activity is low, the

dependency on Riboflavin Metabolism genes increases. Taken together, these results suggest

that there is no generic rule regarding a metabolic pathway's activity and its essentiality. Rather,

these results indicate that metabolic pathway dependency is highly context specific such that

metabolic pathway activity influences metabolic pathway essentiality.

Validation of Genetic PDEA results using other CRISPR data sets and pathway activity metrics

We next wanted to examine whether these findings were reproducible across data sets

and methods. To examine whether these findings were reproducible, we analyzed data from

another large-scale pan-cancer CRISPR-Cas9 gene dependency data set (Sanger Institute) [35]

using the Genetic PDEA pipeline. Since the underlying gene expression profiles were derived

from the CCLE for both Sanger and DepMap Genetic PDEAs, we combined statistical tests using

the harmonic mean p-value (HMP) which combines dependent statistical tests while controlling

the family-wise error rate (FWER) [36]. Applying an HMP threshold of 0.05, we found that 96%

and 90% of significant results in the DepMap Genetic PDEA were recapitulated in the combined

DepMap and Sanger Genetic PDEA results for Adherent RPMI and Adherent DMEM cells,

respectively (**Supp. Fig. 5** and **Supp. Table 3**).

Next, we tested the effect of metrics other than ssGSEA for inference of metabolic pathway

activity. We chose to use normalized weighted average expression (NWAS), which accounts for

overlap between gene sets [37]. To directly compare NWAS with ssGSEA, we re-ran our pipeline

using NWAS to analyze dependency on the Pentose Phosphate Pathway which contains several

enzymes that are shared with other metabolic pathways (e.g., PRKL, PFKM, and PKFP are

present in both Glycolysis-Gluconeogenesis and Pentose Phosphate Pathway gene sets). We

found broad agreement between the metabolic pathway dependencies when using either NWAS

or ssGSEA for both Adherent RPMI and Adherent DMEM cell lines (Spearman r of 0.606 and

0.691, respectively) (**Supp. Fig. 6**). Taken together, these findings demonstrate that our Genetic

PDEA results for metabolic pathway dependencies are reproducible across different data sets

and methods.

Figure 3. Media composition influences metabolic pathway dependency. For adherent cancer cell lines cultured in RPMI (Fig. 2) and DMEM (Supp. Fig. 2), the metabolic pathway dependency NESs from Genetic PDEA analysis were weighted by -$\log_{10}$ FDR. The weighted NESs were then averaged across all 69 KEGG metabolic pathways. Pathways are ranked by the difference between DMEM and RPMI. The relative media composition between RPMI and DMEM are shown on the right on a purple to green heat map with the relevant metabolite(s) indicated. For pathways with multiple metabolites, the average of the metabolites was taken. For example, the concentration of folate in RPMI and DMEM is 1 mg/L and 4 mg/L, respectively. Folate is shown twice because it is both the product of Folate Biosynthesis and the input to One-Carbon Pool by Folate. The dependency on Folate Biosynthesis was much higher in RPMI than in DMEM because these cells must synthesize more folate. Conversely, the dependency on oxidative phosphorylation is much higher in DMEM. This may be due to differences in aspartate levels (RPMI 150 μM, DMEM 0 μM). The indicated pathways are highlighted in bold. Overall, pathways that contained a metabolite which is differentially abundant in cell culture media exhibited a

269   significant difference in pathway essentiality in DMEM and RPMI (p = 3.9×10^{-4} by paired Mann-
270   Whitney U test). In contrast, pathways that do not contain differentially abundant cell culture media
271   metabolites did not exhibit significantly different pathway dependencies (p = 0.545 by paired
272   Mann-Whitney U test).
273

Media Composition Influences Metabolic Pathway Dependency

We next investigated whether cancer cell line metabolic pathway dependencies were

influenced by media composition. To assess the essentiality of a pathway, we weighted each NES

from Genetic PDEA by its -$\log_{10}$ FDR and then took the mean of all weighted NESs. We found

striking differences between DMEM and RPMI metabolic pathway essentialities that can be partly

explained by media composition (**Fig. 3**), consistent with the finding that the essentiality of

individual metabolic genes is influenced by culture medium [23]. For example, cancer cells

cultured in RPMI exhibited a strongly positive average weighted NES for Folate Biosynthesis

(hsa00790) whereas cancer cells cultured in DMEM did not. Notably, DMEM contains four times

the concentration of folate (4 mg/L) compared to RPMI (1 mg/L), suggesting that cancer cells

grown in DMEM need to synthesize less folate, thereby reducing their dependency on Folate

Biosynthesis. Similarly, cancer cells grown in DMEM were more dependent on Oxidative

Phosphorylation (hsa00190) than cancer cells grown in RPMI. One function of oxidative

phosphorylation is to enable aspartate synthesis to accept electrons from the electron transport

chain [38,39]. Since RPMI and DMEM contain 150 µM and 0 µM aspartate, respectively, the

increased dependency on oxidative phosphorylation in DMEM may reflect an increased need for

aspartate synthesis. Interestingly, we did not observe a strong difference in dependency on

Aspartate, Alanine, and Glutamate metabolism (hsa00250) between RPMI and DMEM despite

the difference in aspartate concentrations. We posit that the inclusion of alanine and glutamate

metabolism genes may be a confounding factor for analyzing aspartate dependency.

To test whether media composition globally affected metabolic pathway abundance, we

divided metabolic pathways into two groups: 1) pathways that contain metabolites with differential

abundance in cell culture media (e.g., Glycolysis-Gluconeogenesis (glucose), Folate biosynthesis

(folate)); and 2) pathways that do not contain metabolites with differential abundance in cell

culture media (e.g., fatty acid metabolism, fructose and mannose metabolism). We then tested

whether the metabolic pathway dependency in RPMI and DMEM differed for both groups of

pathways. Notably, we found that pathways that contain differentially abundant metabolites in cell

culture media exhibited significantly different pathway essentialities in DMEM and RPMI (p =

3.9×10^{-4}). In contrast, pathways that do not contain metabolites that are differentially abundant in

cell culture media did not exhibit significantly different pathway dependencies in DMEM and RPMI

(p = 0.545). Taken together, these results suggest that media composition influences cancer cell

line metabolic pathway dependency and that future studies of metabolic vulnerabilities should

take media composition into consideration.

Metabolic pathway activity is correlated with anti-cancer drug sensitivity

We next sought to integrate metabolic pathway activity with large scale pharmacologic

screens. We therefore used the PRISM drug repurposing database [21], which contains 1,448

compounds screened against 499 cell lines at 8 different doses. Once again, cell lines were

separately processed by culture type and culture medium with a focus on adherent cell lines.

Compounds measured in less than 150 cell lines were removed from the analysis, leaving 1,390

compounds. We then correlated drug response area-under-the-curve (AUC) with metabolic

pathway activity after multiplying the response of drugs classified as positive regulators (e.g.,

agonists) by -1 for directional consistency (**Fig. 4A**). Here, the AUC represents the dose

dependent effect of a drug on cell growth, with a lower AUC representing a stronger response to

Figure 4. Metabolic pathway activity identifies anti-cancer drug sensitivity independent of cell culture medium. A) Schematic representing the strategy used to integrate metabolic pathway activity with drug response screens. Cancer cell lines were separately processed by culture type and culture medium with a focus on adherent cell lines. All correlation p-values were FDR corrected using a Benjamini-Hochberg correction. **B-E)** Scatter plots of significant drug:metabolic pathway combinations (FDR < 0.05) in both DMEM and RPMI mediums. Correlation coefficients and FDR corrected p-values are shown for each correlation. The annotated gene target of each drug is listed below the drug name. The remaining significant associations are listed in Supp. Table 2.

drug [21]. First, we asked whether there were any drugs with directional agreement across the

RPMI and DMEM analyses. From a possible 101,360 possible drug:metabolic pathway

combinations, 66 combinations passed FDR-corrected significance thresholds (q < 0.05) and

were of the same sign in both RPMI and DMEM (**Supp. Table 4**). Notably, zero results that passed

FDR correction were of different sign. Many of the common associations in RPMI and DMEM

were tyrosine kinase inhibitors (TKIs), which have been extensively linked to metabolism [40].

Interestingly, we found a strong association between decreased Core Glycolysis (hsa_M00001)

pathway activity and increased sensitivity to AZD8931, an inhibitor of EGFR and ERBB2 (HER2)

(**Fig. 4B**). We also found a strong association between increased α-linoleic acid metabolism

(hsa00592) and sensitivity to afatinib, another EGFR inhibitor (**Fig. 4C**). Of the non-TKI results,

we found a link between decreased phenylalanine metabolism (hsa00360) and increased

sensitivity to atorvastatin, an HMGR inhibitor (**Fig. 4D**). HMGR is the rate limiting enzyme in the

cholesterol biosynthetic pathway [41] and multiple reports have suggested that elevated levels of

phenylalanine inhibit cholesterol biosynthesis [42–44]. Increased response to atorvastatin when

phenylalanine metabolism activity is low suggests that decreased phenylalanine metabolism and

HMGR inhibitors may be redundant. Lastly, we found a known link between decreased mucin

type O-glycan biosynthesis pathway (hsa00512) activity and increased sensitivity to the HSP90

inhibitor NMS-E973 [45] (**Fig. 4E**). Taken together, these results indicate that metabolic pathway

activity can be associated with anti-cancer drug sensitivity independent of cell culture medium.

Pharmacological Pathway Dependency Enrichment Analysis Reveals Common Metabolic

Pathway Vulnerabilities

Having identified strong associations between metabolic pathway activity and individual

drugs, we next asked whether there were commonalities in the response of cancer cell lines to

families of metabolic pathway inhibitors. Drugs were mapped to metabolic pathways using their

annotated target(s) and grouped according to the KEGG metabolic pathways database. Drugs

were also classified as positive or negative regulators based on their annotated mechanism of

action. For example, drugs labeled as "agonists" or "activators" were classified as positive

regulators whereas "blockers" and "antagonists" were classified as negative regulators. To enable

a consistent pathway analysis, correlation coefficients for positive regulators were multiplied by -

1. Pathways with less than 4 drugs were omitted from the analysis, leaving 46 sets of drugs

targeting metabolic pathways. We then analyzed the enrichment of these metabolic pathway

inhibitors in the rank list of drug sensitivity-metabolic pathway activity correlation coefficients

(n=1,390). We termed this approach Pharmacological Pathway Dependency Enrichment Analysis

(Pharmacological PDEA) (**Supp. Fig. 7A**). To test the sensitivity of this approach, we again

performed a simulation study with simulated gene expression data and simulated drug sensitivity

(1,390 drugs). This approach was analogous to the simulation study of Figure 1B with 1,390 drug

sensitivities replacing 16,643 gene dependencies. Like Genetic PDEA, we found that expression

gradients resulted in stronger results than dependency gradients for both individual drug

correlation coefficients and Pharmacological PDEA (**Supp. Fig. 7B-C**).

We then clustered the pharmacological PDEA NESs and found that pathways with similar

function clustered together based on pathway activity (columns) more than with their dependency

profile across pathways (rows) (**Fig. 5A and Supp. Fig. 8A**), similar to our results with Genetic

PDEA (**Fig. 2**). For example, the activities of Core Glycolysis (hsa_M00001), Fructose and

Mannose Metabolism (hsa00051), Starch and Sucrose Metabolism (hsa00500), and Pentose

Phosphate Pathway (hsa00030) were clustered together in the adherent RPMI cell lines. Among

the strongest Pharmacological PDEA results, we found that sensitivity to inhibitors of Terpenoid

Backbone Biosynthesis (hsa00900) was increased in adherent RPMI cancer cells with high

Alanine, Aspartate, and Glutamate Metabolism (hsa00250) (**Fig. 5B**). We also found an

Figure 5. Pharmacological PDEA reveals consistent metabolic pathway vulnerabilities in Adherent RPMI cell lines. A) Pharmacological PDEA (Supp. Fig. 4) was performed on 1,390 anti-cancer drugs from the PRISM database. Drugs were mapped to metabolic pathways by their annotated target(s) and then the enrichment of these metabolic pathway inhibitors was analyzed in the rank list of drug sensitivity-metabolic pathway activity correlation coefficients. Hierarchical

clustering was performed on NES values, and results with FDR < 0.25 are plotted. Dots are
colored according to the NES and sized according to the -log$_{10}$ FDR. Dots with black outline
correspond to results shown in panels B-C and F. **B-C)** Increased Alanine, Aspartate, and
Glutamate metabolism (hsa00250) correlates with increased response to inhibitors of terpenoid
backbone biosynthesis. In contrast, decreased Pentose Phosphate Pathway metabolism
correlates with increased response to inhibitors of Folate Biosynthesis (hsa00790). **D-E)** Inhibitors
of Folate Biosynthesis (hsa00790) are more effective when overall metabolic pathway expression
is low, whereas inhibitors of Ascorbate and Aldarate Metabolism (hsa00053) are more effective
when overall metabolic pathway expression is high. **F)** Representative mountain plots and the
drug(s) driving enrichment of metabolic pathway activities that strongly correlate with response to
inhibitors of Ascorbate and Aldarate metabolism are shown.
397

interesting link between decreased pentose phosphate pathway (PPP) pathway activity and

increased sensitivity to folate biosynthesis inhibitors (**Fig. 5C**). Because folate biosynthesis

inhibitors prevent the generation of NADPH via one-carbon metabolism, these inhibitors may be

more damaging to cellular redox balance when PPP expression is low. In fact, most strong results

for inhibitors of folate biosynthesis occur when overall metabolic pathway activity is low (**Fig. 5D**

and Supp. Fig. 8B). Conversely, inhibitors of Ascorbate and Aldarate metabolism (hsa00053)

are more effective when overall metabolic pathway activity is high (**Fig. 5E**). This may be because

ascorbate (also known as Vitamin C) is an effective antioxidant used to detoxify reactive oxygen

species (ROS). ROS are a byproduct of many metabolic reactions such as oxidative

phosphorylation and methionine metabolism, which indirectly produces ROS by supporting

polyamine synthesis [10]. Indeed, the sensitivity to inhibitors of Ascorbate and Aldarate

metabolism is stronger when expression of these ROS producing pathways is high (**Fig. 5F**)

suggesting that ascorbate's role as an antioxidant is crucial in this context. Taken together, these

results reveal contexts in which pharmacological inhibition of metabolic pathways results in

decreased cell survival.

Integration of pharmacologic and genetic screens reveals consistent metabolic vulnerabilities

Next, we sought to integrate results from genetic and pharmacological screen data to

identify consistent metabolic pathway dependencies found independently in both analyses. First,

we integrated individual gene dependency correlations with their corresponding drug sensitivity

correlations by first annotating each drug with its gene target(s). We then summed the gene

dependency correlation coefficient and the drug sensitivity correlation coefficient for each drug

target and assessed significance by a permutation test with FDR correction (**Fig. 6A**). Out of

187,818 gene+drug:metabolic pathway combinations, we found 176 results that passed an FDR-

corrected significance threshold of 0.01 (**Supp. Table 6**). Interestingly, all significant results

targeted known cancer driver genes such as *EGFR*, *HER2*, *PIK3CA*, and *BRAF* (**Fig. 6B-E**).

These results included a known interaction between HER2 inhibitors and retinol metabolism,

whereby increased retinol metabolism enhances sensitivity to HER2 inhibition [46] (**Fig. 6C**).

Additionally, some results identified well known molecular interactions, such as *BRAF* and

PIK3CA driving sugar metabolism [47,48] (**Fig. 6D,E**). These results demonstrate robust

associations between metabolic pathway activity, gene dependency, and drug response.

Lastly, we sought to integrate the results from Genetic PDEA and Pharmacological PDEA

to identify metabolic pathway vulnerabilities that were consistent between gene dependency and

drug response data (**Fig. 6F, Supp. Table 7**). By applying p-value and q-value filters to each

analysis, we found three consistent vulnerabilities that were significant in both analyses. First, we

found that when tyrosine metabolism is high, there is an increased vulnerability to inhibition or

knockout of terpenoid backbone biosynthesis genes (**Fig. 6G**). Interestingly, the drugs driving the

enrichment (fluvastatin and pitavastatin) target the protein with the largest gene dependency

within the pathway, HMGCR. We also found a common vulnerability between inactivation of the

folate biosynthesis pathway and decreased aminoacyl tRNA biosynthesis (**Fig. 6H**). Here, the top

439
Figure 6. Integration of pharmacological and genetic screens reveals consistent metabolic
pathway vulnerabilities in adherent RPMI cell lines. A) Schematic outlining approach to

identify drug targets and genetic dependencies that are commonly increased or decreased with
metabolic pathway activity. Significance was assessed by permutation testing combined with
Benjamini-Hochberg FDR correction. 176 significant associations of 187,818
gene+drug:metabolic pathway combinations passed the FDR threshold of 0.01 (Supp. Table 6).
B-E) Scatter plots of four drug response and CRISPR gene dependencies associated with
metabolic pathway activity. The gene target of each drug is listed below the drug name. **F)**
Schematic outlining a filtering approach used to identify common pathway-level vulnerabilities in
Genetic PDEA and Pharmacologic PDEA. **G-I)** Mountain plots and leading edge drugs and genes
from the three common pathway vulnerabilities are shown.
451

hits in Genetic PDEA and Pharmacological PDEA did not converge on a single protein product.

Nevertheless, these results indicate that inactivation of the folate biosynthesis pathway is more

effective at slowing cancer cell growth when aminoacyl-tRNA biosynthesis pathway activity is low.

Lastly, we found a strong association between inhibitors and gene knockouts of terpenoid

backbone biosynthesis when pathway activity for biosynthesis of heparan sulfate is low (**Fig. 6I**).

Once again, the targets of the statins driving the Pharmacological PDEA enrichment did not align

with the top gene dependencies (*DHDDS*, *HMGCS1*). Taken together, these results demonstrate

common metabolic pathway vulnerabilities by integrating gene dependency, drug response, and

gene expression data.

Discussion

Traditionally, the analysis of gene essentiality in cancer cells has been limited to

identification of individual genes required by cancer cells. Here, using metabolic pathways as an

example, we have demonstrated an approach for identifying cancer cell dependencies at the level

of pathways rather than individual genes. Illustrating the utility of our approach, we recapitulated

known interactions between metabolic pathway activity, drug response, and gene dependency.

These results build on a strong foundation of research identifying metabolic vulnerabilities in

cancer cells [2,23]. Importantly, our results demonstrate that metabolic dependencies in cancer

cells are highly context specific (**Fig. 2**) and are impacted by the nutritional microenvironment

(**Fig. 3**). By investigating metabolic dependencies at the pathway level using genetic and

pharmacological PDEA, we uncovered novel metabolic crosstalk, identified robust associations

between drug response and metabolic pathway activity, and have discovered interactions

between metabolic pathway activity and essentiality.

A recurring theme from our analyses is the importance of the pathways Folate

Biosynthesis (hsa00790) and One-Carbon Pool by Folate (hsa00670). Folate metabolism

supports two key metabolic phenotypes commonly found in cancer cells by producing one-carbon

units for nucleotide synthesis and maintaining redox balance through production of NADPH [49].

In fact, two of the most widely used chemotherapeutics, methotrexate and 5-fluorouracil, target

folate biosynthesis. In our analysis, we found that inhibitors of folate biosynthesis are highly

effective when activity of other metabolic pathways is low (**Fig. 5D**). This may be because the

drugs classified as folate biosynthesis inhibitors are anti-metabolites that cannot be metabolized

by enzymes like thymidylate synthase or dihydrofolate reductase. When adjacent metabolic

pathway expression is low, compensatory mechanisms cannot offset decreased folate

biosynthesis, causing a crisis in both nucleotide synthesis and redox homeostasis. Our results

suggest that identifying biomarkers of activity for key pathways that control the sensitivity of

antifolate pathways like the PPP (**Fig. 5C**) would enable advances in patient selection for

antifolate chemotherapy.

The approach outlined here also sets the stage for the use of metabolic pathways to guide

patient selection to therapy. Patient selection for targeted therapeutics such as EGFR inhibitors

is often based on mutations and copy number alterations (CNAs), but even for these targeted

therapeutics there exists a need to identify additional features that inform patient selection. Here,

we identified metabolic pathways that strongly correlate with both CRISPR knockout and

pharmacological inhibition of the oncogenes EGFR, HER2, BRAF, and PIK3CA (**Fig. 6**). This

supports that metabolic pathways may be effective biomarkers even when mutations and CNAs

in these oncogenes are not present (e.g., α-linoleic acid metabolism for sensitivity to EGFR

inhibitors). Additionally, within patient populations with mutations or CNAs in these oncogenes,

metabolic pathway activity could serve as a biomarker that predicts response to therapy.

Furthermore, our results support that the statins targeting HMG-CoA reductase (*HMGCR*) might

be effective cancer therapeutics in tumors with high tyrosine metabolism and/or low heparan

sulfate glycosaminoglycan biosynthesis. Taken together, our results demonstrate that there may

be patient benefit in analyzing tumor metabolism to inform patient selection for targeted therapies.

Our analysis identified media composition as a major confounding factor when analyzing

cancer cell metabolic pathway dependencies (**Fig. 3**). This finding is consistent with recent studies

and demonstrates the effect to which metabolism and metabolic vulnerabilities are shaped by the

tumor microenvironment [23,50]. As such, our results highlight the importance of formulating cell

culture mediums that better recapitulate the tumor microenvironment [51]. Furthermore, the data

used in this study comes from adherent cell lines cultured on tissue culture plastic. This removes

environmental stresses such as concentration gradients and physical stimuli that cells experience

in real tumors. Recent efforts have demonstrated that CRISPR-Cas9 screens can be performed

in 3D organoids [50]. As this technology becomes more widely used, computational approaches

such as ours can be applied to identify differences in metabolic pathway dependencies between

2D and 3D culture. Similarly, culture conditions that better reflect the physiological conditions of

tumors will enhance the therapeutic relevance of our approach.

While our study identified robust associations between drug response and metabolic

pathway expression, these analyses (**Figs. 5 and 6**) rely on the annotated targets of these drugs.

Off-target toxicity is a major concern when using small molecule inhibitors. In fact, some recent

studies have found that off-target toxicity drives the anti-tumor effect of these compounds [25]. As

such, we cannot exclude the possibility that off-target effects of these compounds could cause

the associations identified here. Furthermore, some compounds in this study are quite

promiscuous with multiple annotated protein targets. This promiscuity confounds the

Pharmacologic PDEA analysis (**Fig. 5**) since some drugs were mapped to multiple metabolic

pathways. As such, the utility of the Pharmacologic PDEA approach lies in the aggregation of

multiple drugs to arrive at a conclusion, rather than treating each individual drug as significant. In

addition, by integrating the Pharmacologic PDEA and Genetic PDEA results (**Fig. 6**), we

strengthened our confidence in the association between metabolic pathway activity and pathway

dependency.

Another potential weakness of our study is that we rely on the inference of metabolic

pathway activity from gene expression data. Gene expression, however, does not always

accurately reflect cellular metabolism. First, proteomic studies have shown that protein expression

does not always correlate with gene expression [52]. Second, metabolic enzyme activity can be

regulated by post-translational modifications [53,54]. By using gene expression data, we have not

accounted for these factors, and as such our analysis may not reflect pathway activity at the

metabolic flux level. We expect that expanding recent efforts to characterize metabolite

abundance [33] and metabolite flux [55] in panels of cancer cell lines will improve our ability to

identify metabolic pathway vulnerabilities by providing better measures of metabolic pathway

activity.

Taken together, this study serves as a comprehensive characterization of the landscape

of metabolic pathway vulnerabilities. Furthermore, our approach serves as a framework for

integrating gene expression, gene dependency, and drug response data to uncover metabolic

dependencies at the level of pathways rather than individual genes. We anticipate this approach

could be extended to other biologically relevant pathways beyond metabolism. Furthermore, the

utility of our approach will increase as CRISPR-Cas9 and pharmacologic screening expand to

include more cancer cell lines, better measure of cellular metabolism, and physiologically relevant

models like 3D organoids.

Methods

Data Sources

Cancer cell line gene expression data was downloaded from the Cancer Cell Line Encyclopedia

(CCLE) version 19Q4. Gene dependency data was downloaded from the Cancer Dependency

Map (DepMap), Achilles gene effect version 19Q4 was used for the main figures in this study and

the Sanger CRISPR (CERES) gene effect data [35] was used for the comparisons in Supp. Fig

5. Drug response data was downloaded from the PRISM Repurposing database, version 19Q4

with secondary screen with dose response curve parameters was used. Metabolic pathway

annotations were downloaded from the Kyoto Encyclopedia of Genes and Genomes (KEGG). We

removed pathways that either 1) are not expressed in human metabolism or 2) contain less than

five genes in the CCLE gene expression data set.

Simulation Studies

Gene expression data was simulated for 300 cell lines using a normal distribution for each cell

line ($\mu = 0$, $\sigma = 0.5$). Then, a synthetic gene set of 25 genes was perturbed using a normal

distribution gradient, where cell line 1 received a value randomly selected from a normal

distribution with $\mu = -X$, $\sigma = 0.5$ and cell line 300 received a value randomly selected from a normal

distribution of $\mu = +X$, $\sigma = 0.5$, with cell lines 2-299 receiving values randomly selected from normal

distributions with sequentially increasing values μ from $-X$ to X. Single-sample Gene Set

Enrichment Analysis (ssGSEA) was calculated for the synthetic gene set for all 300 cell lines.

Next, gene dependency data was simulated for the same 300 cell lines using the same normal

distribution gradient method. For both gene expression and gene dependency data, values for X

were varied from 0 to 0.5. Next, Spearman correlation coefficients between synthetic gene set

activity (NES) and gene dependency were calculated for all 16,643 genes. Finally, Gene Set

Enrichment Analysis was run to calculate the simulated Genetic Pathway Dependency

Enrichment Analysis values. For the Pharmacological PDEA simulation study in Supp. Figure 7,

a similar approach was used with 200 cell lines and 1,390 drugs to simulate the data used in the

Pharmacological PDEA study. For both simulation studies, the pipeline was run for 50 replicates.

Calculation of Metabolic Pathway Expression

1,019 cancer cell lines from the CCLE were separated by their culture type (adherent or

suspension) and then culture medium (RPMI or DMEM), respectively. Cell lines with missing

information for either culture type or medium were omitted. Gene expression values were unit

normalized across all cell lines of the same culture and medium type (e.g. Adherent RPMI).

Single-sample Gene Set Enrichment Analysis (ssGSEA) across all metabolic pathways in the

KEGG database was run on the normalized gene expression values, giving normalized

enrichment scores (NES) representing relative metabolic pathway activity for 69 metabolic

pathways for each cell line. Normalized Weighted Average Expression (NWAS) was calculated

as in [37]. Briefly, individual cell line gene expression values for each gene within a gene set are

weighted by the corresponding number of pathways with which they overlap. Then, normalized

weighted values are summed to achieve a pathway activity score.

Genetic Pathway Dependency Enrichment Analysis

For each metabolic pathway, the NES was correlated with the -CERES score for all 16,643 genes.

Due to the bimodal nature of NESs, Spearman correlations were used. The resulting correlation

coefficients were ranked and GSEA querying KEGG metabolic pathways was run to calculate

Genetic Pathway Dependency Enrichment Analysis (Genetic PDEA). Positive NES represent

increased essentiality upon increased metabolic pathway activity, whereas negative NES

represent increased essentiality upon decreased metabolic pathway activity.

Drug Response Correlations

For each metabolic pathway, the NES was correlated with the -AUC (area under the curve) for

1,448 anti-cancer drugs in the PRISM repurposing database. Here, the AUC represents the dose

dependent effect of a drug on cell growth, calculated by fitting a four-parameter logistic curve to

viability values for each compound and cell line, with a lower AUC representing a stronger

response to drug [21]. Drugs with less than 150 cell lines were removed, leaving 1,390 drugs.

Spearman correlation p-values were calculated and a Benjamini-Hochberg false discovery rate

correction was applied for each metabolic pathway.

Pharmacological Pathway Dependency Enrichment Analysis

Drugs were mapped to their metabolic pathway using the annotated target(s) and genes from

KEGG metabolic pathways. Since the PRISM database contains both activators and inhibitors,

we annotated all activators by mechanism of action and multiplied their correlation coefficients by

-1. Therefore, a pathway activator would be counted similarly to a pathway inhibitor. Pathways

with 4 or more drugs were kept. Then, GSEA was run on the rank lists of 1,390 correlation

coefficients.

Integration of Individual Drug Response and Gene Dependency

Drug-gene dependency pairs were mapped using the target annotations for each drug.

Correlation coefficients for each drug and gene dependency were summed for each metabolic

pathway, generating 187,818 drug+gene:pathway combinations. An empirical permutation test

was run sampling 1,000 combinations of each drug+gene correlation coefficient. P-values were

calculated by dividing the number of permutations that out-performed the real summed correlation

coefficients by the number of same-signed permutations. P-values were then adjusted using a

Benjamini-Hochberg correction.

Integration of Genetic PDEA and Pharmacological PDEA

Results from Genetic PDEA and Pharmacological PDEA were filtered for same signed NES and

p-values of less than 0.05 and FDR values of less than 0.25 (per the original GSEA algorithm).

Three pathway-drug-gene dependencies were identified out of a possible 3,220 combinations.

Code Availability

635 Our code is freely available for use at: https://github.com/JamesJoly/MetabolicDependencies

Acknowledgments

638 This work was supported by the American Cancer Society Grant IRG-16-181-57, the 2020 AACR-

639 Bayer Innovation and Discovery Grant, Grant Number 20-80-44-GRAH, and the Viterbi School of

640 Engineering.

Conflict of Interest Statement

643 The authors declare no conflicts of interest.

Author Contributions

JHJ conceived the project. JHJ and BTLC performed research. JHJ and NAG analyzed data. JHJ

and NAG wrote the manuscript.

References

1.   Warburg O. On the Origin of Cancer Cells. Science. 1956;123: 309–314.
doi:10.1126/science.123.3191.309

2.   Luengo A, Gui DY, Vander Heiden MG. Targeting Metabolism for Cancer Therapy. Cell
Chem Biol. 2017;24: 1161–1180. doi:10.1016/j.chembiol.2017.08.028

3.   Dang L, White DW, Gross S, Bennett BD, Bittinger MA, Driggers EM, et al. Cancer-
associated IDH1 mutations produce 2-hydroxyglutarate. Nature. 2009;462: 739–744.
doi:10.1038/nature08617

4.   Graham NA, Minasyan A, Lomova A, Cass A, Balanis NG, Friedman M, et al. Recurrent
patterns of DNA copy number alterations in tumors reflect metabolic selection pressures.
Molecular Systems Biology. 2017;13: 914. doi:10.15252/msb.20167159

5.   Muller FL, Colla S, Aquilanti E, Manzo VE, Genovese G, Lee J, et al. Passenger deletions
generate therapeutic vulnerabilities in cancer. Nature. 2012;488: 337–342.
doi:10.1038/nature11331

6.   Sharma AK, Eils R, König R. Copy Number Alterations in Enzyme-Coding and Cancer-
Causing Genes Reprogram Tumor Metabolism. Cancer Res. 2016;76: 4058–4067.
doi:10.1158/0008-5472.CAN-15-2350

7.   Ward PS, Patel J, Wise DR, Abdel-Wahab O, Bennett BD, Coller HA, et al. The Common
Feature of Leukemia-Associated IDH1 and IDH2 Mutations Is a Neomorphic Enzyme
Activity Converting α-Ketoglutarate to 2-Hydroxyglutarate. Cancer Cell. 2010;17: 225–234.
doi:10.1016/j.ccr.2010.01.020

8.   Rohle D, Popovici-Muller J, Palaskas N, Turcan S, Grommes C, Campos C, et al. An
Inhibitor of Mutant IDH1 Delays Growth and Promotes Differentiation of Glioma Cells.
Science. 2013;340: 626–630. doi:10.1126/science.1236062

9.   Yen K, Travins J, Wang F, David MD, Artin E, Straley K, et al. AG-221, a First-in-Class
Therapy Targeting Acute Myeloid Leukemia Harboring Oncogenic IDH2 Mutations. Cancer
Discov. 2017;7: 478–493. doi:10.1158/2159-8290.CD-16-1034

10. Zhang T, Bauer C, Newman AC, Uribe AH, Athineos D, Blyth K, et al. Polyamine pathway
activity promotes cysteine essentiality in cancer cells. Nature Metabolism. 2020; 1–15.
doi:10.1038/s42255-020-0253-2

11. Maddocks ODK, Berkers CR, Mason SM, Zheng L, Blyth K, Gottlieb E, et al. Serine
starvation induces stress and p53-dependent metabolic remodelling in cancer cells.
Nature. 2013;493: 542–546. doi:10.1038/nature11743

12. Holleman A, Cheok MH, den Boer ML, Yang W, Veerman AJP, Kazemier KM, et al. Gene-
Expression Patterns in Drug-Resistant Acute Lymphoblastic Leukemia Cells and
Response to Treatment. New England Journal of Medicine. 2004;351: 533–542.
doi:10.1056/NEJMoa033513

13.   Krall AS, Mullen PJ, Surjono F, Momcilovic M, Schmid EW, Halbrook CJ, et al. Asparagine
couples mitochondrial respiration to ATF4 activity and tumor growth. Cell Metab. 2021.
doi:10.1016/j.cmet.2021.02.001

14.   Gao X, Sanderson SM, Dai Z, Reid MA, Cooper DE, Lu M, et al. Dietary methionine
influences therapy in mouse cancer models and alters human metabolism. Nature.
2019;572: 397–401. doi:10.1038/s41586-019-1437-3

15.   Lien EC, Lyssiotis CA, Juvekar A, Hu H, Asara JM, Cantley LC, et al. Glutathione
biosynthesis is a metabolic vulnerability in PI(3)K/Akt-driven breast cancer. Nat Cell Biol.
2016;18: 572–578. doi:10.1038/ncb3341

16.   Joly JH, Delfarah A, Phung PS, Parrish S, Graham NA. A synthetic lethal drug combination
mimics glucose deprivation–induced cancer cell death in the presence of glucose. J Biol
Chem. 2020;295: 1350–1365. doi:10.1074/jbc.RA119.011471

17.   Liu X, Olszewski K, Zhang Y, Lim EW, Shi J, Zhang X, et al. Cystine transporter regulation
of pentose phosphate pathway dependency and disulfide stress exposes a targetable
metabolic vulnerability in cancer. Nature Cell Biology. 2020;22: 476–486.
doi:10.1038/s41556-020-0496-x

18.   Fan J, Ye J, Kamphorst JJ, Shlomi T, Thompson CB, Rabinowitz JD. Quantitative flux
analysis reveals folate-dependent NADPH production. Nature. 2014;510: 298–302.
doi:10.1038/nature13236

19.   Meyers RM, Bryan JG, McFarland JM, Weir BA, Sizemore AE, Xu H, et al. Computational
correction of copy number effect improves specificity of CRISPR–Cas9 essentiality
screens in cancer cells. Nature Genetics. 2017;49: 1779–1784. doi:10.1038/ng.3984

20.   Tsherniak A, Vazquez F, Montgomery PG, Weir BA, Kryukov G, Cowley GS, et al. Defining
a Cancer Dependency Map. Cell. 2017;170: 564-576.e16. doi:10.1016/j.cell.2017.06.010

21.   Corsello SM, Nagari RT, Spangler RD, Rossen J, Kocak M, Bryan JG, et al. Discovering
the anticancer potential of non-oncology drugs by systematic viability profiling. Nat Cancer.
2020; 1–14. doi:10.1038/s43018-019-0018-6

22.   Ghandi M, Huang FW, Jané-Valbuena J, Kryukov GV, Lo CC, McDonald ER, et al. Next-
generation characterization of the Cancer Cell Line Encyclopedia. Nature. 2019;569: 503–
508. doi:10.1038/s41586-019-1186-3

23.   Lagziel S, Lee WD, Shlomi T. Inferring cancer dependencies on metabolic genes from
large-scale genetic screens. BMC Biology. 2019;17: 37. doi:10.1186/s12915-019-0654-4

24.   Gonçalves E, Segura-Cabrera A, Pacini C, Picco G, Behan FM, Jaaks P, et al. Drug
mechanism-of-action discovery through the integration of pharmacological and CRISPR
screens. Molecular Systems Biology. 2020;16: e9405. doi:10.15252/msb.20199405

25.   Lin A, Giuliano CJ, Palladino A, John KM, Abramowicz C, Yuan ML, et al. Off-target toxicity
is a common mechanism of action of cancer drugs undergoing clinical trials. Science
Translational Medicine. 2019;11. doi:10.1126/scitranslmed.aaw8412

26.   Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The
724   Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity.
725   Nature. 2012;483: 603–607. doi:10.1038/nature11003

726   27.   Garnett MJ, Edelman EJ, Heidorn SJ, Greenman CD, Dastur A, Lau KW, et al. Systematic
727   identification of genomic markers of drug sensitivity in cancer cells. Nature. 2012;483:
728   570–575. doi:10.1038/nature11005

729   28.   Deans RM, Morgens DW, Ökesli A, Pillay S, Horlbeck MA, Kampmann M, et al. Parallel
730   shRNA and CRISPR-Cas9 screens enable antiviral drug target identification. Nature
731   Chemical Biology. 2016;12: 361–366. doi:10.1038/nchembio.2050

732   29.   Subramanian A, Narayan R, Corsello SM, Peck DD, Natoli TE, Lu X, et al. A Next
733   Generation Connectivity Map: L1000 Platform and the First 1,000,000 Profiles. Cell.
734   2017;171: 1437-1452.e17. doi:10.1016/j.cell.2017.10.049

735   30.   Wang T, Yu H, Hughes NW, Liu B, Kendirli A, Klein K, et al. Gene Essentiality Profiling
736   Reveals Gene Networks and Synthetic Lethal Interactions with Oncogenic Ras. Cell.
737   2017;168: 890-903.e15. doi:10.1016/j.cell.2017.01.013

738   31.   Barbie DA, Tamayo P, Boehm JS, Kim SY, Moody SE, Dunn IF, et al. Systematic RNA
739   interference reveals that oncogenic KRAS-driven cancers require TBK1. Nature. 2009;462:
740   108–112. doi:10.1038/nature08460

741   32.   Kanehisa M, Goto S, Sato Y, Kawashima M, Furumichi M, Tanabe M. Data, information,
742   knowledge and principle: back to metabolism in KEGG. Nucl Acids Res. 2014;42: D199–
743   D205. doi:10.1093/nar/gkt1076

744   33.   Li H, Ning S, Ghandi M, Kryukov GV, Gopal S, Deik A, et al. The landscape of cancer cell
745   line metabolism. Nature Medicine. 2019;25: 850–860. doi:10.1038/s41591-019-0404-8

746   34.   May JL, Kouri FM, Hurley LA, Liu J, Tommasini-Ghelfi S, Ji Y, et al. IDH3α regulates one-
747   carbon metabolism in glioblastoma. Science Advances. 2019;5: eaat0456.
748   doi:10.1126/sciadv.aat0456

749   35.   Behan FM, Iorio F, Picco G, Gonçalves E, Beaver CM, Migliardi G, et al. Prioritization of
750   cancer therapeutic targets using CRISPR–Cas9 screens. Nature. 2019;568: 511–516.
751   doi:10.1038/s41586-019-1103-9

752   36.   Wilson DJ. The harmonic mean p-value for combining dependent tests. Proc Natl Acad Sci
753   U S A. 2019;116: 1195–1200. doi:10.1073/pnas.1814092116

754   37.   Xiao Z, Dai Z, Locasale JW. Metabolic landscape of the tumor microenvironment at single
755   cell resolution. Nat Commun. 2019;10: 1–12. doi:10.1038/s41467-019-11738-0

756   38.   Birsoy K, Wang T, Chen WW, Freinkman E, Abu-Remaileh M, Sabatini DM. An Essential
757   Role of the Mitochondrial Electron Transport Chain in Cell Proliferation Is to Enable
758   Aspartate Synthesis. Cell. 2015;162: 540–551. doi:10.1016/j.cell.2015.07.016

39. Sullivan LB, Gui DY, Hosios AM, Bush LN, Freinkman E, Vander Heiden MG. Supporting
Aspartate Biosynthesis Is an Essential Function of Respiration in Proliferating Cells. Cell.
2015;162: 552–563. doi:10.1016/j.cell.2015.07.017

40. Poliaková M, Aebersold DM, Zimmer Y, Medová M. The relevance of tyrosine kinase
inhibitors for global metabolic pathways in cancer. Mol Cancer. 2018;17.
doi:10.1186/s12943-018-0798-9

41. Burg JS, Espenshade PJ. Regulation of HMG-CoA reductase in mammals and yeast. Prog
Lipid Res. 2011;50: 403–410. doi:10.1016/j.plipres.2011.07.002

42. Bhat CS, Ramasarma T. Effect of Phenyl and Phenolic Acids on Mevalonate-5-Phosphate
Kinase and Mevalonate-5-Pyrophosphate Decarboxylase of the Rat Brain. Journal of
Neurochemistry. 1979;32: 1531–1537. doi:10.1111/j.1471-4159.1979.tb11095.x

43. Castillo M, Martínez-Cayuela M, Zafra MF, García-Peregrín E. Effect of phenylalanine
derivatives on the main regulatory enzymes of hepatic cholesterogenesis. Molecular and
Cellular Biochemistry. 1991. doi:10.1007/BF00230371

44. Shefer S, Tint GS, Jean-Guillaume D, Daikhin E, Kendler A, Nguyen LB, et al. Is there a
relationship between 3-hydroxy-3-methylglutaryl coenzyme a reductase activity and
forebrain pathology in the PKU mouse? J Neurosci Res. 2000;61: 549–563.
doi:10.1002/1097-4547(20000901)61:5<549::AID-JNR10>3.0.CO;2-0

45. Zhang F, Snead CM, Catravas JD. Hsp90 regulates O-linked β-N-acetylglucosamine
transferase: a novel mechanism of modulation of protein O-linked β-N-acetylglucosamine
modification in endothelial cells. Am J Physiol Cell Physiol. 2012;302: C1786–C1796.
doi:10.1152/ajpcell.00004.2012

46. Koay DC, Zerillo C, Narayan M, Harris LN, DiGiovanna MP. Anti-tumor effects of retinoids
combined with trastuzumab or tamoxifen in breast cancer cells: induction of apoptosis by
retinoid/trastuzumab combinations. Breast Cancer Research. 2010;12: R62.
doi:10.1186/bcr2625

47. Fruman DA, Chiu H, Hopkins BD, Bagrodia S, Cantley LC, Abraham RT. The PI3K
Pathway in Human Disease. Cell. 2017;170: 605–635. doi:10.1016/j.cell.2017.07.029

48. Parmenter TJ, Kleinschmidt M, Kinross KM, Bond ST, Li J, Kaadige MR, et al. Response
of BRAF-Mutant Melanoma to BRAF Inhibition Is Mediated by a Network of Transcriptional
Regulators of Glycolysis. Cancer Discov. 2014;4: 423–433. doi:10.1158/2159-8290.CD-13-
0440

49. Locasale JW. Serine, glycine and one-carbon units: cancer metabolism in full circle. Nature
Reviews Cancer. 2013;13: 572–583. doi:10.1038/nrc3557

50. Han K, Pierce SE, Li A, Spees K, Anderson GR, Seoane JA, et al. CRISPR screens in
cancer spheroids identify 3D growth-specific vulnerabilities. Nature. 2020; 1–6.
doi:10.1038/s41586-020-2099-x

51. Cantor JR, Abu-Remaileh M, Kanarek N, Freinkman E, Gao X, Louissaint A, et al.
Physiologic Medium Rewires Cellular Metabolism and Reveals Uric Acid as an

Endogenous Inhibitor of UMP Synthase. Cell. 2017;169: 258-272.e17.
doi:10.1016/j.cell.2017.03.023

52. Liu Y, Beyer A, Aebersold R. On the Dependency of Cellular Protein Levels on mRNA
Abundance. Cell. 2016;165: 535–550. doi:10.1016/j.cell.2016.03.014

53. Brunk E, Chang RL, Xia J, Hefzi H, Yurkovich JT, Kim D, et al. Characterizing
posttranslational modifications in prokaryotic metabolism using a multiscale workflow.
PNAS. 2018;115: 11096–11101. doi:10.1073/pnas.1811971115

54. Oliveira AP, Ludwig C, Picotti P, Kogadeeva M, Aebersold R, Sauer U. Regulation of yeast
central metabolism by enzyme phosphorylation. Mol Syst Biol. 2012;8: 623.
doi:10.1038/msb.2012.55

55. Chen P-H, Cai L, Huffman K, Yang C, Kim J, Faubert B, et al. Metabolic Diversity in
Human Non-Small Cell Lung Cancer Cells. Mol Cell. 2019;76: 838-851.e5.
doi:10.1016/j.molcel.2019.08.028