Environmental stressors induced strong small-scale phenotypic differentiation in a wide-dispersing marine snail

Nicolás Bonel[1,2,3*], Jean-Pierre Pointier[4] & Pilar Alda[1,2]

[1] Centro de Recursos Naturales Renovables de la Zona Semiárida (CERZOS—CCT—CONICET Bahía Blanca), Camino de la Carrindanga km 7, Bahía Blanca 8000, Argentina.

[2] Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

[3] Centre d'Écologie Fonctionnelle et Évolutive, UMR 5175, CNRS—Université de Montpellier, Université Paul-Valéry Montpellier—École Pratique des Hautes Études—IRD, 34293 Montpellier Cedex 05, France.

[4] PSL Research University, USR 3278 CNRS–EPHE, CRIOBE Université de Perpignan, Perpignan, France.

* To whom correspondence should be addressed: nicobonel@gmail.com (N. Bonel)

Running head: Small-scale phenotypic differentiation in a wide-dispersing snail

Key words: adaptive plasticity, shell characters, genital morphology, intertidal zonation, contrasting selection pressures, planktotrophic snail, high dispersal potential.

ABSTRACT

Heterogeneous environments pose a particular challenge for organisms because the same phenotype is unlikely to perform best regardless of the types of stress it encounters. To understand how species meet this challenge, we investigated the extent to which contrasting environmental pressures induced ecological and phenotypic responses in a natural population of a wide-dispersing marine snail at a small spatial scale. We analyzed several traits of *Heleobia australis* (Rissooidea: Cochliopidae) collected from heterogeneous, but highly connected, habitats from the intertidal area of the Bahía Blanca estuary, Argentina. We also conducted molecular analyses by amplifying the COI gene in individuals sampled from each habitat. We found that subpopulations living in sympatry of *H. australis* exhibited a strong phenotypic divergence in shell characters and body weight in response to thermal, saline, and dehydration stress, crab predation risk, and parasitic castrators. We proved that this differentiation occurred even early in life as most of the characters observed in juveniles mirrored those found in adults. We also found a divergence in penis size in snails collected from each habitat and raised in common garden laboratory conditions. The molecular analyses confirmed that the individuals studied constituted a single species despite the strong phenotypic differences among subpopulations. The small-scale phenotypic differentiation suggests that *H*. *australis* experienced a fine-grained environment where conditions imposed by different sources of stress favored the expression of beneficial traits. We discuss the role of plasticity in shaping adaptive phenotypic responses that increase the likelihood of persistence of subpopulations facing environmental stressful conditions.

1. INTRODUCTION

If organisms have difficulties in adapting to human-induced global change and fail to track projected environmental changes, populations become vulnerable to decline and extinction (Hill et al. 2011, Hoffmann & Sgrò 2011). Phenotypic plasticity, a common feature in nature across many taxa, is the ability of a single genotype to express different phenotypes in dissimilar biotic or abiotic environments (Travis 1994, West-Eberhard 2003). Many studies indicate a role of adaptive plasticity in shaping phenotypic responses as an effective mechanism that can greatly improve the conditions for persistence of populations facing abrupt environmental shift by facilitating rapid adaptation to the new selection pressures (Price et al. 2003, Ghalambor et al. 2007, Chevin & Lande 2009). Adaptive plasticity that places populations close enough to the new favored phenotypic optimum, for directional selection to act, provides the first step in adaptation to new environments on ecological time scales as long as costs to maintaining and/or producing a plastic response are not substantially large, thus constraining the evolution of plasticity (Price et al. 2003, Ghalambor et al. 2007, Chevin & Lande 2009). In this sense, adaptive phenotypic plasticity is crucial for the long-term persistence of populations facing new environmental stress brought about by human-induced global change (Hill et al. 2011).

Temperate marine habitats, in particular the intertidal zone, exhibit great variability in environmental factors mainly driven by the fall and rise of the tides, creating areas on the shore that are alternately immersed and exposed (Denny & Paine 1998, Bourdeau et al. 2015). The transitional nature of the intertidal habitat from marine to terrestrial conditions strongly influences the physiology and the ecology of intertidal organisms due to, for instance, the increase in environmental harshness (e.g. desiccation, extremes in temperature and salinity) along the vertical zonation of the intertidal area (Vermeij 1972, Denny & Paine 1998, Mouritsen et al. 2018). From this perspective, intertidal organisms, such as aquatic gastropods, are likely to exhibit increased phenotypic plasticity in response to such environmental selective pressures (Berrigan & Scheiner 2004).

Phenotypic variation in shell traits is particularly widespread in gastropods (Bourdeau et al. 2015) and their shells offer an easily measured and permanent record of how the

organism responded to local abiotic (e.g. water chemistry, temperature) and biotic (e.g.

predation, parasitism) agents (Dillon et al. 2013). Major environmental factors as

prolonged submersion times, desiccation, high temperature, and extreme salinity across

the vertical zonation create contrasting pressures that give rise to opposite shell trait

responses (Struhsaker 1968, Johannesson & Johannesson 1996). For instance,

prolonged submersion times enhance foraging times and the absorption of calcium

carbonate from water leading to higher shell growth rates, which result in larger and

narrower shells and increased body mass (Vermeij 1973, review by Chapman 1995). By

contrast, snails from upper intertidal habitats that are periodically exposed to hot and

dry conditions and have only a limited time each day to feed exhibit the opposite shell

and body mass responses and are characterized by having a smaller aperture size, which

has been correlated to an increase in resistance to high desiccation in upper intertidal

habitats (Vermeij 1973, Chapman 1995, Melatunan et al. 2013).

Predators are one component of the heterogeneity across the intertidal zone because

they are often distributed patchily in time and space creating a steep selection gradient

in predation risk, which, in turns, can promote adaptive shell responses (Appleton &

Palmer 1988, Palmer 1990, Boulding & Hay 1993, Hollander et al. 2006, Bourdeau

2011). The adaptive value of predator-induced plasticity primarily linked to shell

defenses are not limited to shell thickening (more resistant to crushing) as snails are

capable of plastically altering different aspects of their shell shape (i.e. increased

globosity; lower shell length to width ratio) or aperture (i.e. more elongated; higher

aperture length to width ratio), which prevents the crab from pulling the soft parts out of

the shell (Johannesson 2003, Dillon et al. 2013, Bourdeau et al. 2015). However, not all

phenotypic plasticity triggered by environmental conditions is adaptive (Ghalambor et

al. 2007). Some plastic trait responses, usually those imposed by the biochemistry and

physiology of the organism, can be reversed over short time scales, which is not the

case for developmental plasticity as it tends to be irreversible or takes longer to be

reversed (Pigliucci et al. 2006). Parasites, for instance, can induce morphological,

behavioral, and physiological change of individual snail hosts thereby influencing

several aspects of host life history that can significantly alter size structure,

demography, resource use, and intra and interspecific interactions of host population

(Fredensborg et al. 2005, Miura et al. 2006, Mouritsen et al. 2018). They have been also

reported to induce changes in microhabitat choice (Curtis 1987) and in body size in snails (Mouritsen & Jensen 1994, Probst & Kube 1999, Levri et al. 2005, Miura et al. 2006, Alda et al. 2010). Such pressure exerted by parasites also differs along the intertidal zone (Smith 2001, Alda et al. 2010, 2019) and can cause important shifts in the expression of host phenotypic traits, creating pronounced phenotypic differences between infected and uninfected hosts (Poulin & Thomas 1999, Fredensborg et al. 2006).

Despite the intertidal area spans a strong selective gradient, small-scale phenotypic differentiation and adaptation of wide-dispersing organisms has been understudied (review by Sanford & Kelly 2011). These contrasting conditions along the vertical distribution pose a particular challenge for wide-dispersing intertidal organisms: the same phenotype is unlikely to perform best regardless of the types of stress it encounters. We can then have as a result (i) a generalist phenotype (probably with maladaptation), (ii) a phenotypic plasticity promoting different phenotypes under different conditions, or (iii) a local genetic differentiation, contributing to the phenotypic variation between habitats, favored by restricted gene flow through habitat selection (and probably partial genetic isolation leading to ecotype formation). For instance, limited gene flow can occur when habitat conditions disrupt larval dispersal thereby increasing localized recruitment (Struhsaker 1968, Johnson et al. 1994, Parsons 1997, Levin et al. 2006), namely 'phenotype-environment mismatch' (reviewed in Marshall et al. 2010), whereat individuals would be subjected to local biotic and abiotic stressors (Johannesson & Johannesson 1996, Westram et al. 2021). Hence, to understand how species meet this challenge, the first step is to observe phenotypic variation and its association in the field with microhabitats characterized by different sources of stress. If we observe different phenotypes associated with different conditions, the next step is to explore whether such phenotypic variation results from a direct impact of stress (e.g. trait shift induced by parasites) or is it because we are dealing with ecotypes that find alternative niches (e.g. different microhabitats) preventing, for instance, reproduction.

Here we tested how contrasting biotic and abiotic conditions across the intertidal gradient affected snail density and induced different phenotypic responses at a small

spatial scale among individuals from different subpopulations living in sympatry of a wide-dispersing intertidal snail. We expected that snail density and most of mean values of phenotypic traits measured shifted to lower values in habitats with high physical stress conditions (thermal, saline, and dehydration stress), crab predation risk, and parasite pressure. To test this, we collected and measured shell traits, weighted, and dissected the mud snail *Heleobia australis* (d'Orbigny, 1835) (Rissooidea: Cochliopidae) over the four seasons throughout a year from three distinct, but highly connected, habitats from the intertidal area of the Bahía Blanca estuary, Southwestern Atlantic, Argentina. Then, we collected juveniles from each habitat and raised them in common garden laboratory conditions until they were adults to investigate if genital morphology varied. Finally, we explored whether individuals living in these contrasting habitats belong to the same species, and not to a species complex, by amplifying the cytochrome oxidase subunit 1 (COI) gene. Altogether, we found clear evidence of small-scale phenotypic differentiation in all traits likely as a result of environmental pressures imposed by a strong biotic and abiotic stressors across the vertical distribution of the intertidal area.

2. MATERIAL AND METHODS

2.1. Study species

The intertidal mud snail *Heleobia australis* (previously known as *Littoridina australis*) has a wide geographic range inhabiting marine and estuarine ecosystems from tropical to temperate regions (from Brazil: 22° 54' S, to Argentina: 40° 84' S; De Francesco & Isla 2003) and is the most common benthic macrofaunal species in the estuaries and coastal lagoons (De Francesco & Isla 2004). In the Bahía Blanca estuary, there are no other competing organism to *H*. *australis* as this snail species is the dominant gastropod (Elías et al. 2004). It is a gonochoristic species with internal fertilization (Neves et al. 2010). Mature females lay egg capsules, containing one fertilized egg (Neves et al. 2010), preferably on shells of conspecific individuals, but they may also be laid on shells of other species, on sand grains, or algae, which later develop into a veliger larva that exhibits a short pelagic larval life (Neves et al. 2010), which is, on average, 10±3 days in standard laboratory conditions (12:12 photoperiod, 25 ºC, water salinity 30 PSU, and *ad libitum* food in the form of boiled ground lettuce; Bonel, *unpublished data*). Adult snails attain a maximum shell size ranging from 7 to 8 mm

with an average shell length of 5.2 mm (0.5 SD; this study) and a lifespan of 2.9 years

in a temperate climate and two well-defined recruitment episodes (De Francesco &

Isla 2004, Carcedo & Fiori 2012). However, in tropical and subtropical areas breeding

and recruitment occur year-round (Neves et al. 2010) suggesting that it is likely that *H.*

australis have shorter generation times. This species is capable of creating an air

bubble inside its pallial cavity, allowing juveniles and adults to temporarily float in the

water column, whereat they are carried along by the tide and wind driven current

(Echeverría et al. 2010), increasing its dispersion potential. Moreover, *H. australis* is

an obligate intermediate host of a large diversity of parasites that infect several other

species hosts. For instance, *H. australis* has been reported to host 18 larval trematode

species and 16 of these were from the Bahía Blanca estuary (Argentina), being all but

one parasitic castrators, with an individual worm eventually filling up the digestive

gland and gonad of the snail host (Alda & Martorelli 2014).

2.2. Study area and (a)biotically heterogeneous habitats

This study was conducted in the Villa del Mar saltmarsh-mudflat located in the middle

reaches of the Bahía Blanca estuary, Argentina (38° 51' S – 62° 07' W). The intertidal

mudflat extends for more than 1 km across the tidal gradient. Its topography is

characterized by a very gently sloping ramp on the seaward side (Pratolongo et al. 2009)

and affected by strong semidiurnal tides and high seasonal variation (Perillo et al.

2001). Close to the mean high tide level, marshes covered by cordgrass (*Spartina*

alterniflora) form a narrow, 150 m wide strip of vegetation followed by a mudflat area

with no vegetation (Pratolongo et al. 2010). During low tide, these upper areas of the

intertidal zone are subjected to high desiccation and temperature and salinity

fluctuations as the tide takes *ca*. 10 hours to cover them (P. Pratolongo, *personal*

communication; 26 February 2019) whereas low areas located close the seaward edge

remain covered by water during low tide.

The existence of a tidal cycle exposes *H. australis* to extreme, but predictable, changes

in abiotic conditions (at least twice every day) likely determining their vertical

distribution (zonation patterns). Snails in the upper intertidal zone must remain

quiescent for long periods of time during daylight hours or during low tide, being

exposed to aerial conditions for longer periods than individuals in the lower intertidal

zone, which remain cover by water during low tide. The intertidal zone can therefore be
characterized in three distinct habitats: flats, marshes, and pans. Flats and marshes are
located in the upper zone and they drain at low tide, though flats are free of vegetation
and marshes are covered by cordgrass. Pans are free of vegetation but remain covered
by water during low tide and are located close the seaward edge. Thermal, saline, and
dehydration stress are strong selective forces occurring mainly in the upper intertidal
area (flats and marshes; Denny & Paine 1998, Bourdeau et al. 2015) whereas these
inducing agents are weaker in pans, which exhibit low environmental stress condition
(Fig. 1).

Biotic agents also vary along the vertical distribution of the intertidal zone (Fig. 1). On
the one hand, the grapsid crab *Neohelice granulata* is one of the most abundant
macroinvertebrates of intertidal areas of the SW Atlantic estuaries where it commonly
inhabits the upper vegetated area (marshes) but uses the entire intertidal zone (Spivak et
al. 1994, Alvarez et al. 2013, Angeletti et al. 2014, Angeletti & Cervellini 2015). This
burrowing crab could exert a strong pressure on *Heleobia australis* since it can
drastically reduce snail density in vegetated areas due to an intense bioturbation activity
(Spivak et al. 1994, Alvarez et al. 2013, Angeletti et al. 2014, Angeletti & Cervellini
2015) and/or through snail predation (D'Incao et al. 1990, Barutot et al. 2011).
Although the main predator in the intertidal mudflat are crabs, there are other potential
predators such as snail-eating birds (e.g. *Charadrius* spp.) that occasionally feed on *H.
australis* (Martorelli 1991, Ieno et al. 2004, Meerhoff et al. 2013). For instance, in the
study area, *H. australis* has been reported to be one of the 39 dietary items consumed by
the White-Backed Stilt (Petracci & Delhey 2005). On the other hand, parasite pressure
also differs along the intertidal zone. Trematode infection not only inevitably leads to
snail castration reducing its fitness to zero (Alda et al. 2019) but also gives rise to
smaller shell-sized morphs for infected snails (Alda et al. 2010). The prevalence
(percentage of individuals infected) of trematodes is higher in the lower area of the
intertidal zone, where parasite infection is predominately caused by one extremely
prevalent trematode, *Microphallus simillimus*. Such pressure is stronger in pans than in
flats and marshes because prolonged submersion times allow snails to increase time
spent foraging and thus ingesting parasite eggs (Alda et al. 2019).

2.3. Field sampling and laboratory procedure

We sampled individuals of the intertidal mud snail *H. australis* in summer (March 6), autumn (July 15), winter (September 20), and spring (December 7) of 2012 from the study area. We performed an intense sampling in three contrasting intertidal habitats from the Villa del Mar saltmarsh-mudflat because our aim was to test for phenotypic variation in *Heleobia australis* at a small-spatial scale (< 1 km) and because marine environmental conditions in this location are representative of conditions found at other sites from the Bahía Blanca estuary (Piccolo & Hoffmeyer 2004). We followed a simple random sampling by area for polygon study areas. We used QGIS 1.7 (https://qgis.org) to define the sampling zone and to randomly distribute the points where the circular quadrats were placed. Specifically, during low tide and from each biological zone from the intertidal area (flats, marshes, and pans), we sampled snails by laying down nine circular quadrats with 10 cm diameter and 2 cm deep (Area $= 78.5$ cm^2) to get a representative sample of each zone (e.g. Milroy 2015). Snails were sieved and washed from the sediment through a 1-mm mesh, and immediately transported alive to the laboratory in 0.5-l plastic bottles filled with *in situ* water where they were kept in aquaria and fed *ad libitum* with flake fish food. We repeated this procedure for each sampling date.

Once in the lab, we counted the total number of snails in each of the nine samples (i.e. samples A to I) taken from each habitat (flats, marshes, or pans) and sampling date. From each sample, we arbitrarily decided to subsample 60% of individuals from each sample to check for parasite infection, 30% for shell and body weight, and 10% for parasite biomass. Because *Heleobia australis* is rather a small snail (mean shell length = 5-6 mm), it is hardly impossible to determine with the naked eye whether individuals are phenotypically different from one another. Thus, as the probability of choosing each individual was equal, we assumed that subsampling was done at random. Overall, our large dataset consisted of 6,250 individuals, which were used to account for infected and uninfected snails (hereafter infection status), and the remaining uncrushed snails were used to estimate snail shell and body weight (n = 3,057; see below) and parasite biomass (n = 1,060; data not showed in this study). All 10,367 snails were used to estimate snail density and were also photographed using a camera attached to a dissecting microscope to analyze shell and aperture morphometrics (see below).

2.4. Snail density and infection status

To test how contrasting habitat conditions affected snail density, we counted all sampled individuals per habitat and sampling date to analyze spatial and temporal variation of snail density. Then, to determine infection status, snails from each habitat were crushed using a mortar and a pestle, tissue was examined under a dissecting microscope, and trematodes were identified under a compound microscope following Alda & Martorelli (2014). Previous studies show that the most prevalent parasite in the study area is *Microphallus simillimus* (Microphallidae), which makes up 87% of the overall prevalence (Alda & Martorelli 2014, Alda et al. 2019). This parasite has life history traits such as abbreviated life cycle (the same snail host can serve as the first- and second-intermediate host), meaning that once snails ingest parasite eggs, the metacercariae encyst within the sporocyst in the infected snail, thereby maximizing transmission success (Alda & Martorelli 2014). One possible approach to the contribution of parasites to the phenotypic variance in host populations is to compare phenotypic responses of uninfected and infected hosts maintained under identical conditions (Poulin & Thomas 1999). Thus, to test for this, we only considered snails infected by the most prevalent trematode *M. simillimus* and also from pans (the habitat with the highest prevalence; Alda et al. 2019), which implies that both infected and uninfected snails grew in the same environmental conditions. By doing so, we can ensure that differences in phenotypic responses can only be attributed to *M. simillimus* effect and not caused by other trematodes or habitat stress conditions.

As older hosts have greater cumulative risk of infection than do young hosts, each sample should be standardized by age, collected at the same time from an area in which hosts are likely to mingle and where they have experienced relatively uniform risks of infection (Lafferty et al. 1994). We therefore identified age cohorts by means of the length-frequency distributions in each sampling date and habitat and removed those individuals outside the lower 95% confidence limit of the cohort with larger individuals (see Text S1 in the appendix for details; Fig. S1).

2.5. Shell and aperture morphometric

To analyze variation in shell morphometric across habitat conditions and infection status, we measured four linear variables from each photograph using ImageJ software:

shell length (SL), shell width (SW), aperture length (AL), and aperture width (AW).
We chose the linear morphometric measurements approach over the geometric
morphometrics analysis because the landmark-based approach has a methodological
limitation that is the small number of individuals (n < 600) than can be analyzed (e.g.
(Carvajal-Rodríguez et al. 2005, Conde-Padín et al. 2007, Guerra-Varela et al. 2009,
Dillon et al. 2013, Larsson et al. 2020) while the classical approach followed in our
study allowed us for measuring more than 10,000 snails.

Similar to some other caenogastropod species (e.g. *Littoridina* sp.), the shell of *H.
australis* is generally small and conical and the axis of coiling lying at an angle of about
45º above the plane of the elliptical aperture (Vermeij 1973, Hershler & Thompson
1992). We estimated the shell volume as proxy of shell size by calculating the volume
of a cone (or double cone) because is a more reliable measure to describe shell size than
only considering a linear measurement such as shell length:

$$\text{Shell size: Volume of a cone} = 1/3\ \pi\ (SW/2)^2\ SL$$

To analyze the shell shape, we estimated the SL to SW ratio. Likewise, we analyzed
aperture shape by calculating the AL to AW ratio. We used AL and AW to calculate
aperture size by calculating the area of an ellipse:

$$\text{Aperture size: Area of ellipse} = (AL/2)\ (AW/2)\ \pi$$

2.6. Shell thickness and body mass

To test whether snails subjected to a stronger predation pressure exhibit thicker shells,
(considered to be a shell defense trait against predators as it is more resistant to
crushing; (Johannesson 2003, Dillon et al. 2013, Bourdeau et al. 2015) and whether
contrasting environmental stress conditions affected body mass, we estimated and
compared shell weight and ash-free dry-weight among habitats. We used a subsample of
individuals (n = 70), representative of the population shell length-frequency
distributions obtained previously for all individuals sampled on each date and habitat.
We removed sediment and epibiota with a scouring pad before weighing and measuring
the organic content of the snail as ash-free dry weight (AFDW, calculated as the

difference between the dry weight and the weight of the incombustible component of

the shell), which was considered as body mass. To achieve this, we dried snails

individually in porcelain crucibles for 48 h at 60 °C, weighed them with a digital scale

(precision 0.1 mg), ashed the snails for 5 h in a muffle furnace at 500 °C and then

reweighed them (Bonel & Lorda 2015). The ash shell weight (the incombustible

component of the shell) was considered as proxy of shell thickness (e.g. Palmer 1981)

and used to estimate and compare its variation among individuals along the vertical

gradient of the intertidal area.

2.7. Variation in penis size

To test for differences in genital morphology, we analyzed variation in the penis size

and morphology of individuals from contrasting habitats conditions living in sympatry

but raised in common garden laboratory conditions. Genital morphology is one of the

most widely-used trait when studying the internal anatomy of gastropods, especially

among cochliopids (Liu et al. 2001). Reproductive anatomy is very useful to delineate

species since they tend to show more variation among than within species and penis

morphology, in particular, is used when delineating *Heleobia* species (Gaillard &

Castellanos 1976). We sampled snails of *H*. *australis* from the three distinct habitats of

the intertidal area of the Villa del Mar saltmarsh-mudflat in November (Spring) of 2017.

Snails were transported to the lab and they were split into adults and juveniles based on

their shell-length distribution (as mentioned above). Adults were preserved in alcohol

96% and used to conduct molecular analysis. Because field-collected snails may retain

the influence of environmental conditions experienced prior to collection even in early

life stages (Sanford & Kelly 2011), we tried to minimize this potential effect by

considering only the smallest individuals sampled from each habitat (mean shell length,

SD: 4.15 ± 0.36 mm). Juveniles were placed in three different aquaria per habitat to

prevent tank effects. The snail abundance at the onset of the experiment was similar in

each set of aquaria to minimize potential density-dependent effects (flats: n = $38.3 \pm$

0.6; marshes: n = 43.7 ± 0.6; pans: n = 46.7 ± 0.6). Juveniles were kept in common

garden laboratory conditions (12:12 photoperiod, 25 ℃, water salinity 30 PSU, and *ad*

libitum food in the form of boiled ground lettuce) for 21 months, which ensured that all

individuals considered in the analysis were adults. Prior to dissection, we collected, on

average, 30 snails from each aquaria and preserved them in the Railliet–Henry's

solution (930 ml distilled water, 6 g NaCl [0.85%], 50 ml formaldehyde [37%], 20 ml

glacial acetic acid). Then, they were all dissected under the dissecting microscope and

the reproductive system of *ca*. 15 males per aquarium was drawn using a camera lucida

attachment (Pointier et al. 2004). These were scanned and the penis surface area was

calculated using the ImageJ software. Shells were photographed and analyzed as

mentioned above to estimate shell size and shape.

2.8. COI analysis

To confirm that individuals living in these contrasting habitats belong to the same

species and not to a species complex, we conducted molecular analyses by amplifying

the cytochrome oxidase subunit 1 (COI) gene in 10 individuals sampled from each of

the three sites. Then, we built a gene tree and a haplotype network to verify if

individuals gather in clusters depending on the habitat they come from. We also applied

a species-delimitation method: Automatic Barcode Gap Detection (ABGD, Puillandre et

al. 2012). ABGD detects barcode gaps, which can be observed whenever the divergence

among organisms belonging to the same species is smaller than divergence among

organisms from different species (see Text S2 in the appendix for specific details on

these procedures).

2.9. Statistical analyses

To test for spatial and temporal variation of snail density, we performed a two-way

ANOVA including habitats (flats, marshes, and pans) and seasons (summer, autumn,

winter, spring) as fixed effect, and their interaction. We transformed density (natural-

log) data to meet assumptions of normality and homoscedasticity.

To test for phenotypic variation in shell characters, we defined three categories (cat.) of

snails that comprised: (*i*) uninfected adults, (*ii*) infected adults, and (*iii*) uninfected

juveniles. These three categories (cat. *i–iii*) were considered to test whether contrasting

(a)biotic stress conditions (environmental stress and crab predation) induced variation in

shell traits (morphometric and thickness) and body mass. To evaluate the effect exerted

by the trematode *Microphallus simillimus* on shell morphometry of *Heleobia australis*

from pans (the habitat with the highest prevalence), we created a fourth category that

included uninfected and infected adult snails (cat. *iv*). We performed four Principal

Components Analyses (PCAs) to visualize the main components of the morphological shell variation for each snail category. Then, we tested for differences in shell morphology (size and shape of the shell and aperture) by fitting independent linear mixed models (LMMs), with a Gaussian error distribution, for the above-mentioned four categories (cat. *i–iv*).

To test the effect of habitat conditions on shell traits (cat. *i–iii*), we considered habitat as a fixed effect. To test for morphometric differences between infected and uninfected (cat. *iv*), we included infection status (infected and uninfected) as a fixed effect as we only considered individuals from one habitat (i.e. same habitat conditions for infected and uninfected snails). For all the four categories (cat. *i–iv*), we included sex (male and female) as a fixed factor (and its interaction with habitat or infection status). We added sampling season as a random factor in all models as it can explain a significant portion of the variance in measured variables. Likewise, we also considered shell size as a covariate when testing for shell and aperture shape, but the models were all ran with and without this effect to check the extent to which the effects were mediated by shell size. We also incorporated shell size as a covariate when testing for shell and body weight (by means of LMMs) and estimates were statistically corrected for variation that could be explained by the covariate. Statistical significance of the fixed effects was obtained from model comparisons using likelihood-ratio tests. Random effects were separately tested using chi-square likelihood-ratio tests with the corrections indicated by Zuur (2009). To test for differences in penis size (area of the penis), we performed a linear model with habitat as a factor and we added shell size (volume) as a covariate, which was used to statistically correct estimates for variation explained by the covariate. All individuals analyzed were free of parasites, meaning that the differences found are not due to parasite effect.

Post-hoc tests were performed when the effects were significant ($P <0.05$), using Holm-Bonferroni correction for multiple testing to compare their effects. Values are given as means $\pm$ 1 SE unless otherwise stated. All analyses and figures were performed with R v.3.3.3 packages *lme4* (Bates et al. 2014), *nlme* (Pinheiro et al. 2017), *car* (Fox et al. 2011), *effects* (Fox 2003), *MASS* (Ripley et al. 2013), *plyr* (Wickham 2011), *dplyr* (Wickham et al. 2019a), *devtools* (Wickham et al. 2019b), *ggbiplot* (Vu 2011), *psych*

(Revelle 2018), *ggplot2* (Wickham & Chang 2008), and *outliers* (Komsta 2011).

3. RESULTS

3.1. Overview

Overall, the mud snail *Heleobia australis* showed a strong variation in snail density and a remarkable variation in several traits linked to contrasting biotic and abiotic pressures (environmental stress, predation, or parasite infection) occurring along the vertical zonation of the intertidal area. Tables 1 to 3 summarize descriptive statistics on density and shell traits measured for the first and fourth snail categories (cat. *i* and *iv*). In the appendix, we reported results of the snail density split by age classes (juvenile and adult; Table S1), principal components analyses (Text S3, Table S2, Fig. S3), descriptive statistics and the linear mixed models for the four snail categories on shell traits (Tables S3–S8) and for shell weight and body mass (Table S9).

3.2. Strong habitat and seasonal effect on snail density

We found spatial differences in total density (including juveniles and adult snails; Table 1; Fig. 2). We observed no significant interaction between habitats and seasons ($F_{(6, 106)}$ = 1.56, $P = 0.167$). Marshes showed the lowest density whereas flats and pans showed a marginal non-significant difference (Table 1). We found significant differences between seasons (Table 1). Density was higher in spring than in autumn, winter, and summer (Table 1). Snail density in autumn did not differ from that of summer or winter, these latter two being not different either (Table 1), but it largely increased from autumn to spring (particularly in pans as compared to flats and marshes; Fig. 2). This increase across seasons was mainly driven by juvenile recruitment (Table S1; Fig. S2). By contrast, the effect of habitat conditions and seasons on adult density showed a significant interaction because density was higher in pans from autumn to winter whereas it decreased in marshes for the same period, and in flats it decreased from summer to autumn and the than remained fairly constant (Table S1; Fig. S2).

3.3. Strong habitat effect on shell morphometrics

The morphometric responses to habitat conditions were similar in all the three snail categories analyzed (cat. *i–iii*). For simplicity, we present results and figures only for

uninfected adult snails (cat. *i*) in the main text and those for infected adults (cat. *ii*) and juvenile snails (cat. *iii*) are shown in the appendix (Tables S4-S7; Figs. S4-S5).

3.3.1. Shell size. We found a significant interaction between fixed effects (habitat and sex; $X^2_2 = 8.42$, $P = 0.015$). This was mainly driven by differences in this trait between sexes in pans, whereas such difference between males and females was not detected in snails from flats and marshes. In other words, males exhibited a larger shell size (mm^3) compared to females in pans, meaning that the sex effect was much stronger in creating size differences between sexes compared to flats and marshes where the sex effect was weaker resulting in no difference in size between sexes. By contrast, the effect of habitat conditions was stronger in both sexes. Both male and female snails from flats and marshes showed a smaller shell size (a decrease of 12 and 22 % in shell size; respectively) relative to individuals from pans (Tables 2 and S3; Fig. 2).

3.3.2. Shell shape. The interaction between fixed effects was significant ($X^2_2 = 8.20$, $P = 0.017$), but when we statistically corrected for shell size, it became marginally non-significant ($X^2_2 = 5.69$, $P = 0.058$). Snails from pans (sex pooled) exhibited the most elongated shells relative to individuals from marshes and flats. At the sex level (habitats pooled), males had a more elongated shell shape than females (Table 2 and S3; Fig. 2).

3.3.3. Aperture size. We found a strong habitat and sex effect on aperture size; the interaction between these variables was not significant ($X^2_2 = 3.05$, $P = 0.217$). Across habitats (sex pooled), snails from pans showed the largest aperture size (mm^2) whereas in flats and marshes it was 9 to 12% smaller, respectively. Moreover, we observed a significant difference in aperture size between sexes (habitats pooled), males had bigger apertures than females and this difference was consistent across habitats (Table 2 and S3; Fig. 2).

3.3.4. Aperture shape. We found no significant difference in aperture shape between sexes ($X^2_1 = 0.65$, $P = 0.724$), and this pattern was similar even after correcting for shell size ($X^2_2 = 2.20$, $P = 0.333$). However, individuals from pans (sex pooled) showed the most rounded shape relative to those from flats and marshes, which showed a more elongated aperture shape (Table 2 and S3; Fig. 2).

3.4. Pronounced trematode effect on shell morphometric

Infection caused by the trematode *M. simillimus* strongly shifted the mean value of most

phenotypic traits of *H. australis* to lower values when comparing infected and

uninfected snails from pans.

3.4.1. Shell size. Shell size of infected snails (sex pooled) decreased 10% relative to

uninfected ones, whereas females (infection status pooled) were 5% smaller than males

(Table 3; Fig. 4). We found no significant interaction between the fixed effects (status

and sex; $X^2_1 = 0.70$, $P = 0.402$).

3.4.2. Shell shape. We found a significant interaction between status and sex ($X^2_1 =$

15.86, $P <0.001$). This was because, at the status level, uninfected males were more

elongated (higher SL to SW ratio) than uninfected females, but infected males and

females showed no difference in shell shape. We found the same pattern at the sex level;

that is, infected male and female snails have a more elongated shell relative to

uninfected male and female individuals (Table 3; Fig. 4).

3.4.3. Aperture size. We observed a strong effect of infection status and sex in aperture

size. Infected snails (sex pooled) and females (status pooled) had a smaller aperture than

uninfected and male snails (15 and 3% respectively; Table 3; Fig. 4). We found no

significant interaction between fixed effects ($X^2_1 = $ 2e-04, $P = 0.990$).

3.4.4. Aperture shape. Uninfected snails exhibited a more elongate aperture shape

(higher AL to AW ratio) than infected individuals whereas we found no sex effect on

this shell trait (Table 3; Fig. 4). We found no significant interaction between fixed

effects and no sex effect on aperture shape, even after correcting by shell size ($X^2_1 =$

0.33, $P = 0.564$).

3.5. Thicker shells and lower body mass in habitats with high (a)biotic stress

We found that snails differed in shell weight (~ thickness) across habitats ($X^2_2 = 48.93$,

$P <0.001$). Snails from pans showed the lightest/thinnest shells (11.53±0.23 mg)

compared to snails from marshes (12.54±0.15 mg; $P_{Pans<Marshes} <0.001$) and flats

(12.00±0.15 mg; $P_{Pans<Flats}$ = 0.001). By contrast, in marshes individuals showed the

heaviest/thickest shells ($P_{Marshes>Flats}$ <0.001; Fig. 4). Body mass also differed across

habitats (X^2_2 = 19.67, P <0.001). Snails from pans exhibited the heaviest body mass

(0.50±0.05 mg) relative to individuals from marshes (0.32±0.03 mg) and flats

(0.33±0.04 mg) ($P_{Pans>Marshes}$ <0.001; $P_{Pans>Flats}$ <0.001), which showed no differences in

body weight ($P_{Marshes<Flats}$ = 0.390; Fig. 5). Further details on observations, corrected and

uncorrected mean values, and statistics are indicated in the appendix (Text S4; Table

S9; Fig. S6).

3.6. Variation in penis size

The penis shape confirmed that individuals from each habitat of the Bahía Blanca

estuary belong to the same species (Fig. S8; Gaillard & Castellanos 1976). We found,

however, a clear difference in penis size ($F_{(2, 39)}$ = 8.42, P <0.001). Individuals from

marshes showed the largest size relative to flats ($P_{flats<marshes}$ <0.001) and pans

($P_{pans<marshes}$ = 0.034); snails collected from these two habitats showed a marginal non-

significant difference ($P_{flats<pans}$ = 0.090; Fig. 6). We found no significant effect of shell

size, meaning that differences in penis size were not due differences in individual's shell

size ($F_{(1, 39)}$ = 0.27, P=0.608). Further, we found no differences in shell shape, even after

controlling by shell size ($F_{(2, 43)}$ = 0.851, P = 0.435).

3.7. Molecular analysis of the cytochrome oxidase subunit 1 (COI) gene

The species-delimitation analysis implemented in ABGD found only one partition (prior

maximal distance, P = 0.001) that confirmed that the individuals here studied constitute

a single *Heleobia* species, which is in agreement with the similar penis shape found

among habitats. In fact, we did not observe any type of structure among individuals.

The gene tree and haplotype network showed that individuals did not gather in clusters

depending on the habitat they come from (Figs. S8 and S9).

4. DISCUSSION

This study provides compelling evidence of ecological and phenotypic trait variation in

response to contrasting biotic and abiotic conditions at a small spatial scale

(summarized in Fig. 7). In the following we discuss these main findings that suggest

that subpopulations living in sympatry of the intertidal snail *Heleobia australis*

experienced a fine-grained environment where the combined effect of plasticity and

environmental pressures favored the expression of potentially beneficial traits at a local

scale.

4.1. Reduced snail density under high environmental and predatory stress

Density of *Heleobia australis* was the lowest in marshes (vegetated area that drains at

low tide) relative to flats (unvegetated area that drains at low tide) and pans (covered by

water at low tide) whereas flats and pans showed no difference. However, when

analyzed by age classes, adult density was higher in pans during winter but remarkably

lower in marshes, even compared to flats (both habitats from the upper area). This is

consistent with the idea that prolonged submersion times could increase the survival

rate of snails during harsh climatic conditions (Fig. S2). As for juvenile snails, their

density increased from autumn to winter in all the three habitats and kept increasing

until spring time where it was 1.6 and 2.4 times higher in pans relative to flats and

marshes, respectively (Table S1; Fig. S2). We found little support for the idea that

intense environmental stress, combined with harsh climatic conditions, could strongly

decrease their survival rate affecting their abundance, especially, in the upper area.

Instead, these findings suggest that conditions at pans might have favored juvenile

recruitment, which could explain the differences observed in spring.

It is puzzling, however, that total density in marshes was lower relative to flats because

H. australis is positively associated with marsh plants (the smooth cordgrass *Spartina*

alterniflora, which dominates the lower marsh) as they buffer physical stress factors

(thermal and dehydration stress) relative to uncovered areas, promoting snail

aggregation (Canepuccia et al. 2007 and references therein). One possible explanation

for the low density in marshes, relative to flats, is if this resulted from a negative

interaction between *H. australis* and the grapsid crab *Neohelice granulata*, whose

aggregation is also facilitated by marsh plants (Spivak et al. 1994, Alvarez et al. 2013,

Angeletti et al. 2014, Angeletti & Cervellini 2015) where it attains density peaks in late

spring and early summer but it is absent throughout the year in unvegetated areas (flats

or pans; (Angeletti & Cervellini 2015). It is therefore likely that the low density of *H.*

australis in vegetated crab-rich habitats resulted from an increased bioturbation activity

caused by *N. granulata*, particularly during that period of time (Alvarez et al. 2013).

Another mutually non-exclusive hypothesis is that this difference between marshes and flats could result from a higher predation pressure caused by this crab (D'Incao et al. 1990, Barutot et al. 2011), in which case it might have induced predatory shell defenses as shell thickening or narrower apertures (Boulding & Hay 1993, Johannesson & Johannesson 1996, Rolán-Alvarez 2007, review by Bourdeau et al. 2015). Accordingly, snails from marshes exhibited a strong variation in shell characters that support this hypothesis, which we discuss later in this section.

4.2. Snails exhibited larger, narrower, and thinner shells and heavier body mass in habitats with low stress conditions

Snails subjected to different levels of stress showed a strong phenotypic variation. In all categories analyzed (cat. *i–iii*), individuals showed a clear shift in their phenotype likely in response to different environmental pressures (Fig. 7). Overall, the distribution of trait values tended to represent two distinct groups, one for flats and marshes and another for pans. Such responses seem to be primarily linked to the level of exposure to physical stress (temperature and salinity fluctuation and desiccation) of individuals during low tide. Prolonged submersion times enhance foraging times and the absorption of calcium carbonate from water increasing shell growth rate, which, in turn, gives rise to larger but more elongated shells (review by Chapman 1995). As there is a maximal rate at which calcium carbonate can be absorbed from the water, rapidly growing individuals produce thinner shells for the same amount of body weight than slower growing individuals. This results in a larger internal volume allowing for accommodating a higher body mass, which is related to higher growth rate favored by longer foraging times (Palmer 1981, Kemp & Bertness 1984, Chapman 1995, Trussell 2000a). Accordingly, snails that remain covered by water during low tide (pans) showed larger, narrower, and thinner shells and heavier body mass relative to individuals from habitats with high environmental stress conditions (flats and marshes). As higher body weight is directly linked to higher fecundity (i.e. egg production; Fredensborg et al. 2006), it is therefore likely that the heaviest body weight of individuals from pans was related to a higher recruitment, which would explain the increased juvenile abundance compared to flats and marshes (Fig. S2).

Juvenile and adult snails from flats and marshes showed a smaller aperture size (smaller aperture surface area) with respect to individuals from pans. It could be that higher environmental stress conditions favored the expression of a smaller aperture size as a potentially beneficial trait, thus increasing resistance to high desiccation in these habitats that drain at low tide. This is consistent with other studies showing that the size of the shell aperture in gastropods is smaller in response to hot and dry conditions (Machin 1967, Vermeij 1973, Chapman 1995, Melatunan et al. 2013, Schweizer et al. 2019). Moreover, individuals from flats and marshes showed thicker shells and lower body mass. Shell thickening could be due to a higher deposition of calcium carbonate at a slow grow rate (likely in response to unfavorable conditions in more exposed habitats from the upper intertidal) that traded off against investment in body mass (Palmer 1981).

4.3. Crab predation pressure would favor the expression of shell defenses in *Heleobia australis*

Snails from marshes showed a more rounded and thicker shell, reduced body mass, and a narrow aperture shape for three snail categories analyzed (cat. *i–iii*; Fig. 7). Such characteristics are consistent with shell defenses probably induced by the high presence of the predatory crab *Neohelice granulata* in that habitat in particular (e.g. Appleton & Palmer 1988, Palmer 1990, Bourdeau et al. 2015). Considering that crushing predators exert a strong selective pressure driving the evolution of behavioral, chemical, and morphological defense traits (Appleton & Palmer 1988, Palmer 1990, Trussell 2000b, Johannesson 2003, Bourdeau et al. 2015), our results, combined with its remarkably low density, support the hypothesis that predation pressure is stronger in marshes than in flats and pans. Yet, specific information on whether crabs actively predates on *H. australis* is limited or inexistent. Earlier studies reported that *N. granulata* consumes mollusks but in a low frequency (D'Incao et al. 1990, Barutot et al. 2011). These studies, however, have used visual examination of gut/stomach contents. This procedure may have under-estimated true predation levels as crabs only eat the soft tissue after crushing/peeling the shell whereby the remains found in guts or stomach become unidentifiable due to maceration and digestion. Thus, further studies using molecular-based tools are needed for detecting the presence of *Heleobia*'s tissue in crab's stomachs as an alternative or complementary approach to visual identification (e.g.

Albaina et al. 2010, Collier et al. 2014). By doing so, we would be able to establish the trophic link between these two species that is highly likely to exist. Evidence found in this study clearly reveals that *H. australis* phenotypically responded to the presence of predatory crabs, even early in life as juveniles from marshes also showed a narrow aperture shape and a more rounded shell. Our results therefore warrant further investigation to understand the adaptive value of the plastic shell responses of gastropods induced by predators.

4.4. Strong parasite effect on shell characters: a trade-off between growth and early reproduction

Infected and uninfected snails from pans showed a clear phenotypic differentiation in shell traits. Trematode infection inevitably leads to snail host castration drastically reducing host fitness, which, in evolutionary terms, is equivalent to death of the host (Lafferty 1993, Fredensborg et al. 2006). In this sense, being castrated by a parasite is similar as being eaten by a predator (Kuris 1974). This means that parasitic castrators can exert a strong selective force that could favor the expression of behavioral, physiological, and morphological adaptations to minimize the negative impact of parasitism on host fitness (Lafferty 1993). Early maturation, which results from fast growing individuals, is an effective strategy that increases current reproductive effort over future reproduction (Cole 1954, Lewontin 1965, Roff 1992), which equals to 0 in castrated hosts. In *Heleobia*, castration caused by *Microphallus* can have a profound impact on snail fitness as this parasite shows an extensive host exploitation occupying the entire gonad and most of digestive gland (see Fig. S2 in Alda et al. 2019).

Our results showed that infected snails had a more elongated shell shape but a smaller size compared to uninfected snails (Fig. 7). Despite infected snails were smaller, they exhibited the same number of whorls (six) as uninfected snails, which indicates that individuals analyzed were adults (Gaillard & Castellanos 1976). It could therefore be that the elongated shape resulted from an initial higher growth rate (review by Chapman, 1995) likely followed by an energy reallocation from growth to early reproduction (Lafferty 1993, Agnew et al. 1999), which could explain the smaller shell size compared to uninfected snails (Alda et al. 2010, 2019, this study). Infected snails also showed a smaller aperture size but a more rounded shape relative to uninfected

individuals (Fig. 7). This change in aperture size and shape could be a side effect of

selection for high rate of shell growth (Boulding & Hay 1993) induced by parasites.

Together, these findings support the idea that infection by trematodes exerted a strong

selective pressure on the snail host causing important shifts in the expression of shell

traits towards lower mean values, creating a pronounced phenotypic difference between

infected and uninfected snails (Poulin & Thomas 1999).

4.5. Difference in penis size in a single *Heleobia* species

Interestingly, individuals from each habitat showed a clear variation in penis size in

spite of they were raised in common garden laboratory conditions (no habitat effect)

and, when dissected, we observed that none of them was infected by parasites (no

parasite effect). Field-collected juveniles might retain the influence of environmental

conditions experienced prior to collection (Sanford & Kelly 2011). Thus, one possible

explanation is that this variation resulted from different growth rates as well as the

differences in shell shape observed for wild snails. We found, however, no differences

in shell shape among male individuals reared in laboratory conditions, even after

controlling by shell size. It is noteworthy that a strong difference in penis size emerged

despite the relatively small number of male individuals analyzed (n = 15 per habitat).

One interpretation of this phenotypic divergence is that it was not related to dissimilar

rates of growth but to genetic differences, suggesting that this is likely to be a non-

plastic trait. Despite the difference in size, its shape indicates that individuals from each

subpopulation belong to the same species, more specifically to *Heleobia australis*

australis (see Gaillard & Castellanos, 1976 for further detail on taxonomy).

Accordingly, the COI gene confirmed that specimens considered in this study constitute

a single species.

Future experimental (laboratory breeding) and non-experimental approaches (multilocus

or genomic analyses) are needed to exclude other processes that may have contributed

to this difference in penis size. These approaches will help us to understand whether this

genital divergence could be an indicative of pre-zygotic reproductive isolation among

subpopulations of *Heleobia australis* living in sympatry (Kameda et al. 2009, Hollander

et al. 2013). If this is true, it could be that individuals studied herein would be part of a

species complex that have recently diverged, which would be in line with the lack of

clusters when analyzing the conserved COI gene (Kemppainen et al. 2009, Janzen et al. 2017, Matos☐Maraví et al. 2019).

4.6. Does the small-scale phenotypic differentiation in *Heleobia australis* reflect adaptive plasticity responses to local environmental pressures?

Theory predicts that adaptive phenotypic plasticity is more common than genetic adaptation in wide-dispersing species relative to species with restricted dispersal (Mather 1955, Levins 1968, Via & Lande 1985, Scheiner 1998, De Jong 1999, 2005). That is because gene flow homogenizes genotype frequencies across space countering the effect of natural selection (review by Lenormand 2002). Our results suggest that *Heleobia australis* experienced a fine-grained environment where conditions imposed by contrasting biotic and abiotic stressors across the vertical gradient favored the expression of beneficial traits. This study was not designed to test for an adaptive basis to phenotypic trait variation, but provides a reasonable ground for advocating that the shifts in the mean trait values could be adaptations in response to the prevailing selection pressure at each habitat. If this is true, adaptive plasticity might have placed populations close enough to the new favored phenotypic optimum for directional selection to act, thus foreshadowing adaptive evolutionary responses over longer periods of time (Ghalambor et al. 2007).

The question that inevitably arises is whether there is scope for local adaptation in this intertidal mud snail. An increasing number of experimental studies indicates that local genetic adaptation may be more common than predicted by theory in marine species with planktonic dispersal, especially in systems characterized by strong selection over relatively fine spatial scale (Hays 2007, Marshall et al. 2010, Sanford & Kelly 2011). One possible explanation for the evolution of locally adapted phenotypes in species with planktonic dispersal is linked to pre- and post-colonization barriers to connectivity (i.e. phenotype-environment mismatch) that increase localized recruitment (Levin et al. 2006, Marshall et al. 2010). That is, local adaptation may be ensured by genetic polymorphism because immigration is rare and the intrapopulation gene pool is highly reliant upon local recruitment whereat individuals are exposed to local selection pressures. Localized recruitment combined with directional selection of adaptive traits may induce rapid morphological and ecological changes favoring genotypes exhibiting

the most extreme phenotypes, the so-called Baldwin Effect, or more commonly known as 'genetic assimilation' (reviewed in Waddington 1953, West-Eberhard 1989, Pigliucci et al. 2006, but see West-Eberhard 2003). Only a few studies reported local genetic differentiation in wide-dispersing marine snails, contributing to the phenotypic differentiation, as a result of local disruption to dispersal of planktotrophic larvae, post-veligers, and juvenile stages (e.g. Struhsaker 1968, Parsons 1997). In this context, it is therefore plausible that the planktotrophic snail *H. australis* benefit from the combined effect of directional selection and plasticity to evolve locally adapted phenotypes to contrasting habitat conditions across the intertidal gradient.

5. CONCLUSION

Overall, our results showed that the same phenotype was unlikely to perform best regardless the stress encountered. Contrasting environmental stressors across the vertical gradient induced important shifts in the expression of phenotypic traits, thus creating pronounced small-scale phenotypic differences. These findings support the standpoint that phenotypic plasticity favored the expression of beneficial traits thereby increasing the likelihood of persistence of subpopulations facing local environmental stressful conditions. The next step is therefore to perform more powerful experimental approaches (including common garden designs and reciprocal transplants) and molecular analyses to unveil the adaptive value of the small-scale phenotypic plasticity in this marine snail. Future studies are clearly needed to understand the spatial scale at which adaptive differentiation can occur and the relative contributions of phenotypic plasticity versus fixed genetic variation in key traits of wide-dispersing marine organisms. Such information is critical for improving the prediction of the biological impacts of climate change and for effective conservation management approaches of marine system populations.

ACKNOWLEDGEMENTS

We would like to express our gratitude to Kevin Lafferty, Patrice David, Antonio A. Vazquez Perera, Philippe Jarne, and Romain Villoutreix for their constructive suggestions on earlier versions of the manuscript. We also thank Leandro A. Hünicken and Micaela Folino for field and laboratory assistance and to Néstor J. Cazzaniga for providing insightful comments on the systematics of *Heleobia* group. This manuscript

was improved by comments from Dr. Sokolova and three anonymous reviewers. This work was partially funded by a research grant from the Universidad Nacional del Sur (UNS 24/B153 and UNS 24/B199). NB was partially supported by the "Programa de Financiamiento Parcial de Estadías en el Exterior para Investigadores Asistentes", National Scientific and Technical Research Council CONICET (Res. Nº 558 1236/08; 4118/16)".

AUTHORS' CONTRIBUTIONS

NB and PA conceived the ideas and designed methodology; NB, JPP, and PA collected the data; NB analyzed the data and led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

REFERENCES

Agnew P, Bedhomme S, Haussy C, Michalakis Y (1999) Age and size at maturity of the mosquito Culex pipiens infected by the microsporidian parasite *Vavraia culicis*. Proc R Soc Lond B 266:947–952.

Albaina A, Fox CJ, Taylor N, Hunter E, Maillard M, Taylor MI (2010) A TaqMan real-time PCR based assay targeting plaice (*Pleuronectes platessa* L.) DNA to detect predation by the brown shrimp (Crangon crangon L.) and the shore crab (*Carcinus maenas* L.)—Assay development and validation. J Exp Mar Biol Ecol 391:178–189.

Alda P, Bonel N, Cazzaniga NJ, Martorelli SR (2010) Effects of parasitism and environment on shell size of the South American intertidal mud snail *Heleobia australis* (Gastropoda). Estuar Coast Shelf Sci 87:305–310.

Alda P, Bonel N, Cazzaniga NJ, Martorelli SR, Lafferty KD (2019) A strong colonizer rules the trematode guild in an intertidal snail host. Ecology:e02696.

Alda P, Martorelli S (2014) Larval trematodes infecting the South-American intertidal mud snail *Heleobia australis* (Rissooidea: Cochliopidae). Acta Parasitol 59.

Alvarez MF, Esquius KS, Addino M, Alberti J, Iribarne O, Botto F (2013) Cascading top-down effects on estuarine intertidal meiofaunal and algal assemblages. J Exp Mar Biol Ecol 440:216–224.

Angeletti S, Cervellini PM (2015) Population structure of the burrowing crab *Neohelice granulata* (Brachyura, Varunidae) in a southwestern Atlantic salt marsh. Lat Am J Aquat Res 43:539–547.

Angeletti S, Lescano L, Cervellini P (2014) Caracterización biosedimentológica y mineralógica de dos sectores intermareales del estuario de Bahía Blanca. GeoActa 39:54–67.

Appleton RD, Palmer AR (1988) Water-borne stimuli released by predatory crabs and damaged prey induce more predator-resistant shells in a marine gastropod. PNAS 85:4387–4391.

Barutot RA, D'Incao F, Fonseca DB (2011) Natural diet of *Neohelice granulata* (Dana, 1851) (Crustacea, Varunidae) in two salt marshes of the estuarine region of the Lagoa dos Patos lagoon. Braz Arch Biol Technol 54:91–98.

Bates D, Mächler M, Bolker B, Walker S (2014) Fitting linear mixed-effects models using lme4. J Stat Softw 67:1–48

Berrigan D, Scheiner S (2004) Modeling the evolution of phenotypic plasticity. In: *Phenotypic plasticity: Functional and conceptual approaches*, DeWitt TJ, Scheiner, SM (eds). Oxford, Univ. Press, New York, p 82–97

Bonel N, Lorda J (2015) Growth and body weight variability of the invasive mussel *Limnoperna fortunei* (Mytilidae) across habitat and season. Malacologia 58:129–145.

Boulding EG, Hay TK (1993) Quantitative genetics of shell form of an intertidal snail: constraints on short-term response to selection. Evolution 47:576.

Bourdeau PE (2011) Constitutive and inducible defensive traits in co‐occurring marine snails distributed across a vertical rocky intertidal gradient. Funct Ecol 25:177–185.

Bourdeau PE, Butlin RK, Brönmark C, Edgell TC, Hoverman JT, Hollander J (2015) What can aquatic gastropods tell us about phenotypic plasticity? A review and meta-analysis. Heredity 115:312–321.

Canepuccia AD, Escapa M, Daleo P, Alberti J, Botto F, Iribarne OO (2007) Positive interactions of the smooth cordgrass *Spartina alterniflora* on the mud snail *Heleobia australis*, in South Western Atlantic salt marshes. J Exp Mar Biol Ecol 353:180–190.

Carcedo MC, Fiori SM (2012) Long-term study of the life cycle and growth of *Heleobia australis* (Caenogastropoda, Cochliopidae) in the Bahía Blanca estuary, Argentina. Ciencias Marinas 38:589–597.

Carvajal-Rodríguez A, Conde-Padín P, Rolán-Alvarez E (2005) Decomposing shell form into size and shape by geometric morphometric methods in two sympatric ecotypes of *Littorina saxatilis*. J Mollus Stud 71:313–318.

Chapman MG (1995) Spatial patterns of shell shape of three species of co-existing littorinid snails in New South Wales, Australia. J Mollus Stud 61:141–162.

Chevin L-M, Lande R (2009) When do adaptive plasticity and genetic evolution prevent extinction of a density-regulated population? Evolution 64:1143–1150.

Cole LC (1954) The population consequences of life history phenomena. Q Rev Biol 29:103–137.

Collier JL, Fitzgerald SP, Hice LA, Frisk MG, McElroy AE (2014) A new PCR-based method shows that blue crabs (*Callinectes sapidus* (Rathbun)) consume winter flounder (*Pseudopleuronectes americanus* (Walbaum)). PLoS ONE 9:e85101.

Conde-Padín P, Grahame JW, Rolán-Alvarez E (2007) Detecting shape differences in species of the Littorina saxatilis complex by morphometric analysis. J Mollus Stud 73:147–154.

Curtis L (1987) Vertical distribution of an estuarine snail altered by a parasite. Science 235:1509–1511.

De Francesco CG, Isla FI (2003) Distribution and abundance of hydrobiid snails in a mixed estuary and a coastal lagoon, Argentina. Estuaries 26:790–797.

De Francesco CG, Isla FI (2004) The life cycle and growth of *Heleobia australis* (d'Orbigny, 1835) and *H. conexa* (Gaillard, 1974) (Gastropoda: Rissooidea) in Mar Chiquita coastal lagoon (Argentina). J Mollus Stud 70:173–178.

De Jong G (2005) Evolution of phenotypic plasticity: patterns of plasticity and the emergence of ecotypes: Research review. New Phytol 166:101–118.

De Jong G (1999) Unpredictable selection in a structured population leads to local genetic differentiation in evolved reaction norms. J Evolution Biol 12:839–851.

Denny MW, Paine RT (1998) Celestial mechanics, sea-level changes, and intertidal ecology. Biol Bull 194:108–115.

Dillon RT, Jacquemin SJ, Pyron M (2013) Cryptic phenotypic plasticity in populations of the freshwater prosobranch snail, *Pleurocera canaliculata*. Hydrobiologia 709:117–127.

D'Incao F, Silva KG, Ruffino ML, Braga AC (1990) Hábito alimentar do caranguejo *Chasmagnathus granulata* Dana, 1851 na barra do Rio Grande, RS (Decapoda, Grapsidae). Atlântica 12:85–93.

Echeverría CA, Neves RAF, Pessoa LA, Paiva PC (2010) Spatial and temporal distribution of the gastropod *Heleobia australis* in an eutrophic estuarine system suggests a metapopulation dynamics. NS 02:860–867.

Elías R, Iribarne O, Bremec CS, Martinez DE (2004) Comunidades bentónicas de fondos blandos. In: *Ecosistema del Estuario de Bahía Blanca*, Piccolo M.C., Hoffmeyer M.S. EDIUNS (UNS), Bahía Blanca, p 179–190

Fox J (2003) Effect displays in R for generalised linear models. J Stat Softw 8:1–27.

Fox J, Weisberg S, Adler D, Bates D, Baud-Bovy G, Ellison S, Heilberger R (2011) Package "car": Companion to applied regression. R package version 2.0-11. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Fredensborg BL, Mouritsen KN, Poulin R (2005) Impact of trematodes on host survival and population density in the intertidal gastropod *Zeacumantus subcarinatus*. Mar Ecol Prog Ser 290:109–117.

Fredensborg BL, Mouritsen KN, Poulin R (2006) Relating bird host distribution and spatial heterogeneity in trematode infections in an intertidal snail—from small to large scale. Mar Biol 149:275–283.

Gaillard MC, Castellanos ZA (1976) Mollusca Gasteropoda Hydrobiidae. In: *Fauna de agua dulce de la República Argentina*, Ringuelet R. A. FECIC, Buenos Aires, p 1–40

Ghalambor CK, McKay JK, Carroll SP, Reznick DN (2007) Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. Funct Ecology 21:394–407.

Guerra-Varela J, Colson I, Backeljau T, Breugelmans K, Hughes RN, Rolán-Alvarez E (2009) The evolutionary mechanism maintaining shell shape and molecular differentiation between two ecotypes of the dogwhelk *Nucella lapillus*. Evol Ecol 23:261–280.

Hays CG (2007) Adaptative phenotypic differentiation across the intertidal gradient in the alga *Silvetia compressa*. Ecology 88:149–157.

Hershler R, Thompson F (1992) A review of the aquatic gastropod subfamily Cochliopinae

(Prosobranchia, Hydrobiidae). Malacological Review 5.

Hill JK, Griffiths HM, Thomas CD (2011) Climate change and evolutionary adaptations at species' range margins. Annu Rev Entomol 56:143–159.

Hoffmann AA, Sgrò CM (2011) Climate change and evolutionary adaptation. Nature 470:479–485.

Hollander J, Collyer ML, Adams DC, Johannesson K (2006) Phenotypic plasticity in two marine snails: constraints superseding life history. J Evolution Biol 19:1861–1872.

Hollander J, Smadja CM, Butlin RK, Reid DG (2013) Genital divergence in sympatric sister snails. J Evol Biol 26:210–215.

Ieno E, Alemany D, Blanco DE, Bastida R (2004) Prey size selection by Red Knot feeding on mud snails at Punta Rasa (Argentina) during migration. Waterbirds 27:493–498.

Janzen DH, Burns JM, Cong Q, Hallwachs W, Dapkey T, Manjunath R, Hajibabaei M, Hebert PDN, Grishin NV (2017) Nuclear genomes distinguish cryptic species suggested by their DNA barcodes and ecology. Proc Natl Acad Sci USA 114:8313–8318.

Johannesson B, Johannesson K (1996) Population differences in behaviour and morphology in the snail *Littorina saxatilis:* phenotypic plasticity or genetic differentiation? J Zool 240:475–493.

Johannesson K (2003) Evolution in *Littorina*: ecology matters. J Sea Res 49:107–117.

Johnson MS, Watts RJ, Black R (1994) High levels of genetic subdivision in peripherally isolated populations of the atherinid fish *Craterocephalus capreoli* in the Houtman Abrolhos Islands, Western Australia. Marine Biology 119:179–184.

Kameda Y, Kawakita A, Kato M (2009) Reproductive character displacement in genital morphology in *Satsuma* land snails. Am Nat 173:689–697.

Kemp P, Bertness MD (1984) Snail shape and growth rates: Evidence for plastic shell allometry in *Littorina littorea*. PNAS 81:811–813.

Kemppainen P, Panova M, Hollander J, Johannesson K (2009) Complete lack of mitochondrial divergence between two species of NE Atlantic marine intertidal gastropods. J Evolution Biol 22:2000–2011.

Komsta L (2011) Outliers: Tests for outliers. R package version 0.13. http://www.komsta.net

Kuris AM (1974) Trophic interactions: similarity of parasitic castrators to parasitoids. Q Rev Biol 49:129–148.

Lafferty KD (1993) The marine snail, *Cerithidea californica*, matures at smaller sizes where parasitism is high. Oikos 68:3.

Lafferty KD, Sammond DT, Kuris AM (1994) Analysis of larval trematode communities. Ecology 75:2275.

Larsson J, Westram AM, Bengmark S, Lundh T, Butlin RK (2020) A developmentally descriptive method for quantifying shape in gastropod shells. J R Soc Interface 17:20190721.

Levin LA, Ziebis W, Mendoza GF, Growney-Cannon V, Walther S (2006) Recruitment response of methane-seep macrofauna to sulfide-rich sediments: An in situ experiment. J Exp

Mar Biol Ecol 330:132–150.

Levins R (1968) Evolution in changing environments: some theoretical explorations. No. 2.
Princeton University Press.

Levri EP, Dillard J, Martin T (2005) Trematode infection correlates with shell shape and
defence morphology in a freshwater snail. Parasitology 130:699–708.

Lewontin RC (1965) Selection for colonizing ability. In: *Proceedings of the First International*
Union of Biological Sciences Symposia on General Biol. Academic Press, p 77–94

Liu H-P, Hershler R, Thompson FG (2001) Phylogenetic relationships of the Cochliopinae
(Rissooidea: Hydrobiidae): an enigmatic group of aquatic gastropods. Mol Phylogenet Evol
21:17–25.

Machin J (1967) Structural adaptation for reducing water loss in three species of terrestrial
snail. J Zool 152:55–65.

Marshall DJ, Monro K, Bode M, Keough MJ, Swearer S (2010) Phenotype–environment
mismatches reduce connectivity in the sea. Ecol Lett 13:128–140.

Martorelli SR (1991) El ciclo biológico abreviado de *Microphallus simillimus* (Travassos,
1920), comb. n. (Digenea; Microphallidae) parásito de *Heleobia conexa* (Mollusca,
Hydrobiidae) y de *Himantopus melanurus* (Aves; Recurvirostridae) en Argentina. Iheringia Ser
Zool 71:91–98.

Mather K (1955) Polymorphism as an outcome of disruptive selection. Evolution 9:52–61.

Matos Maraví P, Wahlberg N, Antonelli A, Penz CM (2019) Species limits in butterflies
(Lepidoptera: Nymphalidae): reconciling classical taxonomy with the multispecies coalescent.
Syst Entomol 44:745–756.

Meerhoff EI, Rodríguez-Gallego LR, Claramunt S (2013) Dieta de ocho especies de aves
costeras en la barra de laguna de Rocha, Uruguay. Boletín de la Sociedad Zoológica del
Uruguay 22:12–22.

Melatunan S, Calosi P, Rundle S, Widdicombe S, Moody A (2013) Effects of ocean
acidification and elevated temperature on shell plasticity and its energetic basis in an intertidal
gastropod. Mar Ecol Prog Ser 472:155–168.

Milroy S (2015) Field methods in marine science: from measurements to models. Garland
Science.

Miura O, Kuris AM, Torchin ME, Hechinger RF, Chiba S (2006) Parasites alter host phenotype
and may create a new ecological niche for snail hosts. Proc R Soc B 273:1323–1328.

Mouritsen KN, Jensen KT (1994) The enigma of gigantism: effect of larval trematodes on
growth, fecundity, egestion and locomotion in *Hydrobia ulvae* (Pennant)(Gastropoda:
Prosobranchia). J Exp Mar Biol Ecol 181:53–66.

Mouritsen KN, Sørensen MM, Poulin R, Fredensborg BL (2018) Coastal ecosystems on a
tipping point: global warming and parasitism combine to alter community structure and
function. Glob Change Biol 24:4340–4356.

Neves RAF, Valentin JL, Figueiredo GM (2010) Morphological description of the gastropod
Heleobia australis (Hydrobiidae) from egg to hatching. Braz j oceanogr 58:247–250.

Palmer AR (1981) Do carbonate skeletons limit the rate of body growth? Nature 292:150–152.

Palmer AR (1990) Effect of crab effluent and scent of damaged conspecifics on feeding, growth, and shell morphology of the Atlantic dogwhelk *Nucella lapillus* (L.). Hydrobiologia 193:155–182.

Parsons KE (1997) Role of dispersal ability in the phenotypic differentiation and plasticity of two marine gastropods. Oecologia 110:461–471.

Perillo GME, Piccolo MC, Parodi E, Freije RH (2001) The Bahia Blanca Estuary, Argentina. In: *Coastal Marine Ecosystems of Latin America*. Seeliger U, Kjerfve B (eds) Springer Berlin Heidelberg, Berlin, Heidelberg, p 205–217

Petracci PF, Delhey K (2005) Guía de aves costeras y marinas de la ría de Bahía Blanca. Harris y Cía S.R.L., Bahía Blanca.

Piccolo MC, Hoffmeyer MS (2004) El ecosistema del estuario de Bahía Blanca. Instituto Argentino de Oceanografía, Bahía Blanca, Argentina. Sapiensa., Bahía Blanca.

Pigliucci M, Murren CJ, Schlichting CD (2006) Phenotypic plasticity and evolution by genetic assimilation. J Exp Biol 209:2362–2367.

Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team (2017) Nlme: linear and nonlinear mixed effects models. R package version 3.1-131. https://cran.r-project.org/web/packages/nlme/index.html

Pointier J, Noya O, Amarista M, Théron A (2004) *Lymnaea cousini* Jousseaume, 1887 (Gastropoda: Lymnaeidae): first record for Venezuela. Mem Inst Oswaldo Cruz 99:567–569.

Poulin R, Thomas F (1999) Phenotypic variabilityinduced by parasites. Parasitol Today 15:28–32.

Pratolongo P, Kirby J, Plater A, Brinson M (2009) Temperate coastal wetlands: morphology, sediment processes, and plant communities. In: *An Integrated Ecosystem Approach, Coastal Wetlands*, Perillo, G.M.E., Wolanski, E., Cahoon, D.R., Brinson, M.M. (Eds.). Elsevier, Amsterdam, p 185–210

Pratolongo P, Perillo GME, Piccolo MC (2010) Combined effects of waves and plants on a mud deposition event at a mudflat-saltmarsh edge in the Bahía Blanca estuary. Estuar Coast Shelf Sci 87:207–212.

Price TD, Qvarnström A, Irwin DE (2003) The role of phenotypic plasticity in driving genetic evolution. Proc R Soc Lond B 270:1433–1440.

Probst S, Kube J (1999) Histopathological effects of larval trematode infections in mudsnails and their impact on host growth: what causes gigantism in *Hydrobia ventrosa* (Gastropoda: Prosobranchia)? J Exp Mar Biol Ecol 238:49–68.

Puillandre N, Lambert A, Brouillet S, Achaz G (2012) ABGD, Automatic Barcode Gap Discovery for primary species delimitation: ABGD, automatic barcode gap discovery. Mol Ecol 21:1864–1877.

Revelle W (2018) Psych: Procedures for Personality and Psychological Research. R package version 1.8.10. https://personality-project.org/r/psych/

Ripley B, Venables B, Bates DM, Hornik K, Gebhardt A, Firth D, Ripley MB (2013) Package 'mass.' R package version 7.3-23. http://www.stats.ox.ac.uk/pub/MASS4/

Roff DA (1992) The evolution of life histories: theory and analysis. Chapman & Hall, New York.

Rolán-Alvarez E (2007) Sympatric speciation as a by-product of ecological adaptation in the Galician *Littorina saxatilis* hybrid zone. J Mollus Stud 73:1–10.

Sanford E, Kelly MW (2011) Local adaptation in marine invertebrates. Annu Rev Mar Sci 3:509–535.

Scheiner SM (1998) The genetics of phenotypic plasticity. VII. Evolution in a spatially-structured environment. J Evol Biol 11:303–320.

Schweizer M, Triebskorn R, Köhler H (2019) Snails in the sun: Strategies of terrestrial gastropods to cope with hot and dry conditions. Ecol Evol 9:12940–12960.

Smith NF (2001) Spatial heterogeneity in recruitment of larval trematodes to snail intermediate hosts. Oecologia 127:115–122.

Spivak E, Anger K, Luppi T, Bas C, Ismael D (1994) Distribution and habitat preferences of two grapsid crab species in Mar Chiquita Lagoon (Province of Buenos Aires, Argentina). Helgol Meeresunters 48:59–78.

Struhsaker JW (1968) Selection mechanisms associated with intraspecific shell variation in *Littorina picta* (Prosobranchia: Mesogastropoda). Evolution 22:459–480.

Travis J (1994) Evaluating the adaptive role of morphological plasticity. In: *Ecological Morphology: Integrative Organismal Biology*, eds P.C. Wainwright&S.M. Reilly. University of Chicago Press, Chicago, p 99–122

Trussell GC (2000a) Phenotypic clines, plasticity, and morphological trade☐offs in an intertidal snail. Evolution 54:151–166.

Trussell GC (2000b) Predator-induced plasticity and morphological trade-offs in latitudinally separated populations of *Littorina obtusata*. Evol Ecol Res 2:803–822.

Vermeij GJ (1972) Intraspecific shore-level size gradients in intertidal molluscs. Ecology 53:693–700.

Vermeij GJ (1973) Morphological patterns in high-intertidal gastropods: Adaptive strategies and their limitations. Mar Biol 20:319–346.

Via S, Lande R (1985) Genotype-environment interaction and the evolution of phenotypic plasticity. Evolution 39:505.

Vu VQ (2011) Ggbiplot: A ggplot2 based biplot. R package version 0.55. https://www.rdocumentation.org/packages/ggbiplot/versions/0.55

Waddington CH (1953) Genetic assimilation of an acquired character. Evolution 7:118–126.

West-Eberhard MJ (2003) Developmental plasticity and evolution. Oxford University Press, Oxford; New York.

West-Eberhard MJ (1989) Phenotypic plasticity and the origins of diversity. Annu Rev Ecol Evol Syst 2:249–278.

Westram AM, Faria R, Johannesson K, Butlin R (2021) Using replicate hybrid zones to understand the genomic basis of adaptive divergence. Mol Ecol: 10.1111/mec.15861.

Wickham H (2011) The split-apply-combine strategy for data analysis. J Stat Softw 40:1–29.

Wickham H, Chang W (2008) Ggplot2: an implementation of the grammar of graphics. R
package version 0.8. https://ggplot2.tidyverse.org

Wickham H, François R, Henry L, Müller K (2019a) Dplyr: a grammar of data manipulation. R
package version 0.8. 0.1. https://cran.r-project.org/web/packages/dplyr/index.htmlc

Wickham H, Hester J, Chang W (2019b) Devtools: tools to make developing R packages easier.
R package version 2.2.1. https://devtools.r-lib.org

Zuur AF (ed) (2009) Mixed effects models and extensions in ecology with R. Springer, New
York, NY.

Tables

Table 1. Summary of means (±SE) and statistical significance of the two-way ANOVA testing for habitat and seasonal effect on total snail density (ind./per sample; one sample being 78.5 cm^{-2}) of the intertidal mud snail *Heleobia australis* from the Bahía Blanca estuary, Argentina. Number of samples are indicated between parentheses. Value in bold indicates overall mean. We reported raw estimates whereas the model fits for Ln-transformed density. See Methods section for details.

Variable	Seasons	Habitats			Mean by Season	Habitat effect	Habitat comparison	Effect size	Season effect	Season comparison	Effect size
		Flats (F)	Marshes (M)	Pans (P)							
Total density	Summer (Su)	132±17 (9)	52±14 (9)	128±20 (9)	104±12 (27)	$F_{(2, 106)} = 15.48^{***}$	P *vs.*M	5.49***	$F_{(3, 106)} = 6.22^{***}$	Au *vs.*Sp	4.26***
	Autumn (Au)	65±13 (9)	87±42 (9)	70±14 (9)	74±15 (27)		M *vs.* F	-3.54***		Wi *vs.*Sp	2.69*
	Winter (Wi)	92±20 (8)	70±24 (9)	206±30 (9)	124±18 (26)		P *vs.* F	1.91		Su *vs.*Sp	2.57*
	Spring (Sp)	187±36 (9)	114±36 (9)	317±34 (9)	206±26 (27)					Su *vs.*Au	1.68
										Au *vs.*Wi	1.53
	Mean by Habitat	120±14 (35)	81±15 (36)	180±20 (36)	**127±10 (107)**					Su *vs.*Wi	0.14

∗ *P* < 0.05; ∗∗ *P* < 0.01; ∗∗∗ *P* < 0.001. We applied the Holm-Bonferroni correction for multiple testing when comparing effect sizes.

Table 2. Uninfected adult snails. Summary of means (±SE) and statistical significance of habitat and sex effect on shell and aperture morphometrics of uninfected adult individuals of the mud snail *Heleobia australis* from the intertidal area of the Bahía Blanca estuary, Argentina. Variables are the same as indicated in table 2. Number of observations are indicated between parentheses, which are only shown for shell size but are the same for other traits measured.

Traits measured	Habitat type	Sex		*Sex pooled*	Habitat effect	Habitat comparison	Effect size	Sex effect	Sex comparison	Effect size
		Males (♂)	Females (♀)							
Shell size Φ	Flats (F)	6.19±0.18 (388)	6.17±0.18 (396)	6.17±0.16 (784)	*Males*	P *vs.* F	17.91***	*Flats*		
	Marshes (M)	6.78±0.13 (266)	6.64±0.11 (245)	6.71±0.11 (511)	$X_2^2 = 346.22$***	P *vs.* M	11.01***	$X_1^2 = 0.20$; n.s.	♂ *vs.* ♀	0.50
	Pans (P)	7.97±0.36 (600)	7.55±0.40 (738)	7.74±0.40 (1338)		M *vs.* F	4.08***	*Marshes*		
	Habitats pooled	7.14±0.22 (1254)	6.95±0.25 (1379)		*Females*	P *vs.* F	15.20***	$X_1^2 = 1.57$; n.s.	♂ *vs.* ♀	1.25
					$X_2^2 = 249.33$***	P *vs.* M	9.01***	*Pans*		
						M *vs.* F	3.62***	$X_1^2 = 24.30$***	♂ *vs.* ♀	4.93***
Shell shape †	Flats	2.27±0.02	2.26±0.02	2.27±0.02	$X_2^2 = 101.02$***	P *vs.* F	10.13***	$X_1^2 = 29.02$***	♂ *vs.* ♀	5.39***
	Marshes	2.32±0.01	2.29±0.01	2.31±0.01		P *vs.* M	5.31***			
	Pans	2.35±0.01	2.31±0.01	2.33±0.01		M *vs.* F	3.61***			
	Habitats pooled	2.32±0.01	2.29±0.01							
Aperture size	Flats	1.82±0.05	1.83±0.06	1.82±0.05	$X_2^2 = 362.62$***	P *vs.* F	17.10***	$X_1^2 = 8.54$**	♂ *vs.* ♀	2.93**
	Marshes	1.90±0.04	1.86±0.03	1.88±0.03		P *vs.* M	12.79***			
	Pans	2.10±0.09	2.05±0.10	2.07±0.09		M *vs.* F	1.63			
	Habitats pooled	1.97±0.06	1.94±0.07							
Aperture shape	Flats	1.73±0.03	1.73±0.03	1.73±0.03	$X_2^2 = 535.73$***	P *vs.* F	-22.91***	$X_1^2 = 0.04$; n.s.	♂ *vs.* ♀	-0.20
	Marshes	1.70±0.02	1.71±0.01	1.70±0.02		P *vs.* M	-15.35***			
	Pans	1.60±0.02	1.60±0.02	1.60±0.02		M *vs.* F	-3.86***			
	Habitats pooled	1.66±0.02	1.66±0.03							

Φ As there was a significant interaction between Habitat and Sex when testing shell size, we report effect sizes separately for this trait.

† We report estimates of shell shape corrected for shell size, as the pattern observed was not the same when removing the covariate from the model.

∗ $P < 0.05$; ∗∗ $P < 0.01$; ∗∗∗ $P < 0.001$. We applied the Holm-Bonferroni correction for multiple testing when comparing effect sizes.

 Table 3. Infected vs. Uninfected adult snails. Summary of means (±SE) and statistical significance of status (infected of uninfected) and sex effect on shell and aperture morphometrics of the intertidal mud snail *Heleobia australis* from the Bahía Blanca estuary, Argentina. In these analyses we only considered individuals from pans, which allowed for preventing habitat effect on morphometry. Shell size estimated as the volume of a cone (mm^3), shell shape as the length to width ratio (SL/SW), aperture size as the area of an ellipse (mm^2), and aperture shape as the ratio between aperture length and width (AL/AW). Number of observations are indicated between parentheses, which are only shown for shell size but are the same for other traits measured.

Traits measured	Habitat type	Sex		*Sex pooled*	Status effect	Status comparison	Effect size	Sex Effect	Sex comparison	Effect size
		Male (♂)	Female (♀)							
Shell size	Uninfected (U)	7.97±0.36 (600)	7.55±0.40 (738)	7.74±0.38 (1338)	$X_1^2 = 149.69$***	U *vs.*I	12.42***	$X_1^2 = 31.68$***	♂ *vs.*♀	5.65***
	Infected (I)	7.01±0.14 (432)	6.76±0.20 (573)	6.86±0.16 (1005)						
	Status pooled	7.59±0.28 (1032)	7.22±0.32 (1311)							
Shell shape Φ	Uninfected	2.36±0.01	2.32±0.01	2.34±0.01	*Males*			*Uninfected*		
	Infected	2.40±0.01	2.39±0.02	2.40±0.01	$X_1^2 = 11.82$***	U *vs.*I	3.43***	$X_1^2 = 37.51$***	♂ *vs.*♀	6.12***
	Status pooled	2.38±0.01	2.35±0.01							
					Females			*Infected*		
					$X_1^2 = 99.50$***	U *vs.*I	9.98***	$X_1^2 = 0.09$; n.s.	♂ *vs.*♀	-0.30
Aperture size	Uninfected	2.10±0.09	2.05±0.10	2.07±0.09	$X_1^2 = 408.65$***	U *vs.*I	21.11***	$X_1^2 = 14.95$***	♂ *vs.*♀	3.88***
	Infected	1.80±0.04	1.76±0.06	1.77±0.05						
	Status pooled	1.98±0.07	1.92±0.08							
Aperture shape	Uninfected	1.60±0.02	1.60±0.02	1.60±0.02	$X_1^2 = 60.57$***	U *vs.*I	7.83**	$X_1^2 = 1.3\mathrm{e}\text{-}3$; n.s.	♂ *vs.*♀	0.03
	Infected	1.56±0.02	1.56±0.02	1.56±0.02						
	Status pooled	1.58±0.02	1.58±0.02							

Φ As there was a significant interaction between Status and Sex when testing shell shape, we report effect sizes separately for this trait.

∗ $P < 0.05$; ∗∗ $P < 0.01$; ∗∗∗ $P < 0.001$. We applied the Holm-Bonferroni correction for multiple testing when comparing effect size

Figure Captions

Figure 1. Contrasting habitat conditions at the intertidal zone of the Bahía Blanca estuary (Villa del Mar), Argentina. Three distinct habitats can be characterized: flats, marshes, and pans. Flats and marshes are located in the upper zone and they drain at low tide, though flats are free of vegetation and marshes covered by cordgrass (*Spartina alterniflora*). Pans are free of vegetation but remain covered by water during low tide and are located close the seaward edge. Thermal, saline, and dehydration stress are strong selective forces occurring mainly in the upper intertidal area (flats and marshes) whereas these inducing agents are weaker in pans, which exhibit low environmental stress condition. Biotic agents also vary along the vertical distribution of the intertidal zone. Predation risk is higher in the upper area (marshes) mainly driven by the grapsid burrowing crab *Neohelice granulata*, which is commonly found in high abundances in marshes, though it uses the entire intertidal zone. Parasite pressure also differs along the intertidal zone. The prevalence (percentage of individuals infected) of trematodes is higher in the lower area of the intertidal zone (pans), where parasite infection is predominately caused by one extremely prevalent trematode, *Microphallus simillimus*.

Figure 2. Variation in total density (juveniles and adults pooled) of the planktotrophic snail *Heleobia australis* across habitats and seasons in the intertidal area of the Bahía Blanca estuary, Argentina. Bars represent ± 1 SE.

Figure 3. Phenotypic variation in shell and aperture morphometrics of uninfected adult snails *Heleobia australis* (cat. *i*) in the Bahía Blanca estuary, Argentina. Blue and red dots indicate mean values of each variable and sex in each habitat (flats, marshes, and pans). Mean values of shell shape were statistically corrected for shell size. Bars represent ± 1 SE.

Figure 4. Phenotypic variation in shell and aperture morphometrics of infected and uninfected adult snails *Heleobia australis* (cat. *iv*) in response to a strong parasite pressure from a habitat with low environmental stress conditions but high parasite prevalence (pans) in the Bahía Blanca estuary, Argentina. Blue and red dots indicate mean values of each variable and sex (males and females, respectively). Mean values of aperture size were statistically corrected for shell size. Bars represent ± 1 SE.

Figure 5. Shell weight (as proxy of thickness) and body mass variation of the intertidal mud snail *Heleobia australis* across habitats with different biotic and abiotic selective pressures in the Bahía Blanca estuary, Argentina. Bars represent ± 1 SE.

Figure 6. Differences in penis size (mm^2) of uninfected adult snails *Heleobia australis* collected from three environmentally distinct habitats from the Bahía Blanca estuary, Argentina and kept in standard laboratory conditions for 21 months, which ensured that individuals analyzed were all adults. Bars represent ± 1 SE.

Figure 7. Summary of the responses observed in snail population density, shell and aperture morphometrics, and shell and body weight of the intertidal mud snail *Heleobia australis* from the Bahía Blanca estuary, Argentina. Such responses were induced by: (i) Environmental pressure, whereby snails experiencing prolonged emersion times in habitats that remain exposed to hot and dry conditions during low tide (marhses and flats) exhibited lower density and lower mean values of morphological traits relative to those from less environmental stress conditions (pans), which showed opposite phenotypic responses; (ii) predation pressure caused by the presence of the burrowing crab *Neohelice granulata* could be linked to the remarkable low density in marshes and could have favored the expression of shell defenses as shell thickening, increased shell globosity (lower SL to SW ratio), narrower apertures (higher AL to AW ratio); and (iii) strong parasite pressure caused by *Microphallus simillimus* where morphological traits of infected (I) snails shifted to lower mean values compared to uninfected (U) ones; note that phenotypic differences are only shown for pans, where parasite prevalence was the highest. Responses observed in this study are indicated by color-filled boxes, white-filled boxes indicate possible mechanisms or processes that might explain the observed responses. Green color refers to responses induced by environmental pressure, orange to phenotypic variation induced by crab predation, and pink by parasite pressure. Upper arrows (↑) indicate increase/higher/longer and down arrows (↓) indicate decrease/lower/shorter.

Figures

102   **Figure 1.**

Figure 2.

Figure 3.

109   **Figure 4.**

111   **Figure 5.**

Figure 6.

115   **Figure 7.**

116