TITLE:

Transmembrane polar relay drives the allosteric regulation for ABCG5/G8 sterol transporter.

AUTHOR NAMES:

Bala M. Xavier[1,‡,#], Aiman A. Zein[1,‡,#], Angelica Venes[1,2], Junmei Wang[3,] and Jyh-Yeuan Lee[1,*]*

AUTHOR ADDRESS:

[1] Department of Biochemistry, Microbiology and Immunology, Faculty of Medicine & [2]

Biomedical Sciences Program, Department of Biology, Faculty of Science, University of Ottawa,

Ontario, Canada, K1H 8M5. [3] Department of Pharmaceutical Sciences, School of Pharmacy,

University of Pittsburgh, Pittsburgh, PA 15261, USA.

KEYWORDS:

ABCG5, ABCG8, ATP-binding cassette transporter, cholesterol, polar relay, sitosterolemia

AUTHOR INFORMATION

* Correspondence should be addressed to Junmei Wang (junmei.wang@pitt.edu) and Jyh-Yeuan

(Eric) Lee (Jyh-Yeuan.Lee@uOttawa.ca).

‡ Contributed equally.

Current address: Department of Cellular and Molecular Medicine, Faculty of Medicine,

University of Ottawa

ABSTRACT:

The heterodimeric ATP-binding cassette (ABC) sterol transporter, ABCG5/G8, is responsible for

the biliary and transintestinal secretion of cholesterol and dietary plant sterols. Missense

mutations of ABCG5/G8 can cause sitosterolemia, a loss-of-function disorder characterized by

plant sterol accumulation and premature atherosclerosis. A new molecular framework was

recently established by a crystal structure of human ABCG5/G8 and reveals a network of polar

and charged amino acids in the core of the transmembrane domains, namely polar relay. In this

study, we utilize genetic variants to dissect the mechanistic role of this transmembrane polar

relay in controlling ABCG5/G8 function. We demonstrated a sterol-coupled ATPase activity of

ABCG5/G8 by cholesteryl hemisuccinate (CHS), a relatively water-soluble cholesterol memetic,

and characterized CHS-coupled ATP catalysis by using three loss-of-function missense variants,

two sitosterolemia mutations (E146Q and R543S; within polar relay) and one sterol-binding

mutation (A540F; distant from the polar relay). The results established an *in vitro* phenotype of

the loss-of-function and missense mutations of ABCG5/G8, showing significantly impaired

ATPase activity and loss of energy sufficient to weaken the polar relay network by all three

mutants. Our data herein provide a biochemical evidence underlying the importance of the polar

relay in regulating the catalytic activity of ABCG5/G8 sterol transporter.

INTRODUCTION

All living cells depend on the ability to translocate nutrients, metabolites, and other molecules across their membranes. One major way to achieve this is through membrane-anchored transporter proteins. The evolutionarily conserved ATP-binding cassette (ABC) transporter superfamily, for example, carry out ATP-dependent and active transport of a wide range of substances across cellular membranes, including both hydrophilic and hydrophobic molecules such as sugars, peptides, antibiotics, or cholesterol (Dean et al., 2001; Dean and Allikmets, 1995; Hwang et al., 2016; Linton and Higgins, 1998). As a key component of cellular membranes, cholesterol constitutes ~50% of cellular lipid content; it is also the precursor of steroid hormones that modulate gene regulation and bile acids that enable nutrient absorption. Translocation of cholesterol molecules on biological membranes plays an essential role in maintaining cellular and whole-body cholesterol homeostasis. Thus, excess cholesterol needs to be eliminated from cells and tissues through either sterol acceptors in the circulation or direct excretion into the bile or the gut (Li et al., 2013; Patel et al., 2018). A large body of evidence indicates that ABC sterol transporters regulate cholesterol metabolism, and their defects are associated with dysregulation of whole-body cholesterol homeostasis, a major risk factor for cardiovascular diseases (Borst et al., 2000; Xavier et al., 2019). Yet we have almost no understanding of how these transporters actually translocate cholesterol molecules and how the sterol-transport process is controlled by ATP catalysis. Given the dysregulation of cholesterol metabolism as a major risk factor for cardiovascular disease, there is a pressing need to elucidate of mechanism of these transporters in moving molecules across the cell membranes.

Recent progress in solving a heterodimeric crystal structure of human ABCG5 and ABCG8 establishes a new molecular framework towards such a mechanistic understanding of

ABC sterol transporters. ABCG5 and ABCG8 are half-sized ABC sterol transporters and co-

expressed on the apical surface of the hepatocytes along the bile ducts and the enterocytes from

the intestinal brush-boarder membranes (Graf et al., 2003, 2002). ABCG5 and ABCG8 function

as obligate heterodimers (ABCG5/G8) and serve as the primary and indispensable sterol-efflux

pump that effectively exports excess cholesterol, non-cholesterol sterols, and dietary plant sterols

into the bile and the intestinal lumen. In mammals, most cholesterol is eliminated by its

metabolism into bile acids or via biliary secretion as free cholesterol. The latter is considered as

the last step of reverse cholesterol transport (RCT), where ABCG5/G8 accounts for more than

75% biliary cholesterol secretion (Bhattacharyya et al., 1991; Klett et al., 2004; Plösch et al.,

2004; Yu et al., 2002). Recent studies have shown that in human subjects and animal models,

ABCG5/G8 is also responsible for eliminating neutral sterols by the transintestinal cholesterol

efflux (TICE), a cholesterol-lowering process independent of RCT (Jakulj et al., 2016).

Physiologically, ABCG5/G8 thus plays an essential role in controlling cholesterol homeostasis in

our bodies.

In general, the smallest functional unit of an ABC transporter consists of two

transmembrane domains (TMD1 and TMD2) and two nucleotide-binding domains (NBD1 and

NBD2), and both NBDs concertedly bind and hydrolyze ATP to provide the energy and drive

substrate transport. The TMDs, on the other hand, have shown to share low sequence similarity

in the amino acid sequences and three-dimensional structural folds, suggesting substrate-specific

mechanisms for individual transporters (Ford and Beis, 2019). Mechanistic analyses of ABC

cholesterol transporters have largely centered on sequence requirement at the canonical ATP-

binding sites (Hirayama et al., 2013; Takahashi et al., 2006; Wang et al., 2011; Zhang et al.,

2006), whereas limited is known about sterol-protein interaction and its relationship with the

ATP catalysis. Recent progress solving a crystal structure of human ABCG5/G8 revealed a

unique TMD fold and several structural motifs (Lee et al., 2016). Of particular, for each subunit,

a network of polar and charged amino acids is present in the core of the TMD, namely polar

relay, whose role remains to be characterized. A triple-helical bundle is located at the

transmission interface between the NBD and the TMD and consists of an elbow connecting

helix, a hot-spot helix (also known as E-helix), and an intracellular loop-1 (ICL1) coupling helix.

However, on the triple-helical bundle or the transmembrane polar relay, several residues have

been shown to bear disease-causing missense mutations from sitosterolemia or other metabolic

disorders with lipid phenotypes (**Figure 1A**). Notably, several disease-causing mutations are

clustered in the membrane-spanning region or at the NBD-TMD interface (Xavier et al., 2019;

Zein et al., 2019). This suggests the unique roles of these structural motifs in regulating the

ABCG5/G8 function; yet, no prior knowledge was available to explain the role of these

structural motifs in the sterol-transport function.

Loss-of-function (LOF) mutations in ABCG5 or ABCG8 are linked to sitosterolemia, a

rare autosomal recessive disease, while several other missense mutations are also associated with

other lipid disorders, such as gallstone formation or elevated LDL cholesterol (Berge et al., 2000;

Buch et al., 2007; Chen et al., 2008; Kajinami et al., 2004; Kuo et al., 2008; Lu et al., 2001). At

the cellular level, many of the missense mutations lead to defects in post-translational trafficking

of ABCG5/G8 from the endoplasmic reticulum (ER), an abnormality commonly observed in

other ABC transporters with missense mutations, *e.g.*, DF508 mutation in the cystic fibrosis

transmembrane conductance regulator (CFTR or ABCC7) (Graf et al., 2004; Lukacs et al.,

1994). However, specific missense mutants of ABCG5/G8 heterodimers have shown no defect in

protein maturation (Graf et al., 2004), suggesting alternative disease-causing mechanisms.

Therefore, studies of these mutants will not only show how they alter the transporter activity, but

will also provide mechanistic insights into the function of wild-type (WT) ABCG5/G8 sterol

transporter.

Disease mutations are instrumental in studying the mechanisms of affected proteins *in*

vitro, *e.g.*, familial hypercholesterolemia mutations for proteins involved in low-density

lipoprotein metabolism (Kizhakkedath et al., 2019). Guided by the structural framework of

ABCG5/G8, we can now investigate its mechanisms using enzymological approaches with

purified proteins. For this, we first need to establish at least one robust and consistent *in vitro*

functional assay. Using ATPase activity as the functional benchmark in this study, we have

optimized an *in vitro* colorimetric ATPase assay that allows high-throughput activity assessment

of detergent-purified ABCG5/G8. Using a soluble cholesterol memetic, cholesteryl

hemisuccinate (CHS), we report here the CHS-stimulated ATP hydrolysis by ABCG5/G8 proteo-

micelles, consisting of phospholipids, cholate, and dodecyl-maltoside (DDM), and present an

enzymatic analysis for the sterol-coupled ATPase activity on ABCG5/G8 sterol transporter.

Using ATPase activity as functional readout of ABCG5/G8, we show differentially inhibition of

the CHS-stimulated ATPase activity by three LOF missense mutants, two sitosterolemia

mutations and one sterol-binding mutation, where residues bearing the two disease mutations are

located along the polar relay. Our data hereby demonstrate the mechanistic basis on regulating

ABCG5/G8 function by the transmembrane polar relay (**Figure 1B**).

RESULTS

CHS stimulates ATP hydrolysis by wild-type (WT) ABCG5/G8. Despite the known physiological role of ABCG5/G8 in biliary and intestinal cholesterol secretion, the direct evidence of sterol-coupled transporter activity was not available. In this study, we investigated sterol-coupled ATPase activity by using CHS, a cholesterol mimetic that is more soluble in aqueous solution. First, to overcome low sensitivity of detecting the ABCG5/G8 ATPase activity by previous protocols, we have optimized the ATPase assay for ABCG5/G8 by adopting a colorimetric bismuth citrate-based detection approach (Cariani et al., 2004), as described in Materials and Methods. This assay significantly reduces the background noise due to cloudiness by phospholipid/cholate/DDM mixtures, which improves the detecting sensitivity of liberated inorganic phosphate within the first few minutes and allows us to calculate more accurate rates of ATP hydrolysis. We show here that CHS can significantly stimulate ABCG5/G8-mediated ATP hydrolysis when co-incubated with sodium cholate (a bile acid) and *E. coli* polar lipids (**Figure 2**). Using 5 mM of ATP, the basal activity of ABCG5/G8 was calculated as 160 ± 15 nmol/min/mg (n = 4), similar to reported values, whereas in the presence of CHS, the specific ATPase activity of ABCG5/G8 reached 565 ± 30 nmol/min/mg (n = 8), three-four times higher than that in the absence of CHS (**Figures 3A & 3B**). The activity was inhibited either by orthovanadate, an ATPase inhibitor (Davies and Hol, 2004) (**Figure 3C**), or by a catalytically-deficient mutant ABCG5$_{WT}$/G8$_{G216D}$ (G8-G216D) (Zhang et al., 2006), which displayed no ATP hydrolysis (**Figures 3A & 3B**).

The lipid environments fine-tune ABCG5/G8 ATPase activity. ABC transporters need to function in phospholipid-embedded environment. However, it is unknown whether the

ABCG5/G8 function is controlled by phospholipids of specific headgroups or in specific lipid

compositions. Because a high concentration of bile acids is required to activate ABCG5/G8

ATPase activity, attempts to use reconstituted proteoliposomes failed due to the immediate

solubilization of the reconstituted proteins. To facilitate the assessment of mutant functions, we

evaluated the lipid environments to obtain the most optimal assay conditions. To study the effect

of lipid conditions and phospholipid species on the ABCG5/G8 function, we analyzed the CHS-

coupled ATPase activity in the presence of two polar lipid extracts under a condition of fixed

concentrations of sodium cholate and CHS (Materials and Methods). Using *E. coli* polar lipids,

we carried out ATP concentration-dependent ATPase assay to determine the Michaelis-Menten

kinetic parameters of CHS-stimulated ATP hydrolysis. We have observed the maximal ATP

hydrolysis by ABCG5/G8 at concentrations slightly over 2.5 mM of ATP with a V_{max} of 677.1 ±

25.6 nmol/min/mg, a K_M(ATP) of 0.60 mM, and a k_{cat} of 1.69 s^{-1}. When using bovine liver polar

lipids, we observed ~3.5-fold lower catalytic rate of ATP hydrolysis and ~50% higher K_M(ATP)

(**Figure 4A & Table 1**). To determine the dependence of phospholipid headgroups, we tested

three most abundant phospholipids in either lipid extract on the ATP hydrolysis by ABCG5/G8,

i.e., phosphatidylethanolamine (PE), phosphatidylcholine (PC), and phosphatidylglycerol (PG)

(see Materials and Methods). Preincubation with egg PE resulted in the highest specific activity,

while the use of soy PC or egg PG only led to slightly higher ATP hydrolysis than the basal

activity (**Figure 4B**). Interestingly, PE alone was sufficient to enhance ATP hydrolysis to almost

the same level as *E. coli* polar lipids, which was higher than that of liver polar lipid mixture.

Missense mutants impair CHS-coupled ATPase Activity of ABCG5/G8. Using the CHS-

coupled ATPase activity as the functional readout, we initiated studies in the catalytic

mechanism of ABCG5/G8 by exploiting the transporter's missense mutations that undergo

proper trafficking to post-ER cell membranes (ER-escaped mutants). In this study, we have used

Pichia pastoris yeast and expressed recombinant proteins of G8-G216D, a catalytically deficient

mutant, $ABCG5_{E146Q}/G8_{WT}$ (G5-E146Q) and $ABCG5_{WT}/G8_{R543S}$ (G8-R543S), two loss-of-

function/sitosterolemia missense mutants, and $ABCG5_{A540F}/G8_{WT}$ (G5-A540F), a loss-of-function

mutant with putative sterol-binding defect (**Figure 1B & Supplementary Figure 1**).The purified

mutants were preincubated with *E. coli* polar lipids and sodium cholate as described above. The

sitosterolemia missense mutants, G5-E146Q and G8-R543S, show a ~80% reduction of the

specific activity in CHS-coupled ATP hydrolysis when compared to WT, 160±15 nmol/min/mg

and 150 ± 5 nmol/min/mg respectively. The sterol-binding mutant G5-A540F shows a ~90%

reduction of ATP hydrolysis, 90 ± 10 nmol/min/mg (**Figure 5**). We then performed ATP

concentration-dependent experiments and analyzed the Michaelis-Menten kinetics for these three

mutants. For all mutants, $K_M(ATP)$ remained nearly the same as compared to WT, but the

mutants displayed a 40-60% reduction in the catalytic rate (**Table 1**). This result suggests that

other molecular events, *e.g.*, sterol-transporter interaction, may have a more profound impact on

ATPase activity than the nucleotide association.

The effects of CHS on ABCG5/G8 WT and mutants were further investigated by measuring the

ATP hydrolysis in the CHS concentration-dependent manner at a saturated ATP concentration

(5mM here). Purified proteins were preincubated with *E. coli* polar lipids, sodium cholate and a

wide range of CHS concentrations (0.064mM - 4.1mM). For WT, we obtained a V_{max} of 702.9 ±

50.7 nmol/min/mg, a $K_M(CHS)$ of 0.80 mM, and a k_{cat} of 1.74 s^{-1} (**Figure 6 & Table 2**). The

catalytic rates of the mutants decreased by 70-90%, but except for G5-A540F, both G5-E146Q

and G8-R543S displayed significantly larger $K_M(CHS)$, up to 200% increase. This suggests a

more profound impact of sitosterolemia mutations on the ABCG5/G8 ATPase activity through

sterol-protein interaction or structural changes.

Missense mutations cause conformational changes at the ATP-binding site. To examine the

relationship between structural changes of missense mutations and their impact on the ATPase

activity, we have performed molecular dynamics (MD) simulations for the WT and three mutants

in this study. We then analyzed the MD structures to understand how the mutations could lead to

different conformation around the hypothetical surrounding residues at the nucleotide-binding

sites (NBS). These residues were obtained through a structural comparison between the crystal

structure of ABCG5/G8 (PDB ID: 5DO7) and a cryo-EM structure of ABCG2 (PDB ID: 6HBU)

for which two ATP were bound in the homodimer (Lee et al., 2016; Manolaridis et al., 2018). In

this study, we have focused on the active nucleotide-binding site (known as NBS2) in

ABCG5/G8 (Lee et al., 2016) and analyzed residues 48-62, 68, 86-94, 134, 218, 250, 251, 267,

268, 302 of ABCG5 and 242-244 of ABCG8. As shown in **Figure 7**, the mutations at the three

sites can lead to global changes on the overall ABCG5/G8 structure, with values of root-mean-

square deviations (RMSDs) larger than 2.0 Å. The difference between the RMSDs of the

secondary structures is smaller, probably because more obvious changes need longer simulation

time to manifest. We are especially interested in the mutational effect on the ATP-binding site

and generated The RMSD *vs*. Simulation Time curves for those hypothetic surrounding residues

(**Supplementary Figure 2**). We observed that the RMSDs with and without least-square (LS)

fitting are very stable for the WT, whereas for G5-E146Q and G5-A540F, both the LS Fitting

and No-Fitting RMSD are significantly larger. However, G8-R543S mutation did not lead to

significantly larger RMSD, probably because the mutation effect needs longer simulation time to

reach the ATP-binding site. Indeed, the RMSD has a trend of getting large along the MD

simulation time for G8-R543S (**Supplementary Figure 2D**). Overall, we observed a significant

perturbation on the conformations of the putative surrounding residues due to the mutations at all

the three sites.

We then identified representative MD conformations for all four protein systems for

comparison (**Figure 8**). It is observed that three positively charged residues, K92, H49 and R50

of ABCG5 are overlaid very well between the crystal and MD structures for the WT (**Figure**

8E), while for the three mutants, at least one residue cannot be well-superimposed. Interestingly,

R52, another charged residue in ABCG5, is more distant from the center of the ATP-binding site.

Therefore, the conformational changes from our molecular modeling can qualitatively explain

why the three mutations can lead to impaired ATPase activity. Of particular note, the K92 is part

of the Walker A motif at the active nucleotide-binding site and required for ABCG5/G8

functions (Johnson et al., 2010; Zhang et al., 2006).

DISCUSSION

In this study, we have shown that CHS stimulates the ATPase activity of the human

ABCG5/G8 sterol transporter (**Figure 3**). The specific activity of ATP hydrolysis by ABCG5/G8

is by far the highest in comparison with the previously reported values (Johnson et al., 2010). In

contrast, when preincubated with sodium cholate and *E. coli* polar lipids, the purified G8-G216D

mutant displayed no ATPase activity, regardless of whether CHS is present or not. CHS is a

relatively water-soluble cholesterol analog and is used to mimic cholesterol in membrane protein

crystallization (Hanson et al., 2008; Lee et al., 2016). Our results showing CHS-stimulated

ATPase activity suggest that the sterol molecules may have played a role in promoting an active

conformation for the ATPase and/or enhancing the stability of ABCG5/G8. This idea of protein

stability is supported by recent findings showing that CHS stabilizes a variety of human

membrane proteins (Neumann et al., 2017). Our results herein provide the first biochemical

evidence of ABCG5/G8 functionality that is directly coupled with sterol substrates. The much

increased CHS-coupled ATPase activity indicates that ABCG5/G8 may need such a high ATP

catalytic rate to achieve the sterol-transport function across the cellular membranes.

We further demonstrated the stimulation of activity by CHS in a concentration-dependent

manner and increasing concentrations of ATP (**Figures 4, 5 & 6**). In the presence of *E. coli* polar

lipids, the catalytic rates were similar between the CHS and ATP-dependent ATPase activities, a

V_{max} of ~700 nmol/min/mg, which is about four times higher than that in the presence of liver

polar lipids (**Tables 1 & 2**) and more than two-fold higher than the previously reported value,

~290 nmol/min/mg (Johnson et al., 2010). Using *E. coli* polar lipids, the K_M(ATP), 0.6 mM, is

about half of the previously reported 1 mM (Johnson et al., 2010), while in the presence of liver

polar lipids, K_M(ATP) remains ~1 mM (**Table 1**). In the current study, polar lipids, cholate (bile

acid) and CHS were also present in the reaction, indicating that the presence of *E. coli* polar

lipids results in higher ATP association and consequently better stimulates ABCG5/G8 ATPase

activity. When comparing the calculated values of k_{cat} and k_{cat}/K_M, we indeed observed an overall

5-fold higher turnover rate in the presence of *E. coli* polar lipids than liver polar lipids.

Interestingly, PE, the phospholipid found in both *E. coli* and liver lipids, is sufficient to

stimulate ATP hydrolysis in ABCG5/G8 to almost the highest activity, as reported in this study.

Using another phospholipid, PC or PG, alone, the specific activity of ABCG5/G8 was higher

than that obtained with the liver polar lipid mixture (**Figure 4**). PE is the major phospholipid of

the *E. coli* polar lipids, ~60%, and the second abundant phospholipid in the bile canalicular

membranes and the small intestine brush-border membranes, ~25% and ~40% respectively of

total phospholipids (Hauser et al., 1980; Kurumi et al., 1991). It has been shown that PE

preferentially fits the headgroup-binding sites on integral membrane proteins (Lee, 2004); thus,

PE may be recruited as better phospholipids to support ABCG5/G8 function in the cell

membranes. The approximate ratio of lipids for either *E. coli* or liver polar lipids may contribute

to the apparent difference in activity, but it remains unknown how phospholipid composition

regulates the transporter function. These individual lipids will be subjected to further

examination to define the lipid specificity on the ABCG5/G8 ATPase activity.

By mapping disease-carrying residues on the apo structure of ABCG5/G8, we have found

that most missense variants occur within or near the structural motifs consisting of several

conserved amino acids (Zein et al., 2019). Several missense mutations (ER-trapped) prevent

protein maturation from the endoplasmic reticulum (ER), but at least five mutations (ER-

escaped) have been shown to undergo proper trafficking to post-ER cell membranes (Graf et al.,

2004). So far, no report has shown the impact of these ER-escaped missense mutants on

ABCG5/G8 function using either *in vitro* or *in vivo* models. In this study, we have used purified

proteins from *Pichia pastoris* to investigate the functional activity of ABCG5/G8 *in vitro* and

aimed to establish the mechanistic basis of ABCG5/G8 through analyzing the structure-function

relationship of its loss-of-function missense mutations. The sitosterolemia missense mutants G5-

E146Q and G8-R543S have shown a reduction of CHS-coupled ATP hydrolysis, but retained

~20% activity as compared to WT, while the putative sterol-binding mutant G5-A540F has

shown further reduction to ~10% of WT ATPase activity (**Figures 5 & 6**). With such activity

reduction, the mutant proteins maintained the ATPase activity similar to the basal level, as

shown by WT, suggesting a remote and allosteric regulation to keep ATPase active during the

reaction. As predicted by MD simulation, the ATP-bound homology model underwent global

conformational changes upon introducing the mutations (**Figure 7**). These mutations, albeit

relatively far away from the nucleotide-binding site, can cause significant structural

rearrangement of the residues within the region that encompass the active NBS2 (**Figure 8**).

Such conformational changes may alter responses to sterol-protein interaction necessary for

maximal ATPase activity.

In the atomic model of ABCG5/G8 (PDB ID: 5D07), G5-E146 is located on the E-helix

of the triple-helical bundle and in proximity to ABCG5's polar relay, while G8-R543 is part of

ABCG8's polar relay in the core of TMD (**Figure 1**). Both triple-helical bundle and polar relay

are believed to form a network of hydrogen bonding and salt bridges and play an important role

in inter-domain communication during the transporter function (Lee et al., 2016). G5-E146 and

G8-R543 are found in the proximity of the hydrogen-bond distance with Arginine 377 of

ABCG5 (G5-R377) and Glutamate 503 of ABCG8 (G8-503), respectively (**Figure 1B**). Based

on the ATP-dependent experiments (**Figure 5 & Table 1**), we obtained the changes of Gibbs

free energy from WT to each mutant ($\Delta\Delta G_{MUT}$) as $\Delta\Delta G_{E146Q}$ = ~11.7 kJ/mol and $\Delta\Delta G_{R543S}$ = ~12.3

kJ/mol. Such energetic loss is in the range of intramolecular hydrogen-bonding potential

observed on transmembrane a-helical bundles (Feldblum and Arkin, 2014). Therefore, the results

support the hypothesis that the E-helix and the polar relay are responsible for transmitting signals

between NBD and TMD. Slightly lower $\Delta\Delta G_{MUT}$ was observed from CHS-dependent

experiments (**Figure 6 & Table 2**), $\Delta\Delta G_{E146Q}$ = ~10.0 kcal/mol and $\Delta\Delta G_{R543S}$ = ~9.0 kcal/mol.

This falls in the range of hydrophobic interaction and argues weakened sterol-transporter

interaction due to these disease mutations. As for the sterol-binding mutant, we obtained higher

energetic loss, but similar $\Delta\Delta G_{MUT}$ from ATP- or CHS-dependent analysis, $\Delta\Delta G_{A540F}$ = ~15.8 or

~16.1 kJ/mol, respectively. This likely indicates a strong hydrophobic interaction between sterols

and the transporter, as no obvious hydrogen donors/acceptors can be found at the putative sterol-

binding site on the crystal structure. In addition, G5-A540 is distant from the polar relay (>10Å

away); thus, this data suggests a remote contact by sterol molecules to control the sterol-coupled

signaling, likely through the polar relay in the transmembrane domains. In the ATP

concentration-dependent experiments, the K_M values for ATP remained almost the same (**Table**

1), suggesting that ATP binding was not affected by these mutants. The K_M values for CHS was

significantly increased in the disease mutants, but not the sterol-binding mutant (**Table 2**),

suggesting that CHS interacts with ABCG5/G8 and remotely regulates the turnover of ATP

hydrolysis in either sequential (Mode 1) or concerted (Mode 2) pathway (**Figure 9**).

Collectively, these results argue that a working network of E-helix and polar relay is essential to

maintain the communication between ATPase and sterol-binding activities in ABCG5/G8, which

are impaired by the loss-of-function missense mutations.

In conclusion, these studies show that CHS stimulates ABCG5/G8 ATPase activity and

may promote an active conformation for ABCG5/G8-mediated sterol transport. The enzymatic

characterization of three loss-of-function missense variants provides a mechanistic basis of how

the polar relay contributes to the inter-domain communication for the sterol-coupled ATPase

activity in ABCG5/G8 and may be directly involved in such ligand-protein interactions. Further

studies will reveal more insight into these molecular events and enable sterol-lowering

therapeutics to treat sitosterolemia and hypercholesterolemia.

MATERIALS AND METHODS

Materials. *E. coli* polar lipids (Cat. #: 100600C) and bovine liver polar lipids (Cat. #: 181108C)

were from Avanti Polar Lipids, Inc. (via MilliporeSigma). Cholesterol, cholesteryl hemisuccinate

(CHS), and n-Dodecyl β-D-maltopyranoside (DDM) were from Anatrace. The Ni-NTA agarose

resin was from Qiagen, the calmodulin (CBP) affinity resin, zeocin, and ampicillin were from

Agilent. Imidazole, ε-aminocaproic acid, sucrose, yeast extract, tryptone, peptone, yeast nitrogen

base (YNB), and ammonium sulfate were obtained from Wisent. ATP disodium trihydrate, Tris-

(2-carboxyethyl)-phosphine (TCEP), sodium chloride, glycerol, ethylene diamine-tetraacetic acid

(EDTA), ethylene glycol-bis(-aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA), sodium

dodecyl sulfate (SDS), Ponceau S solution, sodium azide, Bradford reagents, Tween 20,

magnesium chloride, calcium chloride, and all protease inhibitors were obtained from Bioshop

Canada. Biotin, sodium cholate hydrate, L-ascorbic acid, ammonium molybdate, bismuth citrate,

sodium citrate, methanol, ammonium hydroxide, hydrochloric acid, and acetic acid were

obtained from MilliporeSigma. Dithiothreitol (DTT), Tris base and Tris acetate were obtained

from ThermoFisher. Clarity Western ECL substrates, 30 % acrylamide, agarose and ammonium

persulfates were obtained from Bio-Rad. Restriction enzymes were obtained from New England

Biolabs, Promega and ThermoFisher. YPD: Yeast extract Peptone Dextrose; YPDS: Yeast

extract Peptone Dextrose Sorbitol; MGY: Minimal Glycerol Yeast nitrogen base; mPIB

(minimal protease inhibitor buffer for yeast cell lysis): 0.33 M sucrose, 0.3 M Tris-HCl (pH 7.5),

1 mM EDTA, 1mM EGTA, and 100mM ε-aminocaproic acid, ddH$_2$O to a final volume of 1 L

and stored at 4°C.

Cloning of ABCG5/G8 missense mutants. The expression vectors (pLIC and pSGP18),

carrying human ABCG5 (NCBI accession number NM_022436) and human ABCG8 (NCBI

accession number NM_022437), were derived from pPICZB (Invitrogen) as described (Johnson

et al., 2010; Wang et al., 2006), pLIC-ABCG5 and pSGP18-ABCG8, respectively. A tandem

array of six histidines separated by glycine ($His_6GlyHis_6$) was added to the C terminus of

ABCG5, and A tag encoding a rhinovirus 3C protease site followed by a calmodulin binding

peptide (CBP) was added to the C terminus of ABCG8. To generate the missense mutants in this

study, we performed site-directed mutagenesis by using WT ABCG5 or ABCG8 as the templates

and the following codon-optimized oligonucleotide primers (Eurofins Genomics Canada). G5-

A540F: CCATTTTTGGGGTGCTTGTTGGATCTGGATTCCTCAG   (forward)   and

GCACCCCAAAAATGGACAGCAGAGCCACTACAC   (reverse);   G5-E146Q:

GCGCCAAACGCTGCACTACACCGCGCTGC   (forward)   and

CAGCGTTTGGCGCACGGTGAGGCTGCTCAG   (reverse);   G8-R543S:

GTTGCTCTATTATGGCCCTGGCCGCCGC   (forward)   and

GCCATAATAGAGCAACAGAAGACCACCAGCCAC   (reverse);   G8-G216D:

ACGAGCGCAGGAGAGTCAGCATTGGGGTGCAG   (forward)   and

CTCTCCTGCGCTCGTCCCCCGACAACCCC (reverse). The polymerase chain reaction (PCR)

included 1-unit Phusion High-Fidelity DNA Polymerase (New England Biolabs), 1x Phusion

buffer, 200 mM dNTP, 2% (v/v) DMSO, 100 ng DNA templates, and 0.4 mM forward and

reverse primers. Each mutant-containing plasmid was amplified by the following PCR setting:

initial DNA denaturation (98°C, 2 minutes), followed by 30 cycles of denaturation (98°C, 15

seconds) / primer annealing (55°C, 30 seconds) / DNA extension (72°C, 3 minutes), then final

extension (72°C, 20 minutes). 5μl of the PCR products was run on a 1% agarose gel to confirm

the amplification, and 1µl of Dnp1 restriction enzymes (20 units) was used to digest the WT

templates overnight at 37°C to. The modified plasmids were cleaned up by ethanol acetate

precipitation technique. 5µl of 3M Sodium acetate was added to each 50µl PCR product. 200µl

of 100% Ethanol was added to each tube, vortexed, and left at room temperature for 10 minutes.

At max speed in a table centrifuge for 10 minute the plasmids were pelleted the supernatant was

removed then washed by 75% ethanol. Residual ethanol was dried by a Speed-Vac at the

maximal speed for 20 minutes at room temperature. The pellet was resuspended in ddH$_2$O.

Mutants plasmids were cloned into XL1-Blue competent *E. coli* cells by the heat-shock approach

as described in the supplier's manual (Novagen/Agilent) and by antibiotic selection using Zeocin

(Invitrogen/ThermoFisher). Using PureYield Plasmid Midiprep kit (Promega), DNA

preparations of selected clones were subjected to sequencing at Eurofins Genomics Canada.

Expression of ABCG5/G8 missense mutants in *Pichia pastoris* yeast (Supplementary Figure

1A). Both WT and mutant plasmids (20 mg each plasmid) were linearized using PmeI and co-

transformed into the *Pichia* strain KM71H by electroporation. Immediately the cells were

resuspended with 1-mL ice cold 1M sorbitol and incubated at 30°C for 1 hour. Then 5mL fresh

YPD were added and incubated for 6 hours at 250 rpm and 30°C. The cells were then centrifuged

at 3000xg for 10 minutes and resuspended with 200µL of YPD. 100 µL of transformants were

plated on YPDS plates containing 100 (low), 500 (medium) or 1000 (high) µg/mL of Zeocin to

screen for successful transformation. Seven colonies were picked and grown in 10 mL of MGY

media for 24 hours in sterile 50 mL tubes at 250 rpm and 30°C. The cells were centrifuged for 10

minutes at 3000xg and the resuspended with 10 mL of MM media. 50 µL methanol was added to

the media and once again after 12 hours. The cells were harvested after 24-hour incubation at

250 rpm and 30°C, resuspended in 600 µL mPIB buffer transferred into 1.5 mL Eppendorf tube.

After adding 500 μL glass bead, protease inhibitors, and 10mM DTT, the cells were lysed using

a mini-bead beater (Biospec), 1.5 minutes beating and 1.5 minutes rest on ice for 3 cycles. The

unbroken cells and beads were pelleted by centrifugation at 5000×g for 5 minutes at 4°C,

followed by 21130xg for 5 minutes at 4°C. The supernatant was collected, and the concentration

of the total proteins was estimated by Bradford assay. 1 µl of cell lysate alongside a 0 µg to 10

µg BSA standards were each added to a 200 µl Bradford reagent on a 96-well plate. Absorbance

at 595 nm was used to measure the protein concentrations using a Synergy H1 Hybrid reader.

The cell lysates (20 or 30 µg of total proteins) was resolved by SDS–PAGE, and protein

expression was analyzed by immunoblotting using monoclonal anti-RGSH4 antibodies (Qiagen)

to detect ABCG5 and polyclonal anti-hABCG8 antibodies (Novus Biologicals) to detect

ABCG8. The clones expressing the highest level for both subunits were selected and stored in

20% glycerol at -75°C.

Cell culture and microsomal membrane preparation. The conditions for cell growth and WT

protein induction were as described (14). Briefly, cells were initially grown at 30°C to

accumulate cell mass in an Innova R43 shaker (Eppendorf) at 250 rpm for 24-48 hours with the

pH maintained at pH 5-6. To induce protein expression, cells were left fasting for 6-12 hours,

then incubated with 0.1% (v/v) methanol for 6–12 hours at 20 or 28°C. The methanol

concentration was increased to 0.5% (v/v) by adding methanol every 12 hours for 48-60 hours.

Cell pellets were collected and resuspended in mPIB and stored at -75°C. Approximately 45 ± 10

g of cell mass was typically obtained from 1 L of cultured cells. The frozen cells were thawed

and lysed using a C3-Emulsifier (Avestin) in mPIB in the presence of 10 mM DTT and protease

inhibitors (1 µg/ml leupeptin, 1 µg/ml pepstatin A, 1 µg/ml aprotinin, and 2 mM PMSF). The

microsomal membranes were then prepared, as described (Lee et al., 2016).

Purification of ABCG5/G8 and its mutants. Both WT and mutants were purified following a

protocol as described previously (Lee et al., 2016), with minor modification. Briefly, DDM-

solubilized membranes were subjected to a tandem affinity column chromatography, first using

Ni-NTA and then CBP. The N-linked glycans and the CBP tag remained on the purified

heterodimers, and the CBP eluates were further purified by gel-filtration chromatography using a

Superdex 200 Increase 10/300 GL column on an ÄKTA Pure purification system (Cytiva,

formerly GE Healthcare Life Sciences). The proteins in the peak fractions were collected and

concentrated to 1-3 mg/mL for storage at -75°C. Noticeably, the final yield for mutants was

lower than WT, in a range of 400-800μg per six liters of cells. The expression level of the mutant

proteins in the microsomes and their solubility was slightly lower than WT. Some proteins were

also lost during Ni-NTA binding and imidazole wash. The profile of the gel-filtration

chromatography often showed a higher peak at the void volume than dimeric proteins. These

factors collectively suggest that the mutant proteins are more prone to aggregation, and thus

explain the lower yields.

ATPase assay. We have consistently observed a strong cloudiness in the assay solution when

using previous protocols, consequently resulting in low sensitivity of detecting the ABCG5/G8

ATPase activity. Because a high concentration of bile acids is required, we have reasoned that

the high content of detergents, both in the assay solution and in the protein preparations, may

have caused either high background upon quenching the reaction in the Malachite Green-based

assay (Johnson et al., 2010) or poor organic-aqueous phase separation (Lee et al., 2016). The

measurement of ATPase activity has thus become inconsistent from one protein preparation to

another. To overcome this issue, we first optimized the ATPase assay by adopting a colorimetric

and bismuth citrate-based approach (Cariani et al., 2004), which also allows high-throughput

detection of the liberated inorganic phosphate by a microplate reader. The ATPase assay was

performed in a 65μl final reaction volume containing 2mg/ml *E. coli* or liver polar lipids or

designated phospholipids, 1.5% sodium cholate, 0.2% (4.11mM) CHS, and 2mM DTT in Buffer

A (50mM Tris/Cl pH 7.5, 100mM NaCl, 10% glycerol, 0.1% DDM). The Lipid/CHS/DTT

mixture was thoroughly sonicated and preincubated with ABCG5/G8 proteins (0.3 to 1.5 μg) for

5 minutes at room temperature. The catalytically deficient G8-G216D was used as the negative

control.

The enzymatic activity of ABCG5/G8 was initiated upon the addition of the 10X ATP cocktail

(6.5μl) and incubated at 37°C. Aliquots (8.5μl) were removed every 2 minutes and added to the

pre-chilled quencher wells to stop the reaction. The quencher solution was made of 5% SDS in

5mM HCl. Lipid mixtures were prepared at 30mg/ml (~20mM) in Buffer A containing 7%

sodium cholate. CHS stock solution (1%, w/v) was prepared in a Buffer A and 4.5% sodium

cholate. 10X Mg/ATP cocktail contains 50mM ATP, 75mM $MgCl_2$, 100mM NaN_3 in a buffer

containing 50mM Tris/Cl pH 7.5. To detect the liberated inorganic phosphate, 50μL of freshly-

made Solution II (142mM ascorbic acid, 0.42M HCl, 4.2% Solution I (10% ammonium

molybdate) was added to plate wells and left on ice for 10 minutes. Then 75μl Solution III

(88mM bismuth citrate, 120mM sodium citrate, 1M HCl) was added to plate wells and placed at

37°C for 10 minutes. The absorbance was measured at 695nm using a multi-well plate reader.

For the phosphate standards, 1M monobasic or dibasic sodium or potassium phosphate in 50mM

Tris/Cl pH 7.5 was prepared, and six standard inorganic phosphate solutions (0μM, 12.5μM,

25μM, 50μM, 100μM, 200μM) were used in every experiment. The linear range of each reaction

was used to calculate the initial rate of ATP hydrolysis. GraphPad Prism 8 was used to perform

nonlinear regression and ordinary one-way ANOVA, with a P value of ≤0.05 considered

significant from at least three independent experiments. The kinetic parameters were calculated

by nonlinear Michaelis-Menten curve fitting using GraphPad Prism 8.

Computational Methods. We studied four ABCG5/G8 protein systems including the WT,

E146Q, A540F and R543S mutants. Each MD system consists of one copy of ABCG5/G8

heterodimer, 320 1,2-Dimyristoyl-sn-glycero-3-phosphocholine (DMPC) lipid, 16 Cholesterol,

43,621 TIP3P (Jorgensen et al., 1983) water molecules, 103 Cl^- and 83 Na^+ to neutralize the MD

systems. AMBER ff14SB (Maier et al., 2015), Lipid14 (Dickson et al., 2014) and GAFF (Wang

et al., 2004) force fields were used to model proteins, DMPC lipids and Cholesterols,

respectively. The residue topology of cholesterol was prepared using the Antechamber module

(Dickson et al., 2014). MD simulation was performed to produce isothermal-isobaric ensembles

using the pmemd.cuda program in AMBER 18 (Case et al., 2018). The Particle Mesh Ewald

(PME) method (Darden et al., 1999) was used to accurately calculate the electrostatic energies

with the long-ranged correction taken into account. All bonds were constrained using the

SHAKE algorithm (Miyamoto and Kollman, 1992) in both the minimization and MD simulation

stages following a computational protocol described in our previous publication (Lee et al.,

2016). Briefly, there were three stages in a series of constant pressure and temperature MD

simulations, including the relaxation phase, the equilibrium phase, and the sampling phase. In the

relaxation phase, the simulation system was heated up progressively from 50K to 250K at steps

of 50K and 1-nanosecond MD simulation was run at each temperature. In the next equilibrium

phase, the system was equilibrated at 298K, 1 bar for 10 ns. Finally, a 100-nanosecond MD

simulation was performed at 298K, 1 bar to produce isothermal–isobaric ensemble ensembles. In

total, 1,000 snapshots were recorded from the last phase simulation for post-analysis using

"cpptray"   module   implement   in   the   AMBER   software   package.

ACKNOWLEDGMENTS

We thank William Jennings, Gloria Ihirwe, Midhet Hajira and Chloé van de Panne for technical assistance. We also thank Donna Clary and Hui Li for their technical help in utilizing the common core facilities. We are indebted to critical feedback on reviewing the manuscript from Ms. Vicky Brandt, and Drs. John Baenziger, Jean-François Couture, Gregory Graf and Xiaohui Zha. This work is partially based on the theses that were submitted to fulfill in part the requirement for the degrees of Master of Science (A.A.Z.) and Honors Bachelor of Science (A.V.).

AUTHOR CONTRIBUTIONS

B.M.X. optimized the CHS-stimulated ATPase assay for ABCG5/G8; A.A.Z. generated and validated the mutant constructs; B.M.X., A.A.Z. and A.V. purified the proteins and carried out the ATPase assays and data analysis. J.W. performed the molecular dynamics simulation, J.-Y.L. oversaw the project, and J.W. and J.-Y.L. wrote the manuscript. All authors have read and agreed to the submitted version of the manuscript.

FUNDING

This research was funded by a startup grant from the University of Ottawa, a Discovery Grant from the Natural Sciences and Engineering Research Council (RGPIN 2018-04070), and a National New Investigator Award from the Heart and Stroke Foundation of Canada to J.-Y.L. and the National Institutes of Health grants (R01-GM079383 and R21-GM097617) to J.W. B.M.X. is a recipient of the Travel Awards from the Canadian Society of Molecular Biosciences (2018) and Biophysical Society of Canada (2019).

507 **CONFLICT OF INTEREST**

508   The authors declare no conflict of interest. The funders had no role in the design of the study; in

509   the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the

510   decision to publish the results.

512   **ABBREVIATIONS**

ABC	ATP-binding cassette
ABCC7	ATP-binding cassette sub-family C member 7
ABCG5	ATP-binding cassette sub-family G member 5
ABCG8	ATP-binding cassette sub-family G member 8
ATP	Adenosine triphosphate
CBP	Calmodulin-binding peptide
CFTR	Cystic fibrosis transmembrane conductance regulator
CHS	Cholesteryl hemisuccinate
DDM	Dodecyl maltoside or n-Dodecyl β-D-maltopyranoside
DMPC	1,2-Dimyristoyl-sn-glycero-3-phosphocholine
DNA	Deoxyribonucleic acid
DTT	Dithiothreitol
EDTA	Ethylene diamine-tetraacetic acid
EGTA	Ethylene glycol-bis(-aminoethyl ether)-N,N,N',N'-tetraacetic acid
ER	Endoplasmic reticulum
ICL	Intracellular loop
LDL	Low-density lipoprotein
LOF	Loss of function
LS	Least square
MD	Molecular dynamics
MGY	Minimal glycerol yeast nitrogen base
MM	Minimal methanol
mPIB	Minimum protease inhibitor buffer
NBD	Nucleotide-binding domain
NBS	Nucleotide-binding site
Ni-NTA	Nickle-nitrilotriacetic acid
PC	Phosphatidylcholine
PCR	Polymerase chain reaction
PDB	Protein data bank
PE	Phosphatidylethanolamine
PG	Phosphatidylglycerol
RCT	Reverse cholesterol transport
RMSD	Root mean square deviation
SDS	Sodium dodecyl sulfate
TCEP	Tris-(2-carboxyethyl)-phosphine
TICE	Transintestinal cholesterol efflux
TMD	Transmembrane domain
WT	Wild type
YNB	Yeast nitrogen base
YPD	Yeast extract peptone dextrose
YPDS	Yeast extract peptone dextrose sorbitol

REFERENCES

Berge KE, Tian H, Graf GA, Yu L, Grishin N V., Schultz J, Kwiterovich P, Shan B, Barnes R,

Hobbs HH. 2000. Accumulation of dietary cholesterol in sitosterolemia caused by

mutations in adjacent ABC transporters. *Science (80-)* **290**:1771–1775.

doi:10.1126/science.290.5497.1771

Bhattacharyya AK, Connor WE, Lin DS, McMurry MM, Shulman RS. 1991. Sluggish sitosterol

turnover and hepatic failure to excrete sitosterol into bile cause expansion of body pool of

sitosterol in patients with sitosterolemia and xanthomatosis. *Arterioscler Thromb* **11**:1287–

1294. doi:10.1161/01.atv.11.5.1287

Borst P, Zelcer N, Van Helvoort A. 2000. ABC transporters in lipid transport. *Biochim Biophys*

Acta - Mol Cell Biol Lipids **1486**:128–144. doi:10.1016/S1388-1981(00)00053-6

Buch S, Schafmayer C, Völzke H, Becker C, Franke A, Von Eller-Eberstein H, Kluck C,

Bässmann I, Brosch M, Lammert F, Miquel JF, Nervi F, Wittig M, Rosskopf D, Timm B,

Höll C, Seeger M, Elsharawy A, Lu T, Egberts J, Fändrich F, Fölsch UR, Krawczak M,

Schreiber S, Nürnberg P, Tepel J, Hampe J. 2007. A genome-wide association scan

identifies the hepatic cholesterol transporter ABCG8 as a susceptibility factor for human

gallstone disease. *Nat Genet* **39**:995–999. doi:10.1038/ng2101

Cariani L, Thomas L, Brito J, Del Castillo JR. 2004. Bismuth citrate in the quantification of

inorganic phosphate and its utility in the determination of membrane-bound phosphatases.

Anal Biochem **324**:79–83. doi:10.1016/j.ab.2003.09.008

Case DA, Ben-Shalom IY, Brozell SR, Cerutti DS, Cheatham III TE, Cruzeiro VWD, Darden
TA, Duke RE, Ghoreishi D, Gilson MK. 2018. Amber 2018; 2018, University of California,
San Francisco.

Chen ZC, Shin SJ, Kuo KK, Lin K Der, Yu ML, Hsiao PJ. 2008. Significant association of
ABCG8:D19H gene polymorphism with hypercholesterolemia and insulin resistance. *J*
Hum Genet **53**:757–763. doi:10.1007/s10038-008-0310-2

Darden T, Perera L, Li L, Lee P. 1999. New tricks for modelers from the crystallography toolkit:
The particle mesh Ewald algorithm and its use in nucleic acid simulations. *Structure* **7**:R55–
R60. doi:10.1016/S0969-2126(99)80033-1

Davies DR, Hol WGJ. 2004. The power of vanadate in crystallographic investigations of
phosphoryl transfer enzymes. *FEBS Lett* **577**:315–321. doi:10.1016/j.febslet.2004.10.022

Dean M, Allikmets R. 1995. Evolution of ATP-binding cassette transporter genes. *Curr Opin*
Genet Dev **5**:779–785. doi:10.1016/0959-437X(95)80011-S

Dean M, Hamon Y, Chimini G. 2001. The human ATP-binding cassette (ABC) transporter
superfamily. *J Lipid Res* **42**:1007–1017. doi:10.1101/gr.184901

Dickson CJ, Madej BD, Skjevik ÅA, Betz RM, Teigen K, Gould IR, Walker RC. 2014. Lipid14:
The amber lipid force field. *J Chem Theory Comput* **10**:865–879. doi:10.1021/ct4010307

Feldblum ES, Arkin IT. 2014. Strength of a bifurcated H bond. *Proc Natl Acad Sci U S A*
111:4085–4090. doi:10.1073/pnas.1319827111

Ford RC, Beis K. 2019. Learning the ABCs one at a time: Structure and mechanism of ABC

transporters. *Biochem Soc Trans* **47**:23–36. doi:10.1042/BST20180147

Graf GA, Cohen JC, Hobbs HH. 2004. Missense mutations in ABCG5 and ABCG8 disrupt

heterodimerization   and   trafficking.   *J   Biol   Chem*   **279**:24881–24888.

doi:10.1074/jbc.M402634200

Graf GA, Li W-P, Gerard RD, Gelissen I, White A, Cohen JC, Hobbs HH. 2002. Coexpression

of ATP-binding cassette proteins ABCG5 and ABCG8 permits their transport to the apical

surface. *J Clin Invest* **110**:659–669. doi:10.1172/jci16000

Graf GA, Yu L, Li WP, Gerard R, Tuma PL, Cohen JC, Hobbs HH. 2003. ABCG5 and ABCG8

Are Obligate Heterodimers for Protein Trafficking and Biliary Cholesterol Excretion. *J Biol*

Chem **278**:48275–48282. doi:10.1074/jbc.M310223200

Hanson MA, Cherezov V, Griffith MT, Roth CB, Jaakola VP, Chien EYT, Velasquez J, Kuhn P,

Stevens RC. 2008. A Specific Cholesterol Binding Site Is Established by the 2.8 Å

Structure   of   the   Human   β2-Adrenergic   Receptor.   *Structure*   **16**:897–905.

doi:10.1016/j.str.2008.05.001

Hauser H, Howell K, Dawson RMC, Bowyer DE. 1980. Rabbit small intestinal brush border

membrane. Preparation and lipid composition. *Biochim Biophys Acta - Biomembr* **602**:567–

577. doi:10.1016/0005-2736(80)90335-1

Hirayama H, Kimura Y, Kioka N, Matsuo M, Ueda K. 2013. ATPase activity of human ABCG1

is   stimulated   by   cholesterol   and   sphingomyelin.   *J   Lipid   Res*   **54**:496–502.

doi:10.1194/jlr.M033209

Hwang JU, Song WY, Hong D, Ko D, Yamaoka Y, Jang S, Yim S, Lee E, Khare D, Kim K,

Palmgren M, Yoon HS, Martinoia E, Lee Y. 2016. Plant ABC Transporters Enable Many

Unique Aspects of a Terrestrial Plant's Lifestyle. *Mol Plant* **9**:338–355.

doi:10.1016/j.molp.2016.02.003

Jakulj L, van Dijk TH, de Boer JF, Kootte RS, Schonewille M, Paalvast Y, Boer T, Bloks VW,

Boverhof R, Nieuwdorp M, Beuers UHW, Stroes ESG, Groen AK. 2016. Transintestinal

Cholesterol Transport Is Active in Mice and Humans and Controls Ezetimibe-Induced Fecal

Neutral Sterol Excretion. *Cell Metab* **24**:783–794. doi:10.1016/j.cmet.2016.10.001

Johnson BJH, Lee JY, Pickert A, Urbatsch IL. 2010. Bile acids stimulate atp hydrolysis in the

purified cholesterol transporter ABCG5/G8. *Biochemistry* **49**:3403–3411.

doi:10.1021/bi902064g

Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML. 1983. Comparison of

simple potential functions for simulating liquid water. *J Chem Phys* **79**:926–935.

doi:10.1063/1.445869

Kajinami K, Brousseau ME, Ordovas JM, Schaefer EJ. 2004. Interactions between common

genetic polymorphisms in ABCG5/G8 and CYP7A1 on LDL cholesterol-lowering response

to atorvastatin. *Atherosclerosis* **175**:287–293. doi:10.1016/j.atherosclerosis.2004.03.015

Kizhakkedath P, John A, Al-Sawafi BK, Al-Gazali L, Ali BR. 2019. Endoplasmic reticulum

quality control of LDLR variants associated with familial hypercholesterolemia. *FEBS*

Open Bio **9**:1994–2005. doi:10.1002/2211-5463.12740

Klett EL, Lu K, Kosters A, Vink E, Lee MH, Altenburg M, Shefer S, Batta AK, Yu H, Chen J,

Klein R, Looije N, Oude-Elferink R, Groen AK, Maeda N, Salen G, Patel SB. 2004. A
mouse model of sitosterolemia: Absence of Abcg8/sterolin-2 results in failure to secrete
biliary cholesterol. *BMC Med* **2**:5. doi:10.1186/1741-7015-2-5

Kuo KK, Shin SJ, Chen ZC, Yang YHC, Yang JF, Hsiao PJ. 2008. Significant association of
ABCG5 604Q and ABCG8 D19H polymorphisms with gallstone disease. *Br J Surg*
95:1005–1011. doi:10.1002/bjs.6178

Kurumi Y, Adachi Y, Itoh T, Kobayashi H, Nanno T, Yamamoto T. 1991. Novel high-
performance liquid chromatography for determination of membrane phospholipid
composition of rat hepatocytes. *Gastroenterol Jpn* **26**:628–632. doi:10.1007/BF02781680

Lee AG. 2004. How lipids affect the activities of integral membrane proteins. *Biochim Biophys*
Acta - Biomembr **1666**:62–87. doi:10.1016/j.bbamem.2004.05.012

Lee JY, Kinch LN, Borek DM, Wang Jin, Wang Junmei, Urbatsch IL, Xie XS, Grishin N V.,
Cohen JC, Otwinowski Z, Hobbs HH, Rosenbaum DM. 2016. Crystal structure of the
human sterol transporter ABCG5/ABCG8. *Nature* **533**:561–564. doi:10.1038/nature17666

Li G, Gu HM, Zhang DW. 2013. ATP-binding cassette transporters and cholesterol
translocation. *IUBMB Life* **65**:505–512. doi:10.1002/iub.1165

Linton KJ, Higgins CF. 1998. The Escherichia coli ATP-binding cassette (ABC) proteins. *Mol*
Microbiol **28**:5–13. doi:10.1046/j.1365-2958.1998.00764.x

Lu K, Lee MH, Hazard S, Brooks-Wilson A, Hidaka H, Kojima H, Ose L, Stalenhoef AFH,
Mietinnen T, Bjorkhem I, Bruckert E, Pandya A, Brewer HB, Salen G, Dean M, Srivastava

A, Patel SB. 2001. Two genes that map to the STSL locus cause sitosterolemia: Genomic

structure and spectrum of mutations involving sterolin-1 and sterolin-2, encoded by ABCG5

and ABCG8, respectively. *Am J Hum Genet* **69**:278–290. doi:10.1086/321294

Lukacs GL, Mohamed A, Kartner N, Chang XB, Riordan JR, Grinstein S. 1994. Conformational

maturation of CFTR but not its mutant counterpart (ΔF508) occurs in the endoplasmic

reticulum and requires ATP. *EMBO J* **13**:6076–6086. doi:10.1002/j.1460-

2075.1994.tb06954.x

Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE, Simmerling C. 2015. ff14SB:

Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *J*

Chem Theory Comput **11**:3696–3713. doi:10.1021/acs.jctc.5b00255

Manolaridis I, Jackson SM, Taylor NMI, Kowal J, Stahlberg H, Locher KP. 2018. Cryo-EM

structures of a human ABCG2 mutant trapped in ATP-bound and substrate-bound states.

Nature **563**:426–430. doi:10.1038/s41586-018-0680-3

Miyamoto S, Kollman PA. 1992. Settle: An analytical version of the SHAKE and RATTLE

algorithm for rigid water models. *J Comput Chem* **13**:952–962. doi:10.1002/jcc.540130805

Neumann J, Rose-Sperling D, Hellmich UA. 2017. Diverse relations between ABC transporters

and lipids: An overview. *Biochim Biophys Acta - Biomembr* **1859**:605–618.

doi:10.1016/j.bbamem.2016.09.023

Patel SB, Graf GA, Temel RE. 2018. ABCG5 and ABCG8: more than a defense against

xenosterols. *J Lipid Res* **59**:1103–1113. doi:10.1194/jlr.R084244

Plösch T, Bloks VW, Terasawa Y, Berdy S, Siegler K, Van Der Sluijs F, Kema IP, Groen AK,

Shan B, Kuipers F, Schwartz M. 2004. Sitosterolemia in ABC-Transporter G5-Deficient

Mice Is Aggravated on Activation of the Liver-X Receptor. *Gastroenterology* **126**:290–300.

doi:10.1053/j.gastro.2003.10.074

Takahashi K, Kimura Y, Kioka N, Matsuo M, Ueda K. 2006. Purification and ATPase activity of

human ABCA1. *J Biol Chem* **281**:10760–10768. doi:10.1074/jbc.M513783200

Wang J, Grishin N, Kinch L, Cohen JC, Hobbs HH, Xie XS. 2011. Sequences in the

nonconsensus nucleotide-binding domain of ABCG5/ABCG8 required for sterol transport. *J*

Biol Chem **286**:7308–7314. doi:10.1074/jbc.M110.210880

Wang J, Wolf RM, Caldwell JW, Kollman PA, Case DA. 2004. Development and testing of a

general Amber force field. *J Comput Chem* **25**:1157–1174. doi:10.1002/jcc.20035

Wang Z, Stalcup LD, Harvey BJ, Weber J, Chloupkova M, Dumont ME, Dean M, Urbatsch IL.

2006. Purification and ATP hydrolysis of the putative cholesterol transporters ABCG5 and

ABCG8. *Biochemistry* **45**:9929–9939. doi:10.1021/bi0608055

Xavier BM, Jennings WJ, Zein AA, Wang J, Lee JY. 2019. Structural snapshot of the

cholesterol-transport ATP-binding cassette proteins. *Biochem Cell Biol* **97**:224–233.

doi:10.1139/bcb-2018-0151

Yu L, Hammer RE, Li-Hawkins J, Von Bergmann K, Lutjohann D, Cohen JC, Hobbs HH. 2002.

Disruption of Abcg5 and Abcg8 in mice reveals their crucial role in biliary cholesterol

secretion. *Proc Natl Acad Sci U S A* **99**:16237–16242. doi:10.1073/pnas.252582399

655 Zein AA, Kaur R, Hussein TOK, Graf GA, Lee JY. 2019. ABCG5/G8: A structural view to

656   pathophysiology of the hepatobiliary cholesterol secretion. *Biochem Soc Trans* **47**:1259–

657   1268. doi:10.1042/BST20190130

658 Zhang DW, Graf GA, Gerard RD, Cohen JC, Hobbs HH. 2006. Functional asymmetry of

659   nucleotide-binding domains in ABCG5 and ABCG8. *J Biol Chem* **281**:4507–4516.

660   doi:10.1074/jbc.M512277200

FIGURES AND TABLES.

Figure 1. Disease-causing mutations and SNPs in ABCG5/G8. A, Localization of ABCG5/G8

residues carrying missense mutations. The positions of disorder-related polymorphisms or

mutations are highlighted in black spheres on the structures of ABCG5 (PDB ID: 5D07, chain C)

and ABCG8 (PDB ID 5D07, chain D). Structural motifs are indicated in dashed ovals: triple-

helical bundle (black), TMD polar relay (yellow), and extracellular domain with re-entry helices

(green). **B,** Microenvironment of G5-E146, G5-540, and G8-R543. *(Middle)* The transmembrane

domains (white) and the triple helical bundle (grey) are plotted in tube-styled cartoon

presentation, showing the α-carbons (spheres) of G5-E146 (orange), G8-R543 (blue), and G5-

A540 (black). The polar relays are plotted in dotted yellow spheres. *(Top-left)* Slapped top view

shows G5-A540 situated more than 10Å away from the polar relay of either subunit (red dot-

ended lines). *(Top-right)* Near G5-A540 shows a cholesterol-shaped electron density (mesh) in

the crystal structure of ABCG5/G8. Fo-Fc difference electron density map was contoured at

3.0σ. *(Bottom-left)* At the triple helical bundle of ABCG5, E146 interacts with R377 through

their side-chain termini in a distance of hydrogen bonding, 3.5Å (black dashed line). *(Bottom-*

right) In ABCG8 polar relay, R543 interacts E503 through their side-chain termini in a distance

of hydrogen bonding, 3.1Å (black dashed line).

Figure 2. A, Chemical structures of cholesterol, cholesteryl hemisuccinate (CHS) and cholic

acid (cholate). Source: PubChem. **B,** Schematic illustration of sterol-coupled ATPase activity of

ABCG5/G8. DDM-purified ABCG5/G8 (light/dark grey surface) is preincubated with

phospholipids and cholate. Addition of CHS (four-ringed steroid structure) stimulates hydrolysis

of ATP to ADP and inorganic phosphate (Pi) in the presence of the divalent magnesium ions

(Mg^{2+}). Using the colorimetric and bismuth citrate-based assay, the liberated Pi is then captured

by ammonium molybdate in the presence ascorbic acid. The color is developed upon mixing

with bismuth citrate and sodium citrate, and the absorbance was measured at 695 nm. See details

in Materials and Methods.

Figure 3. ATPase activity of ABCG5/G8. The ATP hydrolysis was used as a measure of

ABCG5/G8 ATPase activity at 37°C in a condition with 5mM ATP and 4.1mM CHS. The

protocol is entailed in Materials and Methods. **A,** Data points are presented as the means ±

standard deviations from 4-8 independent time-course experiments, where not visible, the error

bars are covered by the plot symbols. A linear regression, plotted from the first 12 minutes, is

used to calculate the specific activities. **B,** Bar graphs show the specific activities of ATP

hydrolysis by WT in the presence and absence of CHS and the catalytically deficient mutant G8-

G216D in the presence of CHS. The specific activity of WT in the absence of CHS is regarded as

the basal ABCG5/G8 ATPase activity. **C,** Bar graphs represent percentage inhibition of

ABCG5/G8 ATPase activity by 0.015mM orthovanadate, a P-value of <0.0001 obtained using

ordinary one-way ANOVA (Prism 8).

Figure 4. Lipid dependence of ABCG5/G8 ATPase activity. A, Purified ABCG5/G8 was

assayed in the presence of either *E. coli* or bovine liver polar lipids, and the specific activities of

ATP hydrolysis were obtained by the ATP concentration-dependent experiments (0-5mM ATP).

Both curves are fitted to the Michaelis-Menten equation (Prism 8), and the means of at least

three replicates along with standard deviations are plotted here. The kinetic parameters are listed

in Table 1. **B,** In a condition of 5mM ATP and 4.1mM CHS, ATP hydrolysis of purified

ABCG5/G8 was assayed in the presence of egg PE, soy PC or egg PG, a P-value of 0.0006 and

0.0003, respectively, obtained using ordinary one-way ANOVA (Prism 8).

Figure 5. ATP-Dependence of ABCG5/G8 ATPase Activity. Purified proteins were assayed in the presence of *E. coli* polar lipids, and the specific activities of ATP hydrolysis were obtained by the ATP concentration-dependent experiments (0-5mM ATP). The curves are fitted to the Michaelis-Menten equation (Prism 8), and the means of at least three replicates along with standard deviations are plotted here. The kinetic parameters are listed in Table 1.

Figure 6. CHS-Dependence of ABCG5/G8 ATPase Activity. Purified proteins were assayed in the presence of *E. coli* polar lipids, and the specific activities of ATP hydrolysis were obtained by the CHS concentration-dependent experiments (0-4.1mM CHS). The curves are fitted to the Michaelis-Menten equation (Prism 8), and the means of at least three replicates along with standard deviations are plotted here. The kinetic parameters are listed in Table 2.

729
730

Figure 7. Fluctuation of Root-mean-square deviations (RMSD) along MD simulation time course. RMSD were calculated using the main chain atoms of all residues (black lines) or secondary structures only (red lines). A: wild type, B: E146Q mutant in ABCG5, C: A540F mutant in ABCG5, and D: R543S mutant in ABCG8. G5G8: ABCG5/G8; SS: secondary structure.

Figure 8. Representative structures of the WT ABCG5/G8 and its three missense mutants.

The representative structures (shown as green cartoons, magenta spheres and magenta lines) were aligned to the crystal structure (grey cartoons, blue spheres and blue lines). The three mutation residues, E146Q, A540F and R543S, are shown as spheres. The hypothetical surrounding residues of ATP are shown as lines. A and E: wild type, B and F: E146A, C and F: A540F, D and G: R543S. G5: ABCG5; G8: ABCG8.

Figure 9. Proposed mechanism of sterol-coupled ATP catalysis by ABCG5/G8. *(Mode 1)* A sequential pathway is derived from experiments on the disease mutants, G5-E146Q and G8-R543S. ABCG5/G8 first recruits ATP and Mg^{2+} ions, likely causing a conformational change of the NBD for ATP binding. CHS/sterol then binds the transporter and triggers ATP hydrolysis that may result in its dissociation. *(Mode 2)* A concerted pathway is derived from experiments on the putative sterol-binding mutant, G5-A540F. ABCG5/G8 simultaneously recruits CHS, ATP, and Mg^{2+} ions, induces a transient conformational change of the NBD, and activates ATP hydrolysis and CHS/sterol dissociation from the transporter. G5: ABCG5; G8: ABCG8; E: ABCG5/G8 heterodimer; Pi: inorganic phosphate.

756 **Table 1.** Dependence of ABCG5/G8 ATPase Activity on ATP.

	V_{max}[a] (nmol/min/mg)	K_M (ATP) (mM)	k_{cat}[b] (s^{-1})	k_{cat}/K_M $(M^{-1}s^{-1})$	$\Delta\Delta G_{MUT}$[c] (kJ/mol)
WT (liver polar lipids)	192.8 ± 17.9	0.93	0.48	0.52×10^3	-
WT (*E. coli* polar lipids)	677.1 ± 25.6	0.60	1.69	2.8×10^3	-
G5-E146Q[d]	167.1 ± 0.05	0.51	0.41	0.82×10^3	11.7
G8-R543S[d]	150.7 ± 3.7	0.42	0.38	0.90×10^3	12.3
G5-A540F[d]	101.2 ± 4.2	0.58	0.25	0.43×10^3	15.8

a, Standard errors were calculated from the fits shown in Figures 3A & 5D using GraphPad

Prism 8.

b, Turnover rates, k_{cat}, were calculated using the following formula. $V_{max} = k_{cat} \times [E]$, [E]: protein

concentration of ABCG5/G8 (363.1 nM).

c, Differential Gibbs free energy was calculated according to the formula: $\Delta\Delta G_{MUT} = -$

$RTln(k_{MUT}/k_{WT})$, k_{MUT}: k_{cat} of mutants, k_{WT}: k_{cat} of WT, R = 8.314 $Jmol^{-1}K^{-1}$ (R: Gas Constant, K:

Kelvin), T = 310.15 K (37°C).

d, Mutants were all assayed in the presence of *E. coli* polar lipids.

Table 2. Dependence of ABCG5/G8 ATPase Activity on Cholesteryl Hemisuccinate.

	V_{max}[a] (nmol/min/mg)	K_M (CHS) (mM)	k_{cat}[b] (s^{-1})	k_{cat}/K_M $(M^{-1}s^{-1})$	$\Delta\Delta G_{MUT}$[c] (kJ/mol)
WT[d]	702.9 ± 50.7	0.8	1.74	2.2×10^3	-
G5-E146Q[d]	210.0 ± 33.2	1.13	0.52	0.46×10^3	10.0
G8-R543S[d]	237.1 ± 33.4	2.38	0.59	0.25×10^3	9.0
G5-A540F[d]	99.8 ± 11.4	0.70	0.25	0.36×10^3	16.1

a, Standard errors were calculated from the fits shown in Figures 3A & 5D using GraphPad

Prism 8.

b, Turnover rates, k_{cat}, were calculated using the following formula. $V_{max} = k_{cat} \times [E]$, [E]: protein

concentration of ABCG5/G8 (363.1 nM).

c, Differential Gibbs free energy was calculated according to the formula: $\Delta\Delta G_{MUT} = -$

$RTln(k_{MUT}/k_{WT})$, k_{MUT}: k_{cat} of mutants, k_{WT}: k_{cat} of WT, R = 8.314 Jmol^{-1}K^{-1} (R: Gas Constant, K:

Kelvin), T = 310.15 K (37°C).

d, Both WT and mutants were assayed in the presence of *E. coli* polar lipids.

SUPPLEMENTARY FIGURES

Supplementary Figure 1. Expression and purification of ABCG5/G8 missense mutants. A, Four or six yeast colonies were selected for expression test. Crude microsomal membranes (either WT or mutants), containing 20-30μg total proteins, were resolved by SDS–PAGE. Protein expression was analyzed by Western blotting using a monoclonal anti-RGSH$_4$ antibody to detect ABCG5 and a polyclonal anti-human ABCG8 antibody to detect ABCG8. The clones expressing the highest level for both subunits were selected for protein purification. **B,** Gel-filtration purified mutants were resolved on a 10% SDS-PAGE gel and stained by Coomassie Blue (shown here in greyscale). Lanes 1: molecular weight marker, 2: G8-G216D, 3: G5-E146Q, 4: G8-R543S, 5: G5-A540F, and 6: WT.

Supplementary Figure 2. Fluctuation of Root-mean-square deviations (RMSD) along MD
simulation time course. RMSD were calculated for the hypothetical residues surrounding ATP
under two scenarios. In the first scenario, least-square (LS) fittings were performed for the main
chain atoms of the hypothetic residues which including Residues 48-62, 68, 86-94, 134, 218,
250, 251, 267, 268, 302 of G5 and 242-244 of G8. In the second scenario, the RMSD were
calculated directly for the hypothetical residues without LS Fitting after the MD snapshots were
aligned to the crystal structure using the secondary structures of ABCG5/G8. The fluctuations of
the first and second scenarios were illustrated using black and red curves.