TITLE: North American fireflies host low bacterial diversity.

AUTHORS: Emily A. Green [a], Scott R. Smedley [b,c], Jonathan L. Klassen [a,d]

[a] Department of Molecular and Cell Biology, University of Connecticut, Storrs, Connecticut,

USA

[b] Department of Biology, Trinity College, Hartford, Connecticut, USA

[c] Deceased

[d] To whom correspondence should be addressed: 91 N Eagleville Rd, Unit 3125, Storrs, CT,

06269-3125. jonathan.klassen@uconn.edu

Emily A Green: https://orcid.org/0000-0002-0824-6369

Jonathan Klassen PhD: https://orcid.org/0000-0003-1745-8838

Key Words: Fireflies, microbiome, microbial ecology, insects, low-complexity communities,

Mollicutes

ACKNOWLEDGEMENTS

We would like to thank Erin L. Mostoller and Dr. Craig W Schneider both from Trinity College,

Hartford, CT, and Dr. Steven Deyrup from Siena College, Loudonville, NY for their assistance

with obtaining the firefly samples. We would also like to thank the members of the Klassen Lab

for their thoughtful feedback on this manuscript before submission, and the UConn Microbial

Analysis, Resources, and Services facility for microbiome sequencing.

ABSTRACT

Although there are numerous studies of firefly mating flashes, lantern bioluminescence, and anti-predation lucibufagin metabolites, almost nothing is known about their microbiome. We therefore used 16S rRNA community amplicon sequencing to characterize the gut and body microbiomes of four North American firefly taxa: *Ellychnia corrusca*, the *Photuris versicolor* species complex, *Pyractomena borealis*, and *Pyropyga decipiens*. These firefly microbiomes all have very low species diversity, often dominated by a single species, and each firefly type has a characteristic microbiome. Although the microbiomes of male and female fireflies did not differ from each other, *Ph. versicolor* gut and body microbiomes did, with their gut microbiomes being enriched in *Pseudomonas* and *Acinetobacter*. *Ellychnia corrusca* egg and adult microbiomes were unique except for a single egg microbiome that shared a community type with *E. corrusca* adults, which could suggest microbial transmission from mother to offspring. Mollicutes that had been previously isolated from fireflies were common in our firefly microbiomes. These results set the stage for further research concerning the function and transmission of these bacterial symbionts.

INTRODUCTION:

Beetles (Order: Coleoptera) are one of the most diverse insect groups, containing nearly 400,000 species [1]. The beetles in the family Lampyridae are commonly known by many names: fireflies, lightning bugs, glow worms, lamp-lighters, and night-travelers, all describing the bioluminescent lantern that they use to signal to their mates [2]. Fireflies are found on every continent except Antarctica, living in woods, plains, and marshes. North America is home to over 125 firefly species, including *Ellychnia corrusca*, the *Photuris versicolor* species complex,

Pyractomena borealis and *Pyropyga decipiens* [2]. These 4 types of fireflies occur sympatrically,

ranging from the eastern coast of Canada, south to Florida, and west to the Great Plains [2, 3].

All four of these fireflies are carnivorous or omnivorous as larvae, attacking snails as their prey,

and/or eating nectar and plant sap [4]. *Ellychnia corrusca*, *Pyractomena borealis* and *Pyropyga*

decipiens are all closely related to each other, and both *E. corrusca* and *Pyropyga decipiens* are

diurnal with no lantern [2]. *Ellychnia corrusca*, *Pyractomena borealis*, and many other North

American fireflies use lucibufagins as a chemical defense. Lucibufagins are cardioactive C-24

steroidal pyrenes and a subclass of the bufadienolides [5–8] that create a disagreeable taste to

predators and thereby provide fireflies with protection from bats, birds, and spiders [5].

However, the biosynthetic origin of lucibufagins is not known. At least five genera of fireflies,

including the groups in this study, ingest the nectar of the common milkweed plant (*Asclepias*

syriaca L.). Although eating is not required for adult firefly survival, cardenolides (a type of

toxic steroid) found on milkweed plants are ingested by adult fireflies and may be used to

produce lucibufagins [9].

In this study, we focused especially on two fireflies with unique lifestyles: *E. corrusca*

and the *Photuris versicolor* species complex. *Ellychnia corrusca* is nicknamed the winter firefly

because it has a winter-spring activity cycle, in contrast to the late spring-summer activity cycles

of most other fireflies [2, 10]. *Ellychnia corrusca* larvae in New England (North America)

extend their larval stage across two years instead of one to ensure that they ingest ample calories,

emerging in the late fall of their second year as adults and sexually maturing during the winter to

start mating, which lasts until late spring [11]. Normally, adult fireflies do not need to feed

because they only survive as adults for 2 to 4 weeks [2, 4]. However, *E. corrusca* firefly adults

live for ~9 months, and in Massachusetts USA they ingest interstitial fluid and sap from maple

trees to help them survive during the cold winters [11]. The *Photuris versicolor* species complex

(hereafter: *Ph. versicolor*), contains > 17 firefly species that are differentiated by their male

courtship lantern flashes but that cannot be identified from their morphology with confidence.

The *Ph. versicolor* adult life cycle occurs during the summer months (June-August), but unlike

other fireflies, *Ph. versicolor* females mimic the mating light display of female *Photinus* fireflies

to attract *Photinus* males, earning *Ph. versicolor* the nickname "femme fatale" [12]. Once a male

Photinus firefly is near, the *Photuris versicolor* female will attack and kill the male, gaining

nutrients and lucibufagins that she uses to protect herself, her eggs, and her pupae, which do not

themselves produce lucibufagins [5]. Many *E. corrusca* populations complete their life cycles

before *Ph. versicolor* becomes seasonally abundant, especially where temperatures are colder.

However, *Ph. versicolor* predation of *E. corrusca* can occur, particularly in the Southern USA.

Ph. versicolor will also attack and eat *E. corrusca* to gain protective lucibufagins in the lab [2,

8], which could give insight into the selective pressures that caused *E. corrusca* to mate in the

winter instead of in the summer.

Beetles feed on many different substrates, following herbivorous [13], omnivorous [14],

xylophagous [15], detritivorous [16], and carnivorous diets [17]. Beetles that belong to similar

taxonomic families but that have different diets have distinct microbial communities [17–19].

Diets lacking in nutrients often cause insects to rely on nutritional symbionts to provide vital

nutrients [20], as in dung [21], carabid [14], and carrion beetles [22]. Although there is a vast

amount of research studying beetle microbiomes, only a handful of studies consider firefly-

associated microbes. In those studies, several Mollicute bacteria were isolated from the fireflies

E. corrusca and *Ph. versicolor*, but this does not give insight into the composition of their

microbiome beyond these strains [23–30]. To fill this gap, we used 16S rRNA community

amplicon sequencing to survey the microbiomes of *E. corrusca, Photuris versicolor*,

Pyractomena borealis, and *Pyropyga decipiens* fireflies. Our results show that fireflies have

simple, species-specific microbiomes, and generate hypotheses about how diet and seasonality

may drive firefly microbiome structure and function.

METHODS

Sample collection

Live *E. corrusca* and *Photuris versicolor* fireflies were collected by S. Smedley during

the winter of 2016-17 and spring/summer of 2017 from residential and camping/forest areas

within Vernon, Bolton, and Andover, Connecticut, U.S.A. *Pyractomena borealis* larvae and

adults were raised in the laboratory by S. Smedley. No wild *Pyractomena borealis* samples were

included in this study. After collection, *E. corrusca*, *Ph. versicolor*, and *Pyractomena borealis*

fireflies were stored at -80°C in air-filled vials. *Pyropyga decipiens* fireflies were collected by

Lynn Faust in summer 2016 from Ohio and Tennessee and stored in 95% ethanol. Collection

dates and sample locations are listed in Supplemental File 1.

Sample preparation & DNA extraction

All fireflies were surface-sterilized using 3 rounds of a 10 second submersion in 70%

ethanol, followed by a 10 second submersion in phosphate-buffered saline (PBS) [31]. Surface-

sterilized fireflies were dissected into two separate tissues: a gut sample and the remaining

carcass "body" sample. DNA from the tissue dissections were extracted using a bead beating and

chloroform-isopropanol protocol as described in [32], except using 0.7 g of 1 mm and 0.3 g of

0.1 mm silica/zirconium beads and 5 cycles of bead-beating & chilling on ice. Negative controls

containing only the DNA extraction reagents were processed alongside each batch of tissue

samples. The DNA concentration of each extract and negative control was determined using the

Qubit dsDNA high-sensitivity assay protocol and a Qubit 3.0 fluorimeter (Invitrogen, Carlsbad

California).

16S V4 rRNA PCR screen & community amplicon sequencing

DNA samples were PCR amplified using primers 515F and 806R (targeting the bacterial

16S rRNA gene V4 region) to determine the presence of bacterial DNA [33]. Ten nanograms of

template DNA was added to 5 µl Green GoTaq Reaction Mix Buffer (Promega, Madison,

Wisconsin, USA), 1.25 units of GoTaq DNA Polymerase (Promega, Madison, Wisconsin, USA),

10 µmol of each primer, and 300 ng/µl BSA (New England BioLabs Inc. Ipswitch

Massachusetts), to which nuclease free H_2O was added to a volume of 25 µl. Thermocycling

conditions (BioRad, Hercules, California) were: 3 min at 95°C, 30 cycles of 30 sec at 95°C, 30

sec at 50°C, and 60 sec at 72°C, followed by a 5 min cycle at 72°C and then an indefinite hold at

4°C. Gel electrophoresis was used to confirm the expected band size of 300–350 bp.

All samples that had a gel band of the expected size were prepared for community

amplicon sequencing of the 16S rRNA gene V4 region using an Illumina MiSeq at the

University of Connecticut Microbial Analysis, Resources and Services (MARS) facility.

Approximately 30 ng of DNA from each sample was added to a 96-well plate containing 10

µmol each of the forward and reverse Illumina-barcoded versions of primers 515F and 806R, 5

µl AccuPrime buffer (Invitrogen, Carlsbad, California), 50 mM $MgSO_4$ (Invitrogen, Carlsbad,

California), 300 ng/µl BSA (New England BioLabs Inc. Ipswitch, Massachusetts), a 1µmol

spike-in of both non-barcoded primers 515F and 806R, and 1 unit AccuPrime polymerase

(Invitrogen, Carlsbad, California), to which nuclease-free H_2O was added to a volume of 50 µl.

Reaction mixes were separated into triplicate reactions (each with a volume of 16.7 µl) in a 384

well plate using an epMotion 5075 liquid handling robot (Eppendorf, Hamburg, Germany). The

resulting 384 well plate was then transferred to a thermocycler (Eppendorf, Hamburg, Germany),

which used the following conditions: 2 min at 95°C, 30 cycles of 15 sec at 95°C, 60 sec at 55°C,

and 60 sec at 68°C, followed by a final extension for 5 min at 68°C and then an indefinite hold at

4°C. After PCR, triplicate reactions were re-pooled using the epMotion and DNA concentrations

were quantified using a QIAxcel Advanced capillary electrophoresis system (QIAgen, Hilden,

Germany). Samples that had concentrations > 0.5 ng/µl were pooled using equal weights of

DNA to create the final sequencing libraries. Libraries were then bead-cleaned using Mag-Bind

RXNPure plus beads (OMEGA, Norcross, Georgia) in a 1:0.8 ratio of sequencing library to bead

volume. Cleaned library pools were adjusted to a concentration of 1.1 ng/µl ± 0.1 ng/µl, which

was confirmed using the Qubit dsDNA high-sensitivity assay on a Qubit 3.0 fluorimeter

(Invitrogen, Carlsbad, California). Microbial community sequencing on an Illumina MiSeq

(Illumina, San Diego, California) was completed in 2 batches, the first composed of 52 samples

and the second composed of 195 samples.

Post-sequencing & bioinformatic analyses:

Our DNA sequencing produced 245 16S rRNA community amplicon sequencing

datasets, including 22 negative controls. Reads were analyzed using R v3.5.3 [34] and the dada2

v1.11.1 [35] pipeline for amplicon sequence variants (ASVs)

(https://benjjneb.github.io/dada2/tutorial.html, accessed: November 11, 2017). Read counts

ranged from 2 to 888,868 (Suppl. File S1). Metadata files for the samples were imported into

phyloseq v1.26.1 [36], creating a phyloseq R object that was used for subsequent analyses. Reads

that were not classified as belonging to the kingdom Bacteria using the SILVA database v128

were removed [37, 38]. ASVs that matched to mitochondria were then removed separately,

because SILVA included them in the kingdom Bacteria. Samples were screened for

contamination using the decontam v1.2.1 [39] prevalence protocol with a default threshold value

of 0.1. No reads were flagged as contamination, and 17 samples with 0 reads were removed,

resulting in 1,675 unique ASVs (Suppl. File S1). Negative control samples were not considered

further. All samples in the dataset were then rarefied to 10,000 reads and read counts were

converted to relative abundances. The final phyloseq object contained 133 samples from 20

Photuris versicolor adults, 4 *Pyropyga decipiens* adults, 7 *Pyractomena borealis* larvae, 1

Pyractomena borealis adult, 95 *Ellychnia corrusca* adults, and 6 *E. corrusca* eggs. Despite

amplification during the initial PCR screen, of the initial 223 insect samples, 74 samples were

removed due to low read counts (53 *E. corrusca* adult samples, 3 *E. corrusca* egg samples, 9

Photuris versicolor samples, 2 *Pyropyga decipiens* samples, 4 *Pyractomena borealis* adult

samples & 3 *Pyractomena borealis* larvae samples). These low read counts may have been due

to: 1) high amounts of host DNA acting as a PCR inhibitor; 2) there being a minimal firefly

microbiome, leading to limited template concentrations; or 3) other technical issues such as

inefficient PCR amplification using primers that contained the Illumina barcodes compared to

our initial PCR screen using non-barcoded primers.

Alpha diversity was measured using the phyloseq 'plot_richness' command, and Beta

diversity was measured using weighted and unweighted unifrac distance metrics. Weighted and

unweighted unifrac distances were calculated, ordinated, and viewed using the 'distance',

'ordinate', and 'plot_ordinate' phyloseq commands, respectively. PERMANOVA statistical tests

were calculated using vegan v 2.5-4 [40]. Although weighted unifrac (WUF) and unweighted

unifrac (UUF) distances were used for each test, to keep the text concise only one test is listed in

the text and the complimentary values are presented in Supplemental Tables S1–S4.

Differences in the relative abundances of taxa in firefly microbiomes were compared

using DESeq2 [41], using the Parametric fitType, Wald tests, and an adjusted alpha value of

0.01. A heatmap was constructed for the 26 most abundant genera in the dataset using the gplots

v3.03 (https://www.rdocumentation.org/packages/gplots) command 'heatmap.2' and using the

Euclidean and ward.D distance metrics. To avoid redundancy in the heatmap, ASVs were

grouped by genus name using the phyloseq command 'tax_glom'. Covariance between the top

26 taxa were calculated using SpiecEasi v1.0.7 [42], and the SpiecEasi output file was exported

into Cytoscape v3.7.2 [43] using igraph v1.2.4.2 [44] for visualization.

The most abundant Mollicute ASVs from our dataset included four *Mesoplasma* ASVs,

two *Spiroplasma* ASVs, and one *Entomoplasma* ASV. A phylogenetic tree was constructed to

show the relationships between the 16S rRNA sequences of Mollicutes from this study and those

that had been previously isolated from fireflies. Reference Mollicute sequences from taxa

belonging to the same genera as our firefly ASVs were selected from the SILVA database,

especially those Mollicutes that had been isolated from fireflies and other beetles. Additional

Mollicute reference sequences for fireflies that were not represented in the SILVA database were

downloaded from NCBI. Sequences were aligned using MUSCLE v3.8.31 [45] and trimmed to

the same length. The phylogenetic tree, rooted by the 16S rRNA genes from *Bacillus subtilis* and

Mycoplasma haemominutum, was calculated using a GTRGAMMAI substitution model and 500

bootstrap replicates in RAxML v8.2.11 [46]. The topology of the tree created using these

selected sequences agreed with that constructed using all Mollicute sequences in SILVA,

ensuring that taxon selection did not bias our phylogenetic analysis.

The commands used for all analyses is attached as Suppl. File S2. All data are available

on NCBI under BioProject PRJNA563849. Raw sequencing reads are deposited in SRA under

BioSample numbers SAMN14678004 – SAMN14678257.

RESULTS

Firefly microbiomes are typically dominated by single taxa

We characterized the bacterial communities in 133 firefly gut and body dissections

(Suppl. File S1) using community amplicon sequencing of the 16S rRNA gene. All firefly

microbiomes had low α-diversity (Suppl. Fig. S1). Mean Shannon diversity scores for the firefly

microbiomes ranged from 0.70 to 2.52, and alpha diversities of *Photuris versicolor* and *E.*

corrusca microbiomes differed from each other, as did those from *Pyropyga decipiens* and *E.*

corrusca (Dunn test, $p = 0.001$ and $p = 0.003$ respectively). A heatmap of the top 26 bacteria

genera found in the microbiomes reflects these low levels of α-diversity, with most firefly

microbiomes dominated by a single taxon but with minute amounts of other taxa also present

(Fig. 1). Unsupervised clustering of these data grouped *E. corrusca* samples tightly together at

the left side of the heat map, while the samples from other species clustered together on the right.

Samples from different tissue dissections and sexes did not cluster together for any species of

firefly.

Using these top 26 bacteria genera, we identified taxa whose relative abundances were

correlated with each other to infer potential interactions between them (Suppl. Fig. S2). Of the 26

bacteria genera, the relative abundances of 14 genera did not correlate with those of another

genus in our dataset, and the relative abundances of 11 genera were positively correlated with

those of another genus. The relative abundances of *Ralstonia* and *Cupriavidus*, found in the

Pyropyga decipiens samples, were strongly and positively correlated with each other, as were the

relative abundances of *Tanticharoenia* and *Gluconobacter*, found in *E. corrusca* eggs, and the

relative abundances of *Gordonia* and *Tsukamurella*, found in *Photuris versicolor*. Only the

relative abundances of *Mesoplasma* and *Acinetobacter* were negatively correlated with each

other, and these taxa were not found together in any sample.

In the heatmap, many samples clustered together that were dominated by single taxa.

Based on these clusters, we defined community types defined by the genus or genera of bacteria

that were present in these samples with relative abundances ≥ 30% (Fig. 2). *Ellychnia corrusca*

adults were assigned to 6 distinct community types: T-1 (*Mesoplasma*), T-2 (*Salmonella*), T-3

(*Mesoplasma* & *Salmonella*), T-4 (*Serratia*), T-5 (*Rickettsia*) and T-6 (*Pseudomonas*). *Ellychnia*

corrusca egg microbiomes were all assigned to T-13 (*Gluconobacter*), except for a single egg

microbiome that belonged to T-2 (*Salmonella),* the same as some *E. corrusca* adult microbiomes.

Photuris versicolor microbiomes were assigned to 3 community types: T-8 (*Acinetobacter*), T-9

(*Yersinia*) and T-6 (*Pseudomonas*). *Pyractomena borealis* and *Pyropyga decipiens* were assigned

to T-10 (*Empedobacter*) and T-12 (*Cupriavidus*), respectively.

Firefly species harbor unique microbial communities

The four sampled firefly taxa had distinct bacterial communities (Weighted Unifrac

[WUF] PERMANOVA: R^2=0.215, p=0.001; Fig. 3A, Suppl. Fig. S3, Suppl. Table S1). In a

PCoA ordination of WUF beta-diversity distances, *E. corrusca* samples clustered together,

spanning from left to right (Fig. 3). *Pyropyga decipiens* samples clustered together with some *E.*

corrusca samples and those from *Photuris versicolor* on the left side of Figure 3A, and *Ph*

versicolor samples grouped together down the left side and at the bottom, along with the

Pyractomena borealis samples. Species-specific clustering can be seen in the UUF plot, with *E.*

corrusca having the most variance (Suppl. Fig. S3B). The separation between the *E. corrusca*

and *Ph versicolor* microbiomes was maintained in a better-balanced comparison when the few

Pyropyga decipiens and *Pyractomena borealis* samples were excluded (WUF PERMANOVA:

R^2=0.164, p=0.001; Suppl. Fig S4, Suppl. Table S2).

We identified the bacterial taxa whose relative abundance differed between *E. corrusca*

and *Photuris versicolor* fireflies using DESeq. Although *Pseudomonas* ASVs were found in both

fireflies, one *Pseudomonas* ASV was more abundant in *Ph. versicolor* and two other

Pseudomonas ASVs were more abundant in *E. corrusca* (Fig. 3B). Two *Mesoplasma* ASVs,

Salmonella, *Serratia*, *Rickettsia* and *Rickettsiella* were all more abundant in *E. corrusca* than in

Ph. versicolor, and *Tsukamurella*, *Gordonia*, *Curvibacter* and *Sphingobacterium* were all more

abundant in *Ph. versicolor* than in *E. corrusca* (Fig. 3B).

Gut and body microbiomes are distinct, but these differences are firefly-specific

The above analysis indicated that the four sampled firefly groups all hosted distinct

microbiomes (Fig. 3). However, further analyses using these same data also indicated that

different firefly tissues might also host distinct microbiomes, in a species-specific manner (WUF

PERMANOVA for the interactions between species and tissue: R^2=0.048, p=0.003; Suppl. Fig.

S3, Suppl. Table S1). In contrast, a parallel analysis using only samples for which sex was

determined indicated that microbiomes did not differ between the sexes, regardless of species or

tissue type (all PERMANOVAs testing for differences between sexes: $p > 0.05$; Suppl. Fig. S5,

Suppl. Table S3). In the analysis described above with *Pyractomena borealis* and *Pyropyga*

decipiens samples removed to avoid possible artifacts due to unbalanced sample sizes, firefly

tissue microbiomes again differed in a species-specific manner (WUF PERMAONVA for

interactions between *E. corrusca* and *Photuris versicolor* gut and body microbiomes: R^2=0.053,

p=0.001; Suppl. Fig. S4, Suppl. Table S2).

Because these PERMANOVAs indicated that tissue microbiomes differed in a species-

specific manner, we repeated our analysis for each species separately. *Ellychnia corrusca* eggs,

gut, and body microbiomes differed from each other (WUF PERMANOVA: R^2=0.122,

p=0.001), with most gut and body samples clustered together in the PCoA and most egg samples

clustered separately (Fig. 4A, Suppl. Table S4, Suppl. Fig. S6). When the egg samples were

removed from the analysis, *E. corrusca* gut and body microbiomes differed from each other only

when using the UUF distance metric (UUF PERMANOVA: R^2= 0.072, p=0.001; WUF: Suppl.

Table S4). *Wolbachia* had low relative abundance in all *E. corrusca* samples, but was the only

bacterial genus that was more abundant in *E. corrusca* eggs than in adults. *Mesoplasma*,

Pseudomonas, *Acinetobacter* and several other genera were all more abundant in *E. corrusca*

adults than in eggs (Fig. 4B). *Ellychnia corrusca* eggs and adults also had distinct community

types, with *E. corrusca* egg microbiomes mainly assigned to T-13 (*Gluconobacter*), and only a

single egg microbiome assigned to T-3 (*Salmonella*) like those of *E. corrusca* adults (Fig. 2). In

contrast, *Photuris versicolor* gut and body microbiomes more strongly differed from each other

(WUF PERMANOVA: R^2=0.340, p=0.001; Fig. 4C, Suppl. Fig. S7, Suppl. Table S4), with

Pseudomonas, Acinetobacter, Leucobacter, Elizabethkingia, Curvibacter and *Achromobacter* all

being more abundant in *Ph. versicolor* guts than in bodies (Fig. 4D). *Pyropyga decipiens* gut and

body microbiomes did not differ from each other (WUF PERMANOVA: R^2=0.312, p=0.667;

Suppl. Table S4), and *Pyractomena borealis* gut and body microbiomes differed only when

using the UUF distance metric (UUF PEMANOVA: R^2= 0.616, p=0.008; Suppl. Table S4).

These conclusions should be considered preliminary because of the small sample sizes available

for both *Pyropaga decipiens* and *Pyractomena borealis*.

Fireflies host an abundance of potentially symbiotic Mollicutes

Mollicutes were the most abundant bacteria in our 133 firefly samples, consistent with

the handful of Mollicutes that had previously been isolated from different firefly species [23–30].

We therefore created a phylogenetic tree to discover how our most abundant Mollicute ASVs

were related to the sequences of these known firefly Mollicute isolates (Fig. 5). The most

prevalent *Spiroplasma* ASV (ASV Spiroplasma1), found in many of the *E. corrusca* samples,

was similar to the 16S rRNA sequence of *S. corruscae*, a species that was first isolated from *E.*

corrusca [25]. ASV Spiroplasma2 was found in one *Photuris versicolor* sample, and was similar

to the 16S rRNA sequence of *S. ixodetis*, a male-killing agent in butterflies [47]. *S. ixodetis* has

not been found in fireflies to date. The four *Mesoplasma* ASVs, mainly found in *E. corrusca*, are

all similar to the 16S rRNA sequences for *M. corruscae* and *Entomoplasma ellychniae*, which

had both been isolated previously from *E. corrusca* [26, 28, 30]. Although originally separated

using cell morphology and culture media requirements, the genera *Mesoplasma* and

Entomoplasma are paraphyletic [48]. Our phylogenetic tree showing that both *M. corruscae* and

E. ellychniae cluster tightly together, as seen by Gasparich et al. [48], who suggest that the genus

Mesoplasma should be changed to *Entomoplasma*. This would mean that *M. corruscae* would be

reclassified as a member of the genus *Entomoplasma*, along with all other *Mesoplasma* species.

The single *Entomoplasma* ASV, found in *Pyractomena borealis* adults, closely resembled the

16S rRNA sequence for *E. somnilux*, which had been previously isolated from *Pyractomena*

angulata [24]. This phylogenetic tree shows that the Mollicutes sequences detected in study are

highly similar to strains that had been previously isolated from fireflies, except for that related to

S. ixodetis.

DISCUSSION:

Our results show that all of our sampled fireflies have low-complexity microbiomes. The

alpha diversity of our firefly microbiomes was very low, and all samples had only a few taxa

present in high abundance (Fig 1). A previous survey also suggested that beetles had low-

diversity microbiomes, with carnivorous and herbivorous species hosting the lowest diversity

[17]. Our data is consistent with this, as *Photuris versicolor* fireflies are carnivorous and *E.*

corrusca ingest plant sap. Kolasa et. al [17]also describe large variation in microbiome

composition within a host beetle species, which is consistent with our firefly microbiome data.

However, the Shannon diversities measured for our firefly microbiomes (means ranging from 0.7

to 2.52) is lower than those reported in that beetle survey [17]. The low number of correlations

between the relative abundances of firefly taxa (Suppl Fig. S2) is consistent with this

predominance of simple microbiomes, because (by definition) few interactions can occur when

only a single bacterium is abundant in a microbiome, and correlations will inevitably be weak

when there is high variability between microbial communities. This variance and low diversity

could be explained by microbiomes being acquired via neutral community assembly. Fireflies

may be exposed to environmental bacteria by chance, and these bacteria may then multiply and

form a stable microbiome. This would explain why there is high variance between the

microbiomes from individual insects of the same firefly species.

Samples could be assigned to one of thirteen community types, defined by the taxa in

these samples with >30% abundance. Different firefly taxa rarely shared the same community

type, and most community types were defined by a single taxon, with only one community type

having two taxa that were each >30% abundant (Fig. 2). The presence of multiple community

types in each firefly species raises questions about whether the bacteria are either: 1) resident

members replicating within the firefly, or 2) transient members that replicate more slowly than

the rate of expulsion from the firefly [49]. Although such transient microbiomes are not likely to

be conserved between fireflies, it is possible that they do have some effect on their hosts

(positive or negative). Community types vary within the same firefly species, suggesting that

these microbiomes are transient, but some bacteria were found in extremely high relative

abundances (Fig. 1) that could suggest a stable resident microbiome of unknown function. It is

possible that each community type provides a different function for the firefly, or that the

multiple, highly abundant bacteria in different community types, such as *Mesoplasma*,

Salmonella, and *Pseudomonas*, could provide similar functions for their hosts. However, this

may be unlikely for Mollicute bacteria due to their reduced genomes.

Fireflies from different taxa also had distinct microbiomes (Fig. 3A). *E. corrusca* and

Photuris versicolor had distinct bacterial communities, and samples from the same species

clustered together in PCoA plots of beta-diversity (Suppl. Fig. 4). Differences between the

microbiomes of these two groups might be due to their different lifestyles. North American *E.*

corrusca are active in winter and may feed on tree sap [10, 11], and so *E. corrusca* microbial

communities may therefore be acquired via the ingestion of these fluids. *Ellychnia corrusca* also

live for ~9 months. This much longer life span compared to other the firefly taxa in this study

could explain the greater diversity of community types in *E. corrusca*. Unlike *E. corrusca, Ph.*

versicolor females are predatory and active in the summer. Insect gut microbiomes often vary

based on differences in their host's habitat and diet (e.g., omnivory vs. carnivory) [18], and such

differences might underlie the differences that we observed between *E. corrusca* and *Ph.*

versicolor microbiomes. Both *E. corrusca* and *Ph. versicolor* hosted *Pseudomonas* ASVs that

were similar to those previously found on trees and in soils, which could be explained by *E.*

corrusca and *Ph. versicolor* living in soil, on tree bark, and on plant leaves [2], where

Pseudomonas is common. *Pseudomonas* is also commonly found in beetles from the family

Staphylinidae, which are primarily predatory, but not in other non-predatory beetles [17]. This

could suggest a relationship between diet and microbiome composition.

All of the *E. corrusca* and *Ph. versicolor* samples used in this study were collected in

Connecticut, USA, and all *Pyractomena borealis* samples were-lab raised. *Pyropyga decipiens*

samples were collected ~184 miles apart, but had similar microbiome diversities compared to

fireflies collected in Connecticut, which suggests that *Pyropyga decipiens* microbiomes may not

differ between geographic locations. Whether the microbiomes of *E. corrusca, Ph. versicolor*

and *Pyractomena borealis* vary in other geographic locations needs future research.

For some firefly taxa, eggs, guts, and bodies had distinct microbiomes. *Ellychnia*

corrusca adult and egg microbiomes differed from each other, with samples clustering separately

in PCoAs (Fig. 4A, Suppl. Fig. S6), but an *E. corrusca* egg also had the same community type

(T-2; *Salmonella*) as some adults. Although it is unknown which firefly laid this egg, these

results suggest that it might be possible for *Salmonella* to be transmitted vertically between *E.*

corrusca adults and eggs. The mechanisms that *E. corrusca* may use to acquire these microbes

from the environment are unknown, but there are other insects that rely on horizontal

transmission for their symbionts, such as the bean bug *Riptortus pedestris*, which acquires its

Burkholderia symbiont from soil each generation using a mucus-filled organ that the

Burkholderia specifically penetrates and colonizes [50]. Although *E. corrusca* is not known to

have an organ that selects for a specific bacterial symbiont, *Mesoplasma* was found in many

samples of *E. corrusca*. It is possible that *E. corrusca* may select this bacterium from the

environment, or that *Mesoplasma* selects *E. corrusca* as a preferred host. *E. corrusca* is the most

sampled firefly species in our study, and hosts microbiomes belonging to the most community

types. This increased sampling may therefore have increased the number of community types

detected in this organism relative to the other species in our study.

The microbiomes of the male and female *Photuris versicolor* did not differ from each

other, suggesting that the microbiome is not acquired from the firefly prey because only female

Ph. versicolor are carnivorous, although adult *Ph. versicolor* males do eat milkweed nectar.

Photuris versicolor gut and body microbiomes also differed from each other, with gut

microbiomes having higher amounts of *Pseudomonas* and *Acinetobacter* than body microbiomes

(Figs. 4C and 4D). These ASVs may be acquired from the firefly's environment, because both

ASVs were similar to other strains of bacteria that are common in soil and on plants. The

bacteria present in the *Ph. versicolor* gut may be derived from eating other fireflies or nectar, and

thus the *Pseudomonas* ASVs that are common in *Ph. versicolor* guts may originate from their

last meal. The *Ph. versicolor* microbiome may therefore be transient and assembled from the diet

microbiome. Alternatively, it is possible that the *Ph. versicolor* gut microbiome is not transient,

and the gut provides a niche that is favorable for the growth of *Pseudomonas* and *Acinetobacter*

microbes that might originate from soil, plants, or the diet microbiome and subsequently

colonize the *Ph. versicolor* gut.

The origin of the body microbiome remains unclear. Body dissections include all other

parts of the firefly except for the gut, meaning that other non-gut tissues and hemolymph from

inside of the firefly could have been colonized with bacteria. Alternatively, adult firefly

abdomens are surrounded with sclerotized segments called ventrites and tergites on their ventral

and dorsal sides, respectively [2]. These ventrites and tergites have small gaps between them that

allow for movement. Although our firefly samples were ethanol-washed before dissection, this

sterilization may not be absolutely perfect, and the body microbiome may therefore include

bacteria that are trapped within these segments [51] that were acquired during movement on tree

bark or in the leaf litter, where many bacteria are found.

Mollicutes have been isolated from several different firefly species, but their prevalence

and function is unclear [23–30]. We detected 4 *Mesoplasma* ASVs in North American fireflies,

mainly in *E. corrusca*, that all closely resembled *M. corruscae* and *Entomoplasma ellychniae*,

which were both isolated previously from *E. corrusca*. As mentioned in the results, the tight

clustering of *M. corruscae* and *Entomoplasma ellychniae* in Figure 5 implies that both belong to

the same genus. Gasparich et al. [48] proposed reassigning all *Mesoplasma* species to the genus

Entomoplasma, including changing *Mesoplasma corruscae* to *Entomoplasma corruscae*.

Members of the currently defined genus *Mesoplasma* frequently occur on plants without causing

disease, and therefore it could be horizontally acquired by *E. corrusca* when they feed on

milkweed nectar or nutrient-poor tree sap to aid in survival during cold winter months [10, 11].

All the fireflies in this study are omnivorous or carnivorous predators as larvae. Other than adult

female *Ph. versicolor* species, adult fireflies do not need to eat to survive, although some fireflies

might acquire bacteria while feeding as adults. *Pyractomena borealis* larvae are known to ingest

tree sap, just as *E. corrusca* larvae do [4], and were colonized by *Entomoplasma somnilux.*

Although *Entomoplasma somnilux* has only been found in fireflies, it is possible that it was

acquired via the *Pyractomena borealis* diet. Mollicutes and other bacteria may be acquired by a

larval firefly's diet of tree sap and remain colonized through the adult stage. However, it is

unknown whether bacteria survive firefly metamorphosis from larvae to adults. If not, the

presence or absence of Mollicute in these adult fireflies might be explained by the presence of

their diet. Future work will be required to determine the function, if any, of these Mollicutes in

their firefly hosts.

Our results provide the first description of the microbiomes found in *E. corrusca*,

Photuris versicolor, *Pyropyga decipiens* and *Pyractomena borealis*. These microbiomes have

low alpha diversities and are species-specific. Some firefly species also have distinct egg, gut,

and body microbiomes. Future research will determine the function of these microbiomes and

how they are acquired from their environment or transmitted between hosts.

DECLARATIONS

FUNDING

This work was supported by a University of Connecticut Scholarship Facilitation Fund grant to

J.L.K.

CONFLICTS OF INTEREST

Not applicable

ETHICS APPROVAL

Not applicable

CONSENT TO PARTICIPATE

Not applicable

CONSENT FOR PUBLICATION

Emily A. Green and Jonathan L. Klassen approve this for publication. Scott R. Smedley is

deceased, and therefore was unable to provide explicit consent.

AVAILABILITY OF DATA AND MATERIAL

All data are available on NCBI under BioProject PRJNA563849. Raw sequencing reads are

deposited in SRA under BioSample numbers SAMN14678004 – SAMN14678257.

CODE AVAILABILITY

The commands used for all analyses is attached as Suppl. File S2.

AUTHORS CONTRIBUTIONS

Sample collection: SRS; experiment design: EAG, SRS, and JLK; data analysis: EAG; figure

and table creation: EAG; writing (draft preparation and editing): EAG and JLK.

WORK CITED

1. Slipinski SA, Leschen RAB, Lawrence JF (2011) Order Coleoptera Linnaeus, 1758. In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richnes. Zootaxa 3148:203–208. https://doi.org/10.11646/zootaxa.3148.1.26

2. Faust LF (2017) Fireflies, glow-worms, and lightning bugs!: Identification and natural history of the fireflies of the Eastern and Central United States and Canada. University of Georgia Press, Athens, Georgia

3. Barber HS (1951) Fireflies of the genus *Photuris*. Smithson Misc Collect 117:1–66

4. Faust LF, Hughes LS, Zloba MH, Farrington HL (2019) Life history and updated range extension of *Photinus scintillans* (Coleoptera: Lampyridae) with new Ohio records and regional observations for several firefly species. Ohio Biol Surv Notes 9:16–34

5. Eisner T, Goetz MA, Hill DE, et al (1997) Firefly "femmes fatales" acquire defensive steroids (lucibufagins) from their firefly prey. Proc Natl Acad Sci USA 94:9723–9728. https://doi.org/10.1073/pnas.94.18.9723

6. Goetz MA, Meinwald J, Eisner T (1981) Lucibufagins, IV. New defensive steroids and a pterin from the firefly, *Photinus pyralis* (Coleoptera: Lampyridae). Experientia 37:679–680. https://doi.org/10.1007/BF01967916

7. Smedley SR, Risteen RG, Tonyai KK, et al (2017) Bufadienolides (lucibufagins) from an ecologically aberrant firefly (*Ellychnia corrusca*). Chemoecology 27:141–153. https://doi.org/10.1007/s00049-017-0240-6

8. Deyrup ST, Risteen RG, Tonyai KK, et al (2017) Escape into winter: Does a phenological Shift by *Ellychnia corrusca* (winter firefly) shield it from a specialist predator (*Photuris*)?

Northeast Nat 24:147–166. https://doi.org/10.1656/045.024.s717

9.   Faust L, Faust H (2014) The occurrence and behaviors of North American fireflies

(Coleoptera: Lampyridae) on milkweed, *Asclepias syriaca* L. Coleopt Bull 68:283–291.

https://doi.org/10.1649/0010-065x-68.2.283

10.   Rooney JA, Lewis SM (2000) Notes on the life history and mating behavior of *Ellychnia*

corrusca (Coloeptera: Lampyridae). Florida Entomol 83:324–334.

https://doi.org/10.2307/3496351

11.   Faust L (2012) Fireflies in the snow: Observations on two early-season arboreal fireflies

Ellychnia corrusca and *Pyractomena borealis*. Lampyrid 2:48–71

12.   Lloyd JE (1965) Aggressive Mimicry in *Photuris*: Signal repertoires by femmes fatales.

Am Assoc Adv Sci 149:653–654

13.   Arias-Cordero E, Ping L, Reichwald K, et al (2012) Comparative evaluation of the gut

microbiota associated with the below- and above-ground life stages (larvae and beetles) of

the forest cockchafer, *Melolontha hippocastani*. PLoS One 7:e51557.

https://doi.org/10.1371/journal.pone.0051557

14.   Lundgren JG, Lehman RM (2010) Bacterial gut symbionts contribute to seed digestion in

an omnivorous beetle. PLoS One 5:e10831. https://doi.org/10.1371/journal.pone.0010831

15.   Grünwald S, Pilhofer M, Höll W (2010) Microbial associations in gut systems of wood-

and bark-inhabiting longhorned beetles [Coleoptera: Cerambycidae]. Syst Appl Microbiol

33:25–34. https://doi.org/10.1016/j.syapm.2009.10.002

16.   Kaltenpoth M, Steiger S (2014) Unearthing carrion beetles' microbiome: characterization

of bacterial and fungal hindgut communities across the Silphidae. Mol Ecol 23:1251–

1267. https://doi.org/10.1111/mec.12469

17.   Kolasa M, Ścibior R, Mazur MA, et al (2019) How hosts taxonomy, trophy, and

518   endosymbionts shape microbiome diversity in beetles. Microb Ecol 78:995–1013.

519   https://doi.org/10.1007/s00248-019-01358-y

520   18.   Yun JH, Roh SW, Whon TW, et al (2014) Insect gut bacterial diversity determined by

521   environmental habitat, diet, developmental dtage, and phylogeny of host. Appl Environ

522   Microbiol 80:5254–5264. https://doi.org/10.1128/AEM.01226-14

523   19.   Colman DR, Toolson EC, Takacs-Vesbach CD (2012) Do diet and taxonomy influence

524   insect gut bacterial communities? Mol Ecol 21:5124–5137. https://doi.org/10.1111/j.1365-

525   294X.2012.05752.x

526   20.   Engel P, Moran NA (2013) The gut microbiota of insects - diversity in structure and

527   function. FEMS Microbiol Rev 37:699–735. https://doi.org/10.1111/1574-6976.12025

528   21.   Estes AM, Hearn DJ, Snell-Rood EC, et al (2013) Brood ball-mediated transmission of

529   microbiome members in the dung beetle, *Onthophagus taurus* (Coleoptera: Scarabaeidae).

530   PLoS One 8:e79061. https://doi.org/10.1371/journal.pone.0079061

531   22.   Shukla SP, Plata C, Reichelt M, et al (2018) Microbiome-assisted carrion preservation

532   aids larval development in a burying beetle. Proc Natl Acad Sci U S A 115:11274–11279.

533   https://doi.org/10.1073/pnas.1812808115

534   23.   Hackett KJ, Whitcomb RF, Tully JG, et al (1992) Lampyridae (Coleoptera): A plethora of

535   Mollicute associations. Microb Ecol 23:181–193

536   24.   Williamson DL, Tully JG, Rose DL, et al (1990) *Mycoplasma somnilux* sp. nov.,

537   *Mycoplasma luminosum* sp. nov., and *Mycoplasma lucivorax* sp. nov., new sterol-

538   requiring mollicutes from firefly beetles (Coleoptera: Lampyridae). Int J Syst Bacteriol

539   40:160–164. https://doi.org/10.1099/00207713-40-2-160

25. Hackett KJ, Whitcomb RF, French FE, et al (1996) *Spiroplasma corruscae* sp. nov., from

a firefly beetle (Coleoptera: Lampyridae) and tabanid flies (Diptera: Tabanidae). Int J Syst

Bacteriol 46:947–950. https://doi.org/10.1099/00207713-46-4-947

26. Tully JG, Rose DL, Hackett KJ, et al (1989) *Mycoplasma ellychniae* sp. nov., a sterol-

requiring Mollicute from the firefly beetle *Ellychnia corrusca*. Int J Syst Bacteriol

39:284–289. https://doi.org/10.1099/00207713-39-3-284

27. Stevens C, Tang AY, Jenkins E, et al (1997) *Spiroplasma lampyridicola* sp. nov., from the

firefly beetle *Photuris pennsylvanicus*. Int J Syst Bacteriol 47:709–712.

https://doi.org/10.1099/00207713-47-3-709

28. Tully JG, Whitcomb RF, Hackett KJ, et al (1994) Taxonomic descriptions of eight new

non-sterol-requiring mollicutes assigned to the genus *Mesoplasma*. Int J Syst Bacteriol

44:685–693. https://doi.org/10.1099/00207713-44-4-685

29. Wedincamp J, French FE, Whitcomb RF, Henegar RB (1996) Spiroplasmas and

Entomoplasmas (Procaryotae: Mollicutes) associated with tabanids (Diptera: Tabanidae)

and fireflies (Coleoptera: Lampyridae). J Invertebr Pathol 68:183–186

30. Tully JG, Bove JM, Laigret F, Whitcomb RF (1993) Revised taxonomy of the class

Mollicutes: Proposed elevation of a monophyletic cluster of arthropod-associated

Mollicutes to ordinal rank (Entomoplasmatales ord. nov.), with provision for familial rank

to separate species with nonhelical morphology. Int J Syst Bacteriol 43:630–630.

https://doi.org/10.1099/00207713-43-3-630a

31. Sanders JG, Powell S, Kronauer DJC, et al (2014) Stability and phylogenetic correlation

in gut microbiota: Lessons from ants and apes. Mol Ecol 23:1268–1283.

https://doi.org/10.1111/mec.12611

32.   Lee KM, Adams M, Klassen JL (2019) Evaluation of DESS as a storage medium for

564   microbial community analysis. PeerJ 7:e6414. https://doi.org/10.7717/peerj.6414

565   33.   Caporaso JG, Lauber CL, Walters WA, et al (2011) Global patterns of 16S rRNA

566   diversity at a depth of millions of sequences per sample. Proc Natl Acad Sci U S A

567   108:4516–4522. https://doi.org/10.1073/pnas.1000080107

568   34.   R Core Team (2018) R: A Language and Environment for Statistical Computing

569   35.   Callahan BJ, McMurdie PJ, Rosen MJ, et al (2016) DADA2: High-resolution sample

570   inference from Illumina amplicon data. Nat Methods 13:581–583.

571   https://doi.org/10.1038/nmeth.3869

572   36.   McMurdie PJ, Holmes S (2013) Phyloseq: An R package for reproducible interactive

573   analysis and graphics of microbiome census data. PLoS One 8:e61217.

574   https://doi.org/10.1371/journal.pone.0061217

575   37.   Quast C, Pruesse E, Yilmaz P, et al (2013) The SILVA ribosomal RNA gene database

576   project: Improved data processing and web-based tools. Nucleic Acids Res 41:590–596.

577   https://doi.org/10.1093/nar/gks1219

578   38.   Yilmaz P, Parfrey LW, Yarza P, et al (2014) The SILVA and "all-species Living Tree

579   Project (LTP)" taxonomic frameworks. Nucleic Acids Res 42:643–648.

580   https://doi.org/10.1093/nar/gkt1209

581   39.   Davis NM, Proctor DiM, Holmes SP, et al (2018) Simple statistical identification and

582   removal of contaminant sequences in marker-gene and metagenomics data. Microbiome

583   6:226. https://doi.org/10.1186/s40168-018-0605-2

584   40.   Oksanen J, Blanchet FG, Friendly M, et al (2019) vegan: Community Ecology Package. R

585   package version 2.5-4

586   41.   Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion

587   for RNA-seq data with DESeq2. Genome Biol 15:e550. https://doi.org/10.1186/s13059-

588   014-0550-8

589   42.   Kurtz ZD, Müller CL, Miraldi ER, et al (2015) Sparse and compositionally robust

590   inference of microbial ecological networks. PLOS Comput Biol 11:e1004226.

591   https://doi.org/10.1371/journal.pcbi.1004226

592   43.   Shannon P, Markiel A, Ozier O, et al (2003) Cytoscape: a software environment for

593   integrated models of biomolecular interaction networks. Genome Res 13:2498–2504.

594   https://doi.org/10.1101/gr.1239303

595   44.   Csardi G, Nepusz T (2005) The igraph software package for complex network research.

596   InterJournal Complex Syst 1695

597   45.   Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high

598   throughput. Nucleic Acids Res 32:1792–1797. https://doi.org/10.1093/nar/gkh340

599   46.   Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis

600   of large phylogenies. Bioinformatics 30:1312–1313.

601   https://doi.org/10.1093/bioinformatics/btu033

602   47.   Binetruy F, Bailly X, Chevillon C, et al (2019) Phylogenetics of the *Spiroplasma ixodetis*

603   endosymbiont reveals past transfers between ticks and other arthropods. Ticks Tick Borne

604   Dis 10:575–584. https://doi.org/10.1016/j.ttbdis.2019.02.001

605   48.   Gasparich GE, Kuo C (2019) Genome analysis-based union of the genus *Mesoplasma*

606   with the genus *Entomoplasma*. Int J Syst Evol Microbiol 69:2735–2738.

607   https://doi.org/10.1099/ijsem.0.003548

49.   Hammer TJ, Sanders JG, Fierer N (2019) Not all animals need a microbiome. FEMS

Microbiol Lett 366:fnz117. https://doi.org/10.1093/femsle/fnz117

50.   Ohbayashi T, Takeshita K, Kitagawa W, et al (2015) Insect's intestinal organ for symbiont

sorting. Proc Natl Acad Sci U S A 112:5179–5188.

https://doi.org/10.1073/pnas.1511454112

51.   Hammer TJ, Dickerson JC, Fierer N (2015) Evidence-based recommendations on storing

and handling specimens for analyses of insect microbiota. PeerJ 8:e1190.

https://doi.org/10.7717/peerj.1190

FIGURE LEGENDS

Figure 1) Heat map of the 26 most abundant bacterial genera found in the studied fireflies. To

the left, a dendrogram represents the Euclidian distances between the relative abundances of

reads assigned to each bacterial genus, which are labeled on the right. The top dendrogram

clusters the relative abundances of these genera in each firefly sample using ward.D distances.

On the bottom, 3 different rows indicate the host taxon, tissue type, and sex for each sample,

indicated by the different colors in the key to the left of the figure. The topmost color key

represents the relative abundance of reads in each sample that were assigned to each bacterial

genus, with white and dark blue representing 0 and 100% relative abundance, respectively. n =

133

Figure 2) Firefly microbial community types. The X axis indicates the number of samples that

were assigned to each community type. Colors indicate firefly taxa, as indicated by the key. Note

that some gut and body samples originate from the same individual firefly and so are counted

twice.

Figure 3) A: PCoA of Weighted Unifrac distances between microbial communities for all four

firefly taxa. Species are differentiated using colors. *P. borealis = Pyractomena* borealis, *P.*

decipiens = Pyropyga decipiens. n = 133. B: Over- and underrepresented ASV sequences in *E.*

corrusca and *Photuris versicolor*. adults, including both gut and body samples. The X-axis

indicates the genera of bacteria whose relative abundances differed between firefly taxa, and the

Y-axis indicates the log2 fold change in relative abundance between samples, where the higher

numbers indicate overrepresentation in *Ph. versicolor* and negative numbers indicate

overrepresentation in *E. corrusca*. Colors indicate phyla. n = 114

Figure 4) A) PCoA of Weighted Unifrac distances between microbiomes of *E. corrusca* eggs

and adults. Egg, gut and body samples are differentiated using colors. n = 100. B) Over- and

underrepresented ASV sequences in *E. corrusca* egg and adult samples. The X-axis indicates the

genera of bacteria whose relative abundance differed between *E. corrusca* eggs and adults, and

the Y-axis indicates the log2 fold change in these relative abundances between tissues, where the

higher numbers indicate overrepresentation in eggs and negative numbers indicate

overrepresentation in adults. Colors indicate phyla. C) PCoA of Weighted Unifrac distances

between microbiomes of *Photuris versicolor* gut and body samples. Gut and body samples are

differentiated using colors. D) Over- and underrepresented ASV sequences of *Ph. versicolor* gut

and body samples. The X-axis indicates the genera of bacteria whose relative abundance differed

between *Ph. versicolor* gut and body samples, and the Y-axis indicates the log2 fold change in

these relative abundances between tissues, where the higher numbers indicate an

overrepresentation in the body samples. Colors indicate phyla. n = 20

Figure 5) Phylogenetic tree of the Mollicute 16S rRNA gene ASVs in our dataset (in bold) and

reference 16S rRNA gene sequences, particularly those isolated from fireflies and beetles

(indicated by a ^). The tree was constructed using RAxML with 500 bootstraps and rooted using

Bacillus subtilis and *Mycoplasma haemominutum*. Bootstrap values of 60–79 and 80–100 are

indicated by * and **, respectively. Numbers in parentheses indicate the number of samples in

which that ASV was found, and colors group sequences from the same genera. NCBI accession

numbers are shown to the right of each reference sequence. *Me = Mesoplasma*, *My =*

Mycoplasma.

Figure 1)

670 **Figure 2)**

673   **Figure 3)**

676 **Figure 4)**

Figure 5)