Autoproteolytic Products of the SARS-CoV-2 Nucleocapsid Protein are Primed for Antibody Evasion and Virus Proliferation

Corinne A. Lutomski, Tarick J. El-Baba, Jani R. Bolla, Carol V. Robinson*

Physical and Theoretical Chemistry Laboratory, University of Oxford, South Parks Road, OX13QZ Oxford, UK

*Correspondence: carol.robinson@chem.ox.ac.uk

Abstract

The SARS-CoV-2 nucleocapsid (N) protein is the most immunogenic of the structural proteins and plays essential roles in several stages of the virus lifecycle. It is comprised of two major structural domains: the RNA binding domain, which interacts with viral and host RNA, and the oligomerization domain which assembles to form the viral core. Here, we investigate the assembly state and RNA binding properties of the full-length nucleocapsid protein using native mass spectrometry. We find that dimers, and not monomers, of full-length N protein bind RNA, implying that dimers are the functional unit of ribonucleoprotein assembly. In addition, we find that N protein binds RNA with a preference for GGG motifs which are known to form short stem loop structures. Unexpectedly, we found that N undergoes autoproteolytic processing within the linker region, separating the two major domains. This process results in the formation of at least five proteoforms that we sequenced using electron transfer dissociation, higher-energy collision induced dissociation and corroborated by peptide mapping. The cleavage sites identified are in highly conserved regions leading us to consider the potential roles of the resulting proteoforms. We found that monomers of N-terminal proteoforms bind RNA with the same preference for GGG motifs and that the oligomeric state of a C-terminal proteoform ($N_{156-419}$) is sensitive to pH. We used mass spectrometry to show that N binds to a monoclonal antibody raised against full-length N. No antibody interactions were detected for N proteoforms without C-terminal residues, therefore locating antigenic regions towards the C-terminus. We then tested interactions of the proteoforms with the immunophilin cyclophilin A, a key component in coronavirus replication. We found that N_{1-209} and N_{1-273} bind directly to cyclophilin A, an interaction that is abolished by the approved immunosuppressant drug cyclosporin A. We propose that the proteoforms generated via autoproteolysis evade antibody detection through removal of the antigenic C-terminus and facilitate interactions with structured RNA or cyclophilin thereby enabling the virus to proliferate.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is the etiological agent of coronavirus disease 2019 (COVID-19) which reached pandemic status in fewer than three months following its discovery. As of October 2020, there are >34 million infected and more than 1 million deaths.[1] However, SARS-CoV-2 is not the first coronavirus to cause widespread disease. Two epidemics in the last two decades have been caused by coronaviruses: severe acute respiratory syndrome (SARS) in 2002 and the Middle East respiratory syndrome (MERS) in 2012. Despite belonging to the same viral genus, SARS-CoV-2 has proven far more infectious, implying molecular differences between the viruses. Understanding the molecular components of viral proliferation in SARS-CoV-2 is therefore important to develop new therapeutics to combat COVID-19.

SARS-CoV-2 packages a large RNA genome of ~30 kb which encodes for 25 non-structural and four structural proteins (spike, nucleocapsid, membrane, and envelope proteins). The nucleocapsid (N) protein is one of the most abundant viral proteins; hundreds of copies of N make up the viral core that encapsulates the genomic RNA. In addition to its essential role in RNA replication/transcription and virion assembly, coronavirus N proteins play essential roles in host cell signaling pathways, immune system interference, cell cycle regulation, and chaperone activity.[2] Interestingly, the nucleocapsid and spike proteins are the main immunogens circulating in the blood of COVID-19 patients.[3] However, quantitative measurements of plasma or serum from SARS-CoV-2 patients found that the adaptive immune response to the N protein is more sensitive than the spike protein[4], making it a better indicator of early disease and a good target for antiviral therapies.

N protein has been the subject of much effort in structure elucidation in order to guide the design of novel antivirals.[5,6] Several compounds targeting nucleocapsid proteins of other viruses have proven effective in *in vitro* studies, specifically blocking replication, transcription, and assembly.[7,8,9,10] However, the timeline for developing approved antivirals takes years and the need for treatments to combat COVID-19 is urgent. Therefore, repurposing existing drugs is an attractive and immediate solution to combat COVID-19.[11] On this front, key host interactions have been mapped for the proteins encoded by SARS-CoV-2 to identify host targets for drug repurposing.[12] One particular study revealed 66 potential drug targets, three of which are direct interactors of N protein. The safe and effective use of new and existing therapeutics to target specific viral processes however relies on an understanding of the sequence of interactions between viral and host proteins and their controls.

Here, we present a comprehensive analysis of the SARS-CoV-2 N protein using native mass spectrometry (MS), top-down fragmentation, and bottom-up sequencing. We find that the full-length N protein undergoes autoproteolysis at highly conserved sites to generate at least five unique proteoforms. We identify various stoichiometries of the proteoforms that are influenced by pH and that may be critical targets in drug design. We evaluate the propensity for N and N proteoforms to bind different RNA sequences and find that only the dimeric form of full-length N binds to RNA, suggesting it is the functional unit of ribonucleoprotein assembly. We find that full-length N binds to monoclonal antibodies raised against the full-length protein but interestingly, interactions with N-terminal proteoforms are not detected. Finally, we show that two N proteoforms directly interact with cyclophilin A, a highly abundant cytosolic host protein implicated in viral replication. We found that cyclosporin A, an immunosuppressive drug, abolishes the interaction between N proteoforms and cyclophilin A. Our results indicate that SARS-CoV-2 N protein is a

multifunctional protein and that N proteoforms are primed for additional functions related to viral propagation and survival.

Results and Discussion

N Protein undergoes autoproteolysis in the vicinity of the linker region

Nucleocapsid proteins of coronaviruses share a similar topological organization[13] and show high sequence homology among related coronaviruses.[14] The distribution of residues is characterized by two major structural domains, the RNA binding and oligomerization domain (Figure 1A). [15] Similar to other coronavirus nucleocapsid proteins, the two domains are separated by a long and flexible linker region thought to be devoid of secondary structure.[2] An N-terminal arm and a C-terminal tail flank the RNA binding and oligomerization domains, respectively. We constructed a plasmid consisting of the full-length nucleocapsid protein, an N-terminal purification tag, and a cleavage site (Figure 1A). We expressed and purified this construct in *Escherichia coli* and verified the cleavage of the affinity tag via SDS-PAGE (Figure 1B). Before removal of the purification tag, three distinct protein bands were detected at ~49, ~38 and ~28 kDa. Following removal of the tag, all three protein bands migrated by ~3 kDa, or the mass of the tag (Table 1).

We confirmed that each band corresponded to N protein by in-gel trypsin digestion followed by LC-MS-based bottom-up proteomics. All three bands contained peptides from the N protein, resulting in 78.1%, 49.9%, and 43.2% sequence coverage for bands 1, 2, and 3, respectively. To determine the representation of protein domains in each gel band, we plotted the distribution of the peptides detected across the five protein domains (Figure 1C). As anticipated, we observe an unbiased distribution of peptides across all five domains for band 1, consistent with the expectation that tryptic peptides would be reasonably distributed across the full-length protein. Over 50% of the total peptides detected in bands 2 and 3 were localized to the RNA binding domain, suggesting that the proteins are predominantly N-terminal derivatives. However, in all three bands, peptides located in the 58-residue C-terminal domain were detected, indicating the purified protein is made up of a diverse mixture of N proteoforms[16] and not biased to N-terminal species due to the location of the purification tag.

We recorded a native mass spectrum to identify the masses of the proteoforms present together with the full-length protein (Figure 1D). Two main charge state envelopes centered at 13+ and 20+ species were identified, consistent with coexistence of monomers and dimers. Deconvolution of the *m/z* signals provided experimental masses of 45,769 ± 1 Da and 91,537 ± 2 Da, respectively, which are in excellent agreement with the theoretical monomeric (45,769.83 Da) and dimeric (91,539.66 Da) masses of full-length N protein. The second distribution of monomers and dimers have deconvoluted masses of 28,735 ± 2 Da and 57,534 ± 1 Da, respectively. The SDS-PAGE analysis indicated the presence of additional proteoforms not visible in this spectrum. We separated these lower molecular weight species from the full-length N using size exclusion chromatography (Figure S1). The mass spectrum of the pooled fractions revealed the presence of four additional proteoforms that range in mass from 22,612 to 42,922 Da.

To determine whether these proteoforms are the products of autoproteolysis we incubated N protein in a cocktail of protease inhibitors at 25 °C for 1 week. Aliquots of this solution were withdrawn at regular time intervals and analyzed by SDS-PAGE (Figure S2). The same four bands were detected at two, four, and seven days during incubation. The temporal decrease in the band

corresponding to the full-length N protein, concomitant with the increase in intensity for the truncated proteoforms, confirms that the truncations arise from autoproteolysis. It is also notable that no new bands appear at any of the measured time points over one week which suggests that cleavage of the full-length protein is not the result of a co-purifying non-specific protease but rather is highly specific and therefore likely due to autoproteolysis.[17] Here, we define autoproteolysis as the spontaneous cleavage of the peptide backbone, likely induced by conformational strain.[18,19,20]

After incubation for one week, the mass spectrum shows a series of peaks corresponding to five unique protein distributions (Figure 2A). To determine the identity of each proteoform, we adapted a two-tiered tandem mass spectrometry approach[21] to determine intact mass and amino acid sequence for each series of peaks in the mixture. An individual peak in the mass spectrum was first isolated and subjected to electron transfer dissociation (ETD) under conditions that do not result in the formation of fragments but instead produce a series of charge-reduced peaks (Figure 2B). The charge-reduced spectrum was necessary to confirm the assignment of the charge state series for each proteoform. The assigned charge states were then used to obtain deconvoluted masses of each proteoform present in solution.

We tentatively assigned the proteoforms based on their intact masses and then generated sequence ions to confirm our assignment. Individual proteoforms were subjected to fragmentation by higher-energy collision induced dissociation (HCD), which accelerates the isolated ions into an inert gas to induce fragmentation along the amide backbone. The fragmentation products were then used to determine the molecular composition and to localize the exact site of cleavage. The HCD spectrum results in a series of singly-, doubly-, and triply-charged sequence ions exemplified by the fragmentation of the 9+ charge state at *m/z* 2616.79, Figure 2C. Fragmentation of intact proteins under native conditions is expected to yield 3-10% sequence coverage at the termini[22]. The propensity for fragmentation differs depending on several criteria (e.g. mass, charge, composition, structure) for natively folded proteins[23]. Here, we achieved ~4% sequence coverage of the proteoforms by native top-down MS (Table S1). Fragmentation at the termini was complementary to our goal – to confirm sites of autoproteolytic cleavage of the N-protein.

Cleavage sites are highly conserved in the SARS-CoV-2 genome

Considering the proteoforms identified, we note that three result from cleavage after alanine (N_{1-220}, N_{1-273}, $N_{156-419}$), one from cleavage following an arginine, but preceded by alanine (N_{1-209}), and one results from cleavage C-terminal tail following a valine (N_{1-392}), (Figure 2D). Proteoforms N_{1-209} and N_{1-220} contain primarily the RNA binding domain as they cleave within the flexible linker region. The major component of N_{1-273} is also the RNA binding domain, but in this case is followed by the linker region and a small portion of the oligomerization domain. Conversely, $N_{156-419}$ comprises mainly the oligomerization domain while still retaining a small portion of the RNA binding domain.

Although we find no sequence similarity in the residues that flank each cleavage site, a commonality is that cleavage occurs immediately adjacent to a hydrophobic residue. We were intrigued to discover if these sites were subject to mutation. The locations of the gene and amino acid mutations have been mapped for 38,318 SARS-CoV-2 genome sequences obtained from the China National Center for Bioinformation 2019 Novel Coronavirus Resource.[15] In the sampled population, there are 200 amino acid mutations that occur more than once, and these substitutions are heavily localized at or near the linker region. The proteolytic sites at A220, A273,

A156, T392 are mutated only 4, 1, 9, and 2, times, in >38,000 genomes sequenced, making them highly conserved. By comparison the site at R209 is mutated a total of 36 times; most commonly mutated to T via a G→C missense mutation at position 626 in the gene. The specificity of cleavage at the conserved residues identified here, despite no common motif, suggest that there may be a structural component directing autoproteolysis. This is supported by small angle x-ray scattering measurements which demonstrate that the flexible linker region is not fully extended but contains elements of structure.[24]

The oligomeric states of N proteoforms are influenced by pH

To better understand the role of the individual proteoforms we expressed and purified four individual constructs and recorded native mass spectra of N_{1-209}, N_{1-220}, N_{1-273}, and $N_{156-419}$ from solutions at different pHs (Figures S3-S6). Mass spectra for N_{1-209} and N_{1-220} recorded at pH 5.0, 7.4 and 8.0 reveal highly abundant charge state series centered at 9+ and a low abundance distribution of signals centered at the 13+ charge state corresponding to monomers and dimers, respectively. The mass spectra for N_{1-273} show that it is predominantly monomeric with no significant change in charge state distribution or oligomeric state across the range of pH values tested. However, we observe two low abundant charge state series corresponding to N_{1-209} and N_{1-220}, suggesting that N_{1-273} continues to undergo autoproteolysis at the previously mapped residues (Figure S6).

In contrast to the N-terminal proteoforms, mass spectra for $N_{156-419}$ reveal multiple charge state distributions at pH 5.0, 7.4 and 8.0 (Figure S6). At pH 8, the mass spectrum reveals three charge state distributions centered at 10+, 18+, and 27+ with average masses corresponding to monomers, trimers, and a low population of hexamers (Table S2). At pH 7.4, monomers, dimers, and trimers persist. At the lowest pH (pH 5.0) $N_{156-419}$ is exclusively trimeric. Finally, the mass spectra for full-length N at pH 5.0 and 8.0 reveal broadened and featureless peaks suggesting that N_{FL} is likely aggregated (Figure S7). A low abundance series of highly charged peaks centered at 18+ at pH 8.0 indicates some protein unfolding. Overall, we conclude that N_{1-209}, N_{1-220}, N_{1-273} do not undergo significant pH-dependent changes in oligomeric state while C-terminal fragments are highly sensitive with trimers predominating under both high and low pH conditions.

RNA sequence influences binding stoichiometry

RNA binding and ribonucleoprotein complex formation is the primary function of coronavirus N proteins. The N protein binds nucleic acid nonspecifically[24], however the production of infectious virions relies on N protein forming specific interactions with viral RNA among an abundance of different cellular RNA species**.** With knowledge of the stoichiometries of N_{FL} and N proteoforms, we sought to determine the propensity for these species to bind specific RNA sequences. We created single-stranded RNA oligonucleotides comprised of 20 nucleotides of repeating sequences (4×-GAUGG, 4×-GAGAA). Considering the promiscuity of N protein, we chose sequences shown to interact with the HIV polyprotein (which includes a nucleocapsid domain) at the different stages of virus assembly[25] and hypothesized that N proteins of different oligomeric states would exhibit similar bias toward artificial RNA motifs.

We incubated N proteoforms and RNA oligonucleotides at a molar ratio of 4:1 protein:RNA and recorded native mass spectra for all N protein-RNA complexes. The mass spectrum for N_{1-209} bound to 4×-GAUGG (Figure 3A) reveals three charge state distributions with masses that correspond to apo N_{1-209} monomer and N_{1-209} bound to one and two 4×-GAUGG RNA oligonucleotides. The charge state distribution corresponding to N_{1-209} bound to two 4×-GAUGG RNA predominates over the single RNA bound protein. Conducting the same experiment with a different oligonucleotide (4×-GAGAA) reveals only one additional charge state distribution for N_{1-209} bound to one oligonucleotide (Figure 3B). Similar RNA binding stoichiometries are observed for N_{1-220} and N_{1-273}; the mass spectra for N_{1-220} and N_{1-273} reveal distributions corresponding to the binding of one and two 4×-GAUGG oligonucleotides. Only one additional distribution is observed for N_{1-220} or N_{1-273} incubated with 4×-GAGAA oligonucleotide which corresponds to one oligonucleotide bound, (Figures S8-9, Table S3) confirming the preference for the 4×-GAUGG sequence for all N-terminal constructs.

To examine if this preference was also observed for full-length N, we incubated the protein with 4×-GAUGG oligonucleotide. A series of peaks was identified corresponding to the N protein dimer bound to two 4×-GAUGG oligonucleotides (Figure 3C). Similarly, in the presence of the 4×-GAGAA RNA oligonucleotide, an additional RNA-bound distribution is observed, however the deconvoluted mass indicates that only one 4×-GAGAA oligonucleotide is bound to the N_{FL} dimer (Figure S10, Table S3). No RNA binding to the monomeric form of N_{FL} is observed, regardless of oligonucleotide sequence. Considering the different properties of the two RNA oligonucleotides, the 4×-GAUGG oligonucleotide contains three GGG motifs which form short stem-loops and are known to contribute additively to the efficiency of genome packaging in related viruses.[26] Furthermore, selective RNA packaging has been described as a feature of innate immune response evasion.[27] Our results emphasize that RNA sequence, and likely the secondary structure, is important for interactions with the N-protein. The preference for the RNA sequence known to form stem loops has implications in efficient genome packaging and likely contributes to an optimized packing density in the intact virion. Notably, only N_{FL} dimer binds RNA, suggesting that the dimer is the functional unit of the SARS-CoV-2 ribonucleoprotein assembly.

Antigenic regions of N are located at the C-terminus

Since N protein is detected by antibodies with higher sensitivity than any other structural proteins of SARS-CoV-2[4] we sought to determine if N proteoforms shared similar immunogenic properties as N_{FL}. We first incubated N_{FL} with a monoclonal antibody raised against the full-length protein in molar ratios of 1:1 (Figure 4A). The mass spectrum reveals three distributions >6000 *m/z* with deconvoluted masses that correspond to the apo antibody as well as one and two N_{FL} bound (Table 2, Figure S11). When the same antibody was incubated with three N-terminal proteoforms (N_{1-209}, N_{1-220}, and N_{1-273}) we did not observe any antibody binding in mass spectra (Figure 4B).

To validate these results with conventional methods we turned to protein detection by Coomassie stain and immunoblotting (Figure 4B inset). The Coomassie stain detects N_{FL} and all proteoforms in high abundance as indicated by the dark blue bands. In contrast, the immunoblot reveals dark bands for only N_{FL} and $N_{156-419}$, including higher order oligomers of $N_{156-419}$ that were not detected by Coomassie staining or mass spectrometry. N_{1-209} and N_{1-273} are barely detectable by the antibody, and N_{1-220} completely evades antibody detection. This suggests that N_{1-209}, N_{1-220}, and N_{1-273} are virtually undetected by the antibody responses and therefore maybe able to carry out additional roles in virus proliferation mechanisms.

N proteoforms interact directly with cyclophilin A

Cyclophilin A (CypA) is a highly abundant immunophilin, found in host cells, that plays multifunctional roles in modulating immune responses. In addition, Cyp A has been implicated in the replication cycle of several viruses, including coronaviruses.[28] We sought to determine if CypA plays a role in SARS-CoV-2 infection through monitoring direct interactions with N_{FL} or N proteoforms. We incubated N_{1-209} and CypA in a 1:1 molar ratio and used native mass spectrometry to measure possible interactions. The mass spectrum reveals charge state distributions that correspond to monomeric N_{1-209} and CypA, and three distinct charge state distributions at m/z >3000 (Figure 5A). The three higher-m/z distributions correspond to: (i) heterodimers of N_{1-209} and CypA, (ii) homodimers of CypA, and (iii) a low population of homodimers of N_{1-209} (Table 2). We observe similar interactions for N_{1-273} (Figure S12). Notably, we do not observe evidence of CypA interacting with N_{1-220} or N_{FL} under the same conditions (Figure S13-15). To determine if the interaction between N_{1-209} and CypA could be inhibited by an approved immunosuppressant cyclosporin A (CsA), we incubated the N_{1-209}:CypA-complex with a 2-fold molar excess of the drug (Figure 5B). The mass spectrum reveals three abundant charge state distributions: (i) a distribution centered at 9+ corresponding to N_{1-209} monomers, (ii) a highly abundant distribution centered at a charge state of 8+ that corresponds to CypA bound to CsA, and (iii) a low abundant distribution that corresponds to monomeric CypA (Table 2). Very low abundance distributions of N_{1-209} homodimers, CypA homodimers, and N_{1-209}-CypA heterodimers are barely detected following magnification >3000 m/z. Therefore, we can conclude that the 1:1 interaction between CypA and the N_{1-209} proteoform can be inhibited by CsA binding to CypA.

Conclusion

We present a comprehensive characterization of SARS-CoV-2 N protein and highlight potential features that might influence SARS-CoV-2 infectivity (Figure 6). Specifically, we find that N protein undergoes autoproteolysis at highly conserved residues in the vicinity of the linker region, separating the two major domains (the RNA binding and oligomerization domains). We identify various stoichiometries of N proteoforms that are influenced by pH, explicitly $N_{156-419}$, which forms stable oligomers under both high and low pH solution conditions. We also show that N_{FL} and N proteoforms bind RNA with a preference for GGG-motifs and present evidence for N_{FL} dimers being the functional unit of assembly in ribonucleoprotein complexes. In terms of immunogenicity, we use MS and immunoblot techniques to demonstrate that the antigenic site of SARS-CoV-2 N protein resides towards the C-terminus and that the immunophilin CypA binds directly to N_{1-209} and N_{1-273}, but not N_{FL} or N_{1-220}, an interaction that can be inhibited through addition of the immunosuppressant cyclosporin A. It is interesting that N_{1-220} evades antibody and CypA interactions and therefore must be a structural effect not seen in the other proteoforms.

Interestingly, we find the autoproteolytic products of SARS-CoV-2 N are similar to those observed for SARS-CoV N protein where the sites of cleavage were estimated to be located near the linker domain.[17] The proteolysis of N protein has also been observed in SARS-infected Vero cells[29,30], indicating that cleavage of N protein is likely a feature of coronavirus infection and is not unique to SARS-CoV-2 N protein.

One of the most critical steps of viral proliferation is the packaging of genomic material and its assembly into new virions. Our results provide a means to monitor such interactions between N protein and the elaborate structural features of genomic RNA. We find that N_{FL} and N

proteoforms exhibit preference for distinct RNA sequences. Disruption of RNA viral genomes has proven to inhibit replication in some viruses.[31,32] In the case of SARS-CoV-2, disruption of preferred RNA structures, and therefore disruption of N protein-RNA interactions, presents a promising strategy to intervene in replication and the potential to develop live attenuated vaccines.

One of the first proteoforms generated by autoproteolysis comprises the C-terminal domain $N_{156-419}$ which forms dimers, trimers and hexamers as a function of pH. The absence of detectable monomer and dimer for this proteoform with time and the immunoblot results reveal that $N_{156-419}$ forms higher order oligomers in excess of 100 kDa. Interestingly, this proteoform encompasses the antigenic site in-line with the epitope mapping of the SARS-CoV N protein which also localized a C-terminal antigenic site.[33,34,35,36] We suggest therefore that the ability to fine-tune the oligomeric state of proteoforms, in response to pH, containing these antigenic C-terminal regions may serve to increase their avidity to antibodies. We propose that circulating $N_{156-419}$ oligomers act as decoys for neutralizing antibodies. This proposal is in line with reports for a number of viruses, where antigenic proteins, or protein complexes, are shed in high abundance to consume neutralizing antibodies.[37,38,39] This strategy has also been used in the development of antiviral therapies for SARS-CoV-2 wherein soluble variants of the host receptor (decoy receptors) are used to bind and neutralize the infectious virus.[40] By contrast, our observation that N_{1-209}, a proteoform that encompasses the RNA binding domain, does not bind to the antibody, implies that this proteoform may escape antibody detection and thereby enable unchecked virus proliferation.

Additional immunological roles of N proteoforms have been demonstrated by the highly specific interactions between cyclophilin A and the N_{1-209} and N_{1-273}, but not N_{FL}. Interestingly cyclophilin A is implicated in inflammation and is a mediator of cytokine production.[41] In cases of severe COVID-19 infection, patients have presented with aberrant inflammatory responses, known as "cytokine storms" which can be fatal.[42] The symptoms identifying cytokine storms have been established,[43] however, the viral component(s) that trigger such a response have yet to be elucidated. We can only speculate that specific interactions between N proteoforms and cyclophilin A, as demonstrated here, may play a role in producing such an aberrant immune response. Pathways to intervene with such interactions could therefore prove beneficial as a component of treatment strategies.

Several additional therapeutic avenues are highlighted by this study. Firstly, inhibition of the autoproteolysis reaction would prevent formation of the antigenic decoys and the release of RNA binding proteoforms for virus propagation. While the mechanism of autoproteolysis of N-protein is not yet understood, systematic mutation of cleavage sites defined here could lead to mechanistic and structural insights to enable small molecule screening of potential protease inhibitors. Knowledge of N protein- structured RNA interactions could also aid the design of new therapeutics that would inhibit successful replication. However, since N protein oligomerizes in the absence of RNA,[44] *de novo* drug design of assembly inhibitors is complex as it is necessary to consider oligomerization propensity of the various proteoforms at the pH regimes encountered in the cellular environment. CsA and cyclosporine-derivatives, such as Alisporivir, however are more straightforward to track and have become attractive candidates to treat COVID-19.[45] Disruption of the cyclophilin A- N-proteoform interactions shown here provides a convenient means of screening potential inhibitors for hit to lead optimization.

Finally, although we have uncovered many potential roles of N protein and its proteoforms it interactions with at least nine host proteins involved in RNA processing have yet to be defined.[12]

Furthermore, defining post-translational modifications[46] and interactions with other host proteins is a necessary endeavor that will provide a more complete knowledge of the multifaceted roles of SARS-CoV-2 N protein. The results presented here however do prompt further investigation of a number of therapeutic strategies including inhibition of the autoproteolyis reaction, perturbation of N-protein RNA binding, and prevention of the cyclophilin A proteoform interactions. Given the likelihood that no one intervention is likely to ameliorate the complex symptoms of COVID-19 infection, our findings contribute proteoform-specific information that may guide some of the many therapies currently under investigation.

Methods

Plasmid construction and cell growth. A codon-optimized synthetic gene corresponding to the full length nucleocapsid protein (Thermo GeneArt, Regensburg, Germany) was cloned into a modified pET28a vector using the In-Fusion cloning kit (Takara Bio Saint-Germain-en-Laye, France). The resulting plasmid encoded for an N-terminal His_6 tag followed by thrombin and tobacco etch virus cleavage sequences upstream of the full-length nucleocapsid protein sequence. To generate the nucleocapsid proteoforms, the desired sequences pertaining to the truncated forms of the N protein were subcloned from the synthetic gene using polymerase chain reaction (Phusion polymerase, New England Biolabs, Hertfordshire, UK). All genes were cloned into the modified pet28 vector and gene sequences were confirmed by Sanger Sequencing.

Plasmids were transformed into BL21 (DE3) and streaked onto LB agar plates supplemented with 50 mg mL^{-1} kanamycin. Several colonies were used to inoculate 100 mL of LB broth supplemented with kanamycin and grown at 37 °C overnight. 10 mL of the overnight precultures were used to inoculate 1 L of LB broth supplemented with kanamycin. Cell cultures were grown to $OD_{600} \sim 0.6$ before inducing protein expression with 0.5 μg mL^{-1} IPTG. Cells were grown for an additional 4 hours at 37 °C before harvesting via centrifugation (5000 x g, 10 minutes, 4 °C). Cell pellets were flash frozen in liquid nitrogen and stored at -80 °C until use.

Protein Purification. Cell pellets were resuspended in lysis buffer (25 mM Tris-HCl pH 8.0, 500 mM NaCl, 5 mM $MgCl_2$, 5 mM β-mercaptoethanol, 5 mM imidazole, 10% v/v glycerol) containing EDTA-free protease inhibitor tablets (Roche). Cells were lysed by five passes through a microfluidizer (prechilled to 4 °C) at 20,000 psi. Cell debris was pelleted by centrifugation (20,000 x g, 20 minutes, 4 °C). The supernatant containing soluble nucleocapsid protein was passed through a 0.45 μm filter.

Supernatant was loaded onto a Ni-NTA column pre-equilibrated in loading buffer (25 mM Tris-HCl pH 8.0, 500 mM NaCl, 5 mM MgCl2, 5 mM β-mercaptoethanol, 20 mM imidazole, 10% v/v glycerol) and allowed to pass via gravity flow. To remove common contaminating proteins, a heat-treated BL21 (DE3) *E. coli* lysate in loading buffer containing 10 mM MgATP was passed over the immobilized nucleocapsid protein using a protocol by Rial and Ceccarelli.[47] The resin was washed with 10 column volumes of wash buffer (25 mM Tris-HCl pH 8.0, 500 mM NaCl, 5 mM MgCl2, 5 mM β-mercaptoethanol, 80 mM imidazole, 10% v/v glycerol), then eluted twice with 10 mL of elution buffer (25 mM Tris-HCl pH 8.0, 500 mM NaCl, 5 mM $MgCl_2$, 5 mM β-mercaptoethanol, 400 mM imidazole, 10% v/v glycerol).

The eluted protein was mixed with TEV protease in a 100:1 (w/w) and loaded into a 3 kDa MWCO dialysis cassette (Thermo Fisher Scientific, United Kingdom). Cleavage of the His_6-thrombin-TEV tag was carried out overnight at 4 °C in lysis buffer. The cleaved tag and TEV protease were separated from the untagged protein using reverse immobilized metal affinity chromatography on a Ni-NTA column prepared in loading buffer. The flow-through containing the untagged protein was collected and concentrated in a 10k MWCO centrifugal filter before MS analysis. Protein concentration was determined using UV-VIS spectroscopy by monitoring the absorbance at 280 nm with a theoretical extinction coefficient ($\varepsilon \sim 43890$ M^{-1} cm^{-1}) determined using the Expasy Protparam tool.

Size Exclusion Chromatography. Full-length N protein and N proteoforms were separated using a Superdex 10/300 increase GL column equilibrated in lysis buffer. Fractions corresponding to N_{FL} and N proteoforms were pooled and concentrated to ~10 µM before MS analysis.

Native Mass Spectrometry. RNA oligonucleotides of repeating sequences (4×-GAUGG, 4×-GAGAA) were purchased from Integrated DNA Technologies. Recombinant human cyclophilin A (product ab86219) was purchased from Abcam (Cambridge, United Kingdom). Cyclosporine A purchased from Merck Life Science (Dorset, United Kingdom). N protein and all binding partners were buffer exchanged or diluted into 500 mM NH_4OAc pH 5.0, 7.4, or 8.0. Buffer exchange was carried out using Zeba™ Spin Desalting Columns, 7K MWCO (Thermo Fisher Scientific, United Kingdom).

Measurements were carried out on Qexactive UHMR or Orbitrap Eclipse. The Q-exactive instrument was operated in the positive ion mode using the manufacturer's recommended parameters for native MS. The instrument was operated at a resolving power of 12,500 (at *m/z* 200). An electrospray was generated by applying a slight (~0.5 mbar) backing pressure to an in-house prepared gold-coated electrospray capillary held at ~1.2 kV relative to the instrument orifice (heated to ~100 °C).

An Eclipse Tribrid instrument was also used for native MS and top-down sequencing. The instrument was set to intact protein mode at standard ion routing multipole pressure of 10 mTorr. Ion voltages were set to transmit and detect positive ions at a resolving power of 12,500 (at *m/z* 200). An electrospray voltage of ~1.2 kV and ~0.5 mbar backing pressure were used for ion formation; desolvation was assisted using an instrument capillary temperature of ~100 °C. To identify the accurate mass and sequence of each proteoform, we used a similar approach to that described by Huguet et al.[21] Briefly, a desired signal was isolated using the ion trap (10 *m/z* isolation window, charge state set to 10) and subjected to: (i) electron transfer dissociation (ETD, 3 ms activation time, 1.0×10^6 ETD reagent target) to generate a charge-reduced series for accurate intact mass determination, and (ii) higher energy collisional dissociation (HCD) using ~30 – 50 V HCD collision energy to generate fragment ions.

Monoisotopic masses of fragment ions generated by HCD having a normalized intensity of 10% or higher were fed into Prosite Lite software.[48] A series of candidate sequences that best matched the measured intact masses for each proteoform were generated. The monoisotopic fragment masses were matched to expected ions generated in silico based on the provided

candidate sequence. Comparison of the statistical likelihood for each match compared to a series of candidate sequences (Table S2) localized the cleavage sites to those outlined in Figure 2D.

Liquid Chromatography and Bottom-up Mass Spectrometry. Full-length N was separated from the lower molecular weight proteoforms on a 0.8 mm 4-12% bis-tris SDS-PAGE gel (Invitrogen) and stained with Coomassie Blue. Bands were excised from the gel, minced, and digested with sequencing grade trypsin (Promega, Madison, WI, U.S.A) at 37 °C overnight, extracted with 80% acetonitrile (0.1% formic acid), and dried on a vacuum concentrator. The extracted peptides were resolubilized in buffer A (H_2O, 0.1% FA) and loaded onto a reverse phase trap column (Acclaim PepMap 100, 75µm x 2 cm, nano viper, C18, 3 µm, 100 Å, ThermoFisher, Waltham, MA, U.S.A.) using an Ultimate 3000 for 50 µL at a flow rate of 10 µL min^{-1}. The trapped peptides were then separated using a 15 cm reverse phase analytical column (350 µm x 75 µm) packed in-house (3 µm C18 particles) using a 60 min linear gradient from 5% to 40% buffer B (80% acetonitrile, 20% water, 0.1% formic acid) at a flow rate of 300 nL min^{-1}. The separated peptides were then electrosprayed in the positive ion mode into an Orbitrap Eclipse Tribrid mass spectrometer (ThermoFisher, San Jose, CA, USA) operated in data-dependent acquisition mode (3 s cycle time). Precursor and product mass analysis occurred in the Orbitrap analyzer (120,000 and 60,000 resolving power at m/z 200, respectively). High intensity (threshold: 1.0×10^4) precursors with charge state between $z = 2$ and $z = 7$ were isolated with the quadrupole (0.5 m/z offset, 0.7 m/z isolation window) and fragmented using higher energy collision induced dissociation (HCD collision energy = 30%). Additional MS/MS scans for precursors within 10 ppm were dynamically excluded for 30 s following the initial selection. MS/MS scans were collected using an automated gain control setting of 1.0×10^4 or a maximum fill time of 100 ms. LC-MS data were searched against both the *E. coli* proteome manually annotated with the SARS-CoV-2 nucleocapsid protein sequence using MaxQuant v1.6.17.0.

Western Blot. Recombinant antibody generated from the full-length SARS-CoV-2 nucleocapsid protein (product ab272852) and anti-human secondary antibody were purchased from Abcam. Proteins were resolved on a 4-12% Bis-tris gel using SDS-PAGE and transferred to a PVDF membrane (pore size 0.45 µm). The membrane was blocked in 5% milk in TPBS for 1 hour at RT, incubated with primary antibody in 1:2000 dilution into blocking buffer (also 1 hour, RT) washed with TPBS and incubated with secondary antibody (1:10000) in blocking buffer (also 1 hour, RT). The PVDF membrane was incubated with Horseradish peroxidase chemiluminescent substrate (Pierce ECL Western Blotting Substrate, Thermo Scientific) before detection on photographic film and developed by an X-ray film processor (Xograph Compact X4).

Acknowledgements

We are grateful for generous support provided by the University of Oxford COVID-19 Research Response fund and its donors (BRD00230). C.V.R. is also a part of the COVID-19 mass spectrometry consortium.[49] Work in the C.V.R. laboratory is supported by a Medical Research Council (MRC) program grant (MR/N020413/1), a European Research Council Advanced Grant ENABLE (695511), and a Wellcome Trust Investigator Award (104633/Z/14/Z). C.A.L. is supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement GPCR-MS 836073. T.J.E. is supported by the Royal Society as a Royal Society Newton International Fellow.

Author Contributions

C.A.L, and T.J.E. designed the protein constructs; C.A.L., T.J.E., and J.R.B. expressed and purified protein constructs. C.A.L., T.J.E., and C.V.R., conceived the experiments; C.A.L. performed the experiments; T.J.E, J.R.B., and C.V.R supported the experiments; all authors co-wrote the paper.

Ethics declarations

The authors declare no competing interests. Carol Robinson provides consultancy services for OMass Therapeutics.

Figures and Legends

Figure 1. SARS-CoV-2 N protein exists as an ensemble of proteoforms. (A) Scheme depicting the full-length construct for expression in *E. coli*. The construct contains an N-terminal purification tag that is cleaved at the site indicated (red arrow). (B) SDS-PAGE of the N protein construct in (A) before and after tag cleavage. Three distinct protein bands (denoted 1-3) are observed in lanes labeled +tag and -tag. (C) The protein bands were subjected to in-gel digestion with trypsin followed by LC-MS based proteomics. (C) Histogram displaying percentage of peptides detected, relative to the total peptide count, across the five protein domains: N-terminal domain (NTD), RNA binding, Linker, Oligomerization, and C-terminal Domain (CTD). (D) Mass spectrum of the N protein after enrichment by size exclusion chromatography shows that, despite extensive purification, N protein exists as an ensemble of proteoforms. Four charge state distributions correspond to monomers and dimers of full-length N protein (red circles, MW 45,769 Da) and a proteoform of N protein (blue circles, MW 28,735 Da).

Figure 2. N protein undergoes autoproteolysis in a highly specific manner. (A) Mass spectrum of N protein after incubation with protease inhibitors for one week reveals the coexistence of five distinct charge state distributions corresponding to N proteoforms. The chemical composition of each proteoform was determined using top-down MS (B,C). (B) Charge-reduced mass spectrum resulting from electron-transfer dissociation (ETD) of the selected 9+ charge state at *m/z* 2616.79. (C) Mass spectrum of sequence ions for the same parent ion generated by higher-energy collision induced dissociation (HCD). (D) Scheme representing the composition of protein domains for the observed proteoforms as determined by top-down MS. Five distinct proteoforms are observed: N_{1-209}, N_{1-220}, N_{1-273}, $N_{156-419}$, and N_{1-392}. The exact site of cleavage, including the five residues flanking either side of each cleavage site, is indicated below each construct. (E) Scheme depicting the autoproteolytic cleavage of full-length N protein after virus entry and nucleocapsid disassembly.

Figure 3. The RNA sequence influences binding stoichiometry to N protein. (A) Mass spectrum of N_{1-209} after incubation with 4×-GAUGG RNA oligonucleotides in a molar ratio of 1:4. Two additional charge state distributions are observed that correspond to one and two RNA oligonucleotides bound to N_{1-209}. The mass spectrum at *m/z* >2700 was magnified 3× and offset for clarity (red trace). (B) Mass spectrum of N_{1-209} after incubation with 4×-GAGAA RNA oligonucleotides in a molar ratio of 1:4. One additional charge state distribution is observed that corresponds to one RNA oligonucleotide bound to N_{1-209}. The mass spectrum at *m/z* >2700 was magnified 3× and offset for clarity (red trace). (C) Mass spectrum of N_{FL} after incubation with 4×-GAUGG RNA oligonucleotides in a molar ratio of 1:4. Monomers and dimers of N_{FL} (red circles) and $N_{156-419}$ (blue circles) are observed. An additional peak series between 4300 and 5600 *m/z* corresponds to two 4×-GAUGG RNA oligonucleotides bound to N_{FL} dimer. The scheme in the inset of (C) depicts N_{FL} dimer bound to RNA as the functional unit of ribonucleoprotein assembly.

Figure 4. N$_{FL}$ and N proteoforms have distinct immunological roles. (A) Mass spectrum of N$_{FL}$ after incubation with a monoclonal antibody raised against the full-length N protein in a molar ratio of 1:1. We observe five charge state distributions that correspond to N$_{FL}$ monomers (centered at 13+), N$_{FL}$ dimers (centered at 19+), monomeric antibody (centered at 21+), antibody bound to one NFL (centered at 24+), and a population of antibody bound to N$_{FL}$ dimer (centered at 30+). (B) Mass spectrum of a mixture of N$_{1-209}$, N$_{1-220}$, and N$_{1-273}$ incubated with a monoclonal antibody in a molar ratio of 1:1:1:1. Charge state distributions are observed for antibody monomers and dimers. No binding to the antibody is observed for N proteoforms. This result was confirmed by immunoblot (see inset). SDS-PAGE shows that N$_{FL}$, N$_{1-209}$, N$_{1-220}$, N$_{1-273}$, and N$_{156-419}$ are detected in high abundance by Coomassie stain. The same proteins analyzed by immunoblot show that only N$_{FL}$ and N$_{156-419}$ are detected by the antibody.

Figure 5. N proteoforms directly interact with cyclophilin A. (A) Mass spectrum of N_{1-209} after incubation with cyclophilin A (CypA) in a 1:1 molar ratio. The mass spectrum at *m/z* >3000 was magnified 5✕ and offset for clarity (red trace). Three charge state distributions that correspond to a low population of homodimers of N_{1-209}, homodimers of CypA, and heterodimers of N_{1-209}-CypA. (B) Mass spectrum of N_{1-209} after incubation with CypA and cyclosporin A (CsA) in a molar ratio of 1:1:2. CypA preferentially binds CsA as demonstrated by the charge state distribution centered at 8+ which corresponds to the CypA-CsA complex. The mass spectrum at *m/z* >3500 was magnified 5✕ and offset (red trace) to highlight the exceedingly low abundance of N_{1-209}-CypA heterodimers that persist after CsA treatment. The scheme (inset of B) depicts CsA competitively binding to CypA and abolishing the N_{1-209}-CypA interaction.

Figure 6. Scheme depicting features of SARS-CoV-2 N protein during infection. N protein undergoes autoproteolysis in a highly specific fashion to produce N_{1-209}, N_{1-220}, N_{1-273}, $N_{156-419}$, and N_{1-392}. N_{FL} and N proteoforms bind RNA with a preference for structured RNA and N_{FL} dimers are likely functional unit of assembly in ribonucleoprotein complexes. Immunophilin CypA binds directly to N_{1-209} and N_{1-273}, but not N_{FL} or N_{1-220}, and the interaction can be inhibited through addition of cyclosporin A (CsA). $N_{156-419}$ is readily detectable by antibodies, suggesting a role in immune evasion via the formation of antigenic decoys.

Table 1. Deconvoluted and sequence masses of N_{FL} and N proteoforms.

Protein	Deconvoluted Mass[a] ± s.d. (Da)	Sequence Mass (Da)
N-terminal tagged N_{FL}	--	48,859.21
N_{FL}	45,769 ± 2	45,769.83
N_{FL} dimer	91,537 ± 2	91,539.66
$N_{156-419}$	28,735 ± 2	28,696.12
$N_{156-419}$ dimer	57,534 ± 1	57,392.24
N_{1-209}	22,611 ± 1	22,612.71
N_{1-220}	23,540 ± 0.3	23,541.73
N_{1-273}	29,402 ± 0.6	29,382.43
N_{1-392}	42,922 ± 1	42,918.74

[a] Determined using at least three adjacent charge states

Table 2. Deconvoluted and sequence masses for N proteoform complexes

Protein/Complex	Deconvoluted Mass[a] ± s.d. (Da)	Expected Mass (Da)
4x-GAUGG RNA	--	6,622.00
4x-GAGAA RNA	--	6,650.20
N_{1-209} + one 4x-GAUGG RNA	29,233 ± 0.4	29,234.71
N_{1-209} + two 4x-GAUGG RNA	35,917 ± 10	35,856.71
N_{1-209} + one 4x-GAGAA RNA	29,262 ± 1	29,262.91
N_{FL} dimer + two 4x-GAUGG RNA	105,131 ± 60	104,783.66
Antibody (monomer)	137,990 ± 125 145,193 ± 109	--
N_{FL} + antibody	183,931 ± 102	183,759 190,962
2 N_{FL} + antibody	229,984 ± 36	229,528 236,731
Cyclophilin A (CypA)	20,084 ± 0.4 20,220 ± 0.8	20,175.82
CypA dimers	40,373 ± 62	40,351.64
N_{1-209} + CypA	42,867 ± 15	42,696.71
N_{1-209} dimers	45,356 ± 15	45,225.42
Cyclosporin A	1,202.85 ± 0	1,202.63
CypA + CsA	21,287 ± 0.5	21,286.63

[a] Determined using at least three adjacent charge states

References

[1] Values taken from: https://coronavirus.jhu.edu

[2] McBride, R.; van Zyl, M.; Fielding, B. C. The Coronavirus Nucleocapsid Is a Multifunctional Protein. *Viruses*. **6**, 2991–3018 (2014).

[3] Okba, N. M. A. *et al*. Severe Acute Respiratory Syndrome Coronavirus 2−Specific Antibody Responses in Coronavirus Disease Patients. *Emerg. Infect. Dis*. **26**, 1478-1488 (2020).

[4] Burbelo, P. D. *et al*. Detection of Nucleocapsid Antibody to SARS-CoV-2 is More Sensitive than Antibody to Spike Protein in COVID-19 Patients. *J. Infect. Dis*. **222**, 206–213 (2020).

[5] Kang, S. *et al*, Crystal structure of SARS-CoV-2 nucleocapsid protein RNA binding domain reveals potential unique drug targeting sites. *Acta. Pharm. Sin. B*. **10**, 1228-1238 (2020).

[6] Zhou, R. J.; Zeng, R.; Brunn, A.; Lei, J. Structural characterization of the C-terminal domain of SARS-CoV-2 nucleocapsid protein. *Mol. Biomed*. **1**, 2 (2020).

[7] Musah, R. A. The HIV-1 nucleocapsid zinc finger protein as a target of antiretroviral therapy. *Curr. Top. Med. Chem*. **4**, 1605-1622 (2004).

[8] Kao, R. Y. *et al*. Identification of influenza A nucleoprotein as an antiviral target. *Nat. Biotechno*l. **28**, 600–605 (2010).

[9] Hung, H. C. *et al*. Development of an anti-influenza drug screening assay targeting nucleoproteins with tryptophan fluorescence quenching. *Anal. Chem*. **84**, 6391-6399 (2012).

[10] Lo, Y. S. Oligomerization of the carboxyl terminal domain of the human coronavirus 229E nucleocapsid protein. *FEBS Lett*. **587**, 120-127 (2013).

[11] Riva, L. *et al*. A large-scale drug repositioning survey for SARS-CoV-2 antivirals. Preprint at https://www.biorxiv.org/content/10.1101/2020.04.16.044016v1 (2020).

[12] Gordon, D. E. *et al*. A SARS-CoV-2 protein interaction map reveals targets for drug repurposing. *Nature*. **583**, 459–468 (2020).

[13] Chang, C. K. *et al*. Modular organization of SARS coronavirus nucleocapsid protein. *J. Biomed. Sci*. **13**, 59–72 (2006).

[14] Tilocca, B. *et al*. Comparative computational analysis of SARS-CoV-2 nucleocapsid protein epitopes in taxonomically related coronaviruses. *Microbes Infect*. **22**, 188–194 (2020).

[15] Ye, Q.; West, A. M. V.; Silletti, S.; Corbett, K. D. Architecture and self-assembly of the SARS-CoV-2 nucleocapsid protein. *Protein Sci*. **29**, 1890-1901 (2020).

[16] Smith, L. M.; Kelleher, N. L. Proteoform: a single term describing protein complexity. *Nat. Methods*. **10**, 186–187 (2013).

[17] Mark, J. *et al.* Unusual Lability of SARS Nucleocapsid Protein. *Biochem. Biophys. Res. Commun.* **377**, 429–433 (2008).

[18] Perler, F. B.; Xu, M. Q.; Paulus, H. Protein splicing and autoproteolysis mechanisms. *Curr. Opin. Chem. Biol.* **1**, 292-299 (1997).

[19] Kim, J. K. et al. Insight into autoproteolytic activation from the structure of cephalosporin acylase: A protein with two proteolytic chemistries. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 1732-1737 (2006).

[20] Johansson, D. G. A. et al. Protein Autoproteolysis: Conformational Strain Linked to the Rate of Peptide Cleavage by the pH Dependence of the N → O Acyl Shift Reaction. *J. Am. Chem. Soc.* **131**, 9475-9477 (2009).

[21] Huguet, R. *et al.* Proton transfer charge reduction enables high-throughput top-down analysis of large proteoforms. *Anal. Chem.* **91**, 15732-15739 (2019).

[22] Ives, A. N. *et al.* Using 10,000 Fragment Ions to Inform Scoring in Native Top-down Proteomics. *J. Am. Soc. Mass Spectrom.* **31**, 1398–1409 (2020).

[23] Haverland, N. A. *et al.* Defining Gas-Phase Fragmentation Propensities of Intact Proteins During Native Top-Down Mass Spectrometry. *J. Am. Soc. Mass Spectrom.* **28**, 1203–1215 (2017).

[24] Zeng, W. *et al.* Biochemical characterization of SARS-CoV-2 nucleocapsid protein. *Biochem. Biophys. Res. Commun.* **527**, 618–623 (2020).

[25] Tanwar, H.S. et al. The thermodynamics of Pr55Gag RNA interaction regulate the assembly of HIV. *PLoS Pathog.* **13**, e1006221 (2017).

[26] Kim, D. Y.; Firth, A. E.; Atasheva, S.; Frolova, E.I.; Frolov, I. Conservation of a packaging signal and the viral genome RNA packaging mechanism in alphavirus evolution. *J. Virol.* **85**, 8022–8036 (2011).

[27] Athmer, J. *et al.* Selective packaging in murine coronavirus promotes virulence by limiting type I interferon responses. *mBio.* **9**, e00272-18 (2018).

[28] Watashi, K.; Shimotohno, K. Cyclophilin and Viruses: Cyclophilin as a Cofactor for Viral Infection and Possible Anti-Viral Target. *Drug Target Insights.* **2**, 9–18 (2007).

[29] Ying, W. *et al.* Proteomic analysis on structural proteins of severe acute respiratory syndrome coronavirus. *Proteomics*, **4**, 492-504 (2004).

[30] Dimer, C. *et al.* Cell type-specific cleavage of nucleocapsid protein by effector capsases during SARS coronavirus infection. *J. Mol. Biol.* **376**, 23-34 (2008).

[31] Haller, A. A.; Stewart, S. R.; Semler, B. L. Attenuation stem-loop lesions in the 5' noncoding region of poliovirus RNA: neuronal cell-specific translation defects. *J. Virol.* **70**, 1467–1474 (1996).

[32] Stram, Y.; Kuzntzova, L. Inhibition of Viruses by RNA Interference. *Virus Genes.* **32**, 299–306 (2006).

[33] He, Y. *et al.* Mapping of Antigenic Sites on the Nucleocapsid Protein of the Severe Acute Respiratory Syndrome Coronavirus. *J. Clin. Microbiol.* **42**, 5309-5314 (2004).

[34] Li, S. *et al.* The Epitope Study on the SARS-CoV Nucleocapsid Protein. *Genomics Proteomics Bioinformatics.* **1**, 198–206 (2003).

[35] Shang, B. *et al.* Characterization and application of monoclonal antibodies against N protein of SARS-coronavirus. *Biochem. Biophys. Res. Commun.* **336**, 110–117 (2005).

[36] Lee, H. K. *et al.* Detection of antibodies against SARS-Coronavirus using recombinant truncated nucleocapsid proteins by ELISA. *J. Microbiol. Biotechnol.* **18**, 1717-1721 (2008).

[37] Bukreyev, A. *et al.* The Secreted Form of Respiratory Syncytial Virus G Glycoprotein Helps the Virus Evade Antibody-Mediated Restriction of Replication by Acting as an Antigen Decoy and through Effects on Fc Receptor-Bearing Leukocytes. *J. Virol.* **82**, 12191-12204 (2008).

[38] Zahno, M. L.; Bertoni. G. An Immunodominant Region of the Envelope Glycoprotein of Small Ruminant Lentiviruses May Function as Decoy Antigen. *Viruses.* **10**, 231-242 (2018).

[39] Kanto, T. *et al.* Density analysis of hepatitis C virus particle population in the circulation of infected hosts: implications for virus neutralization or persistence. *J. Hepatol.* **22**, 440-448 (1995).

[40] Chan, K. K. *et al.* Engineering human ACE2 to optimize binding to the spike protein of SARS coronavirus 2. *Science.* **369**, 1261-1265 (2020).

[41] Dawar, F. U. *et al.* Potential Role of Cyclophilin A in Regulating Cytokine Secretion. *J. Leukoc. Biol.* **102**, 989-992 (2017).

[42] Cron, R. Q. Coronavirus is the trigger, but the immune response is deadly. *Lancet.* **2**, 370-371 (2020).

[43] Ragab, D.; Eldin, H. S.; Taeimah, M.; Khattab, R.; Salem, R. The COVID-19 Cytokine Storm; What We Know So Far. *Front. Immunol.* **11**, 1446 (2020).

[44] Cong, Y.; Kriegenburg, F.; de Haan, C. A. M.; Reggiori, F. Coronavirus nucleocapsid proteins assembly constitutively in high molecular oligomers. *Sci. Rep.* **7**, 1-10 (2017).

[46] Bouhaddou, M. *et al.* The Global Phosphorylation Landscape of SARS-CoV-2 Infection. *Cell.* **182**, 1-28 (2020).

[47] Rial, D. V.; Ceccarelli, E. A. Removal of DnaK Contamination During Fusion Protein Purifications. *Protein Expr. Purif.* **25**, 503-507 (2002).

[48] DeHart, C. J.; Fellers, R. T.; Fornelli, L.; Kelleher, N. L.; Thomas, P. M. Bioinformatics Analysis of Top-Down Mass Spectrometry Data with ProSight Lite. *Methods Mol. Biol.* **1558**, 381–394 (2017).

[49] Struwe, W. *et al.* The COVID-19 MS Coalition-accelerating diagnostics, prognostics, and treatment. *The Lancet.* **395**, 1761-1762 (2020).