TITLE:

Attack behaviour in naïve Gyrfalcons is modelled by the same guidance law as in

Peregrines, but at a lower guidance gain

RUNNING TITLE:

Attack behaviour in Gyrfalcons

AUTHORS AND AFFILIATIONS:

Caroline H. Brighton[1], Katherine E. Chapman[2], Nicholas C. Fox[3], and Graham K. Taylor[1]

[1]Department of Zoology, University of Oxford, OX1 3SZ, UK.

[2]Centre for Research in Animal Behaviour, School of Psychology, University of Exeter, EX4

4QG, UK.

[3]Wingbeat Ltd, Carmarthen, SA33 5YL, UK.

CORRESPONIDNG AUTHORS:

graham.taylor@zoo.ox.ac.uk

caroline.brighton@zoo.ox.ac.uk

KEYWORDS: guidance law, optimal guidance, proportional navigation, proportional pursuit,

Falco rusticolus, *Falco peregrinus*

SUMMARY STATEMENT: Naïve Gyrfalcons attacking aerial targets are modelled by the

same proportional navigation guidance law as Peregrines, but with a lower navigation

constant that promotes tail-chasing rather than efficient interception.

ABSTRACT

The aerial hunting behaviours of birds are strongly influenced by their flight morphology and ecology, but little is known of how this variation relates to the behavioural algorithms guiding flight. Here we use onboard GPS loggers to record the attack trajectories of captive-bred Gyrfalcons (*Falco rusticolus*) during their maiden flights against robotic aerial targets, which we compare to existing flight data from Peregrines (*Falco peregrinus*). The attack trajectories of both species are modelled most economically by a proportional navigation guidance law, which commands turning in proportion to the angular rate of the line-of-sight to target, at a guidance gain N. However, Gyrfalcons operate at significantly lower values of N than Peregrines, producing slower turning and a longer path to intercept. Gyrfalcons are less agile and less manoeuvrable than Peregrines, but this physical constraint is insufficient to explain their lower guidance gain. On the other hand, lower values of N promote the tail-chasing behaviour that is typical of wild Gyrfalcons, and which apparently serves to tire their prey in a prolonged high-speed pursuit. Moreover, during close pursuit of fast evasive prey such as Ptarmigan (*Lagopus* spp.), proportional navigation will be less prone to being thrown off by erratic target manoeuvres if N is low. The fact that low-gain proportional navigation successfully models the maiden attack flights of Gyrfalcons suggests that this behavioural algorithm is embedded in a hardwired guidance loop, which we hypothesise is ancestral to the clade containing Gyrfalcons and Peregrines.

INTRODUCTION

Raptorial feeding is a complex mode of foraging behaviour, the success of which hinges on intercepting a target whose own success hinges on evading capture. Aerial pursuit in particular is one of the most challenging behaviours that organisms perform, but also one of the simplest to characterise. A dyadic interaction, for example, is minimally described by the trajectories of a pair of interacting particles representing the predator and its prey. This level of description lends itself to an algorithmic approach, in which a mathematical rule – in this case, a particular kind of behavioural algorithm known as a guidance law – is used to connect sensory input to motor output (Hein et al. 2020). As there are only a limited number of ways in which one particle can be steered to intercept another, this algorithmic approach lends itself in turn to a

rigorous comparative analysis of behaviour across different taxa, locomotor modes, and
spatiotemporal scales. Key research questions include: What sensory information is used to
guide interception, and how? For what function is the attacker's guidance algorithm optimized?
And how is this behavioural algorithm acquired? Here we address these questions for a sample
of naïve Gyrfalcons (*Falco rusticolus*), which are the largest of all falcons, and hence one of
the largest extant predators specialising in aerial interception.

Gyrfalcons are closely related to Peregrines (*Falco peregrinus*), whose attack trajectories are
well modelled (Brighton et al. 2017) by a guidance law called proportional navigation (PN). A
pure PN guidance law commands turning at an angular rate $\dot{\gamma} = N\dot{\lambda}$, where $\dot{\lambda}$ is the angular
rate of the attacker's line-of-sight to target, and where the guidance gain N is called the
navigation constant and assumed to be fixed within an attack. In contrast, the attack trajectories
of Harris' Hawks (*Parabuteo unicinctus*) are best modelled by a mixed PN+PP guidance law,
$\dot{\gamma} = N\dot{\lambda} - K\delta$, which combines a PN element with a proportional pursuit (PP) element, $-K\delta$,
commanding turning in proportion to the deviation angle δ between the attacker's velocity
vector and its line-of-sight to target (Brighton and Taylor 2019). In this case, a PN+PP guidance
law with guidance constants $N = 0.7$ and $K = 1.2$ s^{-1} modelled the observed flights more
closely than a PN or PP guidance law in which N or K was allowed to vary between flights
(Brighton and Taylor 2019). The PN guidance law of Peregrines promotes short-cutting
towards the eventual point of intercept, making it well suited to intercepting non-manoeuvring
targets in open environments (Brighton et al. 2017). In contrast, the PN+PP guidance law of
Harris' Hawks promotes tail-chasing directly after the target, making it better suited to close
pursuit through potentially cluttered environments (Brighton and Taylor 2019). Hawks
(Accipitridae) and falcons (Falconidae) are thought to have diverged >60 mya (Jarvis et al.
2014; Prum et al. 2015), so we would naturally expect the attack behaviour of Gyrfalcons to
be better modelled by the PN guidance law of Peregrines than by the PN+PP guidance law of
Harris' Hawks.

Peregrines and Gyrfalcons are both adapted to open environments, hunting mainly avian prey,
which may be knocked down in flight (Garber et al. 1993), struck on the ground before taking
flight (Bengtson 1971), or forced to the ground after a long chase (Cade 1982; Woodin 1967).
However, whereas Peregrines will often dive from altitude in a high-speed stoop (Cresswell
1996), Gyrfalcons rarely stoop in the wild and almost always hunt close to the ground (Cade

1982; Garber et al. 1993). Gyrfalcons are most often recorded performing low surprise attacks

initiated from a perch or from ridge soaring (Cade 1982; Garber et al. 1993; Potapov and Sale

2005; White 1991; White and Weeden 1966), but if not immediately successful then they will

commonly enter the prolonged tail-chase that is typical of this species (Cade 1982; Pennycuick

et al. 1994). Similar hunting behaviours are also observed in Peregrines (Cresswell 1996), but

the distinct speed advantage that Peregrines acquire when stooping (Mills et al. 2018; Mills et

al. 2019) appears to be lacking in wild Gyrfalcons. Such variation in hunting behaviour may

be expected to be associated with variation in the underlying guidance law.

Peregrine attacks are best modelled by a range of values of N (median: 2.6; 1^{st}, 3^{rd} quartiles:

1.5, 3.2) lower than the interval $3 \leq N \leq 5$ that is typical of missile applications, but close to

the optimum minimizing total steering effort in the classical linear-quadratic formulation of the

optimal guidance problem (Brighton et al. 2017). This theory predicts that $N = 3\, v_c / (v \cos \delta)$

is optimal for attacks on non-manoeuvring targets, where v_c is the speed at which the attacker

closes range on its target, and v is the attacker's groundspeed (Shneydor 1998; Siouris 2004).

In words, the optimal value of N is proportional to the ratio $v_c / (v \cos \delta)$, which expresses how

effectively the attacker closes range on its target (v_c, representing the difference between the

speed of the attacker's approach and the speed of the target's retreat) in relation to its own

motion towards the target ($v \cos \delta$, representing the speed of the attacker's approach). Hence,

whereas $N = 3$ is optimal for attacks on stationary targets (where $v_c = v \cos \delta$), $N < 3$ is

optimal for attacks on retreating targets (where $v_c < v \cos \delta$), at a value which depends on the

relative speeds of target and attacker. In a high-speed stoop, for instance, the target's speed

may be negligible compared to that of its attacker, such that $v_c / (v \cos \delta) \approx 1$ making $N \approx 3$

optimal (Mills et al. 2018; Mills et al. 2019). Conversely, in a prolonged tail chase in which

the target flees at a similar speed to the attacker, much lower values of $N < 3$ may be optimal.

We therefore hypothesise that Gyrfalcons will use PN guidance to intercept targets but will do

so using a lower value of N than Peregrines.

METHODS

To test these hypotheses, we use a combination of empirical measurements and computational

modelling to identify the guidance law used by naïve captive-bred Gyrfalcons chasing robotic

aerial targets. Our observations were recorded on the first attack flights that the birds had ever

made against aerial targets, and therefore reflect as closely as possible the hard-wired form of

the underlying guidance algorithm, with no prior opportunity for reinforcement learning.

Animals

We observed 23 naïve captive-bred Gyrfalcons, comprising 19 pure Gyrs (*F. rusticolus*) and 4

Gyr-Saker hybrids (7/8th *F. rusticolus* × 1/8th *F. cherrug*), chasing robotic aerial targets during

their first flight sessions. This sample contained only naïve first-year birds that had not

previously flown after aerial targets, except for during a single flight against a swung lure

immediately beforehand. This work has received approval from the Animal Welfare and

Ethical Review Board of the Department of Zoology, University of Oxford in accordance with

University policy on the use of protected animals for scientific research, permit no.

APA/1/5/ZOO/NASPA, and is considered not to pose any significant risk of causing pain,

suffering, damage or lasting harm to the animals.

Experimental protocol

Flight trials were carried out on open moorland at Watch Hill, Wealside Farm,

Northumberland, UK, in winds gusting from 4 to 7 m·s⁻¹. The birds were recorded chasing a

remotely piloted, ducted fan "Roprey" model with a food reward strapped to its dorsal surface

(Wingbeat Ltd, Carmarthen, Wales, UK; Fig. 1). Each bird carried a BT-Q1300 GPS receiver

(QStarz International, Taipei, Taiwan) logging position and groundspeed at 5 Hz. The GPS

receiver was carried dorsally on a Trackpack harness (Marshall Radio Telemetry, Salt Lake

City, UT, USA), giving a total load of 0.015 kg. An identical GPS logger was fixed inside the

body compartment of the Roprey, and the flights were filmed using a handheld Lumix DMC-

FZ1000 camera recording 4k video at 25 frames s⁻¹ (Panasonic Corporation, Osaka, Japan).

Each flight trial began as the falconer unhooded the bird on their fist, with the Roprey held ~20

m upwind. The Roprey was launched as soon as the bird took off, or sometimes just

beforehand; if not caught immediately (see e.g. Fig. 2A), it was flown through a series of

evasive turns (see e.g. Fig. 2B). The trial ended when the bird first intercepted the Roprey. If

the bird knocked the Roprey with its talons, then the pilot brought it safely to the ground; if the

bird bound to the Roprey, then the motor was cut, and an airbrake was deployed to prevent the

pair from drifting. In a few cases, the Roprey crash-landed before finally being intercepted by

the bird. After disregarding 3 flights in which the bird did not intercept the target, and another

7 flights in which the GPS logger was lost during the session, we were left with an initial

sample of 28 flights from 19 naïve Gyrfalcons.

Data synchronization and error analysis

We used the GPS time signal to synchronize the two GPS data streams to within ± 0.1 s, having

linearly interpolated a small number of dropped datapoints. We matched the synchronized GPS

data to the video with reference to take-off and landing, and used the video to identify the time

of first intercept. We then shifted the entire GPS trajectory of the bird so as to match its

estimated position at first intercept to that of the target (Fig. 2). This adjustment was necessary

to remove positional bias due to GPS receiver clock error, which is such that at the nominal

horizontal positioning accuracy of our GPS receivers (<3.0 m circular error probable), we

would expect 50% of position estimates of a co-located pair of receivers to be separated by

≥4.8 m under an isotropic gaussian error model. Receiver clock error varies slowly once settled,

so has no significant effect on the measurement of changes in position over short timescales:

in fact, we have shown empirically that our receivers have a precision of order 0.1 m for

changes in position occurring over intervals of order 10 s (Brighton et al. 2017). Even so, there

were 8 flights for which the discrepancy in the horizontal position estimates of the bird and

target at intercept exceeded the 95[th] percentile expected at the nominal accuracy of the

receivers, which indicates a higher-than-expected positioning error in one or both receivers. Of

these 8 flagged flights, 5 were the first flight that we recorded after receiver start-up at the

beginning of a logging session, which suggests that the receiver clock estimate had not been

given sufficient time to settle at the start of all 7 logging sessions (Fisher's exact test: $p =$

0.01). We therefore dropped these 8 flagged flights from the analysis, leaving a final sample

of 20 flights by $n = 13$ Gyrfalcons (Table S1). Among these 20 remaining flights, the

discrepancies in the position estimates of bird and target at intercept (median: 5.5 m; IQR: 8.0-

3.1 m) were distributed as expected at the nominal accuracy of the receivers (median: 4.8m;

IQR: 7.3-2.8 m), albeit with no extreme outliers.

Trajectory modelling

We simulated the birds' measured flight trajectories in MATLAB, by predicting their turn rate

$\dot{\gamma}(t)$ in response to the target's measured trajectory using a guidance law of the form:

$$\dot{\gamma}(t) = N\dot{\lambda}(t) - K\delta(t) \qquad (1)$$

where $\dot{\lambda}(t)$ is the angular rate of the line-of-sight to target, where $\delta(t)$ is the deviation angle

between the target and the attacker's velocity vector, and where N and K are constants. Angles

and angular rates are defined in vector form in the three-dimensional (3D) case, but can be

treated as scalars in the two-dimensional (2D) case. Eq. 1 necessarily ignores the effects of any

sensorimotor delay, on the basis that this is expected to be comparable to the ±0.1 s uncertainty

in the synchronization of the GPS data streams; see (Brighton and Taylor 2019). In the special

case that $K = 0$, Eq. 1 reduces to a pure PN guidance law, whereas in the special case that $N =$

0, Eq. 1 reduces to a pure PP guidance law. We refer to the general case in which $K \neq 0$ and

$N \neq 0$ as mixed PN+PP guidance. We used a forward Euler method to simulate each flight

under either PN, PP, or PN+PP guidance, initializing the simulation using the bird's measured

position and velocity at some given start point, and matching the bird's simulated flight speed

to its known groundspeed. The method is described further in our work with Peregrines and

Harris' Hawks (Shneydor 1998; Siouris 2004), the only difference being the 1/3000th s step

size that was used here to ensure that all parameter estimates were accurate at the reported

precision (for simulation code, see Supporting Data S1).

We used a Nelder-Mead simplex algorithm to find the value of N and/or K that minimized the

mean prediction error for each flight, defined as the mean absolute distance between the

measured and simulated trajectories. However, because the bird did not always start chasing

the target from the moment it was launched, it was necessary to select the start-point of the

simulation by reference to the data. Following the same approach used to model aerial chases

in Peregrines (Brighton et al. 2017), we therefore ran simulations beginning from all possible

start times ≥2.0 s before intercept, reporting the longest 2D simulation (up to a maximum of

20 s) for which the mean prediction error was ≤1.0% of the flight distance modelled. For the

3D case, we used an equivalent error tolerance of 1.2%, to preserve the same tolerance in each

dimension. This process of selecting the start-point of each simulation by reference to the

model's performance on the data is objective, but risks capitalizing on chance. We ensure that

our inferences are robust to the associated risk of overfitting by making contrastive inferences

between alternative guidance models and by focussing our inferences on the population
properties of the estimated parameters. Simulations that failed to model ≥2.0 s of flight at the
specified error tolerance are recorded as unsuccessful, and their parameter estimates are not
reported. Hence, because all of the successful simulations were fitted to within the same
specified error tolerance, the appropriate figure of merit for each simulation is the overall
distance or duration of flight modelled, expressed relative to the number of estimated guidance
parameters.
230
To facilitate direct comparison with our published results from experienced Peregrines
(Brighton et al. 2017), we reanalysed the Peregrine dataset using exactly the same methods
described above. After excluding 33 flights representing attacks on stationary ground targets,
and after dropping another 9 flights that were flagged as having lower-than-expected accuracy
at the point of intercept by the method above, this yielded a refined sub-sample of 13 flights
against aerial targets made by 4 experienced Peregrines (*F. peregrinus*). The mixed PN+PP
guidance law was not considered in the original analysis (Brighton et al. 2017), so was fitted
for the first time here. The only other difference from the original analysis was the refinement
of the integration step size used here. This had only a small effect on the simulated trajectories,
but sometimes caused a different start-point to be selected for the simulations, because of the
thresholding associated with finding the longest simulation fitted at the specified error
tolerance.
243
244
245 **RESULTS**
246
247 **The attack trajectories of naïve Gyrfalcons are well modelled under PN**
248
The 2D simulations under PN successfully modelled 18/20 of the flights by naïve Gyrfalcons,
fitting 1127 m and 111.0 s of flight at 1.0% error tolerance (Figs. 2A-B, 3, S1; Table S2). In
contrast, the PP simulations modelled only 14/20 flights successfully, fitting 824 m and 77.4 s
of flight, for the same number of estimated guidance parameters (Fig. 3A-B; Table S2). PN is
therefore much the better supported of the two pure guidance laws. In contrast, the PN+PP
simulations successfully modelled the terminal phase of all 20 flights, fitting 1358 m and 139.4
s of flight at 1.0% error tolerance (Fig. 3A-B; Table S2). The addition of a PP term therefore
increased the duration of flight fitted by a factor of 1.3 relative to PN, for a doubling in the

number of estimated guidance parameters. It follows that the Gyrfalcons' attack trajectories

are more economically modelled by PN than by PN+PP.

The implied overfitting of the PN+PP simulations is further evidenced by the observation that

the parameter estimates for K were inconsistently signed in the PN+PP simulations (median

K: 0.2 s^{-1}; 1st, 3rd, quartiles: -0.2, 0.7 s^{-1}; sign test: $p = 0.26$; $n = 20$ flights; Fig. 3D), despite

being positive in most of the successful PP simulations (median K: 0.9 s^{-1}; 1st, 3rd quartiles:

0.2, 1.7 s^{-1}; sign test: $p = 0.01$; $n = 14$ flights; Fig. 3D). This volatility in the sign of the

parameter estimates for K indicates that any turning behaviour that is not already modelled by

the PN element of the PN+PP simulations is not being consistently modelled by their PP

element either, so is likely to represent noise. Hence, given that $K > 0$ steers flight towards the

target, whereas $K < 0$ steers flight away from it, there is no evidence to indicate that a PP

element supplements PN in commanding steering towards the target. Conversely, the

observation that the parameter estimates for N were almost always positive in the successful

PN simulations (median N: 1.2; 1st, 3rd quartiles: 0.5, 1.4; sign test: $p < 0.001$; $n = 18$ flights;

Fig. 3C), and were similar in the PN+PP simulations (median N: 1.1; 1st, 3rd quartiles: 0.5, 1.5;

sign test: $p = 0.003$; $n = 20$ flights; Fig. 3C), provides strong positive evidence that steering

towards the target is indeed based on feeding back the line-of-sight rate $\dot{\lambda}$ rather than the

deviation angle δ.

Several of the successful PN simulations had parameter estimates $N \ll 1$, so predict little

turning behaviour at all (Fig. 4). These simulations provide no positive evidence for feedback

of $\dot{\lambda}$, but all 4 simulations with values of N falling below the 1st quartile correspond to nearly

straight sections of flight (Fig. 4D,K,L,N), for which there is little turning behaviour to explain,

and for which parameter estimation is therefore unreliable. Conversely, all 11 simulations with

values of N falling between the 1st and 3rd quartiles were from flights involving a substantial

amount of horizontal turning (Figs. 4C,E-I,M,O-R), which proportional feedback of $\dot{\lambda}$

successfully explains. To check whether PN guidance could also capture the altitudinal

component of the Gyrfalcon flights, we re-fitted all of the PN simulations in 3D (Fig. 5).

Although the number of flights that could be modelled successfully under PN dropped to 12/20

in 3D, comprising 734 m and 69.4 s of flight fitted at 1.2% error tolerance, the parameter

estimates in the $n = 12$ successfully-fitted 3D simulations (median N: 1.0; 1st, 3rd quartiles:

0.2, 1.4) were similar to those of the same flights in 2D (median N: 1.2; 1st, 3rd quartiles: 0.6,

1.8), and were not significantly higher or lower in either case (sign test: $p = 0.39$). We
therefore focus the remainder of our reporting on the 2D simulations.

PN models both short dashes and extended chases in their terminal phase

The 20 Gyrfalcon flights comprised 10 short dashes lasting from 3 to 7 s (Fig. 4A-I), and 10
extended chases lasting from 18 to 223 s (Fig. 4J-R). Typically, these short dashes correspond
to the birds' maiden flights (Table S1), whereas the extended chases correspond to their second
flights (Table S1), which is because the pilot made less of an attempt to evade capture on the
maiden flight. We found no evidence of any systematic increase or decrease in the parameter
estimates for N between the Gyrfalcons' maiden and second flights, for the small subsample
of $n = 6$ individuals for which paired data was available (sign test: $p = 0.69$). Of the 10 short
dashes, 7 were modelled successfully from target launch to intercept (Fig. 4A-G), whilst 2 were
modelled successfully for two-thirds of the total distance flown (Fig. 4H,I). For the 9/10
extended chases that were modelled successfully (Fig. 4J-R), the simulations only captured the
terminal phase of the attacks (median duration fitted: 8.2 s; first, third quartiles: 3.1, 12.3 s).
This nevertheless represents a substantial amount of flight fitted by distance (median distance
fitted: 86.9 m; first, third quartiles: 19.9 m, 146.5 m), because of the high speeds reached by
the end of a chase.

Naïve Gyrfalcons operate at lower navigation constants than experienced Peregrines

All 13 flights recorded previously from experienced Peregrines (Brighton et al. 2017) were
modelled successfully in 2D under PN or PN+PP (Fig. S2; Table S3), whereas only 11/13 of
these flights were modelled successfully under PP (Table S3). Moreover, whereas the PN
simulations fitted 1517 m and 99.6 s of flight at 1.0% error tolerance, the PP simulations fitted
only 927 m and 57.8 s of flight, for the same number of estimated guidance parameters (Table
S3). This confirms our earlier finding (Brighton et al. 2017) that PN is the better supported of
the two pure guidance laws in Peregrines. In comparison, the simulations under PN+PP fitted
1740 m and 131.2 s of flight successfully at 1.0% error tolerance (Table S3). The addition of a
PP term to the PN model for Peregrines therefore increased the duration of flight fitted by a
factor of 1.3, but for a doubling in the number of estimated guidance parameters. It follows that

the Peregrines' attack trajectories are more economically modelled by PN than by PN+PP,

which is the same conclusion as we have reached above for the Gyrfalcons.

The parameter estimates for N in the Gyrfalcons (median N: 1.2; 1st, 3rd quartiles: 0.5, 1.4)

were systematically lower (Fig. 6) than those in the Peregrines (median N: 2.8; 1st, 3rd quartiles:

1.6, 3.1). This difference was statistically significant (Mood's median test), both when treating

repeated measures from the same individual as independent datapoints ($\chi^2(1, n = 31) = 7.30$;

$p = 0.007$), and when analysing only the median values of N for each individual to eliminate

pseudo-replication ($\chi^2(1, n = 16) = 5.33; p = 0.02$). The statistically significant difference

in the distribution of N between the two species has a profound effect on the chase dynamics,

as can be shown by simulating the intercept trajectories that the Gyrfalcons would have

followed had they used the median value of N for the Peregrines, and *vice versa* (Fig. 7). It is

clear by inspection that the Gyrfalcons would have intercepted the target sooner had they

followed the trajectory commanded at $N = 2.8$ typical of Peregrines, and conversely that the

Peregrines would not have intercepted the target as soon as they did had they followed the

trajectory commanded at $N = 1.2$ typical of Gyrfalcons. This begs the question of why the

Gyrfalcons operated at such low values of N, which we tackle from various perspectives in the

Discussion, having first considered our key findings and their limitations.

DISCUSSION

Key scientific findings and their limitations

There are two key findings of this work: first, that naïve Gyrfalcons chase aerial targets as if

using the same PN guidance law found previously in experienced Peregrines (Brighton et al.

2017); and second, that they do so at a lower value of the navigation constant N (Fig. 3). The

first of these key findings is robustly confirmed by showing that PN models the data more

successfully than PP and more economically than PN+PP. It is plausible that the data might be

even better modelled by some alternative guidance law that we have not yet tested, but as there

are only a limited number of variables that can be fed back to command steering in a particle

model of interception, we think it likely that any such guidance law would be a variant of PN.

The second of these key findings is robustly confirmed using non-parametric statistics that

demonstrate a systematic difference in the navigation constant N between the Gyrfalcons

(median N: 1.2; 1st, 3rd quartiles: 0.5, 1.4) and the Peregrines (median N: 2.8; 1st, 3rd quartiles:

1.6, 3.1) for identically-analysed data collected using identical GPS loggers. Its primary

limitations are first that the experiments with Peregrines used a towed lure rather than a Roprey

model, and second that the sample of Peregrines comprised only $n = 4$ individuals. However,

as we have previously reported similar navigation constants (median: $N = 2.6$; 1st, 3rd

quartiles: 1.7, 3.3) in an independent sample of $n = 3$ experienced Peregrines attacking

stationary targets (Brighton et al. 2017), we are confident that this result can be generalised.

Technical limitations

It is important to emphasise that we have not identified a unique navigation constant N for

either species. Rather, we have identified distinct intervals within which their respective values

of N typically fall. Moreover, whilst we have reported a unique estimate of N for each flight,

a different parameter estimate would have been reported had a different start point been chosen

for the simulation. The selected start points are those which maximise the amount of flight

fitted at the specified error tolerance, and therefore represent an objective compromise between

goodness of fit and amount of data modelled. It is uncertain how much of the variability in our

parameter estimates for N is the result of measurement error as opposed to genuine behavioural

flexibility. In particular, the use of GPS loggers recording speed and position at 5 Hz precludes

perfect synchronization of the attacker and target trajectories and hence estimation of the

sensorimotor delay, so it is plausible that data collected at higher spatiotemporal precision

would show lower variability in the parameter estimates for N. Finally, our modelling only

accounts for the influence of wind only insofar as the simulated groundspeed is matched to that

of the measured groundspeed. The simulations therefore treat any distortions of the bird's track

due to the wind as if these were produced by the attacker's own steering commands, although

the effect of this is mitigated in close pursuit by the fact that the attacker and its target are

subject to the same wind. Nevertheless, gusting winds might perhaps explain some of the more

prominent inflections in the measured flight trajectories that are not captured by simulations

fitted over longer distances (e.g. Fig. 4J) or at higher wind speeds (e.g. Fig. 4R).

Physical and physiological constraints

For a given line-of-sight rate, the higher values of N in Peregrines are expected to command
turning at approximately twice the angular rate of the lower values of N in Gyrfalcons.
Minimum turn radius scales linearly with wing loading, W (Taylor and Thomas 2014), which
is ~6.0 kg m^{-2} in Gyrfalcons compared to ~5.3 kg m^{-2} in Peregrines (Pennycuick et al. 1994).
Gyrfalcons are therefore expected to be less manoeuvrable than Peregrines, with a ~13% larger
minimum turn radius. They are also expected to be less agile, because flight speed scales as
$W^{1/2}$ (Taylor and Thomas 2014), such that maximum turn rate scales as $W^{-1/2}$, and is
therefore expected to be ~6% lower in Gyrfalcons than Peregrines. In principle, this physical
constraint could force the navigation constant N to be lower in Gyrfalcons, but the expected
difference in their agility is too small to explain the twofold difference in N that we observe.
Furthermore, there is no evidence that physical constraints on turning influenced the shape of
the recorded trajectories. Rather than turning as tightly as possible on a circular arc, the birds
instead followed a curved trajectory of increasing or decreasing radius, characteristic of the
time history of turning commanded under PN guidance at higher or lower values of N,
respectively (Fig. 4). The observed flight trajectories are similarly inconsistent with an old
hypothesis proposed by Tucker (Tucker 2000), who argued that the curved attack trajectories
of falcons could be generated by steering so as to hold the target's image on the laterally
directed central fovea of the left or right eye whilst holding the head in line with the body
(Tucker et al. 2000). This constraint is expected to produce a trajectory in which the deviation
angle δ between the attacker's velocity and its line-of-sight to target is held constant at
approximately $\pm\,40°$, but there is no evidence of this from the data (see Fig. 2; see also (Kane
and Zamani 2014).
410
411 **A functional account of the observed variation in navigation constant**
412
In principle, the variation in N within and between species (Fig. 6) might be explained as a
behavioural response under linear-quadratic optimal guidance theory (Shneydor 1998; Siouris
2004), which predicts an optimum value of $N = 3\,v_c/(v\cos\delta)$ for attacks on non-
manoeuvring targets minimizing overall steering effort. In practice, the median values of the
ratio $v_c/(v\cos\delta)$ for each flight did not differ significantly between the two species
($\chi^2(1, n = 31) = 1.55$; $p = 0.21$), and nor were they significantly related to N in a bisquare
robust regression controlling for species ($t(28) = 0.46$; $p = 0.69$). We therefore find no
evidence that the statistically significant difference in the parameter estimates for N between

the Peregrines and Gyrfalcons reflects a direct functional response to variation in the rate at
which they closed range on the target relative to their own approach speed. This conclusion is
based on classical theory for non-manoeuvring targets, and therefore takes no account of the
target's manoeuvres. Nevertheless, the tortuosity of the lure's path, defined as its overall path
length divided by the straight line distance from start to finish, was similar in the experiments
with Peregrines (median tortuosity: 1.6; 1st, 3rd quartiles: 1.3, 2.4) and Gyrfalcons (median
tortuosity: 1.7; 1st, 3rd quartiles: 1.3, 2.5), so we conclude that differences in target manoeuvres
are unlikely to explain the variation in N that we observed between species. Even so, an
identical target manoeuvre will produce a greater angular change in the line-of-sight vector the
closer the attacker is to its target, thereby demanding a higher rate of turning under PN. This is
important, because whereas the Gyrfalcons took off together with the target at a range of ~20
m, the Peregrines were launched separately and sometimes at a range of >100 m. When
operating at close range, PN guidance is less prone to being thrown off by erratic target
manoeuvres if N is low (Brighton and Taylor 2019), so it is plausible that the lower values of
N found in the Gyrfalcons might reflect a behavioural response to their proximity to the target
at the initiation of the attack.

An adaptive account of the observed variation in navigation constant

A complementary adaptive argument can also be made at species level. Whereas Peregrines
have a very flexible diet (Cade 1982), Gyrfalcons depend heavily on a single prey type, with
Ptarmigan (*Lagopus* spp.) comprising 74% of the catch on average across 17 studies recording
66,726 individual prey items over most of the Gyrfalcon's range (Nielsen and Cade 2017).
Ptarmigan have a high wing loading of ~9.7 kg m^{-2} (Greenewalt 1962), which is ~62% higher
than that of a Gyrfalcon. They are therefore intrinsically fast fliers, with large flight muscles
that provide rapid acceleration in an explosive take-off, and short wings that make them less
well adapted to sustained aerobic flight (Pennycuick et al. 1994). The prolonged tail-chasing
behaviour that is typical of Gyrfalcons (Cade 1982) is therefore thought to serve to tire their
prey and allow capture (Pennycuick et al. 1994). Such behaviour is promoted by the low values
of N found in Gyrfalcons (Fig. 7), because PN with a low navigation constant $N \approx 1$
commands turning at an angular rate $\dot{\gamma}$ approximately equal to the line-of-sight rate $\dot{\lambda}$. Once a
tail chase has been initiated, this will tend to keep the attacker flying behind its target, and will
not tend to be thrown off too far by the erratic jinking manoeuvres that are characteristic of the

evasive flight of Ptarmigan and many other prey (Mills et al. 2018; Mills et al. 2019). The low

values of N found in Gyrfalcons therefore make sense from an adaptationist perspective if the

function of their PN guidance is to pursue the prey doggedly until it tires. This is in contrast to

the higher values of N found in Peregrines, which make sense if the function of their PN

guidance is to exploit the speed and manoeuvrability acquired through stooping in order to

intercept prey quickly and efficiently (Mills et al. 2018; Mills et al. 2019).

Evolutionary and ontogenetic implications

We found no evidence that the value of N changed systematically between the Gyrfalcons'

first and second flights, but it is reasonable to assume that they would have learned to tune their

guidance over longer timescales. For example, although wild Gyrfalcons do not usually stoop,

falconers commonly train captive birds to do so (Tucker et al. 1998), which implies a degree

of flexibility in their guidance. Likewise, whereas naïve Gyrfalcons tend to fly directly at their

prey, experienced birds seem to anticipate their prey's behaviour. Even so, our finding that the

maiden attack flights of naïve Gyrfalcons are well modelled under PN guidance strongly

suggests that this behavioural algorithm is embedded in a hardwired guidance loop. Moreover,

the fact that the same form of guidance law models the attack flights of experienced Peregrines

is consistent with the hypothesis that PN guidance is ancestral to the clade comprising the

Peregrines and Hierofalcons, of which the Gyrfalcon is a member (Wink 2018). Formal

confirmation of this hypothesis would require the same behavioural algorithm to be identified

in another Hierofalcon, or in a close outgroup such as the Merlin or Hobby falcons, both of

which specialise in aerial pursuit. Because PN guidance commands turning in proportion to the

line-of-sight rate of the target, which is defined in an inertial frame of reference, we would

expect any such birds using PN to share a common sensorimotor architecture that fuses sensory

input from the visual and vestibular systems to obtain the line-of-sight rate, and that uses the

resulting signal to generate motor commands to the flight muscles controlling the wings and

tail. Alternatively, it is possible that the distant landscape might serve as a visual proxy for the

inertial frame of reference in aerial hunters, with target motion being measured directly with

respect to the visual background; see also (Kane and Zamani 2014). It would therefore be of

particular interest to test whether PN guidance also models the attack behaviours of the more

distantly related Kestrels, given their specialism on terrestrial prey with no distant visual

background against which to assess prey motion.

Acknowledgements

We thank Jimmy Robinson and Tom Spink for falconry work, plus their interns, Adam Vacek, Sander Gielen, and Daniel Clark. We thank Remy Van Wijk and Matt Aggett for piloting the Roprey. We thank Barbro Fox for her hospitality, and Sofía Miñano Gonzalez for valuable comments on the manuscript.

Author contributions

CB, GT and NF conceived the study. All authors contributed to field experiments. CB analysed the GPS data. CB and GT performed statistical analysis and wrote the paper. All authors commented on and approved the final version of the manuscript.

Competing interests

CB, KC and GT declare no competing interests. NF is CEO of Wingbeat Ltd, which supplied and piloted the Roprey in this study.

Funding

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant Agreement No. 682501) to GKT.

References

References

Bengtson SA. 1971. Hunting methods and choice of prey of gyrfalcons falco-rusticolus at
myvatn in northeast iceland. Ibis. 113(4):468-&.

Brighton CH, Taylor GK. 2019. Hawks steer attacks using a guidance system tuned for close
pursuit of erratically manoeuvring targets. Nat Commun. 10.

Brighton CH, Thomas ALR, Taylor GK. 2017. Terminal attack trajectories of peregrine falcons
are described by the proportional navigation guidance law of missiles. P Natl Acad Sci
USA. 114(51):13495-13500.

Cade TJ. 1982. The falcons of the world. Ithaca, New York, USA.: Cornell University Press.

Cresswell W. 1996. Surprise as a winter hunting strategy in sparrowhawks accipiter nisus,
peregrines falco peregrinus and merlins f-columbarius. Ibis. 138(4):684-692.

Garber CS, Mutch BD, Platt S. 1993. Observations of wintering gyrfalcons (falco-rusticolus)
hunting sage grouse (centrocercus-urophasianus) in wyoming and montana USA. J
Raptor Res. 27(3):169-171.

Greenewalt CH. 1962. Dimensional relationships for flying animals. Smithsonian
Miscellaneous Collections Smithsonian Institution. 144.

Hein AM, Altshuler DL, Cade DE, Liao JC, Martin BT, Taylor GK. 2020. An algorithmic approach
to natural behavior. Curr Biol. 30(11):R663-R675.

Jarvis ED, Mirarab S, Aberer AJ, Li B, Houde P, Li C, Ho SYW, Faircloth BC, Nabholz B, Howard
JT et al. 2014. Whole-genome analyses resolve early branches in the tree of life of
modern birds. Science. 346(6215):1320-1331.

Kane SA, Zamani M. 2014. Falcons pursue prey using visual motion cues: New perspectives
from animal-borne cameras. J Exp Biol. 217(2):225-234.

Mills R, Hildenbrandt H, Taylor GK, Hemelrijk CK. 2018. Physics-based simulations of aerial
attacks by peregrine falcons reveal that stooping at high speed maximizes catch
success against agile prey. Plos Comput Biol. 14(4):e1006044.

Mills R, Taylor GK, Hemelrijk CK. 2019. Sexual size dimorphism, prey morphology and catch
success in relation to flight mechanics in the peregrine falcon: A simulation study. J
Avian Biol. 50(3).

Nielsen OK, Cade TJ. 2017. Gyrfalcon and ptarmigan predator-prey relationship. In: Anderson
DL, McClure CJW, Franke A, editors. Applied raptor ecology: Essentials from gyrfalcon
research. Boise, Idaho, USA: The Peregrine Fund.

Pennycuick CJ, Fuller MR, Oar JJ, Kirkpatrick SJ. 1994. Falcon versus grouse - flight adaptations
of a predator and its prey. J Avian Biol. 25(1):39-49.

Potapov E, Sale R. 2005. The gyrfalcon. Yale University Press.

Prum RO, Berv JS, Dornburg A, Field DJ, Townsend JP, Lemmon EM, Lemmon AR. 2015. A
comprehensive phylogeny of birds (aves) using targeted next-generation DNA
sequencing. Nature. 526(7574):569-U247.

Shneydor NA. 1998. Missile guidance and pursuit: Kinematics, dynamics and control.
Woodhead Publishing Limited.

Siouris GM. 2004. Missile guidance and control systems. Springer-Verlag.

Taylor GK, Thomas ALR. 2014. Evolutionary biomechanics: Selection, phylogeny, and
constraint. Ox Ecol Ev.1-152.

Tucker VA. 2000. The deep fovea, sideways vision and spiral flight paths in raptors. J Exp Biol.
203(24):3745-3754.

Tucker VA, Cade TJ, Tucker AE. 1998. Diving speeds and angles of a gyrfalcon (falco rusticolus).
J Exp Biol. 201(13):2061-2070.

Tucker VA, Tucker AE, Akers K, Enderson HJ. 2000. Curved flight paths and sideways vision in
peregrine falcons (falco peregrinus). J Exp Biol. 203(24):3755-3763.

White CM, and Nelson, R. W. 1991. Hunting range and strategies of tundra breeding peregrine
and gyrfalcons observed from a helicopter. J Raptor Res. 25:49-62.

White CM, Weeden RB. 1966. Hunting methods of gyrfalcons and behavior of their prey
(ptarmigan). Condor. 68(5):517-&.

Wink M. 2018. Phylogeny of falconidae and phylogeography of peregrine falcons. Ornis
Hungarica. 26(2):27-37.

Woodin N. 1967. Observations on gyrfalcons (falco rusticolus) breeding near lake myvatn,
iceland. Raptor Research. 14(4):97-124.

Figures

Figure 1. Cropped frame from video of a typical chase involving a Gyrfalcon and a "Rokarrowan" Roprey model.

Note the proximity of the attacker to its target, and their similar bank angles, which are characteristic of the tail-

chasing behaviour that we observed.

577
578
Figure 2. Two-dimensional (2D) GPS trajectories for: (A) the entirety of a short dash; and (B) the entirety of an
extended chase, showing the lines-of-sight (cyan lines) between the Gyrfalcon (blue points) and Roprey (magenta
points) at each sample point. Note the small discrepancy in the estimated position of target and attacker at the
point of intercept (enlarged sample points), expected because of the positioning error associated with GPS
receivers (see Methods). (C,D) The terminal phase of the same two flights (see Fig. 4G,M for modelling), after
shifting the attacker's trajectory to correct for this positioning error. Gridlines at 10m spacing.

Figure 3. Box-and-whisker plots comparing model fits and parameter estimates for the 2D simulations of 20 flights from $n = 13$ Gyrfalcons under proportional pursuit (PP), proportional navigation (PN), and mixed (PN+PP) guidance. (A) Relative error of simulation, showing either the relative error for the longest simulation lasting ≥ 2 s that met the 1.0% error tolerance threshold for each flight, or if no simulation met this threshold, then the minimum relative error achieved on any simulation lasting ≥ 2 s. Note that whilst the PN+PP simulations fit the flights more closely than either PN or PP, they are almost certainly overfitted (see Results). (B,C) Parameter estimates for the guidance constants N and K for all successfully modelled flights. Note the variable sign of the parameter estimates for K under PN+PP, which confirms that these simulations are overfitted (one outlier for PN+PP not shown). The centre line of each box denotes the median for all flights; the lower and upper bounds of the box denote the 1st and 3rd quartiles; crosses indicate outliers falling >1.5 times the interquartile range beyond the 1st or 3rd quartile; whiskers extend to the farthest datapoints not treated as outliers.

Figure 4. Two-dimensional (2D) attack trajectories for the 18/20 flights from $n = 13$ Gyrfalcons that were
successfully modelled under proportional navigation (PN) guidance. Panels show the measured trajectories of the
target (magenta points) and attacker (blue points), overlain with the longest simulation fitted to within 1.0% error
tolerance (blue lines) in 2D. The corresponding parameter estimate for N is displayed on each plot. Note that
among the 9 short dashes (panels A-I), 7 flights (panels A-G) are modelled in their entirety from target launch to
intercept; the other 9 flights (panels J-R) each correspond to the terminal phase of an extended chase. Simulations
with values of N falling beneath the 1st quartile ($N < 0.5$) coincide with nearly straight sections of flight (panels
D,K,L,N), for which parameter estimation is unreliable. Simulations with values of N falling between the 1st and
3rd quartiles ($0.5 \le N \le 1.4$) involve a substantial amount of turning that the model successfully explains (panels
C,E-I,M,O-R). Black arrows display mean wind direction; double headed arrows correspond to wind speeds >20
km h^{-1}; gridlines at 10m spacing. See Fig. S1 for the remaining 2/20 flights that were not successfully modelled
under PN.

Figure 5. Three-dimensional (3D) attack trajectories for the subset of 6 Gyrfalcon flights involving the greatest
altitudinal change among the 12/20 that were successfully modelled under proportional navigation (PN) guidance.
Panels show the measured trajectories of the target (magenta points) and attacker (blue points), overlain with the
longest simulation fitted to within 1.2% error tolerance (blue lines) in 3D. The corresponding parameter estimate
for N is displayed on each plot. Panels B, C, F, G correspond to short dashes for which almost the entire flight
was modelled from target launch to intercept. Panel letters match those in Fig. 4; gridlines at 10m spacing.

Figure 6 Box-and-whisker plots comparing parameter estimates for N in proportional navigation (PN) guidance
models fitted independently to the 13 attack flights from $n = 4$ Peregrines, and 18 attack flights from $n = 13$
naïve Gyrfalcons; all in pursuit of manoeuvring targets. The centre line of each box denotes the median for all
flights; the lower and upper bounds of the box denote the 1st and 3rd quartiles; crosses indicate outliers falling >1.5
times the interquartile range beyond the 1st or 3rd quartile (one extreme outlier for the Peregrines not shown);
whiskers extend to the farthest datapoints not treated as outliers.

Figure 7. Effect of the navigation constant N on the dynamics of proportional navigation (PN) guidance. Upper
row: selection of 4 successfully modelled Gyrfalcon flights involving substantial turning, showing the measured
trajectory of the target (magenta dots) and attacker (blue dots) overlain with the best-fitting trajectory under PN
guidance at the value of N displayed on the panel (blue line), and with the trajectory that would have been followed
for the same initial conditions and target motion at the median value of $N = 2.8$ for Peregrines (green line). Green
dot shows the predicted point of intercept had the Gyrfalcon used the median value of N for Peregrines; note that
this is always sooner than the actual point of intercept (magenta dot). Lower row: selection of 4 successfully
modelled Peregrine flights involving substantial turning, showing the measured trajectory of the target (magenta
dots) and attacker (blue dots) overlain with the best-fitting trajectory under PN guidance at the value of N
displayed on the panel (lilac line), and with the trajectory that would have been followed for the same initial
conditions and target motion at the median value of $N = 1.2$ for Gyrfalcons (orange line). Lilac dot shows the
actual point of intercept; black cross shows the predicted position of the bird had the Peregrine used the median
value of N for Gyrfalcons; note that this is always at some significant distance from the target. Panel letters match
those in Figs. 3, S2; gridlines at 10 m spacing.

644