The guidance and adhesion protein FLRT2 dimerizes *in cis* via dual Small-X$_3$-Small transmembrane motifs

Verity Jackson[#,1], Julia Hermann[#,1,6], Christopher J. Tynan[#,2], Daniel J. Rolfe[2], Antreas C. Kalli[3], Anna L. Duncan[1], E. Yvonne Jones[4], Mark S. P. Sansom[1], Marisa L. Martin-Fernandez*[,2], Elena Seiradake*[,1], Matthieu Chavent*[,5]

#These authors contributed equally to this study

1- Department of Biochemistry, University of Oxford, South Parks Road, Oxford, OX1 5RJ, United Kingdom

2- Central Laser Facility, Research Complex at Harwell Science & Technology Facilities Council, Harwell Campus, Didcot, United Kingdom

3- Department of Discovery and Translational Science, Institute of Cardiovascular and Metabolic Medicine, School of Medicine and Astbury Center for Structural Molecular Biology, University of Leeds, United Kingdom

4- Division of Structural Biology, Wellcome Centre for Human Genetics, University of Oxford, Oxford, United Kingdom

5- Institut de Pharmacologie et Biologie Structurale, IPBS, Université de Toulouse, CNRS, UPS, 205 route de Narbonne, 31400, Toulouse, France

6- Present address: German Cancer Research Center (DKFZ), Im Neuenheimer Feld 581, 69120 Heidelberg, Germany

* Correspondence and requests for materials should be addressed to E. S. (email: elena.seiradake@bioch.ox.ac.uk), to M. L. M.F. (email: marisa.martin-fernandez@stfc.ac.uk), or to M. C. (email: matthieu.chavent@ipbs.fr)

Abstract

Fibronectin Leucine-rich Repeat Transmembrane (FLRT 1-3) proteins are a family of broadly expressed single-spanning transmembrane receptors that play key roles in development. Their extracellular domains mediate homotypic cell-cell adhesion and heterotypic protein interactions with other receptors to regulate synapse development and cell guidance. These *in trans* FLRT interactions determine the formation of signaling complexes of varying complexity and function. Whether FLRTs also interact *in cis* remains unknown. Here, we reveal two dimerization motifs in the FLRT2 transmembrane helix. Molecular dynamics simulations and single particle tracking experiments show that these 'Small-X$_3$-Small' motifs synergize with a third dimerization motif encoded in the extracellular domain to permit the *cis* association and diffusion patterns of FLRT2 on cells. The results point to a competitive switching mechanism between *in cis* and *in trans* interactions which suggests that homotypic FLRT interaction mirrors the functionalities of classic adhesion molecules.

Introduction

Fibronectin Leucine-rich Repeat Transmembrane (FLRT) proteins are a family of cell adhesion molecules (CAMs) that are broadly expressed during vertebrate development[1,2]. FLRTs are unusual CAMs as they perform both cell adhesive and repulsive functions, leading to their definition as Repellent CAMs (ReCAMs)[3,4]. In neurons, FLRTs act as repulsive guidance cues during cortical cell migration[3,5-7], whereby they play key role in cortical folding[8] and as adhesion molecules in synaptic complexes[9,10]. Adhesive FLRT functions are elicited by homotypic binding[1,2] or by binding to the G-protein coupled receptor Latrophilin (Lphn 1-3)[6,9,11,12] on opposing cells while cell repulsion results from interaction with Uncoordinated-5 (Unc5A-D)[4,11]. FLRT also interacts with Unc5 *in cis* to regulate Lphn-mediated adhesion, at least *in vitro*[7]. In migrating neurons, FLRT cooperates with the Lphn-binding receptor Teneurin to form a ternary trans-synaptic complex that mediates cell repulsion[13], while the three proteins also function in promoting synapsing[14]. Thus, FLRT acts in a context-dependent manner to determine the formation of different higher order cell-guidance signalling complexes and regulate brain development[15]. Here we ask whether FLRT forms homotypic *cis* complexes and how this may modulate *cis* and *trans* interactions with other partners.

FLRTs share a common architecture[16] (**Fig. 1A**) beginning with an N-terminal Leucine-Rich Repeat (LRR) extracellular domain, which contains a concave surface on which both FLRT and Lphn bind[3,6]. Unc5 binds to an adjacent surface on the LRR domain, which is compatible at least with Lphn-binding[7]. The LRR domain is linked to a type III fibronectin (FN) domain which then leads into the single-spanning transmembrane (TM) domain and a ~100 amino acid long intracellular domain (ICD) of unknown structure. FLRT2 TM domains contained two consecutives "Small-X_3-Small" motifs[17-19] (**Fig. 1B**) which may promote FLRT interactions *in cis*. Indeed, this motif play fundamental roles in other single-spanning transmembrane receptors, including epidermal growth factor receptor (EGFR), fibroblast growth factor receptor (FGFR), integrins, and EphA receptors[20-26].

Characterizing dynamic membrane protein structures is challenging[18,27,28]. However, the recent developments in molecular dynamics (MD) simulations[29] have provided a powerful tool to study membrane receptors and their interactions[30-35]. Coarse-grained molecular dynamics (CG-MD) simulations have emerged as the method of choice to explore the self-assembly of TMs[36-38] and have been exhaustively validated[39-41]. Combining MD simulations with well-established experimental assays such as Single Molecule Tracking (SMT) microscopy[42,43], which tracks the clustering of receptors[44-49] on live cells, is a powerful approach to provide new insights into the dynamic assemblies of cell receptors[21,33,50].

Here, we use molecular dynamics simulations and live cell SMT experiments to reveal how FLRT2 dimerizes *in cis* via two Small-X_3-Small motifs. Unexpectedly, these motifs work synergistically with the extracellular dimerization motif in the ligand-

82 binding domain[3] to produce FLRT-FLRT association. The results suggest a bipartite

83 structural mechanism that underlies the diverse functions of FLRT, and a competitive

84 mechanism for *in cis* versus *in trans* binding via the extracellular domain.

Figure 1: Dimerization of FLRT2 TM domains. A- Schematic of FLRT proteins engaging *in trans* and potentially also *in cis* interactions. **B-**Sequence of the FLRT2 TM helix. The two Small-X_3-Small motifs, key residues for the formation of the helix/helix interface highlighted by the CG-MD simulations, are colored in green and red. Two constructs were used as inputs for MD simulations: the core TM helix of 24 residues (TM_{24}) and an extended version with the four most N- and C-terminal residues (TM_{32}). **C-** The CG-MD protocol to assess TM helix interactions. The two helices are positioned 60 Å apart and diffuse freely in the membrane. The colored bars show, for each simulation, the distance in between the two TM helices as a function of the time. **D-** The TM contact bars of the 30 simulations for the TM_{32} helices in a membrane of 80% POPC and 20% cholesterol. Details for all the simulations can be found in Sup. Fig. 1 and Sup. Fig. 2.

Results

FLRT2 TM dimerization involves two Small-X_3-Small motifs

As no structural information exists for the FLRT2 TM domain, we have used

secondary structure prediction tools (see Methods) to predict the membrane-

embedded helical region of FLRT2 (**Fig. 1B**). We obtained 24 residues as the core TM

helix (denoted TM_{24}). This length is consistent with the average length for a plasma

membrane-spanning TM helix[51]. We also extended the helical segment with four N-

and C-terminal residues, which were modelled as coils (denoted TM_{32}).

We then performed multiple runs of coarse-grained molecular dynamics (CG-

MD) to model the associations of both TM_{24} and TM_{32} in three different types of

membrane: POPC, DPPC, and a lipid mixture of 80% POPC and 20% cholesterol

(**Sup. Table 1**). For each of these three lipid compositions, we positioned the two

helices 60 Å apart allowing them to diffuse freely until there is an encounter which

predominantly leads to the formation of a stable helix dimer (**Fig. 1D and Sup. Fig. 1**

and 2). The TM helices interacted through a network of residues distributed along each
peptide. Among these residues, we identified two consecutive Small-X$_3$-Small motifs
known to favor TM interactions[52,53]: A$_{544}$-X3- G$_{548}$ and G$_{545}$-X3-G$_{549}$ (**Sup. Fig. 3**).

Figure 2: CG-MD simulations uncover two Small-X$_3$-Small motifs as key areas for the dimerization of FLRT2 TM domains. A- TM$_{32}$ helix dimer structural populations for different membrane compositions (see Sup. Fig. 5 to 7 for individual simulations). A positive value for the crossing angle corresponds to a left-handed (LH) dimer, and a negative value to a right-handed (RH) dimer (details for TM$_{24}$ systems are presented in Sup. Fig. 4 and Sup. Fig. 8 to 10). The simulations have highlighted two Right Handed conformations (RH1 and RH2) and one Left Handed (LH). RH1 is defined with a crossing angle between -18° and 0°, RH2 between -36° and -18°, and LH between 0° and 18°. **B-** The three main representative structures for each crossing angle peak with A$_{544}$-G$_{548}$ dimerization motif in green, and A$_{545}$-G$_{549}$ dimerization motif in red. For each conformation (RH1, RH2 and LH), averaged TM contact matrix extracted from simulations of TM$_{32}$ in 80% DPPC and 20% of cholesterol highlight the TM dimerization via the A$_{544}$-G$_{548}$ motif (RH2), G$_{545}$-G$_{549}$ motif (RH1) or a combination of both (LH).

We next performed crossing angle analysis to analyze the geometry of the TM
helices[54,55]. For both the TM$_{24}$ and TM$_{32}$ models, this revealed three dimer populations
(**Fig. 2A and Sup. Fig. 4**): two main right-handed populations with average crossing
angles of approximately -27° (RH1) and -9° (RH2) and one minor left-handed
population with an average crossing angle of around +9° (LH). For each crossing angle
population, a clear interface was identified (**Fig. 2B**). RH1 conformations were driven

by interactions around the G_{545}-X_3-G_{549} motif while RH2 conformations were centered
on the A_{544}-X_3-G_{548} motif. The LH population was constituted by peptides interacting
via both motifs.
To obtain a more detailed view of the dynamical TM dimer association, we
plotted the helix crossing angle against the distance between the two Small-X_3-Small
motifs. This revealed several sub-populations associated with each crossing angle
peak (Fig. 2A). Notably, membrane lipid composition appeared to modulate this
dynamical equilibrium (Fig. 2A and Sup. Fig. 4) as seen in other studies[56-60].
Furthermore, for each membrane composition, the juxtamembrane regions appeared
to modulate TM dimer formation (Fig. 2A and Sup. Fig. 4) as previously reported[61].
Plotting individual simulations (Sup. Fig. 5 to 10) suggested that the RH1
conformation can exchange with the LH conformation. RH2 seemed to also exchange
with RH1 conformation but to a lesser extent. Thus, RH1 may be seen as a transition
state in between RH1 and LH conformations.
We then refined the three main TM dimer structures (RH1, RH2 and LH) by
performing 400 ns of atomistic MD simulations (see Methods) (Sup. Fig. 11-A). The
RH1 and RH2 dimers remained in a right-handed conformation though we observed a
slight increase of the crossing angle for the RH2 structure (with a final crossing angle
of -37°). Interestingly, the LH structure switched to a right-handed conformation at the
beginning of the simulation and then returned to a stable left-handed conformation.
This confirmed the exchanges between RH1 and LH configurations seen in coarse-
grain simulations (Sup. Fig. 5 to 10). For all three structures, the interactions between
the Small-X3-Small motifs were stable throughout the simulation (Sup. Fig. 11-B).
Thus, MD simulations revealed a dynamic equilibrium of dimer structures
involving the two consecutives Small-X_3-Small motifs: A_{544}-X3- G_{548} and G_{545}-X3-G_{549}.
136
Distinctive mutations in the Small-X_3-Small motifs selectively modulate FLRT2
TM dimerization
To assess the individual contributions of the two Small-X_3-Small motifs to the
dimerization, we performed CG-MD simulations with several mutants replacing glycine
residues with isoleucine or valine residues, the larger hydrophobic side chains of which
are expected to disturb the TM dimerization[62-66] (Fig. 3A and Sup. Table 1). For each
mutant, we evaluated both the crossing angle populations and the spatial distributions
for the TM_{32} construct in a DPPC bilayer.

Figure 3: *In silico* mutations in the two Small-X₃-Small motifs affect the TM dimerization and dynamical equilibrium. A- Table of mutations for *in silico* and SMT experiments. The LRR/TM$_{1+2}$ mutant was only used for the SMT experiments. **B-** Spatial distribution profiles of one TM$_{32}$ helix relative to the other for the CG simulations of both WT and mutants. The diagram shows the probability density of finding the backbone particles of one TM$_{32}$ helix at a given point in the bilayer plane around the other helix. Diagrams are colored such that white represents low probability up to blue for higher probabilities. Green (respectively red) circles depict averaged positions of A$_{544}$ and G$_{548}$ (respectively G$_{545}$ and G$_{549}$) residues. **C-** Helix crossing angle distributions for both TM$_{32}$ WT and mutants. **D-** Averaged TM contact matrices of the different mutants based on different representative TM conformations.

In contrast with the wild type, mutations in the A_{544}-X_3-G_{548} motif (mutants TM_0 and TM_1) favored formation of a dimer with a spatial distribution focused on the G_{545}-X_3-G_{549} motif (**Fig. 3B**) and switched of crossing angle populations towards RH1 conformations (**Fig. 3C**). For these conformations, TM helices are packed around the G_{545}-X_3-G_{549} motif (**Fig. 3D**). Mutations in the G_{545}-X_3-G_{549} motif (mutants TM_2 and TM_3) drove interactions through the G_{544}-X_3-G_{548} motif (**Fig. 3B,D**) favoring RH2 crossing angle populations (**Fig. 3C**) and allowing TM domains to explore a wider area around the G_{544}-X_3-G_{548} motif compared to the WT (**Fig. 3B**). Mutations of both motifs (mutant TM_{1+2}) enabled one TM domain to explore the entire bilayer plane surrounding its TM partner (**Fig. 3B**), thereby abolishing the specificity of the TM helix interactions. For this double mutant, the crossing angle density was clearly more diffuse than for the WT or the other mutants, further highlighting a loss of specificity (**Fig. 3C**). Mutating both motifs also limited the interactions in between the TM helices for the two main conformations RH2 and LH (**Fig. 3D**).

Thus, these mutations highlighted two distinct dynamical behaviors of the TMD dimer specific to each motif. Only mutation of both motifs resulted in an almost complete loss of interactions between the two monomers.

Mutations in the Small-X_3-Small motifs affect FLRT2 co-localization in cells

To support the *in silico* results, we performed SMT experiments to assess the contribution of the predicted key residues in the Small-X_3-Small motifs to dimer formation by mutating glycine residues to isoleucine or valine residues (**Fig. 3A**). We tracked FLRT2 receptors on live cells with a sub-pixel accuracy by SMT in two different channels using the dyes Alexa549 and CF640R (**Fig. 4A-C**). Based on receptor frame to frame proximity in each channel (**Fig. 4C**), we then built a distribution of the durations of co-localization events (**Fig. 4D**), referred to as τ_{on}. The duration of co-localization events is a characteristic of the stability of any interaction or association between the tracked receptors, and is independent of expressed receptor concentration[47]. Comparing the τ_{on} distributions of WT and mutants (**Fig. 4E**) revealed that mutations in only one of the two motifs (either TM_1, TM_2 or TM_3 alone) was insufficient to reduce the baseline average τ_{on} of wild-type FLRT2. However, mutation of both Small-X_3-Small motifs (TM_{1+2} mutant) resulted in a significant shift in the τ_{on} distribution towards lower values (**Fig. 4E and Sup. Fig. 12B**). These results are in agreement with a previous study showing that mutation of both Small-X_3-Small transmembrane motifs is necessary to disrupt the EGFR TM dimer and affect receptor function[67]. The significant shift in τ_{on} was also seen for the mutation in the LRR ectodomain, known to abolish FLRT-FLRT trans-interactions[3], and for the triple mutation LRR+TM_{1+2}. This provides experimental support for the *in silico* results demonstrating that the two Small-X_3-Small motifs are required for FLRT interactions *in cis*.

Figure 4: Mutations in the TM domains affect colocalization of FLRT2 monomers at the cell surface. A- Single molecule TIRF image of HeLa cells expressing wtFLRT2 labelled with both SNAP-549 and BG-CF640R. **B-** Single molecule tracks are generated from time series of the molecules under observation. **C-** An example pair of colocalized tracks where the tracks are separated by less than 1 pixel (160 nm) during at least 5 frames (250 ms). **D-** Example τ_{on} distribution for the WT (n represents the number of tracks analyzed). **E-** Distributions of τ_{on} for wtFLRT2 and each of the six FLRT2 mutants tested. Significance analysis of these distributions available in Sup. Fig 12B. **F-** Distributions of diffusion values for wtFLRT2 and each of the six FLRT2 mutants tested. Significance analysis of these distributions available in Sup. Fig 12C.

We next analyzed the diffusion coefficients for the WT and mutants as an
alternative approach to provide further insight into the role of these motifs in dimer
formation (**Fig. 4F**). In line with the τ_{on} results, only diffusion values for the mutants
TM$_{1+2}$, LRR, and LRR+TM$_{1+2}$, increased significantly from the WT (**Sup. Fig. 12C,D**).
Notably, diffusion calculations in MD simulations also indicated a significant difference
in between the double mutant TM$_{1+2}$ and the WT (**Sup. Fig. 12E**). Although the spatial
resolution of single molecule tracking is insufficient to discriminate between direct
pairwise interactions and co-confinement or joint interactions with the same larger
protein complex, this correlation between the decreased duration of co-localization
events and the increased diffusion constant of the tracked receptors is consistent with
the mutations disrupting interactions that usually occur between WT FLRT2 TM
domains.
Taken together, these data indicate that the extracellular motif and both G$_{544}$-
X$_3$-G$_{548}$ and G$_{545}$-X$_3$-G$_{549}$ motifs in the transmembrane region are required for wild-type
FLRT2 homotypic interactions *in cis*.

Discussion

TM dimer association is a highly dynamic process involving multiple states and weak interactions hence challenging structural studies. As a consequence, only a limited number of TM dimer structures are available and these are often restricted to one state of the TM dimer[68]. To complement structural studies, MD simulations can provide details of these dynamic assemblies. Here, we have used MD simulations to gain structural insights into the formation of FLRT2 TM dimers. Our models revealed a dynamical equilibrium between conformations involving two successive Small-X_3-Small motifs: G_{544}-X_3-G_{548} and G_{545}-X_3-G_{549} motifs. To further assess these models and the conformational changes governing the FLRT2 dimer association, one could use

Figure 5: Model of the FLRT *cis*-interaction. A- Sequence alignment of the TM domain for FLRT1-3 in human and for FLRT2 in other species (human:h, chicken:g, frog:x, fish:dr). **B-** Model of FLRT2 cis-interactions that may compete with different FLRT2 trans interactions. The interconversion in between RH1, involving the G_{545}-G_{549} motif (in red), and RH2 interactions, driven by the A_{544}-G_{548} motif (in green), may be modulated by mutations in the TM domain or environmental conditions such as changes in the lipid composition of the membrane.

accelerated MD simulations combined with free energy calculations[69,70]. To enable

this, we are releasing the three main conformations (RH1, RH2 and LH) both in coarse

grained and atomistic representations (https://github.com/MChavent/FLRT).

The TM helices of other receptors such as EGFRs and EPHAs dimerize via

Small-X_3-Small motifs to transduce extracellular signals into intracellular activity[21,22,24].

Unlike these receptors, there is no enzymatic activity associated with FLRT. Instead,

FLRTs act as crucial adaptor proteins that define the architecture of cell surface

signalling hubs[3,7,13], and also regulate the trafficking of diverse receptors. For example,

FGF receptor traffics with FLRT1-3[71,72], while FLRT3 regulates the cell surface levels

of the Deleted in Colorectal Carcinoma (DCC) receptor in thalamic axons[73].

Dimerization of the EGFR Small-X_3-Small motif also regulates EGFR trafficking[65].

Thus*, in cis* dimerization via the Small-X_3-Small motifs in the TM helix of FLRTs may

act similarly and play a central role in receptors trafficking and thereby signal

transduction.

A functional role for the Small-X_3-Small motifs is supported by their

conservation in all three FLRT human homologues (FLRT 1-3) as well as for FLRT2 in

different species (**Fig. 5A**). In addition, the COSMIC database[74] lists a number of

cancer-related mutations targeting the TM domain of FLRT2. Two such mutations

(A544V and G545V) map to the Small-X_3-Small motifs described here, and may affect

FLRT2 function and dynamics as seen in MD simulations (**Fig. 3**). Furthermore, the

lipid environment can modulate WT TM association[75-77] (**Fig. 2A, Sup. Fig. 4**) and may

accentuate deleterious effects of these mutations.

Unexpectedly, our results show that the same mutation in the LRR domain that

disrupts FLRT-FLRT interactions *in trans*[3] also disrupts FLRT-FLRT interaction *in cis*,

posing the question whether FLRT *cis* and *trans* interactions are competitive (**Fig. 5B**).

Adding complexity to this issue is the observation that the same mutation also

abolishes *trans* FLRT-Lphn interactions[3,6]. These findings suggest that Lphn may also

compete with *in cis* FLRT-FLRT dimerization, leading to a mechanism in which FLRTs

switch between *in cis* dimerization and *in trans* interactions (**Fig. 5B**). Interplay

between *cis* and *trans* interactions are essential for the formation of clusters of

adhesion proteins leading to cell-cell recognition[78]. One canonical example is the

formation of adherens junction by the clustering of cadherins for which both *cis* and

trans interactions are required[79,80]. In addition, a competitive switch between *in trans*

and *in cis* interactions occurs for EphA receptors with their ephrin-A ligands[81,82]. Our

results describe the structural basis and nature of *in cis* dimerization of FLRT2

receptors and suggest how these interactions may compete with interactions *in trans*

to determines the formation of specific signalling complexes and regulate fundamental

developmental processes, such as cortical folding and synapsing.

Acknowledgements

A.L.D. is supported by the BBSRC grant BB/R00126X/1 and Pembroke College,

Oxford (BTP Fellowship). E.Y.J. was funded by UK Medical Research Council

programme grant G9900061. The Wellcome Centre for Human Genetics is supported

by Wellcome Trust Centre grant 203141/Z/16/Z. E.S. is funded by the Wellcome Trust
(202827/Z/16/Z), EMBO YIP and supported by the COST action 'Adhere'n Rise'. V.A.J
was funded by the Wellcome Trust DPhil programme in Cellullar Structural Biology.
M.S.P.S. is funded by Wellcome Trust grants WT092970MA and 208361/Z/17/Z. M.
C. is supported by the CNRS-MITI grant "Modélisation du vivant" 2020. This work was
granted access to the HPC resources of CALMIP supercomputing center under the
allocation 2020-17036. The single molecule analysis used computing resources
provided by STFC Scientific Computing Department's SCARF cluster. We
acknowledge Life Science Editors for proofreading the manuscript. M. C. thanks E.
Haanappel and A. Atkinson for fruitful discussions and support.

Competing interests

No competing interests declared

Methods

Modeling Transmembrane domain and Molecular Dynamics Simulations

Results from the PSIpred[83], PRED-TMR2[84], and HMMTOP[85] servers were combined to predict the membrane embedded helical region of FLRT2. Twenty-four residues of human FLRT2 (residues 541 – 564) were selected to form the core of the TM helix (TM_{24}). The transmembrane domain was created using the Pymol secondary structure creation script: build_seq.py (http://pldserver1.biochem.queensu.ca/~rlc/work/pymol/) and then converted into coarse-grained model. For TM_{32}, the four residues both N- and C-terminal of TM24 were modelled as random coils using Modeller 9v9[86].

Coarse-grained MD (CG-MD) simulations were performed using GROMACS 4.6 (www.gromacs.org)[87,88] with the MARTINI 2.1 forcefield[89,90]. The temperature was 323K. Electrostatic interactions were shifted to zero between 0 and 1.2 nm and the Lenard-Jones interactions between 0.9 and 1.2 nm. A Berendsen thermostat in combination with a Berendsen barostat[91] with a coupling constant of 1.0 ps, a compressibility of 5.0 x 10^{-6} bar^{-1}, and a reference pressure of 1 bar were used. The integration timestep was 20 fs. Simulations were run for either 1 or 2 μs (**Sup. Table 1**) over twenty to thirty replicates to ensure exhaustive sampling of TM helix dimer structures.

We then converted the three main representative (RH1, RH2, and LH) coarse grained structures into atomistic models using the CHARMM-GUI MARTINI to All-atom converter (http://www.charmm-gui.org/?doc=input/converter.martini2all). Atomistic simulations were performed with GROMACS 2018 in combination with the CHARMM36 forcefield[92-94] and TIP3P water model. The temperature was held at 310K. A first step of energy minimization was performed using the steepest descent algorithm and was equilibrated with a constant temperature (canonical ensemble, NVT, 310 K) ensemble for 100 picoseconds, followed by a 100 picoseconds equilibration at constant pressure (isothermal-isobaric, NPT, 1 bar). We then ran 100 ns of equilibration by keeping the protein backbone constrained followed by 400 ns of unrestrained production run. We applied a Nosé-Hoover thermostat on the system, coupled with the Parrinello–Rahman barostat, with a compressibility of 4.5x10^{-5} bar^{-1}. Long-range electrostatics were modeled using the Particle-Mesh Ewald method[95]. All bonds were treated using the LINCS algorithm[96]. The integration time step was 1 femtosecond.

Simulation analysis:

Protein and lipid structures were rendered using VMD[97]. Simulations trajectories were analyzed using a combination of Tcl/VMD and Python scripts. Matplotlib was used to create graphs and images of TMD monomer distances, contact matrices, TMD density rendering, and crossing angles analysis. All the scripts used to

perform these analyses are available at: https://github.com/MChavent/FLRT.

Distances were calculated in between the two centers of mass of each TM helix.

Density, TM contacts and crossing angles calculations were performed every

nanosecond for the part of the trajectory where a dimer was formed. In Figure 2-A

(resp. 3-B,C), the values were renormalized to take into account of both the maximum

values and time of interactions to properly compare the different membrane (resp. Wild

Type and mutants) systems.

Diffusion analyses of dimers were carried out by extracting time-averaged

mean squared displacement (MSD) for all simulation replicates, for the part of the

trajectory where a dimer was formed. For each replicate, a MSD vs. Δt was plotted for

Δt in the range 0.2 ns – 200 ns. The diffusion coefficient (D) for each trajectory was

calculated by fitting the equation $MSD = 4D\Delta t$ to the first two points in the plot, in order

to obtain an instantaneous diffusion coefficient. The diffusion analysis script used to

perform the analysis can be found at:

https://github.com/tylerjereddy/diffusion_analysis_MD_simulations.

Cloning

328   SNAP-FLRT2 was cloned into the EcoRI/XhoI restrictrion sites of the pHSec

329   vector[98]. In SNAP-FLRT2 an N-terminal SNAP tag (containing the RPTPσ signal

330   sequence) was fused to murine FLRT2 (residues A35 – T660) via an HA-tag. Mutations

331   were introduced using molecular cloning.

Cell Culture and Transfection

HeLa cells were seeded onto uncoated 4-well μ-Slides, #1.5 polymer coverslips

(Ibidi) at a density of 1.1x105 cells/well in 600 μL phenol red-free DMEM + 10% FBS

+ 1% L-Gln + 1% NEAA (complete medium). After 24 h, each well was transfected with

2.0 μg plasmid DNA using FuGENE6, according to the manufacturer's instructions.

Cells were maintained at 37 °C, 5% CO2 and were prepared for experiments 12-18

hours post-transfection.

BG-CF640R Conjugation

CF640R succinimidyl ester (Biotium) was reacted with BG-NH2 (New England

Biolabs) to produce the benzylguanine functionalised dye BG-CF640R. 1 μmol of

CF640R succinimidyle ester was reconstituted in DMSO and dissolved in 10 ml 0.1 M

sodium bicarbonate buffer at pH 8.4. 1.5 μmol BG-NH2 in DMSO was added to the

dye mixture and vortexed well. The reaction was shaken at room temperature

overnight before dilution with deionised water. For all subsequent dilutions the

conjugation efficiency was assumed to be 100%.

Two-Colour Fluorescent Labelling

To achieve an approximately equal ratio of single molecules labelled with SNAP Dy549 and BG-CF640R a two-step staining procedure was used. Firstly, the medium was removed from each well of the 4-well μ-Slides and the cells washed twice with 300 μL complete medium. BG-CF60R was diluted in complete medium to a final concentration of 10 nM and applied to each well of the μ-Slide for 5 min. The medium was then exchanged for 150 μL 10 nM SNAP-Dy549 (SNAP-Surface 549, New England Biolabs) in complete medium and incubated for a further 5 min. All labelling steps were performed at 37 oC, 5% CO2. Labelled cells were then washed three times with complete medium and the final wash replaced with Live Cell Imaging Solution plus 1:50 ProLong Antifade reagent (both ThermoFisher) and incubated for at least 15 min, at 37 $^{\circ}$C, 5% CO2 before beginning experiments.

Single molecule image acquisition and feature tracking

Single-molecule images were acquired using an Axiovert 200M microscope with an iLas2 TIRF illuminator (Cairn, UK), with a ×100 oil-immersion objective (α-Plan-Fluar, NA = 1.46; Zeiss, UK) and an EMCCD (iXon X3; Andor, UK). The microscope is also equipped with a wrap-around incubator (Pecon XL S1). The 561 and 642 nm lines of a LightHub laser combiner (Omicron-laserage Laserprodukte GmbH) were used to illuminate the sample and an Optosplit Image Splitter (Cairn Research) was used to separate the image into its spectral components as described previously[99]. The field of view of each channel for single-molecule imaging was 80 × 30 μm. Typically, for each condition at least 50 fields of view comprising one or more cells were acquired from a total of 4 independent biological replicates. Single molecules were tracked in each field of view for 30s, by which time the majority of molecules had undergone photobleaching. All single-molecule time series data were analyzed using the multidimensional analysis software described previously[100]. Briefly, this software performs frame-by-frame Bayesian segmentation to detect and measure features to sub-pixel precision, then links these features through time to create tracks using a simple proximity-based algorithm. The software determines cubic polynomial registration transformations between wavelength channels from images of fluorescent beads. Feature detection and tracking was performed independently in each channel.

Calculation of colocalisation and τ$_{ON}$

Two-colour TIRF images of the basolateral surfaces of cells were chromatically separated by a beam splitter and registered using custom-made software to map the relative positions of the probes over the time course of data acquisition[100] and extract single molecule tracks. A colocalisation event was defined as one in which a track in one channel moves within one pixel of a track in the other channel before they move apart again (Sup. Fig 12A). The duration of each such event is one measurement of τ$_{ON}$. This parameter indicates the stability of presumptive receptor interactions while

beingin sensitive to variation in expression of the receptors between cells or different
levels of labelling with the two probes within cells[47]. The track positions were registered
between channels prior to this analysis. To reduce the impact of localisation error on
these results a temporal Gaussian smoothing filter of FWHM 4 frames (200 ms) was
applied to the position traces before the colocalisation analyses. τ_{ON} distributions were
compared between conditions using the two sample Kolmogorov-Smirnov test to
decide which were significantly different (**Sup. Fig 12B**).
399
Mean squared displacement and diffusion calculation
From single particle tracks, mean squared displacement (MSD) curves were calculated
as MSD(ΔT)=<$|\mathbf{r_i}(T+\Delta T)-\mathbf{r_i}(T)|^2$> where $|\mathbf{r_i}(T+\Delta T)-\mathbf{r_i}(T)|$ is the displacement between
position of track i at time T and time $T+\Delta T$ and the average is over all pairs of points
separated by ΔT in each track. The average instantaneous diffusion coefficient (D) for
these tracks was calculated by fitting a straight line to the first two points of the MSD
curve then calculating D directly from the gradient m of the fit, D=m/4. The tracks for
each single molecule field of view (FOV) were pooled into one MSD curve per FOV to
produce a sample of D values, one value per FOV per condition. These D distributions
were compared between conditions using the Kolmogorov-Smirnov test to decide
which were significantly different.

References

1. Karaulanov, E. E., Böttcher, R. T. & Niehrs, C. A role for fibronectin-leucine-rich transmembrane cell-surface proteins in homotypic cell adhesion. *EMBO Rep* **7,** 283–290 (2006).

2. Maretto, S. *et al.* Ventral closure, headfold fusion and definitive endoderm migration defects in mouse embryos lacking the fibronectin leucine-rich transmembrane protein FLRT3. *Dev. Biol.* **318,** 184–193 (2008).

3. Seiradake, E. *et al.* FLRT structure: balancing repulsion and cell adhesion in cortical and vascular development. *Neuron* **84,** 370–385 (2014).

4. Yamagishi, S. *et al.* FLRT2 and FLRT3 act as repulsive guidance cues for Unc5-positive neurons. *EMBO J.* **30,** 2920–2933 (2011).

5. S, Y. *et al.* FLRT2 and FLRT3 act as repulsive guidance cues for Unc5-positive neurons. *EMBO J.* **30,** 2920–2933 (2011).

6. Jackson, V. A. *et al.* Structural basis of latrophilin-FLRT interaction. *Structure (London, England : 1993)* **23,** 774–781 (2015).

7. Jackson, V. A. *et al.* Super-complexes of adhesion GPCRs and neural guidance receptors. *Nat Commun* **7,** 11184 (2016).

8. del Toro, D. *et al.* Regulation of Cerebral Cortex Folding by Controlling Neuronal Migration via FLRT Adhesion Molecules. *Cell* **169,** 621–635.e16 (2017).

9. O'Sullivan, M. L. *et al.* FLRT proteins are endogenous latrophilin ligands and regulate excitatory synapse development. *Neuron* **73,** 903–910 (2012).

10. Sando, R., Jiang, X. & Südhof, T. C. Latrophilin GPCRs direct synapse specificity by coincident binding of FLRTs and teneurins. *Science* **363,** eaav7969 (2019).

11. Lu, Y. C. *et al.* Structural Basis of Latrophilin-FLRT-UNC5 Interaction in Cell Adhesion. *Structure (London, England : 1993)* **23,** 1678–1691 (2015).

12. Ranaivoson, F. M. *et al.* Structural and Mechanistic Insights into the Latrophilin3-FLRT3 Complex that Mediates Glutamatergic Synapse Development. *Structure (London, England : 1993)* **23,** 1665–1677 (2015).

13. del Toro, D. *et al.* Structural Basis of Teneurin-Latrophilin Interaction in Repulsive Guidance of Migrating Neurons. *Cell* **180,** 323–339.e19 (2020).

14. Sando, R., Jiang, X. & Südhof, T. C. Latrophilin GPCRs direct synapse specificity by coincident binding of FLRTs and teneurins. *Science* **363,** eaav7969 (2019).

15. Seiradake, E., Jones, E. Y. & Klein, R. Structural Perspectives on Axon Guidance. *Annu. Rev. Cell Dev. Biol.* **32,** 577–608 (2016).

16. Lacy, S. E., Bönnemann, C. G., Buzney, E. A. & Kunkel, L. M. Identification of FLRT1, FLRT2, and FLRT3: a novel family of transmembrane leucine-rich repeat proteins. *Genomics* **62,** 417–426 (1999).

17. Anderson, S. M., Mueller, B. K., Lange, E. J. & Senes, A. Combination of Cα-H Hydrogen Bonds and van der Waals Packing Modulates the Stability of GxxxG-Mediated Dimers in Membranes. *J. Am. Chem. Soc.* **139,** 15774–15783 (2017).

18. Bugge, K., Lindorff-Larsen, K. & Kragelund, B. B. Understanding single-pass transmembrane receptor signaling from a structural viewpoint-what are we missing? *FEBS J.* **283,** 4424–4451 (2016).

19. Cymer, F., Veerappan, A. & Schneider, D. Transmembrane helix-helix interactions are modulated by the sequence context and by lipid bilayer properties. *Biochim. Biophys. Acta* **1818,** 963–973 (2012).

20. Bocharov, E. V. *et al.* Structure of FGFR3 Transmembrane Domain Dimer: Implications for Signaling and Human Pathologies. *Structure (London, England : 1993)* **21,** 2087–2093 (2013).

21. Brooks, A. J. *et al.* Mechanism of Activation of Protein Kinase JAK2 by the Growth Hormone Receptor. *Science* **344,** 1249783–1249783 (2014).

22. Fleishman, S. J., Schlessinger, J. & Ben-Tal, N. A putative molecular-activation switch in the transmembrane domain of erbB2. *Proc. Natl. Acad. Sci. U.S.A.* **99,** 15937–15940 (2002).

23. Lau, T.-L., Kim, C., Ginsberg, M. H. & Ulmer, T. S. The structure of the integrin alphaIIbbeta3 transmembrane complex explains integrin transmembrane signalling. *EMBO J.* **28,** 1351–1361 (2009).

24. Bocharov, E. V. *et al.* Left-handed dimer of EphA2 transmembrane domain: Helix packing diversity among receptor tyrosine kinases. *Biophys. J.* **98,** 881–889 (2010).

25. Artemenko, E. O., Egorova, N. S., Arseniev, A. S. & Feofanov, A. V. Transmembrane domain of EphA1 receptor forms dimers in membrane-like environment. *Biochim. Biophys. Acta* **1778,** 2361–2367 (2008).

26. Bocharov, E. V. *et al.* Structural basis of the signal transduction via transmembrane domain of the human growth hormone receptor. *Biochimica et Biophysica Acta (BBA) - General Subjects* **1862,** 1410–1420 (2018).

27. Zhang, Q. & Cherezov, V. Chemical tools for membrane protein structural biology. *Curr. Opin. Struct. Biol.* **58,** 278–285 (2019).

28. Cross, T. A., Sharma, M., Yi, M. & Zhou, H.-X. Influence of solubilizing environments on membrane protein structures. *Trends Biochem. Sci.* **36,** 117–125 (2011).

29. Bottaro, S. & Lindorff-Larsen, K. Biophysical experiments and biomolecular simulations: A

perfect match? *Science* **361,** 355–360 (2018).

30. Mravic, M. *et al.* Packing of apolar side chains enables accurate design of highly stable membrane proteins. *Science* **363,** 1418–1423 (2019).

31. Kalli, A. C., Campbell, I. D. & Sansom, M. S. P. Multiscale simulations suggest a mechanism for integrin inside-out activation. *Proc Natl Acad Sci USA* **108,** 11890–11895 (2011).

32. Arkhipov, A. *et al.* Architecture and membrane interactions of the EGF receptor. *Cell* **152,** 557–569 (2013).

33. Wilmes, S. *et al.* Mechanism of homodimeric cytokine receptor activation and dysregulation by oncogenic mutations. *Science* **367,** 643–652 (2020).

34. Chavent, M. *et al.* Interactions of the EphA2 Kinase Domain with PIPs in Membranes: Implications for Receptor Function. *Structure (London, England : 1993)* **26,** 1025–1034.e2 (2018).

35. Kaszuba, K. *et al.* N-Glycosylation as determinant of epidermal growth factor receptor conformation in membranes. *Proc Natl Acad Sci USA* **112,** 4334–4339 (2015).

36. Chavent, M., Chetwynd, A. P., Stansfeld, P. J. & Sansom, M. S. P. Dimerization of the EphA1 Receptor Tyrosine Kinase Transmembrane Domain: Insights into the Mechanism of Receptor Activation. *Biochemistry* **53,** 6641–6652 (2014).

37. Wassenaar, T. A. *et al.* High-Throughput Simulations of Dimer and Trimer Assembly of Membrane Proteins. The DAFT Approach. *J. Chem. Theory Comput.* **11,** 2278–2291 (2015).

38. Hall, B. A., Halim, K. B. A., Buyan, A., Emmanouil, B. & Sansom, M. S. P. Sidekick for Membrane Simulations: Automated Ensemble Molecular Dynamics Simulations of Transmembrane Helices. *J. Chem. Theory Comput.* **10,** 2165–2175 (2014).

39. Kalli, A. C., Hall, B. A., Campbell, I. D. & Sansom, M. S. P. A Helix Heterodimer in a Lipid Bilayer: Prediction of the Structure of an Integrin Transmembrane Domain via Multiscale Simulations. *Structure (London, England : 1993)* **19,** 1477–1484 (2011).

40. Stansfeld, P. J., Jefferys, E. E. & Sansom, M. S. P. Multiscale simulations reveal conserved patterns of lipid interactions with aquaporins. *Structure (London, England : 1993)* **21,** 810–819 (2013).

41. Stansfeld, P. J., Hopkinson, R., Ashcroft, F. M. & Sansom, M. S. P. PIP(2)-binding site in Kir channels: definition by multiscale biomolecular simulations. *Biochemistry* **48,** 10926–10933 (2009).

42. Liu, R., Li, Y. & Liu, L. Single molecule fluorescence spectroscopy for quantitative biological applications. *Quant Biol* **4,** 177–191 (2016).

43. Stone, M. B., Shelby, S. A. & Veatch, S. L. Super-Resolution Microscopy: Shedding Light on the Cellular Plasma Membrane. *Chem. Rev.* **117,** 7457–7477 (2017).

44. Dunne, P. D. *et al.* DySCo: quantitating associations of membrane proteins using two-color single-molecule tracking. *Biophys. J.* **97,** L5–7 (2009).

45. Veya, L., Piguet, J. & Vogel, H. Single Molecule Imaging Deciphers the Relation between Mobility and Signaling of a Prototypical G Protein-coupled Receptor in Living Cells. *Journal of Biological Chemistry* **290,** 27723–27735 (2015).

46. Brandsma, A. M. *et al.* Mechanisms of inside-out signaling of the high-affinity IgG receptor FcγRI. *Sci Signal* **11,** (2018).

47. Zanetti-Domingues, L. C. *et al.* The architecture of EGFR's basal complexes reveals autoinhibition mechanisms in dimers and oligomers. *Nat Commun* **9,** 4325 (2018).

48. Low-Nam, S. T. *et al.* ErbB1 dimerization is promoted by domain co-confinement and stabilized by ligand binding. *Nat. Struct. Mol. Biol.* **18,** 1244–1249 (2011).

49. Steinkamp, M. P. *et al.* erbB3 Is an Active Tyrosine Kinase Capable of Homo- and Heterointeractions. *Molecular and Cellular Biology* **34,** 965–977 (2014).

50. Wang, Y., Bugge, K., Kragelund, B. B. & Lindorff-Larsen, K. Role of protein dynamics in transmembrane receptor signalling. *Curr. Opin. Struct. Biol.* **48,** 74–82 (2018).

51. Sharpe, H. J., Stevens, T. J. & Munro, S. A Comprehensive Comparison of Transmembrane Domains Reveals Organelle-Specific Properties. *Cell* **142,** 158–169 (2010).

52. Russ, W. P. & Engelman, D. M. The GxxxG motif: A framework for transmembrane helix-helix association. *J. Mol. Biol.* **296,** 911–919 (2000).

53. Teese, M. G. & Langosch, D. Role of GxxxG Motifs in Transmembrane Domain Interactions. *Biochemistry* **54,** 5125–5135 (2015).

54. Walters, R. F. S. & DeGrado, W. F. Helix-packing motifs in membrane proteins. *Proc. Natl. Acad. Sci. U.S.A.* **103,** 13658–13663 (2006).

55. Chothia, C., Levitt, M. & Richardson, D. Helix to helix packing in proteins. *J. Mol. Biol.* **145,** 215–250 (1981).

56. Polyansky, A. A., Volynsky, P. E. & Efremov, R. G. Multistate organization of transmembrane helical protein dimers governed by the host membrane. *J. Am. Chem. Soc.* **134,** 14390–14400 (2012).

57. Prakash, A., Janosi, L. & Doxastakis, M. GxxxG motifs, phenylalanine, and cholesterol guide the self-association of transmembrane domains of ErbB2 receptors. *Biophys. J.* **101,** 1949–1958 (2011).

58. Petrache, H. I., Grossfield, A., MacKenzie, K. R., Engelman, D. M. & Woolf, T. B. Modulation of

glycophorin A transmembrane helix interactions by lipid bilayers: molecular dynamics
calculations. *J. Mol. Biol.* **302,** 727–746 (2000).

59. Dominguez, L., Foster, L., Straub, J. E. & Thirumalai, D. Impact of membrane lipid composition
on the structure and stability of the transmembrane domain of amyloid precursor protein. *Proc*
Natl Acad Sci USA **113,** E5281–7 (2016).

60. Hong, H. & Bowie, J. U. Dramatic Destabilization of Transmembrane Helix Interactions by
Features of Natural Membrane Environments. *J. Am. Chem. Soc.* **133,** 11389–11398 (2011).

61. Defour, J.-P. *et al.* Tryptophan at the transmembrane–cytosolic junction modulates
thrombopoietin receptor dimerization and activation. *Proc. Natl. Acad. Sci. U.S.A.* **110,** 2540–
2545 (2013).

62. Dhara, M. *et al.* v-SNARE transmembrane domains function as catalysts for vesicle fusion. *Elife*
5, (2016).

63. Munter, L. M. *et al.* GxxxG motifs within the amyloid precursor protein transmembrane sequence
are critical for the etiology of Abeta42. *EMBO J.* **26,** 1702–1712 (2007).

64. Berger, B. W. *et al.* Consensus motif for integrin transmembrane helix association. *Proc Natl*
Acad Sci USA **107,** 703–708 (2010).

65. Heukers, R. *et al.* Endocytosis of EGFR requires its kinase activity and N-terminal
transmembrane dimerization motif. *J. Cell. Sci.* **126,** 4900–4912 (2013).

66. Huang, Y. *et al.* Molecular basis for multimerization in the activation of the epidermal growth
factor receptor. *Elife* **5,** 18756 (2016).

67. Endres, N. F. *et al.* Conformational coupling across the plasma membrane in activation of the
EGF receptor. *Cell* **152,** 543–556 (2013).

68. Bugge, K., Lindorff-Larsen, K. & Kragelund, B. B. Understanding single-pass transmembrane
receptor signaling from a structural viewpoint-what are we missing? *FEBS J.* **283,** 4424–4451
(2016).

69. Lelimousin, M., Limongelli, V. & Sansom, M. S. P. Conformational Changes in the Epidermal
Growth Factor Receptor: Role of the Transmembrane Domain Investigated by Coarse-Grained
MetaDynamics Free Energy Calculations. *J. Am. Chem. Soc.* **138,** 10611–10622 (2016).

70. Domański, J., Sansom, M. S. P., Stansfeld, P. J. & Best, R. B. Atomistic mechanism of
transmembrane helix association. *PLoS Comput Biol* **16,** e1007919– (2020).

71. Wheldon, L. M. *et al.* Critical Role of FLRT1 Phosphorylation in the Interdependent Regulation
of FLRT1 Function and FGF Receptor Signalling. *PLoS ONE* **5,** e10264– (2010).

72. Haines, B. P., Wheldon, L. M., Summerbell, D., Heath, J. K. & Rigby, P. W. J. Regulated
expression of FLRT genes implies a functional role in the regulation of FGF signalling during
mouse development. *Dev. Biol.* **297,** 14–25 (2006).

73. Leyva-Díaz, E. *et al.* FLRT3 is a Robo1-interacting protein that determines Netrin-1 attraction
in developing axons. *Curr. Biol.* **24,** 494–508 (2014).

74. Forbes, S. A. *et al.* COSMIC: mining complete cancer genomes in the Catalogue of Somatic
Mutations in Cancer. *Nucleic Acids Res.* **39,** D945–D950 (2010).

75. Stone, M. B., Shelby, S. A., Núñez, M. F., Wisser, K. & Veatch, S. L. Protein sorting by lipid
phase-like domains supports emergent signaling function in B lymphocyte plasma membranes.
Elife **6,** 212 (2017).

76. Lorent, J. H. *et al.* Structural determinants and functional consequences of protein affinity for
membrane rafts. *Nat Commun* **8,** 1–10 (2017).

77. Lorent, J. H. *et al.* Plasma membranes are asymmetric in lipid unsaturation, packing and protein
shape. *Nat. Chem. Biol.* **16,** 644–652 (2020).

78. Honig, B. & Shapiro, L. Adhesion Protein Structure, Molecular Affinities, and Principles of Cell-
Cell Recognition. *Cell* **181,** 520–535 (2020).

79. Brasch, J., Harrison, O. J., Honig, B. & Shapiro, L. Thinking outside the cell: how cadherins
drive adhesion. *Trends Cell Biol.* **22,** 299–310 (2012).

80. Wu, Y. *et al.* Cooperativity between trans and cis interactions in cadherin-mediated junction
formation. *Proc Natl Acad Sci USA* **107,** 17592–17597 (2010).

81. Yin, Y. *et al.* EphA receptor tyrosine kinases interact with co-expressed ephrin-A ligands in cis.
Neuroscience Research **48,** 285–295 (2004).

82. Carvalho, R. F. *et al.* Silencing of EphA3 through a cis interaction with ephrinA5. *Nat. Neurosci.*
9, 322–330 (2006).

83. Jones, D. T. Protein secondary structure prediction based on position-specific scoring matrices
1 1Edited by G. Von Heijne. *J. Mol. Biol.* **292,** 195–202 (1999).

84. Pasquier, C. & Hamodrakas, S. J. An hierarchical artificial neural network system for the
classification of transmembrane proteins. *Protein Eng.* **12,** 631–634 (1999).

85. Tusnady, G. E. & Simon, I. The HMMTOP transmembrane topology prediction server.
Bioinformatics **17,** 849–850 (2001).

86. Eswar, N. *et al.* Comparative Protein Structure Modeling Using Modeller. *Current Protocols in*
Bioinformatics **15,** 5.6.1–5.6.30 (2006).

87. Hess, B., Kutzner, C., Van Der Spoel, D. & Lindahl, E. GROMACS 4: Algorithms for Highly
Efficient, Load-Balanced, and Scalable Molecular Simulation. *J. Chem. Theory Comput.* **4,** 435–
447 (2008).

88.   Pronk, S. *et al.* GROMACS 4.5: a high-throughput and highly parallel open source molecular
simulation toolkit. *Bioinformatics* **29,** 845–854 (2013).
89.   Monticelli, L. *et al.* The MARTINI Coarse-Grained Force Field: Extension to Proteins. *J. Chem.*
Theory Comput. **4,** 819–834 (2008).
90.   Marrink, S. J., de Vries, A. H. & Mark, A. E. Coarse Grained Model for Semiquantitative Lipid
Simulations. *J Phys Chem B* **108,** 750–760 (2004).
91.   Berendsen, H., Postma, J., Vangunsteren, W. F., DiNola, A. & Haak, J. R. Molecular-Dynamics
with Coupling to an External Bath. *J. Chem. Phys.* **81,** 3684–3690 (1984).
92.   Lee, J. *et al.* CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and
CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J. Chem. Theory*
Comput. **12,** 405–413 (2016).
93.   Lee, S. *et al.* CHARMM36 United Atom Chain Model for Lipids and Surfactants. *J Phys Chem*
B **118,** 547–556 (2014).
94.   Huang, J. & MacKerell, A. D. CHARMM36 all-atom additive protein force field: Validation based
on comparison to NMR data. *J Comput Chem* **34,** 2135–2145 (2013).
95.   Essmann, U. *et al.* A smooth particle mesh Ewald method. *J. Chem. Phys.* **103,** 8577–8593
(1995).
96.   Hess, B. P-LINCS: A Parallel Linear Constraint Solver for Molecular Simulation. *J. Chem.*
Theory Comput. **4,** 116–122 (2008).
97.   Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J Mol Graph* **14,** 33–
8, 27–8 (1996).
98.   Aricescu, A. R., Lu, W. & Jones, E. Y. A time- and cost-efficient system for high-level protein
production in mammalian cells. *Acta Crystallogr. D Biol. Crystallogr.* **62,** 1243–1250 (2006).
99.   Webb, S. E. D., Needham, S. R., Roberts, S. K. & Martin-Fernandez, M. L. Multidimensional
single-molecule imaging in live cells using total-internal-reflection fluorescence microscopy. *Opt*
Lett **31,** 2157–2159 (2006).
100.   Rolfe, D. J. *et al.* Automated multidimensional single molecule fluorescence microscopy feature
detection and tracking. *Eur Biophys J* **40,** 1167–1186 (2011).
640