**A GLOBAL DATABASE OF PHOTOSYNTHESIS MODEL PARAMETERS, AND MODELLED**

**PHOTOSYNTHETIC RESPONSES FROM EVERY MAJOR TERRESTRIAL PLANT CLADE**

**RUNNING HEADLINE**: Global photosynthesis parameterization

Mina Rostamza[1,2,3], and Gordon G. McNickle[1,2,*]

[1]Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN 47907-2054

[2]Center for Plant Biology, Purdue University, West Lafayette, IN 47907-2054

[3]Current address: Plant Molecular and Cellular Biology Laboratory, Salk Institute for Biological

Studies, 10010 North Torrey Pines Road, La Jolla, CA 92037

*Author for correspondence: gmcnickle@purdue.edu, 765-494-4645

**KEYWORDS**: Photosynthesis, Farquhar model, carbon dioxide, climate change, phylogeny, plant

functional type

**ABSTRACT**

Plant photosynthesis is a major part of the global carbon cycle and climate system. Carbon capture by $C_3$ plants is most often modelled using the Farquhar-von-Caemmerer-Berry (FvCB) equations. We undertook a global synthesis of all parameters required to solve the FvCB model. The publicly available dataset we assembled includes 3663 observations from 336 different $C_3$ plant species among 96 taxonomic families coming from every major vascular plant clade (lycophytes, ferns, gymnosperms, magnoliids, eudicots and monocots). Geographically, the species in the database have distributions that span the majority of the globe. We used the model to predict photosynthetic rates for a hypothetical average plant in each major terrestrial plant clade and find that generally plants have dramatically increased their photosynthetic abilities through evolutionary time, with the average monocot (the youngest clade) achieving maximum rates of photosynthesis almost double that of the average lycophyte (the oldest clade). We also solved the model for different hypothetical average plant functional types (PFTs) and find that herbaceous species generally have much higher rates of photosynthesis compared to woody plants. Indeed, the maximum photosynthetic rate of graminoids is almost three times the rate of the average tree. The resulting functional responses to increasing $CO_2$ in average hypothetical PFTs would suggest that most groups are already at or near their maximum rate of photosynthesis. However, phylogenetic analysis showed that there was no evidence of niche conservatism with most variance occurring within, rather than among clades (K=0.357, p=0.001). This high within-group variability suggests that average PFTs may obscure important plant responses to increasing $CO_2$. Indeed, when we solved the model for each of the 3663 individual observations, we found that, contrary to the predictions of hypothetical average PFTs, that most plants are predicted to be able to increase their photosynthetic rates. These results suggest that global models should seek to incorporate high within-group variability to accurately predict plant photosynthesis in response to a changing climate.

## INTRODUCTION

Plant photosynthesis is a major factor in the global climate system. Indeed, the annual flux of atmospheric carbon (C) through the leaves of terrestrial plants is estimated to be $1 \times 10^{15}$ g yr$^{-1}$ (Beer *et al.*, 2010, Hetherington & Woodward, 2003). Carbon capture by $C_3$ plants is most often modelled using models derived from the Farquhar-von-Caemmerer-Berry (FvCB) equations (Farquhar *et al.*, 1980, Farquhar & Wong, 1984, Sharkey *et al.*, 2007, Von Caemmerer, 2000). The FvCB model is a process based physiological model that accurately describes the rate of photosynthesis across light levels, and across both $CO_2$ and $O_2$ concentrations. In its modern form, the FvCB model also accounts for triose phosphate limitation (Lombardozzi *et al.*, 2018, Mcclain & Sharkey, 2019). Indeed, a version of the FvCB model forms the basis for most physiological, ecological, and earth system models that include plants (Rogers *et al.*, 2017).

Models that incorporate plant photosynthesis require accurate parameter estimates, estimates which are spread across four decades of scientific inquiry and may be difficult to find for specific taxa. There have been several syntheses and meta-analyses that focus on two parameters of the FvCB model, $V_{c,max}$ and $J_{max}$ (E.g. Kattge & Knorr, 2007, Walker *et al.*, 2014, Wullschleger, 1993), as well as syntheses on empirically estimated maximum photosynthetic rates (Gago *et al.*, 2019), but we are unaware of any attempt at a global synthesis of the full suite of at least 12 parameters needed to fully predict photosynthetic rates across the all $C_3$ plants. In addition, the modern FvCB model of photosynthesis is well known to be over-parameterised (Qian *et al.*, 2012), and modern techniques for curve fitting and parameter estimation can benefit from better prior information. For example, Bayesian methods can work from a known prior distribution of parameter values to enhance the ability to accurately estimate parameters (e.g. Patrick *et al.*, 2009). Thus, collecting all available parameter estimates into one database would greatly enhance the ability to model global photosynthesis, as well as our ability to estimate parameters for new taxa.

Here, we describe a synthesis of all FvCB parameters where at least one parameter was

estimated for a given species. The summary includes parameter estimates from 359 different plant

species from 96 taxonomic families coming from every major vascular plant clade (lycophytes, ferns,

gymnosperms, magnoliids, eudicots and monocots) whose distributions span the majority of the

globe. The parameter estimates are presented using a number of summary statistics and probability

density histograms. We also solve the FvCB model using the full range of parameter estimates to

generate predictions about the breadth of plant photosynthetic responses across major vascular

plant clades, plant functional types, and individual leaves. The full dataset containing 3663 unique

rows of data is publicly available.

**MATERIALS AND METHODS**

*The FvCB photosynthesis model*

Here, we briefly describe the equations of the FvCB model we used to seek

parameterizations (Farquhar *et al.*, 1980, Sharkey *et al.*, 2007, Von Caemmerer, 2000). The most

basic modern FvCB photosynthesis approach for C3 plants assumes that the rate of carbon

assimilation $(A)$ in photosynthesis is co-limited by either carbon $(A_c)$, light $(A_j)$ or TPU $(A_p)$ according

to:

$$A = \min\left(A_c, A_j, A_p\right) - R_d, \qquad \text{Eqn 1}$$

84    where $R_d$ is the daytime respiration rate (See Table 1 for units).

86    The carbon-limited portion of assimilation by photosynthesis in Eqn 1 is given by:

$$A_c = \frac{V_{c,max} C_i}{C_i + K_c \left(1 + \dfrac{O_i}{K_o}\right)}, \qquad \text{Eqn 2}$$

where $C_i$ and $O_i$ are the intracellular concentrations of $CO_2$ and $O_2$ at the site of ribulose-1,5-

bisphosphate carboxylase/oxygenase (RuBisCO) activity respectively; $K_c$ and $K_o$ are the RuBisCO

half saturation constants for $CO_2$ and $O_2$, respectively, and; $V_{c,max}$ is the maximum possible rate of

photosynthesis. Half saturation constants are often called the enzyme "affinity" for the substrate,

but in reality they have limited biological meaning and simply describe the shape of the curvature of

the functional response (Mcnickle & Brown, 2014).

The light limited portion of photosynthesis is given by:

$$A_j = \frac{JC_i}{4.5C_i + 10.5\Gamma^*},  \qquad \text{Eqn 3}$$

where $J$ is the realised electron chain transport determined by light; $\Gamma^*$ is the minimum partial

pressure of $CO_2$ where carbon assimilation balances respiration (i.e. $A = R_d$) generally called the $CO_2$

compensation point; and $C_i$ is as above.

For the purposes of a global summary, the variable $J$ has been determined in several ways

over the years. The most common approach following Farquhar and Wong (1984) $J$ was found by

solving for the root of a simple quadratic equation:

$$\theta J^2 - J(J_{max} + \alpha Q) + J_{max}\alpha Q = 0,  \qquad \text{Eqn 4a}$$

102    Where $Q$ is the light photon flux density striking the leaf (μmol photons m$^{-2}$ s$^{-1}$), $J_{max}$ is the light

103    saturated maximum possible rate of electron transport; $\theta$ represents curvature of the light

104    response; and; $\alpha$ is the efficiency of light conversion. Since this is a quadratic equation, and since

105    negative values of $J$ have no biological meaning, we can use the quadratic formula to find the

106    positive root of Eqn 4a where:

$$J = \frac{J_{max} + \alpha Q - \sqrt{(J_{max} + \alpha Q)^2 - 4\theta J_{max}\alpha Q}}{2\theta}.  \qquad \text{Eqn 4b}$$

107    For some reason, most modern papers use different symbols from the original formulation. In

108    addition, though significantly less common (Buckley & Diaz-Espejo, 2015), some authors use an

109    approximation of eqns 4 where $J$ is approximated according to:

$$J = \frac{aQ}{\sqrt{1 + \frac{(aQ)^2}{J_{max}{}^2}}}, \qquad \text{Eqn 5}$$

110    where $a$ represents the efficiency of light conversion (similar to $\alpha$ in eqn 4), and $Q$ and $J_{max}$ have

111    their usual meanings. Note that $a$ in Eqn 5 is more typically written in the literature as the Greek

112    letter '*alpha*', but we altered this to avoid confusion with Eqn 4.

114            Finally, TPU limitation of photosynthesis in its most detailed form is given by:

$$A_p = \frac{3T_p(C_i - \Gamma^*)}{C_i - \Gamma^*(1 + 3r)}, \qquad \text{Eqn 6a}$$

115    where $r$ is a unitless scalar related to the proportion of glycolate recycled in chloroplasts (where

116    $0 < r < 1$); $T_p$ is the rate of TPU, and the other parameters have their usual meanings given above

117    (Table 1).  However, it appears to be most common to assume that no glycolate is recycled (i.e.

118    $r = 0$), and then Eqn 6a simplifies to:

$$A_p = 3T_p. \quad Eqn\ 6b$$

120            We note that there are even more complex versions than we have detailed here in Eqns 1-6.

These more complex versions may include parameters for stomatal conductance of $CO_2$ (Collatz *et*

*al.*, 1992), mesophyll conductance of $CO_2$ (Flexas *et al.*, 2014, Niinemets *et al.*, 2011) and

transpiration (Farquhar & Sharkey, 1982, Farquhar & Wong, 1984). However, leaf resistance to $CO_2$

and transpiration could be considered their own rather large sub-fields independent from many

attempts to estimate FvCB model parameters. Thus, we purposefully omitted resistances from our

survey of the literature to make the scope of the literature synthesis more tractable. Also, models

that include detailed temperature relationships via the Arrhenius equation are also somewhat

common (Kattge & Knorr, 2007). Again however, to limit the scope of our synthesis we did not

collect the parameters of these Arrhenius rate parameters for a more complex temperature

dependent FvCB model. We view these as future updates that could be made to the dataset.

*Literature search*

Briefly, we began with a search for "Farqhuar photosynthesis models" in the Web of Science,

read each paper carefully, and extracted values for all parameters of the FvCB model (Table 1). The

search was expanded by using the bibliographies of these papers. In total data was extracted from

202 papers. Units were standardized across studies for comparison. We also collected as many

methodological details about the conditions in the cuvette as possible to attempt to understand

sources of variation in the data that were not caused by taxonomy (E.g. humidity, temperature,

vapour pressure deficit, light levels, $CO_2$ pressure). The full details of our systematic literature search

and data extraction methods and criteria are given in the Supplementary Information.

We found that studies could generally be broadly organised into three types: temperature

effects, leaf nitrogen effects, or mean plant effects. First, temperature is well known to affect the

biochemical reactions in photosynthesis and we wanted to account for this variability (Kattge &

Knorr, 2007). Thus, when temperature was the treatment parameter, values at each temperature

were recorded separately. These rows are labelled as type '$T$' in the data file. Second, leaf nitrogen

can also be related to photosynthesis parameters and we wanted to account for this variability as

well (Kattge *et al.*, 2009, Walker *et al.*, 2014). Thus, when parameters were estimated separately by

leaf nitrogen, each leaf nitrogen level was recorded separately.  Nitrogen was sometimes reported

as percent dry weight, and sometimes as mass per leaf area. Either were recorded, but in separate

data columns. These rows are labelled as type '$N$' in the data file. Studies that reported just one

value for each parameter and species (i.e. a species mean) are labelled as type '$M$' in the data file.

Thus, there are three types of data: (i) mean values (type $M$); (ii) leaf temperature manipulations

(type $T$), and; (iii) leaf nitrogen differences (type $N$). However, when available, temperature was

recorded for type $M$ and type $N$ studies. Similarly, when available, leaf nitrogen was recorded for

type $T$ and type $M$ studies.

*Taxonomy and biogeography*

Species names reported in the original papers were checked the National Center for

Biotechnology Information's species taxonomy database using the brranching::phylomatic_names

function in R (Webb & Donoghue, 2005). In general, this just updated any outdated species names

to the most modern accepted name. In one case, we had to manually change a species names to a

taxonomic synonym to match to the database: *Echinochloa crus-galli* (L.) Beauv was changed to an

accepted synonym *Digitaria hispidula* (Retz.) Willd. The updated species names were then pruned

from the plant megatree of Zanne *et al.* (2014) to visually represent the taxonomic coverage of the

dataset using the brranching library in R (v. 0.6; Chamberlain, 2020). The literature search produced

a few parameter estimates for 21 C4 graminoid species that others had reported in the literature

and the FvCB model is only appropriate for C3 photosynthesis (Collatz *et al.*, 1992). The final data file

includes these few C4 parameters, but C4 plants are excluded from all analyses described here

because the FvCB model does not apply to them. Species were also assigned to the following broad

plant functional types (PFTs) based on growth form: C3 graminoid, Forb, Vine/Climbing, Shrub, or

Tree (Also, C4 graminoids are labelled in the datafile, but we do not analyse them here). Finally, we

recorded growth habit of each species in the following categories: annual, biennial, perennial, or

crop for herbaceous species; deciduous, evergreen, or crop for woody species; and fern, tree, or club

moss for ferns. Growth habit information was not available for three rare and exotic species and was

recorded as NA. Of course, future users are free to organize species into whatever other categories

are of interest.

Not all studies reported a location of plant material collection, measurement, or the cultivar

examined. Thus, to obtain a sense of the geographic coverage of species in this dataset, we used the

Botanical Information and Ecology Network (BIEN; Maitner *et al.*, 2018) to obtain museum

occurrence records for each species in our global database. We then mapped these occurrence

records. The resulting map shows the global distribution of all species in our dataset. Importantly

this map is not the global distribution of measurement locations. Rather, the resulting map provides

window into the range distributions of all species that have been studied for photosynthesis, and

therefore details what regions of the world have good coverage of at least approximate species level

photosynthetic data. In addition, the resulting map also shows what regions would benefit from

increased empirical attention to improve global models.

*Data summary*

To create a global summary table, we treated the three data types differently. For studies

that report only mean values (type "M"), we used all values in the summary.  For temperature

studies (type "T"), we only used the value nearest to 25°C. This was done because 25°C was the most

common temperature used in studies that did not manipulate leaf temperature and most

temperature studies measure the same leaf across many temperatures.  Our approach of using only

one value per leaf was to avoid pseudoreplication. For leaf nitrogen studies (Type "N"), we averaged

values across leaf N amounts to capture the mean response of plants growing across different soil

fertilities. Because different leaf nitrogen contents represented individual plants in each study, this

method creates a species mean and also seeks to avoid pseudoreplication. To generate a global

summary, we calculated the mean, standard deviation, median, maximum, minimum, skew, and

kurtosis for each parameter from Eqns 1-6.

Data were also summarised by major taxonomic clade and PFT. For these summary rows,

there were generally too few categories in most groups to create density distributions, and we

report only the mean, standard deviation, and sample size. For sample sizes that were n<3, we

report the standard deviation as NA. For these summaries, we did not separate the *T*, *M*, and *N* data

types because many taxa were only studied once.

*Breadth of model outcomes*

It is useful to the modelling community to have a large database of FvCB model parameters,

but the raw parameters do not show the breadth of plant photosynthetic responses on their own,

since 12 parameters (Table 1) may combine in a many different ways to produce the same rate of

photosynthesis. To examine the range of predicted photosynthetic responses, we used the

parameter estimates to actually solve the FvCB model for plant carbon assimilation rate by solving

for model predicted $A - C_i$ and $A - Q$ curves. We did this in three ways. All three of these analysis

uses all three types of data (M, T and N) in order to show the breadth of responses.

First, we solved the model for the average hypothetical plant that describes each major

plant clade. To do this, each parameter value was set equal to the average of the clade, and the

predicted $A - C_i$ and $A - Q$ curves were solved. In addition, all parameters were set to the upper

and lower 95% confidence intervals around the mean, and the equations were solved again to give a

sense of variation within the clade. Missing values were replaced with the global mean in the clade

analysis. This let us compare the average photosynthetic functional response of some hypothetical

average lycophyte, fern, gymnosperm, magnoliid, eudicot, and monocot.

Second, we solved the model for the average hypothetical plant that describes each PFT.

This was done as above with the mean parameter values and the upper and lower confidence

intervals. Here, missing values were replaced with the mean of the appropriate clade (e.g. missing

graminoid parameters were replaced with the monocot mean, while missing vine or shrub

parameters with the eudicot mean).

Third, we went down each of the 3663 rows of our dataset and solved the model for every

individual leaf for which we had data. However, no study in our synthesis estimated all parameters

of the FvCB equation. Thus, to fill in gaps for any row, we used the global mean value for any missing

parameter values. In addition, the average $A_{max}$ for each species was calculated from these model

runs and then drawn onto the phylogeny. We used Bloomberg's K to examine phylogenetic signal in

the species level data using phytools in R (Revell, 2012).

**RESULTS**

*Taxonomy and biogeography*

In total, we obtained at least one parameter estimate from 359 species in 96 families (Fig

1A, Table S1). The data included all major vascular plant clades including: lycophytes (2 species, 1

family), ferns (33 species, 16 families), gymnosperms (23 species, 3 families), and angiosperms (303

species, 77 families). Angiosperms can be further separated into three more sub-clades comprised of

monocots (62 species, 7 families), eudicots (235 species, 68 families), and magnoliids (6 species, 3

families). Tables S1 and S2 include more detailed summaries breaking the available data up among

each of the 96 taxonomic families and within the six clades. In addition, Tables S3 and S4 contain

more detailed summaries breaking the data up by PFT and growth habit.

The occurrence data from BIEN shows that the majority of data comes from North America,

Europe, Australia, New Zealand, and Japan (Fig 1B). Even though much of Africa is desert containing

few to no plants, coverage on the vegetated parts of the continent was patchy, with the best

coverage in South Africa and parts of west Africa.  However, excluding regions in Africa that mostly

do not contain plants (i.e. the Sahara and Namib Deserts), there are entire countries in southern

Africa for which very few species have been studied (e.g. Angola, Morocco, Nigeria). Similarly,

coverage was spotty in northern Asia (E.g. Russia, Kazakhstan, Mongolia), Indochina, Indonesia and

South America (e.g. Chile, Argentina, Uruguay, Paraguay). This is problematic, because many of

these regions of Africa, Asia, and South America with patchy data are known diversity hotspots

where a small number of taxonomic samples may not represent the average plant in those regions

(Myers *et al.*, 2000).  There was, however, a surprising amount of data from species endemic to the

261 southern foothills of the Himalayan mountains, and from Colombia and Ecuador. Regions of Brazil's

262 Amazon rainforest are also sparsely measured, with much of the Brazilian data appearing to have

263 come from the southern grassland regions.

265  Ignoring national borders, the geographic coverage suggests that most named ecosystem

266 'types' (e.g. sensu Whittaker, 1975) also have good coverage. Temperate and boreal forests and

267 grasslands have particularly good coverage (Fig 1B). However, given the high species diversity of

268 tropical ecosystems, there are likely important gaps in our understanding of the diversity of

269 photosynthetic responses in tropical regions across the globe. The arctic regions of Europe have very

270 good coverage, but there is little data from arctic Russia and large gaps in coverage for arctic North

271 America. Similarly, for grasslands, there is very good coverage in Australia and North America, but

272 relatively little for the grasslands of Asia, Africa, and South America.

274 *Summary statistics and parameter distributions*

275  By far, $V_{cmax}$ (n=1364), $J_{max}$ (n=961), and, to a lesser extent, $T_p$ (n=171) were the most

276 frequently estimated and reported parameter values (Table 2, 3). In general, most of the probability

277 density distributions of observed parameters were skewed or possibly multi-modal (Table 2, 3, Fig

278 S1, Fig S2). It is noteworthy that the minimum and maximum reported values of most parameters

279 differed by as much as four orders of magnitude across all vascular plants. Also, the coefficient of

280 variation in most cases was 0.5 or higher, suggesting high dispersion of the data among species.

281 However, the data show that the majority of this within- and among-species variation over orders of

282 magnitude was driven by methodological differences among research groups (Supplementary

283 information). For example, a few species were studied many times by different groups, and variation

284 in reported parameters for these species were largely explained by the nitrogen content of the

285 leaves each group measured, and the VPD tolerance they used when taking measurements (Table

286 S5, Fig S3). Similarly, among all species in our data base, methodological choices of different

research groups explained 60% of the variation in parameter estimates (Fig S4, Fig S5). Thus, once

methods are controlled, the variation is primarily driven by species differences.

Among the six major clades, eudicots had the most data (n=2037 for $V_{c\,max}$) followed by

gymnosperms (n=461 for $V_{c\,max}$) and monocots (n=400 for $V_{c\,max}$). In general, for both $V_{cmax}$ and

$J_{max}$ there was a clear order to the means such that monocots > eudicots > magnoliids >

gymnosperms > lycophytes. This suggests that plants have become increasingly adept at

photosynthesis through evolutionary time. However, ferns do not seem to fit into this schema with

ferns in-between magnoliids and gymnosperms. The half saturation constants ($K_c$ and $K_o$) were only

estimated twice for gymnosperms and never for lycophytes, ferns, and magnoliids. Better in vivo

sampling of lycophyte, fern and magnoliid species may resolve phylogenetic differences of

parameter values in the future.

For PFTs, most researchers studied trees (n=2077 for $V_{c\,max}$) followed by C3 graminoids

(n=388 for $V_{c\,max}$) and forbs (n=222 for $V_{c\,max}$). There were relatively few vines, and shrubs. When

they are included there was a clear order, for both $V_{cmax}$ and $J_{max}$ means such that C3 graminoids >

forbs > shrubs > trees > vines.

*Model outcomes*

We also used the parameters to solve the FvCB model to explore predicted photosynthesis

rates. For comparison among the six major clades represented in our data, we note that $A_{max}$ was

significantly different such that monocots > eudicots > gymnosperms (Fig 2A, B). There were too few

parameter estimates to draw confidence intervals for lycophyte, fern, and magnoliid clades, but

readers should assume they are very wide, and we hesitate to draw many conclusions without more

data. However, if the patterns stand, these predicted photosynthetic rates among clades show a

similar pattern to empirically estimated photosynthetic rates showing that on average land plants

have substantially increased their photosynthetic ability over evolutionary time, matching empirical

observations of $A_{max}$ (Gago *et al.*, 2019).

For PFT photosynthetic rates, $A_{max}$ was significantly different such that C3 graminoids >

shrubs > trees, but the confidence interval around forbs was so large that this group was not

significantly different from any of these groups, suggesting herbaceous forbs fill a wider variety of

photosynthetic niches than other groups (Fig 2C,D). Like magnoliids, there were too few parameter

estimates for vines to draw confidence intervals for these PFTs, and we hesitate to draw conclusions

about these PFTs.

There is reason to think analysing some hypothetical mean clade or PFT member might

obscure diverse responses of genotypes and species. Thus, we also went down our dataset row by

row to generate the entire breadth of photosynthetic responses of every leaf and species for which

we had data. When parameters were missing, we replaced them with the global mean (Table 1). This

created 3663 $A - C_i$ and 3663 $A - Q$ curves (Fig 2E,F). When all 3663 curves were plotted on the

same axis but with slightly transparent lines, the darker regions show the most common responses

and the lighter regions show fewer common responses. Not surprisingly, the darkest lines largely

appear to trace the average monocot and eudicot because these were best taxonomically sampled

groups.  The breadth of maximum photosynthesis values ranged from slightly negative rates of

photosynthesis (-1.6 µmol C m$^{-2}$ s$^{-1}$) to 59.7 µmol C m$^{-2}$ s$^{-1}$.

When these data were averaged by species, there was no phylogenetic niche conservatism

(Fig 1A, K=0.357, p=0.001) indicating that the majority of variation occurred within clades rather

than among clades. Therefore, even though clades do appear to differ on average (Fig 2A,B), each

clade is just as likely to contain some individual species with either low or high modelled $A_{max}$ (Fig

1A).

**DISCUSSION**

Photosynthesis is an important part of the earth climate system. Enormous quantities of $CO_2$

pass through terrestrial plants each year across the globe. As such, accurate parameterisation of

photosynthetic models is key to an understanding of phenomena ranging from simple leaf level

physiology to global atmospheric dynamics. Here, we undertook a synthesis of parameter values in

the literature. The data set we assembled from the literature includes 3663 rows of data across 359

different plant species from 96 taxonomic families that spanned all major vascular plant clades

including lycophytes, ferns, gymnosperms, magnoliids, monocots, and eudicots (Fig 1; Table 2). Our

aim was that modellers with diverse interests and questions could make use of this dataset, either

for some general plant summarised by clade or PFT (Table 2,3). We also hope that empirical

estimations of new parameter estimates can benefit from detailed prior probability distributions for

modern Bayesian model fitting approaches (Fig S1, S2). However, it is also possible with these data

to get more detailed, for example by family (Table S1, Table S2), growth habit (Table S3, Table S4), or

even by individual species.

It has been shown that empirical estimates of $A_{max}$ show a trend towards increasing

photosynthetic efficiency in C3 plants through evolutionary time such that lycophytes > ferns >

gymnosperms > angiosperms (Gago *et al.*, 2019). It is exciting that the FvCB model is accurate

enough to return the same ranking when estimated parameters are used to predict $A_{max}$ (Fig 2).  It's

not surprising that increases in $A_{max}$ are due to evolutionary adaptations that lead to increases in

both $V_{c,max}$, and $J_{max}$ since these parameters control the maximum photosynthetic rate (Table 2).

Our data also show, for gymnosperms, eudicots and monocots that there has also been an increase

in TPU efficiency through evolutionary time (Table 1). However, there are too few data to compare

other parameter values. It would be valuable to have data to compare parameters such as the half

saturation constants to know if there have been evolutionary innovations in RuBisCO affinities. We

restricted our search to *in vivo* parameter estimates. However, analysis of $K_c$ and $K_o$ estimated using

extracted enzyme *in vitro* has recently been done for many diverse lineages ranging from Bacteria,

Archaea and Eukarea (Iñiguez *et al.*, 2020). Interestingly, they find significant phylogenetic

conservation with early groups such as Chorophyta (green algae), Cyanobacteria, and Euglenophyta

(a group of photosynthetic flagellate algae) having RuBisCO affinities for CO2 that were very similar

to vascular land plants, though the actual affinities ranged over an order of magnitude for these

three groups of plants (Iñiguez *et al.*, 2020). Thus, the global understanding of photosynthesis would

benefit from more *in vivo* estimates of enzyme affinities.

Indeed, in terms of actual parameter estimates, we note that $C_i$ (n=112), $\Gamma^*$ (n=64), $K_c$

(n=43), $K_o$ (n=29), and all parameters from eqns 4-5 were rarely estimated (Table 1,2, Fig S1, S2).

Most studies use previously published estimates of these parameters to remove the degrees of

freedom problem caused by over-parameterization of the FvCB model, and then simply estimate

$V_{c,max}$ (n=1294), $J_{max}$ (n=891), and increasingly $T_p$ (n=171). Parameters other than $V_{c,max}$ and $J_{max}$

are more challenging to estimate because of FvCB model over-parameterization. However, these

more rarely estimated parameters are far from constant (Fig S1, S2) despite being required to solve

the model. Thus, there is likely more variation in these more rarely estimated parameters (I.e. $\Gamma^*$,

n=64; $K_c$, n=43; $K_o$, n=29) than is currently known, and we suggest that more attention to these

parameter estimates would greatly improve our ability to accurately model photosynthesis.  Indeed,

all parameters show either a skewed distribution with high kurtosis, or perhaps even a multi-modal

distribution (Fig S1, S2). Currently, there are too few data to know whether these multi-modal

distributions are artefacts of low sample sizes, or if they represent important physiological trade-

offs.

It should also be noted that though we recorded respiration rates ($R_d$, n=468) that were

reported along with the main parameters of the FvCB equations, we did not seek out studies that

report respiration rates of leaves. We are well aware that there are many more estimates of leaf

respiration rates in the literature which are not associated with modelling exercises. However, like

various resistances to $CO_2$ movement into and through the leaf, seeking out leaf respiration rate

measurements would require its own dedicated meta-analysis, which we view as a separate study.

In the future, we seek to continue to update and expand this database of parameters.  Future

updates could also include accounting for stomatal and mesophyll conductance (Flexas *et al.*, 2014,

Niinemets *et al.*, 2011).

399            In addition to the four orders of magnitude variation among species, the data also show that

variation in parameter estimates within a species can be over an order of magnitude (Fig S3).

Importantly, however, most of this variation can be explained by methodological differences among

research groups and are driven by the leaf temperature and VPD inside the cuvette, as well as the

nitrogen content of the leaves (i.e. the growing conditions of the plant). Variation among species is

also partially explained by methods (Fig S4, S5) and clade (Fig 1,2, Table 1). This is well known, and

not particularly surprising, so we do not dwell on these methodological effects here.

*Implications for global models*

408            Most global models rely on simplifying the diversity of plant life by representing plants as

some small number of average PFTs (Wullschleger *et al.*, 2014). However, it is interesting thing to

note that the average plant in a clade (Fig 2A, B), or PFT (Fig 2C, D) tells a different story than the

diversity of results one sees when we model across species, genotypes and even individual leaves

(Fig 2E, F).  For example, from our modelled $A - C_i$ curves, we can predict the average functional

response of each major plant clade (Fig 1A) or PFT (Fig 1C) to rising $CO_2$ levels. Because our data go

back to the 1970s, the average partial pressure of intracellular $CO_2$ across our data set was only 28.4

Pa (~280 ppm), while at the time of this writing the partial pressure of $CO_2$ in the atmosphere was

approximately 42 Pa (~415 ppm). The model output predicts that the average angiosperm has been

able to take advantage of the increased CO2 in the atmosphere in this period from the 1970s to the

present (Fig 2A). Furthermore, the PFT data show that the average angiosperm response is

dominated largely by the average graminoids and average forbs, while the average tree and shrub

shrubs were already at their maximum photosynthetic rate with $CO_2$ at 28.4 Pa. If we continue this

trend and cast forward to the IPCC predictions for the year 2100 (Ipcc, 2012) of 74 – 103.4 Pa (~730-

1020 ppm) $CO_2$, our model results suggest that our hypothetical average plant representing different

clades and PFTs are already near or at their maximum photosynthetic rates at $CO_2$ levels of 42 Pa (Fig

1A, C). Our data using hypothetical mean PFTs suggest that only graminoids, have much capacity for

increased $CO_2$ assimilation beyond their current assimilation rates as we move toward the expected

$CO_2$ composition of the atmosphere by the year 2100. This seems to contradict empirical work

where any plant species nearly always increase photosynthesis rates with increasing $CO_2$ well

beyond 42 Pa (Ainsworth & Long, 2005, Norby & Zak, 2011). What is the cause of this apparent

contradiction?

We suggest, the lack of phylogenetic signal among all the species and clades (Fig 1A) means

modelling plants as some hypothetical average member of a clade or PFT obscures important

underlying species level variability. Indeed, when we examine the individual $A - C_i$ curves for every

leaf for which we had data, the model results show that almost every individual leaf has the capacity

to dramatically increase its photosynthetic rate as we move from current $CO_2$ levels to the IPCC

predictions for the year 2100 (Fig 2E). Thus, our results tell two divergent stories: some hypothetical

average plant from a clade (Fig 2A) or PFT (Fig 2C) is predicted to have little room for additional

photosynthesis with increasing $CO_2$, while actual empirically observed individual leaves almost all

have significant room for additional photosynthesis with increasing $CO_2$ (Fig 2E). Thus, the

simplifying use of PFTs in most climate models may be dramatically underestimating the future

photosynthetic capacity of terrestrial plants because using a mean PFT obscures the fact that most

variation occurs within groups not among groups (Fig 1A). Ecologically, we would expect those

species who can take advantage of increasing $CO_2$ to expand in abundance resulting in an increased

photosynthetic capacity at the community and ecosystem scale.  The clade and PFT means do not

capture this ecological change.

*Future directions: filling in gaps*

Given the divergent results between the response of some mean hypothetical clade or PFT

member which were almost at their maximum rate of photosynthesis in response to rising $CO_2$, and

the individual leaves which almost all had capacity to increase their photosynthetic rate in response

to rising $CO_2$, this seems like a problem for global models. On one hand, the use of a handful of PFTs

is a necessary simplifying assumption in the face of a world with hundreds of thousands of species

(Wullschleger *et al.*, 2014).  But on the other, it obscures the diversity of responses within each

group which may drive future plant-climate feedbacks (Fig 1, 2). Newer models that include

ecosystem demography and ecological competition among more types of plants are already likely

the solution to this problem (Medvigy *et al.*, 2009). We look forward to the increasing use and

development of these ecosystem demography models which are a promising solution to this

problem.

There are some holes in the global data that limit some conclusions that can be drawn.

From a phylogenetic and biogeographic perspective, it seems that the lack of bryophyte and lichen

parameter data (Fig 1A), particularly in the arctic (Fig 1B), is a hole in our ability to predict global

photosynthesis. In some arctic and boreal systems, bryophytes and lichens can represent the bulk of

net primary production by C3 pathways (Limpens *et al.*, 2011). Gago *et al.* (2019) summarized

empirically estimated values of $A_{max}$, and show that mosses and liverworts fit into the evolutionary

hierarchy such that their photosynthetic rates are lower than fern allies like lycophytes. The absence

of FvCB parameters for bryophytes and lichens however, means that it is difficult to build the

important results of Gago *et al.* (2019) into climate models. Since, most variability in the data

occured within groups not among groups (Fig 1A), models would likely benefit from increased

taxonomic coverage of these more rarely studied groups.

472            From a perspective of potential geographic bias, we cannot help but notice that – like many

global datasets (E.g. Díaz *et al.*, 2016, Luyssaert *et al.*, 2007, Wright *et al.*, 2004) – the FvCB

parameter data are very western-centric, primarily coming from North American, Australian, and

European species. This is likely a function of past funding levels, but we should work to correct this

historical pattern. Indeed, many fields in science are well known to have a problem with diversity

among members of the field (Swartz *et al.*, 2019). Such lack of diversity is thought to limit

perspective and cost the field bright minds from underrepresented minorities. Indeed, more diverse

collaborations have been shown to lead to higher impact research (Alshebli *et al.*, 2018). However, in

a field like biology where diversity of life is also a part of the structure of our data, we suggest this

western-centric data bias is also harming our global understanding of ecological systems as much as

it is harming scientists from under-represented groups. There could be much to be gained by

increased coverage of species from the South American, African, and Asian continents. Particularly,

since most terrestrial biodiversity hotspots are in these regions. Data exists for almost every species

of tree in the boreal forest, and arguably the majority of the most common trees in temperate

forests, yet we know comparatively little about the diversity of photosynthetic responses of the

enormous diversity of plant species in the tropics (Fig 1B). However, we do not think the solution to

this dual problem with diversity of scientists and diversity of plant data is for western scientists to

move their research into regions of Africa, South America and Asia that are under sampled. These

regions already have scientists who can become experts or perhaps collaborators. Given that the

cost of instruments required to estimate photosynthesis parameters is unusually high for ecological

research, we recommend an international collaborative approach might be the most useful way to

combine western access to expensive instruments, with local expertise in flora. Some have called

such international collaborations the "fourth age of science" (Adams, 2013).

*Conclusion*

Photosynthesis is a key process in the global climate system and is often modelled with FvCB

type models. However, considering that there are 300,000 estimated plant species on earth, there is

the potential for a large diversity of photosynthetic ability among plants. Thus, accurate models

require large databases of parameter estimates. We have assembled such a database containing all

parameter estimates required to solve the FvCB photosynthesis model. The publicly available

database contains 3663 rows of data where at least one parameter was estimated for a given

species. The summary includes parameter estimates from 359 different plant species from 96

taxonomic families representing all major vascular plant clades. The biogeographic coverage of

species spans the majority of the globe, although there are some important gaps. We find very

different predicted photosynthetic rates depending on whether we examine some hypothetical

average plant that is meant to represent a clade or PFT, compared to when we model individual

leaves in our dataset. Specifically, when hypothetical average plants are modelled we found that

most clades are approaching their maximum photosynthetic rate in response to elevated $CO_2$.

However, when the breadth of responses for individual leaves are modelled, we found that almost

all plants are predicted to increase photosynthetic rates in response to elevated $CO_2$. We hope that

this database can improve our understanding of global carbon flux through the terrestrial biosphere.

**ACKNOWLEDGMENTS**

We thank Mike Mickelbart and Scott McAdam for many helpful discussions about

photosynthesis, and we thank Laura Jessup and Abdel Halloway for comments on the manuscript.

We thank Jaum Flexas and Joesteph Stinziano for detailed comments on the preprint of this

manuscript. This work was funded in part by USDA NIFA Hatch funds to GGM (project number

1010722).  The authors declare no conflicts of interest.

**DATA ACCESSABILITY**

All data will be made publicly available on dryad upon publication at

https://doi.org/10.5061/dryad.3tx95x6dr. All code used to make all figures and any data analysis are

publicly available on GitHub (https://github.com/ggmcnickle/GlobalFvCB).

**AUTHOR CONTRIBUTIONS:**

Both authors contributed equally to data collection, analysis and writing.

**REFERENCES**

Adams J (2013) The fourth age of research. *Nature,* **497**, 557-560.
Ainsworth EA, Long SP (2005) What have we learned from 15 years of free-air co2 enrichment
(face)? A meta-analytic review of the responses of photosynthesis, canopy properties and
plant production to rising co2. *New Phytologist,* **165**, 351-372.
Alshebli BK, Rahwan T, Woon WL (2018) The preeminence of ethnic diversity in scientific
collaboration. *Nature Communications,* **9**, 10.
Beer C, Reichstein M, Tomelleri E *et al.* (2010) Terrestrial gross carbon dioxide uptake: Global
distribution and covariation with climate. *Science,* **329**, 834-838.
Buckley TN, Diaz-Espejo A (2015) Reporting estimates of maximum potential electron transport rate.
*New Phytologist,* **205**, 14-17.
Chamberlain S (2020) Brranching: Fetch 'phylogenies' from many sources. R package version 0.6.0.
pp Page.
Collatz G, Ribas-Carbo M, Berry J (1992) Coupled photosynthesis-stomatal conductance model for
leaves of $c_4$ plants. *Functional Plant Biology,* **19**, 519-538.
Díaz S, Kattge J, Cornelissen JHC *et al.* (2016) The global spectrum of plant form and function.
*Nature,* **529**, 167-171.
Farquhar GD, Caemmerer SV, Berry JA (1980) A biochemical-model of photosynthetic co2
assimilation in leaves of c-3 species. *Planta,* **149**, 78-90.
Farquhar GD, Sharkey TD (1982) Stomatal conductance and photosynthesis. *Annual Review of Plant*
*Physiology and Plant Molecular Biology,* **33**, 317-345.
Farquhar GD, Wong SC (1984) An empirical-model of stomatal conductance. *Australian Journal of*
*Plant Physiology,* **11**, 191-209.
Flexas J, Carriquí M, Coopman RE *et al.* (2014) Stomatal and mesophyll conductances to co2 in
different plant groups: Underrated factors for predicting leaf photosynthesis responses to
climate change? *Plant Science,* **226**, 41-48.
Gago J, Carriquí M, Nadal M, Clemente-Moreno MJ, Coopman RE, Fernie AR, Flexas J (2019)
Photosynthesis optimized across land plant phylogeny. *Trends in Plant Science,* **24**, 947-958.
Hetherington AM, Woodward FI (2003) The role of stomata in sensing and driving environmental
change. *Nature,* **424**, 901-908.
Iñiguez C, Capó-Bauçà S, Niinemets Ü, Stoll H, Aguiló-Nicolau P, Galmés J (2020) Evolutionary trends
in rubisco kinetics and their co-evolution with co2 concentrating mechanisms. *The Plant*
*Journal,* **101**, 897-918.
Ipcc (2012) *Managing the risks of extreme events and disasters to advance climate change*
*adaptation,* Cambridge, England, Cambridge University Press.
Kattge J, Knorr W (2007) Temperature acclimation in a biochemical model of photosynthesis: A
reanalysis of data from 36 species. *Plant, Cell & Environment,* **30**, 1176-1190.
Kattge J, Knorr W, Raddatz T, Wirth C (2009) Quantifying photosynthetic capacity and its relationship
to leaf nitrogen content for global-scale terrestrial biosphere models. *Global Change Biology,*
**15**, 976-991.
Limpens J, Granath G, Gunnarsson U *et al.* (2011) Climatic modifiers of the response to nitrogen
deposition in peat-forming sphagnum mosses: A meta-analysis. *New Phytologist,* **191**, 496-
507.
Lombardozzi DL, Smith NG, Cheng SJ *et al.* (2018) Triose phosphate limitation in photosynthesis
models reduces leaf photosynthesis and global terrestrial carbon storage. *Environmental*
*Research Letters,* **13**, 074025.
Luyssaert S, Inglima I, Jung M *et al.* (2007) Co2 balance of boreal, temperate, and tropical forests
derived from a global database. *Global Change Biology,* **13**, 2509-2537.

Maitner BS, Boyle B, Casler N *et al.* (2018) The bien r package: A tool to access the botanical
information and ecology network (bien) database. *Methods in Ecology and Evolution,* **9**, 373-
379.
Mcclain AM, Sharkey TD (2019) Triose phosphate utilization and beyond: From photosynthesis to
end product synthesis. *Journal of Experimental Botany,* **70**, 1755-1766.
Mcnickle GG, Brown JS (2014) When michaelis and menten met holling: Towards a mechanistic
theory of plant nutrient foraging behaviour. *AoB Plants,* **6**, plu066.
Medvigy D, Wofsy SC, Munger JW, Hollinger DY, Moorcroft PR (2009) Mechanistic scaling of
ecosystem function and dynamics in space and time: Ecosystem demography model version
2. *Journal of Geophysical Research: Biogeosciences,* **114**.
Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GaB, Kent J (2000) Biodiversity hotspots for
conservation priorities. *Nature,* **403**, 853-858.
Niinemets Ü, Flexas J, Peñuelas J (2011) Evergreens favored by higher responsiveness to increased
$CO_2$. *Trends in Ecology & Evolution,* **26**, 136-142.
Norby RJ, Zak DR (2011) Ecological lessons from free-air co2 enrichment (face) experiments. *Annual*
*Review of Ecology, Evolution, and Systematics,* **42**, 181-203.
Patrick LD, Ogle K, Tissue DT (2009) A hierarchical bayesian approach for estimation of
photosynthetic parameters of c3 plants. *Plant, Cell & Environment,* **32**, 1695-1709.
Qian T, Elings A, Dieleman JA, Gort G, Marcelis LFM (2012) Estimation of photosynthesis parameters
for a modified farquhar–von caemmerer–berry model using simultaneous estimation
method and nonlinear mixed effects model. *Environmental and Experimental Botany,* **82**, 66-
73.
Revell LJ (2012) Phytools: An r package for phylogenetic comparative biology (and other things).
*Methods in Ecology and Evolution,* **3**, 217-223.
Rogers A, Medlyn BE, Dukes JS *et al.* (2017) A roadmap for improving the representation of
photosynthesis in earth system models. *New Phytologist,* **213**, 22-42.
Sharkey TD, Bernacchi CJ, Farquhar GD, Singsaas EL (2007) Fitting photosynthetic carbon dioxide
response curves for c3 leaves. *Plant, Cell & Environment,* **30**, 1035-1040.
Swartz TH, Palermo A-GS, Masur SK, Aberg JA (2019) The science and value of diversity: Closing the
gaps in our understanding of inclusion and diversity. *The Journal of Infectious Diseases,* **220**,
S33-S41.
Von Caemmerer S (2000) *Biochemical models of leaf photosynthesis,* Collingwood, Vic, Australia,
CSIRO Publishing.
Walker AP, Beckerman AP, Gu L *et al.* (2014) The relationship of leaf photosynthetic traits – vcmax
and jmax – to leaf nitrogen, leaf phosphorus, and specific leaf area: A meta-analysis and
modeling study. *Ecology and Evolution,* **4**, 3218-3235.
Webb CO, Donoghue MJ (2005) Phylomatic: Tree assembly for applied phylogenetics. *Molecular*
*Ecology Notes,* **5**, 181-183.
Whittaker RH (1975) *Communities and ecosystems,* New York, MacMillan.
Wright IJ, Reich PB, Westoby M *et al.* (2004) The worldwide leaf economics spectrum. *Nature,* **428**,
821-827.
Wullschleger SD (1993) Biochemical limitations to carbon assimilation in c3 plants—a retrospective
analysis of the a/ci curves from 109 species. *Journal of Experimental Botany,* **44**, 907-920.
Wullschleger SD, Epstein HE, Box EO *et al.* (2014) Plant functional types in earth system models: Past
experiences and future directions for application of dynamic vegetation models in high-
latitude ecosystems. *Annals of Botany,* **114**, 1-16.
Zanne AE, Tank DC, Cornwell WK *et al.* (2014) Three keys to the radiation of angiosperms into
freezing environments. *Nature,* **506**, 89-+.

**TABLE 1**: Summary of symbols, associated equation numbers in the text, plain descriptions and units
for the parameters of the FvCB photosynthesis model that are included in this global summary. Note
that those parameters in units of pressure (Pa), are reported differently by many authors. See text
for unit conversions. Note that $\mu mol\ m^{-2}\ s^{-1}$ represents carbon or photons depending on parameter.

| Parameter | Units | Eqn | Description |
|---|---|---|---|
| $V_{c,max}$ | $\mu mol\ m^{-2}\ s^{-1}$ | 2 | The carbon saturated maximum rate of carbon fixation. |
| $J_{max}$ | $\mu mol\ m^{-2}\ s^{-1}$ | 4, 5, 6 | The light saturated maximum possible rate of electron transport. |
| $C_i$ | Pa | 2, 3 | Intercellular partial pressure of $CO_2$ at the site of RUBISCO activity. (When constructing a $CO_2$ response curve, this is considered a variable, not a parameter.) |
| $\Gamma^*$ | Pa | 2, 3 | The partial pressure of the $CO_2$ compensation point. This is equivalent to a giving up density. It is where carbon assimilation (i.e. benefits) equal respiration costs. |
| $K_c$ | Pa | 2 | The half saturation constant for $CO_2$ of RUBISCO. This is sometimes called the enzyme "affinity" for the substrate, though in reality it has limited biological meaning. Rather, it describes the the curvature of the $CO_2$ functional response. It is simply the partial pressure that is half-way to the maximum rate. |
| $K_o$ | Pa | 2 | The half saturation constant for $O_2$ of RUBISCO. This is sometimes called the enzyme "affinity" for the substrate, though in reality it has limited biological meaning. Rather it describes the curvature of the oxygenation functional response of RUBISCO. It is simply the partial pressure of oxygen that is half-way to the maximum rate. |
| $R_d$ | $\mu mol\ m^{-2}\ s^{-1}$ | 1 | The leaf respiration rate during the day of the leaf. |
| $T_p$ | $\mu mol\ m^{-2}\ s^{-1}$ | 7a, b | The rate of TPU production. |
| $r$ | unitless | 7a | The rate of TPU recycling. Often assumed to be 0. |
| $\alpha$ | unitless | 4 | The efficiency of light conversion. |
| $\theta$ | unitless | 4 | A curvature factor of the light response. |
| $a$ | unitless | 5 | The efficiency of light conversion. |
| **Variable** | | | |
| $Q$ | $\mu mol$ photons $m^{-2}$ $s^{-1}$ | 4, 5 | The light photon flux density striking the leaf. (When constructing a $CO_2$ response curve, this can be considered a parameter not a variable) |
| $J$ | $\mu mol\ m^{-2}\ s^{-1}$ | 3, 4, 5 | The actual rate of electron transport going to support NADP$^+$ reduction for RuBP regeneration. |
| $A_c$ | $\mu mol\ m^{-2}\ s^{-1}$ | 1 | Rate of carbon assimilation limited by carbon reactions. |
| $A_j$ | $\mu mol\ m^{-2}\ s^{-1}$ | 1 | Rate of carbon assimilation limited by light and electron transport. |
| $A_p$ | $\mu mol\ m^{-2}\ s^{-1}$ | 1 | Rate of carbon assimilation limited by TPU. |
| $A$ | $\mu mol\ m^{-2}\ s^{-1}$ | 1 | Actual rate of carbon assimilation |

631 **TABLE 2**: Summary statistics and distribution parameters for each parameter of the Farquhar photosynthesis
632 equations (Eqn 1-4). The global density distribution histogram for each parameter is shown in Fig S1. The
633 mean, standard deviation (SD) and sample size is also shown for each major clade. SD is reported as NA for
634 n<3, and empty cells represent no data.

| Clade | Stat | $V_{c\ max}$ (μmol m$^{-2}$ s$^{-1}$) | $J_{max}$ (μmol m$^{-2}$ s$^{-1}$) | $C_i$ (Pa) | $\Gamma^*$ (Pa) | $K_c$ (Pa) | $K_o$ (Pa) | $R_d$ (μmol m$^{-2}$ s$^{-1}$) | $T_p$ (μmol m$^{-2}$ s$^{-1}$) |
|---|---|---|---|---|---|---|---|---|---|
| All | n | 1352 | 953 | 88 | 58 | 43 | 29 | 458 | 171 |
| | Mean | 61.0 | 134.8 | 37.8 | 4.5 | 35.8 | 29169.1 | 1.2 | 12.0 |
| | SD | 39.1 | 81.0 | 6.6 | 2.6 | 12.6 | 9829.1 | 1.7 | 5.4 |
| | CV | 0.6 | 0.6 | 0.17 | 0.58 | 0.35 | 0.33 | 1.4 | 0.45 |
| | Median | 52.9 | 118.0 | 31.8 | 4.0 | 30.5 | 28168.4 | 0.8 | 12 |
| | Min | 0.3 | 10.4 | 15.9 | 0.01 | 17.0 | 16500 | 0.02 | 1.7 |
| | Max | 267.1 | 498.4 | 37.7 | 9.2 | 70.7 | 47926.7 | 20.3 | 29.2 |
| | Skew. | 1.09 | 1.09 | -0.64 | 0.18 | 0.82 | 0.43 | 7.62 | 0.64 |
| | Kurtosis | 4.60 | 4.41 | 2.01 | 2.45 | 3.08 | 2.04 | 75.31 | 3.59 |
| Lycoph. | n | 2 | 2 | | | | | 2 | |
| | Mean | 17.2 | 25.5 | | | | | 0.5 | |
| | SD | NA | NA | | | | | NA | |
| Ferns | n | 33 | 33 | | | | | 33 | |
| | Mean | 51.2 | 59.7 | | | | | 0.5 | |
| | SD | 27.9 | 24.1 | | | | | 0.3 | |
| Gymno. | n | 461 | 301 | 37 | 27 | 2 | 2 | 41 | 49 |
| | Mean | 34.7 | 65.2 | 19.4 | 7.7 | 27.4 | 41543.3 | 0.9 | 2.8 |
| | SD | 32.25 | 42.93 | 5.14 | 3.04 | NA | NA | 0.45 | 2.62 |
| Magno. | n | 18 | 3 | | | | | 1 | |
| | Mean | 21 | 119.3 | | | | | 0.7 | |
| | SD | 12.85 | 29.16 | | | | | NA | |
| Eudic. | n | 2037 | 1060 | 43 | 63 | 42 | 31 | 283 | 48 |
| | Mean | 76.3 | 135.9 | 28.8 | 4.3 | 47.5 | 26442.6 | 1.5 | 6.8 |
| | SD | 56.88 | 64.8 | 5.67 | 1.87 | 31.63 | 11382.05 | 1.76 | 2.6 |
| Monoc. | n | 395 | 342 | 15 | 8 | 22 | 20 | 225 | 236 |
| | Mean | 86.6 | 196.7 | 28.8 | 3.0 | 51.9 | 37944 | 0.79 | 13 |
| | SD | 40.9 | 92.8 | 7.0 | 1.7 | 32.59 | 17120.35 | 0.50 | 5.35 |
| **PFT** | | | | | | | | | |
| C3 gramin. | n | 388 | 342 | 12 | 5 | 21 | 20 | 222 | 236 |
| | Mean | 87.2 | 199.6 | 29.7 | 3.0 | 53.6 | 37944 | 0.8 | 13 |
| | SD | 41.1 | 92.2 | 7.1 | 1.7 | 32.8 | 17120.3 | 0.5 | 5.4 |
| Forb | n | 222 | 123 | 22 | 43 | 23 | 21 | 40 | 6 |
| | Mean | 83.5 | 167.5 | 31.2 | 4.3 | 41.1 | 24574.4 | 1.5 | 9.4 |
| | SD | 50.3 | 75.4 | 6.0 | 1.6 | 32.2 | 10685.8 | 2.2 | 3.4 |
| Vine | n | 55 | 54 | 1 | 1 | | | 5 | 2 |
| | Mean | 55.9 | 95.4 | 20 | 4.1 | | | 0.5 | 9.1 |
| | SD | 23.1 | 37.2 | NA | NA | | | 0.3 | NA |
| Shrub | n | 166 | 143 | 3 | 4 | 3 | | 23 | 21 |
| | Mean | 72.1 | 125.7 | 18.7 | 2.8 | 26.8 | | 2.1 | 6.6 |
| | SD | 58.4 | 63.3 | 0.8 | 3.5 | 5.0 | | 1.7 | 2.5 |
| Tree | n | 2077 | 1048 | 57 | 42 | 19 | 12 | 259 | 68 |
| | Mean | 66.6 | 114.4 | 22.4 | 6.6 | 54.9 | 32073 | 1.3 | 3.7 |
| | SD | 56.1 | 65.4 | 6.0 | 3.0 | 30.1 | 12035.2 | 1.6 | 2.7 |

* only $R_d$ values reported with estimates of FvCB model parameters were sought. This will not be a
complete literature review of plant respiration rates.

**TABLE 3**: Summary statistics across all measurements for 334 species for each parameter of the
Farquhar photosynthesis equations 4-5. The global density distribution histogram for each
parameter is shown in Fig S2. The mean, standard deviation (SD) and sample size is also shown for
each major clade. SD is reported as NA for n<3, and empty cells represent no data. Clades are
summarised by family in Table S1 and by functional type growth habit in Table S4.

| Clade | Stat | $\alpha$ (Eqn 4) | $\theta$ (Eqn 4) | $a$ (Eqn 5) |
|---|---|---|---|---|
| ALL | n | 41 | 192 | 6 |
| | Mean | 0.22 | 0.6 | 0.17 |
| | SD | 0.21 | 0.28 | 0.06 |
| | CV | 0.95 | 0.47 | 0.35 |
| | Median | 0.16 | 0.67 | 0.19 |
| | Min | 0.02 | 0.04 | 0.06 |
| | Max | 0.95 | 1.0 | 0.23 |
| | Skewness | 1.55 | -0.82 | -1.81 |
| | Kurtosis | 5.86 | 2.66 | 6.87 |
| Gymnosperm | n | 14 | 13 | 3 |
| | Mean | 0.23 | 0.53 | 0.19 |
| | SD | 0.05 | 0.17 | 0.02 |
| Magnoliid | n | | 1 | |
| | Mean | | 0.06 | |
| | SD | | NA | |
| Eudicot | n | 26 | 44 | 3 |
| | Mean | 0.27 | 0.51 | 0.16 |
| | SD | 0.22 | 0.41 | 0.09 |
| Monocot | n | 2 | 223 | |
| | Mean | 0.27 | 0.68 | |
| | SD | NA | 0.16 | |
| **PFT** | | | | |
| C3 graminoid | n | | 220 | |
| | Mean | | 0.68 | |
| | SD | | 0.20 | |
| Forb | n | 17 | 23 | |
| | Mean | 0.32 | 0.77 | |
| | SD | 0.20 | 0.30 | |
| Vine | n | | 3 | |
| | Mean | | 0.65 | |
| | SD | | 0.50 | |
| Shrub | n | 4 | 4 | |
| | Mean | 0.02 | 0.33 | |
| | SD | 0.006 | 0.5 | |
| Tree | n | 20 | 30 | 5 |
| | Mean | 0.24 | 0.33 | 0.17 |
| | SD | 0.20 | 0.30 | 0.10 |

Note: Lycophytes and ferns are absent from this table because no data were available for these
parameters.

**FIG 1**: (A) The evolutionary relationships among all 359 species among 6 major terrestrial plant clades for
which photosynthesis parameters were recovered from the literature and included in our database. Scale bar
represents 50 million years of evolution, and colours represent modelled        . Unresolved taxonomy are
drawn as polytomies at the genus level. The raw phylogeny in newick format is given in the supplementary
information to allow digital visualisation of such a large phylogeny. There was no evidence of phylogenetic
niche conservatism across the entire dataset, meaning species with low or high        are equally likely in any
clade. (B) Global distribution data for all 334 species included in the data set (blue points). Points represent
occurrence distribution data for all species retrieved from BIEN, not the locations of the measurements.

**Fig 2:** (**A**) Mean $A - C_i$ and (**B**) $A - Q$ curves among each of the six major plant clades for which we
have parameter estimates (solid lines). We also show (**C**) Mean $A - C_i$ and (**D**) $A - Q$ curves with
species grouped by broad functional type (dashed lines). Shading around the mean curve in panels
A-D represents solving the model with the 95% confidence interval around all parameters. These
95% confidence intervals cross on the $A - C_i$ curve because of the oxygenation behaviour of
RUBISCO at low $CO_2$ partial pressures. For, lycophytes, ferns, magnoliids, and vines the large number
of missing values were replaced with the global mean, and we did not draw confidence intervals for
these groups (readers should assume they are very wide due to low taxonomic sampling). Finally,
each line in the lower panels represent 3663 separate (**E**) $A - C_i$ or (**F**) $A - Q$ curves generated for
every single row in our database and show the breadth of individual species responses across
diverse conditions. For these 3663 lines, missing values were replaced with the global mean (Table
2). In all panels, for $A - C_i$ curves, the black vertical dotted line marks the estimated mean partial
pressure of $CO_2$ inside the leaf in reported in the literature, while the dashed line represents the
approximate current partial pressure of $CO_2$ in the atmosphere at the time of writing. The shaded
rectangle represents the predicted range of $CO_2$ Intergovernmental panel on climate change (IPCC)
predicted partial pressures of $CO_2$ by the year 2100. For $A - Q$ curves, the black dot-dash line
represents the mean saturating Q used among studies in our database.

APPENDIX A – DATA SOURCES

Aalto, T. (1998) Carbon dioxide exchange of scots pine shoots as estimated by a biochemical model
and cuvette field measurements.

Aalto, T. & Juurola, E. (2001) Parametrization of a biochemical co2 exchange model for birch (betula
pendula roth.). Boreal Environment Research, 6, 53-64.

Aalto, T., Hari, P. & Vesala, T. (2002) Comparison of an optimal stomatal regulation model and a
biochemical model in explaining co2 exchange in field conditions.

Aranda, X., Agustí, C., Joffre, R. & Fleck, I. (2006) Photosynthesis, growth and structural
characteristics of holm oak resprouts originated from plants grown under elevated co2. Physiologia
Plantarum, 128, 302-312.

Archontoulis, S.V., Yin, X., Vos, J., Danalatos, N.G. & Struik, P.C. (2011) Leaf photosynthesis and
respiration of three bioenergy crops in relation to temperature and leaf nitrogen: How conserved
are biochemical model parameters among crop species? Journal of Experimental Botany, 63, 895-
911.

Azcón-Bieto, J. & Osmond, C.B. (1983) Relationship between photosynthesis and respiration: The
effect of carbohydrate status on the rate of co2 production by respiration in darkened and
illuminated wheat leaves. Plant Physiology, 71, 574-581.

Bauer, G.A., Berntson, G.M. & Bazzaz, F.A. (2001) Regenerating temperate forests under elevated
co2 and nitrogen deposition: Comparing biochemical and stomatal limitation of photosynthesis. New
Phytologist, 152, 249-266.

Beerling, D.J. & Quick, W.P. (1995) A new technique for estimating rates of carboxylation and
electron transport in leaves of c3 plants for use in dynamic global vegetation models. Global Change
Biology, 1, 289-294.

Berghuijs, H.N.C., Yin, X., Tri Ho, Q., Van Der Putten, P.E.L., Verboven, P., Retta, M.A., Nicolaï, B.M. &
Struik, P.C. (2015) Modelling the relationship between co2 assimilation and leaf anatomical
properties in tomato leaves. Plant Science, 238, 297-311.

Bernacchi, C.J., Pimentel, C. & Long, S.P. (2003) In vivo temperature response functions of
parameters required to model rubp-limited photosynthesis. Plant, Cell & Environment, 26, 1419-
1430.

Bernacchi, C.J., Singsaas, E.L., Pimentel, C., Portis Jr, A.R. & Long, S.P. (2001) Improved temperature
response functions for models of rubisco-limited photosynthesis. Plant, Cell & Environment, 24, 253-
259.

Bernacchi, C.J., Portis, A.R., Nakano, H., Von Caemmerer, S. & Long, S.P. (2002) Temperature
response of mesophyll conductance. Implications for the determination of rubisco enzyme kinetics
and for limitations to photosynthesis in vivo. Plant Physiology, 130, 1992-1998.

Bernacchi, C.J., Rosenthal, D.M., Pimentel, C., Long, S.P. & Farquhar, G.D. (2009) Modeling the
temperature dependence of c3 photosynthesis. Photosynthesis in silico: Understanding complexity
from molecules to ecosystems (ed. by A. Laisk, L. Nedbal and Govindjee), pp. 231-246. Springer
Netherlands, Dordrecht.

Bown, H.E., Watt, M.S., Clinton, P.W., Mason, E.G. & Richardson, B. (2007) Partititioning concurrent
influences of nitrogen and phosphorus supply on photosynthetic model parameters of pinus radiata.
Tree Physiology, 27, 335-344.

Brahim, B.M., Loustau, D., Gaudillère, J. & Saur, E. (1996) Effects of phosphate deficiency on
photosynthesis and accumulation of starch and soluble sugars in 1-year-old seedlings of maritime
pine (pinus pinaster ait). Ann. For. Sci., 53, 801-810.

Braune, H., Müller, J. & Diepenbrock, W. (2009) Integrating effects of leaf nitrogen, age, rank, and
growth temperature into the photosynthesis-stomatal conductance model leafc3-n parameterised
for barley (hordeum vulgare l.). Ecological Modelling, 220, 1599-1612.

Brooks, A. & Farquhar, G.D. (1985) Effect of temperature on the co2/o2 specificity of ribulose-1,5-
bisphosphate carboxylase/oxygenase and the rate of respiration in the light. Planta, 165, 397-406.

Brück, H. & Guo, S. (2006) Influence of n form on growth photosynthesis of phaseolus vulgaris l.
Plants. Journal of Plant Nutrition and Soil Science, 169, 849-856.

Buckley, T.N., Farquhar, G.D. & Mott, K.A. (1999) Carbon-water balance and patchy stomatal
conductance. Oecologia, 118, 132-143.

Bunce, J.A. (2000) Acclimation of photosynthesis to temperature in eight cool and warm climate
herbaceous c3 species: Temperature dependence of parameters of a biochemical photosynthesis
model. Photosynthesis Research, 63, 59-67.

Cai, T. & Dang, Q.-L. (2002) Effects of soil temperature on parameters of a coupled photosynthesis–
stomatal conductance model. Tree Physiology, 22, 819-828.

Calfapietra, C., Tulva, I., Eensalu, E., Perez, M., De Angelis, P., Scarascia-Mugnozza, G. & Kull, O.
(2005) Canopy profiles of photosynthetic parameters under elevated co2 and n fertilization in a
poplar plantation. Environmental Pollution, 137, 525-535.

Carswell, F.E., Whitehead, D., Rogers, G.N.D. & Mcseveny, T.M. (2005) Plasticity in photosynthetic
response to nutrient supply of seedlings from a mixed conifer-angiosperm forest. Austral Ecology,
30, 426-434.

Cernusak, L.A., Hutley, L.B., Beringer, J., Holtum, J.a.M. & Turner, B.L. (2011) Photosynthetic
physiology of eucalypts along a sub-continental rainfall gradient in northern australia. Agricultural
and Forest Meteorology, 151, 1462-1470.

Chen, T., Stützel, H. & Kahlen, K. (2012) Determining photosynthetic limitations under saturated and
non-saturated light conditions.  2012 IEEE 4th International Symposium on Plant Growth Modeling,
Simulation, Visualization and Applications (ed by, pp. 93-95.

Chen, W., Chen, J., Liu, J. & Cihlar, J. (2000) Approaches for reducing uncertainties in regional forest
carbon balance. Global Biogeochemical Cycles, 14, 827-838.

Collatz, G., Ribas-Carbo, M. & Berry, J. (1992) Coupled photosynthesis-stomatal conductance model
for leaves of $c_4$ plants. Functional Plant Biology, 19, 519-538.

Curtis, P.S. & Teeri, J.A. (1992) Seasonal responses of leaf gas exchange to elevated carbon dioxide in
populusgrandidentata. Canadian Journal of Forest Research, 22, 1320-1325.

Dang, Q.-L., Margolis, H.A. & Collatz, G.J. (1998) Parameterization and testing of a coupled
photosynthesis–stomatal conductance model for boreal trees. Tree Physiology, 18, 141-153.

Davey, P.A., Parsons, A.J., Atkinson, L., Wadge, K. & Long, S.P. (1999) Does photosynthetic
acclimation to elevated co2 increase photosynthetic nitrogen-use efficiency? A study of three native
uk grassland species in open-top chambers. Functional Ecology, 13, 21-28.

Deng, X., Ye, W., Feng, H., Yang, Q., Cao, H., Xu, K. & Zhang, Y. (2004) Gas exchange characteristics of
the invasive species mikania micrantha and its indigenous congener m. Cordata (asteraceae) in south
china. Botanical Bulletin of Academia Sinica, 45, 213-220.

Dreyer, E., Le Roux, X., Montpied, P., Daudet, F.A. & Masson, F. (2001) Temperature response of leaf
photosynthetic capacity in seedlings from seven temperate tree species. Tree Physiology, 21, 223-
232.

Dubois, J.-J.B., Fiscus, E.L., Booker, F.L., Flowers, M.D. & Reid, C.D. (2007) Optimizing the statistical
estimation of the parameters of the farquhar–von caemmerer–berry model of photosynthesis. New
Phytologist, 176, 402-414.

Dungan, R.J., Whitehead, D. & Duncan, R.P. (2003) Seasonal and temperature dependence of
photosynthesis and respiration for two co-occurring broad-leaved tree species with contrasting leaf
phenology†. Tree Physiology, 23, 561-568.

Duranceau, M., Ghashghaie, J. & Brugnoli, E. (2001) Carbon isotope discrimination during
photosynthesis and dark respiration in intact leaves of <emph type="2">nicotiana
sylvestris</emph>: Comparisons between wild type and mitochondrial mutant plants. Functional
Plant Biology, 28, 65-71.

Eamus, D., Duff, G.A. & Berryman, C.A. (1995) Photosynthetic responses to temperature, light flux-
density, co2 concentration and vapour pressure deficit in eucalyptus tetrodonta grown under co2
enrichment. Environmental Pollution, 90, 41-49.

Eichelmann, H., et al. (2004) Photosynthetic parameters of birch (betula pendula roth) leaves
growing in normal and in co2- and o3- enriched atmospheres. Plant, Cell & Environment, 27, 479-
495.

Ellsworth, D.S., Laroche, J. & Hendrey, G.R. (1998) Elevated co2 in a prototype free-air co2
enrichment facility affects photosynthetic nitrogen relations in a maturing pine forest. . In, United
States.

Ellsworth, D.S., Oren, R., Huang, C., Phillips, N. & Hendrey, G.R. (1995) Leaf and canopy responses to
elevated co2 in a pine forest under free-air co2 enrichment. Oecologia, 104, 139-146.

Falge, E., Graber, W., Siegwolf, R. & Tenhunen, J.D. (1996) A model of the gas exchange response
of picea abies to habitat conditions. Trees, 10, 277-287.

Farazdaghi, H. (2011) The single-process biochemical reaction of rubisco: A unified theory and model
with the effects of irradiance, co2 and rate-limiting step on the kinetics of c3 and c4 photosynthesis
from gas exchange. Biosystems, 103, 265-284.

Farquhar, G.D., Von Caemmerer, S. & Berry, J.A. (1980) A biochemical model of photosynthetic co2
assimilation in leaves of c3 species. Planta, 149, 78-90.

Flexas, J., Ribas-Carbó, M., Hanson, D.T., Bota, J., Otto, B., Cifre, J., Mcdowell, N., Medrano, H. &
Kaldenhoff, R. (2006) Tobacco aquaporin ntaqp1 is involved in mesophyll conductance to co2in vivo.
The Plant Journal, 48, 427-439.

Gao, Q., Zhang, X., Huang, Y. & Xu, H. (2004) A comparative analysis of four models of
photosynthesis for 11 plant species in the loess plateau. Agricultural and Forest Meteorology, 126,
203-222.

Gerardeaux, E., Saur, E., Constantin, J., Porté, A. & Jordan-Meille, L. (2009) Effect of carbon
assimilation on dry weight production and partitioning during vegetative growth. Plant and Soil, 324,
329-343.

Grassi, G., Meir, P., Cromer, R., Tompkins, D. & Jarvis, P.G. (2002) Photosynthetic parameters in
seedlings of eucalyptus grandis as affected by rate of nitrogen supply. Plant, Cell & Environment, 25,
1677-1688.

Gross, L.J., Kirschbaum, M.U.F. & Pearcy, R.W. (1991) A dynamic model of photosynthesis in varying
light taking account of stomatal conductance, c3-cycle intermediates, photorespiration and rubisco
activation. Plant, Cell & Environment, 14, 881-893.

Gu, L., Pallardy, S.G., Tu, K., Law, B.E. & Wullschleger, S.D. (2010) Reliable estimation of biochemical
parameters from c3 leaf photosynthesis–intercellular carbon dioxide response curves. Plant, Cell &
Environment, 33, 1852-1874.

Guo, L.P., Kang, H.J., Ouyang, Z., Zhuang, W. & Yu, Q. (2015) Photosynthetic parameter estimations
by considering interactive effects of light, temperature and co2 concentration. International Journal
of Plant Production, 9, 321-346.

Hamada, S., Kumagai, T.O., Kochi, K., Kobayashi, N., Hiyama, T. & Miyazawa, Y. (2016) Spatial and
temporal variations in photosynthetic capacity of a temperate deciduous-evergreen forest. Trees,
30, 1083-1093.

Han, Q. & Chiba, Y. (2009) Leaf photosynthetic responses and related nitrogen changes associated
with crown reclosure after thinning in a young chamaecyparis obtusa stand. Journal of Forest
Research, 14, 349-357.

Han, Q., Kawasaki, T., Nakano, T. & Chiba, Y. (2004) Spatial and seasonal variability of temperature
responses of biochemical photosynthesis parameters and leaf nitrogen content within a pinus
densiflora crown. Tree Physiology, 24, 737-744.

Harley, P.C. & Sharkey, T.D. (1991) An improved model of c3 photosynthesis at high co2: Reversed
o2 sensitivity explained by lack of glycerate reentry into the chloroplast. Photosynthesis Research,
27, 169-178.

Harley, P.C. & Baldocchi, D.D. (1995) Scaling carbon dioxide and water vapour exchange from leaf to
canopy in a deciduous forest. I. Leaf model parametrization. Plant, Cell & Environment, 18, 1146-
1156.

Harley, P.C., Weber, J.A. & Gates, D.M. (1985) Interactive effects of light, leaf temperature, co2 and
o2 on photosynthesis in soybean. Planta, 165, 249-263.

Harley, P.C., Tenhunen, J.D. & Lange, O.L. (1986) Use of an analytical model to study limitations on
net photosynthesis in arbutus unedo under field conditions. Oecologia, 70, 393-401.

Harley, P.C., Thomas, R.B., Reynolds, J.F. & Strain, B.R. (1992) Modelling photosynthesis of cotton
grown in elevated co2. Plant, Cell & Environment, 15, 271-282.

Harnos, N., Tuba, Z. & Szente, K. (2002) Modelling net photosynthetic rate of winter wheat in
elevated air $co_2$ concentrations. Photosynthetica, 40, 293-300.

Ho, Q.T., Verboven, P., Yin, X., Struik, P.C. & Nicolaï, B.M. (2012) A microscale model for combined
co2 diffusion and photosynthesis in leaves. PLOS ONE, 7, e48376.

Iio, A., Fukasawa, H., Nose, Y., Naramoto, M., Mizunaga, H. & Kakubari, Y. (2009) Within-branch
heterogeneity of the light environment and leaf temperature in a fagus crenata crown and its
significance for photosynthesis calculations. Trees, 23, 1053-1064.

Jarvis, P.G. (1999) Predicted impacts of rising carbon dioxide and temperature on forests in europe
at stand scale. In: ECOCRAFT ENVIRONMENT R&D ENV4-CT95-0077 IC20-CT96-0028

Juurola, E., Aalto, T., Thum, T., Vesala, T. & Hari, P. (2005) Temperature dependence of leaf-level co2
fixation: Revising biochemical coefficients through analysis of leaf three-dimensional structure. New
Phytologist, 166, 205-215.

Katahata, S.-I., Naramoto, M., Kakubari, Y. & Mukai, Y. (2007) Photosynthetic capacity and nitrogen
partitioning in foliage of the evergreen shrub daphniphyllum humile along a natural light gradient.
Tree Physiology, 27, 199-208.

Kattge, J. & Knorr, W. (2007) Temperature acclimation in a biochemical model of photosynthesis: A
reanalysis of data from 36 species. Plant, Cell & Environment, 30, 1176-1190.

Kattge, J., et al. (2020) Try plant trait database – enhanced coverage and open access. Global Change
Biology, 26, 119-188.

Kellomäki, S. & Wang, K.-Y. (1996) Photosynthetic responses to needle water potentials in scots pine
after a four-year exposure to elevated co2 and temperature. Tree Physiology, 16, 765-772.

Kim, J.H., Lee, J.W., Ahn, T.I., Shin, J.H., Park, K.S. & Son, J.E. (2016) Sweet pepper (capsicum annuum
l.) canopy photosynthesis modeling using 3d plant architecture and light ray-tracing. Frontiers in
Plant Science, 7

Kim, S.H. & Lieth, J.H. (2003) A coupled model of photosynthesis, stomatal conductance and
transpiration for a rose leaf (rosa hybrida l.). Annals of Botany, 91, 771-781.

Kirschbaum, M. & Farquhar, G. (1984) Temperature dependence of whole-leaf photosynthesis in
*eucalyptus pauciflora* sieb. Ex spreng. Functional Plant Biology, 11, 519-538.

Korzukhin, M.D., Vygodskaya, N.N., Milyukova, I.M., Tatarinov, F.A. & Tsel'niker, Y.L. (2004)
Application of a coupled photosynthesis–stomatal conductance model to analysis of carbon
assimilation by spruce and larch trees in the forests of russia. Russian Journal of Plant Physiology, 51,
302-315.

Kosugi, Y. & Matsuo, N. (2006) Seasonal fluctuations and temperature dependence of leaf gas
exchange parameters of co-occurring evergreen and deciduous trees in a temperate broad-leaved
forest. Tree Physiology, 26, 1173-1184.

Kosugi, Y., Shibata, S. & Kobashi, S. (2003) Parameterization of the co2 and h2o gas exchange of
several temperate deciduous broad-leaved trees at the leaf scale considering seasonal changes.
Plant, Cell & Environment, 26, 285-301.

Krall, J.P., Sheveleva, E.V. & Pearcy, R.W. (1995) Regulation of photosynthetic induction state in high-
and low-light-grown soybean and alocasia macrorrhiza (l.) g. Don. Plant Physiology, 109, 307-317.

Kubiske, M.E., Zak, D.R., Pregitzer, K.S. & Takeuchi, Y. (2002) Photosynthetic acclimation of overstory
populus tremuloides and understory acer saccharum to elevated atmospheric co2 concentration:
Interactions with shade and soil nitrogen. Tree Physiology, 22, 321-329.

Kumagai, T.O., et al. (2006) Modeling co2 exchange over a bornean tropical rain forest using
measured vertical and horizontal variations in leaf-level physiological parameters and leaf area
densities. Journal of Geophysical Research: Atmospheres, 111

Lai, C.-T., Katul, G., Butnor, J., Siqueira, M., Ellsworth, D., Maier, C., Johnsen, K., Mckeand, S. & Oren,
R. (2002) Modelling the limits on the response of net carbon exchange to fertilization in a south-
eastern pine forest. Plant, Cell & Environment, 25, 1095-1120.

Le Roux, X., Grand, S., Dreyer, E. & Daudet, F.-A. (1999) Parameterization and testing of a
biochemically based photosynthesis model for walnut (juglans regia) trees and seedlings. Tree
Physiology, 19, 481-492.

Lenz, K.E., Host, G.E., Roskoski, K., Noormets, A., Sôber, A. & Karnosky, D.F. (2010) Analysis of a
farquhar-von caemmerer-berry leaf-level photosynthetic rate model for populus tremuloides in the
context of modeling and measurement limitations. Environmental Pollution, 158, 1015-1022.

Leuning, R. (1990) Modelling stomatal behaviour and and photosynthesis of *eucalyptus grandis*.
Functional Plant Biology, 17, 159-175.

Leverenz, J.W. & Jarvis, P.G. (1979) Photosynthesis in sitka spruce. Viii. The effects of light flux
density and direction on the rate of net photosynthesis and the stomatal conductance of needles.
Journal of Applied Ecology, 16, 919-932.

Li, J.-H., Dijkstra, P., Hinkle, C.R., Wheeler, R.M. & Drake, B.G. (1999) Photosynthetic acclimation to
elevated atmospheric co2 concentration in the florida scrub-oak species quercus geminata and
quercus myrtifolia growing in their native environment. Tree Physiology, 19, 229-234.

Li, L., Luo, G., Chen, X., Li, Y., Xu, G., Xu, H. & Bai, J. (2011) Modelling evapotranspiration in a central
asian desert ecosystem. Ecological Modelling, 222, 3680-3691.

Manter, D.K., Kavanagh, K.L. & Rose, C.L. (2005) Growth response of douglas-fir seedlings to nitrogen
fertilization: Importance of rubisco activation state and respiration rates. Tree Physiology, 25, 1015-
1021.

Maroco, J.P., Breia, E., Faria, T., Pereira, J.S. & Chaves, M.M. (2002) Effects of long-term exposure to
elevated co2 and n fertilization on the development of photosynthetic capacity and biomass
accumulation in quercus suber l. Plant, Cell & Environment, 25, 105-113.

Martin-Stpaul, N.K., Limousin, J.-M., Rodríguez-Calcerrada, J., Ruffault, J., Rambal, S., Letts, M.G. &
Misson, L. (2012) Photosynthetic sensitivity to drought varies among populations of *quercus ilex*
along a rainfall gradient. Functional Plant Biology, 39, 25-37.

Massonnet, C., Costes, E., Rambal, S., Dreyer, E. & Regnard, J.L. (2007) Stomatal regulation of
photosynthesis in apple leaves: Evidence for different water-use strategies between two cultivars.
Annals of Botany, 100, 1347-1356.

Matyssek, R., Günthardt-Goerg, M.S., Keller, T. & Scheidegger, C. (1991) Impairment of gas exchange
and structure in birch leaves (betula pendula) caused by low ozone concentrations. Trees, 5, 5-13.

Medlyn, B.E., Loustau, D. & Delzon, S. (2002) Temperature response of parameters of a
biochemically based model of photosynthesis. I. Seasonal changes in mature maritime pine (pinus
pinaster ait.). Plant, Cell & Environment, 25, 1155-1165.

Meir, P., Levy, P.E., Grace, J. & Jarvis, P.G. (2007) Photosynthetic parameters from two contrasting
woody vegetation types in west africa. Plant Ecology, 192, 277-287.

Meir, P., Kruijt, B., Broadmeadow, M., Barbosa, E., Kull, O., Carswell, F., Nobre, A. & Jarvis, P.G.
(2002) Acclimation of photosynthetic capacity to irradiance in tree canopies in relation to leaf
nitrogen concentration and leaf mass per unit area. Plant, Cell & Environment, 25, 343-357.

Merilo, E., Heinsoo, K., Kull, O., Söderbergh, I., Lundmark, T. & Koppel, A. (2006) Leaf photosynthetic
properties in a willow (salix viminalis and salix dasyclados) plantation in response to fertilization.
European Journal of Forest Research, 125, 93-100.

Midgley, G.F., Wand, S.J.E. & Pammenter, N.W. (1999) Nutrient and genotypic effects on co2-
responsiveness: Photosynthetic regulation in leucadendron species of a nutrient-poor environment.
Journal of Experimental Botany, 50, 533-542.

Montpied, P., Granier, A. & Dreyer, E. (2009) Seasonal time-course of gradients of photosynthetic
capacity and mesophyll conductance to co2 across a beech (fagus sylvatica l.) canopy. Journal of
Experimental Botany, 60, 2407-2418.

Myers, D.A., Thomas, R.B. & Delucia, E.H. (1999) Photosynthetic capacity of loblolly pine (pinus taeda
l.) trees during the first year of carbon dioxide enrichment in a forest ecosystem. Plant, Cell &
Environment, 22, 473-481.

Nijs, I., Behaeghe, T. & Impens, I. (1995) Leaf nitrogen content as a predictor of photosynthetic
capacity in ambient and global change conditions. Journal of Biogeography, 22, 177-183.

Nikolov, N.T., Massman, W.J. & Schoettle, A.W. (1995) Coupling biochemical and biophysical
processes at the leaf level: An equilibrium photosynthesis model for leaves of c3 plants. Ecological
Modelling, 80, 205-235.

Oliver, R.J., Taylor, G. & Finch, J.W. (2012) Assessing the impact of internal conductance to co2 in a
land-surface scheme: Measurement and modelling of photosynthesis in populus nigra. Agricultural
and Forest Meteorology, 152, 240-251.

Onoda, Y., Hikosaka, K. & Hirose, T. (2005) The balance between rubp carboxylation and rubp
regeneration: A mechanism underlying the interspecific variation in acclimation of photosynthesis to
seasonal change in temperature. Functional Plant Biology, 32, 903-910.

Osborne, C.P., Drake, B.G., Laroche, J. & Long, S.P. (1997) Does long-term elevation of co2
concentration increase photosynthesis in forest floor vegetation? (indiana strawberry in a maryland
forest). Plant Physiology, 114, 337-344.

Osborne, C.P., Roche, J.L., Garcia, R.L., Kimball, B.A., Wall, G.W., Pinter, P.J., Morte, R.L.L., Hendrey,
G.R. & Long, S.P. (1998) Does leaf position within a canopy affect acclimation of photosynthesis to
elevated co₂? <span class="subtitle">Analysis of a Wheat Crop under Free-Air
CO$_2$Enrichment</span>, 117, 1037-1045.

Patrick, L.D., Ogle, K. & Tissue, D.T. (2009) A hierarchical bayesian approach for estimation of
photosynthetic parameters of c3 plants. Plant, Cell & Environment, 32, 1695-1709.

Pearcy, R.W., Gross, L.J. & He, D. (1997) An improved dynamic model of photosynthesis for
estimation of carbon gain in sunfleck light regimes. Plant, Cell & Environment, 20, 411-424.

Peltier, D.M.P. & Ibáñez, I. (2015) Patterns and variability in seedling carbon assimilation:
Implications for tree recruitment under climate change. Tree Physiology, 35, 71-85.

Pettersson, R. & Mcdonald, A.J.S. (1992) Effects of elevated carbon dioxide concentration on
photosynthesis and growth of small birch plants (betula pendula roth.) at optimal nutrition. Plant,
Cell & Environment, 15, 911-919.

Porté, A. & Loustau, D. (1998) Variability of the photosynthetic characteristics of mature needles
within the crown of a 25-year-old pinus pinaster. Tree Physiology, 18, 223-232.

Prioul, J.L. & Chartier, P. (1977) Partitioning of transfer and carboxylation components of
intracellular resistance to photosynthetic co2 fixation: A critical analysis of the methods used. Annals
of Botany, 41, 789-800.

Qian, T., Elings, A., Dieleman, J.A., Gort, G. & Marcelis, L.F.M. (2012) Estimation of photosynthesis
parameters for a modified farquhar–von caemmerer–berry model using simultaneous estimation
method and nonlinear mixed effects model. Environmental and Experimental Botany, 82, 66-73.

Rayment, M.B., Loustau, D. & Jarvis, P.J. (2002) Photosynthesis and respiration of black spruce at
three organizational scales: Shoot, branch and canopy. Tree Physiology, 22, 219-229.

Rey, A. & Jarvis, P.G. (1998) Long-term photosynthetic acclimation to increased atmospheric co2
concentration in young birch (betula pendula) trees. Tree Physiology, 18, 441-450.

Robakowski, P., Montpied, P. & Dreyer, E. (2002) Temperature response of photosynthesis of silver
fir (abies alba mill.) seedlings. Ann. For. Sci., 59, 163-170.

Rodeghiero, M., Niinemets, Ü. & Cescatti, A. (2007) Major diffusion leaks of clamp-on leaf cuvettes
still unaccounted: How erroneous are the estimates of farquhar et al. Model parameters? Plant,
Cell & Environment, 30, 1006-1022.

Rodríguez-Calcerrada, J., Reich, P.B., Rosenqvist, E., Pardos, J.A., Cano, F.J. & Aranda, I. (2008) Leaf
physiological versus morphological acclimation to high-light exposure at different stages of foliar
development in oak. Tree Physiology, 28, 761-771.

Rogers, A., Fischer, B.U., Bryant, J., Frehner, M., Blum, H., Raines, C.A. & Long, S.P. (1998)
Acclimation of photosynthesis to elevated co₂under low-nitrogen nutrition is affected by the
capacity for assimilate utilization. Perennial ryegrass under free-air co₂ enrichment. Plant Physiology,
118, 683-689.

Schultz, H.R. (2003) Extension of a farquhar model for limitations of leaf photosynthesis induced by
light environment, phenology and leaf age in grapevines (<emph type="2">vitis</emph> <emph

type="2">vinifera</emph> l. Cvv. White riesling and zinfandel). Functional Plant Biology, 30, 673-
687.

Shipley, B. & Lechowicz, M.J. (2000) The functional co-ordination of leaf morphology, nitrogen
concentration, and gas exchange in40 wetland species. Écoscience, 7, 183-194.

Sholtis, J.D., Gunderson, C.A., Norby, R.J. & Tissue, D.T. (2004) Persistent stimulation of
photosynthesis by elevated co2 in a sweetgum (liquidambar styraciflua) forest stand. New
Phytologist, 162, 343-354.

Sims, D.A., Luo, Y. & Seemann, J.R. (1998) Comparison of photosynthetic acclimation to elevated co2
and limited nitrogen supply in soybean. Plant, Cell & Environment, 21, 945-952.

Su, Y., Zhu, G., Miao, Z., Feng, Q. & Chang, Z. (2009) Estimation of parameters of a biochemically
based model of photosynthesis using a genetic algorithm. Plant, Cell & Environment, 32, 1710-1723.

Sun, J., Sun, J. & Feng, Z. (2015) Modelling photosynthesis in flag leaves of winter wheat (*triticum*
*aestivum*) considering the variation in photosynthesis parameters during development. Functional
Plant Biology, 42, 1036-1044.

Thum, T., Aalto, T., Laurila, T., Aurela, M., Kolari, P. & Hari, P. (2007) Parametrization of two
photosynthesis models at the canopy scale in a northern boreal scots pine forest. Tellus B: Chemical
and Physical Meteorology, 59, 874-890.

Tissue, D.T., Griffin, K.L. & Ball, J.T. (1999) Photosynthetic adjustment in field-grown ponderosa pine
trees after six years of exposure to elevated co2. Tree Physiology, 19, 221-228.

Tissue, D.T., Griffin, K.L., Turnbull, M.H. & Whitehead, D. (2005) Stomatal and non-stomatal
limitations to photosynthesis in four tree species in a temperate rainforest dominated by dacrydium
cupressinum in new zealand. Tree Physiology, 25, 447-456.

Tosens, T., et al. (2016) The photosynthetic capacity in 35 ferns and fern allies: Mesophyll co2
diffusion as a key trait. New Phytologist, 209, 1576-1590.

Turnbull, M.H., Tissue, D.T., Griffin, K.L., Rogers, G.N.D. & Whitehead, D. (1998) Photosynthetic
acclimation to long-term exposure to elevated co2 concentration in pinus radiata d. Don. Is related
to age of needles. Plant, Cell & Environment, 21, 1019-1028.

Turnbull, T.L., Adams, M.A. & Warren, C.R. (2007) Increased photosynthesis following partial
defoliation of field-grown eucalyptus globulus seedlings is not caused by increased leaf nitrogen.
Tree Physiology, 27, 1481-1492.

Uddling, J. & Wallin, G. (2012) Interacting effects of elevated co2 and weather variability on
photosynthesis of mature boreal norway spruce agree with biochemical model predictions. Tree
Physiology, 32, 1509-1521.

Vaz, M., Pereira, J.S., Gazarini, L.C., David, T.S., David, J.S., Rodrigues, A., Maroco, J. & Chaves, M.M.
(2010) Drought-induced photosynthetic inhibition and autumn recovery in two mediterranean oak
species (quercus ilex and quercus suber). Tree Physiology, 30, 946-956.

Von Caemmerer, S., Evans, J.R., Hudson, G.S. & Andrews, T.J. (1994) The kinetics of ribulose-1,5-
bisphosphate carboxylase/oxygenase in vivo inferred from measurements of photosynthesis in
leaves of transgenic tobacco. Planta, 195, 88-97.

Walcroft, A., Le Roux, X., Diaz-Espejo, A., Dones, N. & Sinoquet, H. (2002) Effects of crown
development on leaf irradiance, leaf morphology and photosynthetic capacity in a peach tree. Tree
Physiology, 22, 929-938.

Walcroft, A.S., Whitehead, D., Silvester, W.B. & Kelliher, F.M. (1997) The response of photosynthetic
model parameters to temperature and nitrogen concentration in pinus radiata d. Don. Plant, Cell &
Environment, 20, 1338-1348.

Walker, B., Ariza, L.S., Kaines, S., Badger, M.R. & Cousins, A.B. (2013) Temperature response of in
vivo rubisco kinetics and mesophyll conductance in arabidopsis thaliana: Comparisons to nicotiana
tabacum. Plant, Cell & Environment, 36, 2108-2119.

Wang, K.-Y., Kellomäki, S. & Laitinen, K. (1996) Acclimation of photosynthetic parameters in scots
pine after three years exposure to elevated temperature and co2. Agricultural and Forest
Meteorology, 82, 195-217.

Wang, Q., Fleisher, D.H., Timlin, D., Reddy, V.R. & Chun, J.A. (2012) Quantifying the measurement
errors in a portable open gas-exchange system and their effects on the parameterization of farquhar
et al. Model for $c_3$ leaves. Photosynthetica, 50, 223-238.

Warren, C.R. (2004) The photosynthetic limitation posed by internal conductance to co2 movement
is increased by nutrient supply. Journal of Experimental Botany, 55, 2313-2321.

Warren, C.R. & Adams, M.A. (2001) Distribution of n, rubisco and photosynthesis in pinus pinaster
and acclimation to light. Plant, Cell & Environment, 24, 597-609.

Watanabe, M., Watanabe, Y., Kitaoka, S., Utsugi, H., Kita, K. & Koike, T. (2011) Growth and
photosynthetic traits of hybrid larch f1 (larix gmelinii var. Japonica×l. Kaempferi) under elevated
co2 concentration with low nutrient availability. Tree Physiology, 31, 965-975.

Weise, S.E., Carr, D.J., Bourke, A.M., Hanson, D.T., Swarthout, D. & Sharkey, T.D. (2015) The arc
mutants of arabidopsis with fewer large chloroplasts have a lower mesophyll conductance.
Photosynthesis Research, 124, 117-126.

Whitehead, D., Walcroft, A.S., Scott, N.A., Townsend, J.A., Trotter, C.M. & Rogers, G.N.D. (2004)
Characteristics of photosynthesis and stomatal conductance in the shrubland species mānuka
(leptospermum scoparium) and kānuka (kunzea ericoides) for the estimation of annual canopy
carbon uptake. Tree Physiology, 24, 795-804.

Wilson, K.B., Baldocchi, D.D. & Hanson, P.J. (2000) Spatial and seasonal variability of photosynthetic
parameters and their relationship to leaf nitrogen in a deciduous forest. Tree Physiology, 20, 565-
578.

Wohlfahrt, G., Bahn, M., Haubner, E., Horak, I., Michaeler, W., Rottmar, K., Tappeiner, U. &
Cernusca, A. (1999) Inter-specific variation of the biochemical limitation to photosynthesis and
related leaf traits of 30 species from mountain grassland ecosystems under different land use. Plant,
Cell & Environment, 22, 1281-1296.

Wullschleger, S.D. (1993) Biochemical limitations to carbon assimilation in c3 plants—a retrospective
analysis of the a/ci curves from 109 species. Journal of Experimental Botany, 44, 907-920.

Yarkhunova, Y., Edwards, C.E., Ewers, B.E., Baker, R.L., Aston, T.L., Mcclung, C.R., Lou, P. & Weinig, C.
(2016) Selection during crop diversification involves correlated evolution of the circadian clock and
ecophysiological traits in brassica rapa. New Phytologist, 210, 133-144.

Yin, X., Van Oijen, M. & Schapendonk, A.H.C.M. (2004) Extension of a biochemical model for the
generalized stoichiometry of electron transport limited c3 photosynthesis. Plant, Cell &
Environment, 27, 1211-1222.

Yin, X., Struik, P.C., Romero, P., Harbinson, J., Evers, J.B., Van Der Putten, P.E.L. & Vos, J. (2009) Using
combined measurements of gas exchange and chlorophyll fluorescence to estimate parameters of a
biochemical c3 photosynthesis model: A critical appraisal and a new integrated approach applied to
leaves in a wheat (triticum aestivum) canopy. Plant, Cell & Environment, 32, 448-464.

Zhang, S. & Dang, Q.-L. (2006) Effects of carbon dioxide concentration and nutrition on
photosynthetic functions of white birch seedlings. Tree Physiology, 26, 1457-1467.

Zhu, G.-F., Li, X., Su, Y.-H., Lu, L., Huang, C.-L. & Niinemets, Ü. (2011) Seasonal fluctuations and
temperature dependence in photosynthetic parameters and stomatal conductance at the leaf scale
of populus euphratica oliv. Tree Physiology, 31, 178-195.

Zhu, G.F., Li, X., Su, Y.H. & Huang, C.L. (2010) Parameterization of a coupled $CO_2$ and $H_2O$ gas
exchange model at the leaf scale of *populus euphratica*. Hydrol. Earth Syst. Sci., 14, 419-431.

1077