Detection of a reservoir of bedaquiline / clofazimine resistance associated variants in *Mycobacterium tuberculosis* predating the antibiotic era

Running title: Emergence of bedaquiline resistance in tuberculosis

Lucy van Dorp[1]*, Camus Nimmo[1,2,3]*, Arturo Torres Ortiz[1,4], Juanita Pang[1,2], Mislav Acman[1], Cedric C.S. Tan[1], James Millard[3,5,6], Nesri Padayatchi[7], Alison Grant[3,8], Max O'Donnell[7,9], Alex Pym[3], Ola B Brynildsrud[10], Vegard Eldholm[10], Louis Grandjean[2,11,12], Xavier Didelot[13], François Balloux[1]

*These authors contributed equally.

Correspondence: Camus Nimmo (camus.nimmo@crick.ac.uk), Lucy van Dorp (lucy.dorp.12@ucl.ac.uk) and François Balloux (f.balloux@ucl.ac.uk)

1. UCL Genetics Institute, University College London, London, UK
2. Division of Infection and Immunity, University College London, London, UK
3. Africa Health Research Institute, Durban, South Africa
4. Department of Medicine, Imperial College, London, UK
5. Wellcome Trust Liverpool Glasgow Centre for Global Health Research, Liverpool, UK
6. Institute of Infection and Global Health, University of Liverpool, Liverpool, UK
7. CAPRISA MRC-HIV-TB Pathogenesis and Treatment Research Unit, Durban, South Africa
8. TB Centre, London School of Hygiene & Tropical Medicine, London, UK
9. Department of Medicine & Epidemiology, Columbia University Irving Medical Center, New York, NY, USA
10. Division of Infectious Diseases and Environmental Health, Norwegian Institute of Public Health, Oslo, Norway
11. Laboratorio de Investigacion y Enfermedades Infecciosas/Universidad Peruana Cayetano Heredia, Lima, Peru
12. Department of Infection, Immunity and Inflammation, Institute of Child Health, University College London, London, UK
13. School of Life Sciences and Department of Statistics, University of Warwick, Coventry, UK

Keywords: Tuberculosis, phylogenetics, bedaquiline, drug resistance, AMR

Abstract

Drug resistance in tuberculosis (TB) poses a major ongoing challenge to public health. The recent inclusion of bedaquiline into TB drug regimens has improved treatment outcomes, but this advance is threatened by the emergence of strains of *Mycobacterium tuberculosis* (*Mtb*) resistant to bedaquiline. Clinical bedaquiline resistance is most frequently conferred by off-target resistance-associated variants (RAVs) in the *mmpR5* gene (*Rv0678*), the regulator of an efflux pump, which can also confer cross-resistance to clofazimine, another TB drug. We compiled a dataset of 3,682 *Mtb* genomes, including 157 carrying variants in *mmpR5* that have been linked to resistance. We identified nine cases where RAVs were present in the genomes of strains collected prior to the use of bedaquiline in TB treatment regimes. Phylogenetic reconstruction points to multiple emergence events and circulation of RAVs in *mmpR5*, some estimated to predate the introduction of bedaquiline or clofazimine. However, epistatic interactions can complicate bedaquiline drug-susceptibility prediction from genetic sequence data. Indeed, in one clade of isolates where the RAV Ile67fs is estimated to have emerged prior to the antibiotic era, co-occurrence of mutations in *mmpL5* are found to neutralise bedaquiline resistance. The presence of a pre-existing reservoir of *Mtb* strains carrying bedaquiline RAVs augments the need for rapid drug susceptibility testing and individualised regimen selection to safeguard the use of bedaquiline in TB care and control.

Introduction

Drug-resistant tuberculosis (DR-TB) currently accounts for 450,000 of the 10 million new tuberculosis (TB) cases reported annually[1]. Treatment outcomes for multidrug-resistant TB (MDR-TB), resistant to at least rifampicin and isoniazid, have historically been poor, with treatment success rates of only 50-60% in routine programmatic settings[2,3]. The discovery of bedaquiline, a diarylquinoline antimycobacterial active against ATP synthase, which is highly effective against *Mycobacterium tuberculosis* (*Mtb*)[4], was reported in 2004. Following clinical trials, which confirmed reduced time to culture conversion in patients with DR-TB[5], in 2012 bedaquiline received an accelerated Food and Drug Administration (FDA) licence for use in DR-TB[6].

Cohort studies of patients treated with bedaquiline-containing regimens against MDR-TB report success rates of 70-80%[7,8]. Similar results have been achieved for extensively drug-resistant TB (XDR-TB, traditionally defined as MDR-TB plus by additional resistance to fluoroquinolones and injectables), where treatment outcomes without bedaquiline are even worse[9,10]. In light of these promising results, the World Health Organization (WHO) now recommends that bedaquiline be included in all MDR-TB regimens[11]. It has played a central role in the highly successful ZeNix[12] and TB-PRACTECAL[13] trials of bedaquiline, pretomanid and linezolid (+/- moxifloxacin) six-month all-oral regimens for DR-TB. These are now incorporated in WHO guidance. In addition, bedaquiline is positioned as a key drug in multiple phase III clinical trials for drug-susceptible TB (SimpliciTB, ClinicalTrials.gov NCT03338621; TRUNCATE-TB[14]).

Resistance in *Mtb* is typically reported soon after the introduction of a novel TB drug and often sequentially[15,16]. For example, mutations conferring resistance to isoniazid – one of the first antimycobacterials – tend to have emerged prior to resistance to rifampicin, the other major first-line drug. These also predate resistance mutations to second-line drugs, so termed because they are used clinically to treat patients infected with strains already resistant to first-line drugs. This was observed, for example, in KwaZulu-Natal, South Africa, where resistance-associated mutations accumulated

over decades prior to their identification, leading to a major outbreak of extensively drug-resistant TB

(XDR-TB)[16]. Unlike other major drug-resistant bacteria, *Mtb* reproduces strictly clonally and

systematically acquires resistance by chromosomal mutations rather than via horizontal gene transfer

or recombination[17]. This allows phylogenetic reconstructions, based on whole genome sequencing

data, to be used to infer the timings of emergence and subsequent spread of *Mtb* resistance-associated

variants (RAVs).

A number of mechanisms have been implicated in conferring bedaquiline resistance. For example,

mutations conferring resistance have been selected *in vitro*, located in the *atpE* gene encoding the

target F1F0 ATP synthase, the target of bedaquiline[18]. Off-target resistance-conferring mutations have

also been found in *pepQ* in a murine model and potentially in a small number of patients[19]. However,

the primary mechanism of resistance observed in clinical isolates has been identified in the context of

off-target resistance-associated variants (RAVs) in the *mmpR5 (Rv0678)* gene, a negative repressor of

expression of the MmpL5 efflux pump. Loss of function of mmpR5 leads to pump overexpression[20]

and increased minimum inhibitory concentrations (MIC) to bedaquiline, along with the recently

repurposed antimycobacterial clofazimine, fusidic acid, the azole class of antifungal drugs (which also

have antimycobacterial activity), as well as to the novel therapeutic class of DprE1 inhibitors in

clinical trials[21,22]. Aligned with this mechanism of resistance, coincident mutations leading to loss of

function of the MmpL5 efflux pump can negate the resistance-inducing effect of MmpR5 loss of

function[23].

A range of single nucleotide polymorphisms (SNPs) and frameshift *mmpR5* mutations have been

associated with resistance to bedaquiline, and are often present as heteroresistant alleles in patients[24-

32]. In contrast to most other RAVs in *Mtb,* which often cause many-fold increases in MIC and clear-

cut resistance, *mmpR5* variants may be associated with normal MICs or subtle increases in

bedaquiline MIC, although they may still be clinically important[33]. These increases may not cross the

current WHO critical concentrations used to classify resistant versus susceptible strains (0.25µg/mL

on Middlebrook 7H11 agar, or 1µg/mL in Mycobacteria Growth Indicator Tube [MGIT] liquid

media). The current WHO tuberculosis drug resistance catalogue does not contain any bedaquiline

RAVs, although a meta-analysis has identified two RAVs (a*tpE* Ala63Pro and *mmpR5* Ile67fs)[34].

Bedaquiline has a long terminal half-life of up to 5.5 months[6], leading to the possibility of

subtherapeutic concentrations where adherence is suboptimal or treatment is interrupted, which could

act as a further driver of resistance.

Bedaquiline and clofazimine cross-resistance has now been reported across three continents following

the rapid expansion in usage of both drugs[25,30,35,36], and is associated in some cases with poor

adherence to therapy and inadequate regimens. However, baseline isolates in 8/347 (2.3%) patients

from phase IIb bedaquiline trials demonstrated *mmpR5* RAVs and high bedaquiline MICs in the

absence of prior documented use of bedaquiline or clofazimine[37]. This suggests that bedaquiline

RAVs were in circulation prior to the usage of either of these drugs, which may be expected in the

case of mutations which do not have major fitness consequences[38]. While there have been isolated

clinical reports from multiple geographical regions, the global extent of bedaquiline resistance

emergence and spread has not yet been investigated.

In this study, we characterise and date the emergence of bedaquiline RAVs in the two global *Mtb*

lineage 2 (L2) and lineage 4 (L4) lineages, which include the majority of drug resistance strains[15].

Phylogenetic analyses of two datasets comprising 1,514 *Mtb* L2 and 2,168 L4 whole genome

sequences revealed the emergence and spread of multiple *mmpR5* RAVs prior to the use of

bedaquiline or clofazimine, with some mutations estimated to have been in circulation prior to the

antibiotic era. This pre-existing reservoir of bedaquiline/clofazimine-resistant *Mtb* strains suggests

mmpR5 RAVs exert a relatively low fitness cost which could be rapidly selected for as bedaquiline

and clofazimine are more widely used in the treatment of TB.

Results

The global diversity of *Mtb* lineage L2 and L4

To investigate the global distribution of *Mtb* isolates with variants in *mmpR5*, we curated two large datasets of whole genomes from the two dominant global lineages L2 and L4. Both datasets were enriched for samples with variants in *mmpR5* following a screen for variants in public sequencing repositories (see **Methods**) and retaining those samples uploaded with accompanying full metadata for geolocation and time of sampling (**Figure 1, Supplementary Table S1-S2, Supplementary Figure S1**). The final L2 dataset included 1,514 isolates collected over 24.5 years (between 1994 and 2019) yielding 29,205 SNPs. The final L4 dataset comprised 2,168 sequences collected over 232 years, including three samples from 18[th] century Hungarian mummies[39], encompassing 67,585 SNPs. Both datasets included recently generated data from South Africa (155 L2, 243 L4)[40,41] and new whole genome sequencing data from Peru (9 L2, 154 L4).

Consistent with previous studies[42-44], both datasets are highly diverse and exhibit strong geographic structure (**Figure 2**). As a nonrecombining clonal organism, identification of mutations in *Mtb* can provide a mechanism to predict phenotypic resistance from a known panel of genotypes[45,46]. Based on genotypic profiling[46], 911 strains within the L2 dataset were classified as MDR-TB (60%) and 295 (20%) as XDR-TB. Within the L4 dataset, 911 isolates were classified as MDR-TB (42%) and 115 as XDR-TB (5%). The full phylogenetic distribution of resistance profiles is provided in **Supplementary Figure S2**. As is commonplace with genomic datasets, these percentages of drug-resistant strains exceed their actual prevalence, due to the overrepresentation of drug-resistant isolates in public repositories.

Both the L2 and L4 phylogenetic trees displayed a significant temporal signal following date randomisation (**Supplementary Figure S3**), making them suitable for time-calibrated phylogenetic inference[47]. We estimated the time to the Most Recent Common Ancestor (tMRCA) of both datasets using a Bayesian tip-dating analysis (BEAST2) run on a representative subset of genomes from each

dataset (see **Methods**, **Supplementary Table 3**, **Supplementary Figure S4**). For the final temporal

calibration of the L2 dataset we applied an estimated clock rate of 7.7×10^{-8} (4.9×10^{-8} - 1.03×10^{-7})

substitutions per site per year, obtained from the subsampled BEAST2[48] analysis, to the global

maximum likelihood phylogenetic tree. This resulted in an estimated tMRCA of 1332CE (945CE-

1503CE). Using the same approach for the L4 dataset we estimated a clock rate of 7.1×10^{-8} (6.2×10^{-8} -

7.9×10^{-8}) substitutions per site per year resulting in an estimated tMRCA of 853CE (685CE – 967CE)

(**Figure 2**). We observed a slightly higher, yet statistically not significant, clock rate in L2 compared

to L4 (**Supplementary Table S3**), with all estimated substitution rates falling largely in line with

previously published estimates[48].

Identification of *mmpR5* variants

Since *atpE* and *pepQ* bedaquiline RAVs are found at low prevalence (1 L2 isolate [0.03%] and 18 L4

isolates [0.49%]), we focused on characterising *mmpR5*. In total we identified the presence of non-

synonymous and promoter *mmpR5* variants in 438 sequences (194 L2 [12.8%], 244 L4 [11.3%]). We

classified all identified non-synonymous and promoter mutations in *mmpR5*, based on the literature,

into six phenotypic categories for bedaquiline susceptibility: wild type, hypersusceptible, susceptible,

intermediate, resistant and unknown (full references available in **Supplementary Table S4,**

Supplementary Figures S5-S7). Across both lineages, 240 sequences were initially considered as

bedaquiline resistant (i.e., classified as intermediate or resistant based solely on their *mmpR5*

genotype). The most frequently observed variants are listed in **Table 1**.

We identified a significant relationship between the presence of *mmpR5* variants and drug resistance

status in both the L2 and L4 datasets (**Supplementary Figure S8-S9**), though in both cases we

identified otherwise fully phenotypically susceptible isolates carrying *mmpR5* RAVs (12 L2, 25 L4).

Notably we identified 25 sequenced isolates carrying nonsynonymous variants in *mmpR5* uploaded

with collection dates prior to the first clinical trials for bedaquiline in 2007. This comprised ten L2

isolates collected before 2007, of which eight harboured variants previously associated to phenotypic

bedaquiline resistance (RAVs). For L4 we identified 15 sequences with *mmpR5* variants predating

2007, of which one has previously been associated with phenotypic bedaquiline resistance (RAVs)

and six were classified as conferring an intermediate resistance phenotype (**Figure 1c-d,**

Supplementary Table S5).

Within the datasets, we identified one L2 isolate (ERR2677436 sampled in Germany in 2016) which

already had two *mmpR5* RAVs at low allele frequency – Val7fs (11%) and Val20Phe (20%) – and

also contained two low frequency *atpE* RAVs: Glu61Asp (3.2%) and Alal63Pro (3.7%)[50]. We also

identified three isolates obtained in 2007-08 from separate but neighbouring Chinese provinces

carrying the *Rv1979c* Val52Gly RAV, which has been suggested to be associated with clofazimine

resistance in a study from China[25] but was associated with a normal MIC in another[38], with its role in

resistance remaining unclear[31]. Furthermore, frameshift and premature stop mutations in *pepQ* have

been previously associated with bedaquiline and clofazimine resistance. In this dataset, we identified

18 frameshift mutations in *pepQ* across 11 patients, one of which also had a *mmpR5* frameshift

mutation. In one isolate the *pepQ* frameshift occurred at the Arg271 position previously reported to be

associated with bedaquiline resistance[19].

Sixty-three genomes harboured nonsynonymous *mmpR5* variants of unknown phenotypic effect (12

L2, 51 L4), corresponding to 23 unique mutations or combinations of mutations. To assess properties

associated to RAVs which may be useful predictors of the phenotypic effect of these unknown

variants we employed a machine learning approach, providing a foundation for further exploration of

genomic features associated to RAV status (see **Supplementary Note 1**).

The time to emergence of *mmpR5* variants

To estimate the age of the emergence of different *mmpR5* non-synonymous variants, we identified all

nodes in each of the L2 and L4 global time calibrated phylogenies delineating clades of isolates

carrying a particular *mmpR5* variant (**Figure 3, Supplementary Table S6, Supplementary Table**

S7). For the L2 dataset we identified 58 unique phylogenetic nodes where *mmpR5* RAVs emerged, of

which 40 were represented by a single genome. The point estimates for these nodes ranged from

March 1845 to November 2018. Eight nonsynonymous variants in *mmpR5*, including four confirmed

bedaquiline RAVs, were estimated to have emergence dates (point estimates) predating the first

bedaquiline clinical trial in 2007 (**Supplementary Figure S10**).

For the L4 dataset we identified 85 unique nodes where *mmpR5* RAVs emerged, of which 59 were

represented by a single isolate in the dataset. The point estimates for these nodes ranged from

September 1701 to January 2019 (**Figure 3, Supplementary Figure S11**). Sixteen *mmpR5* mutations,

including six bedaquiline RAVs and two predicted to have an intermediate phenotype, were estimated

to have emerged prior to 2007. This included a *mmpR5* Asp141fs carrying clade predominately

sampled in Uganda we estimate to have emerged in 1857 (1835-1872). We also identified one large

clade of 65 samples, predominantly collected in Peru (henceforth Peruvian clade), which all carry the

Ile67fs *mmpR5* RAV[35,49,50]. While it is not inconceivable that multiple independent emergences of

Ile67fs occurred in this clade, the more parsimonious scenario is a single ancestral emergence. We

estimate the time of this emergence to 1702 (1657-1732). Both these examples significantly predate

the first use of azoles, clofazimine or indeed bedaquiline (**Figure 3, Supplementary Figure S11-**

S12).

Of significance, we identified a frameshift mutation in the adjacent MmpL5 efflux pump (Arg202fs)

in isolates from the Peruvian clade, the protein whose overexpression mediates bedaquiline resistance

following loss-of-function of the MmpR5 regulatory protein. This frameshift, which leads to a

premature stop codon at amino acid 206, is expected to counteract the otherwise resistance-conferring

mutation. This epistatic interaction restoring bedaquiline susceptibility has recently been described

elsewhere[23,38]. The *mmpL5* frameshift mutation was present in all isolates in the Peruvian clade bar

one (ERR7339051) which had *mmpL5* Arg202Leu. This event of reading-frame restoration is likely

explained by a recent secondary duplication of a T downstream of the initial deletion (777876

GGCAT > GGAT, GGAT > GGATT). We considered the phenotype of this strain as unknown. No

other *mmpL5* mutations were found in any isolate containing *mmpR5* mutations within this study

though we did identify a low prevalence of variants in *mmpL5* and *mmpS5* independent of *mmpR5*

mutations across both lineages (**Supplementary Figures S13 and S14**).

While we identified no nodes with a second emergence of *mmpR5* nonsynonymous mutations across

the L4 dataset, eight nodes were identified in the L2 dataset where a clade already carrying a

nonsynonymous variant in *mmpR5* subsequently acquired a second nonsynonymous mutation.

Phenotypic validation of *mmpR5* variants

Given documented epistasis as a modulator of bedaquiline resistance phenotype, we performed MIC

testing on a selection of available isolates and identified further MICs that have been recently

published as part of the Cryptic consortium using microtitre plates (**Supplementary Table S7**)[51,52].

The epidemiological cut-off (ECOFF, defined as MIC of 95-99% of wild-type isolates) for

bedaquiline has been proposed to be 0.12 or 0.25 μg/mL depending on the method used, although the

final decision was to use an ECOFF of 0.25 μg/mL[52].

We were able to identify 30 L4 isolates from Peru (including the Peruvian cluster) for MIC testing,

and a further 9 MICs for L4 that had recently been published by the Cryptic consortium[51]. For the

oldest dated *mmpR5* mutation emergence – the L4 Ile67fs mutation in Peruvian isolates with an

associated MRCA estimated to 1701 - 10/11 (90.9%) had an MIC below the lower proposed ECOFF

of 0.12 μg/mL, presumably due to the co-existing *mmpL5* loss of function mutation. Hence, we

denote isolates from this clade as having a hypersusceptible phenotype. The second oldest predicted

resistance mutation (Arg90Cys, dated to 1940) was however associated with MICs ≥0.12 μg/mL in

6/7 (85.7%) instances, and in 3/4 (75%) instances for the third oldest predicted resistance mutation for

which data were available (Asn98Asp, dated to 1987). These MICs are above the wild-type range, if

not formally classified as resistant. Clades with associated MIC confirmation are highlighted in Figure

3b.

Table 1: Frequency of all *mmpR5* variants occurring ≥ 5 times in dataset and predicted resistance
classification. *where co-existing *mmpL5* mutations were identified this is indicated – only one
mutation was found (*mmpL5* Arg202fs) and it was in the presence of *mmpR5* Ile67fs mutations only,
with no other co-existing *mmpR5* variants.
268
269

Variant	Predicted phenotype	L2	L4	Total
Ile67fs	Resistant	13	18	97
Ile67fs + mmpL5 Arg202fs*	Hypersusceptible	0	66	66
c-11a	Hypersusceptible	93	0	93
Asp5Gly	Susceptible	20	3	23
Met146Thr	Resistant	2	20	22
Leu40Val	Unknown	0	19	19
Glu49fs	Resistant	3	9	12
Arg90Cys	Resistant	2	9	11
Asp47fs	Resistant	8	1	9
Ala59Val	Resistant	7	1	8
Asn98Asp	Resistant	0	7	7
Leu117Arg	Resistant	2	5	7
Val1Ala	Resistant	7	0	7
Val20Ala	Intermediate	1	6	7
Arg156fs	Resistant	0	6	6
Asp141fs	Resistant	0	6	6
Gly121Arg	Resistant	5	1	6
Arg109Leu	Resistant	0	5	5
Arg96Gly	Unknown	0	5	5

Discussion

Our work establishes that the emergence of variants in *mmpR5*, including RAVs, is not solely driven by the use of bedaquiline, clofazimine, azoles or fusidic acid. In particular, we identified 12 cases of emergence of bedaquiline RAVs prior to the first clinical trials of bedaquiline in 2007. Phylogenetic inference estimated the oldest clade containing *mmpR5* mutations, composed mostly of samples from Peru carrying the Ile67fs RAV, to have emerged around 1702 (1657-1732). We identify two further emergences of *mmpR5* mutations occurred prior to the antibiotic era (including the use of clofazimine, fusidic acid and azole antifungals), estimated to between 1871 and 1940, with samples from the latter clade confirmed to have MICs above the wild-type range. Our phylogenetic inference, pointing to multiple emergences of *mmpR5* nonsynonymous variants predating the use of bedaquiline, is also confirmed by the direct observation of 15 *Mtb* genomes carrying *mmpR5* RAVs sampled prior to 2007. We also identified, within the aforementioned Peruvian clade, a consistent frameshift mutation in *mmpL5*, which seemed to counteract the resistance phenotypic through an epistatic interaction (MIC <0.12 μg/mL). While Ile67fs is central for bedaquiline resistance in *Mtb*, its impact is influenced by the strain genetic background.

We identified other localised clusters with *mmpR5* mutations, reinforcing the need for concern even in situations where such mutations are globally rare. This included 19 isolates with a Met146Thr mutation found in lineage 4 isolates from Eswatini and 19 isolates with a Leu40Val mutation found in Peru. Met146Thr mutations have been previously associated with a clade that has a rifampicin-resistance conferring mutation located outside of the canonical rifampicin-resistance determining region, and these isolates exhibit elevated bedaquiline MICs[53]. The emergence of the Met146Thr mutation has previously been dated to have emerged in approximately 2003[23,38,53]. This is in reasonable agreement with our analysis on a much larger dataset which inferred an emergence in 2005.6 (95% confidence intervals 2004.8 – 2006.0). The long-standing presence of bedaquiline RAVs predating the use of the drug and at high prevalence in geographically notable cases is of concern, as it suggests that non-synonymous mutations in *mmpR5* exert little fitness cost. Hence our work

suggests the existence of pre-existent reservoirs of bedaquiline resistant *Mtb*, including in strains

which are otherwise fully susceptible strains. This reservoir is expected to expand under drug pressure

with the increasing use of bedaquiline and clofazimine in TB treatment. Further, these reservoirs may

also pose a threat for other candidate TB agents from different drug classes that are also exported by

mmpS5 and *mmpL5*[19,22,54].

The identification of resistance variants occurring before the use of a drug is not limited to *M.*

*tuberculosi*s and *mmpR5* alone. To illustrate, within *M. bovis*, there is evidence indicating that the

pncA H57D mutation, which is associated with resistance to pyrazinamide (PZA), emerged

approximately 900 years ago, providing inherent resistance to PZA in *M. bovis*[55]. Similarly, variations

in intrinsic susceptibility to pretomanid have been observed across the MTBC, including *Mtb*

lineages, even without prior exposure to nitroimidazoles[56]. It is likely that there are numerous other

instances of such loss of function mutations with minimal or no impact on fitness, similar to the case

of *mmpR5*. Furthermore, the existence of antimicrobial resistance (AMR) in different forms has

persisted throughout the natural history of various bacteria[57].

Nevertheless, it is crucial to determine the age and diversity of resistance variants to gain a better

understanding of the potential for widespread resistance as a contemporary challenge when the

selective pressure of a drug is introduced. For example, we identified a large number of different

mmpR5 nonsynonymous variants across both of our *Mtb* lineage cohorts; 45 in L2 (including 10

unique RAV combinations) and 67 in L4 (nine unique RAV combinations). This suggests the

mutational target leading to bedaquiline resistance is wider than for most other current TB drugs and

raises concerns about the ease with which bedaquiline resistance can emerge during treatment. It is

further concerning that resistance to the new class of nitroimidazole drugs, such as pretomanid and

delamanid, is also conferred by loss of function mutations in any of at least six genes, suggesting that

they may also have a low barrier to resistance[58].

While we identified many non-synonymous variants in *mmpR5*, only one (Ile67fs) has been

previously statistically linked to resistance. We acknowledge that several of our detected variants

have no associated MIC values available in the literature and are thus currently not phenotypically

validated. Determining the phenotypic consequences of *mmpR5* variants that have previously been

described is challenging as there are often only limited reports correlating MICs to genotypes.

Moreover, at least four different methods are used to determine MICs, some of which do not have

associated critical concentrations. Even where critical concentrations have been set, there is an

overlap in MICs of isolates that are genetically wild type and those that have mutations likely to cause

resistance[33]. Furthermore, as we purposefully enriched our dataset for *mmpR5* mutations, the sampling

precludes estimation of the overall prevalence of these mutations in genome sequencing databases.

Prediction of phenotypic bedaquiline resistance from genomic data is further complicated by the

existence of hypersusceptibility variants. For example, the c-11a variant located in the promoter of

MmpR5, which appears to increase susceptibly to bedaquiline[37], was observed to be fixed throughout

a large clade within L2. The early emergence of this variant and its geographical concentration in

South Africa and Eswatini may suggest the role of non-pharmacological influences on MmpR5 which

regulates multiple MmpL efflux systems[20]. Further, analysis of hypersusceptibility is limited by the

truncated lower MIC range of the UKMYC microtitre plates, with many isolates giving MICs below

the lower end of the measured range. While large-scale genotype/phenotype analyses will likely

support the development of rapid molecular diagnostics, targeted or whole genome sequencing, at

reasonable depths, may provide the only opportunity to detect all possible *mmpR5* RAVs, and

possible co-occurring mutations, in clinical settings.

Bedaquiline resistance can also be conferred by other RAVs including in *pepQ* (bedaquiline and

clofazimine), *atpE* (bedaquiline only)[50] and *Rv1979c* (clofazimine only). We only found *atpE* RAVs

at low allele frequency in one patient who also had *mmpR5* variants (sample accession ERR2677436),

which is in line with other evidence suggesting they rarely occur in clinical isolates, likely due to a

high fitness cost. Likewise, we only identified *Rv1979c* RAVs in three patients in China, although

there were other variants in *Rv1979c* for which ability to cause phenotypic resistance has not been

previously assessed. Frameshift *pepQ* mutations that are potentially causative of resistance were

identified in 11 cases, in keeping with its possible role as an additional rare resistance mechanism.

Our findings, of pre-existing reservoirs of *mmpR5* RAVs, are of high clinical relevance as the

presence of *mmpR5* variants during therapy in clinical strains has been associated with substantially

worse outcomes in patients treated with drug regimens including bedaquiline[35]. Although it is

uncertain what the impact of *mmpR5* RAVs are on outcomes when present prior to treatment[59,60], it is

imperative to monitor and prevent the wider transmission of bedaquiline resistant clones, particularly

in high MDR/XDR-TB settings. Early evaluation of new TB drug candidates entering clinical trials

will also be imperative given early data suggesting possible cross-resistance for DprE1 inhibitors such

as macozinone[22]. The large and disparate set of mutations in *mmpR5* we identified, with differing

phenotypes and some being already in circulation before the pre-antibiotic era, adds further urgency

to the development of rapid drug susceptibility testing for bedaquiline to inform effective treatment

choices and mitigate the further spread of DR-TB.

Materials and methods

Sample collection

In this study we curated large representative datasets of *Mtb* whole genome sequences encompassing the global genetic and geographic distribution of lineages 2 (L2) and L4 (**Figure 1, Supplementary Tables S1-S2**). The dataset was enriched to include all available sequenced isolates with *mmpR5* variants, which in some cases included isolates with no, or limited, published metadata. In all other cases samples for which metadata on the geographic location and date of collection was available were retained. To ensure high quality consensus alignments we required that all samples mapped with a minimum percentage cover of 96% and a mean coverage of 30x to the H37Rv reference genome (NC_000962.3). We excluded any samples with evidence of mixed strain infection as identified by the presence of lineage-specific SNPs to more than one sublineage[61] or the presence of a high proportion of heterozygous alleles[62]. The total number of samples included in these datasets, and their source is shown in **Supplementary Table S2.** An index of all samples is available in **Supplementary Table S1**.

A large global dataset of 1,669 L4 *Mtb* sequences has been constructed, which we used as the basis for curating our L4 dataset[43]. We refer to this as the 'base dataset' for L4. For L2, we constructed a 'base dataset' by screening the Sequence Read Archive (SRA) and European Nucleotide Archive (ENA) using BIGSI[63] for the *rpsA* gene sequence containing the L2 defining variant *rpsA* a636c[61] with a 100% match. This search returned 6,307 *Mtb* genomes, of which 1,272 represented unique samples that had the minimum required metadata. Metadata from three studies were also added manually as they were not included in their respective SRA submissions but were available within published studies[64-66].

For isolates with only information on the year of sample collection, we set the date to be equal to the middle of the year. For those with information on the month but not the date of collection we set the

date of collection to the first of the month. For sequenced samples which were missing associated

metadata (32 L2 genomes and 19 L4 genomes) we attempted to estimate an average time of sample

collection in order to impute a sampling date. To do so we computed the average time between date of

collection and sequence upload date for all samples with associated dates separately in each of the L2

and L4 datasets (**Supplementary Figure S1**). For L2 we estimated a mean lag time of 4.7 years (0.5–

12.6 years 95% CI). For L4, having excluded three sequences obtained from 18[th] Century mummies

from Hungary[39], we estimated a mean lag time of 6.9 years (0.6-19.1 years 95% CI). The estimated

dates, where required, are provided in Supplementary Table S1.

To enrich the datasets for isolates with *mmpR5* variants, we included further sequences from our own

studies in KwaZulu-Natal, South Africa[40,41], other studies of drug-resistant TB in southern

Africa[16,43,67-70], and Peru[71,72]. We additionally supplement the Peruvian data with 163 previously

unpublished isolates. In these cases, and to facilitate the most accurate possible estimation of the date

of resistance emergence, we included samples with *mmpR5* variants as well as genetically related

sequences without *mmpR5* variants.

To identify further published raw sequencing data with *mmpR5* variants from studies where

bedaquiline/clofazimine resistance may have been previously unidentified, we screened the NCBI

Sequencing Read Archive (SRA) for sequence data containing 85 previously published *mmpR5*

variants[28-30,40,41,73,74] with BIGSI[63]. BIGSI was employed against a publicly available indexed database

of complete SRA/ENA bacterial and viral whole genome sequences current to December 2016

(available here: http://ftp.ebi.ac.uk/pub/software/bigsi/nat_biotech_2018/all-microbial-index-v03/),

and also employed locally against an updated in-house database which additionally indexed SRA

samples from January 2017 until January 2019. Samples added using this approach are flagged

'BIGSI' in **Supplementary Table S1**. We also used the PYGSI tool[75] to interrogate BIGSI with the

mmpR5 sequence adjusted to include every possible single nucleotide substitution. In each instance

we included 30 bases upstream and downstream of the gene as annotated on the H37Rv *Mtb* reference

genome. For the purpose of this study we only considered coding region, non-synonymous

substitutions and insertions and deletions. Samples added following the PYGSI screen are flagged

'PYGSI' in **Supplementary Table S1**. In the final L2 dataset 194/1514 (12.8%) samples had *mmpR5*

variants, and for L*4* this proportion was 244/2168 (11.3%). A breakdown of the different datasets

used is provided in **Supplementary Table S2**.

Reference mapping and variant calling

Original fastq files for all included sequences were downloaded and paired reads mapped to the

H37Rv reference genome with bwa mem v0.7.17[76]. Mapped reads were sorted and de-duplicated

using Picard Tools v2.20 followed by indel realignment with GATK v3.8[77]. Alignment quality and

coverage was recorded with Qualimap v2.21[78]. Variant calling was performed using bcftools v1.9,

based on reads mapping with a minimum mapping quality of 20, base quality of 20, no evidence of

strand or position bias, a minimum coverage depth of 10 reads, and a minimum of four reads

supporting the alternate allele, with at least two of them on each strand. Moreover, SNPs that were

less than 2bp away from an indel were excluded from the analysis. Similarly, only indels 3bp apart of

other indels were kept.

All sites with insufficient coverage to identify a site as variant or reference were excluded (marked as

'N'), as were those in or within 100 bases of PE/PPE genes, or in insertion sequences or phages. SNPs

present in the alignment with at least 90% frequency were used to generate a pseudoalignment of

equal length to the H37Rv. Samples with more than 10% of the alignment represented by ambiguous

bases were excluded. Those positions with more than 10% of ambiguous bases across all the samples

were also removed. In order to avoid bias on the tree structure, positions known to be associated with

drug resistance were not included.

A more permissive variant calling pipeline was used to identify *mmpR5* variants, as they are often

present at <100% frequency with a high incidence of frameshift mutations. Here we instead employed

FreeBayes v1.2[85] to call all variants present in the *mmpR5* gene (or up to 100 bases upstream) that

were present at ≥5% frequency (alternate allele fraction -*F* 0.05) and supported by at least four reads

including one on each strand. Using this more permissive variant calling strategy we also

systematically screened for all mutations in the efflux pump proteins mmpS5-mmpL5 operon

(**Supplementary Figures S13 and S14**).

Classification of resistance variants

All raw fastq files were screened using the rapid resistance profiling tool TBProfiler[46,79] against a

curated whole genome drug resistance mutations library. This allowed rapid assignment of

polymorphisms associated with resistance to different antimycobacterial drugs and categorisation of

MDR and XDR *Mtb* status (**Supplementary Figure S2, Supplementary Figures S5-S9**).

Classification of *mmpR5* variants

The diverse range of *mmpR5* variants and paucity of widespread MIC testing means that there are

limited data from which to infer the phenotypic consequences of identified *mmpR5* variants. This was

true aside from a subset of data sampled in Peru for which 30 L*4* isolates from Peru were subjected to

MIC testing using the UKMYC6 plate and a further nine were evaluated for MICs reported by the

Cryptic consortium[51]. The approach we used was to assign whether nonsynonymous variants confer a

normal or raised MIC based on published phenotypic tests for strains carrying that variant. A full list

of the literature reports used for each mutation is provided in **Supplementary Table S4**. We also

introduced an intermediate category to describe isolates with MICs at the critical concentration (e.g.,

0.25µg/mL on Middlebrook 7H11 agar), where there is an overlap of the MIC distributions of

mmpR58 mutated and wild type isolates with uncertain clinical implications[33]. We assumed that all

other disruptive frameshift and stop mutations would confer resistance in light of the role of *mmpR5*

as a negative repressor, where loss of function should lead to efflux pump overexpression. All other

promoter and previously unreported missense mutations were categorised as unknown

(**Supplementary Table S4**). However, where *mmpR5* mutations were accompanied by an *mmpS5* or

mmpL5 loss of function mutation, we assumed that would confer susceptibility (or

hypersusceptibility) to bedaquiline[23].

Global phylogenetic inference

The alignments for phylogenetic inference were masked for the *mmpR5* region using bedtools

v2.25.0. All variant positions were extracted from the resulting global phylogenetic alignments using

snp-sites v2.4.1[80], including a L4 outgroup for the L2 alignment (NC_000962.3) and a lineage 3 (L3)

outgroup for the L4 alignment (SRR1188186). This resulted in a 67,585 SNP alignment for the L4

dataset and 29,205 SNP alignment for the L2 dataset. A maximum likelihood phylogenetic tree was

constructed for both SNP alignments using RAxML-NG v0.9.0[81] specifying a GTR+G substitution

model, correcting for the number of invariant sites using the ascertainment flag (ASC_STAM) and

specifying a minimum branch length of $1x10^{-9}$ reporting 12 decimal places (--precision 12).

Estimating the age of emergence of *mmpR5* variants

To test whether the resulting phylogenies can be time-calibrated we first dropped the outgroups from

the phylogeny and rescaled the trees so that branches were measured in unit of substitutions per

genome. We then computed a linear regression between root-to-tip distance and the time of sample

collection using BactDating[82], which additionally assesses the significance of the regression based on

10,000 date randomisations. We obtained a significant temporal correlation for both the L2 and L4

phylogenies, both with and without imputation of dates for samples with missing metadata

(**Supplementary Figure 3**).

We employed the Bayesian method BactDating v1.01[82], run without updating the root

(updateRoot=F), a mixed relaxed gamma clock model and otherwise default parameters to both global

datasets. The MCMC chain was run for $1x10^7$ iterations and $3x10^7$ iterations. BactDating results were

considered only when MCMC chains converged with an Effective Sample Space (ESS) of at least

100. The analysis was applied to the datasets both with and without considering imputed and non-

imputed collection dates (**Supplementary Table 3**).

To independently infer the evolutionary rates associated with each of our datasets, we sub-sampled

both the L4 and L2 datasets to 200 isolates, selected so as to retain the maximal diversity of the tree

using Treemmer v0.3[83]. As before, we excluded all variants currently implicated in drug resistance

from the alignments. This resulted in a dataset for L4 comprising 25,104 SNPs and spanning 232

years of sampling and for L2 comprising 8,221 SNPs and spanning 24 years of sampling. In both

cases the L3 sample SRR1188186 was used as an out-group given this has an associated collection

date. Maximum likelihood trees were constructed using RaXML-NG v0.9.0[81], as previously

described, and a significant temporal regression was obtained for both sub-sampled datasets

(**Supplementary Figure S4**).

BEAST2 v2.6.0[47] was run on both subsampled SNP alignments allowing for model averaging over

possible choices of substitution models[84]. All models were run with either a relaxed or a strict prior on

the evolutionary clock rate for three possible coalescent demographic models: exponential, constant

and skyline. To speed up the convergence, the prior on the evolutionary clock rate was given as a

uniform distribution (limits 0 to 10) with a starting value set to 10^{-7}. In each case, the MCMC chain

was run for 500,000,000 iterations, with the first 10% discarded as burn-in and sampling trees every

10,000 chains. The convergence of the chain was inspected in Tracer 1.7 and through consideration of

the ESS for all parameters (ESS>200). The best-fit model to the data for these runs was assessed

through a path sampling analysis[85] specifying 100 steps, 4 million generations per step, alpha = 0.3,

pre-burn-in = 1 million generations, burn-in for each step = 40%. For both datasets, the best supported

strict clock model was a coalescent Bayesian skyline analysis. The rates (mean and 95% HPD)

estimated under these subsampled analyses (L2 7.7×10^{-8} [4.9×10^{-8} - 1.03×10^{-7}] substitutions per site

per year; L4 7.1×10^{-8} [6.2×10^{-8} - 7.9×10^{-8}] substitutions per site per year) were used to rescale the

maximum likelihood phylogenetic trees generated across the entire L2 and L4 datasets, by

transforming all branch lengths of the tree from per unit substitution to per unit substitutions per site

per year using the R package Ape v5.3[86]. This resulted in an estimated tMRCA of 1332CE (945CE-

1503CE) for L2 and 853CE (685CE – 967CE) for L4 (**Figure 2**).

The resulting phylogenetic trees were visualised and annotated for place of geographic sampling and

mmpR5 variant status using ggtree v1.14.6[87]. All nonsynonymous mutations in *mmpR5* were

considered, with the phenotypic status assigned in **Supplementary Table S4**. For the purpose of this

analysis, and to be conservative, 'unknown' variants classified using XGBoost were still considered

'unknown' (**Supplementary Note 1**). Clades carrying shared variants in *mmpR5* were identified and

the distributions around the age of the node (point estimates – mean - and 95% HPDs) were extracted

from the time-stamped phylogeny. For isolated samples (single emergences) exhibiting variants in

mmpR5, the time of sample collection was extracted together with the date associated with the upper

bound on the age of the next closest node of the tree, allowing for the mutation to have occurred

anywhere over the length of the terminal branch (**Figure 3, Supplementary Figures S11-S12**). For

the Peruvian clade Bayesian skyline analysis was implemented through the skylineplot analysis

functionality available in Ape v5.3[86].

Data availability

Raw sequence data and full metadata for all newly generated isolates are available on NCBI

Sequencing Read Archive under BioProject ID: PRJEB39837.

Footnotes

Author Contributions

LvD, CN and FB conceived and designed the study. JM, NP, AG, MO, AP, OBB, VE and LG

provided sequence data. ATO, JP, MA, CCST and XD performed and advised on computational

analyses. LvD, CN and FB wrote the manuscript with input from all co-authors. All authors read and

approved the final manuscript.

Acknowledgments

CN and JM are supported by the Wellcome Trust (203583/Z/16/Z and 203919/Z/16/Z, respectively).

LvD is supported by a UCL Excellence Fellowship. F.B. acknowledges support from the BBSRC

(equipment grant BB/R01356X/1). FB additionally acknowledges the National Institute for Health

Research University College London Hospitals Biomedical Research Centre. M.A. was supported by

a Ph.D. scholarship from University College London. All authors acknowledge UCL Biosciences Big

Data equipment grant from BBSRC (BB/R01356X/1).

Competing interests

The authors declare no competing financial interests. AP is currently employed by Janssen. Dr Pym's

involvement with the research described herein precedes his employment at Janssen.

Figures

571
572
Figure 1: Compiled global *Mtb* genomic datasets.

Panels a) and b) provide the geographic location of isolates included in the lineage 2 and lineage 4

datasets respectively. Pies are scaled by the number of samples per country (raw data available in

Supplementary Table S1) with the colours providing the fraction of genomes with any

nonsynonymous variants detected in *mmpR5* (coloured as per the legend). Countries comprising

samples with known RAVs are highlighted with a red asterisk. Genomic data for which no associated

metadata on the geographic location of sampling was available are shown in the Pacific Ocean. Panels

c) and d) provide the collection dates associated to each genome in the lineage 2 and lineage 4

datasets respectively highlighting those with any variants in *mmpR5* (colour, as per legend). Lineage 4

Mtb obtained from 18[th] century mummies are excluded from this plot but included in all analyses. The

vertical dashed lines indicate the dates of the first clofazimine clinical trials and bedaquiline clinical

trials and FDA approval.

Figure 2: Global time calibrated *Mtb* phylogenies.

Inferred dated phylogenies (x-axis) for the a) lineage 2 and b) lineage 4 datasets. Tips are coloured by

the geographic region of sampling as given in the legend. The bar provides the assessed phenotype

(colour) based on assignment of nonsynonymous variants in *mmpR5*.

Figure 3: Estimated age of emergence of *mmpR5* nonsynonymous variants.

Inferred point estimates for the age of emergence of clades with *mmpR5* variants for the lineage 2 (a)

and lineage 4 (b) datasets, including a zoomed in reproduction of the period from 2007-2020. Y-axis

provides the absolute number of sequences descending from the identified and dated nodes. The

mmpR5 RAV status is given by the colour as defined in the legend at bottom. *indicates phenotypic

data available that are supportive of MIC classification (see text). The full mutation timelines are

provided in **Supplementary Figures 11-12** and **Supplementary Table S7**.

References

1.   Organization WH. *Global Tuberculosis Report 2022*. 2022. *Global Tuberculosis Reports*.   https://www.who.int/teams/global-tuberculosis-programme/tb-reports/global-tuberculosis-report-2022

2.   Cegielski JP, Kurbatova E, van der Walt M, et al. Multidrug-Resistant Tuberculosis Treatment Outcomes in Relation to Treatment and Initial Versus Acquired Second-Line Drug Resistance. *Clin Infect Dis*. Feb 15 2016;62(4):418-430. doi:10.1093/cid/civ910

3.   Organization WH. *Global Tuberculosis Report 2019*. 2019. *Global Tuberculosis Reports*. https://www.who.int/publications/i/item/9789241565714

4.   Andries K, Verhasselt P, Guillemont J, et al. A diarylquinoline drug active on the ATP synthase of Mycobacterium tuberculosis. *Science*. Jan 14 2005;307(5707):223-7. doi:10.1126/science.1106753

5.   Diacon AH, Pym A, Grobusch MP, et al. Multidrug-resistant tuberculosis and culture conversion with bedaquiline. *N Engl J Med*. Aug 21 2014;371(8):723-32. doi:10.1056/NEJMoa1313865

6.   Spring S, inventor; Sirturo (bedaquiline).

7.   Borisov SE, Dheda K, Enwerem M, et al. Effectiveness and safety of bedaquiline-containing regimens in the treatment of MDR- and XDR-TB: a multicentre study. *Eur Respir J*. May 2017;49(5)doi:10.1183/13993003.00387-2017

8.   Guglielmetti L, Jaspard M, Le Du D, et al. Long-term outcome and safety of prolonged bedaquiline treatment for multidrug-resistant tuberculosis. *Eur Respir J*. Mar 2017;49(3)doi:10.1183/13993003.01799-2016

9.   Olayanju O, Limberis J, Esmail A, et al. Long-term bedaquiline-related treatment outcomes in patients with extensively drug-resistant tuberculosis from South Africa. *Eur Respir J*. May 2018;51(5)doi:10.1183/13993003.00544-2018

10.   Ndjeka N, Schnippel K, Master I, et al. High treatment success rate for multidrug-resistant and extensively drug-resistant tuberculosis using a bedaquiline-containing treatment regimen. *Eur Respir J*. Dec 2018;52(6)doi:10.1183/13993003.01528-2018

11.   Organization WH. *Module 4: treatment - drug-resistant tuberculosis treatment, 2022 update*. WHO consolidated guidelines on tuberculosis. 2022.

12.   Conradie F, Bagdasaryan TR, Borisov S, et al. Bedaquiline-Pretomanid-Linezolid Regimens for Drug-Resistant Tuberculosis. *N Engl J Med*. Sep 1 2022;387(9):810-823. doi:10.1056/NEJMoa2119430

13.   Berry C, du Cros P, Fielding K, et al. TB-PRACTECAL: study protocol for a randomised, controlled, open-label, phase II-III trial to evaluate the safety and efficacy of regimens containing bedaquiline and pretomanid for the treatment of adult patients with pulmonary multidrug-resistant tuberculosis. *Trials*. Jun 13 2022;23(1):484. doi:10.1186/s13063-022-06331-8

14.   Paton NI, Cousins C, Suresh C. Treatment Strategy for Rifampin-Susceptible Tuberculosis. Reply. *N Engl J Med*. Jun 15 2023;388(24):2298. doi:10.1056/NEJMc2304776

15.   Manson AL, Cohen KA, Abeel T, et al. Genomic analysis of globally diverse Mycobacterium tuberculosis strains provides insights into the emergence and spread of multidrug resistance. *Nat Genet*. Mar 2017;49(3):395-402. doi:10.1038/ng.3767

16.   Cohen KA, Abeel T, Manson McGuire A, et al. Evolution of Extensively Drug-Resistant Tuberculosis over Four Decades: Whole Genome Sequencing and Dating Analysis of Mycobacterium tuberculosis Isolates from KwaZulu-Natal. *PLoS Med*. Sep 2015;12(9):e1001880. doi:10.1371/journal.pmed.1001880

17.   Eldholm V, Balloux F. Antimicrobial Resistance in Mycobacterium tuberculosis: The Odd One Out. *Trends Microbiol*. Aug 2016;24(8):637-648. doi:10.1016/j.tim.2016.03.007

18.   Huitric E, Verhasselt P, Koul A, Andries K, Hoffner S, Andersson DI. Rates and
mechanisms of resistance development in Mycobacterium tuberculosis to a novel
diarylquinoline ATP synthase inhibitor. *Antimicrob Agents Chemother*. Mar
2010;54(3):1022-8. doi:10.1128/AAC.01611-09
19.   Almeida D, Ioerger T, Tyagi S, et al. Mutations in pepQ Confer Low-Level
Resistance to Bedaquiline and Clofazimine in Mycobacterium tuberculosis. *Antimicrob*
Agents Chemother. Aug 2016;60(8):4590-9. doi:10.1128/AAC.00753-16
20.   Andries K, Villellas C, Coeck N, et al. Acquired resistance of Mycobacterium
tuberculosis to bedaquiline. *PLoS One*. 2014;9(7):e102135.
doi:10.1371/journal.pone.0102135
21.   Hartkoorn RC, Uplekar S, Cole ST. Cross-resistance between clofazimine and
bedaquiline through upregulation of MmpL5 in Mycobacterium tuberculosis. *Antimicrob*
Agents Chemother. May 2014;58(5):2979-81. doi:10.1128/AAC.00037-14
22.   Poulton NC, Azadian ZA, DeJesus MA, Rock JM. Mutations in rv0678 Confer Low-
Level Resistance to Benzothiazinone DprE1 Inhibitors in Mycobacterium tuberculosis.
Antimicrob Agents Chemother. Sep 20 2022;66(9):e0090422. doi:10.1128/aac.00904-22
23.   Vargas R, Jr., Freschi L, Spitaleri A, et al. Role of Epistasis in Amikacin, Kanamycin,
Bedaquiline, and Clofazimine Resistance in Mycobacterium tuberculosis Complex.
Antimicrob Agents Chemother. Oct 18 2021;65(11):e0116421. doi:10.1128/AAC.01164-21
24.   Bloemberg GV, Keller PM, Stucki D, et al. Acquired Resistance to Bedaquiline and
Delamanid in Therapy for Tuberculosis. *N Engl J Med*. Nov 12 2015;373(20):1986-8.
doi:10.1056/NEJMc1505196
25.   Xu J, Wang B, Hu M, et al. Primary Clofazimine and Bedaquiline Resistance among
Isolates from Patients with Multidrug-Resistant Tuberculosis. *Antimicrob Agents Chemother*.
Jun 2017;61(6)doi:10.1128/AAC.00239-17
26.   Zimenkov DV, Nosova EY, Kulagina EV, et al. Examination of bedaquiline- and
linezolid-resistant Mycobacterium tuberculosis isolates from the Moscow region. *J*
Antimicrob Chemother. Jul 1 2017;72(7):1901-1906. doi:10.1093/jac/dkx094
27.   de Vos M, Ley SD, Wiggins KB, et al. Bedaquiline Microheteroresistance after
Cessation of Tuberculosis Treatment. *N Engl J Med*. May 30 2019;380(22):2178-2180.
doi:10.1056/NEJMc1815121
28.   Ghodousi A, Rizvi AH, Baloch AQ, et al. Acquisition of Cross-Resistance to
Bedaquiline and Clofazimine following Treatment for Tuberculosis in Pakistan. *Antimicrob*
Agents Chemother. Sep 2019;63(9)doi:10.1128/AAC.00915-19
29.   Polsfuss S, Hofmann-Thiel S, Merker M, et al. Emergence of Low-level Delamanid
and Bedaquiline Resistance During Extremely Drug-resistant Tuberculosis Treatment. *Clin*
Infect Dis. Sep 13 2019;69(7):1229-1231. doi:10.1093/cid/ciz074
30.   Mokrousov I, Akhmedova G, Polev D, Molchanov V, Vyazovaya A. Acquisition of
bedaquiline resistance by extensively drug-resistant Mycobacterium tuberculosis strain of
Central Asian Outbreak clade. *Clin Microbiol Infect*. Oct 2019;25(10):1295-1297.
doi:10.1016/j.cmi.2019.06.014
31.   Kadura S, King N, Nakhoul M, et al. Systematic review of mutations associated with
resistance to the new and repurposed Mycobacterium tuberculosis drugs bedaquiline,
clofazimine, linezolid, delamanid and pretomanid. *J Antimicrob Chemother*. Aug 1
2020;75(8):2031-2043. doi:10.1093/jac/dkaa136
32.   Roberts LW, Malone KM, Hunt M, et al. Repeated evolution of bedaquiline
resistance in Mycobacterium tuberculosis is driven by truncation of mmpR5. *bioRxiv*.
2022:2022.12.08.519610. doi:10.1101/2022.12.08.519610
33.   Organization WH. *Technical report on critical concentrations for TB drug*
susceptibility testing of medicines used in the treatment of drug-resistant TB. 2018.

34.   Ismail N, Riviere E, Limberis J, et al. Genetic variants and their association with
phenotypic resistance to bedaquiline in Mycobacterium tuberculosis: a systematic review and
individual isolate data analysis. *Lancet Microbe*. Nov 2021;2(11):E604-E616.
doi:10.1016/S2666-5247(21)00175-0
35.   Nimmo C, Millard J, Brien K, et al. Bedaquiline resistance in drug-resistant
tuberculosis HIV co-infected patients. *Eur Respir J*. Jun
2020;55(6)doi:10.1183/13993003.02383-2019
36.   Martinez E, Hennessy D, Jelfs P, Crighton T, Chen SC, Sintchenko V. Mutations
associated with in vitro resistance to bedaquiline in Mycobacterium tuberculosis isolates in
Australia. *Tuberculosis (Edinb)*. Jul 2018;111:31-34. doi:10.1016/j.tube.2018.04.007
37.   Villellas C, Coeck N, Meehan CJ, et al. Unexpected high prevalence of resistance-
associated Rv0678 variants in MDR-TB patients without documented prior use of
clofazimine or bedaquiline. *J Antimicrob Chemother*. Mar 1 2017;72(3):684-690.
doi:10.1093/jac/dkw502
38.   Merker M, Kohl TA, Barilar I, et al. Phylogenetically informative mutations in genes
implicated in antibiotic resistance in Mycobacterium tuberculosis complex. *Genome Med*.
Mar 6 2020;12(1):27. doi:10.1186/s13073-020-00726-5
39.   Kay GL, Sergeant MJ, Zhou Z, et al. Eighteenth-century genomes show that mixed
infections were common at time of peak tuberculosis in Europe. *Nat Commun*. Apr 7
2015;6:6717. doi:10.1038/ncomms7717
40.   Nimmo C, Millard J, van Dorp L, et al. Population-level emergence of bedaquiline
and clofazimine resistance-associated variants among patients with drug-resistant
tuberculosis in southern Africa: a phenotypic and phylogenetic analysis. *Lancet Microbe*.
Aug 2020;1(4):e165-e174. doi:10.1016/S2666-5247(20)30031-8
41.   Nimmo C, Brien K, Millard J, et al. Dynamics of within-host Mycobacterium
tuberculosis diversity and heteroresistance during treatment. *EBioMedicine*. May
2020;55:102747. doi:10.1016/j.ebiom.2020.102747
42.   O'Neill MB, Shockey A, Zarley A, et al. Lineage specific histories of Mycobacterium
tuberculosis dispersal in Africa and Eurasia. *Mol Ecol*. Jul 2019;28(13):3241-3256.
doi:10.1111/mec.15120
43.   Brynildsrud OB, Pepperell CS, Suffys P, et al. Global expansion of Mycobacterium
tuberculosis lineage 4 shaped by colonial migration and local adaptation. *Sci Adv*. Oct
2018;4(10):eaat5869. doi:10.1126/sciadv.aat5869
44.   Rutaihwa LK, Menardo F, Stucki D, et al. Multiple Introductions of Mycobacterium
tuberculosis Lineage 2-Beijing Into Africa Over Centuries. *Front Ecol Evol*. Apr 16
2019;7doi:ARTN 112
10.3389/fevo.2019.00112
45.   Bradley P, Gordon NC, Walker TM, et al. Rapid antibiotic-resistance predictions
from genome sequence data for Staphylococcus aureus and Mycobacterium tuberculosis. *Nat
Commun*. Dec 21 2015;6:10063. doi:10.1038/ncomms10063
46.   Phelan JE, O'Sullivan DM, Machado D, et al. Integrating informatics tools and
portable sequencing technology for rapid detection of resistance to anti-tuberculous drugs.
Genome Med. Jun 24 2019;11(1):41. doi:10.1186/s13073-019-0650-x
47.   Bouckaert R, Vaughan TG, Barido-Sottani J, et al. BEAST 2.5: An advanced software
platform for Bayesian evolutionary analysis. *PLoS Comput Biol*. Apr 2019;15(4):e1006650.
doi:10.1371/journal.pcbi.1006650
48.   Menardo F, Duchene S, Brites D, Gagneux S. The molecular clock of Mycobacterium
tuberculosis. *PLoS Pathog*. Sep 2019;15(9):e1008067. doi:10.1371/journal.ppat.1008067

49. Ismail N, Peters RPH, Ismail NA, Omar SV. Clofazimine Exposure In Vitro Selects
Efflux Pump Mutants and Bedaquiline Resistance. *Antimicrob Agents Chemother*. Mar
2019;63(3)doi:10.1128/AAC.02141-18
50. Andres S, Merker M, Heyckendorf J, et al. Bedaquiline-Resistant Tuberculosis: Dark
Clouds on the Horizon. *Am J Respir Crit Care Med*. Jun 15 2020;201(12):1564-1568.
doi:10.1164/rccm.201909-1819LE
51. The CC. A data compendium associating the genomes of 12,289 Mycobacterium
tuberculosis isolates with quantitative resistance phenotypes to 13 antibiotics. *PLOS Biology*.
2022;20(8):e3001721. doi:10.1371/journal.pbio.3001721
52. The CC. Epidemiological cutoff values for a 96-well broth microdilution plate for
high-throughput research antibiotic susceptibility testing of M.
tuberculosis. *European Respiratory Journal*. 2022:2200239.
doi:10.1183/13993003.00239-2022
53. Beckert P, Sanchez-Padilla E, Merker M, et al. MDR M. tuberculosis outbreak clone
in Eswatini missed by Xpert has elevated bedaquiline resistance dated to the pre-treatment
era. *Genome Med*. Nov 25 2020;12(1):104. doi:10.1186/s13073-020-00793-8
54. Hariguchi N, Chen X, Hayashi Y, et al. OPC-167832, a Novel Carbostyril Derivative
with Potent Antituberculosis Activity as a DprE1 Inhibitor. *Antimicrob Agents Chemother*.
May 21 2020;64(6)doi:10.1128/AAC.02020-19
55. Loiseau C, Menardo F, Aseffa A, et al. An African origin for Mycobacterium bovis.
Evol Med Public Health. 2020;2020(1):49-59. doi:10.1093/emph/eoaa005
56. Bateson A, Ortiz Canseco J, McHugh TD, et al. Ancient and recent differences in the
intrinsic susceptibility of Mycobacterium tuberculosis complex to pretomanid. *J Antimicrob
Chemother*. May 29 2022;77(6):1685-1693. doi:10.1093/jac/dkac070
57. D'Costa VM, King CE, Kalan L, et al. Antibiotic resistance is ancient. *Nature*. Sep 22
2011;477(7365):457-461. doi:10.1038/nature10388
58. Rifat D, Li SY, Ioerger T, et al. Mutations in fbiD (Rv2983) as a Novel Determinant
of Resistance to Pretomanid and Delamanid in Mycobacterium tuberculosis. *Antimicrob
Agents Ch*. Jan 2021;65(1)doi:ARTN e01948-20
10.1128/AAC.01948-20
59. Liu Y, Gao M, Du J, et al. Reduced Susceptibility of Mycobacterium tuberculosis to
Bedaquiline During Antituberculosis Treatment and Its Correlation With Clinical Outcomes
in China. *Clin Infect Dis*. Nov 2 2021;73(9):e3391-e3397. doi:10.1093/cid/ciaa1002
60. Pym AS, Diacon AH, Tang SJ, et al. Bedaquiline in the treatment of multidrug- and
extensively drug-resistant tuberculosis. *Eur Respir J*. Feb 2016;47(2):564-74.
doi:10.1183/13993003.00724-2015
61. Coll F, McNerney R, Guerra-Assuncao JA, et al. A robust SNP barcode for typing
Mycobacterium tuberculosis complex strains. *Nat Commun*. Sep 1 2014;5:4812.
doi:10.1038/ncomms5812
62. Sobkowiak B, Glynn JR, Houben R, et al. Identifying mixed Mycobacterium
tuberculosis infections from whole genome sequence data. *BMC Genomics*. Aug 14
2018;19(1):613. doi:10.1186/s12864-018-4988-z
63. Bradley P, den Bakker HC, Rocha EPC, McVean G, Iqbal Z. Ultrafast search of all
deposited bacterial and viral genomic data. *Nat Biotechnol*. Feb 2019;37(2):152-159.
doi:10.1038/s41587-018-0010-1
64. Merker M, Blin C, Mona S, et al. Evolutionary history and global spread of the
Mycobacterium tuberculosis Beijing lineage. *Nat Genet*. Mar 2015;47(3):242-9.
doi:10.1038/ng.3195

65.   Luo T, Comas I, Luo D, et al. Southern East Asian origin and coexpansion of
Mycobacterium tuberculosis Beijing family with Han Chinese. *Proc Natl Acad Sci U S A*. Jun
30 2015;112(26):8136-41. doi:10.1073/pnas.1424063112
66.   Norheim G, Seterelv S, Arnesen TM, et al. Tuberculosis Outbreak in an Educational
Institution   in   Norway.   *J   Clin   Microbiol*.   May   2017;55(5):1327-1333.
doi:10.1128/JCM.01152-16
67.   Nimmo C, Shaw LP, Doyle R, et al. Whole genome sequencing Mycobacterium
tuberculosis directly from sputum identifies more genetic diversity than sequencing from
culture. *BMC Genomics*. May 20 2019;20(1):389. doi:10.1186/s12864-019-5782-2
68.   Dheda K, Limberis JD, Pietersen E, et al. Outcomes, infectiousness, and transmission
dynamics of patients with extensively drug-resistant tuberculosis and home-discharged
patients with programmatically incurable tuberculosis: a prospective cohort study. *Lancet
Respir Med*. Apr 2017;5(4):269-281. doi:10.1016/S2213-2600(16)30433-7
69.   Streicher EM, Sampson SL, Dheda K, et al. Molecular Epidemiological Interpretation
of the Epidemic of Extensively Drug-Resistant Tuberculosis in South Africa. *J Clin
Microbiol*. Nov 2015;53(11):3650-3. doi:10.1128/JCM.01414-15
70.   Guerra-Assuncao JA, Crampin AC, Houben RM, et al. Large-scale whole genome
sequencing of M. tuberculosis provides insights into transmission in a high prevalence area.
Elife. Mar 3 2015;4doi:10.7554/eLife.05166
71.   Grandjean L, Gilman RH, Martin L, et al. Transmission of Multidrug-Resistant and
Drug-Susceptible Tuberculosis within Households: A Prospective Cohort Study. *PLoS Med*.
Jun 2015;12(6):e1001843; discussion e1001843. doi:10.1371/journal.pmed.1001843
72.   Grandjean L, Gilman RH, Iwamoto T, et al. Convergent evolution and topologically
disruptive polymorphisms among multidrug-resistant tuberculosis in Peru. *PLoS One*.
2017;12(12):e0189838. doi:10.1371/journal.pone.0189838
73.   Ismail N, Omar SV, Ismail NA, Peters RPH. Collated data of mutation frequencies
and associated genetic variants of bedaquiline, clofazimine and linezolid resistance in
Mycobacterium   tuberculosis.   *Data   Brief*.   Oct   2018;20:1975-1983.
doi:10.1016/j.dib.2018.09.057
74.   Ghajavand H, Kargarpour Kamakoli M, Khanipour S, et al. High Prevalence of
Bedaquiline Resistance in Treatment-Naive Tuberculosis Patients and Verapamil
Effectiveness. *Antimicrob Agents Chemother*. Mar 2019;63(3)doi:10.1128/AAC.02530-18
75.   *pygsi v1.0.0*. 2018. https://zenodo.org/record/1407085
76.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. *arXiv*. 2013;doi:arXiv:1303.3997
77.   Van der Auwera GA, Carneiro MO, Hartl C, et al. From FastQ data to high
confidence variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr Protoc
Bioinformatics*. 2013;43(1110):11 10 1-11 10 33. doi:10.1002/0471250953.bi1110s43
78.   Okonechnikov K, Conesa A, Garcia-Alcalde F. Qualimap 2: advanced multi-sample
quality control for high-throughput sequencing data. *Bioinformatics*. Jan 15 2016;32(2):292-
4. doi:10.1093/bioinformatics/btv566
79.   Coll F, McNerney R, Preston MD, et al. Rapid determination of anti-tuberculosis drug
resistance from whole-genome sequences. *Genome Med*. 2015;7(1):51. doi:10.1186/s13073-
015-0164-0
80.   Page AJ, Taylor B, Delaney AJ, et al. SNP-sites: rapid efficient extraction of SNPs
from   multi-FASTA   alignments.   *Microb   Genom*.   Apr   2016;2(4):e000056.
doi:10.1099/mgen.0.000056
81.   Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: a fast,
scalable and user-friendly tool for maximum likelihood phylogenetic inference.
Bioinformatics. Nov 1 2019;35(21):4453-4455. doi:10.1093/bioinformatics/btz305

82.   Didelot X, Croucher NJ, Bentley SD, Harris SR, Wilson DJ. Bayesian inference of
ancestral dates on bacterial phylogenetic trees. *Nucleic Acids Res*. Dec 14 2018;46(22):e134.
doi:10.1093/nar/gky783
83.   Menardo F, Loiseau C, Brites D, et al. Treemmer: a tool to reduce large phylogenetic
datasets with minimal loss of diversity. *BMC Bioinformatics*. May 2 2018;19(1):164.
doi:10.1186/s12859-018-2164-8
84.   Bouckaert RR, Drummond AJ. bModelTest: Bayesian phylogenetic site model
averaging and model comparison. *BMC Evol Biol*. Feb 6 2017;17(1):42. doi:10.1186/s12862-
017-0890-6
85.   Baele G, Lemey P, Bedford T, Rambaut A, Suchard MA, Alekseyenko AV.
Improving the accuracy of demographic and molecular clock model comparison while
accommodating phylogenetic uncertainty. *Mol Biol Evol*. Sep 2012;29(9):2157-67.
doi:10.1093/molbev/mss084
86.   Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics*. Feb 1 2019;35(3):526-528.
doi:10.1093/bioinformatics/bty633
87.   Yu GC, Smith DK, Zhu HC, Guan Y, Lam TTY. GGTREE: an R package for
visualization and annotation of phylogenetic trees with their covariates and other associated
data. *Methods Ecol Evol*. Jan 2017;8(1):28-36. doi:10.1111/2041-210x.12628
866