Bridging Photosynthesis and Crop Yield Formation with a Mechanistic Model of

Whole Plant Carbon Nitrogen Interaction

Tian-Gen Chang and Xin-Guang Zhu[*]

Key Laboratory for Plant Molecular Genetics, Center of Excellence for Molecular

Plant Sciences, Chinese Academy of Sciences, Shanghai 200032, China

*Correspondence: zhuxg@cemps.ac.cn

ABSTRACT

Crop yield is co-determined by photosynthetic potential of source organs, and pattern
of partitioning and utilization of photosynthate among sink organs. Although
correlation between source sink relation and grain yield has been studied for a
century, a quantitative understanding of the metabolic basis of source sink interaction
is lacking. Here, we describe a mechanistic model of Whole plAnt Carbon Nitrogen
Interaction (WACNI), enabling precise prediction of plant physiological dynamics
during the grain filling period by reconstructing primary metabolic and biophysical
processes in source, sink and transport organs. To get a specified range of parameters
required to quantify the enzymatic kinetics in rice, a data set is established based on
case studies and natural variation surveys in the past decades. The parameterized
model quantitatively predicts plant carbon and nitrogen budget upon various
scenarios, ranging from field management and environmental perturbation to genetic
manipulation, thus enabling dissection of the precise role of such alterations in crop
yield formation. Model simulations further reveal the importance of re-allocating
activity of carbon/nitrogen metabolic and transport processes for a plant physiological
ideotype to maximize crop yield.

Key words: carbon nitrogen metabolism, crop yield, grain filling, ideotype,
photosynthesis, plant physiology, source sink interaction, systems model

INTRODUCTION

As the global climate changes, population increases, arable land decreases, there is a tremendous need for the next green revolution to greatly increase the crop yield once again [1-4]. With the available identification of a large number of molecular markers controlling important traits in plants, molecular breeding by rational design, is becoming a new route for crop improvement [5-9]. In addition to the advances in functional genomics research, other major drivers of this crop improvement route include advances in phenomics technology and genome editing technology [10-13]. To date, one key remaining issue is the quantification of crop ideotype as an objective during crop genetic engineering [14].

Crop ideotype includes both morphological and physiological characteristics, which together determine canopy photosynthesis and whole-plant source sink relation, thus define the crop yield potential [15-18]. Ever since the proposal of the wheat ideotype by Donald [19], improving crop morphology has been extensively studied in genetic research, modeling research, and in crop high-yield breeding, which is largely responsible for the dramatic increase in rice grain yield from around 3 t ha^{-1} to up to 15 t ha^{-1} in the past 60 years [17,20-26]. Although long recognized as the other equally important factor of crop ideotype [27-29], crop physiological parameters, i.e., characteristics associated with photosynthesis, respiration, carbon and nitrogen

metabolism in different organs, and also the transport of metabolites between organs,

have so far generated little impact in crop breeding. This is due to there is still no

consensus on the optimal physiological characteristics associated with a crop

ideotype. In the past century, results from individual field experiments often

contradicted with each other, due to different weather and soil conditions of the

experimental sites, and/or the use of different crops or cultivars or management

practices [30].

Systems modeling approach is recognized as a promising means to guide design of

the optimal crop physiological characteristics to achieve high yield [30-33]. As a result of

research by plant systems modeling community during the past decades, mechanistic

models for a number of individual plant physiological processes, such as leaf

photosynthesis [34,35], phloem long-distance carbon transport transport [36], flowering

time determination [37,38] and root growth and development [39], have been developed.

These mechanistic models have been used to guide improvement of photosynthesis

for greater efficiency [2,40-43] and evaluate the impact of modifying photosynthesis for

crop yields [44]. Missing in the current repertoire of models is a mechanistic framework

model connecting metabolism within individual source/sink organs and transport of

metabolites between these organs, where the consequence on plant growth and

development is predicted as an emergent property.

Carbon and nitrogen metabolism are the basis for plant growth and development [45].

To predict crop yield formation directly from the molecular processes in different

source and sink organs, and the carbon and nitrogen fluxes between organs, we

develop a mechanistic model of **W**hole pl**A**nt **C**arbon **N**itrogen **I**nteraction (WACNI).

As a fully mechanistic model, WACNI not only enables us to quantify plant carbon

and nitrogen budget under different source sink relation alteration scenarios and to

define the physiological features associated with crop ideotype, but also serves to

design molecular manipulations to approach such an ideotype. Thus, a combination of

this theoretical framework and the advanced genome editing technologies [46,47] will

offer unprecedented opportunity in engineering crop physiological features for greater

yield.

RESULTS

Construction of WACNI: a mechanistic model of plant grain yield formation

WACNI is an ordinary differentiation equations (ODEs) based kinetic model

predicting plant-level physiological dynamics from the flowering to the harvest stage.

It comprises five modules, i.e., root, leaf, grain, stem (including culm and sheath) and

vascular transport system (xylem and phloem). Metabolites exchange between

modules by trans-membrane transport (Figure 1). Fourteen types of biochemical and

biophysical processes involved in different source/sink/transport organs are

mathematically represented in WACNI (Figure 1). Briefly, these processes include

assimilation, transport and utilization of six representative primary metabolites, i.e.,

triose phosphates (TP), sucrose (Suc), starch, inorganic nitrogen (I-N, including NH_4^+

and NO_3^-), free form of organic nitrogen (O-N, including amino acids and amides),

and proteins. Further, WACNI also incorporates the interaction between these

metabolites and plant developmental processes, e.g. root growth, grain volume

expansion (endosperm cell division), grain filling (starch and protein cumulation in

endosperm), root senescence and leaf senescence (Figure 1; see Methods). The

apparent kinetic parameters for each process, initial metabolite concentrations and

carbon/nitrogen mass in different organs at the flowering stage are input variables for

the model.

Figure 1. A mechanistic model of plant grain yield formation during rice grain filling period. Abbreviations: Suc, sucrose; TP, triose phosphates; I-N, inorganic nitrogen; O-N, free-form organic nitrogen; TN, total mobile nitrogen, including I-N, O-N and protein; HATS, high-affinity nitrogen transport system; LATS, low-affinity nitrogen transport system; CBB, Calvin–Benson–Bassham cycle. The 14 groups of processes illustrated in the figure are: 1, root nitrogen uptake; 2, root nitrogen assimilation; 3, root growth; 4, root senescence; 5, leaf CO_2 and nitrogen assimilation; 6, leaf triose phosphate, sucrose and starch interconversion; 7, leaf organic nitrogen and protein interconversion; 8, leaf senescence; 9, grain volume growth (cell division); 10, grain starch and protein synthesis; 11, stem sucrose and starch, organic nitrogen and protein interconversion; 12, phloem long distance transport; 13, transporter-dependent short distance transport; 14, symplastic diffusion between phloem and stem.

 Parameterization of WACNI: formation of a primary reference dataset on natural

variation of important rice traits based on bibliomics

Firstly, we have simplified the representation of each process to reduce the required

number of parameters to be estimated (see Methods). Then, for each required

parameter, its value or value range was determined based on the available literature in

rice. Specifically, for most morphological and agronomic parameters, their natural

variations were determined from independent case studies, large-scale surveys on

genetic populations or from the reviews. For determining a certain parameter

controlling capacity of a certain process, maximal activity of the rate limiting enzyme

in this process was used in the first priority; if the maximal activity was not available,

the maximal metabolic flux of this process would be used (see details in

Supplementary Table 1). Finally, a comprehensive dataset for natural variation of

many important parameters determining rice growth and development was generated

(Supplementary Table 1).

Physiological dynamics and carbon & nitrogen economy during rice grain filling

period

After parameterization, WACNI realistically predicted, during the grain filling period,

changes in root weight (Figure 2a, R^2=0.94), leaf area (Figure 2b, R^2=0.96), rates of

canopy photosynthesis (Figure 2c, R^2=0.96), grain volume (Figure 2d, R^2=0.99), grain

dry weight (Figure 2e, R^2=1.00) and stem non-structural carbohydrates (Figure 2f,

R^2=0.99). In addition, a plant-level carbon and nitrogen budget throughout the whole

grain filling period was generated (Figure 2g). For the carbon budget, WACNI

predicted that 21.6% and 52.1% of the photosynthetically fixed carbon could be

respired by the leaf itself and by the entire plant, respectively, which were close to the

previous estimates of 20% and 40-60% in cultivated plants, respectively [48]. WACNI

predicted an 88.0% and 12.0% contribution from photosynthesis-derived carbon and

pre-flowering non-structural carbohydrate storage in stem to grain filling,

respectively. The 'apparent contribution' of pre-flowering stem non-structural

carbohydrate storage to grain yield, i.e., the ratio between the stem weight loss during

grain filling to the grain yield at the harvest stage, was predicted to be 27.3%, which

coincides with the observed value in the range of 0 to 40% [49]. Namely, our simulation

demonstrates that the frequently used 'apparent contribution', which omits respiration

of the grain and the stem, is a substantial over-estimate of the actual contribution of

pre-flowering stem non-structural carbohydrate storage to grain yield (e.g. 27.3%

versus 12.0% in our case). In addition, 19.8% of the total carbohydrates was predicted

to be partitioned to the root from the flowering stage to the harvest stage, while the

remaining 80.2% to the grain.

Figure 2. Prediction of plant physiological dynamics (A-F) and carbon & nitrogen
economy (G) during rice grain filling period. **a**, root dry weight (R2=0.94; data from
[50]); **b**, leaf area (R2=0.96; data from [51]); **c**, diurnal canopy photosynthesis (R2=0.96;
data from [52]); **d**, grain volume (R2=0.99; data from [53]); **e**, grain dry weight (R2=1.00;
data from [53]); **f**, stem non-structural carbohydrates content (R2=0.99; data from [54]).
(A-F), the solid lines are modeled data, the dots are measured data. **g**, carbon and
nitrogen fluxes (mmol) within the whole plant throughout the whole grain filling
period. Abbreviations: R_d, dark respiration; C fluxes, carbon fluxes; N fluxes, nitrogen
fluxes.

For the nitrogen budget, WACNI predicted 38.6% of newly absorbed nitrogen to be

assimilated in the root, while 61.4% to be in the leaf. The total plant nitrogen content

at the flowering stage was 49 mmol (see Methods); thus, the root nitrogen uptake after

flowering accounts for 29.0% of the whole season nitrogen uptake, which is close to

the reported value, i.e., 30% [55]. In addition, the nitrogen harvest index, i.e., the ratio

between the amount of nitrogen stored in the grain at the harvest stage and the total

plant nitrogen uptake during the whole growth season, was 62.8%, which is also in

the known range of 30-77% , based on a large scale survey on irrigated lowland rice

in tropical and subtropical Asia [56].

Carbon and nitrogen dynamics in leaves and grains during grain filling can be

reproduced in silico without re-allocation of enzyme activities across different

nitrogen regimes

We then tested the performance of WACNI under different field management

practices and environmental perturbations. Firstly, we simulated changes in the leaf

nitrogen concentration, the grain nitrogen concentration, and the grain dry weight

dynamics during the grain filling period under different nitrogen application rates

(Figure 3a-l). WACNI reproduced patterns of leaf nitrogen concentration dynamics

from the flowering stage to the harvest stage for all nitrogen treatment groups (Figure

3a, R^2=0.97; Figure 3d, R^2=0.92; Figure 3g, R^2=0.97; Figure 3j, R^2=0.94), except for

an over-estimate of the leaf nitrogen concentration beyond 28 days after flowering for

the low nitrogen and medium nitrogen groups (Figure 3d, g). Remarkably, WACNI

not only precisely predicted the relative values of grain nitrogen concentration and

grain yield at the harvest stage, but also reproduced the dynamic changes of grain

nitrogen concentration (Figure 3b, R^2=0.97; Figure 3e, R^2=0.97; Figure 3h, R^2=0.98;

Figure 3k, R^2=0.95) and the pattern of grain dry weight gain (Figure 3c, R^2=0.94;

Figure 3f, R^2=0.98; Figure 3i, R^2=1.00; Figure 3l, R^2=0.96) throughout the grain

filling period for all treatments, without re-allocating activity of metabolic and

transport processes across different nitrogen regimes.

Figure 3. Predicted relative changes of plant physiological status and agronomic traits
under different nitrogen application rates (a-l), light regimes (m-s), and air CO_2
concentrations (t-u). Dynamic changes of leaf nitrogen concentration, grain nitrogen

concentration and grain dry weight with no nitrogen (**a**, R^2=0.97; **b**, R^2=0.97; **c**,
R^2=0.94), low level nitrogen (**d**, R^2=0.92; **e**, R^2=0.97; **f**, R^2=0.98), medium level
nitrogen (**g**, R^2=0.97; **h**, R^2=0.98; **i**, R^2=1.00) or high level nitrogen (**j**, R^2=0.94; **k**,
R^2=0.95; **l**, R^2=0.96) applied during the ear development period. Grain filling pattern
under normal condition (**m**, R^2=0.98); shading treatments during the first 10 days
after flowering which reduce the solar radiation by 25% (**n**, R^2=0.98), 50% (**o**,
R^2=0.99) and 75% (**p**, R^2=0.98); shading treatments followed by spacing (thinning the
every other plants to reduce the plant density by half; **q**, R^2=0.99; **r**, R^2=1.00; **s**,
R^2=0.99). Grain yield at the harvest stage (**t**) and total root nitrogen uptake during the
grain filling period (**u**) for plants grown under ambient CO_2 concentration with low
nitrogen ('LN'), medium nitrogen ('MN') and high nitrogen ('HN') application rates,
and plants grown under free air CO_2 enrichment (FACE) condition with low nitrogen
('LN+FACE'), medium nitrogen ('MN+FACE') and high nitrogen ('HN+FACE')
application rates. (**a-s**), the dots are measured data, the solid lines are modeled data.
(**t-u**), the black bars are measured data, the white bars are modeled data, the grey bars
are modeled data with additional assumption of the maximal root nitrogen uptake rate
under FACE conditions to be 64% of that under ambient conditions. The experimental
data are from [57] for panels a-l, from [58] for panels m-s, from [59] for panels t-u.
Abbreviations: LNC, leaf nitrogen concentration; GNC, grain nitrogen concentration;
GDW, grain dry weight; DAF, days after flowering.

Low light during early grain filling period affects grain weight gain pattern but does

NOT compromise rice yield potential

When we simulated grain filling under different light regimes (Figure 3m-s), WACNI

predicted an increased penalization to grain dry weight at 10 days after flowering, as

well as at the harvest stage when 25%, 50% or 75% of the incident irradiance was

shaded during the first 10 days after flowering (Figure 3m, R^2=0.98; Figure 3n,

R^2=0.98; Figure 3o, R^2=0.99; Figure 3p, R^2=0.98). In addition, WACNI successfully

predicted an acceleration of grain filling rate beyond 10 days after flowering when

extra light penetrates into the canopy as a result of plant thinning. Thus, it provides

theoretical support to the idea that short-term impaired grain filling by low light-

induced shortage of assimilates can almost be fully compensated if sufficient resource

is provided later (Figure 3q, R^2=0.99; Figure 3r, R^2=1.00; Figure 3s, R^2=0.99)[58].

Inhibited root nitrogen uptake capacity during grain filling under elevated air CO_2

concentration underlies reduced plant nitrogen budget

When we simulated grain yield and root nitrogen uptake under different atmospheric

CO_2 concentration, and soil nitrogen application rates, WACNI correctly predicted

that grain yield increased with nitrogen application rate under both ambient CO_2 level

and elevated CO_2 concentration (Figure 3t). Moreover, simulations using WACNI also

confirmed the observation that grain yield would benefit more from the elevated CO_2

concentration under medium and high nitrogen application rates (10.8% and 11%

enhancement of grain yield, respectively) than that under low nitrogen application rate

(1% enhancement of grain yield; Figure 3t). However, when plants with the same

nitrogen application rates were compared, WACNI predicted higher total root nitrogen

uptake during the grain filling period of plants grown under elevated CO_2

concentration than that of plants grown under ambient CO_2 concentration, which was,

however, against the experimental observation (Figure 3u). Earlier, the nitrogen

uptake capacity per root dry weight was shown to be unaffected under FACE at the

young panicle development period, but reduced by 31-41% from the flowering stage

to the harvest stage compared to the plants grown under ambient CO_2 concentration

[60]. Remarkably, if a mean percentage reduction of maximal root nitrogen uptake rate,

36%, was set under elevated air CO_2 concentration in WACNI, both the relative trend

of grain yield and root nitrogen uptake during the grain filling period could be

reproduced (the 'v_{m_Nupt}=64% under FACE' labeled simulations in Figure 3t, u),

which demonstrates the importance of root nitrogen uptake capacity in characterizing

plant carbon/nitrogen budget under elevated air CO_2 concentration.

Prolonged leaf functional duration greatly benefits grain yield with a slight reduction

in grain nitrogen concentration

We further tested WACNI by predicting the responses of grain yield, plant

physiological traits and agronomic traits to genetic manipulations that affect source

sink relation. We first simulated the impact on grain yield related traits when the

senescence of the major carbon source, leaf, was perturbed. We simulated leaf

nitrogen concentration dynamics during the grain filling period, the grain yield and

grain nitrogen concentration at the harvest stage for rice plants with different activities

of leaf protein degradation. According to Liang, et al. [61], three scenarios could be

examined, and we indeed showed: (1) promoted leaf protein degradation activity

mimicking the *ps1-D* mutant and *OsNAP* overexpression plants (the '*ps1-D*' and 'OE'

groups in Figure 4a-c); (2) medium leaf protein degradation activity mimicking wild

type plants (the 'WT' group in Figure 4a-c); and (3) reduced leaf protein degradation

activity mimicking *OsNAP* RNAi plants (the 'RNAi' group in Figure 4b-c). As a

result, WACNI predicted a faster leaf nitrogen loss of the *ps1-D* mutant plants, as a

result of the accelerated leaf protein degradation rate, which mimicked the observed

faster decrease in chlorophyll content in the *ps1-D* mutant (Figure 4a, R^2=0.86 for WT

and R^2=0.92 for ps1-D mutant). Concurrently, WACNI also predicted an increase of

grain yield at the expense of a slight decrease in grain nitrogen concentration for the

'RNAi' plant and the opposite for the 'OE' group plants, both of which coincided well

with experimental observations (Figure 4b, c). The enhanced grain yield was mainly

contributed by the increased higher canopy photosynthesis in the later grain filling

period as a result of larger leaf area and higher leaf nitrogen concentration, whereas

the reduced grain nitrogen concentration was mainly attributed to the decreased

nitrogen concentration in the phloem as a result of less leaf nitrogen remobilization

(Supplementary Figure 1; Figure 4a).

Figure 4. Predicted relative changes of plant physiological status and agronomic traits under different source (a-c), transport (d-i) or sink (j-l) related genetic manipulations. **a**, leaf nitrogen concentration dynamic change during the grain filling period for the wild type (WT; R^2=0.86) and a *OsNAP* gain-of-function mutant, prematurely senile 1 (*ps1-D* mutant; R^2=0.92). Relative grain yield (**b**) and grain nitrogen concentration (**c**) of *ps1-D* mutant and the *OsNAP* RNAi transgenic plant to the WT. **d**, grain dry weight dynamic change of the wild type (WT; R^2=0.98), a cell-wall invertase gene loss-of-function mutant, grain incomplete filling 1 (*gif1* mutant; R^2=0.98), and the *GIF1* over-expression transgenic line (*GIF1* OE). **e**, relative changes of grain yield, grain nitrogen concentration and harvest index for plants expressing a modified maize AGPase large subunit sequence (*Sh2r6hs*) under control of an endosperm-specific promoter. The experimental data are from [61] for panels a-c, from [62] for panel d, from [63] for panel e. Abbreviations: LNC, leaf nitrogen concentration; GNC, grain nitrogen concentration; N.M., not measured.

Sucrose transport capacity is a major factor determining grain filling rate

We then simulated the consequences of perturbing the sucrose transport system. Here

we tested the effect of changing capacity of sucrose unloading into the grain on grain

filling rate and the final grain yield. Wang, et al. [62] reported that a gene *GIF1*, which

encodes a cell-wall invertase, is required for sucrose transport to the grain. In

agreement with the experimental results, grain filling rate was predicted to be faster

for *GIF1* overexpression plants, which had a higher sucrose unloading capacity.

Further, the grain filling rate was predicted to be slower for *gif1* mutant plants which

had decreased sucrose unloading capacity (Figure 4d, R^2=0.98 for both WT and *gif1*

mutant). Although Wang, et al. [62] had suggested that the *GIF1* gene could increase the

yield potential through improved grain-filling, the grain yield of *GIF1* overexpression

plants was not shown in their experiments. Here, we simulated two scenarios: (1)

plants with parameter values set according to Wang, et al. [62], which represented a

'sink limited' scenario; (2) plants with default parameter values except that the grain

number per ear was set to 190, which represents a 'source limited' scenario. As a

result, when plants were 'sink limited', enhancing maximal rate of sucrose unloading

to the grain ($v_{m\_grain\_Suc\_ul}$) increased both the grain filling rate and the final grain

yield, whereas reducing $v_{m\_grain\_Suc\_ul}$ decreased the grain filling rate , as well as the

final grain yield (Supplementary Figure 2). However, when plants were 'source

limited', enhancing $v_{m\_grain\_Suc\_ul}$ could increase grain filling rate only in the early

phase of grain filling period, whereas the final grain yield even decreased, which was

mainly due to faster sink growth induced accelerated leaf nitrogen remobilization and

the consequently earlier leaf senescence (Supplementary Figure 3). We note that

although larger sugar flow directed towards sinks may relieve 'feedback

downregulation' of photosynthesis, for those 'source limited' cultivars, the unloading

rate of sucrose to grains should be limited to avoid unusually fast draining of nutrients

from the leaves since this causes earlier senescence of the whole plant. Consistent

with this notion, overexpression of *PbSWEET4*, a sugar transporter gene in pear, is

known to cause early senescence in leaves [64]. Similarly, overexpression of

OsSWEET5 in rice, which encodes a sugar transporter protein in leaves, stem, root

and floral organs, causes precocious leaf senescence [65].

Enhanced endosperm ADP-glucose pyrophosphorylase activity is not the mechanism

   for rice with ectopic expression of Sh2r6hs to gain increased grain yield

Here, we present our results on simulating the consequences of altering the grain

storage capacity. Endosperm-specific ectopic expression of a modified maize ADP-

glucose pyrophosphorylase (AGPase) large subunit sequence (*Sh2r6hs*) has been

shown to enhance the activity of starch synthesis in the endosperm of both wheat and

rice [63,66]. In our work, we simulated the effects of increasing the maximal grain starch

synthesis rate ($v_{m\_grain\_Star\_syn}$) to 150% of its default value, on grain yield and yield

related traits; this represented an *in silico Sh2r6hs* ectopic expression plant. As a

reference, in the actual *Sh2r6hs* transgenic plants, the measured AGPase activity in

developing grains was higher (from as small as 9% to as high as 52%) than in wild

type plants, 10-20 days after flowering [63]. We note that WACNI realistically predicted

the increase of grain yield and the maintenance of harvest index, although it predicted

a slight decrease, rather than the observed slight increase, in grain nitrogen

concentration for the *Sh2r6hs* transgenic plants at the harvest time (Figure 4e).

Further, we note that although *Sh2r6hs* was expressed under the control of an

endosperm-specific promoter, the size of transgenic plants at the flowering differed

from that of wild type plants. Thus, we tested whether the increase of the grain yield

was mainly caused by the increased AGPase activity during grain filling, or was it

mainly caused by the plant size change at the flowering stage. To answer this

question, we simulated three scenarios: (1) we changed both the plant size at the

flowering stage and $v_{m\_grain\_Star\_syn}$ during grain filling for *Sh2r6hs* transgenic plant; (2)

we changed only the plant size at the flowering stage for *Sh2r6hs* transgenic plant; (3)

we changed only $v_{\text{m\_grain\_Star\_syn}}$ during grain filling for *Sh2r6hs* transgenic plant.

Simulation results matched experimental observations for both the first and the

second scenarios, but not for the third scenario (Figure 4e). These data demonstrate

that the increased grain yield in *Sh2r6hs* transgenic plants is mainly contributed by

change of the plant size at the flowering stage, rather than being contributed by the

enhancement of endosperm starch synthesis capacity during grain filling.

Mapping genetic variations of capacities of reaction and diffusion processes to

phenotypic variations of grain yield and yield-related agronomic traits

After validating WACNI against experiments under various environmental treatments

and genetic manipulations, we used it to explore influence of the capacity of 28

reaction and diffusion processes on eight agronomic traits (Supplementary Figures 4-

10). Based on the predicted response patterns of grain yield to parameter changes, we

classified the 28 parameters into four categories. Specifically, (1) eleven parameters

that can monotonically enhance grain yield were termed as universal yield enhancers

(UYEs; shown in Figure 5a); (2) four parameters that can monotonically inhibit grain

yield were termed as universal yield inhibitors (UYIs; shown in Figure 5b); (3) six

parameters that can non-linearly influence grain yield were termed as conditional

yield enhancers (CYEs; shown in Figure 5c); (4) seven parameters having little or no

influence on grain yield were termed as weak yield regulators (WYRs; exemplified in

Figure 5d).

Figure 5. Predicting and grouping the effects of manipulating capacities of 28
reaction and diffusion processes on grain yield and yield-related agronomic traits. **a**,
leaf sucrose loading capacity ($v_{m\_leaf\_Suc\_l}$) is an example illustrating a universal yield
enhancer which can monotonically increase grain yield when its value ranges from
[0.1, 10]-fold to its default value. **b**, leaf protein degradation capacity $v_{m\_leaf\_Pro\_deg}$ is
an example illustrating a universal yield inhibitor which can monotonically decrease
grain yield. **c**, grain growth capacity $v_{m\_grain\_grow}$ is an example illustrating a
conditional yield enhancer which can non-linearly influence grain yield. **d**, root
organic nitrogen unloading capacity $v_{m\_root\_ON\_ul}$ is an example illustrating a weak
yield regulator which can hardly influence grain yield. **e**, distribution of false
prediction rate on agronomic traits for 33 of the different genetic manipulation cases
shown in Supplementary Table 2 with at least one agronomic trait measured. **f**,
distribution of consistent, uncertain and inconsistent predictions with experimental
observations on eight agronomic traits for genetic manipulations shown in
Supplementary Table 2.

We further compiled the genes and QTLs in rice controlling the above 28 parameters

published thus far. Thirty-eight genes/QTLs were collected, among which there were

33 manipulations from which at least one of the eight agronomic traits was measured

(Supplementary Table 2; see a brief functional description of each gene/QTL in

Supplementary Table 3). We emphasize that WACNI had a strong predictive power on

the response patterns of agronomic traits by genetic manipulations. For example,

WACNI predicted a strong negative impact of impaired $v_{m\_leaf\_Suc\_l}$ on the grain filling

rate, the total canopy photosynthesis, the harvest index and the final grain yield;

although the grain yield does not decrease with increasing $v_{m\_leaf\_Suc\_l}$, increasing the

$v_{m\_leaf\_Suc\_l}$ to be higher than 60% of the default value can no longer increase grain yield

(Supplementary Figure 5; Figure 5a). Consistently, it was observed that the impaired

function of *OsSUT2*, which exports sugar from the leaves, led to severe growth

retardation and yield decrease, whereas higher expression of *OsSUT2* did not result in

increased yield compared to that of wild type plants [67]. We further calculated the false

prediction rate for each genetic manipulation case. False prediction rate defines the

ratio between number of predictions failed to match the experimental observations

and the number of all experimentally observed agronomic traits. For example,

response patterns of three agronomic traits were measured experimentally ('T1', 'T2'

and 'T3') for *OsFd-GOGAT* mutant plants, among which WACNI correctly predicted

the response patterns of two ('T1' and 'T3'), but failed to predict the response pattern

of 'T2' (Supplementary Table 2); in this case, the false prediction rate was 33.3%. In

all the 33 genetic manipulation cases, the mean and the median false prediction rates

were 11.3% and 0%, respectively (Figure 5e; Supplementary Table 2).

Although none of the above mentioned studies (Supplementary Table 2) measured all

the eight agronomic traits ('T1'-'T8') simultaneously, the response patterns of each

agronomic trait to different genetic manipulations were predicted with high accuracy

across all our measurements (Figure 5f). Specifically, the false prediction rates for

eight agronomic traits were 10%, 17%, 0%, 14%, 0%, 0%, 50%, 0%, respectively

('Inconsistent' in Figure 5f-m). Besides, non-linear effects of certain genetic

manipulations on some agronomic traits exist (Supplementary Figures 4-10). In these

cases, the simulation results cannot be determined as consistent with experimental

observations from the published work or not, due to the lack of information on

numeric values of the manipulated parameters (labelled as 'Uncertain' in Figure 5f).

Designing rice physiological ideotype for super-high yield

Finally, WACNI was combined with a genetic algorithm to search the optimal

parameter values for maximizing the grain yield. Specifically, we first randomly

varied values of specific parameters in a range from 50% to 200% to their default

values to generate five independent populations; then we "evolved" these five

populations using high grain yield as the selection target. At the end of 100

generations of *in silico* evolution, the highest grain yields in these five populations

were found to be similar (Supplementary Figure 11). The top 1% individuals ranking

by the final grain yield in each population were then merged to form an elite plant

group (the 'Elite' group in Figure 6b-k and Supplementary Figure 12). In general,

values of UYEs were up-regulated, values of UYIs were down-regulated, but the

values of CYEs and WYRs did not form any particular pattern (Figure 6a). Although

'Elite' individuals had similar high grain yield, parameter values for these individuals

varied. Some parameters were conserved with small variance among 'Elite'

individuals, e.g. $v_{m\_grain\_Star\_syn}$, $v_{m\_leaf\_Pro\_deg}$, $v_{m\_leaf\_ON\_l}$ and $v_{m\_root\_Suc\_ul}$; on the

contrary, up to a 4-fold differences in values exist for some other parameters, e.g.

$v_{m\_leaf\_N\_ass}$, $v_{m\_leaf\_IN\_ul}$, $v_{m\_grain\_Pro\_syn}$ and $v_{m\_stem\_Pro\_syn}$ (Figure 6a-c). The initial

partitioning of carbon and nitrogen among organs at the flowering stage for 'Elite'

individuals was distinct from the 'Ctrl' individual (individual with default model

parameter values), e.g. the grain number increased while the root size decreased

dramatically for 'Elite' individuals (Supplementary Figure 12). Overall, the

optimization of biochemical and physiological parameters resulted in a remarkable

(54%)increase of the grain yield, at the expense of a 37% decrease in grain nitrogen

concentration (Figure 6e-f).

Figure 6. Designing rice physiological ideotype for super-high yield by optimizing the model parameters. Distribution of relative value of each biochemical/physiological parameter (fold change from its default value) for *in silico* high-yield individuals, including universal yield enhancers (**a**), universal yield inhibitors (**b**), conditional yield enhancers (**c**) and weak yield regulators (**d**). Comparison of grain yield per plant (**e**) and grain nitrogen concentration (**f**) for high-yield individuals (n=500, 'Elite') and grain yield of individual with default parameters ('Ctrl'). **g-n**, comparison of macroscopic physiological dynamic changes of the 'Ctrl' individual, the 'Elite' individuals, the 'Best' individual with highest grain yield in 'Elite', and 'All' other individuals generated during *in silico* evolution. See detailed information of parameter values for 'Elite' individuals (a-d) in Supplementary Datasets 1. Abbreviations: NSC, non-structural carbohydrates; $[O\text{-}N]_p$, phloem O-N concentration.

'Elite' individuals also show distinct patterns with regard to macroscopic

physiological features compared to other individuals. For example, the grain filling

rate of 'Elite' individuals was relatively low during the first 7 days after flowering,

but was moderate between 7 to 25 days after flowering, whereas the rate decreased at

much slower pace, starting from 20 days after flowering to the harvest stage, when

compared to 'Ctrl' and other individuals (Figure 6g). This 'steady and sustained' grain

filling rate resulted in a nearly linear grain weight gain for the 'Elite' individuals

rather than the usual sigmoid curve, as for 'Ctrl' and other individuals (Supplementary

Figure 13). Grain nitrogen concentration was constantly lower during grain filling for

the 'Elite' individuals (Figure 6h). Further, the 'Elite' individuals had slower rate of

decrease in both the leaf area as well as in the rate of canopy photosynthesis (Figure

6i-j), a feature commonly known as 'stay green' [68]. Surprisingly, 'Elite' individuals

had a much smaller root mass throughout the grain filling period, and also a lower

total root nitrogen uptake (Figure 6k-l). Notably, some 'Elite' individuals had similar

root nitrogen uptake as the 'Ctrl' plants did (Figure 6k), as a result of their higher root

nitrogen uptake capacity ($v_{m\_root\_N\_upt}$, Figure 6a). Finally, the 'Elite' individuals had

similar stem non-structural carbohydrate use pattern as 'Ctrl' plants throughout the

grain filling period (Figure 6m); on the other hand, the organic nitrogen concentration

in phloem ($[O-N]_p$) was at a low level in the 'Elite' individuals starting from 10 days

after flowering to the harvest stage; this is in contrast to 'Ctrl' and many other plants,

where $[O-N]_p$ dramatically increases during the late grain filling period (Figure 6n).

DISCUSSION

Despite the descriptive research of source sink interaction in single cells or organs, the

plant level coordination of processes in different source and sink organs has been

largely disregarded during crop improvement research; this has been due to

difficulties in the interpretation of such a complex network [29]. Crop systems models

have been used in designing agronomic practices [69,70], testing hypotheses regarding

the responses of crops to climate change [71,72], guiding selection of traits for breeding

[73] and assessing different strategies to adapt to future climate change [74]. Here, we have

introduced a mechanistic model that accurately predicts plant level source sink

interactions during grain filling, as an emergent property, from basic biochemical and

physiological processes in different plant organs (Figure 1); this clearly differs from

the mainstream approaches using either preset rules on the relationship between the

source and sink organs, or the empirical correlation derived from the measured data.

Considering the numerous processes within different organs and the extensive

metabolite exchange between organs, it is remarkable that WACNI accurately predicts

physiological changes of a rice plant during the grain filling period both under normal

condition as well as under various field practices and/or environmental perturbations

(Figures 2-3). Further analysis using WACNI confirms many conjectures raised over

the past decades on high-yield crop breeding and crop source-sink relations. For

example, functional 'stay green' of leaves, higher leaf sucrose export rate, higher

endosperm starch synthesis capacity and more efficient remobilization of stem starch

reserves are among the major means considered for increasing crop yield (Figure 6a,

i, j, m)[75-78]. Moreover, WACNI systematically evaluates the reported impact on

agronomic traits of genetic manipulations of source, transport and sink related

properties (Figures 4-5; Supplementary Table 2). In particular, WACNI correctly

predicts 64-100% of the responses to all 33 genetic manipulations, for six out of eight

major agronomic traits, with the remaining two traits, harvest index and nitrogen

harvest index, having very low (only 1 to 2) number of reported values (Figure 5f)

and hence cannot be properly evaluated. All these results demonstrate that WACNI is

a robust mechanistic model linking genotypic variation to phenotypic variation under

different environments.

Besides its mechanistic basis, another major feature that confers the high prediction

accuracy of WACNI is its parameterization routine. Parameterization of WACNI relies

on extensive literature survey (Supplementary Table 1) , resulting in a primary

reference dataset on natural variation of important rice traits, which can be constantly

updated to support systems modeling of rice metabolism, growth and development. It

is also worth mentioning here that since most processes described in WACNI are

rather generic among plants, WACNI can be adapted and parameterized for other

plants, especially for cereal crops. In addition, WACNI follows a principle of modular

design, that is, the metabolic processes involved in leaf, grain, root, and stem are

modeled in individual modules, which can easily be replaced by other modules or

extended by other researchers whenever needed. This feature enables WACNI to be a

framework to support future crop systems models development, e.g. in developing

models of plant signaling transduction or hormone regulation [79] and more elaborate

models combining 3D root/shoot morphogenesis, development and functions [80-82].

Plant growth and developmental processes are ultimately supported by the metabolic

processes in different organs and fluxes between organs, establishing a map of the

carbon and nitrogen fluxes within a whole plant has been a major challenge of plant

biology [83]. Here, for the first time, we quantify the entire carbon and nitrogen budget

within a rice plant during the grain filling period using WACNI (Figure 2g). The map

drawn here opens new research opportunities. For example, we can use it in

evaluating how resource partitioning among sink organs influences the overall

efficiency of the whole system; furthermore, it can be used to precisely calculate

contribution from each source tissue to sink development. WACNI therefore provides

a theoretical tool to enable detailed study on controls and even genetic basis of carbon

and nitrogen fluxes in a plant. It is exciting to see the availability of comprehensive

datasets on whole plant level metabolic fluxes based on fluxomics [84].

Model guided enhancement of photosynthetic efficiency has gained major success in

recent years [2,40,42]. Being able to link molecular level processes to physiological

changes all the way up to agronomic traits, WACNI, for the first time, offers an

opportunity to design features associated with super high yield. Optimization of

source sink and transport related processes, based on WACNI prediction, show that

the rice grain yield potential can reach 21 t ha^{-1}, given that the water content is 13.5%

and husks account for 20% of weight in rough rice (Figure 6e). This is 40% higher

than the current highest record of rice grain yield [85]. Analysis based on WACNI

simulations suggests that to reach such a high yield one option is to have a dry weight

of ~22 mg per dehulled grain, a total number of filled spikelets being ~$6.7*10^8$ ha^{-1}

(Supplementary Datasets 1). In such a rice line, a balanced source sink relation is

needed to support 'stay green' of leaves and maximize both instantaneous and long-

term canopy photosynthesis, which together realize the 'steady and sustained' grain

filling rate till 50 days after flowering (Figure 6g). Many previously identified genes,

being either UYEs, UYIs, CYEs, can be used as targets of fine-tuning to control

carbon and nitrogen fluxes between different organs to realize the required

physiological features, ultimately leading to the designed high yield potential

(Supplementary Table 2).

Methods

Model development

The model developed here was depicted diagrammatically in Figure 1. The individual

modules were developed following the basic procedure as in Zhu, et al. [86]. Essentially,

after the reaction diagrams were established, differential equations, rate equations, and

algebraic equations representing conserved quantities were developed. The model was

implemented in MATLAB and solved using ode15s (the MathWorks Inc., Natick, MA,

USA). WACNI computed changes in the metabolite concentrations by the differences

between rates of fluxes generating and consuming metabolites.

Fourteen different types of primary biochemical/biophysical processes were

incorporated in WACNI (Figure 1). The rate equations were described in four different

subsections according to the organs they present in, i.e., root, leaf, grains and stem

(including culm and sheath) in the following (Eqn 1.1-14.1). Respiration for each organ

was described in the subsequent *Respiration* subsection (Eqn 15.1-15.4).

Root

*I-N uptake and assimilation***:** Two I-N uptake systems present in roots, i.e., the high

affinity transport system (HATS) and the low affinity transport system (LATS); which

together ensure I-N uptake at different soil N concentrations [87]. The uptake rates of

HATS and LATS were modeled with Michaelis-Menten kinetics model and linear

model, respectively (Eqn 1.2). Sugar level also influences mineral transport as sugar

provides energy needed for these processes [88,89]. The impact of root sucrose level on I-

N uptake was modelled with a Monod function [90], i.e., the first item of Eqn (1.2). Root

total N concentration has a feedback inhibitory effect ($\alpha_{N\_inhibit}$) on root HATS [87]. Finally,

these regulatory processes were incorporated into a single root I-N uptake equation to

calculate total root I-N uptake rate ($v_{root\_N\_upt}$):

591
$$\alpha_{N\_inhibit} = \begin{cases} 1 - \dfrac{[N]_{root}}{[N]_{inhibit\_up}}, [N]_{root} < [N]_{inhibit\_up} \\ 0, [N]_{root} \geq [N]_{inhibit\_up} \end{cases} \quad (1.1)$$

$$v_{\text{root\_N\_upt}} = \frac{[\text{Suc}]_{\text{root}}}{K_{\text{m1}} + [\text{Suc}]_{\text{root}}} \cdot (\alpha_{\text{N\_inhibit}} \cdot v_{\text{m\_root\_HATS}} \cdot \frac{[\text{I-N}]_{\text{soil}}}{K_{\text{m2}} + [\text{I-N}]_{\text{soil}}} + k_{\text{root\_LATS}} \cdot [\text{I-N}]_{\text{soil}}) \quad (1.2)$$

in which $[\text{I-N}]_{\text{soil}}$ is soil I-N concentration, $[\text{Suc}]_{\text{root}}$ is root sucrose concentration, $[\text{N}]_{\text{root}}$

is total root N concentration (including I-N and O-N). $[\text{N}]_{\text{inhibit\_up}}$ is upper limit of total

root N concentration above which N uptake by HATS ceases, K_{m1} and K_{m2} are

Michaelis-constants, $v_{\text{m\_root\_HATS}}$ is root HATS maximal I-N uptake rate, $k_{\text{root\_LATS}}$ is root

LATS I-N uptake rate coefficient (Supplementary Table 1).

Root N assimilation depends on energy and reducing power derived from sugar [91].

Hence, root I-N assimilation rate ($v_{\text{root\_N\_ass}}$) was modelled based on $[\text{Suc}]_{\text{root}}$ and $[\text{I-}$

$\text{N}]_{\text{root}}$:

$$v_{\text{root\_N\_ass}} = v_{\text{m\_root\_N\_ass}} \cdot \frac{[\text{Suc}]_{\text{root}}}{K_{\text{m3}} + [\text{Suc}]_{\text{root}}} \cdot \frac{[\text{I-N}]_{\text{root}}}{K_{\text{m4}} + [\text{I-N}]_{\text{root}}} \quad (2.1)$$

in which K_{m3} and K_{m4} are Michaelis-constants, $v_{\text{m\_root\_N\_ass}}$ is root maximal I-N

assimilation rate (Supplementary Table 1).

Root growth: As sugar supply is a major factor limiting root growth [92], and organ

growth rate is positively correlated with sugar level in the organ before reaching its

maximal capacity [92-94], a critical sucrose concentration ($[\text{Suc}]_{\text{grow\_low}}$) was set, below

which root growth ceased. A Monod function was applied to describe the relation

between root sucrose level and its growth rate ($v_{\text{root\_grow}}$):

$$v_{\text{root\_grow}} = \begin{cases} 0, [\text{Suc}]_{\text{root}} < [\text{Suc}]_{\text{grow\_low}} \\ v_{\text{m\_root\_grow}} \cdot \frac{[\text{Suc}]_{\text{root}} - [\text{Suc}]_{\text{grow\_low}}}{K_{\text{m5}} + ([\text{Suc}]_{\text{root}} - [\text{Suc}]_{\text{grow\_low}})}, [\text{Suc}]_{\text{root}} \geq [\text{Suc}]_{\text{grow\_low}} \end{cases} \quad (3.1)$$

in which K_{m5} is Michaelis-constant, $v_{\text{m\_root\_grow}}$ is the root maximal relative growth rate

(Supplementary Table 1).

Root senescence: Root loss during senescence was modelled based on two factors,

aging and carbohydrate supply. Specifically, there is a minimal (constant) relative

senescence rate $\alpha_{\text{sene\_root}}$ (Supplementary Table 1) due to aging of root [95,96], but when

sucrose concentration is lower than a critical level ($[\text{Suc}]_{\text{sene\_up}}$), root senescence rate

($v_{\text{root\_sene}}$) would be accelerated as a result of carbohydrate starvation [97]:

618
$$v_{\text{root\_sene}} = \begin{cases} \alpha_{\text{root\_sene}} \cdot e^{-\beta_{\text{root\_sene}} \cdot ([\text{Suc}]_{\text{root}} - [\text{Suc}]_{\text{sene\_up}})}, [\text{Suc}]_{\text{root}} < [\text{Suc}]_{\text{sene\_up}} \\ \alpha_{\text{root\_sene\_}}, [\text{Suc}]_{\text{root}} \geq [\text{Suc}]_{\text{sene\_up}} \end{cases} \quad (4.1)$$

in which $\beta_{\text{root\_sene}}$ is an empirical coefficient (Supplementary Table 1).

Leaf

Photosynthesis and N assimilation: Leaf level light reaction, Calvin-Benson cycle and

photorespiration were modelled based on the Farquhar-von Caemmerer-Berry (FvB)

model [98]. To scale up them to a canopy level, a sun-shade model described in De Pury

and Farquhar [99] was used (see below).

Leaf protein level (especially the level of enzyme Rubisco, which accounts for major

leaf protein) has been shown to be linearly and positively correlated with leaf light

reaction and dark reaction activity before reaching their maximal capacity in many

experiments [100,101], while leaf non-structural carbohydrates (NSC, refer to sucrose and

starch in the current model) accumulation can inhibit photosynthesis [102] through

multiple feedback regulatory pathways [103,104]. These effects were considered by setting

an activation coefficient ($\alpha_{\text{Pro\_promote}}$, Eqn 5.1) of leaf protein content ([Pro]$_{\text{leaf}}$) and an

inhibition coefficient ($\alpha_{\text{NSC\_inhibit}}$, Eqn 5.2) of leaf NSC content ([NSC]$_{\text{leaf}}$) to parameters

involved in both photosynthetic electron transport rate v_J and potential CO_2 assimilation

rate v_{A0} (Eqn 5.3-5.8):

635
$$\alpha_{\text{Pro\_promote}} = \begin{cases} \dfrac{[\text{Pro}]_{\text{leaf}}}{[\text{Pro}]_{\text{promote\_up}}}, [\text{Pro}]_{\text{leaf}} < [\text{Pro}]_{\text{promote\_up}} \\ 1, [\text{Pro}]_{\text{leaf}} \geq [\text{Pro}]_{\text{promote\_up}} \end{cases} \quad (5.1)$$

636
$$\alpha_{\text{NSC\_inhibit}} = \begin{cases} 1, [\text{NSC}]_{\text{leaf}} < [\text{NSC}]_{\text{inhibit\_low}} \\ 1 - \dfrac{[\text{NSC}]_{\text{leaf}} - [\text{NSC}]_{\text{inhibit\_low}}}{[\text{NSC}]_{\text{inhibit\_up}} - [\text{NSC}]_{\text{inhibit\_low}}}, [\text{NSC}]_{\text{inhibit\_low}} \leq [\text{NSC}]_{\text{leaf}} \leq [\text{NSC}]_{\text{inhibit\_up}} \\ 0, [\text{NSC}]_{\text{leaf}} > [\text{NSC}]_{\text{inhibit\_up}} \end{cases}$$

637
$$(5.2)$$

638
$$v_{\text{cmax}} = \alpha_{\text{Pro\_promote}} \cdot \alpha_{\text{NSC\_inhibit}} \cdot v_{\text{cmax0}} \quad (5.3)$$

639
$$J_{\text{max}} = \alpha_{\text{Pro\_promote}} \cdot \alpha_{\text{NSC\_inhibit}} \cdot J_{\text{max0}} \quad (5.4)$$

640
$$\theta = \alpha_{\text{Pro\_promote}} \cdot \theta_0 \quad (5.5)$$

$$\varphi = \alpha_{\text{Pro\_promote}} \cdot \varphi_0 \qquad (5.6)$$

$$v_{A0} = v_{\text{cmax}} \cdot \frac{[CO_2]_i - \Gamma}{K_{m6} + [CO_2]_i} \qquad (5.7)$$

$$v_J = \frac{\varphi \cdot I + J_{\max} - \sqrt{(\varphi \cdot I + J_{\max})^2 - 4 \cdot \varphi \cdot \theta \cdot I \cdot J_{\max}}}{2\theta} \qquad (5.8)$$

in which K_{m6} is Michaelis-constant; $[Pro]_{\text{promote\_up}}$, $[NSC]_{\text{inhibit\_up}}$ and $[NSC]_{\text{inhibit\_low}}$ are critical leaf protein and NSC content; $\varphi_0 = 0.85 \ast 0.5$ and $\theta_0 = 0.7$ are empirical constants [98]; $v_{\text{cmax}0}$ is the potential maximal Rubisco carboxylation rate, $J_{\max 0}$ is the potential maximal electron transport rate (Supplementary Table 1); Γ is CO_2 compensation point in the absence of mitochondrial respiration (Supplementary Table 1). I is irradiance absorbed by leaves. Whole-plant I, $v_{\text{cmax}0}$ and $J_{\max 0}$ were calculated separately for sunlit and shaded leaves, e.g. I_{sun}, I_{sh}, $v_{\text{cmax}0\_\text{sun}}$, $v_{\text{cmax}0\_\text{sh}}$, $J_{\max 0\_\text{sun}}$ and $J_{\max 0\_\text{sh}}$ in Eqn 5.14-5.19, and $v_{\_\text{leaf\_N\_ass}0}$ in Eqn 5.23, based on light extinction profile within the canopy throughout the day following De Pury and Farquhar [99]:

$$\delta = -23.4 \frac{\pi}{180} \cos \frac{2\pi(\text{DOY} + 10)}{365} \qquad (5.9)$$

$$\sin \beta = \sin \lambda \cdot \sin \delta + \cos \lambda \cdot \cos \delta \cdot \cos \frac{\pi \cdot (\tau - 12)}{12} \qquad (5.10)$$

$$f_d = \frac{1 - a^{1/\sin \beta}}{1 + a^{1/\sin \beta}(1/f_a - 1)} \qquad (5.11)$$

$$L_{C\_\text{sun}} = \frac{1 - e^{-k_b \cdot L_C}}{k_b}(1 - e^{-k_b \cdot L_C}) \qquad (5.12)$$

$$I_C = \int_0^{L_C} I_l dl = (1 - \rho_{cb}) \cdot I_b \cdot (1 - e^{-k_b' \cdot L_C}) + (1 - \rho_{cd}) \cdot I_d \cdot (1 - e^{-k_d' \cdot L_C}) \qquad (5.13)$$

$$I_{\text{sun}} = \int_0^{L_C} I_{l\_\text{sun}} \cdot f_{l\_\text{sun}} dl = I_b \cdot (1 - \sigma) \cdot (1 - e^{-k_b \cdot L_C}) + I_d \cdot (1 - \rho_{cd}) \cdot (1 - e^{(-k_d' + k_b) \cdot L_C}) \cdot \frac{k_d'}{k_d' + k_b}$$

$$+ I_b \cdot (1 - \rho_{cb}) \cdot (1 - e^{(-k_b' + k_b) \cdot L_C} \cdot \frac{k_b'}{k_b' + k_b}) - (1 - \sigma) \cdot \frac{1 - e^{-2k_b \cdot L_C}}{2}$$

$$(5.14)$$

$$I_{sh} = I_C - I_{sun} \qquad (5.15)$$

$$v_{cmax0\_sun} = v_{cmax0} \cdot L_{C\_sun} \qquad (5.16)$$

$$J_{max0\_sun} = J_{max0} \cdot L_{C\_sun} \qquad (5.17)$$

$$v_{cmax0\_sh} = v_{cmax0} \cdot (L_C - L_{C\_sun}) \qquad (5.18)$$

$$J_{max0\_sh} = J_{max0} \cdot (L_C - L_{C\_sun}) \qquad (5.19)$$

in which δ is solar declination angle; DOY is current day of year; β is solar elevation

angle; λ is latitude of experimental base; τ is local time of day; f_d is fraction of diffuse

irradiance; f_a=0.425 is forward scattering coefficient of PAR in atmosphere; a=0.75 is

atmospheric transmission coefficient of PAR; L_c is canopy leaf area index; L_{c_sun} is the

sunlit leaf area index; $k_b = k_{b0}/\sin\beta$ is beam radiation extinction coefficient of canopy;

$k'_b = 0.9 k_b$ is beam and scattered beam PAR extinction coefficient; k_d=0.78 is diffuse

PAR extinction coefficient; $k'_d = 0.9 k_d$ is diffuse and scattered diffuse PAR extinction

coefficient; I_0 is total incident PAR intensity at top of canopy; $I_d = I_0 \cdot f_d$ is diffuse PAR

intensity; $I_b = I_0 \cdot (1-f_d)$ is beam PAR intensity; I_C is irradiance absorbed by the canopy;

I_{sun} is irradiance absorbed by the sunlit leaves; I_{sh} is irradiance absorbed by the shaded

leaves; ρ_{cb}=0.029 is canopy reflection coefficient for beam PAR; ρ_{cd}=0.036 is canopy

reflection coefficient for diffuse PAR; σ=0.15 is leaf scattering coefficient of PAR.

More detailed information of these processes and parameters can be found in De Pury

and Farquhar [99].

Intercellular CO_2 concentration ($[CO_2]_i$) is determined by ambient CO_2 concentration

($[CO_2]_a$), leaf stomatal conductance (g_s) and leaf net photosynthetic rate (A - R_{leaf}):

681
$$[CO_2]_i = [CO_2]_a - \frac{A - R_{leaf}}{g_s} \qquad (5.20)$$

in which leaf gross photosynthetic rate A is calculated in Eqn (5.27), and respiration

rate R_{leaf} is calculated in the following *Respiration* subsection.

Leaf stomatal conductance (g_s) is influenced by many factors, e.g. CO_2 concentration,

vapor pressure deficit, light intensity, leaf water potential, temperature and abscisic acid

concentration [105-107]. Diffusion of CO_2 from ambient air to intercellular space was

modeled following Leuning [108]:

$$g_\text{s} = g_0 + a_1 \cdot \frac{A}{[CO_2]_\text{i} - \Gamma} \cdot \frac{1}{1 + \dfrac{VPD_\text{leaf}}{VPD_0}} \quad (5.21)$$

in which $g_0=0.01$ is residual stomatal conductance, $a_1=20$ and $VPD_0=0.35$ are empirical

constants [108]. Leaf vapor pressure deficit (VPD_leaf) is the difference between leaf vapor

pressure (which is assumed to be saturated and determined by leaf temperature T) and

actual air vapor pressure (which is determined by relative air humidity RH_air), and was

calculated following Allen, et al. [109]:

$$VPD_\text{leaf} = 0.6108 \exp[\frac{17.27T}{T + 237.3}] \cdot (1 - RH_\text{air}) \quad (5.22)$$

As photosynthetic light reaction generates abundant ATP and reducing power (e.g.

NADPH and Ferredoxin), which are necessary for I-N assimilation, leaf acts as a major

N assimilator as well as root [110]. Potential I-N assimilation rate ($v_\text{N\_ass0}$) was modeled

based on leaf protein level and I-N concentration ($[I\text{-}N]_\text{leaf}$):

$$v_\text{leaf\_N\_ass0} = \alpha_\text{Pro\_promote} \cdot v_\text{m\_leaf\_N\_ass} \cdot \frac{[I\text{-}N]_\text{leaf}}{K_\text{m7} + [I\text{-}N]_\text{leaf}} \quad (5.23)$$

in which K_m7 is Michaelis-constant, $v_\text{m\_leaf\_N\_ass}$ is leaf maximal N assimilation rate

(Supplementary Table 1).

Calvin cycle, photorespiration and I-N assimilation compete for reducing power

derived from light reaction when light is limiting [111]. Light reaction derived reducing

power producing rate ($v_\text{NADPH\_p}$) was calculated as following:

$$v_\text{NADPH\_p} = \frac{v_\text{J}}{2} \quad (5.24)$$

Calvin cycle and photorespiration consume 2 NADPH per cycle, and average I-N

assimilation consumes 3 NADPH per N (if NO_3^- is the N source, it needs 5 NADPH or

equivalent reduction power; if NH_4^+ is the N source, it needs 1 NADPH or equivalent

reduction power) [112]. Thus the potential NADPH consume rate ($v_\text{NADPH\_c0}$) was

calculated as following:

$$v_\text{NADPH\_c0} = 2v_\text{cmax} \cdot \frac{[CO_2]_\text{i} + \Gamma}{K_\text{m6} + [CO_2]_\text{i}} + 3v_\text{leaf\_N\_ass0} \quad (5.25)$$

The actual CO_2 assimilation rate A and I-N assimilation rate ($v_\text{leaf\_N\_ass}$) were finally

determined by balancing NADPH production and consumption:

$$\delta = \begin{cases} \dfrac{v_{NADPH\_p}}{v_{NADPH\_c0}}, v_{NADPH\_p} < v_{NADPH\_c0} \\ 1, v_{NADPH\_p} \geq v_{NADPH\_c0} \end{cases} \quad (5.26)$$

$$A = \delta \cdot v_{A0} \quad (5.27)$$

$$v_{leaf\_N\_ass} = \delta \cdot v_{leaf\_N\_ass0} \quad (5.28)$$

717   Final CO_2 and I-N assimilation rate were determined by solving the above equations

718   (Eqn 5.1-5.28).

719   *Leaf C and N substrates homeostasis*: During the day, photosynthetic derived triose

720   phosphates (TP) convert to sucrose ($v_{leaf\_Suc\_syn}$) and starch ($v_{leaf\_Star\_syn}$) synchronously:

$$v_{leaf\_Suc\_syn} = v_{m\_leaf\_Suc\_syn} \cdot \frac{[TP]_{leaf} - \dfrac{[Suc]_{leaf}}{K_{e1}}}{K_{m8} + [TP]_{leaf}} \quad (6.1)$$

$$v_{leaf\_Star\_syn} = v_{m\_leaf\_Star\_syn} \cdot \frac{[TP]_{leaf}}{K_{m9} + [TP]_{leaf}} \quad (6.2)$$

in which K_{m8}, K_{m9} and K_{e1} are Michaelis-constants, $v_{m\_leaf\_Suc\_syn}$ is leaf maximal sucrose

synthesis (TP to sucrose conversion) rate and $v_{m\_leaf\_Star\_syn}$ is leaf maximal starch

synthesis (TP to starch conversion) rate (Supplementary Table 1).

In night, sucrose level decreases and daily synthesized starch degrades gradually [113].

There are a number of models simulating nighttime starch degradation, most of which

incorporate a circadian clock control of the process [114-116]. For simplification, we used

a simple substrates feedback regulation mechanism to simulate starch degradation rate

($v_{leaf\_Star\_deg}$):

$$v_{leaf\_Star\_deg} = \begin{cases} v_{m\_leaf\_Star\_deg} \cdot \dfrac{[Star]_{leaf}}{K_{m10} + [Star]_{leaf}} \cdot \dfrac{[Suc]_{leaf\_Star\_deg\_low} - [Suc]_{leaf}}{[Suc]_{leaf\_Star\_deg\_low} + [Suc]_{leaf}}, [Suc]_{leaf} < [Suc]_{leaf\_Star\_deg\_low} \\ 0, [Suc]_{leaf} \geq [Suc]_{leaf\_Star\_deg\_low} \end{cases}$$

733   (6.3)

734   in which K_{m10} is Michaelis-constant, $[Suc]_{leaf\_Star\_deg\_low}$ is critical leaf sucrose

735 concentration below which starch starts to decompose, $v_{m\_leaf\_Star\_deg}$ is leaf maximal

736 starch degradation rate (Supplementary Table 1).

737 Leaf O-N and protein are in dynamic balance by their inter-conversion (conversion rate

738 $v_{leaf\_O\text{-}N2Pro} > 0$ means a conversion from O-N to protein, and *vice versa*):

$$v_{leaf\_O\text{-}N2Pro} = \begin{cases} v_{m\_leaf\_Pro\_syn} \cdot \dfrac{[\text{O-N}]_{leaf} - [\text{O-N}]_{leaf\_Pro\_syn\_low}}{K_{m11} + [\text{O-N}]_{leaf}}, [\text{O-N}]_{leaf} > [\text{O-N}]_{leaf\_Pro\_syn\_low} \\ -v_{m\_leaf\_Pro\_deg} \cdot \dfrac{[\text{O-N}]_{leaf\_Pro\_syn\_low} - [\text{O-N}]_{leaf}}{K_{m11} + [\text{O-N}]_{leaf}}, [\text{O-N}]_{leaf} \leq [\text{O-N}]_{leaf\_Pro\_syn\_low} \end{cases} \tag{7.1}$$

in which K_{m11} is Michaelis-constant, $[\text{O-N}]_{leaf\_Pro\_syn\_low}$ is critical leaf O-N

concentration below which protein synthesis ceases and starts to decompose,

$v_{m\_leaf\_Pro\_syn}$ is leaf maximal protein synthesis rate and $v_{m\_leaf\_Pro\_deg}$ is leaf maximal

protein degradation rate (Supplementary Table 1).

Leaf senescence: Similar to that of root, there is a minimal (constant) relative leaf

photosynthetic area (S_{leaf}) loss rate α_{leaf_sene} (Supplementary Table 1) due to aging [117].

As leaf senescence is closely related to leaf N remobilization during grain filling [118,119],

we proposed that leaf senescence rate (v_{leaf_sene}) would be accelerated as a result of

nitrogen starvation when leaf total nitrogen level ($[\text{N}]_{leaf}$; including I-N, O-N and

protein) decreased below a critical level $[\text{N}]_{sene\_up}$:

$$v_{leaf\_sene} = \begin{cases} \alpha_{leaf\_sene} \cdot e^{-\beta_{leaf\_sene} \cdot ([\text{N}]_{leaf} - [\text{N}]_{sene\_up})}, [\text{N}]_{leaf} < [\text{N}]_{sene\_up} \\ \alpha_{leaf\_sene}, [\text{N}]_{leaf} \geq [\text{N}]_{sene\_up} \end{cases} \tag{8.1}$$

in which β_{leaf_sene} is an empirical coefficient (Supplementary Table 1).

Grains

In WACNI, grain volume expansion and grain filling occur simultaneously rather than

being divided into two distinct phases, as it has been reported in both maize and rice

that time for expression of enzymes envolved in these two processes is overlaped, and

protein/starch granuals are presented in endosperm cells at the very beginning of the

grain filling period [120-124].

Grain volume growth: Grain volume growth, i.e., grain cell division, was modelled

based on current grain surface area since cell division occurs mainly within several

outer layer cells of grain [125]. Assuming that developing grains has an ellipsoid shape

with the width, height and length being w, w and $w \cdot r$ (r is grain length width ratio,

Supplementary Table 1), respectively, grain volume (V_{grain}) and grain surface area (S_{grain})

were estimated as following:

764
$$V_{grain} = \frac{4}{3}\pi \cdot w^3 \cdot r \quad (9.1)$$

765
$$S_{grain} \approx 4\pi \left(\frac{1 + 2r^{1.6075}}{3}\right)^{\frac{1}{1.6075}} \cdot w^2 \quad (9.2)$$

766   Glume size (V_{glume}) and/or and pericarp size can physically restrict expansion of

767   developing grain [126], although the strength of constraint may differ among species [127].

768   This constraint effect ($\alpha_{restrict}$) was modelled as following:

769
$$\alpha_{restrict} = \begin{cases} 1, V_{grain} < (1-\varepsilon) \cdot V_{glume\_max} \\ \dfrac{1}{\varepsilon}\left(1 - \dfrac{V_{grain}}{V_{glume\_max}}\right), V_{grain} \geq (1-\varepsilon) \cdot V_{glume\_max} \end{cases} \quad (9.3)$$

in which $(1-\varepsilon) \cdot V_{glume}$ is a critical grain volume above which grain expansion will be

more and more physically restrained.

In addition, grain volume expansion was modelled based on current grain surface area

(S_{grain}) since cell division occurs mainly within several outer layer cells of grain [125]. We

further proposed that grain volume expansion rate (v_{grain_grow}) was co-determined by

carbohydrate and nitrogen supply, i.e., concentration of sucrose and nitrogen in grain

aqueous space ($[\text{O-N}]_{grain}$ and $[\text{Suc}]_{grain}$, respectively) in WACNI:

777
$$v_{grain\_grow} = \alpha_{restrict} \cdot v_{m\_grain\_grow} \cdot S_{grain} \frac{[\text{O-N}]_{grain}}{K_{m12} + [\text{O-N}]_{grain}} \cdot \frac{[\text{Suc}]_{grain}}{K_{m13} + [\text{Suc}]_{grain}} \quad (9.4)$$

778   in which K_{m12} and K_{m13} are Michaelis-constants, $v_{m\_grain\_grow}$ is grain maximal growth

779   rate (Supplementary Table 1).

780   *Grain starch and protein storage*: The aqueous space volume (V_{grain_aq}) expands with

781   endosperm cells division and reduces by storage starch and protein (which are insoluble)

782   cumulation:

783
$$V_{grain\_aq} = V_{grain} - \frac{m_{grain\_Star}}{\rho_{Star}} - \frac{m_{grain\_Pro}}{\rho_{Pro}} \quad (10.1)$$

784   in which V_{grain} is total grain volume, $m_{\text{grain\_Star}}$ and $m_{\text{grain\_Pro}}$ are biomass and ρ_{Star} and

785   ρ_{Pro} are density of cumulated starch and protein in grain, respectively.

786   For grain filling, storage starch and protein synthesis rate ($v_{\text{grain\_Star\_syn}}$ and $v_{\text{grain\_Pro\_syn}}$)

787   were modeled based on sucrose and O-N concentrations in grain aqueous space

788   following a Michaelis-Menten equation:

789
$$v_{\text{grain\_Star\_syn}} = v_{\text{m\_grain\_Star\_syn}} \cdot \frac{[\text{Suc}]_{\text{grain}}}{K_{\text{m14}} + [\text{Suc}]_{\text{grain}}} \quad (10.2)$$

790
$$v_{\text{grain\_Pro\_syn}} = v_{\text{m\_grain\_Pro\_syn}} \cdot \frac{[\text{O-N}]_{\text{grain}}}{K_{\text{m15}} + [\text{O-N}]_{\text{grain}}} \quad (10.3)$$

in which K_{m14} and K_{m15} are Michaelis-constants, $v_{\text{m\_grain\_Star\_syn}}$ is grain maximal starch

synthesis rate, $v_{\text{m\_grain\_Pro\_syn}}$ is grain maximal protein synthesis rate (Supplementary

Table 1).

Stem

Stem sucrose and starch, O-N and protein homeostasis: Sucrose and starch, O-N and

protein are in dynamic balance in stem (including culm and sheath) as a result of inter-

conversion between them, respectively. The conversion rates between them

($v_{\text{stem\_Suc2Star}}$ and $v_{\text{stem\_O-N2Pro}}$) were modeled as following:

799
$$V_{\text{SSP\_aq}} = \text{SV0} - \frac{m_{\text{SSP\_Star}}}{\rho_{\text{Star}}} - \frac{m_{\text{SSP\_Pro}}}{\rho_{\text{Pro}}} \quad (11.1)$$

801
$$v_{\text{SSP\_Star\_syn}} = \begin{cases} v_{\text{m\_SSP\_Star\_syn}} \cdot \dfrac{[\text{Suc}]_{\text{SSP}} - [\text{Suc}]_{\text{SSP\_Star\_syn\_low}}}{K_{\text{m16}} + [\text{Suc}]_{\text{SSP}} - [\text{Suc}]_{\text{SSP\_Star\_syn\_low}}}, [\text{Suc}]_{\text{SSP}} > [\text{Suc}]_{\text{SSP\_Star\_syn\_low}} \\[2ex] v_{\text{m\_SSP\_Star\_deg}} \cdot \dfrac{[\text{Star}]_{\text{SSP}}}{K_{\text{m17}} + [\text{Suc}]_{\text{SSP}}} \cdot \dfrac{[\text{Suc}]_{\text{SSP}} - [\text{Suc}]_{\text{SSP\_Star\_syn\_low}}}{K_{\text{m16}} + [\text{Suc}]_{\text{SSP}} - [\text{Suc}]_{\text{SSP\_Star\_syn\_low}}}, [\text{Suc}]_{\text{SSP}} \leq [\text{Suc}]_{\text{SSP\_Star\_syn\_low}} \end{cases}$$

802   (11.2)

804
$$v_{\text{SSP\_O-N2Pro}} = \begin{cases} v_{\text{m\_SSP\_Pro\_syn}} \cdot \dfrac{[\text{O-N}]_{\text{SSP}} - [\text{O-N}]_{\text{SSP\_Pro\_syn\_low}}}{K_{\text{m18}} + [\text{O-N}]_{\text{SSP}} - [\text{O-N}]_{\text{SSP\_Pro\_syn\_low}}}, [\text{O-N}]_{\text{SSP}} > [\text{O-N}]_{\text{SSP\_Pro\_syn\_low}} \\[2ex] -v_{\text{m\_SSP\_Pro\_deg}} \cdot \dfrac{[\text{Pro}]_{\text{SSP}}}{K_{\text{m19}} + [\text{Pro}]_{\text{SSP}}} \cdot \dfrac{[\text{O-N}]_{\text{SSP}} - [\text{O-N}]_{\text{SSP\_Pro\_syn\_low}}}{K_{\text{m18}} + [\text{O-N}]_{\text{SSP}} - [\text{O-N}]_{\text{SSP\_Pro\_syn\_low}}}, [\text{O-N}]_{\text{SSP}} \leq [\text{O-N}]_{\text{SSP\_Pro\_syn\_low}} \end{cases}$$

805   (11.3)

in which V_{stem_aq} is aqueous space volume of stem; m_{stem_Star} and m_{stem_Pro} are biomass

of starch and protein in stem, respectively; K_{m16}, K_{m17} K_{m18} and K_{m19} are Michaelis-

constants; $v_{m\_stem\_Star\_syn}$ and $v_{m\_stem\_Star\_deg}$ are stem maximal starch synthesis and starch

degradation rate, $v_{m\_stem\_Pro\_syn}$ and $v_{m\_stem\_Pro\_deg}$ are stem maximal protein synthesis and

protein degradation rate, respectively (Supplementary Table 1).

Xylem and phloem transport: Phloem transport includes short distance transport that

occurs at source or sink ends, i.e., material loading or unloading, and long distance

transport along phloem sieve tubes. According to Münch's theory, phloem long distance

transport resistance is mainly determined by the physical property of the sieve tube and

fluid in it. However, transporter proteins, whose efficiency is mainly determined by

Michaelis-Menten kinetics, usually control the resistance of short distance transport. As

solutes in xylem move with transpiration flow, which is not explicitly considered in our

current model, solutes were assumed to be uniformly distributed along xylem path

without concentration gradient. For phloem, we divided it into three different

compartments and considered the transport resistance between them (see below).

Flux between different phloem compartments: Phloem path was divided into three

compartments (see Figure 1), i.e., leaf phloem, grain phloem and root phloem. Flux in

phloem is driven by osmotic pressure difference between source and sink according to

Münch theory. Sucrose and amino acids are the main osmotic components in phloem

[128]. Flux rate ($v_{phloem\_flux\_leaf\_X}$) between phloem compartments of leaf and organ X (X

can be grain or root) was modeled as following:

$$v_{phloem\_flux\_leaf\_X} = \frac{([Suc]_{leaf\_phloem} + [O\text{-}N]_{leaf\_phloem}) - ([Suc]_{X\_phloem} + [O\text{-}N]_{X\_phloem})}{R_{phloem\_leaf\_X}} \quad (12.1)$$

in which $R_{phloem\_leaf\_X}$ is transport resistance between leaf phloem and X phloem

(Supplementary Table 1).

Loading and unloading: Rate of substrate *s* loading and unloading at each source or

sink organ X ($v_{X\_s\_l}$ and $v_{X\_s\_ul}$) were modeled with a simple equation which describes

832   metabolites transport across membranes, similar to that in Wang, et al. [35]:

$$v_{\text{X\_s\_l}} = v_{\text{m\_X\_s\_l}} \cdot \frac{[s]_{\text{X}} - \dfrac{[s]_{\text{X}}}{K_{\text{e\_X\_s\_l}}}}{K_{\text{m\_X\_s\_l}} + [s]_{\text{X}}} \quad (13.1)$$

$$v_{\text{X\_s\_ul}} = v_{\text{m\_X\_s\_ul}} \cdot \frac{[s]_{\text{tube}} - \dfrac{[s]_{\text{X}}}{K_{\text{e\_X\_s\_ul}}}}{K_{\text{m\_X\_s\_ul}} + [s]_{\text{tube}}} \quad (13.2)$$

in which $K_{\text{m\_X\_s\_l}}$ and $K_{\text{m\_X\_s\_ul}}$ are Michaelis-constants; $v_{\text{m\_X\_s\_l}}$ is maximal loading rate

and $v_{\text{m\_X\_s\_ul}}$ is maximal unloading rate of substrate s at organ X; $[s]_{\text{X}}$ and $[s]_{\text{tube}}$ are the

concentrations of substrate s in organ X and in xylem or phloem tube, respectively.

These loading and unloading processes include root-to-shoot I-N and O-N loading (to

xylem), shoot-to-root sucrose and O-N unloading (from phloem) at root; photosynthetic

sucrose and O-N loading (to phloem), I-N and O-N unloading (from xylem) at leaf;

sucrose and O-N unloading (from phloem) at grain; and xylem-to-phloem O-N transfer

(see Figure 1, process 13).

Phloem-stem diffusion: As a simplification, stem was set to locate around leaf phloem

only. Substrates (sucrose and O-N in this model) difffusion rate ($v_{\text{phloem\_stem\_s\_d}}$) between

leaf phloem and stem were determined by their diffusion property:

$$v_{\text{phloem\_SSP\_s\_d}} = \frac{[s]_{\text{phloem}} - [s]_{\text{SSP}}}{R_{\text{\_phloem\_SSP\_s}}} \quad (14.1)$$

847   in which $[s]_{\text{phloem}}$ and $[s]_{\text{stem}}$ are substrate s concentration in leaf phloem and stem,

848   respectively; $R_{\text{\_phloem\_stem\_s}}$ is transfer resistance of substrate s (Supplementary Table 1).

849   ***Respiration***

850   Plant dark respiration rate was calculated for each organ ($R_{\text{d\_root}}$, $R_{\text{d\_leaf}}$, $R_{\text{d\_grain}}$, $R_{\text{d\_stem}}$),

851   following the concept proposed by Cannell and Thornley [129]:

$$R_{\text{d\_root}} = [\eta_{\text{l}} \cdot (v_{\text{root\_IN\_l}} + v_{\text{root\_ON\_l}} + v_{\text{root\_ON\_ul}} + v_{\text{root\_Suc\_ul}}) + \eta_{\text{N\_upt}} \cdot v_{\text{root\_N\_upt}}$$
$$+ \eta_{\text{N\_ass}} \cdot v_{\text{root\_N\_ass}} + \eta_{\text{grow}} \cdot v_{\text{root\_grow}}] \cdot \text{RW} + \gamma_{\text{root\_residual}} \cdot V_{\text{root\_aq}} \quad (15.1)$$

$$R_{\text{d\_leaf}} = [\eta_{\text{l}} \cdot (v_{\text{leaf\_ON\_l}} + v_{\text{leaf\_Suc\_l}} + v_{\text{leaf\_IN\_ul}} + v_{\text{leaf\_ON\_ul}}) + \eta_{\text{store}} \cdot v_{\text{leaf\_Star\_deg}}$$
$$+ \eta_{\text{Pro\_syn}} \cdot v_{\text{leaf\_Pro\_syn}} + \eta_{\text{Pro\_deg}} \cdot v_{\text{leaf\_Pro\_deg}}] \cdot \text{LA} + \gamma_{\text{residual}} \cdot V_{\text{leaf\_aq}} \quad (15.2)$$

$$R_{\text{d\_grain}} = [\eta_{\text{ul}} \cdot (v_{\text{grain\_ON\_ul}} + v_{\text{grain\_Suc\_ul}}) + \eta_{\text{grow}} \cdot v_{\text{grain\_grow}}] \cdot S_{\text{grain}} + (\eta_{\text{store}} \cdot v_{\text{grain\_Star\_syn}}$$
$$+ \eta_{\text{Pro\_syn}} \cdot v_{\text{grain\_Pro\_syn}}) \cdot V_{\text{grain\_aq}} + \gamma_{\text{residual}} \cdot V_{\text{grain}} \tag{15.3}$$

$$R_{\text{d\_stem}} = [\eta_{\text{store}} \cdot (v_{\text{stem\_Star\_deg}} + v_{\text{stem\_Star\_syn}}) + \eta_{\text{Pro\_syn}} \cdot v_{\text{stem\_Pro\_syn}} + \eta_{\text{Pro\_deg}} \cdot v_{\text{stem\_Pro\_deg}}$$
$$+ \gamma_{\text{residual}}] \cdot V_{\text{stem\_aq}} \tag{15.4}$$

in which η_{l}, η_{ul}, $\eta_{\text{N\_upt}}$, $\eta_{\text{N\_ass}}$, η_{grow}, η_{store}, $\eta_{\text{Pro\_deg}}$ and $\eta_{\text{Pro\_syn}}$ are respiration coefficients of loading, unloading, soil I-N uptake, N assimilation, growth, starch degdaration or synthesis, protein degradation and protein synthesis processes, respectively; $v_{\text{X\_IN\_l}}$ and $v_{\text{X\_ON\_l}}$ and $v_{\text{X\_Suc\_l}}$, $v_{\text{X\_IN\_ul}}$ and $v_{\text{X\_ON\_ul}}$ and $v_{\text{X\_Suc\_ul}}$, $v_{\text{X\_N\_upt}}$, $v_{\text{X\_N\_ass}}$, $v_{\text{X\_grow}}$, $v_{\text{X\_Star\_deg}}$ and $v_{\text{X\_Star\_syn}}$, $v_{\text{X\_Pro\_deg}}$ and $v_{\text{X\_Pro\_syn}}$ are the corresponding reaction rates in organ X; $V_{\text{root\_aq}}$, $V_{\text{leaf\_aq}}$, $V_{\text{grain\_aq}}$ and $V_{\text{stem\_aq}}$ are the aqueous space volume of root, leaf, grain and stem, respectively; γ_{residual} is the residual respiration (respiration for other processes) coefficient for leaf, stem and grain; $\gamma_{\text{root\_residual}}$ is the residual respiration coefficient for root, which is twice larger than γ_{residual} as a compensation for potential root exudation that is not explicitly represented in the model (Supplementary Table 1).

Model parameterization

The maximal velocities (v_{max}) for processes modeled in WACNI were derived from a vast amount of genetic, molecular, biochemical and plant physiological research in rice; while most of the apparent Michaelis-constants (K_{m}) and equilibrium constants (K_{e}) were estimated based on initial metabolite concentrations in each organ. A detailed description of parameters, their values, and the comprehensive parameterization procedure used to derive or estimate their values are given in Supplementary Table 1.

Model validation on multiple datasets

Simulating typical plant physiological changes during rice grain filling period (Figure 2a-f)

WACNI was firstly validated by comparing the relative changes of physiological traits

in different organs during the grain filling period between model predictions with

default parameter settings and experimental data obtained from literatures under normal

conditions (Figure 2a-f). Specifically, data of root weight dynamic change (Figure 2a)

were extracted from *Table 2* in Tang, et al. [50], and the average values of seven

combinations of two-line hybrid rice were used; data of leaf area dynamic change

(Figure 2b) were extracted from *Figure 1(d)* in Haque, et al. [51], and the average values

of three hybrid and inbred varieties in 2008-09 were used; data of canopy

photosynthesis (Figure 2c) were extracted from *Fig. b* in Zhao, et al. [52], and values of

treatment F3 were used; data of grain volume (Figure 2d) and grain dry weight (Figure

2e) were extracted from *Fig. 1c* and *Fig. 2c* in Yang, et al. [53], respectively, and values

of spikelets of cultivar Yangdao 4 that flowered on the fourth day were used; data of

stem non-structural carbohydrate (Figure 2f) were extracted from *Fig. 4B* in Yang, et

al. [54], and the average values of normal nitrogen with well-watered and water deficit

stressed treatments of cultivar Yangdao 4 were used. All the extracted experimental

data are tabulated in Supplementary Datasets 2.

Simulating plant carbon and nitrogen fluxes during rice grain filling period (Figure

2g)

Specifically, total carbon content in a plant at flowering (PTC0) was calculated as:

$$
\begin{aligned}
\text{PTC0}=&\text{RW0+SSW0+LA0}\,/\,\text{SLA+SStar0}+\text{SPro0+LPro0}+\\
&\text{RV0}\cdot(c_{\text{root\_ON}}+c_{\text{root\_IN}}+c_{\text{root\_Suc}})+\text{SV0}\cdot(c_{\text{stem\_ON}}+c_{\text{stem\_Suc}})+\\
&\text{LV0}\cdot(c_{\text{leaf\_ON}}+c_{\text{leaf\_IN}}+c_{\text{leaf\_Suc}})+\text{PV0}\cdot(c_{\text{phloem\_ON}}+c_{\text{phloem\_Suc}})+\\
&\text{XV0}\cdot(c_{\text{xylem\_ON}}+c_{\text{xylem\_IN}})
\end{aligned}
$$

in which RW0, SSW0, LA0, SLA, SStar0, SPro0 and LPro0 are root weight, stem

structural weight, leaf area, specific leaf area, stem stored starch weight, stem stored

protein weight and leaf protein weight at flowering, respectively; RV0, SV0, LV0, PV0

and XV0 are root volume, stem volume, leaf volume, phloem volume and xylem

volume, respectively; $c_{\text{x\_y}}$ are concentration of metabolite y in organ x.

Total nitrogen content in a plant at flowering (PTN0) was calculated as:

$$PTN0 = RW0 \cdot c_{root\_strucN} + SSW0 \cdot c_{stem\_strucN} + LA0 / SLA \cdot c_{leaf\_strucN} +$$
$$RV0 \cdot (c_{root\_ON} + c_{root\_IN}) + SV0 \cdot c_{stem\_ON} + LV0 \cdot (c_{leaf\_ON} + c_{leaf\_IN}) +$$
$$PV0 \cdot c_{phloem\_ON} + XV0 \cdot (c_{xylem\_ON} + c_{xylem\_IN}) +$$
$$SPro0 + LPro0$$

For default model parameters, PTN0=49 mmol/plant.

For the carbon budget, we calculated several types of carbon fluxes. They are:

● leaf respiration to total photosynthetically fixed carbon = 504/2328*100% =
21.6%;

● entire plant respiration to total photosynthetically fixed carbon =
[420+504+84+204]/2328*100% = 52.1%;

● contribution of photosynthesis-derived carbon to grain filling =
1764/[1764+240]*100% = 88.0%;

● contribution of stem non-structural carbohydrate storage to grain filling =
240/[1764+240]*100% = 12.0%;

● ratio between stem weight loss during grain filling to grain yield at harvest =
[240+84]/[1608-420]*100% = 27.3%;

● ratio of carbohydrates partitioning to root from flowering to harvest =
396/[396+1608]*100% = 19.8%;

● ratio of carbohydrates partitioning to ear from flowering to harvest = 100% -
19.8% = 80.2%.

For nitrogen budget, we calculated several types of nitrogen fluxes. They are:

● ratio of nitrogen assimilation in root = 7.8/[7.8+12.4]*100% = 38.6%;

● ratio of nitrogen assimilation in leaf = 100% - 38.6% = 61.4%;

● ratio of root nitrogen uptake after flowering to whole season nitrogen uptake =

20/[20+49]*100% = 29.0%;

● nitrogen harvest index = 43.3/[20+49]*100% = 62.8%.

Simulating effects of different nitrogen fertilizer application rates on rice grain filling

(Figure 3a-l)

To examine effects of different nitrogen fertilizer application rates on rice grain filling,

experimental data were collected from Zhao, et al. [57] (Supplementary Datasets 2). The

authors reported four nitrogen topdressing treatments during the ear development

period, i.e., no nitrogen was applied; low level (0.6 g nitrogen/pot), medium level (1.2

g nitrogen/pot), and high level (1.8 g nitrogen/pot) nitrogen were applied. In simulation,

soil nitrogen concentrations were set to be 20%, 50%, 100% and 150% of the default

value for the four nitrogen treatments, respectively, with the assumption that there was

still 20% of nitrogen in soil when no nitrogen was applied. According to *Table 1* in

Zhao, et al. [57], ratio of leaf dry matter and grains dry weight per plant among the four

nitrogen treatments at 7 days after anthesis were 0.60 : 0.90 : 1.00 : 1.23, and 0.43 :

0.75 : 1.00 : 1.18, respectively. Therefore, we set ratio of tiller number between these

groups as that of leaf dry matter, i.e., 0.60 : 0.90 : 1.00 : 1.23, and set ratio of grain

number per ear between these groups by dividing grains dry weight by tiller number,

i.e., 0.43/0.60 : 0.75/0.90 : 1.00 : 1.18/1.23. According to *Table 2* of Zhao, et al. [57], ratio

of leaf protein amino acids per plant between the four nitrogen treatments at 7 days

after anthesis were 0.40 : 0.75 : 1.00 : 1.21. Therefore, we set ratio of leaf protein

concentration between groups by dividing leaf protein amino acids by leaf dry matter,

i.e., 0.40/0.60 : 0.75/0.90 : 1.00 : 1.21/1.23. We further set the same ratio of stem protein

concentration between groups as that of leaf. In addition, tiller number per plant for the

medium nitrogen treatment was set to 11.

For the sake of comparing physiological traits during the grain filling period between

model simulation and experimental data, grain nitrogen concentration data were

extracted from *Table 2* in Zhao, et al. [57], and values of total amino acids were used; leaf

nitrogen concentration data were calculated by dividing amount of leaf total amino

acids by leaf dry matter from *Table 1* and *Table 2* in Zhao, et al. [57], grain dry weight

data were extracted from *Table 1* in Zhao, et al. [57]. For both model simulation and

experimental data, values of leaf nitrogen concentration, grain nitrogen concentration

at 7 days after flowering and grain dry weight at harvest of the medium nitrogen

treatment were used as a reference, and data at all time points and in all nitrogen

treatment groups were normalized by dividing these values, respectively. Extracted

experimental data are tabulated in Supplementary Datasets 2.

Simulating effects of shading and thinning on rice grain filling (Figure 3m-s)

To examine effects of different light regimes on rice grain filling, experimental data

were collected from Kobata, et al. [58] (Supplementary Datasets 2). The authors reported

seven different shading/thinning treatments after flowering. Firstly, control plots were

grown under normal condition throughout the entire grain filling period. Concurrently,

three shading treatments were applied immediately after flowering, i.e., the heavy

shaded treatment using double black cloth, the moderate shaded treatment using single

black cloth, and the light shaded treatment using white cloth, which reduced full sun

radiation by 74.4%, 48.1% and 25.1%, respectively. The shade frames were removed

10 days later. Then, half the plots were left untouched (Figure 3n-p), while the other

half were thinned to every other plant to reduce the plant density by half for the rest of

the grain filling period (Figure 3q-s).

Default model parameters were used during simulation. In WACNI, incident solar light

intensity during the firstly 10 days after flowering were reduced to 75%, 50% and 25%

for light, medium and heavy shaded treatments, respectively; and the planting density

beyond 10 days after flowering were halved for the thinning treatment groups.

Experimental data of grain dry weight changes during the grain filling period for all

seven treatments were extracted from *Fig. 2* in Kobata, et al. [58]. For both model

simulation and experimental data, values of grain dry weight at harvest of the 'normal

condition' group were used as a reference, and data at all time points and for all

treatments were normalized by dividing them, respectively. Extracted experimental

data are tabulated in Supplementary Datasets 2.

Simulating interaction effects of different nitrogen fertilizer application rates and air

CO_2 concentrations on rice grain filling (Figure 3t-u)

For simulating interaction effects of different nitrogen fertilizer application rates and

air CO_2 concentrations on rice grain filling, experimental data were collected from Kim,

et al. [59] (Supplementary Datasets 2). The authors reported six different combinations of

nitrogen fertilizer application rate and air CO_2 concentration treatments. Namely,

nitrogen was supplied as ammonium sulphate at three application rates: 4 g m^{-2} (low;

'LN'), 8-9 g m^{-2} (medium; 'MN') and 12-15 g m^{-2} (high; 'HN'), respectively; air CO_2

concentration was controlled at two levels: ambient level of around 390 μmol mol^{-1},

free-air CO_2 enrichment of 230-365 μmol mol^{-1} higher than that under ambient level

('FACE'). Correspondingly, in the simulation, we set the soil nitrogen concentrations

to be 50%, 100% and 150% of the default value for 'LN', 'MN' and 'HN' treatments,

respectively; the air CO_2 concentrations were set to 390 and 690 μmol mol^{-1} for ambient

and 'FACE' treatments, respectively.

According to *Table 1* in Kim, et al. [59], it was known that ratio of plant total nitrogen,

plant root dry matter and plant total dry matter between 'LN', 'MN', 'HN', 'LN+FACE',

'MN+FACE' and 'HN+FACE' groups at ear initiation were 0.83 : 1.00 : 1.25 : 0.85 :

1.15 : 1.53, 1.00 : 1.00 : 1.03 : 1.18 : 1.46 : 1.35 and 0.87 : 1.00 : 1.06 : 1.21 : 1.45 :

1.51, respectively. Therefore, ratio of root weight between these groups at flowering

was set as that of plant root dry matter measured at ear initiation; ratio of leaf area and

pre-flowering stem stored starch at flowering was set to be the same as that of measured

plant total dry matter at ear initiation; and ratio of leaf protein concentration and pre-

flowering stem stored protein at flowering was set by dividing plant total nitrogen by

plant total dry matter, i.e., 0.83/0.87 : 1.00 : 1.25/1.06 : 0.85/1.21 : 1.15/1.45 : 1.53/1.51.

According to *Table 2* in Kim, et al. [59], it was known that ratio of ear number, fertile

spikelets and individual grain mass between 'LN', 'MN', 'HN', 'LN+FACE',

'MN+FACE' and 'HN+FACE' groups at harvest were 0.87 : 1.00 : 1.07 : 0.89 : 1.09 :

1.15, 0.83 : 1.00 : 1.13 : 0.85 : 1.09 : 1.30, and 1.07 : 1.00 : 0.98 : 1.09 : 1.04 : 0.93,

respectively. Therefore, ratio of tiller number and grain size between these groups was

set as that of ear number and individual grain mass measured at harvest, respectively;

ratio of grain number per ear between these groups was set by dividing fertile spikelets

by tiller number, i.e., 0.83/0.87 : 1.00 : 1.13/1.07 : 0.85/0.89 : 1.09/1.09 : 1.30/1.15.

Ratio of leaf nitrogen concentration at flowering was set by dividing plant nitrogen

uptake by plant total dry matter at panicle initiation, which was 0.95 : 1.00 : 1.18 : 0.70 :

0.79 : 1.01. In addition, grain number per ear was set to 145 for the 'MN' group.

During comparison of model prediction and experimental data, grain yield at harvest

was extracted from *Table 2* in Kim, et al. [59], root nitrogen uptake throughout the grain

filling period was approximated by the difference of plant total nitrogen at harvest and

plant total nitrogen at ear initiation from *Table 1* in Kim, et al. [59]. We simulated two

scenarios, one was using adjusted parameters as above mentioned, with other

parameters using the default values (the 'Simulation' group in Figure 3t, u); the other

one was with a further assumption that maximal root nitrogen uptake rate under 'FACE'

was reduced by 36% compared to that under ambient condition (the 'Simulation

(v_{m_Nupt}=64% under FACE)' group in Figure 3t, u). For both model simulation and

experimental data, values of grain yield and root nitrogen uptake in the 'MN' group

were used as a reference, and data at all time points and for all treatments were

normalized by dividing them, respectively. Extracted experimental data are tabulated

in Supplementary Datasets 2.

Simulating effects of genetic manipulation of leaf senescence rate on rice grain

filling (Figure 4a-c)

To examine effects of genetic manipulation of leaf senescence rate on rice grain filling,

experimental data were collected from Liang, et al. [61] (Supplementary Datasets 2). The

authors reported a gain-of-function mutant prematurely senile 1 (*ps1-D*) that

demonstrated significant premature leaf senescence. Further genetic study shows that

PS1 encodes a plant-specific NAC transcription activator, *OsNAP*, overexpression of

which significantly promoted senescence, whereas knockdown of which would delay

leaf senescence. Default model parameters were used for simulating wild type plants

('WT'), except that the $v_{m\_leaf\_Pro\_deg}$ was set to 50% of the default value. According to

Fig. 1C in Liang, et al. [61], expression of chlorophyll degradation and leaf senescence

process related genes in '*ps1-D* mutant' increased to 3-5 folds to that of 'WT'; and

according to *Fig. 4B-C* in Liang, et al. [61], *OsNAP* gene expression level in leaf of

'*OsNAP* RNAi' decreased to 0.2-0.3 fold, while ABA content increased to 1.4-1.9 folds

to that of 'WT'. Therefore, we set $v_{m\_leaf\_Pro\_deg}$ of '*ps1-D* mutant' and '*OsNAP* RNAi'

to be 0.5-fold and 4-folds of 'WT', respectively.

According to *Table S4* in Liang, et al. [61], ratio of dry weight of vegetative organs among

'WT', '*ps1-D* mutant' and '*OsNAP* RNAi' groups was 1 : 0.74 : 1.11. Therefore, ratio

of plant size between these groups, including leaf area, root weight, stem stored

starch/protein and grain number per ear, were set the same as that for dry weight of

vegetative organs, i.e., 1 : 0.74 : 1.11.

To enable comparison between model predictions and experimental data, we extracted

leaf chrolophyll content dynamic changes of 'WT' and '*ps1-D* mutant' during the grain

filling period from *Fig. 1B* in Liang, et al. [61], which was used to represent relative

change of leaf nitrogen concentration. Grain yield and grain nitrogen concentration at

harvest were extracted from *Fig. 4F* and *Fig. 5B* in Liang, et al. [61], respectively. For

both model simulation and experimental data, values of leaf nitrogen concentration at

flowering, and grain yield and grain nitrogen concentration at harvest of 'WT' were

used as a reference, i.e., data at all time points and for other plants were normalized by

dividing them, respectively. Extracted experimental data are tabulated in

Supplementary Datasets 2.

Simulating effects of genetic manipulation of grain sucrose unloading capacity on

rice grain filling (Figure 4d)

To examine effects of genetic manipulation of grain sucrose unloading capacity on rice

grain filling, experimental data were collected from Wang, et al. [62] (Supplementary

Datasets 2). The authors reported a loss-of-function mutant grain incomplete filling 1

('*gif1* mutant') that showed slower grain-filling rate than wild-type rice ('WT'). Further

genetic study shows that *GIF1* encodes a cell-wall invertase, which is responsible for

sucrose unloading into grain, particularly during early grain-filling. As shown in Figure

1g of Wang, et al. [62], the grain filling duration was as short as less than 30 days, we

halved the leaf area, root weight and stem stored starch/protein from default values, set

grain number per ear to 90, and set the maximal rate of sucrose unloading to grain as

200% of the default value for 'WT'.

According to *Figure 1h-j* in Wang, et al. [62], sugar content at 5 days after pollination in

'*gif1* mutant' decreased to 0.4-0.7 fold to that of 'WT'. Therefore, the maximal grain

sucrose unloading rate of '*gif1* mutant' was set to be 0.5-fold of 'WT'; the maximal

grain sucrose unloading rate of plants overexpressing *GIF1* from its native promoter

('*GIF1* OE') was set to be 2-fold of 'WT'.

According to *Figure 1g* and *Figure 4b* in Wang, et al. [62], ratio of grain dry weight at

harvest between 'WT', '*gif1* mutant' and '*GIF1* OE' groups was 1 : 0.76 : 1.11.

Therefore, ratio of grain size between these groups was set to be the same as that for

measured grain dry weight at harvest, i.e., 1 : 0.76 : 1.11. Extracted experimental data

are tabulated in Supplementary Datasets 2.

For both model simulation and experimental data, grain dry weights at harvest of 'WT'

were used as a reference, i.e., data at all time points and for other plants were normalized

by dividing them, respectively.

To test the effects of changing grain sucrose unloading capacity on plant physiological

changes during the grain filling period for 'source limited' plants (with default model

parameter settings except that the grain number per ear is set to 190), we further

simulated grain dry weight, leaf nitroge concentration, canopy photosynthesis, root dry

weight and root nitrogen uptake throughout the grain filling period for plants with

maximal grain sucrose unloading rates of 200%, 100% and 50% to that of the default

value.

Simulating effects of genetic manipulation of grain starch synthesis capacity on rice

grain filling (Figure 4e)

To examine effects of genetic manipulation of grain starch synthesis capacity on rice

grain filling, experimental data were collected from Smidansky, et al. [63] (Supplementary

Datasets 2). The authors reported that transforming rice with a modified maize AGPase

large subunit sequence (*Sh2r6hs*) in an endosperm-specific manner could specifically

enhance capacity for starch synthesis in endosperm. According to *Table 4* in Smidansky,

et al. [63], the grain number per ear was only 90 for negative homozygotes of *Sh2r6hs*

transgene ('WT') plants. Therefore, the leaf area, root weight and stem stored

starch/protein were halved from default values, and grain number per ear was set to 90

for 'WT' plants. According to *Table 4* in Smidansky, et al. [63], ratio of grain number per

ear and tiller number between 'WT' and *Sh2r6hs* transgene groups were 1 : 1.13 and 1 :

1.05, respectively. Ratio of grain size between two groups was set to be the same as that

for grain weight, i.e., 1: 1.02. Ratio of leaf area, root weight, stem structural weight,

stem starch storage and stem protein storage was set by dividing total plant weight by

tiller number, i.e., 1 : 1.22/1.05. According to *Table 2* in Smidansky, et al. [63], the

measured AGPase activity was 9-52% higher in *Sh2r6hs* transgenic grains than in

untransformed grains between 10-20 days after anthesis. Therefore, the maximal grain

starch synthesis rate $v_{m\_grain\_Star\_syn}$ of *Sh2r6hs* transgenic plants was set to be 1.5-fold

of 'WT'.

In total three scenarios were simulated, the first one was using adjusted plant size and $v_{m\_grain\_Star\_syn}$ as above mentioned, with other parameters using the default values (the grey bars in Figure 4e); the second one was using only adjusted plant size with unchanged $v_{m\_grain\_Star\_syn}$ (the black bars Figure 4e); the third one was using only adjusted $v_{m\_grain\_Star\_syn}$ with unchanged plant size (the white bars Figure 4e).

To enable comparison between model predictions and experimental data, we extracted grain yield, grain nitrogen concentration and harvest index at harvest of positive homozygotes and negative homozygotes of *Sh2r6hs* transgenic plants from *Table 4* in Smidansky, et al. [63]. For both model simulation and experimental data, grain yield, grain nitrogen concentration and harvest index at harvest of negative homozygotes of *Sh2r6hs* transgenic plants were used as a reference, i.e., data at all time points and for other plants were normalized by dividing them, respectively. Extracted experimental data are tabulated in Supplementary Datasets 2.

Sensitivity analysis of model parameters

To conduct parameter perturbations, 26 model parameters controlling capacities of biochemical processes (v_{max_*}) and two phloem resistance parameters ($R_{\_phloem\_leaf\_root}$ and $R_{\_phloem\_leaf\_grain}$) were perturbed within [0.1, 10] folds to their default values individually. Grain number per ear was set to 190 for each group to allow sufficient sink capacity. Grain yield and values of other seven agronomic traits were calculated for each simulation, i.e., whole grain-filling season total canopy photosynthesis, whole grain-filling season total root nitrogen uptake, active grain filling duration, grain nitrogen concentration at harvest, average grain filling rate from flowering to harvest, dry matter harvest index and nitrogen harvest index. The active grain filling duration (GFD) was determined as the time interval between flowering day and the day when total grain dry weight ($TGDW_1$) reaches 95% of its maximal value (at the end of the simulation), and average grain filling rate (GFR) was calculated as:

$$GFR = \frac{TGDW_1}{GFD}$$

1141

Identification of optimal parameter combinations to maximize grain yield using a genetic algorithm

A genetic algorithm was used to identify the optimal value combinations of parameters to maximize grain yield. In brief, this algorithm mimics the process of natural biological evolution. the above mentioned 28 molecular biochemical/biophysical parameters, together with partitioning of initial mass between three organs at flowering, i.e., root weight, leaf area and grain number, were perturbed independently by randomly multiplying a scaling coefficient ranging from 0.5 to 2. Grain yield was used as the selection pressure in the algorithm.

To avoid a local optimum at the end of the evolution, five independent evolutionary populations under normal condition (ambient CO_2 concentration is 400 µbar; soil I-N concentration is 0.14 mol m^{-3}; grain-filling season is 50 days; see light intensity, air temperature and humidity information in Supplementary Figure 14) were constructed. Each of the evolutionary populations has 100*100 individuals, i.e., 100 generations with 100 individuals in each generation. As mentioned above, each parameter was restricted to vary between 0.5 to 2-fold of its default value. For the sake of comparison between lines, for each individual in the virtual population, we restrict the total plant equivalent carbon and nitrogen at flowering to be the same. Specifically, for total plant equivalent carbon balance, we have:

$$\frac{\Delta SStar0}{\eta_{store}} = -\frac{\Delta RW0 + \Delta LA0/SLA + \Delta GN0 \cdot HW}{\eta_{grow}}$$

in which Cstar0, RW0, LA0 and GN0 are stem stored starch weight, root weight, leaf area and grain number at flowering, respectively; η_{grow} is growth efficiency, i.e., the ratio of biomass production to the sucrose consumption (g g^{-1}); η_{store} is storage efficiency, i.e., the ratio of carbon fixed in starch to the total amount of carbon used by sucrose [mol C (mol C)$^{-1}$]; SLA is specific leaf area; HW is husk weight per spikelet.

Amount of total nitrogen in leaf is a constant:

$$\Delta LP0 = -\Delta LA0 / SLA \cdot c_{leaf\_strucN} - \Delta LV0 \cdot (c_{leaf\_ON} + c_{leaf\_IN})$$

in which LP0 and LV0 are total leaf protein and leaf volume at flowering, respectively;

c_{leaf_ON} and c_{leaf_IN} are concentrations of leaf organic nitrogen and inorganic nitrogen,

respectively; c_{leaf_strucN} is nitrogen concentration for leaf structural matter.

Amount of total nitrogen in root, stem and grain is a constant:

$$\Delta SPro0 = -(\Delta RW0 \cdot c_{root\_strucN} + \Delta RV0 \cdot (c_{root\_ON} + c_{root\_IN}) + \Delta GN0 \cdot HW \cdot c_{husk\_strucN})$$

in which SPro0 and RV0 are stem stored protein and total root volume at flowering,

respectively; c_{root_ON} and c_{root_IN} are concentrations of root organic nitrogen and

inorganic nitrogen, respectively; c_{root_strucN} and c_{husk_strucN} are nitrogen concentration for

root and husk structural matter, respectively.

1178

1179   **Data and software availability**

1180   Experimental data extracted from literature that used in model parameterization and

1181   model-data comparison were given in Supplementary Datasets 2.

1182   Source code is free for use for academic and non-commercial applications upon request.

1183

References

1   Ray, D. K., Mueller, N. D., West, P. C. & Foley, J. A. Yield trends are insufficient to double
global crop production by 2050. *PLoS One* **8**, e66428 (2013).
2   Long, S. P., Marshall-Colon, A. & Zhu, X.-G. Meeting the global food demand of the future
by engineering crop photosynthesis and yield potential. *Cell* **161**, 56-66 (2015).
3   Zhao, C. *et al.* Temperature increase reduces global yields of major crops in four independent
estimates. *Proc Natl Acad Sci USA* **114**, 9326-9331, doi:10.1073/pnas.1701762114 (2017).
4   Bailey-Serres, J., Parker, J. E., Ainsworth, E. A., Oldroyd, G. E. D. & Schroeder, J. I. Genetic
strategies for improving crop yields. *Nature* **575**, 109-118, doi:10.1038/s41586-019-1679-0
(2019).
5   Peleman, J. D. & van der Voort, J. R. Breeding by design. *Trends Plant Sci* **8**, 330-334,
doi:10.1016/S1360-1385(03)00134-1 (2003).
6   Wang, Y. *et al.* Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers
heritable resistance to powdery mildew. *Nat Biotechnol* **32**, 947-951 (2014).

1198	7	Mickelbart, M. V., Hasegawa, P. M. & Bailey-Serres, J. Genetic mechanisms of abiotic stress
1199		tolerance that translate to crop yield stability. *Nature Reviews. Genetics* **16**, 237 (2015).
1200	8	Qian, Q., Guo, L. B., Smith, S. M. & Li, J. Y. Breeding high-yield superior quality hybrid
1201		super rice by rational design. *Natl Sci Rev* **3**, 283-294, doi:10.1093/nsr/nww006 (2016).
1202	9	Zeng, D. L. *et al.* Rational design of high-yield and superior-quality rice. *Nat Plants* **3**, 17031,
1203		doi:10.1038/Nplants.2017.31 (2017).
1204	10	Hartung, F. & Schiemann, J. Precise plant breeding using new genome editing techniques:
1205		opportunities, safety and regulation in the EU. *The Plant Journal* **78**, 742-752 (2014).
1206	11	Araus, J. L. & Cairns, J. E. Field high-throughput phenotyping: the new crop breeding
1207		frontier. *Trends Plant Sci* **19**, 52-61 (2014).
1208	12	Wallace, J. G., Rodgers-Melnick, E. & Buckler, E. S. On the road to breeding 4.0: unraveling
1209		the good, the bad, and the boring of crop quantitative genomics. *Annual Review of Genetics*
1210		**52**, 421-444, doi:10.1146/annurev-genet-120116-024846 (2018).
1211	13	Watt, M. *et al.* Phenotyping: new windows into the plant for breeders. *Annu Rev Plant Biol*
1212		**71**, 689-712, doi:10.1146/annurev-arplant-042916-041124 (2020).
1213	14	Chang, T.-G., Chang, S., Song, Q.-F., Perveen, S. & Zhu, X.-G. Systems models, phenomics
1214		and genomics: three pillars for developing high-yielding photosynthetically efficient crops. *In*
1215		*Silico Plants* **1**, doi:10.1093/insilicoplants/diy003 (2019).
1216	15	White, A. C., Rogers, A., Rees, M. & Osborne, C. P. How can we make plants grow faster? A
1217		source–sink perspective on growth rate. *J Exp Bot* **67**, 31-45, doi:10.1093/jxb/erv447 (2015).
1218	16	Xing, Y. & Zhang, Q. Genetic and molecular bases of rice yield. *Annu Rev Plant Biol* **61**, 421-
1219		442 (2010).
1220	17	Wang, Y. & Li, J. Molecular basis of plant architecture. *Annu Rev Plant Biol* **59**, 253-279
1221		(2008).
1222	18	Li, Z., Pinson, S. R. M., Stansel, J. W. & Paterson, A. H. Genetic dissection of the source-sink
1223		relationship affecting fecundity and yield in rice (shape *Oryza sativa* L.). *Mol Breed* **4**, 419-
1224		426 (1998).
1225	19	Donald, C. M. The breeding of crop ideotypes. *Euphytica* **17**, 385-403 (1968).
1226	20	Chen, W., Xu, Z., Zhang, W., Zhang, L. & Yang, S. Creation of new plant type and breeding
1227		rice for super high yield. *Acta Agronomica Sinica* **27**, 665-672 (2000).
1228	21	Peng, S., Khush, G. S., Virk, P., Tang, Q. & Zou, Y. Progress in ideotype breeding to increase
1229		rice yield potential. *Field Crops Res* **108**, 32-38 (2008).
1230	22	Yuan, L. Progress in super-hybrid rice breeding. *Crop J* **5**, 100-102 (2017).
1231	23	Jiao, Y. Q. *et al.* Regulation of *OsSPL14* by OsmiR156 defines ideal plant architecture in rice.
1232		*Nat Genet* **42**, 541-544, doi:10.1038/ng.591 (2010).
1233	24	Song, Q. F., Zhang, G. L. & Zhu, X. G. Optimal crop canopy architecture to maximise canopy
1234		photosynthetic CO_2 uptake under elevated CO_2 - a theoretical study using a mechanistic model
1235		of canopy photosynthesis. *Funct Plant Biol* **40**, 109-124, doi:10.1071/FP12056 (2013).
1236	25	Zhang, L. *et al.* A natural tandem array alleviates epigenetic repression of IPA1 and leads to
1237		superior yielding rice. *Nat Commun* **8**, doi:10.1038/ncomms14789 (2017) (2017).
1238	26	Pan, Q. *et al.* The genetic basis of plant architecture in 10 maize recombinant inbred line
1239		populations. *Plant Physiol* **175**, 858-873, doi:10.1104/pp.17.00709 (2017).
1240	27	Long, S. P., Zhu, X. G., Naidu, S. L. & Ort, D. R. Can improvement in photosynthesis
1241		increase crop yields? *Plant Cell Environ* **29**, 315-330, doi:10.1111/j.1365-3040.2005.01493.x
1242		(2006).
1243	28	Simkin, A. J., López-Calcagno, P. E. & Raines, C. A. Feeding the world: improving
1244		photosynthetic efficiency for sustainable crop production. *J Exp Bot* **70**, 1119–1140,
1245		doi:10.1093/jxb/ery445 (2019).
1246	29	Sonnewald, U. *et al.* The Cassava Source–Sink project: opportunities and challenges for crop
1247		improvement by metabolic engineering. *The Plant Journal* **n/a**, doi:10.1111/tpj.14865 (2020).
1248	30	Chang, T.-G. & Zhu, X.-G. Source-sink interaction: a century old concept under the light of
1249		modern molecular systems biology. *J Exp Bot* **68**, 4417-4431, doi:10.1093/jxb/erx002 (2017).
1250	31	Fernie, A. R. *et al.* Synchronization of developmental, molecular and metabolic aspects of
1251		source-sink interactions. *Nat Plants* **6**, 55-66, doi:10.1038/s41477-020-0590-x (2020).
1252	32	Marshall-Colon, A. *et al.* Crops *in silico*: generating virtual crops using an integrative and
1253		multi-scale modeling platform. *Front Plant Sci* **8** (2017).
1254	33	Hammer, G., Messina, C., Wu, A. & Cooper, M. Biological reality and parsimony in crop
1255		models—why we need both in crop improvement! *In Silico Plants* **1**,
1256		doi:10.1093/insilicoplants/diz010 (2019).

1257	34	Zhu, X. G., Wang, Y., Ort, D. R. & Long, S. P. e‑photosynthesis: a comprehensive dynamic
1258		mechanistic model of C$_3$ photosynthesis: from light capture to sucrose synthesis. *Plant Cell*
1259		*Environ* **36**, 1711-1727 (2013).
1260	35	Wang, Y., Long, S. P. & Zhu, X.-G. Elements required for an efficient NADP-malic enzyme
1261		type C$_4$ photosynthesis. *Plant Physiol* **164**, 2231-2246 (2014).
1262	36	Minchin, P. E. H. & Lacointe, A. New understanding on phloem physiology and possible
1263		consequences for modelling long-distance carbon transport. *New Phytol* **166**, 771-779,
1264		doi:10.1111/j.1469-8137.2005.01323.x (2005).
1265	37	Valentim, F. L. *et al.* A quantitative and dynamic model of the Arabidopsis flowering time
1266		gene regulatory network. *PLoS One* **10**, e0116973 (2015).
1267	38	Buckler, E. S. *et al.* The genetic architecture of maize flowering time. *Science* **325**, 714-718,
1268		doi:10.1126/science.1174276 (2009).
1269	39	Lynch, J. P., Nielsen, K. L., Davis, R. D. & Jablokow, A. G. SimRoot: modelling and
1270		visualization of root systems. *Plant Soil* **188**, 139-151 (1997).
1271	40	Kromdijk, J. *et al.* Improving photosynthesis and crop productivity by accelerating recovery
1272		from photoprotection. *Science* **354**, 857-861 (2016).
1273	41	Zhu, X. G., Ort, D. R., Whitmarsh, J. & Long, S. P. The slow reversibility of photosystem II
1274		thermal energy dissipation on transfer from high to low light may cause large losses in carbon
1275		gain by crop canopies: a theoretical analysis. *J Exp Bot* **55**, 1167-1175 (2004).
1276	42	South, P. F., Cavanagh, A. P., Liu, H. W. & Ort, D. R. Synthetic glycolate metabolism
1277		pathways stimulate crop growth and productivity in the field. *Science* **363**, eaat9077,
1278		doi:10.1126/science.aat9077 (2019).
1279	43	Xin, C., Tholen, D. & Zhu, X. G. The benefits of photorespiratory bypasses: How can they
1280		work? *Plant Physiol* **167**, 574-585 (2015).
1281	44	Wu, A., Hammer, G. L., Doherty, A., von Caemmerer, S. & Farquhar, G. D. Quantifying
1282		impacts of enhancing photosynthesis on crop yield. *Nat Plants* **5**, 380-388,
1283		doi:10.1038/s41477-019-0398-8 (2019).
1284	45	Zhang, N. *et al.* Genome-wide association of carbon and nitrogen metabolism in the maize
1285		nested association mapping population. *Plant Physiol* **168**, 575-583, doi:10.1104/pp.15.00025
1286		(2015).
1287	46	Li, C. *et al.* Targeted, random mutagenesis of plant genes with dual cytosine and adenine base
1288		editors. *Nat Biotechnol*, doi:10.1038/s41587-019-0393-7 (2020).
1289	47	Walton, R. T., Christie, K. A., Whittaker, M. N. & Kleinstiver, B. P. Unconstrained genome
1290		targeting with near-PAMless engineered CRISPR-Cas9 variants. *Science* **368**, 290-296,
1291		doi:10.1126/science.aba8853 (2020).
1292	48	Tcherkez, G. G. B. & Ribas-Carbo, M. in *Terrestrial Photosynthesis in a Changing*
1293		*Environment: A Molecular, Physiological, and Ecological Approach* (eds Francesco
1294		Loreto, Hipólito Medrano, & Jaume Flexas) 1-6 (Cambridge University Press, 2012).
1295	49	Yoshida, S. *Fundamentals of rice crop science*. (Los Banos, Philippines: International Rice
1296		Research Institute., 1981).
1297	50	Tang, W.-B. *et al.* Root characteristics of high-yield C Liangyou rice combinations of two-line
1298		hybrid rice. *Sci Agric Sinica* **43**, 2859-2868, doi:10.3864/j.issn.0578-1752.2010.14.004
1299		(2010).
1300	51	Haque, M. M., Pramanik, H. R., Biswas, J. K., Iftekharuddaula, K. & Hasanuzzaman, M.
1301		Comparative performance of hybrid and elite inbred rice varieties with respect to their source-
1302		sink relationship. *The Scientific World Journal* **2015** (2015).
1303	52	Zhao, Q., Huang, P. & Ling, Q. Relations between canopy apparent photosynthesis and store
1304		matter in stem and sheath between and yield and nitrogen regulations in rice. *Sci Agric Sinica*
1305		**34**, 304-310 (2001).
1306	53	Yang, J., Zhang, J., Wang, Z., Liu, K. & Wang, P. Post-anthesis development of inferior and
1307		superior spikelets in rice in relation to abscisic acid and ethylene. *J Exp Bot* **57**, 149-160
1308		(2006).
1309	54	Yang, J., Zhang, J., Wang, Z., Zhu, Q. & Wang, W. Remobilization of carbon reserves in
1310		response to water deficit during grain filling of rice. *Field Crops Res* **71**, 47-55 (2001).
1311	55	Ida, M., Ohsugi, R., Sasaki, H., Aoki, N. & Yamagishi, T. Contribution of nitrogen absorbed
1312		during ripening period to grain filling in a high-yielding rice variety, Takanari. *Plant Prod Sci*
1313		**12**, 176-184, doi:10.1626/pps.12.176 (2009).

56   Witt, C. *et al.* Internal nutrient efficiencies of irrigated lowland rice in tropical and subtropical
Asia. *Field Crops Res* **63**, 113-138, doi:10.1016/S0378-4290(99)00031-3 (1999).
57   Zhao, Y. *et al.* Nitrogen effect on amino acid composition in leaf and grain of japonica rice
during grain filling stage. *J Cereal Sci* **64**, 29-33 (2015).
58   Kobata, T., Sugawara, M. & Takatu, S. Shading during the early grain filling period does not
affect potential grain dry matter increase in rice. *Agron J* **92**, 411-417 (2000).
59   Kim, H., Lieffering, M., Miura, S., Kobayashi, K. & Okada, M. Growth and nitrogen uptake
of CO_2 - enriched rice under field conditions. *New Phytol* **150**, 223-229 (2001).
60   Shimono, H. & Bunce, J. A. Acclimation of nitrogen uptake capacity of rice to elevated
atmospheric CO_2 concentration. *Ann Bot* **103**, 87-94, doi:10.1093/aob/mcn209 (2009).
61   Liang, C. *et al.* OsNAP connects abscisic acid and leaf senescence by fine-tuning abscisic acid
biosynthesis and directly targeting senescence-associated genes in rice. *Proc Natl Acad Sci*
USA **111**, 10013-10018 (2014).
62   Wang, E. *et al.* Control of rice grain-filling and yield by a gene with a potential signature of
domestication. *Nat Genet* **40**, 1370-1374 (2008).
63   Smidansky, E. D., Martin, J. M., Hannah, L. C., Fischer, A. M. & Giroux, M. J. Seed yield
and plant biomass increases in rice are conferred by deregulation of endosperm ADP-glucose
pyrophosphorylase. *Planta* **216**, 656-664, doi:10.1007/s00425-002-0897-z (2003).
64   Ni, J. *et al.* Overexpression of sugar transporter gene *PbSWEET4* of pear causes sugar reduce
and early senescence in leaves. *Gene* **743**, 144582, doi:10.1016/j.gene.2020.144582 (2020).
65   Zhou, Y. *et al.* Overexpression of *OsSWEET5* in rice causes growth retardation and
precocious senescence. *PLoS One* **9**, e94210-e94210, doi:10.1371/journal.pone.0094210
(2014).
66   Smidansky, E. D. *et al.* Enhanced ADP-glucose pyrophosphorylase activity in wheat
endosperm increases seed yield. *Proc Natl Acad Sci USA* **99**, 1724-1729 (2002).
67   Eom, J.-S. *et al.* Impaired function of the tonoplast-localized sucrose transporter in rice,
OsSUT2, limits the transport of vacuolar reserve sucrose and affects plant growth. *Plant*
Physiol **157**, 109-119 (2011).
68   Thomas, H. & Ougham, H. The stay-green trait. *J Exp Bot* **65**, 3889-3900,
doi:10.1093/jxb/eru037 (2014).
69   Wang, Y. *et al.* Development of a three-dimensional ray-tracing model of sugarcane canopy
photosynthesis and its application in assessing impacts of varied row spacing. *Bioenergy Res*
10, 626-634, doi:10.1007/s12155-017-9823-x (2017).
70   Deng, J. *et al.* Models and tests of optimal density and maximal yield for crop plants. *Proc*
Natl Acad Sci USA **109**, 15823-15828, doi:10.1073/pnas.1210955109 (2012).
71   Song, Q., Srinivasan, V., Long, S. P. & Zhu, X.-G. Decomposition analysis on soybean
productivity increase under elevated CO_2 using 3-D canopy model reveals synergestic effects
of CO_2 and light in photosynthesis. *Ann Bot*, doi:10.1093/aob/mcz163 (2019).
72   Chenu, K. *et al.* Contribution of crop models to adaptation in wheat. *Trends Plant Sci* **22**, 472-
490, doi:10.1016/j.tplants.2017.02.003 (2017).
73   Tuberosa, R. Phenotyping for drought tolerance of crops in the genomics era. *Front Physiol* **3**,
347-347, doi:10.3389/fphys.2012.00347 (2012).
74   Peng, B., Guan, K., Tang, J., Ainsworth, E. A. & Zhou, W. Towards a multiscale crop
modelling framework for climate change adaptation assessment. *Nat Plants* **6**, 338-348
(2020).
75   Fu, J.-D., Yan, Y.-F. & Lee, B.-W. Physiological characteristics of a functional stay-green rice
"SNU-SG1" during grain-filling period. *J Crop Sci Biotechnol* **12**, 47-52,
doi:10.1007/s12892-009-0078-8 (2009).
76   Braun, D. M., Wang, L. & Ruan, Y.-L. Understanding and manipulating sucrose phloem
loading, unloading, metabolism, and signalling to enhance crop yield and food security. *J Exp*
Bot **65**, 1713-1735, doi:10.1093/jxb/ert416 (2014).
77   Yang, J. & Zhang, J. Grain filling of cereals under soil drying. *New Phytol* **169**, 223-236
(2006).
78   Cakir, B., Tuncel, A., Hwang, S.-K. & Okita, T. W. in *Starch: Metabolism and Structure*
(ed Yasunori Nakamura)   371-395 (Springer Japan, 2015).
79   Yu, S.-M., Lo, S.-F. & Ho, T.-H. D. Source–sink communication: regulated by hormone,
nutrient, and stress cross-signaling. *Trends Plant Sci* **20**, 844-857,
doi:10.1016/j.tplants.2015.10.009 (2015).

1372	80	Postma, J. A. *et al.* OpenSimRoot: widening the scope and application of root architectural
1373		models. *New Phytol* **215**, 1274-1286 (2017).
1374	81	Watanabe, T. *et al.* Rice morphogenesis and plant architecture: Measurement, specification
1375		and the reconstruction of structural development by 3D architectural modelling. *Ann Bot* **95**,
1376		1131-1143, doi:Doi 10.1093/Aob/Mci136 (2005).
1377	82	Chang, T.-G. *et al.* A three-dimensional canopy photosynthesis model in rice with a complete
1378		description of the canopy architecture, leaf physiology, and mechanical properties. *J Exp Bot*
1379		**70**, 2479–2490, doi:10.1093/jxb/ery430 (2019).
1380	83	Sonnewald, U. & Fernie, A. R. Next-generation strategies for understanding and influencing
1381		source–sink relations in crop plants. *Curr Opin Plant Biol* **43**, 63-70,
1382		doi:10.1016/j.pbi.2018.01.004 (2018).
1383	84	Salon, C. *et al.* Fluxomics links cellular functional analyses to whole-plant phenotyping. *J Exp*
1384		*Bot* **68**, 2083-2098, doi:10.1093/jxb/erx126 (2017).
1385	85	Cheung, F. Yield: The search for the rice of the future. *Nature* **514**, S60-S61 (2014).
1386	86	Zhu, X. G., de Sturler, E. & Long, S. P. Optimizing the distribution of resources between
1387		enzymes of carbon metabolism can dramatically increase photosynthetic rate: A numerical
1388		simulation using an evolutionary algorithm. *Plant Physiol* **145**, 513-526, doi:DOI
1389		10.1104/pp.107.103713 (2007).
1390	87	Glass, A. D. M. *et al.* The regulation of nitrate and ammonium transport systems in plants. *J*
1391		*Exp Bot* **53**, 855-864, doi:DOI 10.1093/jexbot/53.370.855 (2002).
1392	88	Pitman, M. & Cram, W. Regulation of inorganic ion transport in plants. *Ion Transport in*
1393		*Plants*, 465-481 (2013).
1394	89	Henry, L. T. & Raper Jr, C. D. Soluble carbohydrate allocation to roots, photosynthetic rate of
1395		leaves, and nitrate assimilation as affected by nitrogen stress and irradiance. *Botanical*
1396		*Gazette*, 23-33 (1991).
1397	90	Monod, J. The growth of bacterial cultures. *Annual Reviews in Microbiology* **3**, 371-394
1398		(1949).
1399	91	Miflin, B. & Lea, P. Ammonia assimilation. *The Biochemistry of Plants* **5**, 169-202 (1980).
1400	92	Radin, J. W., Parker, L. L. & Sell, C. R. Partitioning of sugar between growth and nitrate
1401		reduction in cotton roots. *Plant Physiol* **62**, 550-553 (1978).
1402	93	Muller, B. *et al.* Water deficits uncouple growth from photosynthesis, increase C content, and
1403		modify the relationships between C and growth in sink organs. *J Exp Bot*, erq438 (2011).
1404	94	Willaume, M. & Pagès, L. Correlated responses of root growth and sugar concentrations to
1405		various defoliation treatments and rhythmic shoot growth in oak tree seedlings (*Quercus*
1406		*pubescens*). *Ann Bot*, mcq270 (2011).
1407	95	Johnson, I. R. & Thornley, J. H. M. Dynamic model of the response of a vegetative grass crop
1408		to light, temperature and nitrogen. *Plant Cell Environ* **8**, 485-499, doi:10.1111/j.1365-
1409		3040.1985.tb01684.x (1985).
1410	96	Asseng, S., Richter, C. & Wessolek, G. Modelling root growth of wheat as the linkage
1411		between crop and soil. *Plant Soil* **190**, 267-277, doi:10.1023/A:1004228201299 (1997).
1412	97	Fanello, D. D., Bartoli, C. G. & Guiamet, J. J. Qualitative and quantitative modifications of
1413		root mitochondria during senescence of above-ground parts of *Arabidopis thaliana*. *Plant Sci*
1414		**258**, 112-121, doi:10.1016/j.plantsci.2017.01.013 (2017).
1415	98	Von Caemmerer, S. *Biochemical models of leaf photosynthesis*. (Csiro publishing, 2000).
1416	99	De Pury, D. & Farquhar, G. Simple scaling of photosynthesis from leaves to canopies without
1417		the errors of big-leaf models. *Plant, Cell & Environment* **20**, 537-557 (1997).
1418	100	Tazoe, Y., Noguchi, K. & Terashima, I. Effects of growth light and nitrogen nutrition on the
1419		organization of the photosynthetic apparatus in leaves of a C_4 plant, Amaranthus cruentus.
1420		*Plant Cell Environ* **29**, 691-700 (2006).
1421	101	San-oh, Y., Sugiyama, T., Yoshita, D., Ookawa, T. & Hirasawa, T. The effect of planting
1422		pattern on the rate of photosynthesis and related processes during ripening in rice plants. *Field*
1423		*Crops Res* **96**, 113-124 (2006).
1424	102	Azcón-Bieto, J. Inhibition of photosynthesis by carbohydrates in wheat leaves. *Plant Physiol*
1425		**73**, 681-686 (1983).
1426	103	Stitt, M. Rising CO_2 levels and their potential significance for carbon flow in photosynthetic
1427		cells. *Plant Cell Environ* **14**, 741-762 (1991).
1428	104	Paul, M. J. & Pellny, T. K. Carbon metabolite feedback regulation of leaf photosynthesis and
1429		development. *Journal of experimental botany* **54**, 539-547, doi:10.1093/jxb/erg052 (2003).

105   Jarvis, P. The interpretation of the variations in leaf water potential and stomatal conductance
found in canopies in the field. *Philosophical Transactions of the Royal Society of London B:*
Biological Sciences **273**, 593-610 (1976).
106   Farquhar, G. D. & Sharkey, T. D. Stomatal conductance and photosynthesis. *Annu Rev Plant*
Physiol **33**, 317-345 (1982).
107   Zhang, J. & Davies, W. Changes in the concentration of ABA in xylem sap as a function of
changing soil water status can account for changes in leaf conductance and growth. *Plant Cell*
Environ **13**, 277-285 (1990).
108   Leuning, R. A critical appraisal of a combined stomatal-photosynthesis model for C_3 plants.
Plant Cell Environ **18**, 339-355 (1995).
109   Allen, R. G., Pereira, L. S., Raes, D. & Smith, M. Crop evapotranspiration-Guidelines for
computing crop water requirements-FAO Irrigation and drainage paper 56. *FAO, Rome* **300**,
D05109 (1998).
110   Andrews, M. The partitioning of nitrate assimilation between root and shoot of higher plants.
Plant Cell Environ **9**, 511-519 (1986).
111   Nunes-Nesi, A., Fernie, A. R. & Stitt, M. Metabolic and signaling aspects underpinning the
regulation of plant carbon nitrogen interactions. *Mol Plant* **3**, 973-996, doi:10.1093/mp/ssq049
(2010).
112   Walker, B. J., Strand, D. D., Kramer, D. M. & Cousins, A. B. The response of cyclic electron
flow around photosystem I to changes in photorespiration and nitrate assimilation. *Plant*
Physiol **165**, 453-462 (2014).
113   Pilkington, S. M. *et al.* Relationship between starch degradation and carbon demand for
maintenance and growth in *Arabidopsis thaliana* in different irradiance and temperature
regimes. *Plant Cell Environ* **38**, 157-171 (2015).
114   Feugier, F. G. & Satake, A. Dynamical feedback between circadian clock and sucrose
availability explains adaptive response of starch metabolism to various photoperiods. *Front*
Plant Sci **3**, 305 (2013).
115   Scialdone, A. *et al.* Arabidopsis plants perform arithmetic division to prevent starvation at
night. *Elife* **2**, e00669 (2013).
116   Seaton, D. D., Ebenhöh, O., Millar, A. J. & Pokhilko, A. Regulatory principles and
experimental approaches to the circadian control of starch turnover. *Journal of the Royal*
Society Interface **11**, 20130979 (2014).
117   Yin, X. & van Laar, H. Crop Systems Dynamics: An Ecophysiological Model of Genotype-
by-Environment Interactions (GECROS). *Wageningen Academic Pub., Wageningen* (2005).
118   Sinclair, T. & De Wit, C. Analysis of the carbon and nitrogen limitations to soybean yield.
Agron J **68**, 319-324 (1976).
119   Masclaux‑Daubresse, C., Reisdorf‑Cren, M. & Orsel, M. Leaf nitrogen remobilisation for
plant development and grain filling. *Plant Biol* **10**, 23-36 (2008).
120   Ou-Lee, T. M. & Setter, T. L. Enzyme activities of starch and sucrose pathways and growth of
apical and basal maize kernels. *Plant Physiol* **79**, 848-851, doi:10.1104/pp.79.3.848 (1985).
121   Ober, E. S., Setter, T. L., Madison, J. T., Thompson, J. F. & Shapiro, P. S. Influence of water
deficit on maize endosperm development : enzyme activities and RNA transcripts of starch
and zein synthesis, abscisic acid, and cell division. *Plant Physiol* **97**, 154-164,
doi:10.1104/pp.97.1.154 (1991).
122   Ohdan, T. *et al.* Expression profiling of genes involved in starch synthesis in sink and source
organs of rice. *J Exp Bot* **56**, 3229-3244, doi:10.1093/jxb/eri292 (2005).
123   Toyosawa, Y. *et al.* Deficiency of starch synthase IIIa and IVb alters starch granule
morphology from polyhedral to spherical in rice endosperm. *Plant Physiol* **170**, 1255-1270,
doi:10.1104/pp.15.01232 (2016).
124   Cai, Y. *et al. OsPK2* encodes a plastidic pyruvate kinase involved in rice endosperm starch
synthesis, compound granule formation and grain filling. *Plant Biotechnol J* **16**, 1878-1891,
doi:10.1111/pbi.12923 (2018).
125   Olsen, O. A. Nuclear endosperm development in cereals and *Arabidopsis thaliana. Plant Cell*
16 Suppl, S214-227, doi:10.1105/tpc.017111 (2004).
126   Yang, Z., Van Oosterom, E. J., Jordan, D. R. & Hammer, G. L. Pre-anthesis ovary
development determines genotypic differences in potential kernel weight in sorghum. *J Exp*
Bot, erp019 (2009).

127   Tashiro, T. & Wardlaw, I. F. A comparison of the effect of high temperature on grain
development in wheat and rice. *Ann Bot* **64**, 59-65 (1989).
128   Lalonde, S., Tegeder, M., Throne‐Holst, M., Frommer, W. & Patrick, J. Phloem loading and
unloading of sugars and amino acids. *Plant Cell Environ* **26**, 37-56 (2003).
129   Cannell, M. G. R. & Thornley, J. H. M. Modelling the components of plant respiration: Some
guiding principles. *Ann Bot* **85**, 45-54, doi:DOI 10.1006/anbo.1999.0996 (2000).

1493

1494
1495

Acknowledgements

This research was financially supported by the CAS (Chinese Academy of Science)

strategic leading project (grant no. XDB27020105), the open research fund of the

state key laboratory of hybrid rice (Hunan Hybrid Rice Research Center), and the

national key research and development project (2018YFA0900600). The authors

thank Prof. Govindjee for constructive comments and improving the language of the

manuscript.

Author contributions

X.Z. designed the study; T.C. performed the analysis; T.C. and X.Z. wrote the

manuscript.

Competing interests

The authors declare no competing financial interests

Supplementary Information

The following material is available in the online version of this article.

Supplementary Figure 1. Predicted relative changes of plant physiological traits for

the wild type, the *ps1-D* mutant and the *OsNAP* RNAi plants.

Supplementary Figure 2. Predicted relative changes of plant physiological traits for

the wild type, the *gif1* mutant and the *GIF1* overexpression plants.

Supplementary Figure 3. Predicted relative changes of plant physiological traits for

'source limited' plants with $v_{\text{m\_grain\_Suc\_ul}}$ set to 100%, 50% and 200% of default

values.

Supplementary Figure 4. The responses of major agronomic traits to the change in

model parameters $v_{\text{m\_grain\_Star\_syn}}$, $R_{\text{phloem\_leaf\_root}}$, $v_{\text{m\_root\_N\_upt}}$ and $v_{\text{m\_leaf\_N\_ass}}$,

respectively.

Supplementary Figure 5. The responses of major agronomic traits to the change in

model parameters $v_{\text{m\_root\_N\_ass}}$, $v_{\text{m\_leaf\_Suc\_l}}$, $v_{\text{m\_leaf\_IN\_ul}}$ and $v_{\text{m\_root\_IN\_l}}$, respectively.

Supplementary Figure 6. The responses of major agronomic traits to the change in

model parameters $v_{\text{m\_leaf\_Star\_deg}}$, $v_{\text{m\_stem\_Star\_deg}}$, $v_{\text{m\_leaf\_Suc\_syn}}$ and $v_{\text{m\_leaf\_Pro\_deg}}$,

respectively.

Supplementary Figure 7. The responses of major agronomic traits to the change in

model parameters $v_{\text{m\_leaf\_ON\_l}}$, $v_{\text{m\_root\_grow}}$, $v_{\text{m\_leaf\_Star\_syn}}$ and $v_{\text{m\_grain\_Pro\_syn}}$, respectively.

Supplementary Figure 8. The responses of major agronomic traits to the change in

model parameters $v_{m\_grain\_ON\_ul}$, $v_{m\_grain\_grow}$, $v_{m\_grain\_Suc\_ul}$ and $R_{\_phloem\_leaf\_grain}$,

respectively.

Supplementary Figure 9. The responses of major agronomic traits to the change in

model parameters $v_{m\_root\_Suc\_ul}$, $v_{m\_root\_ON\_l}$, $v_{m\_ON\_Vbtransfer}$ and $v_{m\_stem\_Star\_syn}$,

respectively.

Supplementary Figure 10. The responses of major agronomic traits to the change in

model parameters $v_{m\_leaf\_Pro\_syn}$, $v_{m\_stem\_Pro\_deg}$, $v_{m\_root\_ON\_ul}$ and $v_{m\_stem\_Pro\_syn}$,

respectively.

Supplementary Figure 11. The evolution of total grain yield throughout generations

in the *in silico* 'evolutionary populations'.

Supplementary Figure 12. Distribution of relative values of plant initial size related

parameters at the flowering stage for *in silico* high-yield 'Elite' individuals.

Supplementary Figure 13. WACNI predictions of grain filling patterns for plants in

in silico evolutionary populations.

Supplementary Figure 14. Daily weather data of the normal condition used in

WACNI.

Supplementary Table 1. Detailed description and comprehensive parameterization

procedure of the parameters used in WACNI.

Supplementary Table 2. Effects on eight agronomic traits, obtained by manipulating

candidate genes and/or QTLs responsible for capacities of 28 reaction and diffusion

processes in WACNI.

Supplementary Table 3. Functional description of genes in Table S2.

Supplementary Datasets 1. 'Supplementary Datasets 1 paras_of_elites.xlsx'.

Parameter values (relative fold change from default values) of 'Elite' individuals and

the final grain weight and grain nitrogen concentration.

Supplementary Datasets 2. 'Supplementary Datasets 2

experimental_data_from_literature.xlsx'. Details of experimental data extracted from

literature that used in model parameterization and model-data comparison (Figures 2-

4).

1560