EVALUATION OF ANTIMICROBIAL AND ANTIPROLIFERATIVE ACTIVITIES OF ACTINOBACTERIA ISOLATED FROM THE SALINE LAGOONS OF NORTHWEST PERU.

Rene Flores Clavo[1,2,8*], Nataly Ruiz Quiñones[1,2,7], Álvaro Tasca Hernandez[3¶], Ana Lucia Tasca Gois Ruiz[4&], Lucia Elaine de Oliveira Braga[4&], Zhandra Lizeth Arce Gil[6¶], Luis Miguel Serquen Lopez[7,8¶], Jonas Henrique Costa[5&], Taícia Pacheco Fill[5&], Marcos José Salvador[3¶], Fabiana Fantinatti Garboggini[2].

Graduate Program in Genetics and Molecular Biology, Institute of Biology, University of Campinas (UNICAMP), Campinas, SP, Brazil.

Chemical, Biological and Agricultural Pluridisciplinary Research Center (CPQBA), University of Campinas (UNICAMP), Paulínia, SP, Brazil.

University of Campinas, Department of plant Biology Bioactive Products, Institute of Biology Campinas, Sao Paulo, Brazil.

University of Campinas, Faculty Pharmaceutical Sciences, Campinas, Sao Paulo, Brazil.

University of Campinas, Institute of Chemistry, Campinas, Sao Paulo, Brazil.

Private University Santo Toribio of Mogrovejo, Facultity of Human Medicine, Chiclayo, Lambayeque Perú.

Direction of Investigation Hospital Regional Lambayeque, Chiclayo, Lambayeque, Perú.

Research Center and Innovation and Sciences Actives Multidisciplinary (CIICAM), Department of Biotechnology, Chiclayo, Lambayeque, Perú.

24   * Correspond author: renefloresclavo@gmail.com

26   ¶ These authors contributed equally to this work.

27   & These authors contributed equally to this work.

Abstract

The unexplored saline lagoons of the north of Peru harbor a rich microbiome, due to reported

studies of different extreme environments around the world. In these regions, there are

several ecosystems and microhabitats not yet explored, and little is known about the diversity

of actinobacteria and other microorganisms. We suggest that the endemic bacteria present in

this extreme environment could be source of active molecules with anticancer, antimicrobial,

antiparasitic properties. Using phenotypic and genotypic characterization techniques

including the 16S rRNA were identified into the genera *Streptomyces* 39 (78%),

Pseudonocardia 3 (6%), *Staphylococcus* 4 (8%), *Bacillus* 2 (4%), and *Pseudomonas* 2 (4%).

All isolated bacteria for the genotypic data were preliminarily identified. Actinobacteria

strains were found dominantly in both sites (Lagoon1-3 = 16 isolates and lagoon 4 = 12

isolates). Phylogenetic analysis revealed that 28 isolates were exclusively affiliated to eleven

different clusters of Actinobacteria of the major genus *Streptomyces*. Three *Streptomyces* sp.

strains M-92, B-146, and B-81, were tested for antibacterial and antiproliferative activities.

The results showed antiproliferative activities against three tumor cell lines, U251 glioma;

MCF7 breast; NCI-H460 lung non-small type of cells, and the antibacterial activity to

Staphylococcus aureus ATCC 6538, *E. coli* ATCC 10536, and *Acinetobacter baumanni* AC-

972 which is resistant to multiple drugs. The promising results belong to *Streptomyces* sp. B-

81 strain in the R2A medium using a doxorubicin with control positive, the best result was

from the latter (TGI = 0,57 µg/mL) for glioma; NCI-H460 lung of type non-small cells (TGI

= 0,61 µg/mL), and breast cancer (TGI =0,80 µg/mL), this strain was selected to be

fractionated because it had better antiproliferative and antibacterial activity, and its fractions

were evaluated concerning antiproliferative activity against nine types of tumor cells and one

non-tumor. The methanolic fraction showed a better result in the antiproliferative activity

and was able to inhibit U251 (glioma) (TGI = 38.3 µg/mL), OVCAR-03 (ovary) (TGI = 62.1

µg/mL), and K562 (leukemia) (TGI = 81.5 µg/mL). The methanol 50% - acetate 50% fraction

(Fraction 4) inhibited U251 (glioma) (TGI = 73.5 µg/mL) and UACC-62 (melanoma) (TGI

= 89.4 µg/mL). Moreover, the UHPLC-MS/MS data and molecular networking of

Streptomyces sp. B-81 isolate extract revealed the production cholic acid, Lobophorin A,

Lobophorin B, Lobophorin E, Lobophorin K and compound 6. Extremophilic environments

such as the Mórrope and Bayovar Salt Flats are promising sources of new bacteria with

promising pharmaceutical potential; These compounds could be useful to treat various

infectious diseases or even some type of cancer.

Keywords: *Streptomyces* sp. B-81; saline lagoons; Morrope salt flats; Bayovar salt flats;

antimicrobial activity; antiproliferative activity.

65 1. Introduction

Perú has extreme environments such as the salt marshes located on the coast, center, and

south of the country. In these regions, there are several ecosystems and microhabitats, whose

studies are still restricted and report the diversity of bacteria and other organisms (Caceres

and Legendre, 2009; Flores and Pisfil, 2014; Leon et al. 2016). Among other

microorganisms, the members of the phylum Actinobacteria can be found in all kinds of

extreme environments. *Micromonospora*, *Actinomadura*, and *Nocardiopsis* were reported

from saline soils of the ephemeral salty lakes in Buryatiya (Lubsanova et al. 2014).

Streptomyces, *Nocardiopsis*, and *Nocardioides* were isolated from the Regions Western

Ghats of India (Siddhart et al. 2020). Concurrent to this, there is still little information

documented on isolated substances and their biological activities of the microorganisms

isolated from saline environments. *Micromonospora*, *Streptomyces*, *Salinispora*, and *Dietzia*

were isolated from the coastal zone of the geographically remote young volcanic Easter

Island Chile. Besides, actinobacterial halophilic and halotolerant strains show heterogeneous

physiological characteristics among different genera because these bacteria have the ability

to synthesize secondary metabolites to cope with conditions of high salinity and the different

temperatures present in their environments (Binayke et al. 2018; Oueriaghli et al. 2018;

Boyadzhieva et al. 2018). These extreme conditions favor the development of metabolically

competitiveness to the production enzymes, which can help the bacterial population to adapt

to high salinity (Kim et al. 2019). Furthermore, these microorganisms can also perform

essential processes such as carbon cycle, metal transfer, and the removal of organic pollutants

to higher trophic levels (Fiedler et al. 2005; Blunt et al. 2007). In addition, this microbial

group presents a unique ability to produce new products, mainly antibiotics (Berdy, 2005;

Strohl, 2004).

The search for these microorganisms has been mainly related to the production of antibiotics

and antitumor substances (Newman and Cragg, 2007). There are about 22.500 biologically

active substances obtained from microorganisms, 45% of which are represented by

actinomycetes, with *Streptomyces* being the producer of 70% of them (Hayakawa et al. 2007

and Valliappan et al. 2014). According to Lam (2006), there are new actinobacteria from

habitats not yet explored or little explored to be the source of new bioactive secondary

metabolites.

The bioguided study strategy helps us find a wide variety of microorganisms that have

biotechnological potential (Maldonado et al. 2005). This study reports the isolation and

phylogenetic analysis of a collection of bacterial isolates from the saline lagoons from the

Northern Peru, and their potential as producers of secondary metabolites with antimicrobial

and antiproliferative activities.

102   2. Materials and methods

103   2.1 Site description and sampling

Sample collection at Mórrope saline lagoons was carried out in December 2012 the zone "1",

in January 2013 the zone "2", and in July 2014 the zone "3", while samples were collected at

Bayovar saline lagoons in March 2015 the zone "4" (Figure 1). The present work did not

need permits from a competent authority, since it does not include any protected species,

because the standard is currently in the process of being implemented to achieve its

registration, likewise the sampling areas are lagoons that due to their nature are in the process

of adjudication to the competent governmental organism through the Peruvian Ministry of

the Environment.

Fig 1. a = Geographic location of sampling sites; b and c = lagoons 1, 2 and 3 (State Mórrope); and e = lagoon 4 (State Bayovar).

116 The sources and places from which samples were obtained are detailed in Table 1. Samples

117 were collected aseptically, placed in sterile plastic bags, and kept refrigerated at 4 °C for

118 further processing in the Laboratory of Bacteriology the Regional Hospital of Lambayeque,

119 a flowchart depicting the methodological strategy adopted in this work is shown in Figure 2.

121 **Table 1. Data related to saline lagoons samples.**

Mórrope lagoons				
Collection zone	**Sample type**	**Geographic location**		
Zone 1 (December 2012)	A; AES	6°10S;80°35W (3 samples x triplicate)	6°11S;80°37W (3 samples x triplicate)	6°9S;80°39W (3 samples x triplicate)
Zone 2 (January 2013)	AES; SE	6°19S;80°28W (3 samples x triplicate)	6°17S;80°26W (3 samples x triplicate)	6°18S;80°27W (3 samples x triplicate)
Zone 3 (December 2014)	A; SE	6°08S;80°50W (3 samples x triplicate)	6°08S;80°51W (3 samples x triplicate)	6°08S;80°40W (3 samples x triplicate)
Bayovar lagoons				
Zone 4 (March 2015)	A; AES; SE	6°23S;80°66W (3 samples x triplicate)	6°25S;80°26W (3 samples x triplicate)	6°23S;80°78W (3 samples x triplicate)

122   A= (saline water), AES= (water and sediment), SE= (sediment).

Fig. 2. Flowchart depicting the methodological strategy adopted in this study.

2.2 Isolation of cultivable actinomyces

An aliquot of 10.0 mL of saline lagoons sample was transferred to an Erlenmeyer containing 10.0 mL of salt broth (0.6% yeast extract, 2% glucose, 5% peptone, 3% meat extract supplemented with chloramphenicol and 1% fluconazole at pH 7.0). All Erlenmeyer were homogenized and maintained at 50 °C in a water bath for 60 min to reduce the pollutant load (Pisano et al. 1989 and Takizawa et al. 1993). Then, they were incubated at 28 °C for 7 to 30 days under aerobic conditions; after, the microorganisms were isolated in the media: modified saline agar (1.0 g yeast extract, 5.0 g peptone, and 15.0 g of agar at pH 7.6 ± 0.2) and trypticase soy agar (Bacto™) supplemented with saline water. The isolates were preserved in 20% glycerol and kept at -80 °C cryopreserved. The isolates were reactivated

and stored in a refrigerated chamber at -20 ° C to the transport to the Microbial Resources

Division of the Pluridisciplinary Center for Chemical, Biological and Agricultural Research

(CPQBA) in August 2015.

2.3 Identification of bacterial isolates

2.3.1 Reactivation and morphological phenotypic characterization of bacterial

strains

The reactivation of the isolates was carried out in R2A broth (Difco ref. 234000),

supplemented with artificial seawater (ASW) NaCl 5% (0.1 g potassium bromide, 23.48 g

sodium chloride, 10.61 g of magnesium chloride, 1.47 g of calcium chloride, 0.66 g of

potassium chloride, 0.04 g of strontium chloride, 3.92 g of sodium sulfate, 0.19 g of sodium

bicarbonate, 0.03 g of boric acid all in 1 L of distilled water) at 28 °C for 28 days, the colonies

were observed under a microscope and stereoscope for macro morphological characterization

and after Gram staining, the cells were observed under an optical microscope. The bacteria

were preserved in 20% glycerol and kept at -80 °C.

2.3.2 DNA extraction and 16S rRNA gene sequencing

An isolated colony was used for extraction of genomic DNA, according to the method

described by Pospiech and Neuman, 1995, with some modifications. Amplification of the

16S ribosomal RNA gene was done by means of a polymerase chain reaction (PCR) using

the pair of primers p10f (5' GAG TTT GAT CCT GGC TCA G 3') and p1401r (5' CGG

TGT GTA CAA GGC CCG GGA ACG 3') or p1492r (5'- ACC TTG TTA CGA CTT 3')

(Lane et al. 1985) homologous to conserved regions of the bacterial 16S ribosomal RNA

gene. Conditions used in the amplification reaction were: 2.5 μL of 10X buffer (without

Mg^{+2}), 0.75 μL of $MgCl_2$ (2.5 mM), 0.2 μL dNTP (2.5 mM each), 0.5 μL of each primer (20

μM), 0.2 μL of Taq DNA polymerase (Invitrogen) and 5.0 μL of the DNA template (50-100

ng/ μL) and deionized and sterilized water by filling the volume to 25 μL of reaction. The

PCR amplification was an initial denaturation step of 5 min at 95°C, followed by 30 cycles

of 1 min at 94°C for denaturation, 1 min at 55°C for annealing, 3 min at 72°C for extension

and 3 min at 72°C for final extension in an Eppendorf thermal cycler. The obtained products

were analyzed and visualized in agarose gel electrophoresis, purified with mini columns

(GFX PCR DNA & gel band purification kit, GE Healthcare), and sequenced in ABI3500XL

Series automatic sequencer (Applied Biosystems). Sequencing reactions were performed

with the Big Dye Terminator Cycle Sequencing Ready Reaction Kit (Applied Biosystems)

according to the manufacturer's specifications.

Partial sequences of the 16S ribosomal RNA gene obtained from each isolate were assembled

into a contig and then compared to the sequences of organisms represented in EZBioCloud

16S Database (https://www.ezbiocloud.net/) using the "Identify" service (Yoon et al. 2017),

and species assignment were based on closest hits (Jeon et al. 2014). 16S ribosomal RNA

gene sequences retrieved from the database and related to the unknown organism gene were

selected for alignment in the Clustal X program (Thompson et al. 1997), and phylogenetic

analyzes were performed using the Mega version 6.0 program (Tamura et al. 2013). The

evolutionary distance matrix was calculated with the model of Kimura-2 parameters

(Kimura, 1980), and the phylogenetic tree constructed from the evolutionary distances

calculated by the Neighbor-Joining method (Saitou and Nei, 1987), with bootstrap values

from 1000 resampling.

2.4 Activity approach of bacterial isolates

2.4.1 Crude extract producing extremophilic bacterial

Secondary metabolites of bacterial isolates were extracted with ethyl acetate. A pre-inoculum of the bacteria was performed in 5.0 mL of R2A Broth medium (Himedia ref. 1687) with AWS NaCl 5% and incubated at 28 °C for 5 to 7 days. After the growth of the culture, the total volume was transferred to an Erlenmeyer containing 500.0 mL of the same culture medium and incubated for 7 days under agitation at 150 rpm, followed by 23 days without agitation. After this period, 500.0 mL of ethyl acetate were added to the medium containing the bacterial growth and ruptured in ultra turrax basic (IKA ref. 02H2063.08.CC) at 7,000 rpm for 15 min. The organic fraction was obtained and the crude extract was concentrated in the rotary evaporator (R-215 Buchi), under vacuum and temperature of 40 °C, until the solvent was completely dried and stored at 4°C. The crude extracts of three strains representatives were taken for antimicrobial and antiproliferative activity tests. Simultaneously, the crude extracts also were analyzed by liquid chromatography coupled to the mass spectrometer (Versamax, molecular manufacturer Devices).

2.4.2 Crude extract fractionation of the extremophilic bacterial.

The crude extracts of the isolates were fractionated using a vacuum chromatographic column with C18. The initial crude extract was solubilized in methanol. The mobile phases used were fraction 1 water (H_2O); fraction 2 water: methanol (H_2O: MeOH 1:1 v/v); fraction 3 methanol (MeOH); fraction 4 MeOH: ethylacetate (EtOAc) (1:1 v/v); fraction 5 EtOAc 100%; fraction 6 EtOAc: Glacial acetic acid 1%. These fractions were dried in rotary evaporator at 45 °C,

206   under vacuum weighted and used for activity antibacterial and antiproliferative tests.

2.4.3 *In vitro* antibacterial activity assay

Three crude bacterial extracts (M-92, B-81 and B-146) were tested as antimicrobial producers

using the minimum inhibitory concentration (MIC) assay, following the protocol reported by

Siddharth and Rai 2018. The crude extracts partially diluted in 1% Dimethylsulfoxide

(DMSO) 1 mg/mL – 3.9 µg/mL, and sterile broth were added into pre-coated microbial

cultures, completed a total volume of 200 µL. The plate was incubated at 37 °C and room

temperature; the lowest concentration of extract, which completely inhibited the bacterial

growth was considered as MIC. Each biological assay was performed in triplicate. The

pathogenic bacteria used in this test were *Escherichia coli* ATCC 10536, *Staphylococcus*

aureus ATCC 6538, and *Acinetobacter baumannii* with code AC-972, the strain was not

acquired prospectively, the source of origin comes from the bank of MDRs isolated from a

patient with pneumonia from the UCI of the Hospital Regional Lambayeque, details such as

patient data linked to the sample were anonymized in order to obtain access

2.4.4 *In vitro* antiproliferative activity assay

This assay aimed to detect anticancer activities by evaluating antiproliferative action in

human tumor cells (Monks et al. 1991). *In vitro* tests with the crude extract of *Streptomyces*

sp. M-92, B-81, and B-146 on human tumor cell lines of different origins and a non-tumor

cell line. Human tumor cell lines U251 (glioma), UACC-62 (melanoma), MCF-7 (breast),

NCI-ADR/RES (ovarian expressing phenotype with multiple drugs resistance), 786-0

(renal), NCI-H460 (lung, non-small cells), PC-3 (prostate), OVCAR-03 (ovarian), K562

(leukemia) and a non-tumor cell line HaCaT (keratinocyte) were obtained from the National

Cancer Institute Frederick, Molecules (MD, USA). Stock cultures were grown in medium

containing 5 mL of RPMI 1640 (GIBCO BRL, Gaithers-Burg, MD, USA) supplemented with

5 % fetal bovine serum (FBS, GIBCO) at 37 °C with 5% CO_2. Penicillin: streptomycin (1000

µg·L−1:1000 U·L−1, 1 mL·L−1) was added to the experimental cultures. Cells in 96-well

plates (100 µL cells well−1) were exposed to the extracts in DMSO (Sigma-Aldrich) /RPMI

(0.25, 2.5, 25, and 250 µg·mL−1) at 37 °C, and 5% CO_2 in the air for 48 h. The final DMSO

concentration (0.2% in higher concentration) did not affect cell viability. Before (T_0) and

after (T_1) sample application, cells were fixed with 50% trichloroacetic acid (Merck), and

cell proliferation was determined by the spectrophotometric quantification (540 nm) of

cellular protein content using sulforhodamine B assay. Using the concentration-response

curve for each cell line, the values of the sample concentration required to produce total

growth inhibition or cytostatic effect (TGI) were determined through non-linear regression

analysis using software ORIGIN 8.6® (OriginLab Corporation, Northampton, MA, USA).

2.5 Mass Spectrometry analysis

Streptomyces sp. B-81 extract was resuspended in 1 mL of methanol (HPLC grade) and 100

µL were diluted in 900 µL of methanol. The final solution was filtered through 0.22 µm into

vials. Ultra-high pressure liquid chromatography-mass spectrometry (UHPLC-MS) analyses

were performed in a Thermo Scientific QExactive® Hybrid Quadrupole-Orbitrap Mass

Spectrometer. Parameters: positive mode, capillary voltage at +3.5 kV; capillary temperature

at 250 °C; S-lens of 50 V and *m/z* range of 133.40-2000.00. Tandem mass spectrometry

(MS/MS) was performed using normalized collision energy (NCE) of 30 eV and 5 precursors

per cycle were selected. Stationary phase: Thermo Scientific Accucore C18 2.6 μm (2.1 mm

x 100 mm) column. Mobile phase: 0.1% formic acid (A) and acetonitrile (B). Eluent profile

(A: B) 0-10 min, gradient from 95:5 up to 2:98; held for 5 min; 15-16.2 min gradient up to

95:5; held for 8.8 min. Flow rate: 0.2 mL min^{-1}. The injection volume of the samples was 3

μL. Operation and spectra analyses were conducted using Xcalibur software (version 3.0.63)

developed by Thermo Fisher Scientific.

2.6 Molecular MS/MS network

A molecular network for *Streptomyces sp*. B-81 was created using the online workflow at

GNPS (https://gnps.ucsd.edu/). The data were filtered by removing all MS/MS peaks within

± 17 Da of the precursor *m/z*. MS/MS spectra were window filtered by choosing only the top

6 peaks in the ± 50 Da window throughout the spectrum. The data was then clustered with

MS-Cluster with a parent mass tolerance of 0.02 Da and a MS/MS fragment ion tolerance of

0.02 Da to create consensus spectra. A network was then created where edges were filtered

to have a cosine score above 0.5 and more than 5 matched peaks. Further edges between two

nodes were kept in the network only if each of the nodes appeared in each other's respective

top 10 most similar nodes. The spectra in the network were then searched against GNPS'

spectral libraries. The library spectra were filtered in the same manner as the input data. All

matches kept between network spectra and library spectra were required to have a score

above 0.5 and at least 5 matched peaks (Wang et al. 2016).

3. Results and discussion

3.1 Isolation, identification and selective of bacterial species from the northern

saline lagoons of Perú

In total, 166 pure cultures showing different colony morphologies were obtained grown on

R2A medium from saline water (A), water and sediment (AES), and sediment (SE). The

morphological characteristics such as aerial mycelium, the morphological spore mass color,

pigmentation of vegetative or substrate mycelium, and the production of diffusible pigment

were used to classify 42 filamentous and 8 non filamentous to the genera *Streptomyces* 39

(78%), *Pseudonocardia* 3 (6%), *Staphylococcus* 4 (8%), *Bacillus* 2 (4%), and *Pseudomonas*

2 (4%) in Table 2, (Flores, 2017).

Table 2. 16S rRNA gene-based identification of bacteria isolate from lagoons Morrope

and Bayovar.

Sample type	Number of isolate (%)	Phylum	Family	Genus
AES, SE	39 (78%)	*Actinobacteria*	*Streptomycetaceae*	*Streptomyces*
SE	3 (6%)	*Actinobacteria*	*Pseudonocardiaceae*	*Pseudonocardia*
A	2 (4%)	*Firmicutes*	*Bacillaceae*	*Bacillus*
A	4 (8%)	*Firmicutes*	*Staphylococcaceae*	*Staphylococcus*
AES	2 (4%)	*Proteobacteria*	*Pseudomonadaceae*	*Pseudomonas*

287   The isolates were regrouped into 28 isolates due to their similar phenotypic characteristics

288   and were identified based on the sequencing and alignment, from the 16S rRNA gene

289   analyzes. They belong to the genus *Streptomyces* 27 isolates and 1 *Pseudonocardia* isolate,

290   and 23 isolates showed a 16S rRNA gene sequence similarity below the 98.7% threshold for

291   proposing as novel species (Table 3).

Table 3. Strain Actinobacterial isolate from different sample Bayovar and Mórrope saline lagoons, indicating the top hit taxon, sequences length nucleotides, and similarity percent of the isolates the "Identify" service best-hit classifier in the EZBioCloud 16S Database.

	EZBioCloud Hit Strain name	Top Hit taxon	16S rRNA sequences length (nt)	Similarity %
Mórrope Zona 1				
M-92	TRM 45540	*S. luteus*	1313	99.16
M-142	NBRC 15864	*S. labedae*	1440	97.58
Mórrope Zona 2				
M-239	TRM 45540	*S. luteus*	1472	99.19
M-239.2	TRM 45540	*S. luteus*	1508	99.19
M-261	LMG 19316	*S. griseoincarnatus*	1468	97.58
M-262	NBRC 12780	*S. griseorubens*	1440	97.57
M-266	NBRC 12825	*S. variabilis*	1345	97.58
M-270	NBRC 13433	*S. pactum*	1471	97.98
M-274.2	LMG 19316	*S. griseoincarnatus*	1471	97.58
M-275	NRRL B-3009	*S. olivaceus*	1380	97.98
M-282.1	NRRL B-3009	*S. olivaceus*	1412	97.98
M-285	NRRL B-3009	*S. olivaceus*	1507	97.98
M-289.1	NBRC 13433	*S. pactum*	1441	97.98
M-290	OU-40	*S. hyderabadensis*	1481	98.06
M-290.1	NRRL B-3009	*S. olivaceus*	1398	97.98
Mórrope Zona 3				
M-323	NRRL B-3009	*S. olivaceus*	1500	97.98
Bayovar Zona 4				
B-81	NRRL B-3009	*S. olivaceus*	1345	99.93

B-93.1	ISP 5183	*S. violascens*	1339	97.37
B-93.3	LMG 19316	*S. griseoincarnatus*	1498	97.58
B-95	LMG 19316	*S. griseoincarnatus*	1456	97.58
B-95.1	NBRC 15864	*S. labedae*	1359	97.58
B-236.1	YIM 63235	*Pseudonocardia antimicrobica*	1500	98.41
B-144.2	NRRL B-3009	*S. olivaceus*	1519	97.98
B-146	NRRL B-3009	*S. olivaceus*	1287	99.84
B-236	LMG 19316	*S. griseoincarnatus*	1516	97.58
B-244	NRRL B-3009	*S. olivaceus*	1463	97.98
B-265	LMG 19316	*S. griseoincarnatus*	1391	97.58
B-379	NBRC 13475	*S. hydrogenans*	1493	97.27

Therefore, these strains should be designed for further taxonomic and analytical chemistry

analyses to confirm their novelty at species rank and as a source of novel chemical entities.

Two isolates were identified in lagoon 1 of the Morrope salt flats, and according to the

EZBioCloud the isolate M-92 presented similarity with *Streptomyces luteus* TRM 45540

(99.16%), while the isolate M-142 presented similarity with *Streptomyces labedae* NBRC

15864 (97.58%). In lagoon 2 of the Morrope salt flats, 13 isolates were identified, of which

M-239 and M-239.2 were similar to *S. luteus* TRM 45540 (99.19%), the isolates M-261 and

M-274.2 presented similar to *S. griseoincarnatus* LMG 19316 (97.58%), the isolate M-262

showed similarity to *S. griseorubens* NBRC 12780 (97.57%), the isolate M-266 was similar

to *S. variabilis* NBRC 12825 (97.58%), the isolates M-270 and M- 289.1 were similar to *S.*

pactum NBRC 13433 (97.98%), the isolate M-290 presented similar to *S. hyderabadensis*

OU-40 (98.06%), and finally four isolates, M-275, M-282.2, M-285 and M- 290.1 were

similar to *S. olivaceus* NRRL B-3009 (97.98%). From the lagoon 3, also belongs to the

Morrope salt flats, it was identified one isolate M-323 similar to *S. olivaceus* NRRL B-3009

(97.98 %). Finally, from lagoon 4, belongs to the Bayovar salt flats, 12 isolates were

identified, four isolates were similar to *S. olivaceus* NRRL B-3009, B-81 (99.93%0, B-144.2

(97.98%), B-146 (99.84%), and B-244 (97.98%). Also, in the lagoon 4 the isolate B-93, was

similar to *S. violascens* ISP 5183 (97.37%), and other four isolates presented similar *S.*

griseoincarnatus LMG 19316, B-93.3, B-95, B-236, B-265 (97.58%), the isolate B-95.1 was

similar to *S. labedae* NBRC 15864 (97.58%), and the isolate B-379 was similar to *S.*

hydrogenans NBRC 13475 (97.27%). Besides Streptomyces, another of the actinobacteria

genus was recovered to the Bayovar salt flats. The isolate B-236.1 was similar to

Pseudonocardia antimicrobica YIM 63235 (97.27%) and our work is the first to report this

genus present in the Bayovar salt flats (Table 3).

In the Morrope salt flats, eight different groups of *Streptomyces* were identified, whereas in

the Bayovar saline only five groups of *Streptomyces* were recovered, however, a different

genus like *Pseudonocardia*, whichhas been reported by Zhang et al. (2016) as producers of

γ-Butyrolactones that previously had only been reported for Streptomyces genera and in

plants was recovered in this study (Table 3). Our culture-based approach using a pre-

enrichment to decrease the bacterial load agrees with the reports described in the literature,

in which the isolation of halophilic bacteria reveals low species richness and the dominance

of *Streptomyces* genus Parthasarathi et al. (2012). Ballav et al. (2015) reported *Streptomyces*

spp. as the most predominant group contributing to 46% of the total isolates in crystallizer

pond sediments of Ribandar saltern, Goa, India. In this study the isolation was performed

with three different types of culture media and four different concentrations of salt because

in lagoons were found the concentration of salts between 0-300 psu units (Ballav et al. 2015).

Cortes et al. (2019) in the Salar de Huasco report similar results, they were isolated

Streptomyces (86%), *Nocardiopsis*, (9%), *Micromonospora* (3%), *Bacillu*s (1%) and

Pseudomonas (1%).

The phylogenetic analysis based on their nearly complete 16S rRNA gene sequences (>1206

bp) was constructed with the isolates M-92, B-81 and B-146 because them presented

antibacterial and antiproliferative activities. The phylogenetic tree showed clades well

supported to the three isolates and the reference sequences (Figure 3).

Fig 3. Phylogenetic tree based on 16S rRNA gene sequences (1206 positions in the final

dataset) inferred using the neighbour-joining method in MEGA7, the evolutionary

distances were computed using the Kimura 2-parameter method, showing the

phylogenetic positions of different saline isolates and type strains within the genus

Streptomyces and Pseudonocardia. Numbers at branching points refer to percentages

of bootstrap values from 1000 replicates. The scale bar indicates 0.02 indicated substitutions per nucleotide. *Kitasospora kazusensis* SK60^T (NR_041539) was used as outgroup.

Isolate M-92 forms a distinct branch closely related to the type strain of *Streptomyces olivoverticulatus* NRRL-1994^T, *Streptomyces swartbergensis* HMC13^T, and *Streptomyces caelestis* NRRL-2418^T. Isolates B-81 and B-146 form a well-supported sub-clade closely associated to *Streptomyces olivaceus* NBRC 12805^T, *Streptomyces litmocidini* NRRL B-3635^T, and *Streptomyces pactum* NBRC 13433^T (Figure 3). The length of the branch of all *Streptomyces* isolates in the phylogenetic tree and the assignation to these isolates to completely different clades from each other, highlight the divergence of them from their closely related neighbors. Further studies need to be performed to confirm the right affiliations of these isolates to the novel species within the evolutionary radiation of the genus *Streptomyces*.

3.2 Antibacterial activity Screening

Minimum Inhibitory Concentration (MIC) assay was used to screen antibacterial metabolites producer strains against the indicator strains *Escherichia coli* ATCC 10536, *Staphylococcus aureus* ATCC 6538, and *Acinetobacter baumannii* AC-972 MDR; the crude extracts tested from the three extremophilic bacterial reported positive results against the pathogenic bacteria tested. The antimicrobial activity of the crude extract of *Streptomyces* sp. B-81 strain present the highest inhibitory activity with the lowest MIC with 7.82 µg/mL against the three pathogens, followed by *Streptomyces* sp. M-92 showed MIC of 7.82 µg/mL against *E. coli* ATCC 10536 and *S. aureus ATCC 6538*, and 15.63 µg/mL against *A. baumannii AC-972*

370 MDR. *Streptomyces* sp. B-146 showed MIC of 15.63 µg/mL against the pathogens tested. It

371 is important to highlight that all crude extracts of the strains presented low MIC in

372 comparison with the reference antibiotic (Table 4).

374 **Table 4. Minimum inhibitory concentration MIC antibacterial of three *Streptomyces* sp.**

375 **crude extracts, by broth dilution method.**

Isolates	Minimum Inhibition Concentration(ug/mL)			
	E. coli ATCC 10536	*S. aureus* *ATCC 6538*	*A. baumannii* *AC-972* MDR	Antibiotic Chloramphenicol
M-92	7.82	7.82	15.63	22,2
B-146	15.63	15.63	15.63	23,1
B-81	7.82	7.82	7.82	22,3

Similar results were reported in a marine *Streptomyces* sp. S2A with MIC 31.25 µg/mL

against *Klebsiella pneumoniae*, 15.62 µg/mL against *Staphylococcus epidermidis*,

Staphylococcus aureus, *Bacillus cereus, Escherichia coli* and *Micrococcus luteus* with 7.8

µg/mL (Saket Siddharth and Ravishankar R. Vittal, 2018). The crude extract of the

Streptomyces sp. YBQ59 strain against nine pathogens, obtaining MIC results between 10.5

to 22.5 µg/mL, their results being similar to those we obtained (Vu. H et al. 2018).

3.3 Antiproliferative activity Screening

The antiproliferative activity of the crude extracts of *Streptomyces* sp. M-92, B-81, and B-

146 against three human tumor cell lines were expressed as TGI (Total Growth Inhibition-

concentration that inhibited cell growth by 100%) (Table 5).

Table 5. Antiproliferative activity of three Streptomyces extracts against human

tumor cell lines.

Crude extract and positive control	TGI (µg/mL)[c]		
	U251	MCF-7	NCI-H460
Doxorubicin[a]	2.46	6.11	<0,025
M-92	5.06	7.96	9.33
B-146	5.41	8.03	5.06
Doxorubicin[b]	1.50	>25	1.80
B-81	0.57	0.80	0.61

391 Human tumor cell lines: U251 (glioma), MCF-7 (breast), NCI-H460 (lung of type non-small cells)
392 [a, b] Doxorubicin used as positive control;
393 [c] TGI Concentration that totally inhibited cell growth
394

According to Abreu et al. (2014), TGI values higher than 50 µg/mL represent inactive

samples. In comparison to the others *Streptomyces* tested and doxorrubicin, *Streptomyces* sp.

B-81 shown the best result against glioma (TGI =0,57 µg/mL), lung cancer (TGI =0,61

µg/mL) and breast cancer (TGI =0,80 µg/mL). Due to the result obtained with the crude

extract of *Streptomyces* sp. B-81 strain, the extracts from three different media were selected

for further fractionation, to have access to the metabolic profile. It was possible to verify that

the highest activity detected was in the extracts from the fermentation of *Streptomyces* sp. B-

81 in the R2A medium (Table 5), which inhibited the nine different types of tumor cell lines.

However, it is observed that the fractionation decreases the antitumor activity presented in

the crude extract and the putative active compounds are recovered in fractions 3 and 4.

Fraction 3 seemed to contain antiproliferative substances in lower concentration and was able

to inhibit U251 (glioma) (TGI = 38.3 µg/mL), OVCAR-03 (ovary) (TGI = 62.1 µg/mL), and

K562 (leukemia) (TGI = 81.5 µg/mL), and fraction 4 inhibited U251 (glioma) (TGI =

73.5 µg/mL) and UACC-62 (melanoma) (TGI = 89,4 µg/mL) (Table 6).

Table 6. Antiproliferative evaluation of isolate *Streptomyces* sp. B-81 fermented in three

different medium and fractions of the R2A medium against nine tumors and one non-

tumor cell line.

	TGI (µg/mL)									
Crude extract	**2**	**u**	**m**	**a**	**7**	**4**	**o**	**h**	**k**	**q**
Doxorubicin	0,03	0,76	>25	>25	0,22	>25	14,4	>25	6,3	>25
B-81 NB	70,9	250	*	*	*	*	*	*	*	*
B-81 ISP2	14,1	17,2	*	155,9	*	38,9	55,2	155,7	*	*
B-81 R2A	1,6	21,3	8,0	31,0	39,0	11,2	19,4	89,7	11,6	158,1
B-81 R2A fractions	**2**	**u**	**m**	**a**	**7**	**4**	**o**	**h**	**k**	**q**
Fraction 1	*	*	*	*	*	*	*	*	*	*
Fraction 2	*	*	*	*	*	*	*	*	*	*
Fraction 3	38,3	127,2	154,6	189,8	172,6	108,0	62,1	167,4	81,5	*
Fraction 4	73,5	89,4	184,5	143,2	*	235,1	81,6	195,2	100,4	*
Fraction 5	*	*	*	*	*	*	*	*	*	*
Fraction 6	163,0	*	*	*	*	*	*	*	*	*

B-81 NB: *Streptomyces sp*. Crude extract in nutrient broth growth medium. B-81 ISP2: *Streptomyces sp*. Crude
extract in yeast extract-malt. B-81 R2A: *Streptomyces sp*. Crude extract in R2A growth medium. B-81 R2A
fractions: 1 = H_2O; 2 = MeOH-H_2O; 3 = MeOH; 4 = MeOH-EtOAc; 5 = 100% EtOAc; 6 = EtOAc -acid: Human
tumor lines: 2 = U251 (glioma), u = UACC-62 (melanoma); m = MCF-7 (breast); a = NCI-ADR / RES (ovary
with multiple drug resistance phenotype); 7 = 786-O (kidney adenocarcinoma); 4 = NCI-H460 (lung, non-small
cell type); o = OVCAR-03 (ovary); h = HT-29 (colon); k = K562 (leukemia). Human non-tumor lineage: q =
HaCaT (keratinocyte).
Doxorubicin (Reference chemotherapy).
TGI: Total Growth Inhibition- Concentration necessary to completely inhibit cell proliferation.
* TGI >250 µg/mL

It is common to find this antiproliferative activity in the *Streptomyces* genus due to its

metabolic and genetic capacity to produce secondary metabolites (Tan and Liu, 2017).

Additionally, actinomycetes isolated from marine environments have a greater capacity to

express these metabolites in response to the pressure of environmental selection to which

they have been subjected throughout the evolutionary history of aquatic organisms (Lam,

2006). This pressure has generated high specificity and a complex three-dimensional

conformation of the compounds to act in the marine environment (Song et al, 2018). In this

particular case, the geographical isolation, the geological conditions of formation of the

saline lagoon and the extreme environmental conditions generate a greater selection

environmental pressure, which makes it possible to observe antibiotic and antiproliferative

activity in the isolates analyzed.

The anticancer potential of halophilic actinomycetes has been extensively studied. Secondary

metabolites Salternamides A-D isolated from halophilic actinomycetes of saline on the island

of Shinui (Republic of Korea), have potential cytotoxicity against the human colon cancer

cell line (HCT116) and gastric cancer cell line (SNU638) (Kim et al. 2015). Another

moderately halophilic *Streptomyces sp*. W1 with number accession GenBank JN187420.1,

isolated from Weihai Solar Saltern in China, has potent cytotoxicity against human lung

cancer cell line (A549) with IC_{50} 78.6 µM; human cervical epithelial cell line (HeLa) IC_{50}

56.6 µM, human liver cancer cell line (BEL-7402) with IC_{50} 47.1 µM, and human colon

cancer cell line (HT-29) with IC_{50} 94.3 µM (Liu et al. 2013). Four compounds called

shellmicin A-D isolated from the *Streptomyces* sp. shell-016 were described by (Yong Han

et al. 2020) and studied concerning cytotoxicity against five human tumor cells lines: non-

small cell lung cancer (H1299, ATCC CRL-5803), malignant melanoma (A375, ATCC CRL-

1619), hepatocellular carcinoma (HepG2, ATCC HB-8065), colorectal adenocarcinoma

(HT29 ATCC HTB-38) and breast cancer (HCC1937, ATCC CRL-2336). The study

indicated that the compounds Shellmicin A, B and D showed greater cytotoxicity than

Shellmicin C compound, with an IC_{50} ranging from 0.69 µM to 3.11 µM at 72 h. An

interesting fact was that shellmicin C and D are a pair of stereoisomers and their biological

activity was significantly different. It is worth mentioning that in our study we determined

that the activity of the crude extract was better than that of the fractions, the same ones that

presented a different activity, therefore, we can deduce that we would be facing compounds

with synergism.

An actinomycete strain designated as *Streptomyces* sp. KML-2 isolated from a saline soil

mine in Khewra, Pakistan, showed antitumor activity against Hela cells (cervical cancer) and

MCF-7 (breast cancer) cell lines with IC_{50} values of 8.9 and 7.8 µg/mL respectively in the

initial screening, then they carried out a higher production (20 L) for the purification of the

compounds, confirming a wonderful anti-tumor potential against MCF-7 cells with an IC_{50}

value of only 0.97 µg /mL (Usman et al. 2015). This study also revealed that the Khewra salt

mine from which the KML-2 strain was isolated is a powerful ecological niche with

inimitable strain diversity that has yet to be discovered, a fact that reinforces the approach of

our study in the search for new compounds with biotechnological activity.

3.4 Secondary Metabolite Analysis of *Streptomyces sp*. B-81 isolate.

The crude extract obtained from *Streptomyces* sp. B-81 was investigated through UHPLC-

MS/MS analyses and demonstrated a broad profile of biomolecules (Fig. S1A). Furthermore,

a metabolite screening in the GNPS platform was performed and molecular networking

revealed two clusters (A and B) that exhibited secondary metabolites produced by

Streptomyces sp. B-81 extract (pink) (Fig. 4).

Fig 4. Molecular networking obtained for *Streptomyces* sp. B-81 extract. Cholic acid and

derivatives were grouped in cluster A, and lobophorins were grouped in cluster B.

Nodes circled in blue indicate molecules identified by comparison with the GNPS

platform database.

Metabolites were identified as a hit in the GNPS database or manually identified by accurate

mass analyses, which showed mass errors below 5 ppm (Table 7).

Table 7. MS data obtained for secondary metabolites detected in *Streptomyces* sp. B-81

extract.

Compound	Ion formula	Calculated *m/z*	Experimental *m/z*	Error (ppm)
Cholic acid	$C_{24}H_{44}NO_5$	426.3219	426.3212	-1.6
Lobophorin A	$C_{61}H_{93}N_2O_{19}$	1157.6372	1157.6373	0.1
Lobophorin B	$C_{61}H_{91}N_2O_{21}$	1187.6114	1187.6108	-0.5
Lobophorin E	$C_{61}H_{91}N_2O_{20}$	1171.6165	1171.6160	-0.4
Lobophorin K	$C_{61}H_{93}N_2O_{20}$	1173.6322	1173.6321	-0.1
Compound 6	$C_{11}H_{17}O_3$	197.1177	197.1173	-2.0

The observed signals corresponded to cholic acid (1), lobophorin A (2), lobophorin B (3),

lobophorin E (4) and lobophorin K (5).

In cluster A, GNPS database indicated the presence of cholic acid in the *Streptomyces* sp. B-

81 extract (Fig. S2 and S3). In the molecular networking, each MS/MS spectra are

represented by nodes, which are grouped in clusters based on their fragmentation pattern

similarity. Thus, compounds of the same class of molecules are grouped in the same cluster

(Wang et al. 2016). As observed in Figure 4, the cholic acid cluster is composed by a range

of nodes, suggesting the presence of other analogues. Cholic acid and other bile acids have

been reported as secondary metabolites of specific members of the genus *Streptomyces* and

Myroides (Kim et al. 2007). In the literature, cholic acid-derivatives and bile acids are known

as antimicrobials agents, displaying activity against gram-negative and positive bacteria or

improving the antimicrobial effect of antibiotics (Schimidt et al. 2001, Rasras et al 2010,

Darkoh et al. 2010, Savage et al. 2000).

In cluster B, GNPS database indicated the production of lobophorin A by *Streptomyces* sp.

B-81 strain (Fig. S4 and S5). Also, MS/MS profile of compound **2** presented *m/z* 883.49 and

753.43 (Fig. S5), related to the loss of the sugar units in the structure, according with the

literature (Nguyen et al. 2020). Similar as observed in cluster A, cluster B also exhibited

other lobophorin compounds such as lobophorins B, E and K that were identified based on

their accurate masses and fragmentation profiles with typical fragments at *m/z* 184.09,

157.09, 108.08 and 97.06 (Fig. S6-S8). Lobophorins were discovered by Jiang et. al. (1999)

in the expedition to Belize on board of the Columbus research ship. In the occasion, a new

strain of actinomycete, named as # CNC-837, was isolated from the surface of a brown algae

from the Caribbean called *Lobophora variegata*, and was reported to produce lobophorins A

and B, potent anti-inflammatory agents (Jiang et al. 1999). Besides lobophorins A and B,

other compounds of this class have been reported in the literature. Lobophorins C and D have

similar structures to A and B, while Lobophorin C exhibited potent cytotoxic activity against

human liver cancer cells, lobophorin D displayed significant inhibitory effect on human

breast cancer cells (Wei et al. 2011). Lobophorins E and F were reported by Niu et al. 2011,

that also described antibacterial and cytotoxic activities for lobophorin F. In other study, Pan

et al. 2013 reported lobophorins H and I and described that lobophorin H exhibited

antibacterial activities against *Bacillus subtilis*. Most recent, Lobophorin K was reported

(Braña et al. 2017) and also displays antimicrobial and cytotoxic activities.

Besides cholic acid and lobophorins, a furan-type compound (6) were also detected in the

Streptomyces sp. B-81 extract (Fig. S9). The structures of the metabolites **1-6** are shown in

Fig. 5.

R_1 = NH$_2$, R_2 = CH$_2$OH
R_1 = NO$_2$, R_2 = CH$_2$OH
R_1 = NO$_2$, R_2 = CH$_3$
R_1 = NHOH, R_2 = CH$_2$OH

Fig 5. Structures of secondary metabolites identified in *Streptomyces* sp. B-81 extract.

Compound 6 exhibited fragments at m/z 168.07, 167.07 and 153.05, in agreement with the

literature (Nguyen et al. 2020). Compound 6 was reported as a secondary metabolite of the

Streptomyces sp. VN1 strain, a microorganism isolated of sea sediment from a coastal region

of Viet Nam (Nguyen et al. 2020). In addition, *Streptomyces sp*. VN1 was also described to

produce lobophorin A (Nguyen et al. 2020). Nguyen et al 2020, reported that compound 6

displayed cytotoxic activity, *in vitro*, against 5 types of tumor cell lines. Furthermore, Wang

et al. 2008 reported that other furan compound named HS071, produced by *Streptomyces sp*.

HS-HY-071, displayed anticancer activity.

Our data deserves further investigation in order to correlate the antimicrobial and cytotoxic

activities, observed for the *Streptomyces* sp. B-81 extract, to the bile acids, lobophorins and

furan compounds. In addition, the extremophilic environments, in this particular case the

Salinas de Bayovar, are prolific sources of microorganisms, such as unique *Streptomyces*.

Due the extreme conditions, these microorganisms produce different compounds in order to

adapt to survive in the environment, which makes *Streptomyces* sp. B-81 an attractive source

of bioactive compounds.

Conclusion

The present study was successful in determining the diversity and bioactive potential of the

actinobacterial isolates present in this type of environment (Mórrope and Bayovar salt flats)

located in northern Peru, which are environments that had not been explored for this type of

study. this being the first study. Furthermore, it has been found that the isolates of

actinobacetria *Streptomyces sp*. M-92, B-146 and B-81 may be new species, their importance

for these isolates lies in their antibacterial and antiproliferative potential, being the most

promising for their best activity *Streptomyces sp*. B-81. Six biomolecules (colic acid,

Lobophorin A, B, E and K in addition to a sixth compound) were detected in this cultivable

actinobacterial isolated from Bayovar's salt flats, but not yet named from the cured base

GNPS. With this study it is determined that the actinobacteria that live in these environmental

conditions present an enormous wealth of microorganisms that are new species that produce

new and biologically active compounds and contribute to the study of extreme environments

such as the saline lagoons of northern Peru with antibacterial potential. and antiproliferative.

Further studies will be carried out in the identification of these molecules in order to purify

and characterize them since this can result in the economic production of bioactive

compounds for future pharmaceutical applications.

Acknowledgements

At the research agency of Peru (Concytec-Fondecyt) financially supported this work within

the framework of the 041-01 call with the 190-2018 contract; as well as the National Council

for Scientific and Technological Development of Brazil (CNPq), it was the authors are

grateful to the cooperation of the postgraduate programs in Genetics and Molecular Biology

and Biosciences of the Biology Institute of the State University of Campinas. Also,

Coordination for the Improvement of Higher Education Personnel - Brazil (CAPES) -

(Scholarship-code 001). We thank the research group of the Pluridisciplinary Research

Center for Chemistry, Biology and Agriculture to the laboratory infrastructure. Finally, we

are also grateful to the anonymous reviewers whose constructive criticisms have significantly

improved the quality of the manuscript.

References

1. Abreu PA, Sousa TS, Jimenez PC, Wilke DV, Rocha DD, Freitas HP, Pessoa OD, La Clair JJ, Costa-Lotufo LV. Identification of pyrroloformamide as a cytokinesis modulator. Chembiochem. 2014;15:501–506.

2. Ballav S, Kerkar S, Thomas S, Augustine N. Halophilic and halotolerant actinomycetes from a marine saltern of Goa, India producing antibacterial metabolites. Journal of bioscience and bioengineering. 2015;119(3):323-330.

3. Berdy J. Bioactive microbial metabolites. Journal of Antibiotics. 2005;58:1-26.

4. Binayke A, Ghorbel S, Hmidet N, Raut A, Gunjal A, Uzgare A, Patil N, Waghmode M, Nawani N. Analysis of diversity of actinomycetes from arid and saline soils at Rajasthan, India. Environmental Sustainability. 2018;1(1):61-70.

5. Boyadzhieva S, Georgieva S, Angelov G. Optimization of the extraction of natural antioxidants from avocado seeds. Bulg Chem Commun. 2018;50:80-84.

6. Braña AF, Vizcaíno AS, Osset M, Victoria IP, Martín J, Pedro DN, Mercedes de la Cruz M, Díaz C, Vicente J, Reyes F, García LA, Blanco G. Lobophorin K a new natural product with cytotoxic activity produced by *Streptomyces sp*. M-207, associated with the deep-sea coral *Lophelia pertusa*. Marine Drugs. 2017;15(5):144.

7. Blunt WJ, Cop BR, Hu WP, Munro MH, Northcote PT, Prinsep MR. Marine natural products. This journal is the Royal Society of Chemistry Nat. Prod. Rep. 2007;24:31–86.

8. Cáceres MD and Legendre P. Associations between species and groups of sites: indices and statistical inference. Ecology. 2009; 90(12):3566-3574.

9. Cortés-Albayay C, Silber J, Imhoff JF, Asenjo JA, Andrews B, Nouioui I, Dorador C. The

polyextreme ecosystem, salar de Huasco at the chilean altiplano of the Atacama desert

houses diverse *Streptomyces spp*. with promising pharmaceutical potentials. Diversity.

2019;11(5):69.

10. Rene Flores Clavo, Molecular characterization of bacteria the lagoons of northern and

evaluation of potential of new biomolecules against different types of tumor cells. 2017.

1 recurso online (201 p.). Dissertação (mestrado) - Universidade Estadual de Campinas,

Instituto de Biologia, Campinas, SP.

11. Darkoh C, Lichtenberger LM, Ajami N, Dial EJ, Jiang ZD, DuPont HL. Bile acids

improve the antimicrobial effect of rifaximin. Antimicrobial agents and chemotherapy.

2010;54(9):3618-3624.

12. Fiedler HP, Bruntner C, Bull AT, Ward AC, Goodfellow M, Potterat O, Puder C, Mihm

G. Marine actinomycetes as a source of novel secondary metabolites. Antonie Van

Leeuwenhoek. 2005;87:37-42.

13. Flores CR, Pisfil CE, Aislamiento, caracterización y actividad antibacteriana de

actinomicetos halófilos de las salinas de mórrope frente a *Staphylococcus aureus*

meticilino-resistente y *Escherichia coli* productor de betalactamasas de espectro

extendido (BLEE). Course completion work - Thesis, Pedro Ruiz Gallo University

Lambayeque, Peru (http://repositoro.unprg.edu.pe/handle/UNPRG/791). 2014;1-77

14. Han Y, Wang Y, Yang Y, Chen H. Shellmycin A–D novel bioactive Tetrahydroanthra-

γ-Pyrone antibiotics from marine *Streptomyces sp*. Shell-016. Marine Drugs. 2020;

18(1):58.

15. Hayakawa Y, Shirasaki S, Shiba S, Kawasaki T, Matsuo Y, Adachi K, Shizuri Y.

Piericidins C7 and C8, new cytotoxic antibiotics produced by a marine *Streptomyces sp*.

J. Antibiot. 2007;60:196-200.

16. Jeon YS, Lee K, Park SC, Kim BS, Cho YJ, Ha SM, Chun J. A versatile sequence

alignment editor for both rRNA and protein coding genes. International journal of

systematic and evolutionary microbiology. 2014;64(2):689-691.

17. Jiang ZD, Jensen PR, Fenical W. Lobophorins A and B, new anti-inflammatory

macrolides produced by a tropical marine bacterium. Bioorganic and Medicinal

Chemistry Letters. 1999;9(14):2003-2006.

18. Kim D, Lee JS, Kim J, et al. Biosynthesis of bile acids in a variety of marine bacterial

taxa. J Microbiol Biotechnol. 2007;17(3):403407.

19. Kim SH, Shin Y, Lee SH, Oh KB, Lee SK, Shin J, Oh DC. Salternamides A–D from a

halophilic *Streptomyces sp*. Actinobacterium. J. Nat. Prod. 2015;78:836–843.

20. Kim K, Samaddar S, Chatterjee P, Krishnamoorthy R, Jeon S, Sa T. Structural and

functional responses of microbial community with respect to salinity levels in a coastal

reclamation land. Applied soil ecology. 2019;137:96-105.

21. Kimura M. A simple model for estimating evolutionary rates of base substitutions

through comparative studies of nucleotide sequences. Journal of Molecular Evolution.

1980;16:111-120.

22. Lane DJ, Pace B, Olsen GJ, Stahl DA, Sogin ML, Pace NR. Rapid determination of 16S

ribosomal RNA sequences for phylogenetic analysis. Proceedings of the National

Academy of Sciences. 1985;82:6955-6959.

23. Lam Kim S. Discovery of novel metabolites from marine actinomycetes. Cur Opine

Microbiol. 2006;9:245–251.

24. Leon J, Aponte JJ, Cuadra DL, Galindo N, Jaramillo L, Vallejo M, Marguet E.

Extracellular enzymes production and pathogen inhibitory activity of actinomycetes

isolated from *Argopecten purpuratus*. Journal of Biology Marin and Oceanographic.

2016;51(1):69-80.

25. Liu H, Xiao L, Wei J, Schmitz JC, Liu M, Wang C, Lin X, et al. Identification of

Streptomyces sp. nov. WH26 producing cytotoxic compounds isolated from marine solar

saltern in China. World journal of microbiology and biotechnology. 2013;29(7):1271-

1278.

26. Lubsanova DA, Zenova GM, Kozhevin PA, Manucharova NA, Shvarov AP. Filamentous

Actinobacteria of the saline soils of arid territories. Moscow University soil science

bulletin. 2014;69(2):88-92.

27. Maldonado LA, Stach JE, Pathom-aree W, Ward AC, Bull AT, Goodfellow M. Diversity

of cultivable actinobacteria in geographically widespread marine sediments. Antonie Van

Leeuwenhoek. 2005;87(1):11-18.

28. Monks A, Scudiero D, Skehan P, Shoemaker R, Paull K, Vistica D, Gray-Goodrich M.

Feasibility of a high-flux anticancer drug screen using a diverse panel of cultured human

tumor cell lines. Journal of the National Cancer Institute. 1991;83(11):757-766.

29. Newman DJ, Cragg GM. Natural products as sources of new drugs over the last 25 years.

J Nat Prod. 2007;70:461-477.

30. Niu S, Li S, Chen Y, Tian X, Zhang H, Zhang G, Zhang W, Yang X, Zhang S, Ju J, Zhang

C. Lobophorins E and F, new spirotetronate antibiotics from a South China Sea-derived

Streptomyces sp. SCSIO 01127. The Journal of Antibiotics. 2011;64:711-716

31. Nguyen HT, Pokhrel AR, Nguyen CT, Dhakal D, Lim HN, Jung HJ, Sohng JK, et al.

Streptomyces sp. VN1, a producer of diverse metabolites including non-natural furan-type

anticancer compound. Scientific reports. 2020;10(1):1-14.

32. Oueriaghli N, Castro DJ, Llamas I, Béjar V, Martínez-Checa F. Study of bacterial

community composition and correlation of environmental variables in Rambla Salada, a

hypersaline environment in South-Eastern Spain. Frontiers in microbiology. 2018;9:

1377.

33. Pan HQ, Zhang SY, Wang N, Li ZL, Hua HM, Hu JC, Wang SJ. New spirotetronate

antibiotics, lobophorins H and I, from a South China Sea-derived Streptomyces sp.

12A35. Marine drugs. 2013;11(10):3891-3901.

34. Parthasarathi S, Sathya S, Bupesh G, Samy RD, Mohan MR, Kumar GS, Manikandan M,

Kim CJ, Balakrishnan K. Isolation and characterization of antimicrobial compound from

marine *Streptomyces hygroscopicus* BDUS 49. World J. Fish Mar. Sci. 2012;4:268-277.

35. Pisano MA, Sommer MJ, Brancaccio L. Isolation of bioactive actinomycetes from marine

sediments using rifampicin. Applied microbiology and Biotechnology. 1989;31(6):609-

612.

36. Pospiech A, Neumann B. A versatile quick-prep of genomic DNA from Gram-positive

bacteria. Trends Genetics. 1995;11(6):217-218.

37. Rasras AJ, Al-Tel TH, Al-Aboudi AF, Al-Qawasmeh RA. Synthesis and antimicrobial

activity of cholic acid hydrazone analogues. European journal of medicinal chemistry.

2010;45(6):2307-2313.

38. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing

phylogenetics trees. Mol. Biol. Evol. 1987;21:105-121.

39. Savage PB, Li C. Cholic acid derivatives: novel antimicrobials. Expert opinion on

investigational drugs. 2000;9(2):263-272.

40. Siddharth S, Vittal RR. Evaluation of antimicrobial, enzyme inhibitory, antioxidant and

cytotoxic activities of partially purified volatile metabolites of marine *Streptomyces sp*.

S2A. microorganisms. 2018;6:72.

41. Siddharth S, Vittal RR, Wink J, Steinert M. Diversity and Bioactive Potential of

Actinobacteria from Unexplored Regions of Western Ghats, India. Microorganisms.

2020;8(2):225.

42. Song X, Xiong Y, Qi X, Tang W, Dai J, Gu Q, Li J. Molecular targets of active anticancer

compounds derived from marine sources. Marine drugs. 2018;16(5):175.

43. Schmidt E J, Boswell JS, Walsh JP, Schellenberg MM, Winter TW, Li C, Savage PB, et

al. Activities of cholic acid-derived antimicrobial agents against multidrug-resistant

bacteria. Journal of Antimicrobial Chemotherapy. 2001;47(5):671-674.

44. Strohl WR. Antimicrobials. In Microbial diversity and bioprospecting. American Society

of Microbiology. 2004;336-355.

45. Tamura K, Stecher G, Peterson, Filipski A, Kumar S. MEGA6: Molecular Evolutionary

Genetics Analysis Version 6.0. Mol Biol Evol. 2013;2725-2729.

46. Takizawa M, Colwell RR, Hill RT. Isolation and Diversity of Actinomycetes in the

Chesapeake Bay. Applied and environmental Microbiology. 1993;997-1002.

47. Tan GY, Liu T. Rational synthetic pathway refactoring of natural products biosynthesis

in actinobacteria. Metabolic engineering. 2017;39:228-236.

48. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. The Clustal_X

windows interface: flexible strategies for multiple sequence alignment aided by quality

analysis tools. Nucleic Acids Res. 1997;25:4876-4882.

49. Aftab U, Zechel DL, Sajid, I. Antitumor compounds from Streptomyces sp. KML-2,

isolated from Khewra salt mines, Pakistan. Biological research. 2015;48(1):58.

50. Valliappan K, Sun W, Li Z. Marine actinobacteria associated with marine organisms and

their potentials in producing pharmaceutical natural products. Applied microbiology and

biotechnology. 2014; 98(17):7365-7377.

51. Vu HN, Nguyen DT, Nguyen HQ, Chu HH, Chu SK, Van Chau M, Phi QT.

Antimicrobial and cytotoxic properties of bioactive metabolites produced by *Streptomyces*

cavourensis YBQ59 isolated from *Cinnamomum cassia* Prels in Yen Bai Province of

Vietnam. Current microbiology. 2018;75(10):1247-1255.

52. Wang J, Zhang H, Yang X, Zhou Y, Wang H, Bai H. HS071 a new furan-type cytotoxic

metabolite from *Streptomyces sp*. HS-HY-071. The Journal of antibiotics.

2008;61(10):623-626.

53. Wang M, Carver JJ, Phelan VV, Sanchez LM, Garg N, Peng Y, Porto C, et al. Sharing

and community curation of mass spectrometry data with Global Natural Products Social

Molecular Networking. Nature biotechnology. 2016;34(8):828-837.

54. Wei RB, Xi T, Li J, Wang P, Li FC, Lin YC, Qin S. Lobophorin C and D, new kijanimicin

derivatives from a marine sponge-associated actinomycetal strain AZS17. Marine drugs.

2011;9(3):359-368.

55. Yoon SH, Ha SM, Kwon S, Lim J, Kim Y, Seo H, Chun J. Introducing EzBioCloud: a

taxonomically united database of 16S rRNA gene sequences and whole-genome

assemblies. International journal of systematic and evolutionary microbiology.

2017;67(5):1613.

56. Zhang XM, Zhang DF, Li WJ, Lu CH. Pseudonocardides A–G, New c-Butyrolactones

from Marine-derived *Pseudonocardia sp*. YIM M13669. Helv. Chim. Acta. 2016;99:191-

196.