The *Trypanosoma brucei* Cytoskeletal Protein KHARON

Associates with Partner Proteins to Mediate Both

Cytokinesis and Trafficking of Flagellar Membrane Proteins

Marco A. Sanchez, and Scott M. Landfear*

Department of Molecular Microbiology & Immunology, Oregon Health & Science

University, Portland, Oregon, USA

16 *Author for correspondence (landfear@ohsu.edu)

19 **Summary Statement:** This study investigates the essential role in African

20 trypanosomes of the KHARON protein and its molecular partners in trafficking of

21 membrane proteins to the flagellum.

ABSTRACT

In the African trypanosome *Trypanosoma brucei*, the cytoskeletal protein *Tb*KHARON is required for trafficking of a putative Ca^{2+} channel to the flagellar membrane, and it is essential for parasite viability in both the mammalian stage bloodstream forms and the tsetse fly procyclic forms. This protein is located at the base of the flagellum, in the pellicular cytoskeleton, and in the mitotic spindle in both life cycle forms, and it likely serves multiple functions for these parasites. To begin to deconvolve the functions of KHARON, we have investigated partners associated with this protein and their roles in parasite biology. One KHARON associated protein, *Tb*KHAP1, is a close interaction partner that can be crosslinked to KHARON by formaldehyde and pulled down in a molecular complex, and it colocalizes with *Tb*KHARON in the basal body at the base of the flagellum. Knockdown of *TbKHAP1* mRNA has similar phenotypes to knockdown of its partner *Tb*KHARON, impairing trafficking of the Ca^{2+} channel to the flagellar membrane and blocking cytokinesis, implying that the *Tb*KHARON/*Tb*KHAP1 complex mediates trafficking of flagellar membrane proteins. Two other KHAPs, *Tb*KHAP2 and *Tb*KHAP3, are in close proximity to *Tb*KHARON, but knockdown of their mRNAs does not affect trafficking of the Ca^{2+} channel. Two different flagellar membrane proteins, which are extruded from the flagellar membrane into extracellular vesicles, are also dependent upon *Tb*KHARON for flagellar trafficking. These studies confirm that *Tb*KHARON acts in complexes with other proteins to carry out various biological functions, and that some partners are involved in the core activity of targeting membrane proteins to the flagellum.

KEYWORDS *Trypanosoma brucei*, cytoskeleton, cytokinesis, flagellar membrane protein trafficking, protein complexes, virulence

INTRODUCTION

African trypanosomes of the species *Trypanosoma brucei* are parasitic protists that cause human African trypanosomiasis and the disease nagana in cattle and are thus of great medical and veterinary importance (Kennedy, 2013). In addition, these parasites have been recognized as valuable models for probing fundamental questions in cell and molecular biology (Cayla et al., 2019). *T. brucei* and related kinetoplastid parasites such as *Trypanosoma cruzi* and *Leishmania* species are flagellated and offer novel insights into the structure and function of flagella (Langousis and Hill, 2014) and the roles of these organelles in infection (Kelly et al., 2020a). Furthermore, the cell division cycle of African trypanosomes has been studied extensively (Farr and Gull, 2012; Vaughan and Gull, 2008; Wheeler et al., 2019), identifying various processes that are important for both proliferative and differentiation-linked cell division.

In previous work on trafficking of integral membrane proteins to flagella, we identified a kinetoplastid-specific protein designated KHARON (KH) that plays a critical role in flagellar targeting of the putative Ca^{2+} channel *Tb*CaCh (Tb927.10.2880) in *T. brucei*, originally identified as a flagellar surface protein FS179 (Oberholzer et al., 2011), and the flagellar glucose transporter *Lmx*GT1 in *L. mexicana* (Tran et al., 2013) (see Table S1 for tabulation of gene IDs and names of *T. brucei* proteins investigated in this study). In both species of parasite (Sanchez et al., 2016), KHARON was localized to three distinct subcellular compartments: the base of the flagellum (Fig. 1A), the subpellicular microtubules that subtend the plasma membrane around the cell body, and the mitotic spindle (Fig. 1B). Application of RNA interference (RNAi) to knock down *TbKH* (Tb927.10.8940) revealed that, in addition to preventing flagellar trafficking of *Tb*CaCh/FS179, the flagellum attachment zone (FAZ) was disrupted, resulting in detachment of flagella from the cell body (Sanchez et al., 2016), leaving this organelle adherent only through its connection at the flagellar pocket. Interfering with trafficking of *Tb*CaCh/FS179 to the flagellar membrane is likely to induce disruption of flagellar attachment, as RNAi directed against this channel also results in a similar flagellar detachment phenotype (Oberholzer et al., 2011). In addition, as found for many genetic alterations that disrupt flagellar attachment, these parasites are unable to initiate cell division, generating trypanosomes in which nuclei, kinetoplasts (mitochondrial DNA-containing structures), basal bodies, and flagella have replicated but cytokinesis has not occurred. This phenotype was apparent in both mammalian bloodstream form (BF) and insect stage procyclic form (PF) parasites and was thus lethal to both life cycle stages.

Similar studies in *L. mexicana* have also established a critical role for *Lmx*KH (LmxM.36.5850) in the life cycle of *Leishmania* parasites. Thus a Δ*lmxkh* null mutant was

generated in insect stage promastigotes, where trafficking of *Lmx*GT1 to the flagellum was
strongly impaired (Tran et al., 2013), but cell division and replication of this life cycle stage was
not affected. In contrast, Δ*lmxkh* null mutants were unable to undergo cytokinesis after invading
host macrophages, resulting in formation of multinucleate multiflagellated amastigotes that died
over the course of several days (Tran et al., 2013). These null mutants were also avirulent
following injection into BALB/c mice (Tran et al., 2015), and studies by others have shown that
KHARON null mutants in *L. infantum* have potential as a live attenuated vaccine (Santi et al.,
2018).

KHARON exhibits both similarities and striking differences between *T. brucei* (*Tb*KH) and *L.*
mexicana (*Lmx*KH). Thus, the two orthologs are relatively divergent in sequence, sharing 27%
amino acid identity and differing significantly in length (411 amino acids for *Tb*KH versus 520
amino acids for *Lmx*KH). *Tb*KH is critical for cell division of both mammalian BF and insect
stage PF parasites, whereas *Lmx*KH is only essential for division of disease-causing
amastigotes. Nonetheless, the three subcellular locations for KHARON are shared between the
two parasites, as are functions in cytokinesis and formation of the flagellar membrane.

KHARON proteins do not share significant sequence similarity to proteins outside the
Kinetoplastida, nor do they contain conserved sequence motifs that are suggestive of specific
biochemical or cellular functions. Furthermore, their residence at multiple subcellular locations
suggests that KHARON proteins are likely to be multifunctional, participating in flagellar
membrane trafficking, cytokinesis, and spindle function. Additionally, it is possible that
distinctions in function could be conferred by association of KHARON with different partner
proteins at each of its three subcellular loci. Thus, we hypothesize that there could exist three
distinct KHARON Complexes, Complex 1 at the base of the flagellum, Complex 2 at the
subpellicular cytoskeleton, and Complex 3 at the mitotic spindle. Furthermore, given the
apparent differences in *Tb*KH and *Lmx*KH noted above, there may be similarities and
differences between these putative complexes between the two species of parasite.

To initiate a study of putative KHARON Complexes and their functions, we carried out
biotinylation proximity labeling (BioID) (Roux et al., 2012) and tandem affinity purification-mass
spectrometry (TAP-MS) (Kaiser et al., 2008) on *Lmx*KH, resulting in the identification of two
KHARON Associated Proteins, *Lmx*KHAP1 and *Lmx*KHAP2 (LmxM.32.2440 and
LmxM.05.0380, respectively; (Kelly et al., 2020b)). In parallel, we investigated these two
KHARON partners in *T. brucei* and report the results of those studies here. As anticipated,
*Tb*KHAP1 and *Tb*KHAP2 exhibit both similarities and notable differences compared to their
orthologs in *L. mexicana*. In addition, *T. brucei* expresses another KHARON partner related to

*Tb*KHAP2 that we designate *Tb*KHAP3. Furthermore, characterization of additional flagellar

membrane proteins suggests that *Tb*KH expression is important for flagellar targeting of multiple

such proteins in African trypanosomes, whereas the role of *Lmx*KH in trafficking of flagellar

membrane proteins appears to be more restricted. These studies confirm that KHARON

proteins in both parasites exist in complexes with various partners and that these partner

proteins can play distinct roles in the functions of different KHARON Complexes.

RESULTS

Localization of *Tb*KHAP1, *Tb*KHAP2, TbKHAP3, and *Tb*KH in bloodstream and procyclic African trypanosomes. To facilitate studies on *Tb*KH and its partners, we raised and affinity purified a polyclonal antibody against this protein, anti-*Tb*KH pAb. Western blot analysis indicated that anti-*Tb*KH pAb detects a single protein of ~49 kDa molecular weight, and that an additional band of ~62 kDa appears in parasites also expressing a BirA* fusion on the N-terminus of *Tb*KH (Fig. 1 C), establishing that this antibody is of suitable specificity to employ in localization and biochemical characterization of *Tb*KH. To determine whether *Tb*KHAP1 (Tb927.11.2610), *Tb*KHAP2 (Tb927.10.10360) and *Tb*KHAP3 (Tb927.10.10280) are associated with *Tb*KH in a complex, several complementary approaches were applied. First, each KHAP was tagged at its N-terminus with the triple hemagglutinin peptide tag HA_3, and formaldehyde-fixed BF and PF trypanosomes were examined by immunofluorescence deconvolution microscopy (Fig. 2). HA_3::*Tb*KHAP1 (Fig. 2A,B, green) overlaps with *Tb*KH (red) at the cell periphery, as demonstrated by the yellow color in this region of both BF and PF parasites. In contrast, there was no apparent overlap of the two signals in the mitotic spindle (central red oval or line marked with a white arrow), indicating that this protein could be associated with *Tb*KH in the subpellicular cytoskeleton but not at the mitotic spindle. Similarly, *Tb*KHAP2::HA_3 (Figs. 2C,D) and $V5_3$::*Tb*KHAP3 or HA_3::*Tb*KHAP3 (Fig. 2E,F) overlap with *Tb*KH at the cell periphery but not at the mitotic spindle. For each of the *Tb*KHAPs, there is also green fluorescence that does not coincide with *Tb*KH so that there is not complete overlap of the signals, and there may thus be populations of each protein that are not associated with each other. However overall, these three *Tb*KHAPs are candidates for *Tb*KH partners that are selective for the subpellicular cytoskeleton versus the mitotic spindle.

To determine whether *Tb*KHAP1, *Tb*KHAP2, or *Tb*KHAP3 might associate with *Tb*KH at the base of the flagellum, flagella were isolated from parasites expressing each HA_3-tagged *Tb*KHAP and imaged by deconvolution microscopy. Fig. 3 shows that HA_3::*Tb*KHAP1 (Fig. 3A), $HA_{3::}$TbKHAP2 (Fig. 3B) and HA_3::TbKHAP3 (Fig. 3C) overlap significantly with *Tb*KH in the region of the flagellum immediately adjacent to the kinetoplast DNA (kDNA, blue, Fig. 3A), which is close to and physically attached to (Robinson and Gull, 1991) the flagellar basal body. Additional images showing overlap of *Tb*KH with each of the three TbKHAPs at the base of the flagellum are provided in Fig. S1. Overall, these results indicate that *Tb*KHAP1, *Tb*KHAP2, and *Tb*KHAP3 could be partners for *Tb*KH at both the pellicular cytoskeleton and the base of the flagellum. It is noteworthy that *Tb*KHAP3 was previously identified as a basal body protein in a

proteomic study of that subcellular structure (Dang et al., 2017). Hence, it is likely that *Tb*KH,

*Tb*KHAP1, *Tb*KHAP2, and *Tb*KHAP3 are all basal body components.

Molecular association of *Tb*KHAP1, *Tb*KHAP2, and *Tb*KHAP3 with *Tb*KH. To determine

whether the observed subcellular overlap of the fluorescence signals from *Tb*KHAPs and *Tb*KH

could indicate physical association in molecular complexes, *Tb*KH was endogenously tagged at

its N-terminus with a His_{10} affinity tag (His_{10}::*Tb*KH) to allow pulldown of this protein, and

associated partners, with Ni-NTA magnetic beads, and this affinity tagged protein was

expressed in a BF cell line also expressing either HA_3::*Tb*KHAP1, *Tb*KHAP2:: HA_3, or

$V5_3$::*Tb*KHAP3. Because *Tb*KH is an integral component of the parasite cytoskeleton (Sanchez

et al., 2016) and would pulldown many cytoskeletal proteins in an experiment performed under

native conditions, we first crosslinked with formaldehyde parasites expressing each pair of

tagged proteins. Formaldehyde crosslinks proteins that are in very close proximity (~2-3 Å,

reference (Hoffman et al., 2015)), so this treatment will covalently attach close molecular

partners of *Tb*KH but not proteins that are more distant partners in a complex or proteins that

are in the cytoskeleton but distant from *Tb*KH. Subsequent treatment with strongly denaturing

reagents will dissociate peripheral proteins from His_{10}::*Tb*KH while retaining crosslinked

partners, and the closely associated partners will thus be purified along with His_{10}::*Tb*KH,

following binding and elution from the Ni-NTA beads, and released upon heat-induced reversal

of the crosslinks.

Fig. 4A demonstrates that HA_3::*Tb*KHAP1 is pulled down with His_{10}::*Tb*KH when parasites

are formaldehyde crosslinked (EF*) but not when they are not subjected to crosslinking (EF). As

a negative control, another subpellicular cytoskeletal protein, CAP15 (Vedrenne et al., 2002),

was HA_3 tagged and expressed in His_{10}::*Tb*KH expressing BF parasites, but this protein was not

pulled down even from formaldehyde-crosslinked parasites. Parallel experiments were

performed using BF parasites expressing *Tb*KHAP2::HA_3, or $V5_3$::*Tb*KHAP3 and His_{10}::*Tb*KH.

However, for reasons that are not clear but possibly having to do with the highly repetitive

nature of both proteins (see below), treatment of parasites with formaldehyde followed by

immediate dissolution in the strongly denaturing urea buffer employed to disrupt protein

interactions led to massive degradation of both TbKHAP2::HA_3 and TbKHAP3::HA_3. Indeed, the

susceptibility of these proteins to degradation has been noted previously (Schneider et al.,

1988). Hence, for technical reasons we have not been able to observe pulldowns of either of

these two tagged protein with His_{10}::*Tb*KH, thus preventing us from directly demonstrating

interactions between these proteins at the 2-3 Å level. Nonetheless, these experiments

establish that *Tb*KHAP1 is in very close proximity to *Tb*KH.

Proximity ligation assay (PLA) confirms close proximity of *Tb*KH with *Tb*KHAP1,
***Tb*KHAP2, and *Tb*KHAP3.** To provide another independent examination of whether *Tb*KH is in
close physical proximity with each *Tb*KHAP, we performed the PLA in parasites expressing
HA$_3$::*Tb*KHAP1, *Tb*KHAP2::HA$_3$, and HA$_3$::*Tb*KHAP3. In this assay (Fredriksson et al., 2002;
Soderberg et al., 2006), cells expressing two partner proteins are first reacted with primary
antibodies from different species. Parasites are subsequently incubated with species-specific
secondary antibodies directed against each primary antibody, and each of these secondary
antibodies contains a unique, covalently attached oligonucleotide. Only if the two target proteins
are within ~400 Å of each other, these oligonucleotides can base pair to another linker
oligonucleotide and be covalently ligated into a circular substrate that can participate in rolling
circle DNA amplification of the cognate sequence. The amplified sequence is then hybridized to
a fluorescently labeled DNA probe, resulting in fluorescent puncta within the cell.

Fig. 4B,C shows a positive PLA signal (left panels) for BF and PF trypanosomes expressing
HA$_3$::*Tb*KHAP1 and probed with anti-*Tb*KH rabbit and anti-HA murine mAb. In contrast, when
the anti-HA murine antibody is not employed (right panels), the PLA signal is absent,
demonstrating the dependency of the signal on detection of both closely associated proteins.
Similar results confirm that *Tb*KH is in close physical proximity to *Tb*KHAP2 (Fig. 4D,E) and
*Tb*KHAP3 (Fig. 4F,G).

Predicted properties of *Tb*KHAP1, *Tb*KHAP2, and *Tb*KHAP3. Bioinformatic analysis of
the 50.9 kDa *Tb*KHAP1 sequence indicates that it is a protein apparently unique to kinetoplastid
protists for which there are orthologs widely distributed among Kinetoplastida. A BLASTP
search (tritrypdb.org) revealed several coiled-coil proteins such as neurofilament proteins and
tropomyosin as being significantly, although not closely, related (E values of 2.6e-08 – 5.4e-12).
Prediction of protein disorder using the PrDOS web server (Ishida and Kinoshita, 2007)
(http://prdos.hgc.jp/cgi-bin/top.cgi) generated a strong prediction of disorder (probability >0.9)
over the C-terminal region from amino acids 314 – 461. Indeed, this sequence is rich in E
residues, which predispose such regions to intrinsic disorder (Uversky, 2013), and this property
suggests that this region could be involved in protein-protein interactions through induced
folding (Zhang et al., 2013). InterPro (Mitchell et al., 2015)
(https://www.ebi.ac.uk/interpro/search/sequence/) predicted coils between amino acids 7 – 31
and 196 – 241 and a disordered region from residue 332 - 461 , and PSIPRED V4.0 (McGuffin
et al., 2000) (http://bioinf.cs.ucl.ac.uk/psipred/) predicted the sequence to be largely helix or coil.
Overall, computational analyses suggest that TbKHAP1 is a coiled-coil protein with an

intrinsically disordered C-terminus, both properties that are consistent with formation of multi-protein complexes.

*Tb*KHAP2 is the 374 kDa microtubule-associated repetitive protein 1, MARP-1, and *Tb*KHAP3 is the 267 kDa MARP-2 that have been studied previously by Seebeck and colleagues (Affolter et al., 1994; Hemphill et al., 1992; Schneider et al., 1988) and will hereafter be referred to as *Tb*KHAP2/MARP-1 and *Tb*KHAP3/MARP-2 to indicate both their association with *Tb*KH and their previously demonstrated roles in microtubule binding. Each sequence contains short unique N- and C-terminal domains, and the remainder of the sequence consists of 38-amino acid repeats that are largely conserved within each sequence but ~50% identical between the two proteins. The unique C-terminal domains (95% identical between the two proteins) bind to microtubules (Affolter et al., 1994), and the proteins decorate the subpellicular cytoskeleton (Schneider et al., 1988), but their specific biological functions have not been elucidated. In addition, these proteins have also been localized to the basal body and *Tb*KHAP3/MARP-2 was designated *Tb*BBP268 (Dang et al., 2017).

Phenotypes of BF trypanosomes following knockdown of *TbKHAP1* RNA. We have previously demonstrated that knockdown of *TbKH* RNA by inducible RNAi results in a lethal phenotype on both BF and PF trypanosomes (Sanchez et al., 2016). In these parasites, the flagellum detaches from the cell body along the flagellum attachment zone (FAZ), and the parasites are blocked in cytokinesis, resulting in accumulation of multi-nucleated, multi-flagellated 'monster cells' that are not viable in the long term. To assess the roles of *Tb*KHAP1, *Tb*KHAP2/MARP-1, and *Tb*KHAP3/MARP-2 in the biology of BF parasites, we targeted by RNAi *TbKHAP1* mRNA, using a unique RNAi probe, and *TbKHAP2/TbKHAP3* mRNAs jointly, using a 500 nt probe covering the conserved C-termini, and assessed the consequent phenotypes.

Induction of RNAi against *Tb*KHAP1 in BFs using doxycycline resulted in rapid reduction in the level of this protein (Fig. 5A). Furthermore, RNAi-induced parasites stopped growing almost immediately and were largely dead by 72 h (Fig. 5B). Quantification via microscopy of the percentage of cells with different numbers of nuclei and kinetoplasts (Fig. 5C) showed that following induction of RNAi over 48 h, the percentage of 1N1K parasites dropped dramatically, while those with multiple nuclei and kinetoplasts (XNYK) increased and began to predominate the population. In comparison to the normal morphology of pre-induced parasites (Fig. 5D), induction of RNAi for 20 h (Fig. 5E) or 48 h (Fig. 5F) resulted in parasites with multiple nuclei and/or kinetoplasts and tadpole-like morphology (white arrowhead) or duplicated flagella located at opposite poles of the cell body (yellow arrowhead).

Notable among cells in RNAi induced populations are those with multiple flagella located at
various relative positions around the cell (e.g., the two parasites in Fig. 5F). Such parasites
have initiated cytokinesis and cleavage furrow formation, as the two duplicated flagella have
moved apart from the initial position they would have following flagellar duplication. However,
the cleavage furrow did not progress to separate the duplicated nuclei and kinetoplasts as it
would in normal cell division. In summary, these observations suggest that loss of *Tb*KHAP1
protein from BF parasites results in a block in progression of cleavage furrow rather than an
inability to initiate cleavage furrow ingression.

These results are further enhanced by more refined time course studies shown in Fig. S2
following the progression of nuclear content and cell morphologies between 0 – 48 h after
induction of RNAi against *TbKHAP1* RNA. At 4 h (Fig. S2B) and 8 h (Fig. S2C) most parasites
had morphologies similar to that preceding induction of RNAi (0 h, Fig. S2A). However, by 12 h
(Fig. S2D), multi-flagellated parasites with ingression furrows appeared, and by 24 h and 48 h
(Fig. S2E,F), many parasites had incompletely resolved ingression furrows and multiple nuclei.

Phenotypes of parasites following knockdown of *TbKHAP2/MARP-1* and
***TbKHAP3/MARP-2* RNAs.** Induction of RNAi jointly against *TbKHAP2/MARP-1* and
TbKHAP3/MARP-2 resulted in complete loss of HA$_3$::*Tb*KHAP2/MARP-1 protein by 24 h (Fig.
6A. Depletion of *Tb*KHAP2/MARP-1 and *Tb*KHAP3/MARP-2 impaired growth of BF parasites,
resulting in an ~50-fold reduction in parasite number by 120 h (Fig. 6B) post-induction, but
growth inhibition was not nearly as pronounced as it is for *Tb*KHAP1 RNAi (Fig. 5B). Compared
to uninduced parasites (Fig. 6C), images of parasites following 4 d RNAi (Fig. 6D) still showed
many parasites with normal morphology similar to that of uninduced parasites, but some
parasites rounded up and showed multiple flagella (Fig. 6D, white arrowhead, flagella on
opposite sides of the cell body in DIC image). By 4 d post-RNAi, parasites with 1N2K and XNYK
began to accumulate (Fig. 6E), but the proportion was not nearly as great as for RNAi directed
against TbKHAP1 (Fig. 5C).

***Tb*KHAP1, but not *Tb*KHAP2/MARP-1 or *Tb*KHAP3/MARP-2, is required for targeting**
***Tb*CaCh/FS179 to the flagellar membrane.** The observation that *Tb*KHAP1 is located at the
base of the flagellum (Fig. 3A), likely in the basal body, raises the question of whether it could
play a role in the function of *Tb*KH in mediating trafficking of the putative Ca^{2+} channel,
*Tb*CaCh/FS179, to the flagellar membrane. To test this possibility, we induced RNAi against
TbKHAP1 RNA in parasites expressing *Tb*CaCh/FS179::HA$_3$ tagged at the C-terminus, which
localized to the flagellar membrane prior to RNAi (Fig. 7A). BF parasites induced for RNAi for 24
h (Fig. 7B) or 48 h (Fig. 7C) exhibited flagella that were devoid of *Tb*CaCh::HA$_3$ (white

arrowheads). These results suggest that a complex of *Tb*KH/*Tb*KHAP1, and potentially other
currently unknown partners located at the base of the flagellum, is involved in trafficking this
channel to the flagellar membrane. Since both proteins are also located in the subpellicular
cytoskeleton, it is not possible to definitively ascribe this flagellar trafficking phenotype to the
complex at the base of the flagellum; complexes at both locations will be downregulated by
TbKHAP1 RNAi. However, integral membrane proteins are first delivered to the flagellar pocket
membrane during biosynthesis (Manna et al., 2014). Hence, the presence of a protein complex
located close to the interface between the flagellar pocket and flagellar membrane, and for
which downregulation of both known partners inhibits trafficking of a protein into the flagellar
membrane, suggests that this complex may mediate trafficking of *Tb*CaCh/FS179 from the
flagellar pocket membrane into the flagellar membrane. Such trafficking would presumably be
mediated by a direct interaction between *Tb*KH and the cargo, *Tb*CaCh/FS179, and such a
molecular interaction has been demonstrated to occur (Sanchez et al., 2016).

In contrast, RNAi directed against *TbKHAP2/MARP-1* and *TbKHAP3/MARP-2* RNAs did not
prevent trafficking of *Tb*CaCh/FS179::HA$_3$ into the flagellar membrane, where it is located prior
to RNAi (Fig. 7D). At both 4 d (Fig. 7E) and 7 d (Fig. 7F) post RNAi, BF parasites with multiple
flagella still trafficked this channel into the flagellar membrane (green arrowheads).

_Tb_KH-dependent trafficking of other flagellar membrane proteins. *Tb*KH is required for
trafficking of *Tb*CaCh/FS179 to the flagellar membrane of BF trypanosomes, and the two protein
interact with each other, as demonstrated by crosslinking pulldown assays (Sanchez et al.,
2016). Is *Tb*KH important for flagellar trafficking of other membrane proteins? To address this
question, we monitored the dependency of other flagellar membrane proteins on *Tb*KH for
targeting to that organelle. The TrypTag project (http://tryptag.org/) has defined the subcellular
location of a large number of trypanosome proteins in PF parasites (Dean et al., 2017),
employing live cell microscopy of parasites expressing mNeonGreen fluorescent protein
fusions, and this endeavor has identified a cohort of flagellar membrane proteins.

One such flagellar membrane protein is Tb927.7.4270, a 25 kDa protein predicted to have a
N-terminal signal sequence and a single transmembrane domain (TMD) near its C-terminus.
This protein is one of four paralogous proteins (Tb927.7.4230, 4260, 4270, and 4280) studied
previously by Shimogawa *et al*. (Shimogawa et al., 2015) and designated the Fam79 proteins
(Fam79.1, 79.2, 79.3, and 79.4, respectively). To address potential dependency upon *Tb*KH for
flagellar targeting, we monitored the localization of the C-terminal mNG fusion of
Tb927.7.4270/Fam97.3 in both formaldehyde fixed and live PF parasites immobilized in CyGEL
(MacLean et al., 2013). As shown in Fig. 8A (0 h RNAi), this fusion protein is present in flagella

and also in filaments and vesicles that emerge from the flagella, and we designate
Tb927.7.4270 as extracellular vesicle membrane protein 1 or *Tb*EVMP1/Fam79.3. Multiple
investigators have observed filaments and vesicles emerging from various parts of
trypanosomes (Baudieri and Tomassini, 1962; Ellis et al., 1976; Molloy and Ormerod, 1964;
Schepilewsky, 1912; Vickerman and Luckins, 1969; Wright and Lumsden, 1970), including the
flagella, and a recent study by Szempruch *et al.* (Szempruch et al., 2016a; Szempruch et al.,
2016b) has investigated such structures from BF parasites in detail and concluded that the
flagellum-derived nanotubes and resulting extracellular vesicles (EVs) incorporate a cohort of
parasite proteins. Furthermore, delivery of parasite-derived EVs to host red blood cells or to
other trypanosomes can mediate pathogenic processes, such as erythrocyte clearance and
anemia in the mammalian host or delivery of innate immune factors from a resistant to a
sensitive strain of trypanosome. Hence, understanding the process for delivery of parasite
proteins to these EVs is of importance for deciphering mechanisms of parasite virulence.
Notably, when PF parasites expressing *Tb*EVMP1::mNG were subjected to RNAi directed
against *Tb*KH for 24 h, they were strongly impaired in trafficking of this fusion protein to flagella
or nanotubes (Fig. 8A, 24 h RNAi), and fluorescence often accumulated within the parasite cell
body. White arrows indicate flagella that are devoid of fluorescence and which thus exhibit a
trafficking defect. This result indicates that *Tb*EVMP1/Fam79.3 is dependent upon *Tb*KH, either
directly or indirectly, for trafficking to the flagellum and subsequently for release into EVs.
Furthermore, this trafficking defect occurs after 24 h of RNAi directed against *Tb*KH, but these
PF parasites do not exhibit significant loss of viability until ~10 days of continuous RNAi
(Sanchez et al., 2016), indicating that the effects of *TbKH* RNAi upon flagellar trafficking are not
due to global loss of cellular functions. Notably, Fam79.1 (Tb927.7.4230) was also detected by
proteomic analysis in EVs of BF trypanosomes by Szempruch *et al.* (Szempruch et al., 2016b),
indicating that multiple members of this family are delivered to the membranes of EVs during the
parasite life cycle.

A second flagellar membrane protein localized in TrypTag is Tb927.11.1830, designated
here as *Tb*EVMP2. This 62 kDa protein has 6 predicted TMDs and is widely distributed among
the kinetoplastid protists but does not have obvious orthologs outside that order, nor does it
possess conserved Pfam domains (Sonnhammer et al., 1998), except for the TMDs and one
predicted coiled coil. The *Tb*EVMP2::mNG fusion protein is also localized to the flagellar
membrane, nanotubes, and extracellular vesicles in PF parasites (Fig. 8B, 0 h RNAi), but
induction of RNAi directed against *Tb*KH also inhibits trafficking to these flagellar structures (Fig.
8B, 24 h RNAi), albeit less strongly than for *Tb*EVMP1. Although fluorescence is visible in some

flagella after induction of RNAi (green arrows, righthand image for 24 h RNAI in Fig. 8B), there

are some flagella that exhibit little if any fluorescence (white arrows). We designate this protein

*Tb*EVMP2 and suggest that it is a second such protein that is dependent upon *Tb*KH for efficient

trafficking to the surface of the flagellum.

In contrast, the mNG tagged FLA1 binding protein *Tb*FLA1BP::mNG, which is in the flagellar

membrane component of the FAZ (Sun et al., 2013), traffics efficiently to the FM both before

and after induction of RNAi (Fig. 8C, 0 h, 24 h, and 48 h RNAi), even in PF parasites that have

multiple flagella. Hence, *Tb*FLA1BP does not require *Tb*KH for targeting to the FM, implying that

there are both *Tb*KH-dependent and *Tb*KH-independent FM proteins in this parasite.

DISCUSSION

KHARON is a cytoskeletal protein that plays multiple roles in the biology of both *T. brucei* and *L. mexicana*. In trypanosomes, *Tb*KH is critical for flagellar trafficking of *Tb*CaCh/FS179 to the FM, and failure to traffic this channel to the flagellar component of the FAZ causes flagellar detachment and arrest of cytokinesis. Nonetheless, *Tb*KH may affect cell viability by other mechanisms as well. In addition to its localization at the base of the flagellum, the protein is also present in the subpellicular cytoskeleton and mitotic spindles where it could also play roles vital to parasite viability.

In the current study, we have investigated three potential partners of *Tb*KH: *Tb*KHAP1 and the two related proteins *Tb*KHAP2/MARP1 and *Tb*KHAP3/MARP2. *Tb*KHAP1 can be crosslinked to *Tb*KH by formaldehyde indicating that the two proteins associate within 2-3 Å of each other. *Tb*KHAP2 and *Tb*KHAP3 are close enough to *Tb*KH to give a consistently positive signal using the PLA, that is within 400 Å, but the degradation of these two *Tb*KHAPs under conditions of crosslinking and solubilization has prevented us from definitively demonstrating their interaction at the near atomic level. The knockdown of *TbKHAP1* RNA has the most pronounced phenotype, strongly arresting division of BF parasites, inhibiting progression of the cleavage furrow during cytokinesis, and impairing trafficking of *Tb*CaCh/FS179 to the FM. Hence, the phenotypes of RNAi for both *TbKH* and *TbKHAP1* are similar. In contrast, efficient knockdown of *TbKHAP2/3* RNAs slows growth of BF parasites but has a much less severe effect on cell division than knockdown of *TbKHAP1* RNA. In addition, trafficking of *Tb*CaCh/FS179 to the FM is not impaired, even after 7 days of knockdown in BFs. These results imply that although both types of protein likely associate with *Tb*KH in the cytoskeleton, they play different roles. These distinctions in functions could either result from separate complexes between *Tb*KH and each partner or from different roles that each partner plays in the same complex. The association of all partners with the cytoskeleton complicates this issue, as one cannot readily separate different complexes from each other, as would be possible for cytosolic multi-protein complexes. Nonetheless, these studies confirm that *Tb*KH associates with other partner proteins that mediate its activities in different ways.

One central activity for *Tb*KH is to traffic *Tb*CaCh/FS179 to the FM, a process that is critical for integrity of the FAZ and for parasite viability. The localization of many PF proteins to their subcellular sites achieved in the TrypTag.org project (Dean et al., 2017) has identified some additional FM proteins, along with some others that were identified previously as FM components from various targeted studies (Kelly et al., 2020a), and one question of relevance is how many of these flagellar surface components rely upon *Tb*KH for organellar trafficking. For

*Tb*CaCh/FS179, *Tb*KH appears to be directly involved in trafficking, since the two proteins can

be crosslinked by formaldehyde and isolated as molecular partners (Sanchez et al., 2016), but it

is possible that others depend upon *Tb*KH either directly or indirectly via the ability of this

protein to affect various cellular processes. In this study, we have shown that two additional FM

proteins, *Tb*EVMP1/Fam79.3 (Tb927.7.4270) and *Tb*EVMP2 (Tb927.11.1830) are present in

both the FM and in EVs secreted from the FM in PF trypanosomes. Both proteins exhibit

dependency upon *Tb*KH for trafficking to the FM, as RNAi directed against *TbKH* reduces the

efficiency of their localization to this organelle. In contrast, FLA1BP, which participates in

adhesion of the flagellum to the cell body by binding to the FLA1 protein in the cell body

component of the FAZ (Sun et al., 2013), is not dependent upon *Tb*KH to reach the FM,

confirming that both *Tb*KH-dependent and *Tb*KH-independent FM proteins exist.

EVs released from the cell body and FM of BF trypanosomes play important roles in

virulence of African trypanosomes, including lysis of host erythrocytes leading to anemia, a

major mechanism of trypanosome-mediated pathogenesis (Szempruch et al., 2016a;

Szempruch et al., 2016b). One might anticipate that surface components of EVs could play

important roles in either formation of the EV membrane or interaction of EVs with mammalian or

tsetse fly tissues. *TbEVMP1* and *TbEVMP2* mRNAs are both preferentially expressed in PF

trypanosomes (tritrypdb.org), but paralogs of *Tb*EVMP1, such as Tb927.7.4230 and

Tb927.7.4260, are expressed at higher levels in BFs compared to PFs, suggesting potential

roles for such EVMPs in both life cycle stages.

The *Tb*KH partners discovered in this study are associated primarily with the subpellicular

microtubules, but there are likely to be other partners that may reside principally at the base of

the flagellum or in the mitotic spindles and could be associated with distinct activities at those

sites. A BioID study by Zhou *et al*. (Zhou et al., 2018) identified five spindle-associated proteins,

NuSAP1, NuSAP2, Kif13-1, *Tb*MIP2, and *Tb*AUK1, that are in proximity to *Tb*KH and are thus

candidates for molecular partners at the mitotic spindle. In addition, Akiyoshi and Gull (Akiyoshi

and Gull, 2014) identified kinetoplast kinetochore proteins (KKTs) that associate with KHARON

by CoIP/MS experiments, suggesting a possible role of KHARON in faithful chromosome

segregation in kinetoplastid parasites. Molecular interaction studies of the type carried out here

will be required to determine which of these proteins may be *bona fide* molecular partners with

*Tb*KH and what roles KHARON complexes may be playing at the spindle. Similarly, the ability to

isolate flagella with associated kinetoplast DNA and basal bodies (Oberholzer et al., 2011;

Robinson and Gull, 1991; Subota et al., 2014) should facilitate identification by either BioID or

TAP-MS of additional *Tb*KH partners at the base of the flagellum. Thus, it should be possible to

achieve a comprehensive understanding of the role of KHARON in the cytoskeleton and the

distinct functions it carries out in association with different partner proteins.

MATERIALS AND METHODS

Growth and transfection of *T. brucei* cell lines. BF and PF *T. brucei* cell lines were grown as described previously (Sanchez, 2013). For T7 RNA polymerase-independent driven expression the BF/pHD1313 or PF/pHD1313 clones were employed (Sanchez et al., 2016). For T7 RNA polymerase driven expression in BF, *T. brucei* 427 parasites transfected with the pSmOx (Poon et al., 2012) plasmid expressing the tetracycline repressor and T7 RNA polymerase were generated and grown in 0.1 µg/ml puromycin. For T7 RNA polymerase driven expression in PF, *T. brucei* 427 13-6 clone expressing TETR and T7 RNA polymerase was used (Wirtz et al., 1999). Linear plasmid or PCR DNA amplicons were used to transfect mid-log phase parasites as described (Sanchez et al., 2016). Transfected clones were obtained by limiting dilution according to published protocols (Burkard et al., 2007; McCulloch et al., 2004) .

Primary amino acid sequence analysis. For DNA and amino acid sequence analysis of *TbKHAPs*, ExPASy, via the SIB Bioinformatics Resource Portal (http://expasy.org), and GeneBank (http://blast.ncbi.nlm.nih.gov/Blast.cgi) or TritrypDB   (http://tritrypdb.org/tritrypdb) were used.

Inhibition of gene expression by RNAi. To inhibit the expression of *Tb*KH, *Tb*KHAPs or *Tb*FLA1 a number of cell lines were generated employing different genetic backgrounds. The BF and PF TbKH^{RNAi} and *Tb*FLA1^{RNAi} clones, for which expression is T7 polymerase-independent, were previously described (Sanchez et al., 2016), and RNAi was induced by adding doxycycline (1 µg/ml) to the culture medium. The BF and PF *Tb*KHAP1^{RNAi} clones were generated by subcloning the first 500 bp of the *TbKHAP1* (Tb927.10.1026) ORF into the pZJM RNAi vector, where expression is driven by two opposing T7 promoters (Wang et al., 2000). Similarly, BF and PF TbKHAP2/3^{RNAi} clones were generated by subcloning the last 500 bp of *TbKHAP2* (Tb927.10.10360) that is almost identical to the *TbKHAP3* (Tb927.10.10280) ORF into the pZJM RNAi vector. RNAi clones were selected by resistance to 2.5 µg/ml phleomycin and 0.1 µg/ml puromycin, and expression of dsRNA was induced by addition of 1 µg/ml doxycycline. To verify inhibition of *Tb*KHAP1-3 expression, total cell lysates were obtained from parasite cultures induced for RNAi and subjected to Western blot experiments as indicated below.

Endogenous epitope tagging. For endogenous tagging of *Tb*KH, *Tb*KHAP1, *Tb*KHAP2, *Tb*KHAP3, CAP-15, *Tb*CaCh (FS179), *Tb*EVMP1, *Tb*EVMP2 and *Tb*FLA1BP at the N-terminus or C-terminus, different epitopes were employed as indicated in the text, following the protocol described (Dean et al., 2015). Briefly, epitope-tagging cassettes were generated by using two specific 100 nt oligonucleotides containing ~80 nt each that are homologous to the ORF region

to be tagged and ~20 nt homologs to the plasmid pPOTV4 template flanking the drug resistance

marker cassette, and using the universal PCR settings. Epitope tagging PCR cassettes were

ethanol precipitated and resuspended in 10 μl of nucleofection buffer (Wang et al., 2000), then

parasites were transfected with the purified tagging cassettes as described (Dean et al., 2015)

and selected using 1.5 μg/ml G418 (15 μg ml^{-1} for PF) or 0.1 μg/ml puromycin (1 μg/ml for PF),

and cloned by limiting dilution.

Generation of Rabbit anti-*Tb*KH antibody. A custom rabbit anti-*Tb*KH polyclonal antibody

(pAb) was generated by GenScript, using their 49-day antibody generation protocol. Briefly, two

rabbits were injected with 200 μg of His$_6$::*Tb*KH, representing amino acids 43 - 411, emulsified

in Freund's complete adjuvant. The rabbit was boosted 3 times at 14-day intervals with 200 μg

of His$_6$::*Tb*KH emulsified in Freund's incomplete adjuvant. Antibody specificity for *Tb*KH was

evaluated by western blot comparing the reactivities of the rabbit serum from immunized rabbits

to *T. brucei* protein lysates from wild-type cells and N-BirA*::*Tb*KH cell line (Fig 3A). Proteins

were immunodetected using 1:2500 dilution of the rabbit anti-*Tb*KH polyclonal antibody and

1:15,000 dilution of goat anti-rabbit-HRP antibody (Sigma-Aldrich). The chemiluminescent

protocol was used for developing as indicated below.

Immunofluorescence microscopy. For immunofluorescence microscopy, 5 X 10^6

parasites were centrifuged at 1000 X g for 5 min and washed twice at room temperature with

phosphate buffered saline pH 7.2 (PBS) containing 10 mM glucose. The cell pellet was

resuspended in 4% paraformaldehyde in PBS, pH 7.2 and incubated for 15 min at room

temperature, cells were centrifuged as descried above and washed once with PBS,

resuspended in 100 μl PBS, spotted onto poly-L-lysine coated coverslips, permeabilized with

0.1% Triton X-100 in PBS for t min, and washed 3X with PBS. Then parasites were blocked with

2% goat serum, 0.01 sodium azide, 0.01 saponin in PBS (blocking solution) for 1 h at room

temperature, rinsed 3 X with PBS, and incubated with primary antibodies for 1 h at room

temperature. The following primary antibodies were employed: 1:250 dilution rabbit anti-*Tb*KH

pAb (reported in this work), 1:500 dilution mouse anti-HA monoclonal antibody (mAb)

(BioLegend, Cat # MMS-101R), and 1:1000 dilution mouse anti-α-tubulin mAb (Sigma-Aldrich,

Cat. # T5168). Subsequently, cells were rinsed as before and incubated with a 1:1000 dilution of

secondary antibodies coupled to Alexa Fluor dyes (Molecular Probes) as follows: Alexa Fluor®

488 goat anti-mouse IgG (H+L) (Cat. # A11001), Alexa Fluor® 594 goat anti-mouse IgG (H+L)

(Cat. # A11005), Alexa Fluor® 594 goat anti-rabbit IgG (H+L) (Cat. # A11012) and Alexa Fluor®

488 goat anti-rabbit IgG (H+L) (Cat. # A11008), as indicated, in blocking solution for 1h at room

temperature in the dark. Cover slips were rinsed 3 X with PBS and mounted onto slides using

DAPI Fluoromount-G (SouthernBiotech). Fluorescence images were obtained using a wide field

deconvolution system (Applied Precision Instruments, Inc.) consisting of an inverted Nikon TE

200 Eclipse microscope, a Kodak CH350 CCD camera, and the Deltavision operating system.

Images were acquired using a 60 X objective and 1.25 X magnification in a 1024 X 1024 format,

and deconvolved using SoftWoRx software. Adobe Photoshop CC and Adobe Illustrator CC

(Adobe Systems Inc.) were used to create image compositions.

Flagellar protein purification, formaldehyde crosslinking, pull down, and Western blot

assay. Flagellar purification was performed from BF parasites (Fig. 3A,B and Fig. S1A,B) as

described (Subota et al., 2014) with slight changes. Briefly, BF TbKH1^{RNAi}, TbFLA1^{RNAi},

TbKHAP1^{RNAi} and TbKHAP2/3^{RNAi} clones were grown in 1 µg/ml doxycycline for the indicated

times and pelleted at 420 X g for 10 min. Cell pellets were washed with buffer A (25 mM Na$^+$-

tricine, pH 7, 1% BSA, 0.1 mM CaCl$_2$, 0.2 mM EDTA, 5 mM MgCl$_2$ and 12 mM β-

mercaptoethanol) containing 0.32 M sucrose and centrifuged at 420 X g for 10 min. Cell pellets

were gently resuspended at 3 X 10^8 parasites/ml in buffer A plus 0.3 M sucrose, transferred into

Eppendorf tubes and vortexed for 5 min or until microscopic verification of flagellum

detachment, followed by centrifugation at 420 X g for 10 min. Supernatants were recovered and

centrifuged at 16,000 X g for 20 min at 4 °C. Pellets containing the isolated flagella were

resuspended in 200 µl of PBS and used for immunofluorescence microscopy. Also, flagellar

purification from PF clones without induction of RNAi (Fig. 3C and Fig. S1C) was performed by

isolation of cytoskeletons followed by treatment with 1 mM CaCl$_2$, according to Imhof et al.

(Imhof et al., 2019).

For formaldehyde crosslinking and pull down, cell lines were prepared that expressed

TbKH1 endogenously tagged with the His$_{10}$ epitope at the N-terminus (His$_{10}$::TbKH1) and also

co-expressed with a protein of interest endogenously tagged at its N- or C-terminus with a HA$_3$

or V5$_3$ epitope (HA$_3$::TbKHAP1, HA$_3$::TbKHAP2, TbKHAP2::HA$_3$, V5$_3$::TbKHAP3,

HA$_3$::TbKHAP3 and TbCAP15::HA$_3$). Parasites were washed once with PBS and pelleted at

1000 X g for 10 min, and cell pellets resuspended in 9.37 ml PBS plus 0.63 ml 16%

formaldehyde-EM grade (Polysciences, Inc) and incubated at room temperature for 10 min. For

non-crosslinked control samples, PBS was added instead of formaldehyde. Subsequently, 1 ml

of 2.5 M glycine in PBS was added to the crosslinked samples and incubated at room

temperature for 5 min, crosslinked parasites were pelleted at 1000 X g for 10 min and washed

twice with PBS. Then, cell pellets with or without crosslinking were resuspended in 1 ml of

Buffer 1 (8 M urea, 300 mM NaCl, 0.5% Nonidet P-40, 50 mM NaH$_2$PO$_4$, 50 mM Tris, pH 7.0, 10

mM imidazole) on ice. Samples were sonicated 3 times on ice at 50% max amplitude (Sonic

Dismembrator, 500W, Fisher Scientific) for 10 s with 30 s between pulses. A 2.5% aliquot of this

protein lysate was saved as the protein lysate fraction (LF).

The remainder of the protein was incubated with 50 µl of Ni-NTA Magnetic Beads (New

England Biolabs, Inc.) on a rocker for 45 min at room temperature. The beads were washed

three times according to the manufacturer's instructions using Buffer 1 plus 20 mM imidazole.

Finally, bound protein complexes were eluted with 100 µl Buffer 1 containing 500 mM imidazole.

Crosslinking was reversed by boiling protein samples for 30 min in 1X Bolt™-LDS sample buffer

(Life Technologies) containing 10 mM DTT.

Protein extracts were prepared and analyzed by Western blot employing Bolt™ 4-12% Bis-

Tris Mini Protein Gels or NuPage™ 3-8% Tris-acetate gels, Mini Gel Tank and Mini Blot Module

following the manufacturer's instructions (Life Technologies). Proteins were transferred onto

PVDF membranes (Millipore). Protein immunodetection was done using rabbit anti-*Tb*KH pAb at

1:2500 dilution, mouse anti-HA mAb (BioLegend, Cat # MMS-101R) 1:2500 dilution, mouse

anti-V5 mAb (Invitrogen, Cat # MA5-15253) 1:2500 dilution, and mouse anti-α tubulin mAb

(Sigma-Aldrich, Cat. # T5168) 1:10,000 dilution. Goat anti-rabbit-HRP (Sigma-Aldrich) 1:15,000

dilution and goat anti-mouse-HRP (Jackson ImmunoResearch Laboratories, Cat. # 115-03-174,

Lot # 117119) were used as secondary antibodies and Western blots were developed using the

SuperSignal™ West Pico Plus Chemiluminescent Substrate (Thermo Fisher Scientific) and an

Image Quant LAS 400 (GE Healthcare) scanner was employed to acquire luminescent images.

Adobe Photoshop CC and Adobe Illustrator CC (Adobe Systems Inc.) were used to create

image compositions.

Proximity ligation assay (PLA). Mid-log BF or PF parasites were harvested by

centrifugation at 1000 x g for 10 min, washed once in PBS and fixed with 4% paraformaldehyde

in PBS for 15 min at room temperature. Fixed cells were attached to cover slips and

permeabilized as indicated above for immunofluorescence analysis. Subsequently the PLA

protocol was followed according to the Duolink In Situ Red Starter Kit Mouse/Rabbit (Millipore

Sigma) instructions. Briefly, after blocking cells were incubated with 1:250 dilution of rabbit anti-

*Tb*KH pAb and 1:500 dilution of mouse anti-HA mAb (BioLegend, Cat# MMS-101R). As a

negative control, cells were incubated only with 1:250 dilution of the rabbit anti-*Tb*KH pAb. Then

PLA species-specific secondary antibodies with minus and plus oligonucleotide probes were

added, followed by ligation, amplification and hybridization with specific red-fluorescent

oligonucleotides to allow detection by fluorescence microscopy. Samples were mounted and

imaged as described for immunofluorescence analysis.

Acknowledgements

We appreciate the expert advice and support of the staff of the Advanced Light Microscopy

Core in the Jungers Center for Neurosciences at Oregon Health & Science University. We

acknowledge discussions with Dr. Samuel Dean (University of Oxford) concerning new flagellar

membrane proteins identified by the TrypTag project.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: M.A.S. and S.M.L.; Data collection: M.A.S.; Data analysis: M.A.S. and

S.M.L.; Writing and editing: M.A.S. and S.M.L.; Funding acquisition and project administration:

S.M.L.

Funding

This work was supported by National Institutes of Health grant AI121160 to S.M.L. The content

is solely the responsibility of the authors and does not necessarily represent the official views of

the National Institutes of Health.

References

Affolter, M., Hemphill, A., Roditi, I., Muller, N. and Seebeck, T. (1994). The repetitive microtubule-associated proteins MARP-1 and MARP-2 of *Trypanosoma brucei. J Struct Biol* **112**, 241-51.

Akiyoshi, B. and Gull, K. (2014). Discovery of unconventional kinetochores in kinetoplastids. *Cell* **156**, 1247-1258.

Baudieri, B. and Tomassini, N. (1962). Fine struttura dei tripanosomi. *Parassitologia* **4**, 89-95.

Burkard, G., Fragoso, C. M. and Roditi, I. (2007). Highly efficient stable transformation of bloodstream forms of *Trypanosoma brucei. Mol Biochem Parasitol* **153**, 220-3.

Cayla, M., Rojas, F., Silvester, E., Venter, F. and Matthews, K. R. (2019). African trypanosomes. *Parasit Vectors* **12**, 190.

Dang, H. Q., Zhou, Q., Rowlett, V. W., Hu, H., Lee, K. J., Margolin, W. and Li, Z. (2017). Proximity Interactions among basal body components in *Trypanosoma brucei* identify novel regulators of basal body biogenesis and inheritance. *mBio* **8**.

Dean, S., Sunter, J., Wheeler, R. J., Hodkinson, I., Gluenz, E. and Gull, K. (2015). A toolkit enabling efficient, scalable and reproducible gene tagging in trypanosomatids. *Open Biol* **5**, 140197.

Dean, S., Sunter, J. D. and Wheeler, R. J. (2017). TrypTag.org: A trypanosome genome-wide protein localisation resource. *Trends Parasitol* **33**, 80-82.

Ellis, D. S., Ormerod, W. E. and Lumsden, W. H. R. (1976). Filaments of "*Trypanosoma brucei*": some notes on differences in origin and structure in two stains of "*Trypanosoma (Trypanozoon) brucei rhodesiense*". *Acta Tropica* **33**, 151-168.

Farr, H. and Gull, K. (2012). Cytokinesis in trypanosomes. *Cytoskeleton (Hoboken)* **69**, 931-41.

Fredriksson, S., Gullberg, M., Jarvius, J., Olsson, C., Pietras, K., Gustafsdottir, S. M., Ostman, A. and Landegren, U. (2002). Protein detection using proximity-dependent DNA ligation assays. *Nat Biotechnol* **20**, 473-7.

Hemphill, A., Affolter, M. and Seebeck, T. (1992). A novel microtubule-binding
motif identified in a high molecular weight microtubule-associated protein from
Trypanosoma brucei. J Cell Biol **117**, 95-103.

Hoffman, E. A., Frey, B. L., Smith, L. M. and Auble, D. T. (2015).
Formaldehyde crosslinking: a tool for the study of chromatin complexes. *J Biol Chem*
290, 26404-11.

Imhof, S., Zhang, J., Wang, H., Bui, K. H., Nguyen, H., Atanasov, I., Hui, W.
H., Yang, S. K., Zhou, Z. H. and Hill, K. L. (2019). Cryo electron tomography with Volta
phase plate reveals novel structural foundations of the 96-nm axonemal repeat in the
pathogen *Trypanosoma brucei. Elife* **8**.

Ishida, T. and Kinoshita, K. (2007). PrDOS: prediction of disordered protein
regions from amino acid sequence. *Nucleic Acids Res* **35**, W460-4.

Kaiser, P., Meierhofer, D., Wang, X. and Huang, L. (2008). Tandem affinity
purification combined with mass spectrometry to identify components of protein
complexes. *Methods Mol Biol* **439**, 309-26.

Kelly, F. D., Sanchez, M. A. and Landfear, S. M. (2020a). Touching the
surface: Diverse roles for the flagellar membrane in Kinetoplastid parasites. *Microbiol*
Mol Biol Rev **84**, e00079-19.

Kelly, F. D., Tran, K. D., Hatfield, J., Schmidt, K., Sanchez, M. A. and
Landfear, S. M. (2020b). A cytoskeletal protein complex is essential for division of
intracellular amastigotes of *Leishmania mexicana. J Biol Chem* **295**, 13106-13122.

Kennedy, P. G. (2013). Clinical features, diagnosis, and treatment of human
African trypanosomiasis (sleeping sickness). *Lancet Neurol* **12**, 186-94.

Langousis, G. and Hill, K. L. (2014). Motility and more: the flagellum of
Trypanosoma brucei. Nat Rev Microbiol **12**, 505-18.

MacLean, L., Myburgh, E., Rodgers, J. and Price, H. P. (2013). Imaging
African trypanosomes. *Parasite Immunol* **35**, 283-94.

Manna, P. T., Boehm, C., Leung, K. F., Natesan, S. K. and Field, M. C. (2014).
Life and times: synthesis, trafficking, and evolution of VSG. *Trends Parasitol* **30**, 251-8.

McCulloch, R., Vassella, E., Burton, P., Boshart, M. and Barry, J. D. (2004).
Transformation of monomorphic and pleomorphic *Trypanosoma brucei*. *Methods Mol*
Biol **262**, 53-86.

McGuffin, L. J., Bryson, K. and Jones, D. T. (2000). The PSIPRED protein
structure prediction server. *Bioinformatics* **16**, 404-5.

Mitchell, A., Chang, H. Y., Daugherty, L., Fraser, M., Hunter, S., Lopez, R.,
McAnulla, C., McMenamin, C., Nuka, G., Pesseat, S. et al. (2015). The InterPro
protein families database: the classification resource after 15 years. *Nucleic Acids Res*
43, D213-21.

Molloy, J. O. and Ormerod, W. E. (1964). A fibril emerging from the posterior
end of *Trypanosoma rhodesiense*. *Philos Trans R Soc Lond B Biol Sci* **58**, 2.

Oberholzer, M., Langousis, G., Nguyen, H. T., Saada, E. A., Shimogawa, M.
M., Jonsson, Z. O., Nguyen, S. M., Wohlschlegel, J. A. and Hill, K. L. (2011).
Independent analysis of the flagellum surface and matrix proteomes provides insight
into flagellum signaling in mammalian-infectious *Trypanosoma brucei*. *Mol Cell*
Proteomics **10**, M111 010538.

Poon, S. K., Peacock, L., Gibson, W., Gull, K. and Kelly, S. (2012). A modular
and optimized single marker system for generating *Trypanosoma brucei* cell lines
expressing T7 RNA polymerase and the tetracycline repressor. *Open Biol* **2**, 110037.

Robinson, D. R. and Gull, K. (1991). Basal body movements as a mechanism
for mitochondrial genome segregation in the trypanosome cell cycle. *Nature* **352**, 731-3.

Roux, K. J., Kim, D. I., Raida, M. and Burke, B. (2012). A promiscuous biotin
ligase fusion protein identifies proximal and interacting proteins in mammalian cells. *J*
Cell Biol **196**, 801-10.

Sanchez, M. A. (2013). Molecular identification and characterization of an
essential pyruvate transporter from *Trypanosoma brucei*. *J Biol Chem* **288**, 14428-37.

Sanchez, M. A., Tran, K. D., Valli, J., Hobbs, S., Johnson, E., Gluenz, E. and
Landfear, S. M. (2016). KHARON Is an essential cytoskeletal protein involved in the
trafficking of fagellar membrane proteins and cell division in African trypanosomes. *J*
Biol Chem **291**, 19760-73.

Santi, A. M. M., Lanza, J. S., Tunes, L. G., Fiuza, J. A., Roy, G., Orfano, A. D.
S., de Carvalho, A. T., Frezard, F., Barros, A. L. B., Murta, S. M. F. et al. (2018).
Growth arrested live-attenuated *Leishmania infantum* KHARON1 null mutants display
cytokinesis defect and protective immunity in mice. *Sci Rep* **8**, 11627.

Schepilewsky, E. (1912). Fadenförmige anhängsel bei den trypanosomen. *Zbl.*
Bakt. **65**, 79-83.

Schneider, A., Hemphill, A., Wyler, T. and Seebeck, T. (1988). Large
microtubule-associated protein of *T. brucei* has tandemly repeated, near-identical
sequences. *Science* **241**, 459-62.

Shimogawa, M. M., Saada, E. A., Vashisht, A. A., Barshop, W. D.,
Wohlschlegel, J. A. and Hill, K. L. (2015). Cell surface proteomics provides insight
into stage-specific remodeling of the host-parasite interface in *Trypanosoma brucei*. *Mol*
Cell Proteomics **14**, 1977-88.

Soderberg, O., Gullberg, M., Jarvius, M., Ridderstrale, K., Leuchowius, K. J.,
Jarvius, J., Wester, K., Hydbring, P., Bahram, F., Larsson, L. G. et al. (2006). Direct
observation of individual endogenous protein complexes *in situ* by proximity ligation. *Nat*
Methods **3**, 995-1000.

Sonnhammer, E. L., Eddy, S. R., Birney, E., Bateman, A. and Durbin, R.
(1998). Pfam: multiple sequence alignments and HMM-profiles of protein domains.
Nucleic Acids Res **26**, 320-2.

Subota, I., Julkowska, D., Vincensini, L., Reeg, N., Buisson, J., Blisnick, T.,
Huet, D., Perrot, S., Santi-Rocca, J., Duchateau, M. et al. (2014). Proteomic analysis
of intact flagella of procyclic *Trypanosoma brucei* cells identifies novel flagellar proteins
with unique sub-localization and dynamics. *Mol Cell Proteomics* **13**, 1769-86.

Sun, S. Y., Wang, C., Yuan, Y. A. and He, C. Y. (2013). An intracellular
membrane junction consisting of flagellum adhesion glycoproteins links flagellum
biogenesis to cell morphogenesis in *Trypanosoma brucei*. *J Cell Sci* **126**, 520-31.

Szempruch, A. J., Dennison, L., Kieft, R., Harrington, J. M. and Hajduk, S. L.
(2016a). Sending a message: extracellular vesicles of pathogenic protozoan parasites.
Nat Rev Microbiol **14**, 669-675.

Szempruch, A. J., Sykes, S. E., Kieft, R., Dennison, L., Becker, A. C.,
Gartrell, A., Martin, W. J., Nakayasu, E. S., Almeida, I. C., Hajduk, S. L. et al.
(2016b). Extracellular vesicles from *Trypanosoma brucei* mediate virulence factor
transfer and cause host anemia. *Cell* **164**, 246-257.

Tran, K. D., Rodriguez-Contreras, D., Vieira, D. P., Yates, P. A., David, L.,
Beatty, W., Elferich, J. and Landfear, S. M. (2013). KHARON1 mediates flagellar
targeting of a glucose transporter in *Leishmania mexicana* and is critical for viability of
infectious intracellular amastigotes. *J. Biol. Chem.* **288**, 22721-22733.

Tran, K. D., Vieira, D. P., Sanchez, M. A., Valli, J., Gluenz, E. and Landfear,
S. M. (2015). *Kharon1* null mutants of *Leishmania mexicana* are avirulent in mice and
exhibit a cytokinesis defect within macrophages. *PLoS One* **10**, e0134432.

Uversky, V. N. (2013). The alphabet of intrinsic disorder: II. Various roles of
glutamic acid in ordered and intrinsically disordered proteins. *Intrinsically Disord*
Proteins **1**, e24684.

Vaughan, S. and Gull, K. (2008). The structural mechanics of cell division in
Trypanosoma brucei. *Biochem Soc Trans* **36**, 421-4.

Vedrenne, C., Giroud, C., Robinson, D. R., Besteiro, S., Bosc, C., Bringaud,
F. and Baltz, T. (2002). Two related subpellicular cytoskeleton-associated proteins in
Trypanosoma brucei stabilize microtubules. *Mol Biol Cell* **13**, 1058-70.

Vickerman, K. and Luckins, A. G. (1969). Localization of variable antigens in
the surface coat of *Trypanosoma brucei* using ferritin conjugated antibody. *Nature* **224**,
1125-1126.

Wang, Z., Morris, J. C., Drew, M. E. and Englund, P. T. (2000). Inhibition of
Trypanosoma brucei gene expression by RNA interference using an integratable vector
with opposing T7 promoters. *J Biol Chem* **275**, 40174-9.

Wheeler, R. J., Gull, K. and Sunter, J. D. (2019). Coordination of the Cell Cycle
in Trypanosomes. *Annu Rev Microbiol* **73**, 133-154.

Wirtz, E., Leal, S., Ochatt, C. and Cross, G. A. M. (1999). A tightly regulated
inducible expression system for conditional gene knock-outs and dominant-negative
genetics in *Trypanosoma brucei. Mol. Biochem. Parasitol.* **99**, 89-101.

Wright, K. A. and Lumsden, W. H. R. (1970). The formation of filopodium-like
processes by *Trypanosoma*. *J. Cell Sci.* **6**, 285-297.

Zhang, T., Faraggi, E., Li, Z. and Zhou, Y. (2013). Intrinsically semi-disordered
state and its role in induced folding and protein aggregation. *Cell Biochem Biophys* **67**,
1193-205.

Zhou, Q., Lee, K. J., Kurasawa, Y., Hu, H., An, T. and Li, Z. (2018). Faithful
chromosome segregation in *Trypanosoma bruce*i requires a cohort of divergent spindle-
associated proteins with distinct functions. *Nucleic Acids Res* **46**, 8216-8231.

749
750

FIGURE LEGENDS

Fig. 1. *Tb*KH localizes in three distinct subcellular compartments. (A) Flagellar preparation from *Tb*FLA1[RNAi] cell line, isolated after 72 h of induction with 1 µg/ml doxycycline (dox), stained with DAPI (*blue*) and immunostained with anti-*Tb*KH pAb (*Tb*KH, *red*). *Tb*KH localization at the base of the flagellum is indicated by the *red arrowhead*. (B) Wild type BF parasites were immunostained with anti-*Tb*KH pAb (*Tb*KH, *red*) and stained with DAPI. These images display a parasite in which *Tb*KH staining is associated with both the subpellicular microtubules (*right arrowheads*) and the mitotic spindle that connects the two nuclei late during mitosis (*left arrowheads*). DIC indicates images collected by differential interference contrast microscopy. (C) Western blot of total protein lysates from BF trypanosomes that are either wild type (*WT*) or expressing BirA*::TbKH fusion protein (*A**). Blot was probed with anti-*Tb*KH pAb and developed by chemiluminescence. Specific immunodetected proteins corresponding to the native *Tb*KH and the BirA*::*Tb*KH fusion protein are shown. Relative protein molecular weights are shown in kDa, as determined by mobility relative to molecular weight markers.

Fig. 2. Subcellular localization of *Tb*KHAPs in BF and PF *T. brucei*. (A, B) The HA$_3$::TbKHAP1 cell line was immunostained with anti-HA mAb (HA$_3$::*Tb*KHAP1*, green*) and anti-*Tb*KH pAb (*Tb*KH*, red*). (C, D) The TbKHAP2::HA$_3$ cell line was stained with anti-HA mAb (*Tb*KHAP2::HA$_3$*, green*) and anti-*Tb*KH pAb (*Tb*KH*, red*). (E) The BF 3V5::*Tb*KHAP3 and (F) PF HA$_3$::*Tb*KHAP3 cell lines were immunostained with (E) anti-V5 mAb (3V5::*Tb*KHAP3, *green)* or (F) anti-HA mAb (HA$_3$::*Tb*KHAP3*, green*) and anti-*Tb*KH pAb (*Tb*KH*, red*). All preparations were stained with DAPI, which detects both nuclear and kinetoplast DNA (*blue*). DIC images were also acquired from all samples.

Fig. 3. KHARON and KHAPs colocalize at the base of the flagellum in *T. brucei*. (A) Whole isolated flagella from the BF HA$_3$::*Tb*KHAP1/*Tb*FLA1[RNAi] clone induced for RNAi for 72 h, were immunostained with anti-HA mAb (HA$_3$::*Tb*KHAP1, *green*) and anti-*Tb*KH pAb (*Tb*KH, *red*) and stained with DAPI (*blue*). (A, left panel) shows an immunofluorescence/DIC image of an isolated flagellum, *white punctuated* box indicates the magnified area depicted in (A, center-right, center-left and left) panels. *Green arrowheads* indicate HA$_3$::*Tb*KHAP1 localization, *red arrowheads* indicate *Tb*KH localization and *yellow arrowheads* indicate overlapping signals, when channels are merged, near the kinetoplast (*blue*). (B) Flagella were isolated from BF trypanosomes expressing HA$_3$::TbKHAP2 and imaged as described in (A). (C) Flagellar cytoskeletons were isolated from PF trypanosomes expressing HA$_3$::TbKHAP3 and imaged as described in (A).

Fig. 4. *Tb*KH is closely associated with *Tb*KHAP1, *Tb*KHAP2, and *Tb*KHAP3. (A) Western blot of protein samples from pull down of HA$_3$::*Tb*KHAP1 using His$_{10}$::*Tb*KH as a bait and

CAP15::HA$_3$ as negative control. *LS* and *EF*, lysate and elution fractions without formaldehyde

crosslinking; *LS** and *EF**, lysate and elution fraction with formaldehyde crosslinking. Protein

blots were developed by chemiluminescence, bands are indicated by protein names, and

migration of molecular weight markers are designated in kDa. (B, C) PLA employing BF or PF

HA$_3$::TbKHAP1/His$_{10}$::TbKH cell lines, as indicated. For these and all subsequent sections of

this figure, the *left panel* shows results using both anti-HA mAb and anti-*Tb*KH pAb, and the

right panel shows results using only the anti-*Tb*KH pAb, as the negative control. *Red puncta*

indicate positive HA$_3$::*Tb*KHAP1-*Tb*KH interaction. DIC images were acquired from all samples.

(D, E) PLA employing BF or PF *Tb*KHAP2::HA$_3$ cell lines, as indicated. *Red puncta* indicate

positive *Tb*KHAP2-*Tb*KH interaction. (F, G) PLA employing BF or PF HA$_3$::*Tb*KHAP3 cell lines,

as indicated using. *Red puncta* indicate positive *Tb*KHAP3-*Tb*KH interaction.

Figure 5. Depletion of *Tb*KHAP1 is lethal for *T. brucei* parasites. (A) Western blot of total

protein lysates from TbKHAP1^{RNAi} BF parasites grown in the presence of doxycycline (+dox)

and immunodetected with anti-HA mAb, anti-*Tb*KH pAb. The approximate molecular weights in

kDa of HA$_3$::*Tb*KHAP1 and *Tb*KH are indicated as determined by mobility compared to weight

markers. *Numbers* under the blot represent the relative intensity (*RI*) of HA$_3$::*Tb*KHAP1 protein

normalized to the *Tb*KH protein level. (B) Cell density for induced (*empty circles*) and non-

induced (*filled circles*) *Tb*KHAP1^{RNAi} BF cell lines. Parasite density was quantified by phase

contrast microscopy using a hemocytometer. Data represent the averages and standard

deviations of two experiments, each employing an independently isolated *Tb*KHAP1^{RNAi} clonal

cell line, and technical replicates were also performed for each biological replicate. Standard

deviations are too small to be visible. (C) Microscopic analysis of nuclei and kinetoplast

numbers of BF *Tb*KHAP1^{RNAi} parasites following induction of RNAi. Data represent frequency

(*%*) of cells with different numbers of DAPI stained nuclei (*N*) and kinetoplasts (*K*). Results

represent the average and range of two independent experiments. (D-F) *Tb*KHAP1^{RNAi} BF

parasites were stained with DAPI (*blue*) and immunostained with anti-α-tubulin mAb (Tub, *red*)

at (D) 0 h, (E) 20 h and (F) 48 h post-RNAi induction. In (E) the *white arrowhead* indicates a cell

with a tadpole-like morphology and the *yellow arrowhead* indicates a cell with two flagella at

opposite poles of the cell body.

Fig. 6. Phenotypes resulting from RNAi directed against *TbKHAP2* and *TbKHAP3*. (*A*)

Western blot of total protein lysates of BF parasites expressing HA$_3$::*Tb*KHAP2 following

induction of RNAi (1 µg ml^{-1} doxycycline, *dox*) directed against *TbKHAP2/3* RNAs. Protein blots

were probed with anti-HA mAb and anti-*Tb*KH pAb as loading control. Molecular weight markers

are indicated in kDa. (*B*) Growth curve of induced (*empty circles*) and non-induced (*filled circles*)

TbKHAP2/3^{RNAi} cell line. Parasite density was quantified by phase contrast microscopy using a

hemocytometer. The data represent two biological replicate experiments, but the standard

deviations are too small to see in the figure. Representative *Tb*KHAP2/3^{RNAi} cells induced for

RNAi were stained with DAPI (*blue*) and immunostained with anti-*Tb*KH pAb (*Tb*KH, *red*) at (*C*)

0 h and (*D)* 4 days post-RNAi induction. *White arrowhead* in (*D*) indicates a cell showing

aberrant morphology. DIC images were obtained for all samples. (*E*) Microscopic analysis of NK

in BF parasites undergoing RNAi. Frequency (*%*) of cells with different numbers of nuclei (*N*)

and kinetoplasts (*K*) at different times following induction of RNAi against *TbKHAP2/3* mRNA.

Results represent the average and range of two independent experiments.

Fig. 7. Knockdown of *Tb*KHAP1, but not *Tb*KHAP2 or *Tb*KHAP3, impairs trafficking of

***Tb*CaCh::HA$_3$ to the flagellum.** (A-C)*Tb*CaCh::HA$_3$/*Tb*KHAP1^{RNAi} BF cell line was induced with

1 µg/ml doxycycline (dox). Parasites were stained with DAPI (*blue*) and immunostained with

anti-HA mAb (*Tb*CaCh::HA$_3$*, green*) and anti-*Tb*KH pAb (*Tb*KH, *red*). (A) Non-induced

TbCaCh::HA$_3$/TbKHAP1^{RNAi} parasites (*0 h dox*), (B) parasites induced for 24 h (2*4 h dox*) and

(C*)* parasites induced for 48 h (*48 h dox*). *Green arrowheads* in (A-C) indicate flagella where

*Tb*CaCh::HA$_3$ is present, and *white arrowheads* in *(*B-C*)* indicate flagella where *Tb*CaCh::HA$_3$ is

absent. (D-F) *Tb*CaCh::mNG/*Tb*KHAP2/3^{RNAi} BF cell line was induced with 1 µg/ml doxycycline

(dox). Parasites were stained with DAPI (*blue*) and anti-*Tb*KH pAb (*Tb*KH, *red*), and mNG

endogenous fluorescence was also acquired (*Tb*CaCh::mNG*, green*). Parasites were induced

with doxycycline for (D) 0 days (*0 d dox*), (E) 4 days (*4d dox*), or (F) 7 days (*7d dox*). *Green*

arrowheads indicate flagella where *Tb*CaCh::mNG is present. DIC images are indicated. The

scale bar shown in (A) applies to all images.

Fig. 8. Trafficking of *Tb*EVMP1, *Tb*EVMP2, and *Tb*FLA1BP in *Tb*KH^{RNAi} BF parasites. (A)

Parasites expressing *Tb*EVMP1:mNG before (*0 h RNAi*) and after (*24 h RNAi*) induction of RNAi

against *TbKH* mRNA. *Green arrowheads* indicate mNG fluorescence in the FM (all panels) or in

extracellular vesicles (*0 h RNAi*, right panel). *White arrowheads* indicate flagella without mNG

fluorescence. (B) Parasites expressing *Tb*EVMP2::mNG before (*0 h RNAi*) and after (*24 h*

RNAi, 48 h RNAi) induction of RNAi against *TbKH* mRNA. (C) Parasites expressing

*Tb*FLA1BP::mNG before (*0 h RNAi*) and after (*24 h RNAi, 48 h RNAi*) induction of RNAi against

TbKH mRNA. The left-most images in A, B, and C represent formaldehyde fixed parasites,

whereas for the other images, live parasites were suspended in CyGel, which facilitates

visualization of secreted extracellular vesicles.

Table 1. List of Proteins with Gene IDs.
853

Gene ID	Protein
Tb927.10.8940	*Tb*KH
Tb927.10.2880	*Tb*CaCh/FS179
Tb927.11.2610	*Tb*KHAP1
Tb927.10.10360	*Tb*KHAP2/MARP-1
Tb927.10.10280	*Tb*KHAP3/MARP-2
Tb927.7.4270	*Tb*EVMP1/Fam79.3
Tb927.11.1830	*Tb*EVMP2
Tb927.8.4050	*Tb*FLA1BP

854
855

Figure 2

Figure 3

Figure 4

*Tb*CaCh::HA₃ *Tb*KH DAPI

Figure 8