The *Trypanosoma brucei* Cytoskeletal Protein KHARON Associates with Partner Proteins to Mediate Both Cytokinesis and Trafficking of Flagellar Membrane Proteins

Marco A. Sanchez, and Scott M. Landfear*

Department of Molecular Microbiology & Immunology, Oregon Health & Science University, Portland, Oregon, USA

*Author for correspondence (landfear@ohsu.edu)

Summary Statement: This study investigates the essential role in African trypanosomes of the KHARON protein and its molecular partners in trafficking of membrane proteins to the flagellum.

ABSTRACT

In the African trypanosome *Trypanosoma brucei*, the cytoskeletal protein *Tb*KHARON is

required for trafficking of a putative Ca^{2+} channel to the flagellar membrane, and it is

essential for parasite viability in both the mammalian stage bloodstream forms and the

tsetse fly procyclic forms. This protein is located at the base of the flagellum, in the

pellicular cytoskeleton, and in the mitotic spindle in both life cycle forms, and it likely

serves multiple functions for these parasites. To begin to deconvolve the functions of

KHARON, we have investigated partners associated with this protein and their roles in

parasite biology. One KHARON associated protein, *Tb*KHAP1, is a close interaction

partner that can be crosslinked to KHARON by formaldehyde and pulled down in a

molecular complex, and it colocalizes with *Tb*KHARON at the base of the flagellum.

Knockdown of *TbKHAP1* mRNA has similar phenotypes to knockdown of its partner

*Tb*KHARON, impairing trafficking of the Ca^{2+} channel to the flagellar membrane and

blocking cytokinesis, implying that the *Tb*KHARON/*Tb*KHAP1 complex mediates

trafficking of flagellar membrane proteins. Two other KHAPs, *Tb*KHAP2 and *Tb*KHAP3,

are in close proximity to *Tb*KHARON but may not be direct interaction partners, and

knockdown of their mRNAs does not affect trafficking of the Ca^{2+} channel. Two different

flagellar membrane proteins, which are extruded from the flagellar membrane into

extracellular vesicles, are also dependent upon *Tb*KHARON for flagellar trafficking.

These studies confirm that *Tb*KHARON acts in complexes with other proteins to carry

out various biological functions, and that some partners are involved in the core activity

of targeting membrane proteins to the flagellum.

KEYWORDS   *Trypanosoma brucei*, cytoskeleton, cytokinesis, flagellar membrane

protein trafficking, protein complexes, virulence

INTRODUCTION

African trypanosomes of the species *Trypanosoma brucei* are parasitic protists that cause human

African trypanosomiasis and the disease nagana in cattle and are thus of great medical and

veterinary importance (Kennedy, 2013). In addition, these parasites have been recognized as

valuable models for probing fundamental questions in cell and molecular biology (Cayla et al.,

2019). *T. brucei* and related kinetoplastid parasites such as *Trypanosoma cruzi* and *Leishmania*

species are flagellated and offer novel insights into the structure and function of flagella

(Langousis and Hill, 2014) and the roles of these organelles in infection (Kelly et al., 2020a).

Furthermore, the cell division cycle of African trypanosomes has been studied extensively (Farr

and Gull, 2012; Vaughan and Gull, 2008; Wheeler et al., 2019), identifying various processes that

are important for both proliferative and differentiation-linked cell division.

In previous work on trafficking of integral membrane proteins to flagella, we identified a

kinetoplastid-specific protein designated KHARON (KH) that plays a critical role in flagellar

targeting of the putative Ca^{2+} channel *Tb*CaCh (Tb927.10.2880) in *T. brucei*, originally identified

as a flagellar surface protein FS179 (Oberholzer et al., 2011), and the flagellar glucose transporter

*Lmx*GT1 in *L. mexicana* (Tran et al., 2013) (see Table S1 for tabulation of gene IDs and names

of *T. brucei* proteins investigated in this study). In both species of parasite (Sanchez et al., 2016),

KHARON was localized to three distinct subcellular compartments: the base of the flagellum (Fig.

1A), the subpellicular microtubules that subtend the plasma membrane around the cell body, and

the mitotic spindle (Fig. 1B). Application of RNA interference (RNAi) to knock down *TbKH*

(Tb927.10.8940) revealed that, in addition to preventing flagellar trafficking of *Tb*CaCh/FS179,

the flagellum attachment zone (FAZ) was disrupted, resulting in detachment of flagella from the

cell body (Sanchez et al., 2016), leaving this organelle adherent only through its connection at

the flagellar pocket. Interfering with trafficking of *Tb*CaCh/FS179 to the flagellar membrane is

likely to induce disruption of flagellar attachment, as RNAi directed against this channel also

results in a similar flagellar detachment phenotype (Oberholzer et al., 2011). In addition, as found

for many genetic alterations that disrupt flagellar attachment, these parasites are unable to initiate

cell division, generating trypanosomes in which nuclei, kinetoplasts (mitochondrial DNA-

containing structures), basal bodies, and flagella have replicated but cytokinesis has not occurred.

This phenotype was apparent in both mammalian bloodstream form (BF) and insect stage

procyclic form (PF) parasites and was thus lethal to both life cycle stages.

Similar studies in *L. mexicana* have also established a critical role for *Lmx*KH (LmxM.36.5850)

in the life cycle of *Leishmania* parasites. Thus a Δ*lmxkh* null mutant was generated in insect stage

promastigotes, where trafficking of *Lmx*GT1 to the flagellum was strongly impaired (Tran et al.,

2013), but cell division and replication of this life cycle stage was not affected. In contrast, Δ*lmxkh*

null mutants were unable to undergo cytokinesis after invading host macrophages, resulting in

formation of multinucleate multiflagellated amastigotes that died over the course of several days

(Tran et al., 2013). These null mutants were also avirulent following injection into BALB/c mice

(Tran et al., 2015), and studies by others have shown that *KHARON* null mutants in *L. infantum*

have potential as a live attenuated vaccine (Santi et al., 2018).

KHARON exhibits both similarities and striking differences between *T. brucei* (*Tb*KH) and *L.*

mexicana (*Lmx*KH). Thus, the two orthologs are relatively divergent in sequence, sharing 27%

amino acid identity and differing significantly in length (411 amino acids for *Tb*KH versus 520

amino acids for *Lmx*KH). *Tb*KH is critical for cell division of both mammalian BF and insect stage

PF parasites, whereas *Lmx*KH is only essential for division of disease-causing amastigotes.

Nonetheless, the three subcellular locations for KHARON are shared between the two parasites,

as are functions in cytokinesis and formation of the flagellar membrane.

KHARON proteins do not share significant sequence similarity to proteins outside the

Kinetoplastida, nor do they contain conserved sequence motifs that are suggestive of specific

biochemical or cellular functions. Furthermore, their residence at multiple subcellular locations

suggests that KHARON proteins are likely to be multifunctional, participating in flagellar

membrane trafficking, cytokinesis, and spindle function. Additionally, it is possible that distinctions

in function could be conferred by association of KHARON with different partner proteins at each

of its three subcellular loci. Thus, we hypothesize that there could exist three distinct KHARON

Complexes, Complex 1 at the base of the flagellum, Complex 2 at the subpellicular cytoskeleton,

and Complex 3 at the mitotic spindle. Furthermore, given the apparent differences in *Tb*KH and

*Lmx*KH noted above, there may be similarities and differences between these putative complexes

between the two species of parasite.

To initiate a study of putative KHARON Complexes and their functions, we carried out

biotinylation proximity labeling (BioID) (Roux et al., 2012) and tandem affinity purification-mass

spectrometry (TAP-MS) (Kaiser et al., 2008) on *Lmx*KH, resulting in the identification of two

KHARON Associated Proteins, *Lmx*KHAP1 and *Lmx*KHAP2 (LmxM.32.2440 and LmxM.05.0380,

respectively; (Kelly et al., 2020b)). In parallel, we investigated these two KHARON partners in *T.*

brucei and report the results of those studies here. As anticipated, *Tb*KHAP1 and *Tb*KHAP2

exhibit both similarities and notable differences compared to their orthologs in *L. mexicana*. In

addition, *T. brucei* expresses another KHARON partner related to *Tb*KHAP2 that we designate

*Tb*KHAP3. Furthermore, characterization of additional flagellar membrane proteins suggests that

*Tb*KH expression is important for flagellar targeting of multiple such proteins in African

trypanosomes, whereas the role of *Lmx*KH in trafficking of flagellar membrane proteins appears

to be more restricted. These studies confirm that KHARON proteins in both parasites exist in

complexes with various partners and that these partner proteins can play distinct roles in the

functions of different KHARON Complexes.

RESULTS

Localization of *Tb*KHAP1, *Tb*KHAP2, TbKHAP3, and *Tb*KH in bloodstream and procyclic African trypanosomes. To facilitate studies on *Tb*KH and its partners, we raised and affinity purified a polyclonal antibody against this protein, anti-*Tb*KH pAb. Western blot analysis indicated that anti-*Tb*KH pAb detects a single protein of ~49 kDa molecular weight, and that an additional band of ~62 kDa appears in parasites also expressing a BirA* fusion on the N-terminus of *Tb*KH (Fig. 1 C), establishing that this antibody is of suitable specificity to employ in localization and biochemical characterization of *Tb*KH. To determine whether *Tb*KHAP1 (Tb927.11.2610), *Tb*KHAP2 (Tb927.10.10360) and *Tb*KHAP3 (Tb927.10.10280) are associated with *Tb*KH in a complex, several complementary approaches were applied. First, each KHAP was tagged at its N-terminus with the triple hemagglutinin peptide tag HA$_3$, and formaldehyde-fixed BF and PF trypanosomes were examined by immunofluorescence deconvolution microscopy (Fig. 2). HA$_3$::*Tb*KHAP1 (Fig. 2A,B, green) overlaps with *Tb*KH (red) at the cell periphery, as demonstrated by the yellow color in this region of both BF and PF parasites. In contrast, there was no apparent overlap of the two signals in the mitotic spindle (central red oval or line marked with a white arrow), indicating that this protein could be associated with *Tb*KH in the subpellicular cytoskeleton but not at the mitotic spindle. Similarly, *Tb*KHAP2::HA$_3$ (Figs. 2C,D) and V5$_3$::*Tb*KHAP3 or HA$_3$::*Tb*KHAP3 (Fig. 2E,F) overlap with *Tb*KH at the cell periphery but not at the mitotic spindle. For each of the *Tb*KHAPs, there is also green fluorescence that does not coincide with *Tb*KH so that there is not complete overlap of the signals, and there may thus be populations of each protein that are not associated with each other. However overall, these three *Tb*KHAPs are candidates for *Tb*KH partners that are selective for the subpellicular cytoskeleton versus the mitotic spindle.

To determine whether *Tb*KHAP1, *Tb*KHAP2, or *Tb*KHAP3 might associate with *Tb*KH at the base of the flagellum, flagella were isolated from parasites expressing each HA$_3$- or V5$_3$-tagged *Tb*KHAP and imaged by deconvolution microscopy. Fig. 3 shows that HA$_3$::*Tb*KHAP1 (Fig. 3A), TbKHAP2::HA$_3$ (Fig. 3B) and HA$_3$::TbKHAP3 (Fig. 3C) overlap significantly with *Tb*KH in the region of the flagellum immediately adjacent to the kinetoplast DNA (kDNA, blue, Fig. 3A), which is close to and physically attached to (Robinson and Gull, 1991) the flagellar basal body. Overall, these results indicate that *Tb*KHAP1, *Tb*KHAP2, and *Tb*KHAP3 could be partners for *Tb*KH at both the pellicular cytoskeleton and the base of the flagellum. It is noteworthy that *Tb*KHAP3 was previously identified as a basal body protein in a proteomic study of that subcellular structure (Dang et al., 2017). Hence, it is likely that *Tb*KH, *Tb*KHAP1, *Tb*KHAP2, and *Tb*KHAP3 are all basal body components.

Molecular association of *Tb*KHAP1, *Tb*KHAP2, and *Tb*KHAP3 with *Tb*KH. To determine whether the observed subcellular overlap of the fluorescence signals from *Tb*KHAPs and *Tb*KH could indicate physical association in molecular complexes, *Tb*KH was endogenously tagged at its N-terminus with a His_{10} affinity tag (His_{10}::*Tb*KH) to allow pulldown of this protein, and associated partners, with Co^{2+} agarose resin, and this affinity tagged protein was expressed in a BF cell line also expressing either HA_3::*Tb*KHAP1, *Tb*KHAP2:: HA_3, or $V5_3$::*Tb*KHAP3. Because *Tb*KH is an integral component of the parasite cytoskeleton (Sanchez et al., 2016) and would pulldown many cytoskeletal proteins in an experiment performed under native conditions, we first crosslinked with formaldehyde parasites expressing each pair of tagged proteins. Formaldehyde crosslinks proteins that are in very close proximity (~2-3 Å, reference (Hoffman et al., 2015)), so this treatment will covalently attach close molecular partners of *Tb*KH but not proteins that are more distant partners in a complex or proteins that are in the cytoskeleton but distant from *Tb*KH. Subsequent treatment with strongly denaturing reagents will dissociate peripheral proteins from His_{10}::*Tb*KH while retaining crosslinked partners, and the closely associated partners will thus be purified along with His_{10}::*Tb*KH, following binding and elution from Co^{2+} resin, and released upon heat-induced reversal of the crosslinks.

Fig. 4A demonstrates that HA_3::*Tb*KHAP1 is pulled down with His_{10}::*Tb*KH when parasites are formaldehyde crosslinked (EF*) but not when they are not subjected to crosslinking (EF). As a negative control, another subpellicular cytoskeletal protein, CAP15 (Vedrenne et al., 2002), was HA_3 tagged and expressed in His_{10}::*Tb*KH expressing BF parasites, but this protein was not pulled down even from formaldehyde-crosslinked parasites. Parallel experiments were performed using BF parasites expressing *Tb*KHAP2::HA_3, or $V5_3$::*Tb*KHAP3 and His_{10}::*Tb*KH. Despite extensive efforts, we have not been able to observe pulldowns of either tagged protein with His_{10}::*Tb*KH, raising the possibility that these proteins are not in close enough contact to *Tb*KH to crosslink. These experiments establish that *Tb*KHAP1 is in very close proximity to *Tb*KH and these two proteins are thus molecular partners.

Proximity ligation assay (PLA) confirms close proximity of *Tb*KH with *Tb*KHAP1, *Tb*KHAP2, and *Tb*KHAP3. To provide another independent examination of whether *Tb*KH is in close physical proximity with each *Tb*KHAP, we performed the PLA in parasites expressing HA_3::*Tb*KHAP1, *Tb*KHAP2::HA_3, and HA_3::*Tb*KHAP3. In this assay (Fredriksson et al., 2002; Soderberg et al., 2006), cells expressing two partner proteins are first reacted with primary antibodies from different species. Parasites are subsequently incubated with species-specific secondary antibodies directed against each primary antibody, and each of these secondary antibodies contains a unique, covalently attached oligonucleotide. Only if the two target proteins

are within ~400 Å of each other, these oligonucleotides can base pair to another linker
oligonucleotide and be covalently ligated into a circular substrate that can participate in rolling
circle DNA amplification of the cognate sequence. The amplified sequence is then hybridized to
a fluorescently labeled DNA probe, resulting in fluorescent puncta within the cell.

Fig. 4B,C shows a positive PLA signal (left panels) for BF and PF trypanosomes expressing
HA$_3$::*Tb*KHAP1 and probed with anti-*Tb*KH rabbit and anti-HA murine mAb. In contrast, when the
anti-*Tb*KH antibody is not employed (right panels), the PLA signal is absent, demonstrating the
dependency of the signal on detection of both closely associated proteins. Similar results confirm
that *Tb*KH is in close physical proximity to *Tb*KHAP2 (Fig. 4D,E) and *Tb*KHAP3 (Fig. 4F,G).

Predicted properties of *Tb*KHAP1, *Tb*KHAP2, and *Tb*KHAP3. Bioinformatic analysis of the
50.9 kDa *Tb*KHAP1 sequence indicates that it is a protein apparently unique to kinetoplastid
protists for which there are orthologs widely distributed among Kinetoplastida. A BLASTP search
(tritrypdb.org) revealed several coiled-coil proteins such as neurofilament proteins and
tropomyosin as being significantly, although not closely, related (E values of 2.6e-08 – 5.4e-12).
Prediction of protein disorder using the PrDOS web server (Ishida and Kinoshita, 2007)
(http://prdos.hgc.jp/cgi-bin/top.cgi) generated a strong prediction of disorder (probability >0.9)
over the C-terminal region from amino acids 314 – 461. Indeed, this sequence is rich in E
residues, which predispose such regions to intrinsic disorder (Uversky, 2013), and this property
suggests that this region could be involved in protein-protein interactions through induced folding
(Zhang et al., 2013). InterPro (Mitchell et al., 2015)
(https://www.ebi.ac.uk/interpro/search/sequence/) predicted coils between amino acids 7 – 31
and 196 – 241 and a disordered region from residue 332 - 461 , and PSIPRED V4.0 (McGuffin et
al., 2000) (http://bioinf.cs.ucl.ac.uk/psipred/) predicted the sequence to be largely helix or coil.
Overall, computational analyses suggest that TbKHAP1 is a coiled-coil protein with an intrinsically
disordered C-terminus, both properties that are consistent with formation of multi-protein
complexes.

*Tb*KHAP2 is the 374 kDa microtubule-associated repetitive protein 1, MARP-1, and *Tb*KHAP3
is the 267 kDa MARP-2 that have been studied previously by Seebeck and colleagues (Affolter
et al., 1994; Hemphill et al., 1992; Schneider et al., 1988) and will hereafter be referred to as
*Tb*KHAP2/MARP-1 and *Tb*KHAP3/MARP-2 to indicate both their association with *Tb*KH and their
previously demonstrated roles in microtubule binding. Each sequence contains short unique N-
and C-terminal domains, and the remainder of the sequence consists of 38-amino acid repeats
that are largely conserved within each sequence but ~50% identical between the two proteins.
The unique C-terminal domains (95% identical between the two proteins) bind to microtubules

(Affolter et al., 1994), and the proteins decorate the subpellicular cytoskeleton (Schneider et al.,
1988), but their specific biological functions have not been elucidated. In addition, these proteins
have also been localized to the basal body and *Tb*KHAP3/MARP-2 was designated *Tb*BBP268
(Dang et al., 2017).

Phenotypes of BF trypanosomes following knockdown of *TbKHAP1* RNA. We have
previously demonstrated that knockdown of *TbKH* RNA by inducible RNAi results in a lethal
phenotype on both BF and PF trypanosomes (Sanchez et al., 2016). In these parasites, the
flagellum detaches from the cell body along the flagellum attachment zone (FAZ), and the
parasites are blocked in cytokinesis, resulting in accumulation of multi-nucleated, multi-flagellated
'monster cells' that are not viable in the long term. To assess the roles of *Tb*KHAP1,
*Tb*KHAP2/MARP-1, and *Tb*KHAP3/MARP-2 in the biology of BF parasites, we targeted by
RNAi *TbKHAP1* mRNA, using a unique RNAi probe, and *TbKHAP2/TbKHAP3* mRNAs jointly,
using a 500 nt probe covering the conserved C-termini, and assessed the consequent
phenotypes.

Induction of RNAi against *Tb*KHAP1 in BFs using doxycycline resulted in rapid reduction in
the level of this protein (Fig. 5A). Furthermore, RNAi-induced parasites stopped growing almost
immediately and were largely dead by 72 h (Fig. 5B). Quantification via microscopy of the
percentage of cells with different numbers of nuclei and kinetoplasts (Fig. 5C) showed that
following induction of RNAi over 48 h, the percentage of 1N1K parasites dropped dramatically,
while those with multiple nuclei and kinetoplasts (XNYK) increased and began to predominate the
population. In comparison to the normal morphology of pre-induced parasites (Fig. 5D), induction
of RNAi for 20 h (Fig. 5E) or 48 h (Fig. 5F) resulted in parasites with multiple nuclei and/or
kinetoplasts and tadpole-like morphology (white arrowhead) or duplicated flagella located at
opposite poles of the cell body (yellow arrowhead).

Notable among cells in RNAi induced populations are those with multiple flagella located at
various relative positions around the cell (e.g., the two parasites in Fig. 5F). Such parasites have
initiated cytokinesis and cleavage furrow formation, as the two duplicated flagella have moved
apart from the initial position they would have following flagellar duplication. However, the
cleavage furrow did not progress to separate the duplicated nuclei and kinetoplasts as it would in
normal cell division. In summary, these observations suggest that loss of *Tb*KHAP1 protein from
BF parasites results in a block in progression of cleavage furrow rather than an inability to initiate
cleavage furrow ingression.

These results are further enhanced by more refined time course studies shown in Fig. S1
following the progression of nuclear content and cell morphologies between 0 – 48 h after

induction of RNAi against *TbKHAP1* RNA. At 4 h (Fig. S1B) and 8 h (Fig. S1C) most parasites

had morphologies similar to that preceding induction of RNAi (0 h, Fig. S1A). However, by 12 h

(Fig. S1D), multi-flagellated parasites with ingression furrows appeared, and by 24 h and 48 h

(Fig. S1E,F), many parasites had incompletely resolved ingression furrows and multiple nuclei.

Phenotypes of parasites following knockdown of *TbKHAP2/MARP-1* and

***TbKHAP3/MARP-2* RNAs.** Induction of RNAi jointly against *TbKHAP2/MARP-1* and

TbKHAP3/MARP-2 resulted in complete loss of HA$_3$::*Tb*KHAP2/MARP-1 protein by 24 h (Fig.

6A). Depletion of *Tb*KHAP2/MARP-1 and *Tb*KHAP3/MARP-2 impaired growth of BF parasites,

resulting in an ~50-fold reduction in parasite number by 120 h (Fig. 6B) post-induction, but growth

inhibition was not nearly as pronounced as it is for *Tb*KHAP1 RNAi (Fig. 5B). Compared to

uninduced parasites (Fig. 6C), images of parasites following 4 d RNAi (Fig. 6D) still showed many

parasites with normal morphology similar to that of uninduced parasites, but some parasites

rounded up and showed multiple flagella (Fig. 6D, white arrowhead, flagella on opposite sides of

the cell body in DIC image). By 4 d post-RNAi, parasites with 1N2K and XNYK began to

accumulate (Fig. 6E), but the proportion was not nearly as great as for RNAi directed against

TbKHAP1 (Fig. 5C).

***Tb*KHAP1, but not *Tb*KHAP2/MARP-1 or *Tb*KHAP3/MARP-2, is required for targeting**

***Tb*CaCh/FS179 to the flagellar membrane.** The observation that *Tb*KHAP1 is located at the

base of the flagellum (Fig. 3A), likely in the basal body, raises the question of whether it could

play a role in the function of *Tb*KH in mediating trafficking of the putative Ca^{2+} channel,

*Tb*CaCh/FS179, to the flagellar membrane. To test this possibility, we induced RNAi against

TbKHAP1 RNA in parasites expressing *Tb*CaCh/FS179::HA$_3$ tagged at the C-terminus, which

localized to the flagellar membrane prior to RNAi (Fig. 7A). BF parasites induced for RNAi for 24

h (Fig. 7B) or 48 h (Fig. 7C) exhibited flagella that were devoid of *Tb*CaCh::HA$_3$ (white

arrowheads). These results suggest that a complex of *Tb*KH/*Tb*KHAP1, and potentially other

currently unknown partners located at the base of the flagellum, is involved in trafficking this

channel to the flagellar membrane. Since both proteins are also located in the subpellicular

cytoskeleton, it is not possible to definitively ascribe this flagellar trafficking phenotype to the

complex at the base of the flagellum; complexes at both locations will be downregulated by

TbKHAP1 RNAi. However, integral membrane proteins are first delivered to the flagellar pocket

membrane during biosynthesis (Manna et al., 2014). Hence, the presence of a protein complex

located close to the interface between the flagellar pocket and flagellar membrane, and for which

downregulation of both known partners inhibits trafficking of a protein into the flagellar membrane,

suggests that this complex may mediate trafficking of *Tb*CaCh/FS179 from the flagellar pocket

membrane into the flagellar membrane. Such trafficking would presumably be mediated by a direct interaction between *Tb*KH and the cargo, *Tb*CaCh/FS179, and such a molecular interaction has been demonstrated to occur (Sanchez et al., 2016).

In contrast, RNAi directed against *TbKHAP2/MARP-1* and *TbKHAP3/MARP-2* RNAs did not prevent trafficking of *Tb*CaCh/FS179::HA$_3$ into the flagellar membrane, where it is located prior to RNAi (Fig. 7D). At both 4 d (Fig. 7E) and 7 d (Fig. 7F) post RNAi, BF parasites with multiple flagella still trafficked this channel into the flagellar membrane (green arrowheads).

_Tb_KH-dependent trafficking of other flagellar membrane proteins. *Tb*KH is required for trafficking of *Tb*CaCh/FS179 to the flagellar membrane of BF trypanosomes, and the two protein interact with each other, as demonstrated by crosslinking pulldown assays (Sanchez et al., 2016). Is *Tb*KH important for flagellar trafficking of other membrane proteins? To address this question, we monitored the dependency of other flagellar membrane proteins on *Tb*KH for targeting to that organelle. The TrypTag project (http://tryptag.org/) has defined the subcellular location of a large number of trypanosome proteins in PF parasites (Dean et al., 2017), employing live cell microscopy of parasites expressing mNeonGreen fluorescent protein fusions, and this endeavor has identified a cohort of flagellar membrane proteins.

One such flagellar membrane protein is Tb927.7.4270, a 25 kDa protein predicted to have a N-terminal signal sequence and a single transmembrane domain (TMD) near its C-terminus. This protein is one of four paralogous proteins (Tb927.7.4230, 4260, 4270, and 4280) studied previously by Shimogawa *et al*. (Shimogawa et al., 2015) and designated the Fam79 proteins (Fam79.1, 79.2, 79.3, and 79.4, respectively). To address potential dependency upon *Tb*KH for flagellar targeting, we monitored the localization of the C-terminal mNG fusion of Tb927.7.4270/Fam97.3 in both formaldehyde fixed and live PF parasites immobilized in CyGEL (MacLean et al., 2013). As shown in Fig. 8A (0 h RNAi), this fusion protein is present in flagella and also in filaments and vesicles that emerge from the flagella, and we designate Tb927.7.4270 as extracellular vesicle membrane protein 1 or *Tb*EVMP1/Fam79.3. Multiple investigators have observed filaments and vesicles emerging from various parts of trypanosomes (Baudieri and Tomassini, 1962; Ellis et al., 1976; Molloy and Ormerod, 1964; Schepilewsky, 1912; Vickerman and Luckins, 1969; Wright and Lumsden, 1970), including the flagella, and a recent study by Szempruch *et al*. (Szempruch et al., 2016a; Szempruch et al., 2016b) has investigated such structures from BF parasites in detail and concluded that the flagellum-derived nanotubes and resulting extracellular vesicles (EVs) incorporate a cohort of parasite proteins. Furthermore, delivery of parasite-derived EVs to host red blood cells or to other trypanosomes can mediate pathogenic processes, such as erythrocyte clearance and anemia in the mammalian host or

delivery of innate immune factors from a resistant to a sensitive strain of trypanosome. Hence,
understanding the process for delivery of parasite proteins to these EVs is of importance for
deciphering mechanisms of parasite virulence. Notably, when PF parasites expressing
*Tb*EVMP1::mNG were subjected to RNAi directed against *Tb*KH for 24 h, they were strongly
impaired in trafficking of this fusion protein to flagella or nanotubes (Fig. 8A, 24 h RNAi), and
fluorescence often accumulated within the parasite cell body. White arrows indicate flagella that
are devoid of fluorescence and which thus exhibit a trafficking defect. This result indicates that
*Tb*EVMP1/Fam79.3 is dependent upon *Tb*KH, either directly or indirectly, for trafficking to the
flagellum and subsequently for release into EVs. Furthermore, this trafficking defect occurs after
24 h of RNAi directed against *Tb*KH, but these PF parasites do not exhibit significant loss of
viability until ~10 days of continuous RNAi (Sanchez et al., 2016), indicating that the effects of
TbKH RNAi upon flagellar trafficking are not due to global loss of cellular functions. Notably,
Fam79.1 (Tb927.7.4230) was also detected by proteomic analysis in EVs of BF trypanosomes by
Szempruch *et al*. (Szempruch et al., 2016b), indicating that multiple members of this family are
delivered to the membranes of EVs during the parasite life cycle.

A second flagellar membrane protein localized in TrypTag is Tb927.11.1830, designated here
as *Tb*EVMP2. This 62 kDa protein has 6 predicted TMDs and is widely distributed among the
kinetoplastid protists but does not have obvious orthologs outside that order, nor does it possess
conserved Pfam domains (Sonnhammer et al., 1998), except for the TMDs and one predicted
coiled coil. The *Tb*EVMP2::mNG fusion protein is also localized to the flagellar membrane,
nanotubes, and extracellular vesicles in PF parasites (Fig. 8B, 0 h RNAi), but induction of RNAi
directed against *Tb*KH also inhibits trafficking to these flagellar structures (Fig. 8B, 24 h RNAi),
albeit less strongly than for *Tb*EVMP1. Although fluorescence is visible in some flagella after
induction of RNAi (green arrows, righthand image for 24 h RNAI in Fig. 8B), there are some
flagella that exhibit little if any fluorescence (white arrows). We designate this protein *Tb*EVMP2
and suggest that it is a second such protein that is dependent upon *Tb*KH for efficient trafficking
to the surface of the flagellum.

In contrast, the mNG tagged FLA1 binding protein *Tb*FLA1BP::mNG, which is in the flagellar
membrane component of the FAZ (Sun et al., 2013), traffics efficiently to the FM both before and
after induction of RNAi (Fig. 8C, 0 h, 24 h, and 48 h RNAi), even in PF parasites that have multiple
flagella. Hence, *Tb*FLA1BP does not require *Tb*KH for targeting to the FM, implying that there are
both *Tb*KH-dependent and *Tb*KH-independent FM proteins in this parasite.

DISCUSSION

In the current study, we have investigated three partners of *Tb*KH: *Tb*KHAP1 and the two related proteins *Tb*KHAP2/MARP1 and *Tb*KHAP3/MARP2. *Tb*KHAP1 can be crosslinked to *Tb*KH by formaldehyde indicating that the two proteins associate within 2-3 Å of each other. The failure to demonstrate crosslinking of *Tb*KHAP2 or *Tb*KHAP3 to *Tb*KH could be due to their more distant location from each other, or to technical difficulties associated with these large highly repetitive proteins, but they are in close enough proximity to give a consistently positive signal using the PLA, that is within ~400 Å. One possibility is that they are part of a multiprotein complex but not in immediate contact with each other. The knockdown of *TbKHAP1* RNA has the most pronounced phenotype, strongly arresting division of BF parasites, inhibiting progression of the cleavage furrow during cytokinesis, and impairing trafficking of *Tb*CaCh/FS179 to the FM. Hence, the phenotypes of RNAi for both *TbKH* and *TbKHAP1* are similar. In contrast, efficient knockdown of *TbKHAP2/3* RNAs slows growth of BF parasites but has a much less severe effect on cell division than knockdown of *TbKHAP1* RNA. In addition, trafficking of *Tb*CaCh/FS179 to the FM is not impaired, even after 7 days of knockdown in BFs. These results imply that although both types of protein likely associate with *Tb*KH in the cytoskeleton, they play different roles. These distinctions in functions could either result from separate complexes between *Tb*KH and each partner or from different roles that each partner plays in the same complex. The association of all partners with the cytoskeleton complicates this issue, as one cannot readily separate different complexes from each other, as would be possible for cytosolic multi-protein complexes. Nonetheless, these studies confirm that *Tb*KH associates with other partner proteins that mediate its activities in different ways.

One central activity for *Tb*KH is to traffic *Tb*CaCh/FS179 to the FM, a process that is critical for integrity of the FAZ and for parasite viability. The localization of many PF proteins to their subcellular sites achieved in the TrypTag.org project (Dean et al., 2017) has identified some additional FM proteins, along with some others that were identified previously as FM components from various targeted studies (Kelly et al., 2020a), and one question of relevance is how many of these flagellar surface components rely upon *Tb*KH for organellar trafficking. For *Tb*CaCh/FS179, *Tb*KH appears to be directly involved in trafficking, since the two proteins can be crosslinked by formaldehyde and isolated as molecular partners (Sanchez et al., 2016), but it is possible that others depend upon *Tb*KH either directly or indirectly via the ability of this protein to affect various cellular processes. In this study, we have shown that two FM proteins, *Tb*EVMP1/Fam79.3 (Tb927.7.4270) and *Tb*EVMP2 (Tb927.11.1830) are present in both the FM and in EVs secreted from the FM in PF trypanosomes. Both proteins exhibit dependency upon *Tb*KH for trafficking to

the FM, as RNAi directed against *TbKH* reduces the efficiency of their localization to this

organelle. In contrast, FLA1BP, which participates in adhesion of the flagellum to the cell body by

binding to the FLA1 protein in the cell body component of the FAZ (Sun et al., 2013), is not

dependent upon *Tb*KH to reach the FM, confirming that both *Tb*KH-dependent and *Tb*KH-

independent FM proteins exist.

EVs released from the cell body and FM of BF trypanosomes play important roles in virulence

of African trypanosomes, including lysis of host erythrocytes leading to anemia, a major

mechanism of trypanosome-mediated pathogenesis (Szempruch et al., 2016a; Szempruch et al.,

2016b). One might anticipate that surface components of EVs could play important roles in either

formation of the EV membrane or interaction of EVs with mammalian or tsetse fly tissues.

TbEVMP1 and *TbEVMP2* mRNAs are both preferentially expressed in PF trypanosomes

(tritrypdb.org), but paralogs of *Tb*EVMP1, such as Tb927.7.4230 and Tb927.7.4260, are

expressed at higher levels in BFs compared to PFs, suggesting potential roles for such EVMPs

in both life cycle stages.

The *Tb*KH partners discovered in this study are associated primarily with the subpellicular

microtubules, but there are likely to be other partners that may reside principally at the base of

the flagellum or in the mitotic spindles and could be associated with distinct activities at those

sites. A BioID study by Zhou *et al*. (Zhou et al., 2018) identified five spindle-associated proteins,

NuSAP1, NuSAP2, Kif13-1, *Tb*MIP2, and *Tb*AUK1, that are in proximity to *Tb*KH and are thus

candidates for molecular partners at the mitotic spindle. In addition, Akiyoshi and Gull (Akiyoshi

and Gull, 2014) identified kinetoplast kinetochore proteins (KKTs) that associate with KHARON

by CoIP/MS experiments, suggesting a possible role of KHARON in faithful chromosome

segregation in kinetoplastid parasites. Molecular interaction studies of the type carried out here

will be required to determine which of these proteins may be *bona fide* molecular partners with

*Tb*KH and what roles KHARON complexes may be playing at the spindle. Similarly, the ability to

isolate flagella with associated kinetoplast DNA and basal bodies (Oberholzer et al., 2011;

Robinson and Gull, 1991; Subota et al., 2014) should facilitate identification by either BioID or

TAP-MS of *Tb*KH partners at the base of the flagellum. Thus, it should be possible to achieve a

comprehensive understanding of the role of KHARON in the cytoskeleton and the distinct

functions it carries out in association with different partner proteins.

MATERIALS AND METHODS

Growth and transfection of *T. brucei* cell lines. BF and PF *T. brucei* cell lines were grown as described previously (Sanchez, 2013). For T7 RNA polymerase-independent expression, the BF/pHD1313 or PF/pHD1313 clones were employed (Sanchez et al., 2016). For T7 RNA polymerase driven expression in BF, *T. brucei* 427 parasites transfected with the pSmOx (Poon et al., 2012) plasmid expressing the tetracycline repressor and T7 RNA polymerase were generated and grown in 0.1 μg/ml puromycin. For T7 RNA polymerase driven expression in PF, *T. brucei* 427 13-6 clone expressing TETR and T7 RNA polymerase was used (Wirtz et al., 1999). Linear plasmid or PCR DNA amplicons were used to transfect mid-log phase parasites as described (Sanchez et al., 2016). Transfected clones were obtained by limiting dilution according to published protocols (Burkard et al., 2007; McCulloch et al., 2004) .

Primary amino acid sequence analysis. For DNA and amino acid sequence analysis of *TbKHAPs*, ExPASy, via the SIB Bioinformatics Resource Portal (http://expasy.org), and GeneBank (http://blast.ncbi.nlm.nih.gov/Blast.cgi) or TritrypDB (http://tritrypdb.org/tritrypdb) were used.

Inhibition of gene expression by RNAi. To inhibit the expression of *Tb*KH, *Tb*KHAPs or *Tb*FLA1, a number of cell lines were generated employing different genetic backgrounds. The BF and PF TbKH[RNAi] and *Tb*FLA1[RNAi] clones, for which expression is T7 polymerase-independent, were previously described (Sanchez et al., 2016), and RNAi was induced by adding doxycycline (1 μg/ml) to the culture medium. The BF and PF *Tb*KHAP1[RNAi] clones were generated by subcloning the first 500 bp of the *TbKHAP1* (Tb927.10.1026) ORF into the pZJM RNAi vector, where expression is driven by two opposing T7 promoters (Wang et al., 2000). Similarly, BF and PF TbKHAP2/3[RNAi] clones were generated by subcloning the last 500 bp of *TbKHAP2* (Tb927.10.10360) that is almost identical to the *TbKHAP3* (Tb927.10.10280) ORF into the pZJM RNAi vector. RNAi clones were selected by resistance to 2.5 μg/ml phleomycin and 0.1 μg/ml puromycin, and expression of dsRNA was induced by addition of 1 μg/ml doxycycline. To verify inhibition of *Tb*KHAP1-3 expression, total cell lysates were obtained from parasite cultures induced for RNAi and subjected to Western blot experiments as indicated below.

Endogenous epitope tagging. For endogenous tagging of *Tb*KH, *Tb*KHAP1, *Tb*KHAP2, *Tb*KHAP3, CAP-15, *Tb*CaCh (FS179), *Tb*EVMP1, *Tb*EVMP2 and *Tb*FLA1BP at the N-terminus or C-terminus, different epitopes were employed as indicated in the text, following the protocol described (Dean et al., 2015). Briefly, epitope-tagging cassettes were generated by using two specific 100 nt oligonucleotides containing ~80 nt each that are homologous to the ORF region to be tagged and ~20 nt homologs to the plasmid pPOTV4 template flanking the drug resistance

marker cassette, and using the universal PCR settings. Epitope tagging PCR cassettes were

ethanol precipitated and resuspended in 10 µl of nucleofection buffer (Wang et al., 2000), then

parasites were transfected with the purified tagging cassettes as described (Dean et al., 2015)

and selected using 1.5 µg/ml G418 (15 µg ml^{-1} for PF) or 0.1 µg/ml puromycin (1 µg/ml for PF),

and cloned by limiting dilution.

Generation of Rabbit anti-*Tb*KH antibody. A custom rabbit anti-*Tb*KH polyclonal antibody

(pAb) was generated by GenScript, using their 49-day antibody generation protocol. Briefly, two

rabbits were injected with 200 µg of His$_6$::*Tb*KH, representing amino acids 43 - 411, emulsified in

Freund's complete adjuvant. The rabbit was boosted 3 times at 14-day intervals with 200 µg of

His$_6$::*Tb*KH emulsified in Freund's incomplete adjuvant. Antibody specificity for *Tb*KH was

evaluated by Western blot comparing the reactivities of the rabbit serum from immunized rabbits

to *T. brucei* protein lysates from wild-type cells and N-BirA*::*Tb*KH cell line (Fig 3A). Proteins were

immunodetected using 1:2500 dilution of the rabbit anti-*Tb*KH polyclonal antibody and 1:15,000

dilution of goat anti-rabbit-HRP antibody (Sigma-Aldrich). The chemiluminescent protocol was

used for developing as indicated below.

Immunofluorescence microscopy. For immunofluorescence microscopy, 5 X 10^6 parasites

were centrifuged at 1000 X g for 5 min and washed twice at room temperature with phosphate

buffered saline pH 7.2 (PBS) containing 10 mM glucose. The cell pellet was resuspended in 4%

paraformaldehyde in PBS, pH 7.2 and incubated for 15 min at room temperature, cells were

centrifuged as descried above and washed once with PBS, resuspended in 100 µl PBS, spotted

onto poly-L-lysine coated coverslips, and blocked with 2% goat serum, 0.01 sodium azide, 0.01

saponin in PBS (blocking solution) for 1 h at room temperature, rinsed 3 X with PBS, and

incubated with primary antibodies for 1 h at room temperature. The following primary antibodies

were employed: 1:250 dilution rabbit anti-*Tb*KH pAb (reported in this work), 1:500 dilution mouse

anti-HA monoclonal antibody (mAb) (BioLegend, Cat # MMS-101R), and 1:1000 dilution mouse

anti-α-tubulin mAb (Sigma-Aldrich, Cat. # T5168). Subsequently, cells were rinsed as before and

incubated with a 1:1000 dilution of secondary antibodies coupled to Alexa Fluor dyes (Molecular

Probes) as follows: Alexa Fluor® 488 goat anti-mouse IgG (H+L) (Cat. # A11001), Alexa Fluor®

594 goat anti-mouse IgG (H+L) (Cat. # A11005), Alexa Fluor® 594 goat anti-rabbit IgG (H+L)

(Cat. # A11012) and Alexa Fluor® 488 goat anti-rabbit IgG (H+L) (Cat. # A11008), as indicated,

in blocking solution for 1h at room temperature in the dark. Cover slips were rinsed 3 X with PBS

and mounted onto slides using DAPI Fluoromount-G (SouthernBiotech). Fluorescence images

were obtained using a wide field deconvolution system (Applied Precision Instruments, Inc.)

consisting of an inverted Nikon TE 200 Eclipse microscope, a Kodak CH350 CCD camera, and

the Deltavision operating system. Images were acquired using a 60 X objective and 1.25 X

magnification in a 1024 X 1024 format, and deconvolved using SoftWoRx software. Adobe

Photoshop CC and Adobe Illustrator CC (Adobe Systems Inc.) were used to create image

compositions.

Flagellar protein purification, formaldehyde crosslinking, pull down, and Western blot

assay. Flagellar purification was performed as described (Subota et al., 2014) with slight

changes. Briefly, BF or PF $TbKH1^{RNAi}$, $TbFLA1^{RNAi}$, $TbKHAP1^{RNAi}$ and $TbKHAP2/3^{RNAi}$ clones were

grown in 1 µg/ml doxycycline for the indicated times and pelleted at 420 X g for 10 min. Cell pellets

were washed with buffer A (25 mM Na^+-tricine, pH 7, 1% BSA, 0.1 mM $CaCl_2$, 0.2 mM EDTA, 5

mM $MgCl_2$ and 12 mM β-mercaptoethanol) containing 0.32 M sucrose and centrifuged at 420 X g

for 10 min. Cell pellets were gently resuspended at 3 X 10^8 parasites/ml in buffer A plus 0.3 M

sucrose, transferred into Eppendorf tubes and vortexed for 5 min or until microscopic verification

of flagellum detachment, followed by centrifugation at 420 X g for 10 min. Supernatants were

recovered and centrifuged at 16,000 X g for 20 min at 4 °C. Pellets containing the isolated flagella

were resuspended in 200 µl of PBS and used for immunofluorescence microscopy. Also,

purification of flagellar cytoskeletons from BF or PF clones was performed by isolation of whole

cytoskeletons followed by treatment with 1 mM $CaCl_2$ to solubilize the pellicular cytoskeletons,

according to Imhof et al. (Imhof et al., 2019).

For formaldehyde crosslinking and pull down, cell lines were prepared that expressed $TbKH1$

endogenously tagged with the His_{10} epitope at the N-terminus (His_{10}::$TbKH1$) and also co-

expressed with a protein of interest endogenously tagged at its N- or C-terminus with a HA_3 or

$V5_3$ epitope (HA_3::$TbKHAP1$, HA_3::$TbKHAP2$, $TbKHAP2$::HA_3, $V5_3$::$TbKHAP3$, HA_3::$TbKHAP3$

and $TbCAP15$::HA_3). Parasites were washed once with PBS and pelleted at 1000 X g for 10 min,

and cell pellets resuspended in 9.37 ml PBS plus 0.63 ml 16% formaldehyde-EM grade

(Polysciences, Inc) and incubated at room temperature for 10 min. For non-crosslinked control

samples, PBS was added instead of formaldehyde. Subsequently, 1 ml of 2.5 M glycine in PBS

was added to the crosslinked samples and incubated at room temperature for 5 min, crosslinked

parasites were pelleted at 1000 X g for 10 min and washed twice with PBS. Then, cell pellets with

or without crosslinking were resuspended in 1 ml of Buffer 1 (8 M urea, 300 mM NaCl, 0.5%

Nonidet P-40, 50 mM NaH_2PO_4, 50 mM Tris, pH 7.0) on ice. Samples were sonicated 3 times on

ice at 50% max amplitude (Sonic Dismembrator, 500W, Fisher Scientific) for 10 s with 30 s

between pulses. A 2.5% aliquot of this protein lysate was saved as the protein lysate fraction (LF).

The remainder of the protein was incubated with 750 µl of HisPur^{TM} Cobalt Resin (Thermo

Scientific Pierce, Rockford, IL) on a rocker for 45 min at room temperature. The resins were

washed 5 times with 1 ml of Buffer 1 (5 min each), 5 times with 1 ml of Buffer 1 at pH 6.4 (5 min

each), and 5 times with 1 ml of Buffer 1 pH 6.4 plus 10 mM imidazole (5 min each). Cobalt bound

protein complexes were eluted with 1 ml of Buffer 2 (45 mM NaH_2PO_4, 8 M urea, 270 mM NaCl,

150 mM imidazole). Eluate fractions (EF) were concentrated using Amicon Ultra-0.5 Centrifugal

Filter Units, 10K NMWL (Millipore Sigma). Crosslinking was reversed by boiling protein samples

for 30 min in 1X Bolt™-LDS sample buffer containing 10 mM DTT.

Protein extracts were prepared and analyzed by Western blot employing Bolt™ 4-12% Bis-Tris

Mini Protein Gels or NuPage™ 3-8% Tris-acetate gels, Mini Gel Tank and Mini Blot Module

following the manufacturer's instructions (Life Technologies). Proteins were transferred onto

PVDF membranes (Millipore). Protein immunodetection was done using rabbit anti-*Tb*KH pAb at

1:2500 dilution, mouse anti-HA mAb (BioLegend, Cat # MMS-101R) 1:2500 dilution, mouse anti-

V5 mAb (Invitrogen, Cat # MA5-15253) 1:2500 dilution, and mouse anti-α tubulin mAb (Sigma-

Aldrich, Cat. # T5168) 1:10,000 dilution. Goat anti-rabbit-HRP (Sigma-Aldrich) 1:15,000 dilution

and goat anti-mouse-HRP (Jackson ImmunoResearch Laboratories, Cat. # 115-03-174, Lot #

117119) were used as secondary antibodies and Western blots were developed using the

SuperSignal™ West Pico Plus Chemiluminescent Substrate (Thermo Fisher Scientific) and an

Image Quant LAS 400 (GE Healthcare) scanner was employed to acquire luminescent images.

Adobe Photoshop CC and Adobe Illustrator CC (Adobe Systems Inc.) were used to create image

compositions.

Proximity ligation assay (PLA). Mid-log BF or PF parasites were harvested by

centrifugation at 1000 x g for 10 min, washed once in PBS and fixed with 4% paraformaldehyde

in PBS for 15 min at room temperature. Fixed cells were attached to cover slips and

permeabilized as indicated above for immunofluorescence analysis. Subsequently the PLA

protocol was followed according to the Duolink In Situ Red Starter Kit Mouse/Rabbit (Millipore

Sigma) instructions. Briefly, after blocking cells were incubated with 1:250 dilution of rabbit anti-

*Tb*KH pAb and 1:500 dilution of mouse anti-HA mAb (BioLegend, Cat# MMS-101R). As a

negative control, cells were incubated only with 1:250 dilution of the rabbit anti-*Tb*KH pAb. Then

PLA species-specific secondary antibodies with minus and plus oligonucleotide probes were

added, followed by ligation, amplification and hybridization with specific red-fluorescent

oligonucleotides to allow detection by fluorescence microscopy. Samples were mounted and

imaged as described for immunofluorescence analysis.

Acknowledgements

We appreciate the expert advice and support of the staff of the Advanced Light Microscopy

Core in the Jungers Center for Neurosciences at Oregon Health & Science University. We

acknowledge discussions with Dr. Samuel Dean (University of Oxford) concerning new flagellar

membrane proteins identified by the TrypTag project.

Competing interests

563 The authors declare no competing or financial interests.

Author contributions

566 Conceptualization: M.A.S. and S.M.L.; Data collection: M.A.S.; Data analysis: M.A.S. and

567 S.M.L.; Writing and editing: M.A.S. and S.M.L.; Funding acquisition and project administration:

568 S.M.L.

Funding

571 This work was supported by National Institutes of Health grant AI121160 to S.M.L. The content is

572 solely the responsibility of the authors and does not necessarily represent the official views of the

573 National Institutes of Health.

References

Affolter, M., Hemphill, A., Roditi, I., Muller, N. and Seebeck, T. (1994). The repetitive
microtubule-associated proteins MARP-1 and MARP-2 of *Trypanosoma brucei*. *J Struct Biol* **112**,
241-51.

Akiyoshi, B. and Gull, K. (2014). Discovery of unconventional kinetochores in
kinetoplastids. *Cell* **156**, 1247-1258.

Baudieri, B. and Tomassini, N. (1962). Fine struttura dei tripanosomi. *Parassitologia* **4**,
89-95.

Burkard, G., Fragoso, C. M. and Roditi, I. (2007). Highly efficient stable transformation
of bloodstream forms of *Trypanosoma brucei*. *Mol Biochem Parasitol* **153**, 220-3.

Cayla, M., Rojas, F., Silvester, E., Venter, F. and Matthews, K. R. (2019). African
trypanosomes. *Parasit Vectors* **12**, 190.

Dang, H. Q., Zhou, Q., Rowlett, V. W., Hu, H., Lee, K. J., Margolin, W. and Li, Z. (2017).
Proximity Interactions among basal body components in *Trypanosoma brucei* identify novel
regulators of basal body biogenesis and inheritance. *mBio* **8**.

Dean, S., Sunter, J., Wheeler, R. J., Hodkinson, I., Gluenz, E. and Gull, K. (2015). A
toolkit enabling efficient, scalable and reproducible gene tagging in trypanosomatids. *Open Biol*
5, 140197.

Dean, S., Sunter, J. D. and Wheeler, R. J. (2017). TrypTag.org: A trypanosome genome-
wide protein localisation resource. *Trends Parasitol* **33**, 80-82.

Ellis, D. S., Ormerod, W. E. and Lumsden, W. H. R. (1976). Filaments of "*Trypanosoma
brucei*": some notes on differences in origin and structure in two stains of "*Trypanosoma
(Trypanozoon) brucei rhodesiense*". *Acta Tropica* **33**, 151-168.

Farr, H. and Gull, K. (2012). Cytokinesis in trypanosomes. *Cytoskeleton (Hoboken)* **69**,
931-41.

**Fredriksson, S., Gullberg, M., Jarvius, J., Olsson, C., Pietras, K., Gustafsdottir, S.
M., Ostman, A. and Landegren, U.** (2002). Protein detection using proximity-dependent DNA
ligation assays. *Nat Biotechnol* **20**, 473-7.

Hemphill, A., Affolter, M. and Seebeck, T. (1992). A novel microtubule-binding motif
identified in a high molecular weight microtubule-associated protein from *Trypanosoma brucei*. *J
Cell Biol* **117**, 95-103.

Hoffman, E. A., Frey, B. L., Smith, L. M. and Auble, D. T. (2015). Formaldehyde
crosslinking: a tool for the study of chromatin complexes. *J Biol Chem* **290**, 26404-11.

Imhof, S., Zhang, J., Wang, H., Bui, K. H., Nguyen, H., Atanasov, I., Hui, W. H., Yang,
S. K., Zhou, Z. H. and Hill, K. L. (2019). Cryo electron tomography with Volta phase plate reveals
novel structural foundations of the 96-nm axonemal repeat in the pathogen *Trypanosoma brucei*.
Elife **8**.

Ishida, T. and Kinoshita, K. (2007). PrDOS: prediction of disordered protein regions from
amino acid sequence. *Nucleic Acids Res* **35**, W460-4.

Kaiser, P., Meierhofer, D., Wang, X. and Huang, L. (2008). Tandem affinity purification
combined with mass spectrometry to identify components of protein complexes. *Methods Mol Biol*
439, 309-26.

Kelly, F. D., Sanchez, M. A. and Landfear, S. M. (2020a). Touching the surface: Diverse
roles for the flagellar membrane in Kinetoplastid parasites. *Microbiol Mol Biol Rev* **84**, e00079-19.

Kelly, F. D., Tran, K. D., Hatfield, J., Schmidt, K., Sanchez, M. A. and Landfear, S. M.
(2020b). A cytoskeletal protein complex is essential for division of intracellular amastigotes of
Leishmania mexicana. *J Biol Chem* **295**, 13106-13122.

Kennedy, P. G. (2013). Clinical features, diagnosis, and treatment of human African
trypanosomiasis (sleeping sickness). *Lancet Neurol* **12**, 186-94.

Langousis, G. and Hill, K. L. (2014). Motility and more: the flagellum of *Trypanosoma*
brucei. *Nat Rev Microbiol* **12**, 505-18.

MacLean, L., Myburgh, E., Rodgers, J. and Price, H. P. (2013). Imaging African
trypanosomes. *Parasite Immunol* **35**, 283-94.

Manna, P. T., Boehm, C., Leung, K. F., Natesan, S. K. and Field, M. C. (2014). Life and
times: synthesis, trafficking, and evolution of VSG. *Trends Parasitol* **30**, 251-8.

McCulloch, R., Vassella, E., Burton, P., Boshart, M. and Barry, J. D. (2004).
Transformation of monomorphic and pleomorphic *Trypanosoma brucei*. *Methods Mol Biol* **262**,
53-86.

McGuffin, L. J., Bryson, K. and Jones, D. T. (2000). The PSIPRED protein structure
prediction server. *Bioinformatics* **16**, 404-5.

Mitchell, A., Chang, H. Y., Daugherty, L., Fraser, M., Hunter, S., Lopez, R., McAnulla,
C., McMenamin, C., Nuka, G., Pesseat, S. et al. (2015). The InterPro protein families database:
the classification resource after 15 years. *Nucleic Acids Res* **43**, D213-21.

Molloy, J. O. and Ormerod, W. E. (1964). A fibril emerging from the posterior end of
Trypanosoma rhodesiense. *Philos Trans R Soc Lond B Biol Sci* **58**, 2.

Oberholzer, M., Langousis, G., Nguyen, H. T., Saada, E. A., Shimogawa, M. M.,
Jonsson, Z. O., Nguyen, S. M., Wohlschlegel, J. A. and Hill, K. L. (2011). Independent analysis

of the flagellum surface and matrix proteomes provides insight into flagellum signaling in
mammalian-infectious *Trypanosoma brucei*. *Mol Cell Proteomics* **10**, M111 010538.

Poon, S. K., Peacock, L., Gibson, W., Gull, K. and Kelly, S. (2012). A modular and
optimized single marker system for generating *Trypanosoma brucei* cell lines expressing T7 RNA
polymerase and the tetracycline repressor. *Open Biol* **2**, 110037.

Robinson, D. R. and Gull, K. (1991). Basal body movements as a mechanism for
mitochondrial genome segregation in the trypanosome cell cycle. *Nature* **352**, 731-3.

Roux, K. J., Kim, D. I., Raida, M. and Burke, B. (2012). A promiscuous biotin ligase
fusion protein identifies proximal and interacting proteins in mammalian cells. *J Cell Biol* **196**, 801-
10.

Sanchez, M. A. (2013). Molecular identification and characterization of an essential
pyruvate transporter from *Trypanosoma brucei*. *J Biol Chem* **288**, 14428-37.

Sanchez, M. A., Tran, K. D., Valli, J., Hobbs, S., Johnson, E., Gluenz, E. and Landfear,
S. M. (2016). KHARON Is an essential cytoskeletal protein involved in the trafficking of fagellar
membrane proteins and cell division in African trypanosomes. *J Biol Chem* **291**, 19760-73.

Santi, A. M. M., Lanza, J. S., Tunes, L. G., Fiuza, J. A., Roy, G., Orfano, A. D. S., de
Carvalho, A. T., Frezard, F., Barros, A. L. B., Murta, S. M. F. et al. (2018). Growth arrested
live-attenuated *Leishmania infantum* KHARON1 null mutants display cytokinesis defect and
protective immunity in mice. *Sci Rep* **8**, 11627.

Schepilewsky, E. (1912). Fadenförmige anhängsel bei den trypanosomen. *Zbl. Bakt.* **65**,
79-83.

Schneider, A., Hemphill, A., Wyler, T. and Seebeck, T. (1988). Large microtubule-
associated protein of *T. brucei* has tandemly repeated, near-identical sequences. *Science* **241**,
459-62.

Shimogawa, M. M., Saada, E. A., Vashisht, A. A., Barshop, W. D., Wohlschlegel, J.
A. and Hill, K. L. (2015). Cell surface proteomics provides insight into stage-specific remodeling
of the host-parasite interface in *Trypanosoma brucei*. *Mol Cell Proteomics* **14**, 1977-88.

Soderberg, O., Gullberg, M., Jarvius, M., Ridderstrale, K., Leuchowius, K. J., Jarvius,
J., Wester, K., Hydbring, P., Bahram, F., Larsson, L. G. et al. (2006). Direct observation of
individual endogenous protein complexes *in situ* by proximity ligation. *Nat Methods* **3**, 995-1000.

Sonnhammer, E. L., Eddy, S. R., Birney, E., Bateman, A. and Durbin, R. (1998). Pfam:
multiple sequence alignments and HMM-profiles of protein domains. *Nucleic Acids Res* **26**, 320-
2.

Subota, I., Julkowska, D., Vincensini, L., Reeg, N., Buisson, J., Blisnick, T., Huet, D.,
Perrot, S., Santi-Rocca, J., Duchateau, M. et al. (2014). Proteomic analysis of intact flagella of
procyclic *Trypanosoma brucei* cells identifies novel flagellar proteins with unique sub-localization
and dynamics. *Mol Cell Proteomics* **13**, 1769-86.

Sun, S. Y., Wang, C., Yuan, Y. A. and He, C. Y. (2013). An intracellular membrane
junction consisting of flagellum adhesion glycoproteins links flagellum biogenesis to cell
morphogenesis in *Trypanosoma brucei. J Cell Sci* **126**, 520-31.

Szempruch, A. J., Dennison, L., Kieft, R., Harrington, J. M. and Hajduk, S. L. (2016a).
Sending a message: extracellular vesicles of pathogenic protozoan parasites. *Nat Rev Microbiol*
14, 669-675.

Szempruch, A. J., Sykes, S. E., Kieft, R., Dennison, L., Becker, A. C., Gartrell, A.,
Martin, W. J., Nakayasu, E. S., Almeida, I. C., Hajduk, S. L. et al. (2016b). Extracellular vesicles
from *Trypanosoma brucei* mediate virulence factor transfer and cause host anemia. *Cell* **164**,
246-257.

Tran, K. D., Rodriguez-Contreras, D., Vieira, D. P., Yates, P. A., David, L., Beatty, W.,
Elferich, J. and Landfear, S. M. (2013). KHARON1 mediates flagellar targeting of a glucose
transporter in *Leishmania mexicana* and is critical for viability of infectious intracellular
amastigotes. *J. Biol. Chem.* **288**, 22721-22733.

Tran, K. D., Vieira, D. P., Sanchez, M. A., Valli, J., Gluenz, E. and Landfear, S. M.
(2015). *Kharon1* null mutants of *Leishmania mexicana* are avirulent in mice and exhibit a
cytokinesis defect within macrophages. *PLoS One* **10**, e0134432.

Uversky, V. N. (2013). The alphabet of intrinsic disorder: II. Various roles of glutamic acid
in ordered and intrinsically disordered proteins. *Intrinsically Disord Proteins* **1**, e24684.

Vaughan, S. and Gull, K. (2008). The structural mechanics of cell division in
Trypanosoma brucei. Biochem Soc Trans **36**, 421-4.

Vedrenne, C., Giroud, C., Robinson, D. R., Besteiro, S., Bosc, C., Bringaud, F. and
Baltz, T. (2002). Two related subpellicular cytoskeleton-associated proteins in *Trypanosoma*
brucei stabilize microtubules. *Mol Biol Cell* **13**, 1058-70.

Vickerman, K. and Luckins, A. G. (1969). Localization of variable antigens in the surface
coat of *Trypanosoma brucei* using ferritin conjugated antibody. *Nature* **224**, 1125-1126.

Wang, Z., Morris, J. C., Drew, M. E. and Englund, P. T. (2000). Inhibition of
Trypanosoma brucei gene expression by RNA interference using an integratable vector with
opposing T7 promoters. *J Biol Chem* **275**, 40174-9.

Wheeler, R. J., Gull, K. and Sunter, J. D. (2019). Coordination of the Cell Cycle in
Trypanosomes. *Annu Rev Microbiol* **73**, 133-154.

Wirtz, E., Leal, S., Ochatt, C. and Cross, G. A. M. (1999). A tightly regulated inducible
expression system for conditional gene knock-outs and dominant-negative genetics in
Trypanosoma brucei. Mol. Biochem. Parasitol. **99**, 89-101.

Wright, K. A. and Lumsden, W. H. R. (1970). The formation of filopodium-like processes
by *Trypanosoma. J. Cell Sci.* **6**, 285-297.

Zhang, T., Faraggi, E., Li, Z. and Zhou, Y. (2013). Intrinsically semi-disordered state and
its role in induced folding and protein aggregation. *Cell Biochem Biophys* **67**, 1193-205.

Zhou, Q., Lee, K. J., Kurasawa, Y., Hu, H., An, T. and Li, Z. (2018). Faithful chromosome
segregation in *Trypanosoma bruce*i requires a cohort of divergent spindle-associated proteins
with distinct functions. *Nucleic Acids Res* **46**, 8216-8231.

FIGURE LEGENDS

Fig. 1. *Tb*KH localizes in three distinct subcellular compartments. (A) Flagellar preparation

from *Tb*FLA1^{RNAi} cell line, isolated after 72 h of induction with 1 µg/ml doxycycline (dox), stained

with DAPI (*blue*) and immunostained with anti-*Tb*KH pAb (*Tb*KH, *red*). *Tb*KH localization at the

base of the flagellum is indicated by the *red arrowhead*. (B) Wild type BF parasites were

immunostained with anti-*Tb*KH pAb (*Tb*KH, *red*) and stained with DAPI. These images display a

parasite in which *Tb*KH staining is associated with both the subpellicular microtubules (*right*

arrowheads) and the mitotic spindle that connects the two nuclei late during mitosis (*left*

arrowheads). DIC indicates images collected by differential interference contrast microscopy.

(C) Western blot of total protein lysates from BF trypanosomes that are either wild type (*WT*) or

expressing BirA*::TbKH fusion protein (*A**). Blot was probed with anti-*Tb*KH pAb and developed

by chemiluminescence. Specific immunodetected proteins corresponding to the native *Tb*KH

and the BirA*::*Tb*KH fusion protein are shown. Relative protein molecular weights are shown in

kDa, as determined by mobility relative to molecular weight markers.

Fig. 2. Subcellular localization of *Tb*KHAPs in BF and PF *T. brucei*. (A, B) The HA$_3$::TbKHAP1

cell line was immunostained with anti-HA mAb (HA$_3$::*Tb*KHAP1*, green*) and anti-*Tb*KH pAb

(*Tb*KH*, red*). (C, D) The TbKHAP2::HA$_3$ cell line was stained with anti-HA mAb (*Tb*KHAP2::HA$_3$*,*

green) and anti-*Tb*KH pAb (*Tb*KH*, red*). (E) The BF 3V5::*Tb*KHAP3 and (F) PF HA$_3$::*Tb*KHAP3

cell lines were immunostained with (E) anti-V5 mAb (3V5::*Tb*KHAP3, *green)* or (F) anti-HA mAb

(HA$_3$::*Tb*KHAP3*, green*) and anti-*Tb*KH pAb (*Tb*KH*, red*). All preparations were stained with DAPI,

which detects both nuclear and kinetoplast DNA (*blue*). DIC images were also acquired from all

samples.

Fig. 3. KHARON and KHAPs colocalize at the base of the flagellum in *T. brucei*. (A) Whole

isolated flagella from the BF HA$_3$::*Tb*KHAP1/*Tb*FLA1^{RNAi} clone induced for RNAi for 72 h, were

immunostained with anti-HA mAb (HA$_3$::*Tb*KHAP1, *green*) and anti-*Tb*KH pAb (*Tb*KH, *red*) and

stained with DAPI (*blue*). (A, left panel) shows an immunofluorescence/DIC image of an isolated

flagellum, *black punctuated* box indicates the magnified area depicted in (A, center-right, center-

left and left) panels. *Green arrowheads* indicate HA$_3$::*Tb*KHAP1 localization, *red arrowheads*

indicate *Tb*KH localization and *yellow arrowheads* indicate overlapping signals, when channels

are merged, near the kinetoplast (*blue*). (B) Flagellar cytoskeletons were isolated from BF

trypanosomes expressing HA$_3$::TbKHAP2 and imaged as described in (A). (C) Flagellar

cytoskeletons were isolated from PF trypanosomes expressing HA$_3$::TbKHAP3 and imaged as

described in (A).

Fig. 4. *Tb*KH is closely associated with *Tb*KHAP1, *Tb*KHAP2, and *Tb*KHAP3. (A) Western blot of protein samples from pull down of HA$_3$::*Tb*KHAP1 using His$_{10}$::*Tb*KH as a bait and CAP15::HA$_3$ as negative control. *LS* and *EF*, lysate and elution fractions without formaldehyde crosslinking; *LS** and *EF**, lysate and elution fraction with formaldehyde crosslinking. Protein blots were developed by chemiluminescence, bands are indicated by protein names, and migration of molecular weight markers are designated in kDa. (B, C) PLA employing BF or PF HA$_3$::TbKHAP1/His$_{10}$::TbKH cell lines, as indicated. For these and all subsequent sections of this figure, the *left panel* shows results using both anti-HA mAb and anti-*Tb*KH pAb, and the *right panel* shows results using only the anti-*Tb*KH pAb, as the negative control. *Red puncta* indicate positive HA$_3$::*Tb*KHAP1-*Tb*KH interaction. DIC images were acquired from all samples. (D, E) PLA employing BF or PF *Tb*KHAP2::HA$_3$ cell lines, as indicated. *Red puncta* indicate positive *Tb*KHAP2-*Tb*KH interaction. (F, G*)* PLA employing BF or PF HA$_3$::*Tb*KHAP3 cell lines, as indicated using. *Red puncta* indicate positive *Tb*KHAP3-*Tb*KH interaction.

Figure 5. Depletion of *Tb*KHAP1 is lethal for *T. brucei* parasites. (A) Western blot of total protein lysates from TbKHAP1^{RNAi} BF parasites grown in the presence of doxycycline (+dox) and immunodetected with anti-HA mAb, anti-*Tb*KH pAb. The approximate molecular weights in kDa of HA$_3$::*Tb*KHAP1 and *Tb*KH are indicated as determined by mobility compared to weight markers. *Numbers* under the blot represent the relative intensity (*RI*) of HA$_3$::*Tb*KHAP1 protein normalized to the *Tb*KH protein level. (B) Cell density for induced (*empty circles*) and non-induced (*filled circles*) *Tb*KHAP1^{RNAi} BF cell lines. Parasite density was quantified by phase contrast microscopy using a hemocytometer. Data represent the averages and standard deviations of two experiments, each employing an independently isolated *Tb*KHAP1^{RNAi} clonal cell line, and technical replicates were also performed for each biological replicate. Standard deviations are too small to be visible. (C) Microscopic analysis of nuclei and kinetoplast numbers of BF *Tb*KHAP1^{RNAi} parasites following induction of RNAi. Data represent frequency (*%*) of cells with different numbers of DAPI stained nuclei (*N*) and kinetoplasts (*K*). Results represent the average and range of two independent experiments. Brackets compare 0 h to 48 h doxycycline induction, with *** representing $P<0.001$, as determined by 2-way ANOVA using Dunnett's multiple comparison test, where all time points were compared to 0 h doxycycline but only statistical significance for 48 h is shown. (D-F) *Tb*KHAP1^{RNAi} BF parasites were stained with DAPI (*blue*) and immunostained with anti-α-tubulin mAb (Tub, *red*) at (D) 0 h, (E) 20 h and (F) 48 h post-RNAi induction. In (E) the *white arrowhead* indicates a cell with a tadpole-like morphology and the *yellow arrowhead* indicates a cell with two flagella at opposite poles of the cell body.

Fig. 6. Phenotypes resulting from RNAi directed against *TbKHAP2* and *TbKHAP3*. (*A*)

Western blot of total protein lysates of BF parasites expressing HA$_{3::}$*Tb*KHAP2 following

induction of RNAi (1 μg ml^{-1} doxycycline, *dox*) directed against *TbKHAP2/3* RNAs. Protein blots

were probed with anti-HA mAb and anti-*Tb*KH pAb as loading control. Molecular weight markers

are indicated in kDa. (*B*) Growth curve of induced (*empty circles*) and non-induced (*filled circles*)

TbKHAP2/3^{RNAi} cell line. Parasite density was quantified by phase contrast microscopy using a

hemocytometer. The data represent two biological replicate experiments, but the standard

deviations are too small to see in the figure. Representative *Tb*KHAP2/3^{RNAi} cells induced for

RNAi were stained with DAPI (*blue*) and immunostained with anti-*Tb*KH pAb (*Tb*KH, *red*) at (*C*)

0 h and (*D)* 4 days post-RNAi induction. *White arrowhead* in (*D*) indicates a cell showing

aberrant morphology. DIC images were obtained for all samples. (*E*) Microscopic analysis of NK

in BF parasites induced for RNAi. Frequency (*%*) of cells with different numbers of nuclei (*N*)

and kinetoplasts (*K*) at different times following induction of RNAi against *TbKHAP2/3* mRNA.

Results represent the average and range of two independent experiments. Statistical

comparisons were performed as in Fig. 5C, with *** representing P<0.001, ** P<0.01, and *

P<0.05.

Fig. 7. Knockdown of *Tb*KHAP1, but not *Tb*KHAP2 or *Tb*KHAP3, impairs trafficking of

_Tb_CaCh::HA$_3$ to the flagellum. (A-C)*Tb*CaCh::HA$_3$/*Tb*KHAP1^{RNAi} BF cell line was induced with

1 μg/ml doxycycline (dox). Parasites were stained with DAPI (*blue*) and immunostained with

anti-HA mAb (*Tb*CaCh::HA$_3$*, green*) and anti-*Tb*KH pAb (*Tb*KH*, red*). (A) Non-induced

TbCaCh::HA$_3$/TbKHAP1^{RNAi} parasites (*0 h dox*), (B) parasites induced for 24 h (2*4 h dox*) and

(C*)* parasites induced for 48 h (*48 h dox*). *Green arrowheads* in (A-C) indicate flagella where

*Tb*CaCh::HA$_3$ is present, and *white arrowheads* in *(*B-C*)* indicate flagella where *Tb*CaCh::HA$_3$ is

absent. (D-F) *Tb*CaCh::mNG/*Tb*KHAP2/3^{RNAi} BF cell line was induced with 1 μg/ml doxycycline

(dox). Parasites were stained with DAPI (*blue*) and anti-*Tb*KH pAb (*Tb*KH*, red*), and mNG

endogenous fluorescence was also acquired (*Tb*CaCh::mNG*, green*). Parasites were induced

with doxycycline for (D) 0 days (*0 d dox*), (E) 4 days (*4d dox*), or (F) 7 days (*7d dox*). *Green*

arrowheads indicate flagella where *Tb*CaCh::mNG is present. Merged DIC images are

presented in the right panel of each image pair. The scale bar shown in (A) applies to all

images.

Fig. 8. Trafficking of *Tb*EVMP1, *Tb*EVMP2, and *Tb*FLA1BP in *Tb*KH^{RNAi} BF parasites. (A)

Parasites expressing *Tb*EVMP1:mNG before (*0 h RNAi*) and after (*24 h RNAi*) induction of RNAi

against *TbKH* mRNA. *Green arrowheads* indicate mNG fluorescence in the FM (all panels) or in

extracellular vesicles (*0 h RNAi*, right panel). *White arrowheads* indicate flagella without mNG

fluorescence. (B) Parasites expressing *Tb*EVMP2::mNG before (*0 h RNAi*) and after (*24 h*

RNAi) induction of RNAi against *TbKH* mRNA. (C) Parasites expressing *Tb*FLA1BP::mNG

before (*0 h RNAi*) and after (*24 h RNAi, 48 h RNAi*) induction of RNAi against *TbKH* mRNA. The

left-most images in A, B, and C represent formaldehyde fixed parasites, whereas for the other

images, live parasites were suspended in CyGel, which facilitates visualization of secreted

extracellular vesicles.

Figure 2

Figure 3

Figure 6

A 0 h dox DIC 0 h dox

7 μm

B *Tb*CaCh::HA₃ *Tb*KH DAPI 24 h dox DIC 24 h dox

C 48 h dox DIC 48 h dox

D 0 h dox DIC 0 h dox

E 4 d dox DIC 4 d dox

F 7 d dox 7 d dox

A *Tb*EVMP1 DAPI   *Tb*EVMP1

0 h RNAi   24 h RNAi

B *Tb*EVMP2   *Tb*EVMP2   *Tb*EVMP2
DAPI

0 h RNAi   24 h RNAi

C *Tb*FLA1BP   *Tb*FLA1BP
DAPI

0 h RNAi   24 h RNAi   48 h RNAi