Breeding Program Optimization for Genomic Selection in Winter Wheat

Megan Calvert[a], Byron Evers[b], Xu Wang[b], and Jesse Poland[b]

AFFILIATION

5   a Interdepartmental Genetics, Kansas State University, Manhattan Kansas 66502

b Department of Plant Pathology, Kansas State University, Manhattan, KS, 66506

KEYWORDS

Triticum aestivum, Breeding, Genomic Prediction

Abbreviations

10   AYN, advanced yield nurseries; BEL, Belleville; BLUPs, Best Linear Unbiased Predictions; GEBVs, genomic estimated breeding values; GNDVI, Green Normalized Difference Vegetation Index; GP, genomic prediction; GRVI, green-red vegetation indices; GRYLD, grain yield; GYP, Gypsum; GxE, genotype x environment; HTP, high-throughput phenotyping; HUTCH, Hutchinson; KSU, Kansas State University; MANH,

15   Manhattan; MP, McPherson; NDRE, normalized difference red edge; NDVI, normalized difference vegetation indices; PTHT, plant height; PC, principal component; PYN, preliminary yield nurseries; RKHS, reproducing kernel Hilbert spaces regression; RR, ridge regression; SNPs, Single nucleotide polymorphisms; UAS, uncrewed aerial system; VI, vegetation indices

20   **Abstract**

Developing methodologies in the fields of phenomics and genomic prediction

have the potential to increase the production of crop species by influencing germplasm

improvement. The integration of these technologies into germplasm improvement and

breeding programs requires evidence that there will be a direct economic benefit to the

25   program. We determined a basic set of parameters, such as prediction accuracy greater

than 0.3, the ability to genotype over 7 lines for the cost of one phenotypic evaluation,

and heritability levels below 0.4, at which the use of genomic selection would be of

economic benefit in terms of genetic gain and operational costs to the Kansas State

University (KSU) winter wheat breeding program. The breeding program was then

30   examined to determine whether the parameters benefitting genomic selection were

observed or achievable in a practical sense. Our results show that the KSU winter wheat

breeding program is at a decision point with regards to their primary means of selection.

A few operational changes to increase prediction accuracy would place the program in

the parameter space where genomic selection would be of greatest benefit to the

35   program.

Introduction

Technologies are constantly evolving and growing in today's ever-changing

world. The introduction of new technologies into an essential service and the

development of new crop varieties can be challenging and costly (Moose and Mumm

40   2008). New technologies must be carefully evaluated to determine if they provide a

significant enough advantage to adjust years of traditional practice. There are many

technologies that appeared to have great promise, such as quantitative trait loci (QTL),

and marker-assisted selection (MAS), and yet were not beneficial in a practical sense to

breeding programs at the time (Bernardo 2008). Yet with a growing global population,

45   erratic environmental conditions, and a finite-amount of arable land, the introduction of

new technologies into our crop development systems is essential. The determination of

which technologies to implement and the most efficient way to do so is a constant

challenge facing plant breeders.

Wheat (*Triticum aestivum*) is one of the top 3 field crops planted in the USA,

50   behind soybeans and corn, with 1.9 billion bushels produced in 2019 (Bond 2020).

Wheat acres and production in the USA have been in a decline since the 1980s as a

result of international competition and changing economic practices. This is despite the

increasing demand for wheat due to growing global populations, which is only expected

to increase over the coming century. It is estimated that a 2% increase in yearly grain

55   yields are required to meet these demands (Bassi et al. 2016). Currently there are no

reports of wheat breeding programs achieving this level of gain, making wheat an

important candidate for breeding technologies designed to accelerate variety

development.

Genomic prediction (GP) is one of the new technologies that is showing a great

60   deal of promise to assist in crop-variety production. GP involves the use of genome-wide

markers to predict the breeding value of individuals in a population (Meuwissen, Hayes,

and Goddard 2001). This is done by genotyping and phenotyping a training population

which is used to establish a model for the trait of interest. This model is then used to

calculate genomic estimated breeding values (GEBVs) of a population for which only

65   genotype information is available. GP is already a common technique used in animal

breeding due to the benefit of being able to predict a phenotype without having to

observe the phenotype, for example the milk yield of a bull's offspring (Hayes et al.

2009; Georges, Charlier, and Hayes 2019).

Plant breeding programs have yet to fully utilize GP for a number of reasons.

70   Breeding programs have to test the same experimental line in multiple locations due to

the need to select varieties that are stable across environments. This negates some of

the benefit that GP supplies to animal breeding, in which the same genotype cannot be

tested under multiple conditions. The genotype x environment (GxE) variation is often

high in plant breeding populations due in part to the large weather differences

75   experienced between locations. This significantly decreases the accuracies of most GP

models (Dawson et al. 2013; Heslot et al. 2012). Previous GP models did not account for

GxE interactions and were not of as much use to plant breeding. Newer genomic

prediction models can now take into account these GxE interactions and are showing

greater accuracy in plant breeding situations (Burgueño et al. 2012; Pérez-Rodríguez et

80   al. 2017; Montesinos-López et al. 2016).

The accuracy of GP models has shown some improvement when a covariate is

included (Crain et al. 2018; Rutkoski et al. 2016). This has had some success when

including high-throughput phenotyping (HTP) data that is taken throughout the course

of the growing season. HTP techniques have the ability to measure thousands of

85   phenotypes accurately in a short period of time. Many of the HTP techniques used in

breeding programs take advantage of new developments in remote sensing, uncrewed

aerial system (UAS), and sensor technology to measure reflectance and temperature

phenotypes. These phenotypic traits have been shown to be correlated with grain yield

in wheat (Elliott and Regan 1993; Blackmer et al. 1996; Curran et al. 1983).

90   In addition to the use of correlated traits, the efficiency of GP is also affected by

he stage of the breeding program when it is implemented. Primarily, GP is used to

predict the breeding value which is comprised of the additive genetic variation. For the

model to perform accurately the additive genetic variation for the phenotype of interest

needs to be segregating in the population (Bassi et al. 2016). This is more likely to occur

95   in earlier stages of a breeding program as it is easily selected for, while later stages of a

breeding program are more likely to select for other epistatic genetic variations.

Another factor in addition to stage of implementation that has hindered the

adoption of GP in plant breeding programs are the costs associated with it. The

development and establishment of genotyping practices in a plant breeding program

100 requires an investment in infrastructure and training (Moose and Mumm 2008). Along

with genetic marker costs, the establishment and maintenance of an adequate training

population adds additional costs to the GP protocols. These costs may not be offset by

the gains that the program could potentially achieve using GP (Bassi et al. 2016; Jarquín

et al. 2017).

105   In this study we examined the implementation of GP in a wheat breeding

program to specifically examine 1) what parameters/infrastructure is required to

implement GP as a primary selection strategy, and 2) are those parameters being met in the KSU public breeding program?

Method and Materials

110   ***Cost-Benefit Simulation***

A simulated breeding program was used to determine the expected genetic gain when basing selection of material exclusively on observed breeding values, or exclusively on prediction of breeding values. The simulation assumes that a breeding program does not use a combination of phenotypic selection and genomic selection at

115   the same time. In consultation with the KSU hard winter wheat breeding program an appropriate range of cost estimates were determined, as well as sizes of programs (Supplementary Table 1). The simulated values assume that the cycle length for each selection option is the same and covers the period of the program before the Advanced Yield Nursery (AYN) stage. It also assumes that the number of lines that will be selected

120   for the AYN stage are the same regardless of which selection method is used.

The cost of line developments was estimated to be between \$4-\$30 per line which covers the initial cross and formation of an inbred line. This can either be done by several years of inbreeding or by the formation of double haploid lines. This was applied as a fixed cost needed to develop the initial population for selection regardless of which

125   selection method was used.

Phenotypic Selection

The costs of phenotypic observation plots are estimated to be between $12-$40 per plot and evaluated within this range at $5 increments. To obtain an accurate phenotypic measurement, replications of the experimental line need to be planted and phenotyped. Depending on the structure of the program this may mean several replications at a single site, or fewer replications at several sites. The maximum number of experimental breeding lines was calculated as:

$$ExperimentalLines = Budget \Big/ (LineDevelopment + (PhenoCost * rep)) \quad (1)$$

where $ExperimentalLines$ is the number of lines that would be advanced to the phenotyping stage of selection, the $Budget$ is the total monetary budget for that stage of the breeding program, $LineDevelopment$ is the cost of advancing a single cross to an inbred experimental line for evaluation, $PhenoCost$ is the total cost to obtain the phenotype of interest including labor and other miscellaneous operation costs, and rep is the number of replications of each experimental line that are planted in field and will require phenotyping.

The number of experimental breeding lines was used as the population size when estimating other population parameters.

Genomic Selection

The cost of genotyping a single line for prediction was estimated to be between $1-$10 and evaluated within this range at $1 increments. The maximum number of experimental lines that can be genotyped for prediction was calculated as:

$$GenotypedLines = Budget \Big/ LineDevelopment + GenoCost \quad (2)$$

Where $GenotypedLines$ is the number of lines that would be evaluated by genotyping, the $Budget$ is the total monetary budget for that stage of the breeding program, $LineDevelopment$ is the cost of advancing a single cross to an inbred experimental line for evaluation, and $GenoCost$ which is the cost of genotyping a single line including labor and other operational costs. It is assumed that the genotyping is only performed once, and that replication is not required.

The number of possible genotypes was then used as the population size when estimating other population parameters.

Simulation Details

The selection methods were compared for every possible combination of number of experimental lines phenotyped and number of experimental lines genotyped, based on the ratio between the correlated response to selection, and the response to selection (Falconer and Mackay 2009).

The expected response to selection for the primary trait, grain yield, was calculated by:

$$R = ih\sigma_A \tag{3}$$

Where i is the intensity of selection, h is the square-root if the narrow-sense heritability of the primary trait, and σ_A is the standard deviation of the additive genetic variance (Falconer and Mackay 2009).

For this study, the GEBVs are assumed to be the secondary trait that is correlated to the response trait of grain yield, which would be through experimental observation plots. The correlated response to selection is calculated by:

$$170 \qquad CR = i * \sqrt{h_x^2} * \sqrt{h_y^2} * r_g * \sigma_{py} \qquad (4)$$

Where i is the intensity of selection, $\sqrt{h_x^2}$ is the square-root of the narrow-sense

heritability of the response trait, $\sqrt{h_y^2}$ is the square-root of the narrow-sense heritability

of the secondary trait, r_g is the additive genetic correlation between the traits, and σ_{py}

is the standard deviation of the phenotypic variation for the secondary trait (Falconer

175 and Mackay 2009).

A comparison between the indirect response to selection and the expected

response to selection is best demonstrated as (Falconer and Mackay 2009):

$$\frac{CR_x}{R_x} = \frac{i_Y r_g \sqrt{h_y^2}}{i_x \sqrt{h_x^2}} \qquad (5)$$

Where i_Y is the selection intensity on the secondary trait, r_g is the additive

180 genetic correlation between the traits, $\sqrt{h_y^2}$ is the square-root of the narrow-sense

heritability of the secondary trait, i_x is the selection intensity on the direct trait, and

$\sqrt{h_x^2}$ is the square-root of the narrow-sense heritability of the response trait.

It was assumed that the individual phenotypes were made up of a genetic portion

and an environmental portion. The distribution of each of these portions was assumed to

185 be a random normal with a mean of 0, and a standard deviation of $\sqrt{h^2}$ or $\sqrt{(1 - h^2}$ for

the genotypic and environmental proportions respectively. A sample the size of the number

of lines that were possibly genotyped was taken from this distribution for each individual to

give the overall phenotypic distribution. This overall phenotypic distribution was used to

determine the intensity of selection (i) in terms of the standard deviation and the selection

190differential with the msm package (Jackson 2011).

The narrow sense heritabilities for the response trait and the additive genetic

correlation were set with testing the ranges of 0 and 1 at increments of 0.1 each. The

narrow-sense heritability of the secondary trait, the genotyping, was assumed to be 0.9.

This is under the assumption that the genotypes are inherited almost exactly as they are

195sequenced and that there are only a few genotyping errors. The ratio between the

correlated response and the expected response to selection was plotted against the ratio

between number of experimental lines phenotyped and number of lines genotyped.

Visualizations were created using ggplot2 (Wickham 2016).

Plant Material

200   The KSU Breeding Program breeds hard red bread wheat for a large area which

contains different mega-environmental conditions. A subset of 5 locations within the

same Kansas mega-environment, Belleville (BEL), Gypsum (GYP), McPherson (MP),

Hutchinson (HUTCH) and Manhattan (MANH), were selected for analysis in Kansas

between 2016 and 2019. This resulted in 1989 experimental lines being examined over

205   the course of 4 years. These sites contain trials from the Preliminary Yield Trials (PYN,

primary $F_{5:7}$) and the advance yield nursery (AYN, primarily $F_{5:8}$). The planting, harvest

dates and trial size are provided in Table 1. These locations, excluding Manhattan and

Hutchinson, are located in farmers' fields under typical grower management practices.

The PYN and AYN trials were all planted in six-row plots of 1.5m by 4.5m. The PYN are

210   planted in a modified augmented design with one replicate of the experimental line per

location (Federer and Raghavarao 1975). Plant checks are planted across whole rows and columns in the trial, and sub-block checks are assigned randomly within each block. The AYN is made up of lines selected from the PYN trials. The lines are planted using two replicated α-lattice designs (Patterson and Williams 1976). All 5 locations are planted

215   each year but if a site experiences extreme environmental variation from the normal climate it may not be harvested, providing an unbalanced set of data.

Phenotyping

Phenotypic information was collected either by combine for grain yield (GRYLD), by UAS, or by hand using the Field Book application (Rife and Poland 2014) for plant

220   height (PTHT). A DJI Matrice100 (DJI, USA) quadcopter UAS was equipped with a 5-band multi-spectral RedEdge camera (MicaSense Inc. USA) to collect plot-level reflectance values for each year, based on the standard protocols developed by The Wheat Genetics Lab at Kansas State University as laid out in X. Wang et al. (2018). The UAS data was collected throughout the course of the growing season, once every 7-10 days depending

225   on weather conditions. Plant height was collected manually during the grain ripening stage before harvest. Vegetation indices (VIs) were calculated from the reflectance values based on the protocols laid out in X. Wang et al. (2018) and given by Table 2.

Genotyping

The 1991 lines in the study years were sequenced using the genotype-by-

230   sequencing on an Illumina Hi Seq2000 or Hi Seq2500 (Elshire et al. 2011). Single nucleotide polymorphisms (SNPs) were called with the Tassel software with the Chinese

Spring wheat assembly v1.0 as a reference (Bradbury et al. 2007; Glaubitz et al. 2014; International Wheat Genome Sequencing Consortium et al. 2018). The final data set included 8182 SNPs that were selected for use passed one of three filtering criteria

235   optimized for the wheat genome by Shrestha et al. (2020) that include Chi-square, Fisher's test for independence, and the inbreeding coefficient, as well as having a minor allele frequency greater than 0.05 and missing less than 20% of the data. Missing SNPs were imputed with Beagle 5.1 (Browning, Zhou, and Browning 2018).

Data Analysis

240   The Best Linear Unbiased Predictions (BLUPs) for each line were calculated for each individual year and for multiple years, including or excluding locations as needed. Where multiple years and locations are included, such as for GRYLD, the BLUPs were calculated by:

$$y_{ijklt} = \mu + G_i + S_j + M_{l(j)} + GM_{il(j)} + R_{k(lj)} + P_{t(lj)} + \varepsilon_{ijklt} \quad (6)$$

245   Where y_{ijklt} is the phenotypic response variable, μ is the fixed overall mean, G_i accounts for the random genotype effect N(0,σ^2), S_j is the random effect for year j N(0,σ^2), $M_{l(j)}$ is the random effect nested for location l within year j N(0,σ^2), $GM_{il(j)}$ is the random genotype by location effect nested within year j N(0,σ^2), $R_{k(lj)}$ is the random effect of replication k within location and year N(0,σ^2), $P_{t(lj)}$ is the random

250   treatment effect t nested within year-location N(0,σ^2), and ε_{ijklt} as the residual effect.

When only a single year with multiple locations is included the BLUPs are calculated by:

$$y_{ijkl} = \mu + G_i + L_j + GL_{ij} + R_{k(j)} + T_{l(j)} + \varepsilon_{ijkl} \qquad (7)$$

Where y_{ijkl} is the phenotypic response variable, μ is the fixed overall mean, G_i is

the random genotype effect $N(0,\sigma^2)$, L_j is the random effect of location j $N(0,\sigma^2)$, GL_{ij}

is the random effect of genotype by location $N(0,\sigma^2)$, $R_{k(j)}$ is the random rep effect

nested in location $N(0,\sigma^2)$, $T_{l(j)}$ is the random treatment effect nested within location

$N(0,\sigma^2)$, and ε_{ijkl} is the residual effect.

The broad-sense heritability for all years and multiple year-locations was

calculated as:

$$H^2 = \frac{\sigma_g}{\sigma_g + \frac{\sigma_{r(ly)}}{y*l} + \frac{\sigma_e}{y*l*r}} \qquad (8)$$

Where σ_g is the total genetic variance, $\sigma_{r(ly)}$ is the variance contributed by the

rep-location effect nested within year, σ_e is the residual environmental variance. As the

data is unbalanced, y is the harmonic mean of the number of years planted, l is the

harmonic mean of the number of locations planted within year, and r is the harmonic

mean of the number of replications per location per year (Holland, Nyquist, and

Cervantes-Martínez 2010).

The broad-sense heritability for individual year AYN and individual year-multiple

location PYN was calculated by:

$$H^2 = \frac{\sigma_g^2}{\sigma_g^2 + \frac{\sigma_{r(l)}^2}{l} + \frac{\sigma_e^2}{l*r}} \qquad (9)$$

Where σ_g^2 is the total genetic variance, $\sigma_{r(l)}^2$ is the variance contributed by the

replication nested in location, and σ_e^2 is the residual environmental variance. Similar to

equation 8, l is the harmonic mean of the number of locations planted and r is the

harmonic mean of the number of replications per location (Holland, Nyquist, and

275 Cervantes-Martínez 2010).

The VI phenotypes were mainly considered on a year-location-trial basis. As such
the BLUPs for the VI phenotypes for the AYN trials are calculated by:

$$y_{irb} = \mu + G_i + M_r + B_{r(b)} + \varepsilon_{irb} \qquad (10)$$

Where y_{irb} is the phenotypic response variable, μ is the fixed overall mean, G_i is

280 the random genotypic effect N(0,σ^2), M_r is the random effect of the replicate N(0,σ^2),

$B_{r(b)}$ is the random effect of the experimental block nested within replicate N(0,σ^2),

and ε_{irb} is the residual effect. The broad-sense heritability was calculated for each year-

location's AYN that had VI phenotypes by:

$$H^2 = \frac{\sigma_g^2}{\sigma_g^2 + \frac{\sigma_{error}^2}{r}} \qquad (11)$$

285 Where σ_g^2 is the genetic variance, σ_{error}^2 is the residual environmental variance

and r is the number of replicates.

The BLUPs for the VI phenotypes for the PYN trials are calculated by:

$$y_{ib} = \mu + G_i + B_b + \varepsilon_{ib} \qquad (12)$$

Where y_{ib} is the phenotypic response variable, μ is the fixed overall mean, G_i is

290 the random genotypic effect N(0,σ^2), B_b is the random effect of the experimental block

N(0,σ^2), and ε_{ib} is the residual effect.

Genome-wide Association Analysis

A principal component (PC) analysis of the genotypic information was conducted
with the pcaMethods package (Stacklies et al. 2007). A genome-wide association

295   analysis was performed for the VI BLUPs for each year-location-trial using the rrBLUP

package (Endelman 2011). The kinship matrix was calculated using rrBLUP was included

with 4 PCs to account for kinship and population structure respectively (Endelman and

Jannink 2012). A Bonferroni correction was applied with α = 0.05 to determine

significance.

300   ***Genomic Prediction***

Genomic prediction was performed with the rrBLUP package and the BGLR

statistical package in the R software environment (Pérez and de los Campos 2014). The

rrBLUP package performs ridge regression (RR), and the BGLR package was used to

perform a BayesC based prediction and a reproducing kernel Hilbert spaces regression

305   (RKHS). A cross-validation strategy of 100 replications with 80% of the population in the

training data set and 20% of the population in the predicted data set was used. The

accuracy of the prediction was determined by the correlation between the predicted

and observed values. Across year and location predictions were also performed where

years, year-trial, and year-location combinations that were not included in the training

310   population are predicted from the remaining data. An example of this would be to use

all years excluding the 2016 season as the training data set and the 2016 season as the

prediction data set, or all years excluding 2016 season except for the 2016-HUTCH data

as the training data set and the remaining three 2016 locations as the prediction data

set.

315   For the VI data only rrBLUP was used for genomic prediction. The VI were used

as a cofactor when predicting grain yield for that specific year-location-trial

combination. The same cross-validation procedure as before was used.

Unless otherwise stated, all analysis took place in the R software environment

using the Tidyverse suite of packages (Core Team 2020; Wickham et al. 2019).

320   **Results**

Simulation results

To determine the optimal parameters for the operation of the KSU winter wheat

breeding program to favor genomic prediction a simulation was created based on

economic decisions. The correlated response based on the ratio of lines experimental

325   lines to be phenotyped to experimental lines to be genotypes is given in Figure 1. Once

ratio between the correlated response and the expected response is greater than 1,

then selection on the correlated trait is expected give greater gain for the primary trait.

For the KSU wheat breeding program a primary focus on genotyping is beneficial in

terms of genetic gain when the narrow-sense heritability of grain yield is below 0.4,

330   approximately 7.5 lines can be genotyped for every one line that can be phenotyped,

and the prediction accuracy of the genomic prediction models is greater than 0.3.

Heritability of traits

Heritability was calculated to identify the phenotypic variance that could be

partitioned to the genetic component across the experimental trials. A summary of the

335   grain yield phenotypes is given in Table 3. The broad sense heritabilities for each of the

seasons ranged from 0.022 for the AYN to 0.441 for the PYN in the 2015/2016 season (Figure 2). The broad-sense heritability for the 2018/2019 PYN cannot be calculated as only one rep was planted at one location (HUTCH).

The VIs show a moderate heritability across the seasons, with the majority being above 0.5 (Figure 3) The heritabilities are variable across the course of the season and locations but the same date-location combination shows similarity across the various VIs, indicating a large environmental influence.

The correlations between grain yield for a location-season-trial combination and the VI's for that location-season-trial combination over the course of the season are given in Figure 4. The correlations ranged between -0.6 in the 2017/2018 season PYN at HUTCH for the NDRE VI, to 0.69 in the 2017/2018 season AYN at MP for the NDRE VI.

GWAS

To try and determine if any of the VI had genomic regions associated with a large effect size, a GWAS was conducted. The PCA of the 1989 experimental lines genotypic data shows no distinct population structure and the plot is bounded by the expected founder genotypes, Figure 5. There are no major QTLs for grain yield in any year, yet several of the VIs showed significant associations with regions of the genome. When these associations were further examined it was shown that they had an influence on the VI value but not GRYLD (Supplementary Figure 1). The regions that showed an association with the VI were often shared between different VI.

Genomic Prediction

Based on the GWAS, GP was evaluated to predict the highly quantitative trait of grain yield. All of the tested genomic prediction models produced similar accuracies (Supplementary Table 2), therefore we focused on rrBLUP models for computational

360   efficiently for further analysis which will be reported. The genomic prediction accuracies based on cross-validation range between 0.311 (SD = 0.079) for the 2017/2018 season and 0.469 (SD = 0.105) for the 2016/2017 season (Figure 6, Supplementary Table 2).

When making blind predictions the strongest correlation, -0.164, was achieved using all seasons excluding the 2016/2017 season as the training population, and the

365   2016/2017 season as the prediction population (Table 4). When predicting all locations in a single season except for HUTCH, using the data from other seasons and the data from HUTCH as the training population, the greatest accuracy achieved was 0.572 (95% CI [0.503, 0.634]) for 2018/2019 PYN in MANH and BEL. HUTCH was chosen as the location to include in the blind predictions as it is the first location that is normally

370   harvested in Kansas. This would be similar to harvesting one location and then using the predictions to prioritize which locations to harvest.

The VI were used as covariates in the prediction of grain yield. The prediction of grain yield was not improved by the addition of a covariate (Figure 7). The prediction of grain yield without a covariate was performed at the same time for comparison and was

375   comparable to the best estimate using VI as covariate.

Discussion

Changing global conditions require current food production systems to be resilient against unexpected events in the face of global population growth. This will require the development of crop varieties that are adapted to the changing climatic, land-usage and population conditions. The speed at which these varieties are required will require the adoption of new technologies that have been proven to show a positive economic investment.

This study examined the validity of using GP and other HTP techniques in the Kansas State University winter wheat breeding program based on several population and model parameters, such as heritability of the primary trait, the prediction accuracy and the selection intensity. These parameters give an indication of whether a new GP method would provide more genetic gain than traditional phenotypic selection under differing financial and population conditions.

Heritability

The heritability of a trait has a significant impact on whether the trait can be selected for and the possibility for genetic gain of that trait. Traits with a lower heritability are difficult to select for regardless of which selection method is used. However, our simulation showed that traits with a low to moderate narrow-sense heritability (0-0.4) favor the use of GP. For the Kansas State University winter wheat breeding program, the overall broad-sense heritability of grain yield is below 0.3. As the

narrow-sense heritability is the always less than the broad-sense heritability, the

heritability favors GP under the operation cost parameters for the program.

Prediction Accuracy

The prediction accuracy for GP when making blind predictions in the breeding

400   program do not meet the criteria to favor GP unless very large populations are used.

This is a possibility in a breeding program setting as historical grain yield data and

genotypes are available for previous seasons. The experimental lines in these seasons

are likely to be less related to the current lines as the parental lines in the crossing block

are updated which may lower prediction accuracy, but the sheer amount of data is likely

405   to be able to account for the lack of relatedness.

HTP Prediction Accuracy

Utilizing the VI to increase the prediction accuracy of the GP models requires

more optimization before it is commonly adopted. The VI need to be able to be used

across locations and years for them to really have an impact on the prediction accuracy

410   of the GP models. This will require either the manual measurement of growth stages to

standardize growth phases across fields or the use of another method to account for

differences in growth phase in the training populations and prediction populations. This

additional labor makes the use of VIs in GP models more costly than utilizing just the GP

model. It is an additional factor that needs to be taken into account before the decision

415   to transition to GP is made.

Conclusion

When all parameters are considered it appears that the Kansas State University winter wheat breeding program is on the edge of a large decision. The heritability of grain yield in the breeding program as well as the cost of phenotyping compared to

420   genotyping favor genomic prediction as the way forward for the breeding program. Yet the low accuracy of blind predictions favors the use of phenotypic selection. The blind prediction accuracy can be increased as seen in the 2018/2019 season (Table 4), but this requires a much larger training population. With a few adjustments to the experimental design such as allowing for more replicates of the training population, and changes to

425   the program workflow that allows for the loss of the PYN, genomic prediction could allow the KSU winter wheat breeding program to make larger genetic gains.

Authors' contributions

MC and BE collected data. MC and XW analyzed data. JP conceived and planned the

430   experiments. All authors read and approved the final manuscript.

Acknowledgements

We would like to acknowledge all of the undergraduate assistants who assisted with data collection. Daljit Singh, Richard Brown and Grant Williams who assisted with

435   UAS flights and data collection.

MC was supported in part by fellowship funds from the Interdepartmental Genetics program at Kansas State University. This material is based upon work

supported by the National Science Foundation under Grant No. 1238187 project "A

Field-Based High-Throughput Phenotyping Platform for Plant Genetics" and the USDA

440   NIFA International Wheat Yield Partnership grant no. 2017-67007-25933/project

accession no. 1011391 project "Wheat Yield Prediction and Advanced Selection

Methodologies through Field-Based High-Throughput Phenotyping with UAVs". Any

opinions, findings, and conclusions or recommendations expressed in this material are

those of the author(s) and do not necessarily reflect the views of the National Science

445   Foundation or the U.S. Dept. of Agriculture.

References

Bassi, Filippo M., Alison R. Bentley, Gilles Charmet, Rodomiro Ortiz, and Jose Crossa. 2016. "Breeding Schemes for the Implementation of Genomic Selection in Wheat (*Triticum* Spp .)." Plant Science 242 (January): 23–36. https://doi.org/10.1016/j.plantsci.2015.08.021.

Bernardo, Rex. 2008. "Molecular Markers and Selection for Complex Traits in Plants: Learning from the Last 20 Years." Crop Science 48 (5): 1649. https://doi.org/10.2135/cropsci2008.03.0131.

Blackmer, Tracy M., James S. Schepers, Gary E. Varvel, and Elizabeth A. Walter-Shea. 1996. "Nitrogen Deficiency Detection Using Reflected Shortwave Radiation from Irrigated Corn Canopies." Agronomy Journal 88 (1): 1–5. https://doi.org/10.2134/agronj1996.00021962008800010001x.

Bond, Jennifer K. 2020. "USDA ERS - Wheat Sector at a Glance." June 26, 2020. https://www.ers.usda.gov/topics/crops/wheat/wheat-sector-at-a-glance/.

Bradbury, P. J., Z. Zhang, D. E. Kroon, T. M. Casstevens, Y. Ramdoss, and E. S. Buckler. 2007. "TASSEL: Software for Association Mapping of Complex Traits in Diverse Samples." Bioinformatics 23 (19): 2633–35. https://doi.org/10.1093/bioinformatics/btm308.

Browning, Brian L., Ying Zhou, and Sharon R. Browning. 2018. "A One-Penny Imputed Genome from Next-Generation Reference Panels." The American Journal of Human Genetics 103 (3): 338–48. https://doi.org/10.1016/j.ajhg.2018.07.015.

Burgueño, Juan, Gustavo de los Campos, Kent Weigel, and José Crossa. 2012. "Genomic Prediction of Breeding Values When Modeling Genotype × Environment Interaction Using Pedigree and Dense Molecular Markers." Crop Science 52 (2): 707–19. https://doi.org/10.2135/cropsci2011.06.0299.

Core Team, R. 2020. R: A Language and Environment for Statistical Computing (version 4.0.0). Vienna, Austria: R Foundation for Statistical Computing. https://www.R-project.org/.

Crain, Jared, Suchismita Mondal, Jessica Rutkoski, Ravi P. Singh, and Jesse Poland. 2018. "Combining High-Throughput Phenotyping and Genomic Information to Increase Prediction and Selection Accuracy in Wheat Breeding." The Plant Genome 11 (1): 0. https://doi.org/10.3835/plantgenome2017.05.0043.

Curran, P. J., M. D. Steven, John Theodore Houghton, Arthur Herbert Cook, and Henry Charnock. 1983. "Multispectral Remote Sensing for the Estimation of Green Leaf Area Index." Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences 309 (1508): 257–70. https://doi.org/10.1098/rsta.1983.0039.

485 Dawson, Julie C., Jeffrey B. Endelman, Nicolas Heslot, Jose Crossa, Jesse Poland, Susanne Dreisigacker, Yann Manès, Mark E. Sorrells, and Jean-Luc Jannink. 2013. "The Use of Unbalanced Historical Data for Genomic Selection in an International Wheat Breeding Program." Field Crops Research 154 (December): 12–22. https://doi.org/10.1016/j.fcr.2013.07.020.

490 Elliott, G. A., and K. L. Regan. 1993. "Use of Reflectance Measurements to Estimate Early Cereal Biomass Production on Sandplain Soils." Australian Journal of Experimental Agriculture 33 (2): 179–83. https://doi.org/10.1071/ea9930179.

Endelman, Jeffrey B. 2011. "Ridge Regression and Other Kernels for Genomic Selection with R Package RrBLUP." The Plant Genome Journal 4 (3): 250. https://doi.org/10.3835/plantgenome2011.08.0024.

495 Falconer, Douglas S., and Trudy F. C. Mackay. 2009. Introduction to Quantitative Genetics. 4. ed., [16. print.]. Harlow: Pearson, Prentice Hall.

Federer, W. T., and D. Raghavarao. 1975. "On Augmented Designs." Biometrics 31 (1): 29–35. https://doi.org/10.2307/2529707.

Georges, Michel, Carole Charlier, and Ben Hayes. 2019. "Harnessing Genomic
500 Information for Livestock Improvement." Nature Reviews Genetics 20 (3): 135–56. https://doi.org/10.1038/s41576-018-0082-2.

Glaubitz, Jeffrey C., Terry M. Casstevens, Fei Lu, James Harriman, Robert J. Elshire, Qi Sun, and Edward S. Buckler. 2014. "TASSEL-GBS: A High Capacity Genotyping by Sequencing Analysis Pipeline." Edited by Nicholas A. Tinker. PLoS ONE 9 (2):
505 e90346. https://doi.org/10.1371/journal.pone.0090346.

Hayes, B. J., P. J. Bowman, A. J. Chamberlain, and M. E. Goddard. 2009. "Invited Review: Genomic Selection in Dairy Cattle: Progress and Challenges." Journal of Dairy Science 92 (2): 433–43. https://doi.org/10.3168/jds.2008-1646.

Heslot, Nicolas, Hsiao-Pei Yang, Mark E. Sorrells, and Jean-Luc Jannink. 2012. "Genomic
510 Selection in Plant Breeding: A Comparison of Models." Crop Science 52 (1): 146–60. https://doi.org/10.2135/cropsci2011.06.0297.

Holland, James B., Wyman E. Nyquist, and Cuauhtemoc T. Cervantes-Martínez. 2010. "Estimating and Interpreting Heritability for Plant Breeding: An Update." In Plant Breeding Reviews, 9–112. John Wiley & Sons, Ltd.
515 https://doi.org/10.1002/9780470650202.ch2.

International Wheat Genome Sequencing Consortium, Michael Alaux, Jane Rogers, Thomas Letellier, Raphaël Flores, Françoise Alfama, Cyril Pommier, et al. 2018. "Linking the International Wheat Genome Sequencing Consortium Bread Wheat Reference Genome Sequence to Wheat Genetic and Phenomic Data." Genome
520 Biology 19 (1): 111. https://doi.org/10.1186/s13059-018-1491-4.

Jackson, Christopher. 2011. "Multi-State Models for Panel Data: The Msm Package for R." Journal of Statistical Software 38 (1): 1–28. https://doi.org/10.18637/jss.v038.i08.

Jarquín, Diego, Cristiano Lemes da Silva, R. Chris Gaynor, Jesse Poland, Allan Fritz, Reka Howard, Sarah Battenfield, and Jose Crossa. 2017. "Increasing Genomic-Enabled Prediction Accuracy by Modeling Genotype × Environment Interactions in Kansas Wheat." The Plant Genome 10 (2): plantgenome2016.12.0130. https://doi.org/10.3835/plantgenome2016.12.0130.

Meuwissen, T. H. E., B. J. Hayes, and M. E. Goddard. 2001. "Prediction of Total Genetic Value Using Genome-Wide Dense Marker Maps." Genetics 157 (4): 1819–29.

Montesinos-López, Osval A., Abelardo Montesinos-López, José Crossa, Fernando H. Toledo, Oscar Pérez-Hernández, Kent M. Eskridge, and Jessica Rutkoski. 2016. "A Genomic Bayesian Multi-Trait and Multi-Environment Model." G3: Genes|Genomes|Genetics 6 (9): 2725–44. https://doi.org/10.1534/g3.116.032359.

Moose, Stephen P., and Rita H. Mumm. 2008. "Molecular Plant Breeding as the Foundation for 21st Century Crop Improvement." Plant Physiology 147 (3): 969–77. https://doi.org/10.1104/pp.108.118232.

Patterson, H. D., and E. R. Williams. 1976. "A New Class of Resolvable Incomplete Block Designs." Biometrika 63 (1): 83–92. https://doi.org/10.2307/2335087.

Pérez, Paulino, and Gustavo de los Campos. 2014. "Genome-Wide Regression and Prediction with the BGLR Statistical Package." Genetics 198 (2): 483–95. https://doi.org/10.1534/genetics.114.164442.

Pérez-Rodríguez, Paulino, José Crossa, Jessica Rutkoski, Jesse Poland, Ravi Singh, Andrés Legarra, Enrique Autrique, Gustavo de los Campos, Juan Burgueño, and Susanne Dreisigacker. 2017. "Single-Step Genomic and Pedigree Genotype × Environment Interaction Models for Predicting Wheat Lines in International Environments." The Plant Genome 10 (2): 0. https://doi.org/10.3835/plantgenome2016.09.0089.

Poland, Jesse A., Patrick J. Brown, Mark E. Sorrells, and Jean-Luc Jannink. 2012. "Development of High-Density Genetic Maps for Barley and Wheat Using a Novel Two-Enzyme Genotyping-by-Sequencing Approach." Edited by Tongming Yin. PLoS ONE 7 (2): e32253. https://doi.org/10.1371/journal.pone.0032253.

Rife, Trevor W., and Jesse A. Poland. 2014. "Field Book: An Open-Source Application for Field Data Collection on Android." Crop Science 54 (4): 1624–27. https://doi.org/10.2135/cropsci2013.08.0579.

Rutkoski, Jessica, Jesse Poland, Suchismita Mondal, Enrique Autrique, Lorena González Pérez, José Crossa, Matthew Reynolds, and Ravi Singh. 2016. "Canopy Temperature and Vegetation Indices from High-Throughput Phenotyping Improve Accuracy of Pedigree and Genomic Selection for Grain Yield in Wheat."

560   G3: Genes|Genomes|Genetics 6 (9): 2799–2808.
https://doi.org/10.1534/g3.116.032888.

Stacklies, W., H. Redestig, M. Scholz, D. Walther, and J. Selbig. 2007. "PcaMethods a Bioconductor Package Providing PCA Methods for Incomplete Data." Bioinformatics 23 (9): 1164–67. https://doi.org/10.1093/bioinformatics/btm069.

565   Wang, Xu, Daljit Singh, Sandeep Marla, Geoffrey Morris, and Jesse Poland. 2018. "Field-Based High-Throughput Phenotyping of Plant Height in Sorghum Using Different Sensing Technologies." Plant Methods 14 (1): 53. https://doi.org/10.1186/s13007-018-0324-5.

Wickham, Hadley, Mara Averick, Jennifer Bryan, Winston Chang, Lucy D'Agostino
570   McGowan, Romain François, Garrett Grolemund, et al. 2019. "Welcome to the Tidyverse." Journal of Open Source Software 4 (43): 1686. https://doi.org/10.21105/joss.01686.

Figures

Figure 0.1. Simulation results comparing phenotypic selection as a direct trait to selection on the genotype as a secondary trait.

The narrow-sense heritability of the direct trait is in the panel title and the y-axis is the Correlated Response / Response of selection for the direct trait. When this ratio is above 1, represented by the dotted line, selection on the secondary trait is favored. The x-axis is the number of experimental lines that can be genotyped and predicted for every line that is phenotyped based on cost. The trend of each prediction accuracy is given by the color of the line.

Figure 0.2. Broad sense heritabilities for grain yield across 4 years of phenotypic data.

The broad sense heritabilities are presented by season on the x-axis with each trial type denoted by the character shape. There is no PYN for the 2019 season as it was only planted in one location with one replication.

Figure 0.3. The broad sense heritabilities for the VIs for the AYN trials planted at each location.

The title of each plot gives the season-VI combination, with the date on which the VI was taken given by the x-axis, the y-axis is the broad sense heritability, while the color of the point determines the location. NDVI = Normalized Difference Vegetation Index, GNDVI = Green Normalized Difference Vegetation Index, and NDRE = Red Edge Normalized Difference Vegetation Index.

Figure 0.4. The correlations between grain yield and VI by year-trial combinations.

The title of each plot gives the season-VI combination, with the date on which the VI was taken given by the x-axis, the y-axis is the Pearson correlation coefficient, while the color of the point determines the location. NDVI = Normalized Difference Vegetation Index, GNDVI = Green Normalized Difference Vegetation Index, and NDRE = Red Edge Normalized Difference Vegetation Index.

Figure 0.5. Population structure based on genotypic information in the Kansas State winter wheat breeding program nurseries between the 2016-2019 seasons.

The top panel shows the first two principal components (PC), while the bottom panel displays the 2nd and 3rd PC. The variance contributed by each PC is given next to the PC name. Several "founder" lines are highlighted.

Figure 0.6. Genomic prediction accuracies for grain yield determined by cross-validation.

The season or seasons used in each analysis are given on the x-axis. The y-axis shows the range of possible correlations between the predicted phenotype and the observed phenotype. The color of the point determines if the individual line had been in the training or testing population of the analysis. 100 cross-validations were performed in each analysis.

Figure 0.7. Genomic prediction accuracies for PYN grain yield when a VI is used as a covariate determined by cross-validation.

The season and VI are given in the strip title and the points are colored by the location. The GRYLD measurements are those for the prediction of grain yield without a covariate. The large point is the mean and the error bars give the standard deviation. NDVI = Normalized Difference Vegetation Index, GNDVI = Green Normalized Difference Vegetation Index, and NDRE = Red Edge Normalized Difference Vegetation Index.

Tables

Table 0.1. Summary of trials planted between the 2016 and 2019 seasons across locations. A summary is given by season, trial and location with the number of individual plots, experimental lines, date of planting, date of harvest, the mean grain yield in bu/ac and the standard deviation of the grain yield.

Season	Trial	Location	Plots	Lines	Planted	Harvested	Mean	SD
2015/2016	AYN	MP	40	75			5.050	0.9
		MANH	240	75			3.206	0.651
		HUTCH	320	109			4.139	0.92
		GYP	40	109			5.739	0.726
	PYN	MP	504	439			5.113	1.21
		MANH	504	379			3.720	0.657
		HUTCH	594	516			3.770	0.886
		GYP	593	506			5.421	0.702
2016/2017	AYN	MP	315	91	10/11/16	20/06/17	5.933	1.00
		MANH	320	91	10/18/16	22/06/17	5.822	0.768
		HUTCH	168	97	10/12/16	21/06/17	6.242	0.845
		BEL	48	36	10/19/16	28/06/17	5.110	0.776
	PYN	MP	126	120	10/11/16	20/06/17	4.805	1.015
		HUTCH	108	102	10/12/16	21/06/17	6.319	0.925
		BEL	8	8	10/19/16	28/06/17	4.5	0.385
2017/2018	AYN	MP	199	90	10/19/17	29/06/18	3.203	0.412
		MANH	200	90	10/20/17	23/06/18	2.644	0.402
		HUTCH	280	125	10/18/17	28/06/18	3.345	0.489
		BEL	280	125	10/17/17	30/06/18	2.561	0.430
		GYP	279	125	10/18/17	18/06/18	2.343	0.318
	PYN	MP	323	282	10/19/17	29/06/18	3.048	0.490

		MANH	414	366	10/20/17	23/06/18	2.880	0.431
		HUTCH	414	366	10/18/17	28/06/18	3.236	0435
		BEL	287	251	10/17/17	30/06/18	2.198	0.412
		GYP	286	250	10/18/17	18/06/18	1.973	0.278
2018/2019	AYN	MANH	80	40	11/01/18		5.798	0.535
		HUTCH	212	99	10/24/18	01/07/19	4.925	0.563
		BEL	130	63	10/24/18	17/07/19	4.965	0.897
	PYN	HUTCH	504	454	10/24/18	01/07/19	4.623	0.794

Table 0.2. Summary of vegetation indices used in the KSU breeding program across 5 years. For each index the formula and reference are provided.

	Index	Equation	Reference
GNDVI	Green Normalized Difference Vegetation Index	$GNDVI = \dfrac{NIR - Green}{NIR + Green}$	(Blackmer et al. 1996)
NDRE	Red Edge Normalized Difference Vegetation Index	$NDRE = \dfrac{NIR - RedEdge}{NIR + RedEdge}$	(Elliott and Regan 1993)
NDVI	Normalized Difference Vegetation Index	$NDVI = \dfrac{NIR - Red}{NIR + Red}$	(Curran et al. 1983)

Table 0.3. Genomic Prediction Accuracies as the correlation between BLUPs and GEBVs. The Pearson correlation coefficients is given with a 95% CI.

Training Population	Testing Population	Correlation	95% CI
Excluding 2015/2016	2015/2016	0.09	[-0.004, 0.182]
Excluding 2015/2016 PYN	2015/2016 PYN	0.078	[-0.026, 0.180]
Excluding 2015/2016 MP MANH GYP	2015/2016 MP MANH GYP	0.024	[-0.08, 0.127]
Excluding 2016/2017	2016/2017	-0.164	[-0.336, 0.018]
Excluding 2016/2017 PYN	2016/2017 PYN	-0.055	[-0.249, 0.144]
Excluding 2016/2017 MP MANH BEL	2016/2017 MP MANH BEL	-0.009	[-0.206, 0.189]
Excluding 2017/2018	2017/2018	0.048	[-0.049, 0.145]
Excluding 2017/2018 PYN	2017/2018 PYN	0.061	[-0.046, 0.165]
Excluding 2017/2018 MP MANH BEL	2017/2018 MP MANH BEL	0.009	[-0.097, 0.114]
Excluding 2018/2019	2018/2019	-0.091	[-0.186, 0.005]
Excluding 2018/2019 PYN	2018/2019 PYN	-0.039	[-0.135, 0.058]
Excluding 2018/2019 MANH BEL	2018/2019 MANH BEL	0.572	[0.502, 0.634]

References

620   Álvarez, María F., Teresa Mosquera, and Matthew W. Blair. 2014. "The Use of Association Genetics Approaches in Plant Breeding." In *Plant Breeding Reviews: Volume 38*, 17–68. John Wiley & Sons, Ltd. https://doi.org/10.1002/9781118916865.ch02.

         Ammiraju, J. S. S., B. B. Dholakia, D. K. Santra, H. Singh, M. D. Lagu, S. A. Tamhankar, H.
625   S. Dhaliwal, V. S. Rao, V. S. Gupta, and P. K. Ranjekar. 2001. "Identification of Inter Simple Sequence Repeat (ISSR) Markers Associated with Seed Size in Wheat:" *Theoretical and Applied Genetics* 102 (5): 726–32. https://doi.org/10.1007/s001220051703.

         Anand, Shivani. 2019. "India Smartphone Market Registers Highest Second-Quarter
630   Shipments in 2Q19, IDC India Reports." IDC: The Premier Global Market Intelligence Company. August 13, 2019. https://www.idc.com/getdoc.jsp?containerId=prAP45442419.

         Andrade-Sanchez, Pedro, Michael A. Gore, John T. Heun, Kelly R. Thorp, A. Elizabete Carmo-Silva, Andrew N. French, Michael E. Salvucci, and Jeffrey W. White. 2014.
635   "Development and Evaluation of a Field-Based High-Throughput Phenotyping Platform." *Functional Plant Biology* 41 (1): 68. https://doi.org/10.1071/FP13126.

         Araus, José Luis, and Jill E. Cairns. 2014. "Field High-Throughput Phenotyping: The New Crop Breeding Frontier." *Trends in Plant Science* 19 (1): 52–61. https://doi.org/10.1016/j.tplants.2013.09.008.

640   Bajarin, Ben. 2014. "Why India Will Be the World's Second Biggest Smartphone Market." Fortune. December 2, 2014. https://fortune.com/2014/12/02/why-india-will-be-the-worlds-second-biggest-smartphone-market/.

         Barker, Jared, Naiqian Zhang, Joshua Sharon, Ryan Steeves, Xu Wang, Yong Wei, and Jesse Poland. 2016. "Development of a Field-Based High-Throughput Mobile
645   Phenotyping Platform." *Computers and Electronics in Agriculture* 122 (March): 74–85. https://doi.org/10.1016/j.compag.2016.01.017.

         Bassi, Filippo M., Alison R. Bentley, Gilles Charmet, Rodomiro Ortiz, and Jose Crossa. 2016. "Breeding Schemes for the Implementation of Genomic Selection in Wheat (Triticum Spp .)." *Plant Science* 242 (January): 23–36.
650   https://doi.org/10.1016/j.plantsci.2015.08.021.

         Bean, S. R., O. K. Chung, M. R. Tuinstra, J. F. Pedersen, and J. Erpelding. 2006. "Evaluation of the Single Kernel Characterization System (SKCS) for Measurement of Sorghum Grain Attributes." *Cereal Chemistry Journal* 83 (1): 108–13. https://doi.org/10.1094/CC-83-0108.

655   Beavis, William D. 1998. "QTL Analyses: Power, Precision, and Accuracy." *Molecular Dissection of Complex Traits*, 145–62.

Bernardo, Rex. 2008. "Molecular Markers and Selection for Complex Traits in Plants: Learning from the Last 20 Years." *Crop Science* 48 (5): 1649. https://doi.org/10.2135/cropsci2008.03.0131.

660   Blackmer, Tracy M., James S. Schepers, Gary E. Varvel, and Elizabeth A. Walter-Shea. 1996. "Nitrogen Deficiency Detection Using Reflected Shortwave Radiation from Irrigated Corn Canopies." *Agronomy Journal* 88 (1): 1–5. https://doi.org/10.2134/agronj1996.00021962008800010001x.

Blake, Victoria C., Clay Birkett, David E. Matthews, David L. Hane, Peter Bradbury, and
665   Jean-Luc Jannink. 2016. "The Triticeae Toolbox: Combining Phenotype and Genotype Data to Advance Small-Grains Breeding." *The Plant Genome* 9 (2): 0. https://doi.org/10.3835/plantgenome2014.12.0099.

Bond, Jennifer K. 2020. "USDA ERS - Wheat Sector at a Glance." June 26, 2020. https://www.ers.usda.gov/topics/crops/wheat/wheat-sector-at-a-glance/.

670   Bradbury, P. J., Z. Zhang, D. E. Kroon, T. M. Casstevens, Y. Ramdoss, and E. S. Buckler. 2007. "TASSEL: Software for Association Mapping of Complex Traits in Diverse Samples." *Bioinformatics* 23 (19): 2633–35. https://doi.org/10.1093/bioinformatics/btm308.

Bradski, Gary. 2000. *The OpenCV Library*. https://www.drdobbs.com/open-source/the-
675   opencv-library/184404319.

Breseghello, Flavio, and Mark E. Sorrells. 2006. "Association Mapping of Kernel Size and Milling Quality in Wheat (Triticum Aestivum L.) Cultivars." *Genetics* 172 (2): 1165–77. https://doi.org/10.1534/genetics.105.044586.

Breseghello, Flavio, and Mark E. Sorrells. 2007. "QTL Analysis of Kernel Size and Shape
680   in Two Hexaploid Wheat Mapping Populations." *Field Crops Research* 101 (2): 172–79. https://doi.org/10.1016/j.fcr.2006.11.008.

Broman, Karl W., Hao Wu, Saunak Sen, and Gary A. Churchill. 2003. "R/Qtl: QTL Mapping in Experimental Crosses." *Bioinformatics (Oxford, England)* 19 (7): 889–90. https://doi.org/10.1093/bioinformatics/btg112.

685   Browning, Brian L., Ying Zhou, and Sharon R. Browning. 2018. "A One-Penny Imputed Genome from Next-Generation Reference Panels." *The American Journal of Human Genetics* 103 (3): 338–48. https://doi.org/10.1016/j.ajhg.2018.07.015.

Browning, Brian L., and Sharon R. Browning. 2016. "Genotype Imputation with Millions of Reference Samples." *The American Journal of Human Genetics* 98 (1): 116–26.
690   https://doi.org/10.1016/j.ajhg.2015.11.020.

Burgueño, Juan, Gustavo de los Campos, Kent Weigel, and José Crossa. 2012. "Genomic Prediction of Breeding Values When Modeling Genotype × Environment Interaction Using Pedigree and Dense Molecular Markers." *Crop Science* 52 (2): 707–19. https://doi.org/10.2135/cropsci2011.06.0299.

695 Busemeyer, Lucas, Daniel Mentrup, Kim Möller, Erik Wunder, Katharina Alheit, Volker Hahn, Hans Peter Maurer, et al. 2013. "BreedVision — A Multi-Sensor Platform for Non-Destructive Field-Based Phenotyping in Plant Breeding." *Sensors* 13 (3): 2830–47. https://doi.org/10.3390/s130302830.

Butler, David. 2020. *Asreml: Fits the Linear Mixed Model*. www.vsni.co.uk.

700 Cabrera-Bosquet, Llorenç, José Crossa, Jarislav von Zitzewitz, María Dolors Serret, and José Luis Araus. 2012. "High-Throughput Phenotyping and Genomic Selection: The Frontiers of Crop Breeding Converge: High-Throughput Phenotyping and Genomic Selection." *Journal of Integrative Plant Biology* 54 (5): 312–20. https://doi.org/10.1111/j.1744-7909.2012.01116.x.

705 Campbell, Kim Garland, Christine J. Bergman, Daisy G. Gualberto, James A. Anderson, Micheal J. Giroux, Gary Hareland, R. Gary Fulcher, Mark E. Sorrells, and Patrick L. Finney. 1999. "Quantitative Trait Loci Associated with Kernel Traits in a Soft × Hard Wheat Cross." *Crop Science* 39 (4): 1184. https://doi.org/10.2135/cropsci1999.0011183X003900040039x.

710 Chaerle, L., and D. Van Der Straeten. 2001. "Seeing Is Believing: Imaging Techniques to Monitor Plant Health." *Biochimica Et Biophysica Acta* 1519 (3): 153–66. https://doi.org/10.1016/s0167-4781(01)00238-x.

Core Team, R. 2020. *R: A Language and Environment for Statistical Computing* (version 4.0.0). Vienna, Austria: R Foundation for Statistical Computing. https://www.R-
715 project.org/.

Crain, Jared L., Yong Wei, Jared Barker, Sean M. Thompson, Phillip D. Alderman, Matthew Reynolds, Naiqian Zhang, and Jesse Poland. 2016. "Development and Deployment of a Portable Field Phenotyping Platform." *Crop Science* 56 (3): 965–75. https://doi.org/10.2135/cropsci2015.05.0290.

720 Crain, Jared, Suchismita Mondal, Jessica Rutkoski, Ravi P. Singh, and Jesse Poland. 2018. "Combining High-Throughput Phenotyping and Genomic Information to Increase Prediction and Selection Accuracy in Wheat Breeding." *The Plant Genome* 11 (1): 0. https://doi.org/10.3835/plantgenome2017.05.0043.

Curran, P. J., M. D. Steven, John Theodore Houghton, Arthur Herbert Cook, and Henry
725 Charnock. 1983. "Multispectral Remote Sensing for the Estimation of Green Leaf Area Index." *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences* 309 (1508): 257–70. https://doi.org/10.1098/rsta.1983.0039.

Dawson, Julie C., Jeffrey B. Endelman, Nicolas Heslot, Jose Crossa, Jesse Poland,
730 Susanne Dreisigacker, Yann Manès, Mark E. Sorrells, and Jean-Luc Jannink. 2013. "The Use of Unbalanced Historical Data for Genomic Selection in an International Wheat Breeding Program." *Field Crops Research* 154 (December): 12–22. https://doi.org/10.1016/j.fcr.2013.07.020.

735 Dekkers, Jack C. M., and Frédéric Hospital. 2002. "The Use of Molecular Genetics in the Improvement of Agricultural Populations." *Nature Reviews Genetics* 3 (1): 22–32. https://doi.org/10.1038/nrg701.

Dholakia, B. B., J. S. S. Ammiraju, H. Singh, M. D. Lagu, M. S. Roder, V. S. Rao, H. S. Dhaliwal, P. K. Ranjekar, and V. S. Gupta. 2003. "Molecular Marker Analysis of Kernel Size and Shape in Bread Wheat." *Plant Breeding* 122 (5): 392–95.
740 https://doi.org/10.1046/j.1439-0523.2003.00896.x.

Dhondt, Stijn, Nathalie Wuyts, and Dirk Inzé. 2013. "Cell to Whole-Plant Phenotyping: The Best Is yet to Come." *Trends in Plant Science* 18 (8): 428–39. https://doi.org/10.1016/j.tplants.2013.04.008.

Dunckel, Sandra M. 2015. "Whole Genome Approaches for Characterizing and Utilizing
745 Synthetic Wheat." https://krex.k-state.edu/dspace/handle/2097/20518.

Eathington, Sam R., Theodore M. Crosbie, Marlin D. Edwards, Robert S. Reiter, and Jason K. Bull. 2007. "Molecular Markers in a Commercial Breeding Program." *Crop Science* 47 (S3): S-154-S-163. https://doi.org/10.2135/cropsci2007.04.0015IPBS.

750 Elliott, G. A., and K. L. Regan. 1993. "Use of Reflectance Measurements to Estimate Early Cereal Biomass Production on Sandplain Soils." *Australian Journal of Experimental Agriculture* 33 (2): 179–83. https://doi.org/10.1071/ea9930179.

Elshire, Robert J., Jeffrey C. Glaubitz, Qi Sun, Jesse A. Poland, Ken Kawamoto, Edward S. Buckler, and Sharon E. Mitchell. 2011. "A Robust, Simple Genotyping-by-
755 Sequencing (GBS) Approach for High Diversity Species." Edited by Laszlo Orban. *PLoS ONE* 6 (5): e19379. https://doi.org/10.1371/journal.pone.0019379.

Endelman, Jeffrey B. 2011. "Ridge Regression and Other Kernels for Genomic Selection with R Package RrBLUP." *The Plant Genome Journal* 4 (3): 250. https://doi.org/10.3835/plantgenome2011.08.0024.

760 Endelman, Jeffrey B., and Jean-Luc Jannink. 2012. "Shrinkage Estimation of the Realized Relationship Matrix." *G3: Genes|Genomes|Genetics* 2 (11): 1405–13. https://doi.org/10.1534/g3.112.004259.

Falconer, Douglas S., and Trudy F. C. Mackay. 2009. *Introduction to Quantitative Genetics*. 4. ed., [16. print.]. Harlow: Pearson, Prentice Hall.

765 Federer, W. T., and D. Raghavarao. 1975. "On Augmented Designs." *Biometrics* 31 (1): 29–35. https://doi.org/10.2307/2529707.

Furbank, Robert T., and Mark Tester. 2011. "Phenomics – Technologies to Relieve the Phenotyping Bottleneck." *Trends in Plant Science* 16 (12): 635–44. https://doi.org/10.1016/j.tplants.2011.09.005.

770 Georges, Michel, Carole Charlier, and Ben Hayes. 2019. "Harnessing Genomic Information for Livestock Improvement." *Nature Reviews Genetics* 20 (3): 135–56. https://doi.org/10.1038/s41576-018-0082-2.

775 Gitelson, Anatoly A., Yoram J. Kaufman, and Mark N. Merzlyak. 1996. "Use of a Green Channel in Remote Sensing of Global Vegetation from EOS-MODIS." *Remote Sensing of Environment* 58 (3): 289–98. https://doi.org/10.1016/S0034-4257(96)00072-7.

Giura, A., and N. N. Saulescu. 1996. "Chromosomal Location of Genes Controlling Grain Size in a Large Grained Selection of Wheat (Triticum Aestivum L.)." *Euphytica* 89 (1): 77–80. https://doi.org/10.1007/BF00015722.

780 Glaubitz, Jeffrey C., Terry M. Casstevens, Fei Lu, James Harriman, Robert J. Elshire, Qi Sun, and Edward S. Buckler. 2014. "TASSEL-GBS: A High Capacity Genotyping by Sequencing Analysis Pipeline." Edited by Nicholas A. Tinker. *PLoS ONE* 9 (2): e90346. https://doi.org/10.1371/journal.pone.0090346.

Haghighattalab, Atena, Lorena González Pérez, Suchismita Mondal, Daljit Singh, Dale
785 Schinstock, Jessica Rutkoski, Ivan Ortiz-Monasterio, Ravi Prakash Singh, Douglas Goodin, and Jesse Poland. 2016. "Application of Unmanned Aerial Systems for High Throughput Phenotyping of Large Wheat Breeding Nurseries." *Plant Methods* 12 (1): 35. https://doi.org/10.1186/s13007-016-0134-6.

Hayes, B. J., P. J. Bowman, A. J. Chamberlain, and M. E. Goddard. 2009. "Invited Review:
790 Genomic Selection in Dairy Cattle: Progress and Challenges." *Journal of Dairy Science* 92 (2): 433–43. https://doi.org/10.3168/jds.2008-1646.

Heslot, Nicolas, Hsiao-Pei Yang, Mark E. Sorrells, and Jean-Luc Jannink. 2012. "Genomic Selection in Plant Breeding: A Comparison of Models." *Crop Science* 52 (1): 146–60. https://doi.org/10.2135/cropsci2011.06.0297.

795 Holland, James B., Wyman E. Nyquist, and Cuauhtemoc T. Cervantes-Martínez. 2010. "Estimating and Interpreting Heritability for Plant Breeding: An Update." In *Plant Breeding Reviews*, 9–112. John Wiley & Sons, Ltd. https://doi.org/10.1002/9780470650202.ch2.

Houle, David, Diddahally R. Govindaraju, and Stig Omholt. 2010. "Phenomics: The next
800 Challenge." *Nature Reviews Genetics* 11 (12): 855–66. https://doi.org/10.1038/nrg2897.

Huang, Ning, Arnold Parco, Teresita Mew, Gerard Magpantay, Susan McCouch, Emmanuel Guiderdoni, Jichen Xu, Prasanta Subudhi, Enrique R. Angeles, and Gurdev S. Khush. 1997. "RFLP Mapping of Isozymes, RAPD and QTLs for Grain
805 Shape, Brown Planthopper Resistance in a Doubled Haploid Rice Population." *Molecular Breeding* 3 (2): 105–13. https://doi.org/10.1023/A:1009683603862.

International Wheat Genome Sequencing Consortium, Michael Alaux, Jane Rogers, Thomas Letellier, Raphaël Flores, Françoise Alfama, Cyril Pommier, et al. 2018. "Linking the International Wheat Genome Sequencing Consortium Bread Wheat
810 Reference Genome Sequence to Wheat Genetic and Phenomic Data." *Genome Biology* 19 (1): 111. https://doi.org/10.1186/s13059-018-1491-4.

Jackson, Christopher. 2011. "Multi-State Models for Panel Data: The Msm Package for R." *Journal of Statistical Software* 38 (1): 1–28. https://doi.org/10.18637/jss.v038.i08.

815 Jarquín, Diego, Cristiano Lemes da Silva, R. Chris Gaynor, Jesse Poland, Allan Fritz, Reka Howard, Sarah Battenfield, and Jose Crossa. 2017. "Increasing Genomic-Enabled Prediction Accuracy by Modeling Genotype × Environment Interactions in Kansas Wheat." *The Plant Genome* 10 (2): plantgenome2016.12.0130. https://doi.org/10.3835/plantgenome2016.12.0130.

820 Kang, Hyun Min, Noah A. Zaitlen, Claire M. Wade, Andrew Kirby, David Heckerman, Mark J. Daly, and Eleazar Eskin. 2008. "Efficient Control of Population Structure in Model Organism Association Mapping." *Genetics* 178 (3): 1709–23. https://doi.org/10.1534/genetics.107.080101.

Korkmaz, Selçuk, Dincer Goksuluk, and Gokmen Zararsiz. 2014. "MVN: An R Package for
825 Assessing Multivariate Normality." https://doi.org/10.32614/RJ-2014-031.

Lenaerts, Bert, Bertrand C.Y. Collard, and Matty Demont. 2019. "Review: Improving Global Food Security through Accelerated Plant Breeding." *Plant Science* 287 (October): 110207. https://doi.org/10.1016/j.plantsci.2019.110207.

Li, Jiming, Jinhua Xiao, Silvana Grandillo, Longying Jiang, Yizhen Wan, Qiyun Deng,
830 Longping Yuan, and Susan R. McCouch. 2004. "QTL Detection for Rice Grain Quality Traits Using an Interspecific Backcross Population Derived from Cultivated Asian (O. Sativa L.) and African (O. Glaberrima S.) Rice." *Genome* 47 (4): 697–704. https://doi.org/10.1139/g04-029.

Liu, Ying, Liwei Wang, Chuanlong Sun, Zuxin Zhang, Yonglian Zheng, and Fazhan Qiu.
835 2014. "Genetic Analysis and Major QTL Detection for Maize Kernel Size and Weight in Multi-Environments." *Theoretical and Applied Genetics* 127 (5): 1019–37. https://doi.org/10.1007/s00122-014-2276-0.

Marshall, D. R., D. J. Mares, H. J. Moss, and F. W. Ellison. 1986. "Effects of Grain Shape and Size on Milling Yields in Wheat. II. Experimental Studies." *Australian Journal*
840 *of Agricultural Research* 37 (4): 331–42. https://doi.org/10.1071/ar9860331.

Meuwissen, T. H. E., B. J. Hayes, and M. E. Goddard. 2001. "Prediction of Total Genetic Value Using Genome-Wide Dense Marker Maps." *Genetics* 157 (4): 1819–29.

Moles, A. T. 2005. "A Brief History of Seed Size." *Science* 307 (5709): 576–80. https://doi.org/10.1126/science.1104863.

845 Montesinos-López, Osval A., Abelardo Montesinos-López, José Crossa, Fernando H. Toledo, Oscar Pérez-Hernández, Kent M. Eskridge, and Jessica Rutkoski. 2016. "A Genomic Bayesian Multi-Trait and Multi-Environment Model." *G3: Genes|Genomes|Genetics* 6 (9): 2725–44. https://doi.org/10.1534/g3.116.032359.

850   Moose, Stephen P., and Rita H. Mumm. 2008. "Molecular Plant Breeding as the
           Foundation for 21st Century Crop Improvement." *Plant Physiology* 147 (3): 969–
           77. https://doi.org/10.1104/pp.108.118232.

       Neilsen, Mitchell, Shravan D Gangadhara, and Trevor W Rife. 2016. "Extending
           Watershed Segmentation Algorithms for High-Throughput Phenotyping." In .
855   Denver USA.
           https://www.researchgate.net/publication/322369315_Extending_Watershed_S
           egmentation_Algorithms_for_High-Throughput_Phenotyping.

       Ortiz, Rodomiro, Richard Trethowan, Guillermo Ortiz Ferrara, Masa Iwanaga, John H.
           Dodds, Jonathan H. Crouch, Jose Crossa, and Hans-Joachim Braun. 2007. "High
860   Yield Potential, Shuttle Breeding, Genetic Diversity, and a New International
           Wheat Improvement Strategy." *Euphytica* 157 (3): 365–84.
           https://doi.org/10.1007/s10681-007-9375-9.

       Pask, A., J. Pietragalla, D. Mullan, and M. P. Reynolds. 2012. "Physiological Breeding II: A
           Field Guide to Wheat Phenotyping."
865   https://repository.cimmyt.org/handle/10883/1288.

       Patterson, H. D., and E. R. Williams. 1976. "A New Class of Resolvable Incomplete Block
           Designs." *Biometrika* 63 (1): 83–92. https://doi.org/10.2307/2335087.

       Peng, Jinrong, Donald E. Richards, Nigel M. Hartley, George P. Murphy, Katrien M.
           Devos, John E. Flintham, James Beales, et al. 1999. "'Green Revolution' Genes
870   Encode Mutant Gibberellin Response Modulators." *Nature* 400 (6741): 256–61.
           https://doi.org/10.1038/22307.

       Pérez, Paulino, and Gustavo de los Campos. 2014. "Genome-Wide Regression and
           Prediction with the BGLR Statistical Package." *Genetics* 198 (2): 483–95.
           https://doi.org/10.1534/genetics.114.164442.

875   Pérez-Rodríguez, Paulino, José Crossa, Jessica Rutkoski, Jesse Poland, Ravi Singh, Andrés
           Legarra, Enrique Autrique, Gustavo de los Campos, Juan Burgueño, and Susanne
           Dreisigacker. 2017. "Single-Step Genomic and Pedigree Genotype × Environment
           Interaction Models for Predicting Wheat Lines in International Environments."
           The Plant Genome 10 (2): 0. https://doi.org/10.3835/plantgenome2016.09.0089.

880   Poland, Jesse, Jeffrey Endelman, Julie Dawson, Jessica Rutkoski, Shuangye Wu, Yann
           Manes, Susanne Dreisigacker, et al. 2012. "Genomic Selection in Wheat Breeding
           Using Genotyping-by-Sequencing." *The Plant Genome Journal* 5 (3): 103.
           https://doi.org/10.3835/plantgenome2012.06.0006.

       Ramya, P., A. Chaubal, K. Kulkarni, L. Gupta, N. Kadoo, H. S. Dhaliwal, P. Chhuneja, M.
885   Lagu, and V. Gupt. 2010. "QTL Mapping of 1000-Kernel Weight, Kernel Length,
           and Kernel Width in Bread Wheat (Triticum Aestivum L.)." *Journal of Applied
           Genetics* 51 (4): 421–29. https://doi.org/10.1007/BF03208872.

       Rasheed, Awais, Xianchun Xia, Francis Ogbonnaya, Tariq Mahmood, Zongwen Zhang,
           Abdul Mujeeb-Kazi, and Zhonghu He. 2014. "Genome-Wide Association for Grain

890   Morphology in Synthetic Hexaploid Wheats Using Digital Imaging Analysis." *BMC Plant Biology* 14 (1): 128. https://doi.org/10.1186/1471-2229-14-128.

Ray, Deepak K., Nathaniel D. Mueller, Paul C. West, and Jonathan A. Foley. 2013. "Yield Trends Are Insufficient to Double Global Crop Production by 2050." *PLoS ONE* 8 (6): e66428. https://doi.org/10.1371/journal.pone.0066428.

895   Revelle, William. 2019. *Psych: Procedures for Psychological, Psychometric, and Personality Research*. Evanston, Illinois: Northwestern University. https://CRAN.R-project.org/package=psych.

Reynolds, Matthew, Scott Chapman, Leonardo Crespo-Herrera, Gemma Molero, Suchismita Mondal, Diego N. L. Pequeno, Francisco Pinto, et al. 2020. "Breeder

900   Friendly Phenotyping." *Plant Science*, Food Security under Climate Change, 295 (June): 110396. https://doi.org/10.1016/j.plantsci.2019.110396.

Rife, Trevor W., and Jesse A. Poland. 2014. "Field Book: An Open-Source Application for Field Data Collection on Android." *Crop Science* 54 (4): 1624–27. https://doi.org/10.2135/cropsci2013.08.0579.

905   Rutkoski, Jessica, Jesse Poland, Suchismita Mondal, Enrique Autrique, Lorena González Pérez, José Crossa, Matthew Reynolds, and Ravi Singh. 2016. "Canopy Temperature and Vegetation Indices from High-Throughput Phenotyping Improve Accuracy of Pedigree and Genomic Selection for Grain Yield in Wheat." *G3: Genes|Genomes|Genetics* 6 (9): 2799–2808.

910   https://doi.org/10.1534/g3.116.032888.

Sankaran, Sindhuja, Lav R. Khot, and Arron H. Carter. 2015. "Field-Based Crop Phenotyping: Multispectral Aerial Imaging for Evaluation of Winter Wheat Emergence and Spring Stand." *Computers and Electronics in Agriculture* 118 (October): 372–79. https://doi.org/10.1016/j.compag.2015.09.001.

915   Sankaran, Sindhuja, Lav R. Khot, Carlos Zúñiga Espinoza, Sanaz Jarolmasjed, Vidyasagar R. Sathuvalli, George J. Vandemark, Phillip N. Miklas, et al. 2015. "Low-Altitude, High-Resolution Aerial Imaging Systems for Row and Field Crop Phenotyping: A Review." *European Journal of Agronomy* 70 (October): 112–23. https://doi.org/10.1016/j.eja.2015.07.004.

920   Schwieters, T, and H Saleem. 2013. "Smartphone Uptake Gaining Pace in Africa as IDC Tips Shipments to Double over next Four Years." October 25, 2013. http://itersnews.com/?p=55430.

Sorrells, Mark E., J. Perry Gustafson, Daryl Somers, Shiaoman Chao, David Benscher, Gina Guedira-Brown, Eric Huttner, et al. 2011. "Reconstruction of the Synthetic

925   W7984 × Opata M85 Wheat Reference Population." Edited by A. Van Deynze. *Genome* 54 (11): 875–82. https://doi.org/10.1139/g11-054.

Stacklies, W., H. Redestig, M. Scholz, D. Walther, and J. Selbig. 2007. "PcaMethods a Bioconductor Package Providing PCA Methods for Incomplete Data." *Bioinformatics* 23 (9): 1164–67. https://doi.org/10.1093/bioinformatics/btm069.

930   Subramanian, Ram, Edgar P. Spalding, and Nicola J. Ferrier. 2013. "A High Throughput
        Robot System for Machine Vision Based Plant Phenotype Studies." *Machine
        Vision and Applications* 24 (3): 619–36. https://doi.org/10.1007/s00138-012-
        0434-4.

        Tanabata, Takanari, Taeko Shibaya, Kiyosumi Hori, Kaworu Ebana, and Masahiro Yano.
935   2012. "SmartGrain: High-Throughput Phenotyping Software for Measuring Seed
        Shape through Image Analysis." *Plant Physiology* 160 (4): 1871–80.
        https://doi.org/10.1104/pp.112.205120.

        Tester, M., and P. Langridge. 2010. "Breeding Technologies to Increase Crop Production
        in a Changing World." *Science* 327 (5967): 818–22.
940   https://doi.org/10.1126/science.1183700.

        United Nations, Department of Economic and Social Affairs, and Population Division.
        2019. *World Population Prospects 2019: Highlights*. Vol. (ST/ESA/SER.A/423).

        Wang, Shichen, Debbie Wong, Kerrie Forrest, Alexandra Allen, Shiaoman Chao, Bevan E.
        Huang, Marco Maccaferri, et al. 2014. "Characterization of Polyploid Wheat
945   Genomic Diversity Using a High-Density 90 000 Single Nucleotide Polymorphism
        Array." *Plant Biotechnology Journal* 12 (6): 787–96.
        https://doi.org/10.1111/pbi.12183.

        Wang, Xu, Daljit Singh, Sandeep Marla, Geoffrey Morris, and Jesse Poland. 2018. "Field-
        Based High-Throughput Phenotyping of Plant Height in Sorghum Using Different
950   Sensing Technologies." *Plant Methods* 14 (1): 53.
        https://doi.org/10.1186/s13007-018-0324-5.

        Watson, Amy, Sreya Ghosh, Matthew J. Williams, William S. Cuddy, James Simmonds,
        María-Dolores Rey, M. Asyraf Md Hatta, et al. 2018. "Speed Breeding Is a
        Powerful Tool to Accelerate Crop Research and Breeding." *Nature Plants* 4 (1):
955   23–29. https://doi.org/10.1038/s41477-017-0083-8.

        Whan, Alex P, Alison B Smith, Colin R Cavanagh, Jean-Philippe F Ral, Lindsay M Shaw,
        Crispin A Howitt, and Leanne Bischof. 2014. "GrainScan: A Low Cost, Fast
        Method for Grain Size and Colour Measurements." *Plant Methods* 10 (1): 23.
        https://doi.org/10.1186/1746-4811-10-23.

960   White, Jeffrey W., Pedro Andrade-Sanchez, Michael A. Gore, Kevin F. Bronson, Terry A.
        Coffelt, Matthew M. Conley, Kenneth A. Feldmann, et al. 2012. "Field-Based
        Phenomics for Plant Genetics Research." *Field Crops Research* 133 (July): 101–12.
        https://doi.org/10.1016/j.fcr.2012.04.003.

        Wickham, Hadley. 2016. *Ggplot2: Elegant Graphics for Data Analysis*. New York:
965   Springer-Verlag. https://ggplot2.tidyverse.org.

        Wickham, Hadley, Mara Averick, Jennifer Bryan, Winston Chang, Lucy D'Agostino
        McGowan, Romain François, Garrett Grolemund, et al. 2019. "Welcome to the
        Tidyverse." *Journal of Open Source Software* 4 (43): 1686.
        https://doi.org/10.21105/joss.01686.

970   Wilhelm, Edward P., Ian J. Mackay, Robert J. Saville, Andrey V. Korolev, Francois Balfourier, Andy J. Greenland, Margaret I. Boulton, and Wayne Powell. 2013. "Haplotype Dictionary for the Rht-1 Loci in Wheat." *Theoretical and Applied Genetics* 126 (7): 1733–47. https://doi.org/10.1007/s00122-013-2088-7.

Williams, Keith, and Mark E. Sorrells. 2014. "Three-Dimensional Seed Size and Shape
975   QTL in Hexaploid Wheat (L.) Populations." *Crop Science* 54 (1): 98. https://doi.org/10.2135/cropsci2012.10.0609.

Xu, Yunbi, and Jonathan H. Crouch. 2008. "Marker-Assisted Selection in Plant Breeding: From Publications to Practice." *Crop Science* 48 (2): 391–407. https://doi.org/10.2135/cropsci2007.04.0191.

980   Yang, Wanneng, Lingfeng Duan, Guoxing Chen, Lizhong Xiong, and Qian Liu. 2013. "Plant Phenomics and High-Throughput Phenotyping: Accelerating Rice Functional Genomics Using Multidisciplinary Technologies." *Current Opinion in Plant Biology* 16 (2): 180–87. https://doi.org/10.1016/j.pbi.2013.03.005.

Yu, Jianming, Gael Pressoir, William H Briggs, Irie Vroh Bi, Masanori Yamasaki, John F
985   Doebley, Michael D McMullen, et al. 2006. "A Unified Mixed-Model Method for Association Mapping That Accounts for Multiple Levels of Relatedness." *Nature Genetics* 38 (2): 203–8. https://doi.org/10.1038/ng1702.

Zhang, Zhanhui, Zonghua Liu, Yanmin Hu, Weihua Li, Zhiyuan Fu, Dong Ding, Haochuan Li, Mengmeng Qiao, and Jihua Tang. 2014. "QTL Analysis of Kernel-Related Traits
990   in Maize Using an Immortalized F2 Population." Edited by James C. Nelson. *PLoS ONE* 9 (2): e89645. https://doi.org/10.1371/journal.pone.0089645.