Breeding Program Optimization for Genomic Selection in Winter Wheat

Megan Calvert[a], Byron Evers[b], Xu Wang[b], Allan Fritz[c], Jesse Poland[b]

AFFILIATION

a Interdepartmental Genetics, Kansas State University, Manhattan Kansas 66506

b Department of Plant Pathology, Kansas State University, Manhattan, KS, 66506

c Department of Agronomy, Kansas State University, Manhattan, KS, 66506

KEYWORDS

Triticum aestivum, Plant Breeding, Genomic Prediction, Optimization

Abbreviations

AYN, advanced yield nurseries; BEL, Belleville; BLUPs, Best Linear Unbiased Predictions; GEBVs, genomic estimated breeding values; GNDVI, Green Normalized Difference Vegetation Index; GP, genomic prediction; GRVI, green-red vegetation indices; GRYLD, grain yield; GYP, Gypsum; GxE, genotype x environment; HTP, high-throughput phenotyping; HUTCH, Hutchinson; KSU, Kansas State University; MANH, Manhattan; MP, McPherson; NDRE, normalized difference red edge; NDVI, normalized difference vegetation indices; PTHT, plant height; PC, principal component; PYN, preliminary yield nurseries; RKHS, reproducing kernel Hilbert spaces regression; RR, ridge regression; SNPs, Single nucleotide polymorphisms; UAS, uncrewed aerial system; VI, vegetation indices

Abstract

Developing methodologies in the fields of phenomics and genomic prediction have the potential to increase the production of crop species by accelerating germplasm improvement. The integration of these technologies into germplasm improvement and breeding programs requires evidence that there will be a direct economic benefit to the program. We determined a basic set of parameters, such as prediction accuracy greater than 0.3, the ability to genotype over 7 lines for the cost of one phenotypic evaluation, and heritability levels below 0.4, at which the use of genomic selection would be of economic benefit in terms of genetic gain and operational costs to the Kansas State University (KSU) winter wheat breeding program. The breeding program was then examined to determine whether the parameters benefitting genomic selection were observed or achievable in a practical sense. Our results show that the KSU winter wheat breeding program is at a decision point with regards to their primary means of selection. A few operational changes to increase prediction accuracy would place the program in the parameter space where genomic selection is expected to outpace the current phenotypic selection methodology at a parity of the operation cost and would be of greatest benefit to the program.

Introduction

Technologies are constantly evolving and growing in today's ever-changing world. The introduction of new technologies into an essential service and the development of new crop varieties can be challenging and costly (Moose and Mumm

2008). New technologies must be carefully evaluated to determine if they provide a significant enough advantage to adjust proven current practice. There are many technologies that appeared to have great promise, such as quantitative trait loci (QTL), and marker-assisted selection (MAS), and yet were not beneficial in a practical sense to breeding programs at the time (Bernardo 2008). Yet with a growing global population, erratic environmental conditions, and a finite-amount of arable land, the introduction of new technologies into our crop development systems is essential. The determination of which technologies to implement and the most efficient way to do so is a constant challenge facing plant breeders.

Wheat (*Triticum aestivum*) is one of the top three field crops planted in the USA, behind soybeans and corn, with 1.9 billion bushels produced in 2019 (Bond 2020). Wheat acres and production in the USA have been in a decline since the 1980s as a result of international competition, changing economic conditions and production practices. This is despite the increasing demand for wheat due to growing global populations, which is only expected to increase over the coming century. It is estimated that a 2% increase in yearly grain yields are required to meet these demands (Bassi et al. 2016). Currently there are no reports of wheat breeding programs achieving this level of gain, making wheat an important candidate for breeding technologies designed to accelerate variety development.

Genomic prediction (GP) is one of the new technologies that is showing a great deal of promise to assist in crop-variety production. GP involves the use of genome-wide markers to predict the breeding value of individuals in a population (Meuwissen, Hayes,

and Goddard 2001). This is done by genotyping and phenotyping a training population which is used to establish a model for the trait of interest. This model is then used to calculate genomic estimated breeding values (GEBVs) of a population for which only genotype information is available. GP is already a common technique used in animal breeding due to the benefit of being able to predict a phenotype without having to observe the phenotype, for example the milk yield of a bull's offspring (Hayes et al. 2009; Georges, Charlier, and Hayes 2019).

Plant breeding programs have yet to fully utilize GP for a number of reasons. Breeding programs have to test the same experimental line in multiple locations due to the need to select varieties that are stable across environments. This negates some of the benefit that GP supplies to animal breeding, in which the same genotype cannot be tested under multiple conditions. The genotype x environment (GxE) variation is often high in plant breeding populations due in part to the large weather differences experienced between locations. This significantly decreases the accuracies of most GP models (Dawson et al. 2013; Heslot et al. 2012). Previous GP models did not account for GxE interactions which limited inference and selection decisions in plant breeding. Newer genomic prediction models can now take into account these GxE interactions and are showing greater accuracy in plant breeding situations (Burgueño et al. 2012; Pérez-Rodríguez et al. 2017; Montesinos-López et al. 2016).

The accuracy of GP models has shown some improvement when a covariate is included (Crain et al. 2018; Rutkoski et al. 2016). This has had some success when including high-throughput phenotyping (HTP) data that is taken throughout the course

of the growing season. HTP techniques have the ability to measure thousands of phenotypes accurately in a short period of time. Many of the HTP techniques used in breeding programs take advantage of new developments in remote sensing, uncrewed aerial system (UAS), and sensor technology to measure reflectance and temperature phenotypes. These are all phenotypes that have been shown to be correlated with yield and as such could be effective secondary targets for high-throughput phenotyping and indirect selection (Elliott and Regan 1993; Blackmer et al. 1996; Curran et al. 1983).

In addition to the use of correlated traits, the efficiency of GP is also affected by the stage of the breeding program when it is implemented. Primarily, GP is used to predict the breeding value which is comprised of the additive genetic variation. The greatest advantages of GS will be found when the selection candidates still encompass the most additive genetic variation for the phenotype of interest (Bassi et al. 2016). This is more likely to occur in earlier stages of a breeding program as it is easily selected for, while later stages of a breeding have progressed through strong selection and are more likely to select for other epistatic genetic variation on the performance of the line *per se*.

Another factor in addition to stage of implementation that has hindered the adoption of GP in plant breeding programs are the costs and complex logistics associated with it. The development and establishment of genotyping practices in a plant breeding program requires an investment in infrastructure and training (Moose and Mumm 2008). Along with genetic marker costs, the establishment and maintenance of an adequate training population adds additional costs to the GP protocols. These

costs may not be offset by the gains that the program could potentially achieve using GP (Bassi et al. 2016; Jarquín et al. 2017).

In this study we examined the implementation of GP in a wheat breeding program to specifically examine 1) what parameters/infrastructure is required to implement GP as a primary selection strategy, and 2) are those parameters being met in the KSU public breeding program?

Method and Materials

Cost-Benefit Simulation

A simulated breeding program was used to determine the expected genetic gain when basing selection of material exclusively on observed line performance *per se*, or exclusively on prediction of breeding values. The simulation assumes that a breeding program does not use a combination of phenotypic selection and genomic selection at the same time. In consultation with the Kansas State Hard Winter Wheat Breeding Program, referred to as the KSU breeding program from here on, an appropriate range of cost estimates were determined, as well as sizes of the program as determined by the number of lines developed and evaluated (Supplementary Table 1). For simplicity looking at a single stage of selection in the program, the simulated values assume that the cycle length for each selection scheme is the same and covers the period of the program before the Advanced Yield Nursery (AYN) stage. It also assumes that the number of lines that will be selected for the AYN stage are the same regardless of which selection method is used.

The cost of line developments was estimated to be between $4-$30 per line which covers the initial cross and formation of an inbred line. This can either be done by several years of inbreeding or by the formation of double haploid lines. This was applied as a fixed cost needed to develop the initial population for selection regardless of which selection method was used.

Phenotypic Selection

The costs of phenotypic observation plots are estimated to be between $12-$40 per plot and evaluated within this range at $5 increments. To obtain an accurate phenotypic measurement, replications of the experimental line need to be planted and phenotyped. Depending on the structure of the program this may mean several replications at a single site, or fewer replications at several sites. The maximum number of experimental breeding lines was calculated as:

$$ExperimentalLines = {Budget}\Big/{(LineDevelopment + (PhenoCost * rep))} \quad (1)$$

where $ExperimentalLines$ is the number of lines that would be advanced to the phenotyping stage of selection, the $Budget$ is the total monetary budget for that stage of the breeding program, $LineDevelopment$ is the cost of advancing a single cross to an inbred experimental line for evaluation, $PhenoCost$ is the total cost to obtain the phenotype of interest including labor and other miscellaneous operation costs, and rep is the number of replications of each experimental line that are planted in field and will require phenotyping.

The number of experimental breeding lines was used as the population size when estimating other population parameters.

Genomic Selection

The cost of genotyping a single line for prediction was estimated to be between $1-$10 and evaluated within this range at $1 increments. The maximum number of experimental lines that can be genotyped for prediction was calculated as:

$$GenotypedLines = \frac{Budget}{LineDevelopment + GenoCost} \tag{2}$$

Where $GenotypedLines$ is the number of lines that would be evaluated by genotyping, the $Budget$ is the total monetary budget for that stage of the breeding program, $LineDevelopment$ is the cost of advancing a single cross to an inbred experimental line for evaluation, and $GenoCost$ which is the cost of genotyping a single line including labor and other operational costs. It is assumed that the genotyping is only performed once, and that replication is not required.

The number of possible genotyped lines was then used as the population size when estimating other population parameters.

Simulation Details

The selection methods were compared for every possible combination of number of experimental lines phenotyped and number of experimental lines genotyped, based on the ratio between the correlated response to selection, and the response to selection (Falconer and Mackay 2009).

The expected response to selection for the primary trait was calculated by:

$$R = ih\sigma_A \tag{3}$$

Where i is the intensity of selection, h is the square-root if the narrow-sense heritability of the primary trait, and σ_A is the standard deviation of the additive genetic variance (Falconer and Mackay 2009).

For this study, the GEBVs are assumed to be the secondary trait that is correlated to the primary trait, which would be through experimental observation plots. The correlated response to selection is calculated by:

$$CR = i * \sqrt{h_x^2} * \sqrt{h_y^2} * r_g * \sigma_{py} \tag{4}$$

Where i is the intensity of selection, $\sqrt{h_x^2}$ is the square-root of the narrow-sense heritability of the response trait, $\sqrt{h_y^2}$ is the square-root of the narrow-sense heritability of the secondary trait, r_g is the additive genetic correlation between the traits, and σ_{py} is the standard deviation of the phenotypic variation for the secondary trait (Falconer and Mackay 2009).

A comparison between the indirect response to selection and the expected response to selection is best demonstrated as (Falconer and Mackay 2009):

$$\frac{CR_x}{R_x} = \frac{i_Y r_g \sqrt{h_y^2}}{i_x \sqrt{h_x^2}} \tag{5}$$

Where i_Y is the selection intensity on the secondary trait, r_g is the additive genetic correlation between the primary and secondary trait, $\sqrt{h_y^2}$ is the square-root of the narrow-sense heritability of the secondary trait, i_x is the selection intensity on the direct trait, and $\sqrt{h_x^2}$ is the square-root of the narrow-sense heritability of the response trait.

It was assumed that the individual phenotypes were made up of a genetic portion and an environmental portion. The distribution of each of these portions was assumed to be a random normal with a mean of 0, and a standard deviation of $\sqrt{h^2}$ or $\sqrt{(1 - h^2)}$ for the genotypic and environmental proportions, respectively. A sample the size of the number of lines that were possibly genotyped was taken from this distribution for each individual to give the overall phenotypic distribution. This overall phenotypic distribution was used to determine the intensity of selection (i) in terms of the standard deviation and the selection differential with the msm package (Jackson 2011).

The narrow sense heritabilities for the response trait and the additive genetic correlation were set, with testing the ranges of 0 and 1 at increments of 0.1 each. The narrow-sense heritability of the secondary trait, the genotyping, was assumed to be 0.95. This is under the assumption that the genotypes are inherited almost exactly as they are sequenced and that there are only a few genotyping errors. The ratio between the correlated response and the expected response to selection was plotted against the ratio between number of experimental lines phenotyped and number of lines genotyped.

Plant Material

The KSU Breeding Program breeds hard red winter wheat for a large area which contains different mega-environmental conditions. A subset of 5 locations within the same Kansas mega-environment based on breeder knowledge, Belleville (BEL), Gypsum (GYP), McPherson (MP), Hutchinson (HUTCH) and Manhattan (MANH), were selected for analysis in Kansas between 2016 and 2019. This resulted in 1989 experimental lines being examined over the course of 4 years. These sites contain trials from the

Preliminary Yield Trials (PYN, primary $F_{5:7}$) and the advance yield nursery (AYN, primarily

$F_{5:8}$). The planting, harvest dates and trial size are provided in Table 1. These locations,

excluding Manhattan and Hutchinson, are located in farmers' fields under typical grower

management practices. The PYN and AYN trials were all planted in six-row plots of 1.5m

by 4.5m. The PYN are planted in a modified augmented design with one replicate of the

experimental line per location (Federer and Raghavarao 1975). Plant checks are planted

across whole rows and columns in the trial, and sub-block checks are assigned randomly

within each block. The AYN is made up of lines selected from the PYN trials. The lines are

planted using two replicated α-lattice designs (Patterson and Williams 1976). All 5

locations were planted each year but if a site experienced extreme environmental

variation from the normal climate it was not harvested, providing an unbalanced set of

data.

Phenotyping

Phenotypic information was collected either by combine for grain yield (GRYLD),

by UAS for vegetation indices (VIs), or by hand using the Field Book application (Rife

and Poland 2014) for plant height (PTHT). A DJI Matrice100 (DJI, USA) quadcopter UAS

was equipped with a 5-band multi-spectral RedEdge camera (MicaSense Inc. USA) to

collect plot-level reflectance values for each year, based on the standard protocols

developed by The Wheat Genetics Lab at Kansas State University as laid out in Wang et

al. (2018). The UAS data was collected throughout the course of the growing season,

once every 7-10 days depending on weather conditions. Plant height was collected

manually during the grain ripening stage before harvest. VIs were calculated from the

reflectance values based on the protocols laid out in Wang et al. (2018) and given in

Table 2.

Genotyping

The 1989 lines in the study years were sequenced using genotype-by-sequencing

on an Illumina Hi Seq2000 or Hi Seq2500 (Elshire et al. 2011). Single nucleotide

polymorphisms (SNPs) were called with the Tassel software with the Chinese Spring

wheat assembly v1.0 as a reference (Bradbury et al. 2007; Glaubitz et al. 2014;

International Wheat Genome Sequencing Consortium et al. 2018). The final data set

included 8182 SNPs that were selected for use passed one of three filtering criteria

optimized for the wheat genome by Shrestha et al. (2020) that include Chi-square,

Fisher's test for independence, and the inbreeding coefficient, as well as having a minor

allele frequency greater than 0.05 and missing less than 20% of the data. Missing SNPs

were imputed with Beagle 5.1 (Browning, Zhou, and Browning 2018).

Data Analysis

The Best Linear Unbiased Predictions (BLUPs) for each line were calculated for

each individual year and for multiple years, including or excluding locations as needed.

Where multiple years and locations are included, such as for GRYLD, the BLUPs were

calculated by:

$$y_{ijklt} = \mu + G_i + S_j + M_{l(j)} + GM_{il(j)} + R_{k(lj)} + P_{t(lj)} + \varepsilon_{ijklt} \quad (6)$$

where y_{ijklt} is the phenotypic response variable, μ is the fixed overall mean, G_i is the

random genotype effect for line i distributed as iid $G_i \sim N(0, \sigma_i^2)$, S_j is the random

effect for year j distributed as iid $S_j \sim N(0, \sigma_j^2)$, $M_{l(j)}$ is the random effect for location l within year j distributed as iid $M_{l(j)} \sim N(0, \sigma_l^2)$, $GM_{il(j)}$ is the random genotype by location effect nested within year j distributed as iid $GM_{il(j)} \sim N(0, \sigma_{il}^2)$, $R_{k(lj)}$ is the random effect of replication k within location and year distributed as iid $R_{k(lj)} \sim N(0, \sigma_k^2)$, $P_{t(lj)}$ is the random fungal treatment effect t nested within year-location distributed as iid $P_{t(lj)} \sim N(0, \sigma_t^2)$, and ε_{ijklt} as the residual effect distributed as iid $\varepsilon_{ijklt} \sim N(0, \sigma_{\partial}^2)$.

When only a single year with multiple locations is included the BLUPs are calculated by:

$$y_{ijkl} = \mu + G_i + L_j + GL_{ij} + R_{k(j)} + T_{l(j)} + \varepsilon_{ijkl} \tag{7}$$

where y_{ijkl} is the phenotypic response variable, μ is the fixed overall mean, G_i is the random genotype effect for line i distributed as iid $G_i \sim N(0, \sigma_i^2)$, L_j is the random effect of location j distributed as iid $L_j \sim N(0, \sigma_j^2)$, GL_{ij} is the random effect of genotype by location distributed as iid $GL_{ij} \sim N(0, \sigma_{ij}^2)$, $R_{k(j)}$ is the random effect of replication k nested in location j distributed as iid $R_{k(j)} \sim N(0, \sigma_k^2)$, $T_{l(j)}$ is the random effect of fungal treatment l nested within location j distributed as iid $T_{l(j)} \sim N(0, \sigma_T^2)$, , and ε_{ijkl} is the residual effect distributed as iid $\varepsilon_{ijkl} \sim N(0, \sigma_{\partial}^2)$.

The broad-sense heritability for all years and multiple year-locations was calculated as:

$$H^2 = \frac{\sigma_g^2}{\sigma_g^2 + \frac{\sigma_{gxe}^2}{y*l} + \frac{\sigma_e^2}{y*l*r}} \tag{8}$$

Where σ_g is the total genetic variance, σ_{gxe} is the variance contributed by the

location and variety nested within year, σ_e is the residual environmental variance. As

the data is unbalanced, y is the harmonic mean of the number of years planted, l is the

harmonic mean of the number of locations planted within year, and r is the harmonic

mean of the number of replications per location per year (Holland, Nyquist, and

Cervantes-Martínez 2010).

The broad-sense heritability for individual year AYN and individual year-multiple

location PYN was calculated by:

$$H^2 = \frac{\sigma_g^2}{\sigma_g^2 + \frac{\sigma_{gxe}^2}{l} + \frac{\sigma_e^2}{l*r}} \tag{9}$$

where σ_g^2 is the total genetic variance, σ_{gxe}^2 is the variance contributed by the

genotype-location combination, and σ_e^2 is the residual environmental variance. Similar

to equation 8, l is the harmonic mean of the number of locations planted and r is the

harmonic mean of the number of replications per location (Holland, Nyquist, and

Cervantes-Martínez 2010).

The VI phenotypes were mainly considered on a year-location-trial basis. As such

the BLUPs for the VI phenotypes for the AYN trials are calculated by:

$$y_{irb} = \mu + G_i + M_r + B_{r(b)} + \varepsilon_{irb} \tag{10}$$

where y_{irb} is the phenotypic response variable, μ is the fixed overall mean, G_i is the

random genotypic effect for line i distributed as iid $G_i \sim N(0, \sigma_i^2)$, M_r is the random

effect of the replicate r distributed as iid $M_r \sim N(0, \sigma_r^2)$, $B_{r(b)}$ is the random effect of

the experimental block nested within replicate distributed as iid $B_{r(b)} \sim N(0, \sigma_b^2)$, and

ε_{irb} is the residual effect distributed as iid $\varepsilon_{irb} \sim N(0, \sigma_{\delta}^2)$. The broad-sense heritability was calculated for each year-location's AYN that had VI phenotypes by:

$$H^2 = \frac{\sigma_{\hat{g}}^2}{\sigma_{\hat{g}}^2 + \frac{\sigma_{error}^2}{r}} \tag{11}$$

where σ_g^2 is the genetic variance, σ_{error}^2 is the residual environmental variance and r is the number of replicates.

The BLUPs for the VI phenotypes for the PYN trials are calculated by:

$$y_{ib} = \mu + G_i + B_b + \varepsilon_{ib} \tag{12}$$

where y_{ib} is the phenotypic response variable, μ is the fixed overall mean, G_i is the random genotypic effect N(0,σ^2), B_b is the random effect of the experimental block N(0,σ^2), and ε_{ib} is the residual effect.

Genome-wide Association Analysis

A principal component (PC) analysis of the genotypic information was conducted with the pcaMethods package (Stacklies et al. 2007). A genome-wide association analysis was performed for the VI BLUPs for each year-location-trial using the rrBLUP package (Endelman 2011). The kinship matrix was calculated using rrBLUP was included with 4 PCs to account for kinship and population structure respectively (Endelman and Jannink 2012). A Bonferroni correction was applied with $\alpha = 0.05$ to determine significance.

Genomic Prediction

Genomic prediction was performed with the rrBLUP package and the BGLR statistical package (Pérez and de los Campos 2014) in the R software environment. The rrBLUP package performs ridge regression (RR), and the BGLR package was used to perform a BayesC based prediction and a reproducing kernel Hilbert spaces regression (RKHS). A cross-validation strategy of 100 replications with 80% of the population in the training data set and 20% of the population in the predicted data set was used. The accuracy of the prediction was determined by the correlation between the predicted and observed values. Across year and location predictions were also performed where years, year-trial, and year-location combinations that were not included in the training population are predicted from the remaining data. An example of this would be to use all years excluding the 2016 season as the training data set and the 2016 season as the prediction data set, or all years excluding 2016 season except for the 2016-HUTCH data as the training data set and the remaining three 2016 locations as the prediction data set.

For the VI data only rrBLUP was used for genomic prediction. The VI were used as a cofactor when predicting grain yield for that specific year-location-trial combination as follows:

$$y = \mu + X\beta + Zu + \varepsilon \tag{13}$$

where y is the BLUP for grain yield, μ is the overall mean, X is a ($n\ x\ 1$) matrix of the individual observations of the VI, β is the fixed effects of the VI measurements, Z is an ($n\ x\ m$) matrix assigning markers to genotypes, u is a ($1\ x\ n$) array of the random

effects of the markers and ε is the residual error. The same cross-validation procedure as before was used.

Unless otherwise stated, all analysis took place in the R software environment using the Tidyverse suite of packages (Core Team 2020; Wickham et al. 2019) and visualised with ggplot2 (Wickham 2016). The required code can be found at: https://github.com/megzcalvert/ProgramBreeding

Results

Simulation

To determine the optimal parameters for the operation of the KSU winter wheat breeding program comparing current phenotypic selection methodology verse genomic prediction a simulation was created based on economic decisions. We observed that at low heritabilities of the primary trait, prediction is favored even at low prediction accuracies (Figure 1). Once the ratio between the correlated response and the expected response is greater than 1, then selection on the correlated trait is expected to give greater genetic gain than selection on the primary trait. For the KSU wheat breeding program a primary focus on genotyping is beneficial in terms of genetic gain when the narrow-sense heritability of grain yield is below 0.4, approximately 7.5 lines can be genotyped for every one line that can be phenotyped, and the prediction accuracy of the genomic prediction models is greater than 0.3.

Heritability of traits

Heritability was calculated to identify the phenotypic variance that could be partitioned to the genetic component across the experimental trials. The broad sense heritabilities for each of the seasons ranged from 0.022 for the AYN to 0.441 for the PYN in the 2016 season (Figure 2). The broad-sense heritability for the 2019 PYN cannot be calculated as only one rep was planted at one location (HUTCH).

The VIs show a moderate heritability across the seasons, with the majority being above 0.5 (Figure 3) The heritabilities are variable across the course of the season and locations but the same date-location combination shows similarity across the various VIs, indicating a large influence of ambient conditions of various days within the growing season on the accuracy of VI measurements.

The correlations between grain yield for a location-season-trial combination and the VI's for that location-season-trial combination over the course of the season are given in Figure 4. The correlations ranged between -0.6 in the 2018 PYN at HUTCH for NDRE, to 0.69 in the 2018 AYN at MP for NDRE.

GWAS

To try and determine if any of the VI had genomic regions associated with a large effect size, a GWAS was conducted. As population structure is known to have an effect on GWAS results a PCA of the 1989 experimental lines genotypic data was performed. The PCA shows no distinct population structure and the plot is bounded by the expected founder genotypes, Figure 5. There were no major QTLs for grain yield in any year, yet several of the VIs showed significant associations with regions of the genome. When

these associations were further examined it was shown that they had an influence on the VI value but not GRYLD (Supplementary Figure 1). The regions that showed an association with the VI were often shared between different VI (3/4 in the 2018 season and 3/3 in the 2018 season, Supplementary Figure 1).

Genomic Prediction

Based on the GWAS, GP was evaluated to predict the highly quantitative trait of grain yield. All of the tested genomic prediction models produced similar accuracies (Supplementary Table 2), we therefore focused on rrBLUP models for computational efficiently for further analysis which will be reported. The genomic prediction accuracies based on cross-validation range between 0.311 (SD = 0.079) for the 2018 season and 0.469 (SD = 0.105) for the 2017 season (Figure 6, Supplementary Table 2).

When making forward predictions the strongest correlation, -0.164, was achieved using all seasons excluding the 2017 season as the training population, and the 2017 season as the prediction population (Table 4). When predicting all locations in a single season except for HUTCH, using the data from other seasons and the data from HUTCH as the training population, the greatest accuracy achieved was 0.572 (95% CI [0.503, 0.634]) for 2019 PYN in MANH and BEL. HUTCH was chosen as the location to include in the forward predictions as it is the first location that is normally harvested in Kansas. This would be similar to harvesting one location and then using the predictions to make selections.

To evaluate a prediction approach using secondary traits, the VI were used as covariates in the prediction of grain yield. The prediction of grain yield was not

improved by the addition of a covariate (Figure 7). The prediction of grain yield without a covariate was performed at the same time for comparison and was comparable to the best estimate using VI as covariate.

Discussion

Changing global conditions require current food production systems to be resilient against unexpected events in the face of global population growth. This will require the development of crop varieties that are adapted to hotter and dryer climates. The speed at which these varieties are developed will require the adoption of new technologies that have been proven to show a positive economic investment.

This study examined the validity of using GP and other HTP techniques in the Kansas State University winter wheat breeding program based on several population and model parameters, such as heritability of the primary trait, the prediction accuracy and the selection intensity. These parameters give an indication of whether a new GP method would provide more genetic gain than traditional phenotypic selection equal resource expenditures.

Heritability

The heritability of a trait has a significant impact on whether the trait can be selected for and the possibility for genetic gain of that trait. Traits with a lower heritability are difficult to select for regardless of which selection method is used. However, our simulation showed that traits with a low to moderate narrow-sense heritability (0-0.4) favor the use of GP. For the Kansas State University winter wheat

breeding program, the overall broad-sense heritability of grain yield is below 0.3. As the

narrow-sense heritability is the always less than the broad-sense heritability, the

heritability favors GP under the operation cost parameters for the program.

Prediction Accuracy

The prediction accuracy for GP when making forward predictions in the breeding

program do not meet the criteria to favor GP unless very large populations are used.

This is a possibility in a breeding program setting as historical grain yield data and

genotypes are available for previous seasons. The experimental lines in these seasons

are likely to be less related to the current lines as the parental lines in the crossing block

are updated which may lower prediction accuracy, however, this is the critical

assessment of prediction accuracy that is needed for implementation in the breeding

program as all selections will be focused on new breeding lines into new year's.

HTP Prediction Accuracy

Utilizing the VI to increase the prediction accuracy of the GP models requires

more optimization before it is commonly adopted. The VI need to be able to be used

across locations and years for them to really have an impact on the prediction accuracy

of the GP models. This will require either the manual measurement of growth stages to

standardize measurements to growth stage across locations and years, or the use of

another method such as thermal units to account for differences in growth stage in the

training populations and prediction populations. This additional labor makes the use of

VIs in GP models more costly than utilizing just the GP model. It is an additional factor that needs to be taken into account before the decision to transition to GP is made.

Conclusion

When all parameters are considered it appears that the Kansas State University winter wheat breeding program is on the edge of a large decision. The heritability of grain yield in the breeding program as well as the cost of phenotyping compared to genotyping favor genomic prediction as the way forward for the breeding program. Yet the low accuracy of forward predictions favors the use of phenotypic selection. The forward prediction accuracy can be increased as seen in the 2019 season (Table 4), but this requires a much larger training population. With a few adjustments to the experimental design such as allowing for more replicates of the training population, and changes to the program workflow that allows for the loss of the PYN, genomic prediction could allow the KSU winter wheat breeding program to make larger genetic gains.

Authors' contributions

MC and BE collected data. MC and XW analyzed data. JP conceived and planned the experiments. All authors read and approved the final manuscript.

Acknowledgements

We would like to acknowledge the undergraduate assistants, Quentin Ediger, Kyle Laessig, Jacob Ramsey, Jamie Clark, Molly Smith, Erika Kringen, Nicholas VanPelt,

Bryce Teaford, who assisted with data collection. Daljit Singh, Richard Brown and Grant Williams who assisted with UAS flights and data collection.

MC was supported in part by fellowship funds from the Interdepartmental Genetics program at Kansas State University. This material is based upon work supported by the National Science Foundation under Grant No. 1238187 project "A Field-Based High-Throughput Phenotyping Platform for Plant Genetics" and the USDA NIFA International Wheat Yield Partnership grant no. 2017-67007-25933/project accession no. 1011391 project "Wheat Yield Prediction and Advanced Selection Methodologies through Field-Based High-Throughput Phenotyping with UAVs". Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation or the U.S. Dept. of Agriculture.

References

Bassi, Filippo M., Alison R. Bentley, Gilles Charmet, Rodomiro Ortiz, and Jose Crossa. 2016. "Breeding Schemes for the Implementation of Genomic Selection in Wheat (*Triticum* Spp .)." Plant Science 242 (January): 23–36. https://doi.org/10.1016/j.plantsci.2015.08.021.

Bernardo, Rex. 2008. "Molecular Markers and Selection for Complex Traits in Plants: Learning from the Last 20 Years." Crop Science 48 (5): 1649. https://doi.org/10.2135/cropsci2008.03.0131.

Blackmer, Tracy M., James S. Schepers, Gary E. Varvel, and Elizabeth A. Walter-Shea. 1996. "Nitrogen Deficiency Detection Using Reflected Shortwave Radiation from Irrigated Corn Canopies." Agronomy Journal 88 (1): 1–5. https://doi.org/10.2134/agronj1996.00021962008800010001x.

Bond, Jennifer K. 2020. "USDA ERS - Wheat Sector at a Glance." June 26, 2020. https://www.ers.usda.gov/topics/crops/wheat/wheat-sector-at-a-glance/.

Bradbury, P. J., Z. Zhang, D. E. Kroon, T. M. Casstevens, Y. Ramdoss, and E. S. Buckler. 2007. "TASSEL: Software for Association Mapping of Complex Traits in Diverse

Samples." Bioinformatics 23 (19): 2633–35.
https://doi.org/10.1093/bioinformatics/btm308.

Browning, Brian L., Ying Zhou, and Sharon R. Browning. 2018. "A One-Penny Imputed
Genome from Next-Generation Reference Panels." The American Journal of
Human Genetics 103 (3): 338–48. https://doi.org/10.1016/j.ajhg.2018.07.015.

Burgueño, Juan, Gustavo de los Campos, Kent Weigel, and José Crossa. 2012. "Genomic
Prediction of Breeding Values When Modeling Genotype × Environment
Interaction Using Pedigree and Dense Molecular Markers." Crop Science 52 (2):
707–19. https://doi.org/10.2135/cropsci2011.06.0299.

Core Team, R. 2020. R: A Language and Environment for Statistical Computing (version
4.0.0). Vienna, Austria: R Foundation for Statistical Computing. https://www.R-
project.org/.

Crain, Jared, Suchismita Mondal, Jessica Rutkoski, Ravi P. Singh, and Jesse Poland. 2018.
"Combining High-Throughput Phenotyping and Genomic Information to Increase
Prediction and Selection Accuracy in Wheat Breeding." The Plant Genome 11 (1):
0. https://doi.org/10.3835/plantgenome2017.05.0043.

Curran, P. J., M. D. Steven, John Theodore Houghton, Arthur Herbert Cook, and Henry
Charnock. 1983. "Multispectral Remote Sensing for the Estimation of Green Leaf
Area Index." Philosophical Transactions of the Royal Society of London. Series A,
Mathematical and Physical Sciences 309 (1508): 257–70.
https://doi.org/10.1098/rsta.1983.0039.

Dawson, Julie C., Jeffrey B. Endelman, Nicolas Heslot, Jose Crossa, Jesse Poland,
Susanne Dreisigacker, Yann Manès, Mark E. Sorrells, and Jean-Luc Jannink. 2013.
"The Use of Unbalanced Historical Data for Genomic Selection in an International
Wheat Breeding Program." Field Crops Research 154 (December): 12–22.
https://doi.org/10.1016/j.fcr.2013.07.020.

Elliott, G. A., and K. L. Regan. 1993. "Use of Reflectance Measurements to Estimate
Early Cereal Biomass Production on Sandplain Soils." Australian Journal of
Experimental Agriculture 33 (2): 179–83. https://doi.org/10.1071/ea9930179.

Endelman, Jeffrey B. 2011. "Ridge Regression and Other Kernels for Genomic Selection
with R Package RrBLUP." The Plant Genome Journal 4 (3): 250.
https://doi.org/10.3835/plantgenome2011.08.0024.

Falconer, Douglas S., and Trudy F. C. Mackay. 2009. Introduction to Quantitative
Genetics. 4. ed., [16. print.]. Harlow: Pearson, Prentice Hall.

Federer, W. T., and D. Raghavarao. 1975. "On Augmented Designs." Biometrics 31 (1):
29–35. https://doi.org/10.2307/2529707.

Georges, Michel, Carole Charlier, and Ben Hayes. 2019. "Harnessing Genomic
Information for Livestock Improvement." Nature Reviews Genetics 20 (3): 135–
56. https://doi.org/10.1038/s41576-018-0082-2.

Glaubitz, Jeffrey C., Terry M. Casstevens, Fei Lu, James Harriman, Robert J. Elshire, Qi Sun, and Edward S. Buckler. 2014. "TASSEL-GBS: A High Capacity Genotyping by Sequencing Analysis Pipeline." Edited by Nicholas A. Tinker. PLoS ONE 9 (2): e90346. https://doi.org/10.1371/journal.pone.0090346.

Hayes, B. J., P. J. Bowman, A. J. Chamberlain, and M. E. Goddard. 2009. "Invited Review: Genomic Selection in Dairy Cattle: Progress and Challenges." Journal of Dairy Science 92 (2): 433–43. https://doi.org/10.3168/jds.2008-1646.

Heslot, Nicolas, Hsiao-Pei Yang, Mark E. Sorrells, and Jean-Luc Jannink. 2012. "Genomic Selection in Plant Breeding: A Comparison of Models." Crop Science 52 (1): 146–60. https://doi.org/10.2135/cropsci2011.06.0297.

Holland, James B., Wyman E. Nyquist, and Cuauhtemoc T. Cervantes-Martínez. 2010. "Estimating and Interpreting Heritability for Plant Breeding: An Update." In Plant Breeding Reviews, 9–112. John Wiley & Sons, Ltd. https://doi.org/10.1002/9780470650202.ch2.

International Wheat Genome Sequencing Consortium, Michael Alaux, Jane Rogers, Thomas Letellier, Raphaël Flores, Françoise Alfama, Cyril Pommier, et al. 2018. "Linking the International Wheat Genome Sequencing Consortium Bread Wheat Reference Genome Sequence to Wheat Genetic and Phenomic Data." Genome Biology 19 (1): 111. https://doi.org/10.1186/s13059-018-1491-4.

Jackson, Christopher. 2011. "Multi-State Models for Panel Data: The Msm Package for R." Journal of Statistical Software 38 (1): 1–28. https://doi.org/10.18637/jss.v038.i08.

Jarquín, Diego, Cristiano Lemes da Silva, R. Chris Gaynor, Jesse Poland, Allan Fritz, Reka Howard, Sarah Battenfield, and Jose Crossa. 2017. "Increasing Genomic-Enabled Prediction Accuracy by Modeling Genotype × Environment Interactions in Kansas Wheat." The Plant Genome 10 (2): plantgenome2016.12.0130. https://doi.org/10.3835/plantgenome2016.12.0130.

Meuwissen, T. H. E., B. J. Hayes, and M. E. Goddard. 2001. "Prediction of Total Genetic Value Using Genome-Wide Dense Marker Maps." Genetics 157 (4): 1819–29.

Montesinos-López, Osval A., Abelardo Montesinos-López, José Crossa, Fernando H. Toledo, Oscar Pérez-Hernández, Kent M. Eskridge, and Jessica Rutkoski. 2016. "A Genomic Bayesian Multi-Trait and Multi-Environment Model." G3: Genes|Genomes|Genetics 6 (9): 2725–44. https://doi.org/10.1534/g3.116.032359.

Moose, Stephen P., and Rita H. Mumm. 2008. "Molecular Plant Breeding as the Foundation for 21st Century Crop Improvement." Plant Physiology 147 (3): 969–77. https://doi.org/10.1104/pp.108.118232.

Patterson, H. D., and E. R. Williams. 1976. "A New Class of Resolvable Incomplete Block Designs." Biometrika 63 (1): 83–92. https://doi.org/10.2307/2335087.

Pérez, Paulino, and Gustavo de los Campos. 2014. "Genome-Wide Regression and Prediction with the BGLR Statistical Package." Genetics 198 (2): 483–95. https://doi.org/10.1534/genetics.114.164442.

Pérez-Rodríguez, Paulino, José Crossa, Jessica Rutkoski, Jesse Poland, Ravi Singh, Andrés Legarra, Enrique Autrique, Gustavo de los Campos, Juan Burgueño, and Susanne Dreisigacker. 2017. "Single-Step Genomic and Pedigree Genotype × Environment Interaction Models for Predicting Wheat Lines in International Environments." The Plant Genome 10 (2): 0. https://doi.org/10.3835/plantgenome2016.09.0089.

Poland, Jesse A., Patrick J. Brown, Mark E. Sorrells, and Jean-Luc Jannink. 2012. "Development of High-Density Genetic Maps for Barley and Wheat Using a Novel Two-Enzyme Genotyping-by-Sequencing Approach." Edited by Tongming Yin. PLoS ONE 7 (2): e32253. https://doi.org/10.1371/journal.pone.0032253.

Rife, Trevor W., and Jesse A. Poland. 2014. "Field Book: An Open-Source Application for Field Data Collection on Android." Crop Science 54 (4): 1624–27. https://doi.org/10.2135/cropsci2013.08.0579.

Rutkoski, Jessica, Jesse Poland, Suchismita Mondal, Enrique Autrique, Lorena González Pérez, José Crossa, Matthew Reynolds, and Ravi Singh. 2016. "Canopy Temperature and Vegetation Indices from High-Throughput Phenotyping Improve Accuracy of Pedigree and Genomic Selection for Grain Yield in Wheat." G3: Genes|Genomes|Genetics 6 (9): 2799–2808. https://doi.org/10.1534/g3.116.032888.

Stacklies, W., H. Redestig, M. Scholz, D. Walther, and J. Selbig. 2007. "PcaMethods a Bioconductor Package Providing PCA Methods for Incomplete Data." Bioinformatics 23 (9): 1164–67. https://doi.org/10.1093/bioinformatics/btm069.

Wang, Xu, Daljit Singh, Sandeep Marla, Geoffrey Morris, and Jesse Poland. 2018. "Field-Based High-Throughput Phenotyping of Plant Height in Sorghum Using Different Sensing Technologies." Plant Methods 14 (1): 53. https://doi.org/10.1186/s13007-018-0324-5.

Wickham, Hadley, Mara Averick, Jennifer Bryan, Winston Chang, Lucy D'Agostino McGowan, Romain François, Garrett Grolemund, et al. 2019. "Welcome to the Tidyverse." Journal of Open Source Software 4 (43): 1686. https://doi.org/10.21105/joss.01686.

Figures

Figure 0.1. Simulation results comparing phenotypic selection as a direct trait to selection on the genotype as a secondary trait.

The narrow-sense heritability of the direct trait is in the panel title and the y-axis is the Correlated Response / Response of selection for the direct trait. When this ratio is above 1, represented by the dotted line, selection on the secondary trait is favored. The x-axis is the number of experimental lines that can be genotyped and predicted for every line that is phenotyped based on estimated costs. The trend of each prediction accuracy is given by the color of the line.

Figure 0.2. Broad sense heritabilities for grain yield across 4 years of phenotypic data.

The broad sense heritabilities are presented by season on the x-axis with each trial type denoted by the character shape. There is no PYN for the 2019 season as it was only planted in one location with one replication.

Figure 0.3. The broad sense heritabilities for the VIs for the AYN trials planted at each location.

The title of each plot gives the season-VI combination, with the date on which the VI was taken given by the x-axis, the y-axis is the broad sense heritability, while the color of the point determines the location. NDVI = Normalized Difference Vegetation Index, GNDVI = Green Normalized Difference Vegetation Index, and NDRE = Red Edge Normalized Difference Vegetation Index.

Figure 0.4. The correlations between grain yield and VI by year-trial combinations.

The title of each plot gives the season-VI combination, with the date on which the VI was taken given by the x-axis, the y-axis is the Pearson correlation coefficient, while the color of the point determines the location. NDVI = Normalized Difference Vegetation Index, GNDVI = Green Normalized Difference Vegetation Index, and NDRE = Red Edge Normalized Difference Vegetation Index.

Figure 0.5. Population structure based on genotypic information in the Kansas State winter wheat breeding program nurseries between the 2016-2019 seasons.

The top panel shows the first two principal components (PC), while the bottom panel displays the 2nd and 3rd PC. The variance contributed by each PC is given next to the PC name. Several "founder" lines are highlighted.

Figure 0.6. Genomic prediction accuracies for grain yield determined by cross-validation.

The season or seasons used in each analysis are given on the x-axis. The y-axis shows the range of possible correlations between the predicted phenotype and the observed phenotype. The color of the point determines if the individual line had been in the training or testing population of the analysis. 100 cross-validations were performed in each analysis. The modelTraining results give an indication of the model-fit whereas the modelTesting results give an indication of the predictive ability of the model. The AYN and PYN trials are included in the analysis.

Figure 0.7. Genomic prediction accuracies for PYN grain yield when a VI is used as a covariate determined by cross-validation.

The season and VI are given in the strip title and the points are colored by the location. The GRYLD measurements are those for the prediction of grain yield without a covariate. The large point is the mean and the error bars give the standard deviation. NDVI = Normalized Difference Vegetation Index, GNDVI = Green Normalized Difference Vegetation Index, and NDRE = Red Edge Normalized Difference Vegetation Index.

Tables

Table 0.1. Summary of trials planted between the 2016 and 2019 seasons across locations. A summary is given by season, trial and location with the number of individual plots, experimental lines, date of planting, date of harvest, the mean grain yield in t/ha and the standard deviation of the grain yield.

Season	Trial	Location	Plots	Lines	Planted	Harvested	Mean	SD
2015/2016	AYN	MP	40	75			5.050	0.9
		MANH	240	75			3.206	0.651
		HUTCH	320	109			4.139	0.92
		GYP	40	109			5.739	0.726
	PYN	MP	504	439			5.113	1.21
		MANH	504	379			3.720	0.657
		HUTCH	594	516			3.770	0.886
		GYP	593	506			5.421	0.702
2016/2017	AYN	MP	315	91	10/11/16	20/06/17	5.933	1.00
		MANH	320	91	10/18/16	22/06/17	5.822	0.768
		HUTCH	168	97	10/12/16	21/06/17	6.242	0.845
		BEL	48	36	10/19/16	28/06/17	5.110	0.776
	PYN	MP	126	120	10/11/16	20/06/17	4.805	1.015
		HUTCH	108	102	10/12/16	21/06/17	6.319	0.925
		BEL	8	8	10/19/16	28/06/17	4.5	0.385
2017/2018	AYN	MP	199	90	10/19/17	29/06/18	3.203	0.412
		MANH	200	90	10/20/17	23/06/18	2.644	0.402
		HUTCH	280	125	10/18/17	28/06/18	3.345	0.489
		BEL	280	125	10/17/17	30/06/18	2.561	0.430
		GYP	279	125	10/18/17	18/06/18	2.343	0.318
	PYN	MP	323	282	10/19/17	29/06/18	3.048	0.490

		MANH	414	366	10/20/17	23/06/18	2.880	0.431
		HUTCH	414	366	10/18/17	28/06/18	3.236	0435
		BEL	287	251	10/17/17	30/06/18	2.198	0.412
		GYP	286	250	10/18/17	18/06/18	1.973	0.278
2018/2019	AYN	MANH	80	40	11/01/18		5.798	0.535
		HUTCH	212	99	10/24/18	01/07/19	4.925	0.563
		BEL	130	63	10/24/18	17/07/19	4.965	0.897
	PYN	HUTCH	504	454	10/24/18	01/07/19	4.623	0.794

Table 0.2. Summary of vegetation indices used in the KSU breeding program across 5 years. For each index the formula and reference are provided.

Abbr.	Index	Equation	Reference
GNDVI	Green Normalized Difference Vegetation Index	$GNDVI = \dfrac{NIR - Green}{NIR + Green}$	(Blackmer et al. 1996)
NDRE	Red Edge Normalized Difference Vegetation Index	$NDRE = \dfrac{NIR - RedEdge}{NIR + RedEdge}$	(Elliott and Regan 1993)
NDVI	Normalized Difference Vegetation Index	$NDVI = \dfrac{NIR - Red}{NIR + Red}$	(Curran et al. 1983)

Table 0.3. Genomic Prediction Accuracies as the correlation between BLUPs and GEBVs. The Pearson correlation coefficients is given with a 95% CI and the rank correlation between the GEBVs and the BLUPs.

Training Population	Testing Population	Correlation	95% CI	Rank Correlation
Excluding 2015/2016	2015/2016	0.09	[-0.004, 0.182]	0.093
Excluding 2015/2016 PYN	2015/2016 PYN	0.078	[-0.026, 0.180]	0.093
Excluding 2015/2016 MP MANH GYP	2015/2016 MP MANH GYP	0.024	[-0.08, 0.127]	0.019
Excluding 2016/2017	2016/2017	-0.164	[-0.336, 0.018]	-0.125
Excluding 2016/2017 PYN	2016/2017 PYN	-0.055	[-0.249, 0.144]	0.010
Excluding 2016/2017 MP MANH BEL	2016/2017 MP MANH BEL	-0.009	[-0.206, 0.189]	0.059
Excluding 2017/2018	2017/2018	0.048	[-0.049, 0.145]	0.030
Excluding 2017/2018 PYN	2017/2018 PYN	0.061	[-0.046, 0.165]	0.019
Excluding 2017/2018 MP MANH BEL	2017/2018 MP MANH BEL	0.009	[-0.097, 0.114]	-0.026
Excluding 2018/2019	2018/2019	-0.091	[-0.186, 0.005]	-0.13
Excluding 2018/2019 PYN	2018/2019 PYN	-0.039	[-0.135, 0.058]	-0.079
Excluding 2018/2019 MANH BEL	2018/2019 MANH BEL	0.572	[0.502, 0.634]	0.537

Supplementary Material

Figure C.1. Results of GWAS Analysis.

The effect of the significant SNPs identified in the GWAS analysis for the specific phenotype and grain yield are given in the boxplot below the GWAS results. The points are colored by chromosome.

GWAS ayn18 NDRE_2018.05.17_HUTCH

Figure C.1. Results of GWAS Analysis Continued.

The effect of the significant SNPs identified in the GWAS analysis for the specific phenotype and grain yield are given in the boxplot below the GWAS results. The points are colored by chromosome.

GWAS ayn18 NDVI_2018.05.17_HUTCH

GWAS ayn18 NDVI_2018.06.06_MP

Figure C.1. Results of GWAS Analysis Continued.

The effect of the significant SNPs identified in the GWAS analysis for the specific phenotype and grain yield are given in the boxplot below the GWAS results. The points are colored by chromosome.

Figure C.1. Results of GWAS Analysis Continued.

The effect of the significant SNPs identified in the GWAS analysis for the specific phenotype and grain yield are given in the boxplot below the GWAS results. The points are colored by chromosome.

Figure C.1. Results of GWAS Analysis Continued.

The effect of the significant SNPs identified in the GWAS analysis for the specific phenotype and grain yield are given in the boxplot below the GWAS results. The points are colored by chromosome.

Table C.1. Summary of simulation costs. The range over which the parameter was tested as well as the step change for each are given.

Parameter	Range	Step
Line Development	$4-40	$1
Genotyping Costs	$1-20	$1
Phenotyping Costs	$35-70	$5
Phenotyping Replications	6	N/A
Heritability of Primary Trait	0-1	0.1
Heritability of Correlated Trait	0.95	N/A
Prediction Accuracy	0-1	0.1
AYN Selected	100	N/A

Table C.2. Accuracy of different GP models. The average correlation between the observed and GEBVs for the different genomic prediction models for 100 CV's is given with the standard deviation in brackets.

Season	Ridge Regression	Reproducing Kernel Hilbert Space	BayesC
All	0.3274 (0.0494)	0.3279 (0.0492)	0.3274 (0.0496)
2015/2016	0.4505 (0.0700)	0.4495 (0.0700)	0.4500 (0.0703)
2016/2017	0.4834 (0.1075)	0.4843 (0.1075)	0.4840 (0.1067)
2017/2018	0.2963 (0.0702)	0.2962 (0.0704)	0.2971 (0.0702)
2018/2019	0.3700 (0.0802)	0.3704 (0.0793)	0.3704 (0.0801)
Excluding_2016	0.3197 (0.0516)	0.3201 (0.0518)	0.3201 (0.0526)
Excluding_2017	0.3342 (0.0551)	0.3345 (0.0551)	0.3341 (0.0555)
Excluding_2018	0.3619 (0.0533)	0.3634 (0.0539)	0.3623 (0.0534)
Excluding_2019	0.3682 (0.0610)	0.3683 (0.0611)	0.3678 (0.0612)