A fluorescent sensor for spatiotemporally resolved endocannabinoid dynamics *in*
vitro* and *in vivo

Ao Dong[1,2,3], Kaikai He[1,2], Barna Dudok[4], Jordan S Farrell[4], Wuqiang Guan[5], Daniel J
Liput[6,7], Henry L Puhl[7], Ruyi Cai[1,2], Jiali Duan[1,2], Eddy Albarran[8], Jun Ding[9], David M
Lovinger[6], Bo Li[5], Ivan Soltesz[4], & Yulong Li[1,2,3,10,*]

[1]State Key Laboratory of Membrane Biology, Peking University School of Life Sciences,
Beijing, China;
[2]PKU-IDG/McGovern Institute for Brain Research, Beijing, China;
[3]Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary
Studies, Peking University, Beijing, China;
[4]Department of Neurosurgery, Stanford University, Palo Alto, CA, USA;
[5]Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, USA;
[6]Laboratory for Integrative Neuroscience, National Institute on Alcohol Abuse and
Alcoholism, Bethesda, MD, USA;
[7]Laboratory of Molecular Physiology, National Institute on Alcohol Abuse and Alcoholism,
Bethesda, MD, USA;
[8]Neuroscience PhD Program, Stanford University, Palo Alto, CA, USA;
[9]Department of Neurosurgery, Department of Neurology, Stanford University School of
Medicine, Palo Alto, CA, USA;
[10]Chinese Institute for Brain Research, Beijing, China.

*Correspondence should be addressed to Y.L. (yulongli@pku.edu.cn).

Endocannabinoids (eCBs) are retrograde lipid neuromodulators involved in many
physiologically important processes. However, their release and dynamics in the
brain remain largely unknown, in part due to lack of probes capable of reporting
real-time eCBs with sufficient spatiotemporal resolution. Here, we developed a new
G protein-coupled receptor Activation Based eCB sensor GRAB$_{eCB2.0}$ using the
human CB1 cannabinoid receptor and a circular-permutated EGFP. GRAB$_{eCB2.0}$
exhibited proper cell membrane trafficking, ~seconds kinetics, high specificity and
robust fluorescence response to eCBs at physiological concentrations. Using
GRAB$_{eCB2.0}$, we detected evoked eCB dynamics in both cultured neurons and acute
brain slices. Interestingly, we also observed spontaneous compartmental eCB
transients that spread ~11 μm in cultured neurons, suggesting locally-restricted eCB
signaling. By expressing GRAB$_{eCB2.0}$ *in vivo*, we readily observed foot-shock elicited
and running triggered eCB transients in mouse amygdala and hippocampus,
respectively. Lastly, using GRAB$_{eCB2.0}$ in an epilepsy model, we observed a
spreading eCB wave following a calcium wave in mouse hippocampus. In summary,
GRAB$_{eCB2.0}$ is a powerful new probe to resolve eCB release and dynamics under both
physiological and pathological conditions.

Cannabis derivatives have long been used for medicinal and recreational purposes across

cultures in formulations such as marijuana and hashish[1]. Bioactive compounds in cannabis,
phytocannabinoids, exert their function by hijacking the endocannabinoid system in our
body. The biological function of endocannabinoids (eCBs), majorly two lipid metabolites 2-
arachidonoylglycerol (2-AG) and anandamide (AEA), is primarily mediated by the
activation of two G protein-coupled receptors (GPCRs), cannabinoid receptor type 1
(CB1R) and type 2 (CB2R)[2]. As important neuromodulators, eCBs are widely distributed in
the peripheral and central nervous system. Interestingly, distinct from other classical
neurotransmitters that are stored in synaptic vesicles and released from presynaptic
terminals, eCBs are typically produced and released from the postsynaptic site in a
neuronal activity-dependent manner, then retrogradely travel to the presynaptic terminal
and activate the CB1R, activation of which often results in an inhibition of presynaptic
neurotransmitter release[3,4]. In addition, eCBs can also function in glial cells and
intracellular organelles[5-9]. eCBs participate in the short-term and long-term synaptic
plasticity of glutamatergic and gamma-aminobutyric acid (GABA)-ergic synapses in a
variety of brain regions such as cerebral cortex, hippocampus, striatum, ventral tegmental
area, amygdala and cerebellum[4,10], and play important roles in many physiological
processes including development, emotional state, pain, sleep/wake cycle, energy
metabolism, reward, learning and memory[11-15]. Given the broad distribution and functions
of eCBs, dysregulation of eCB system is tightly linked with a variety of disorders, including
neuropsychiatric and neurodegenerative diseases, epilepsy, cancer, and others[16-18].
Therefore, the eCB system has become a therapeutic target for treating multiple
neurological diseases[19,20].

In contrast to the increasing knowledge about the eCB biochemistry and physiology,
the spatiotemporal dynamics of eCBs in the brain remains largely unknown. Synaptic
transmission mediated by classic neurotransmitters such as glutamate and GABA and their
ionotropic receptors can happen in milliseconds timescale and is confined to the
nanometer dimensions of the synaptic cleft[21], while neuropeptides are thought to be
secreted in seconds or minutes after stimulation and act at longer distances[22]. However,
our understanding of eCB signaling is limited due to drawbacks of existing eCB detection
methods. For example, qualitative and quantitative measurement of eCBs in brain tissues
can provide valuable information on eCB levels. However, this usually relies on lipid
extraction, purification and analysis by chromatography and mass spectrometry[23,24],
therefore, has poor spatial and temporal resolution and cannot detect eCBs *in vivo*.
Electrophysiology, together with genetics and pharmacology, is frequently used to provide
indirect measurement of eCB action by studying eCB-mediated synaptic modulation[25-28].
However, this method is mostly used in *in vitro* preparations with reduced physiological
relevance and has poor spatial resolution. Microdialysis, while challenging for hydrophobic
lipid molecules, has been used to monitor eCB abundance in the brain during
pharmacological manipulations and behaviors[29,30], but it has a long sampling interval
(~minutes) that is well beyond the time scale of synaptic plasticity mediated by eCBs (~sub-
second to seconds), preventing the accurate detection of eCBs in real time *in vivo*.
Therefore, development of an *in vivo* eCB detection tool with satisfactory spatiotemporal
resolution would meet a clear need in this field.

Recently, several GPCR- and circular-permutated (cp) fluorescent protein-based

sensors for neurotransmitters and neuromodulators were successfully developed[31-39].
Following this strategy, here we report a novel GPCR Activation Based eCB sensor
GRAB$_{eCB2.0}$ (eCB2.0 for simplicity) based on the human CB1R and cpEGFP. eCB2.0
exhibits proper membrane trafficking, high specificity, ~second-scale kinetics, ~800%
fluorescence response to 2-AG and ~550% to AEA in cultured neurons. After validating the
performance of eCB2.0 in cultured cells and acute brain slices, we are able to detected
foot-shock evoked eCB signals in the amygdala in freely moving mice and eCB dynamics
in the mouse hippocampus during running and seizures.
97
RESULTS
Development and *in vitro* characterization of GRAB$_{eCB}$ sensors
Among the two cannabinoid receptors, we chose CB1R as the eCB sensor scaffold,
because it has a higher affinity towards eCBs than CB2R[40]. To start, we inserted the
intracellular loop 3 (ICL3)-cpEGFP module from our recently developed GRAB$_{NE}$ sensor[34]
into the ICL3 of human CB1R (**Fig. 1a**). After several rounds of screening for insertion site
and GRAB$_{NE}$ ICL3 truncation, we generated the first generation eCB sensor named
GRAB$_{eCB1.0}$ (eCB1.0 for simplicity), which showed a moderate response (ΔF/F$_0$ ~100%)
and 3 μM apparent affinity for 2-AG (**Fig. 1b and Extended Data Fig. 1a**). To further
improve the dynamic range of the eCB sensor, we selected 8 sites in cpEGFP for individual
random mutation based our the experience gained through the development of previous
GRAB sensors[31,33-35,37-39] (**Extended Data Fig. 1b**). A combination of mutations from
single-mutation candidates with improved performance resulted in GRAB$_{eCB1.5}$ (eCB1.5),
which showed a higher response (ΔF/F$_0$ ~200%) than eCB1.0 (**Extended Data Fig. 1a**).
We next focused on the CB1R ligand binding pocket, aiming to further improve the
response and affinity of the sensor. Residues F177$^{2.64}$, V196$^{3.32}$ and S383$^{7.39}$ were selected
for targeted screening based on studies of CB1R structure[41-46] (**Extended Data Fig. 1c**).
Interestingly, we found that eCB1.5 S383$^{7.39}$T showed an increased response and a similar
apparent affinity to 2-AG compared to eCB1.5, while eCB1.5 S383$^{7.39}$T F177$^{2.64}$A showed
no response to 2-AG (**Extended Data Fig. 1a**). We thus named eCB1.5 S383$^{7.39}$T as
GRAB$_{eCB2.0}$ (eCB2.0), and eCB1.5 S383$^{7.39}$T F177$^{2.64}$A as GRAB$_{eCBmut}$ (eCBmut)
(**Extended Data Fig. 2**), which is an eCB non-binding mutant sensor to be used as a
negative control.
Both eCB2.0 and eCBmut exhibited proper cell membrane trafficking when expressed
in HEK293T cells (**Fig. 1c**). Upon application of ligands, eCB2.0 showed concentration-
dependent fluorescence increases to both 2-AG and AEA, with the maximum ΔF/F$_0$
~210%, and half maximal effective concentrations (EC$_{50}$) of ~7.2 μM for 2-AG and ~0.5 μM
for AEA. On the contrary, eCBmut showed no response to 2-AG or AEA at all
concentrations tested (**Fig. 1d**). We then tested whether eCB2.0 signal was specific to
eCBs but not other neurotransmitters. eCB2.0 responses to 10 μM 2-AG and AEA were
completely blocked by the CB1R inverse agonist AM251 (10 μM), and eCB2.0 showed no
response to other common neurotransmitters or neuromodulators (**Fig. 1e**). Next, we
measured the kinetics of eCB2.0 by local micropressure (puff) application (**Fig. 1f**).
HEK293T cells expressing eCB2.0 were exposed to puffs of 100 μM 2-AG to measure the
"on" kinetics, and cells expressing eCB2.0 and incubated in 10 μM 2-AG were exposed to

puffs of 100 μM AM251 to measure the "off" kinetics at room temperature. Time constants
of on and off kinetics were ~1.6 s and ~11.2 s, respectively (**Fig. 1g**). To examine whether
or not there is any potential downstream coupling of eCB sensors with intracellular
signaling pathways, we first measured G-protein activation using a $G_{\beta\gamma}$ BRET sensor based
on the $G_{\beta\gamma}$ binding region of phosducin fused to NanoLuc luciferase. This unified BRET
sensor was based upon similar systems[47,48]. 20 μM 2-AG significantly increased the BRET
signal in wild-type CB1R expressing cells, indicating that G-protein signaling was activated,
but not in cells expressing eCB2.0, eCBmut or in blank control cells (**Fig. 1h**). We then
measured β-arrestin recruitment using the Tango assay[49]. We found that AEA potently
recruited β-arrestin in CB1R expressing cells but not in either eCB2.0, eCBmut or blank
control cells (**Fig. 1i**). These data together demonstrate that eCB2.0 and eCBmut have no
detectable coupling with the two main GPCR downstream effectors, implying that the
expression and activation of eCB sensors themselves may have minimal perturbation to
cell physiology.
To examine the performance of eCB sensors in neurons, we first sparsely expressed
eCB2.0 in cultured rat cortical neurons. We found eCB2.0 targeted to the cell membrane
throughout the neuron, including axons and dendrites, as indicated by the colocalization of
eCB2.0 with the axonal presynaptic marker synaptophysin-mScarlet and the postsynaptic
marker PSD95-mScarlet, respectively (**Fig. 2a**). To measure the response of eCB sensors,
we infected cultured rat cortical neurons by adeno-associated viruses (AAVs) carrying
eCB2.0 and eCBmut under the control of the human synapsin promoter, which enables
efficient labeling of all neurons in the culture. eCB2.0 showed fluorescence responses to
bath-applied 2-AG and AEA in a concentration-dependent manner. In contrast, eCBmut
showed no response to 2-AG or AEA (**Fig. 2b,c**). The maximum responses of eCB2.0 to
2-AG and AEA were ~800% and 550%, and EC_{50}s were ~17.2 μM and ~0.7 μM,
respectively. eCB2.0 responses in neurites were higher than in somata (**Fig. 2d**). Bath
application of the CB1R agonist WIN55212-2, which can activate eCB2.0 (**Extended Data**
Fig. 3a), induced a fluorescence increase of eCB2.0 on neuronal membrane that was
stable for as long as 2 hours (**Fig. 2e**). These results indicate minimal arrestin-mediated
internalization or desensitization of eCB2.0 sensor in neurons, suggesting the utility of the
sensor for long-term eCB imaging.
164
Imaging endogenous eCBs in primary cultured neurons using eCB2.0
Cultured neurons are commonly used for studying eCB mediated synaptic modulation[28,50].
Thus, we examined whether eCB2.0 can detect endogenous eCB release from primary rat
cortical neurons. Firstly, we expressed eCB2.0 and a red glutamate sensor R^{ncp}-
iGluSnFR[51] in neurons using AAVs to simultaneously record eCB and glutamate dynamics.
Electrical field stimulation (100 pulses at 50 Hz) elicited robust eCB and glutamate signal
increases (**Fig. 3a**), demonstrating that eCB2.0 is able to report endogenous eCB release
and can be used together with red indicators. We then expressed eCB2.0 in neurons using
AAVs and loaded neurons with red calcium dye Calbryte590 to simultaneously record
eCBs and neuronal calcium activity. Again, 100 pulses at 50 Hz elicited strong calcium and
eCB signals (**Fig. 3b**). The rise and decay kinetics of the calcium signal were faster than
those of the eCB signal, consistent with the notion that eCB production is dependent on

calcium[52]. When we stimulated neurons by 1, 5, 10 and 20 pulses at 20 Hz, we observed
progressively increased peak calcium and eCB signals. The peak eCB and peak calcium
signals were highly correlated ($R^2 = 0.99$, **Fig. 3c**). Importantly, when we removed calcium
from the extracellular solution, a 20-pulse-stimulation was unable to elicit either calcium or
eCB signal (**Fig. 3c**), confirming the requirement of calcium activity on eCB release.
Next, we asked which specific eCB(s), namely, 2-AG and/or AEA, was released in
cultured rat cortical neurons. 2-AG is mainly produced from diacylglycerol (DAG) by
diacylglycerol lipase (DAGL) in postsynaptic neurons (**Fig. 3d**). After applying DO34, a
potent and selective DAGL inhibitor[53], the electrically evoked eCB signals were gradually
decreased and were almost abolished after 30 min incubation (**Fig. 3e,f**), indicating 2-AG
is mobilized via DAGL in cultured neurons. We further validated this by manipulating the
eCB degradation pathways: JZL184, an inhibitor of the main 2-AG degrading enzyme
monoacylglycerol lipase (MAGL)[54], significantly prolonged the decay phase of evoked eCB
signals (**Fig. 3g-i**); in contrast, URB597, an inhibitor of the main AEA degrading enzyme
fatty acid amide hydrolase (FAAH)[55], only slightly increased the decay time constant.
These data together demonstrate that 2-AG, but not AEA, is the major eCB released from
cultured rat cortical neurons in response to trains of electrical stimulation.
In addition to the stimuli-evoked eCB signals, we also observed spontaneous, local
eCB transients in neurons in the absence of external stimulation (**Fig. 3j**). The amplitude
and rise kinetics of the transient eCB signal were smaller and slower than that of a single-
pulse evoked eCB signal from the same region of interest (ROI) (**Fig. 3k,l**), suggesting
different releasing patterns of eCBs in these two conditions. The diameter of the transient
signal was ~11.3 μm quantified by the full width at half maximum (FWHM) (**Fig. 3m**),
consistent with previous speculations that eCB functions in local compartments[56,57]. These
spontaneous transients were specific because no such transients were observed in the
presence of AM251 (**Fig. 3l,n,o**).
203
Validation of eCB2.0 in acute mouse brain slices
To further test the ability of eCB2.0 to detect endogenous eCBs in more physiological
relevant conditions, we next expressed eCB sensors in acute mouse brain slices. We first
injected AAVs carrying eCB2.0 or eCBmut into mouse dorsolateral striatum (DLS, **Fig. 4a**),
where eCBs mediate short- and long-term depression and regulates motor behavior[58-60].
Four weeks after AAV injection, acute brain slices were prepared and the expression of
eCB sensors could be visualized (**Fig. 4b**). The fluorescence signals evoked by electrical
stimuli in the DLS was recorded by photometry. Two-pulse stimulation was sufficient to
elicit robust fluorescence increases in a stimulation frequency-dependent manner, while 5-
and 10-pulse stimuli elicited even larger signals (**Fig. 4c,d**). The rise and decay kinetics of
eCB2.0 signals were ~0.8–1.2 s and ~5.2–8.5 s respectively, as quantified by half rise time
and decay time constant (**Fig. 4d**). When we pre-treated brain slices with 10 μM AM251,
the same electrical stimuli failed to evoke an eCB2.0 signal change, indicating the eCB
signals were specific (**Fig. 4e**). eCBmut showed no response to electrical stimuli, further
demonstrating the signal specificity (**Fig. 4e**). 2-photon (2P) microscopic images of eCB2.0
in striatal slices showed the expression of eCB2.0 and its response to bath-applied AEA
(**Extended Data Fig. 4**). We also expressed the eCB2.0 in mouse hippocampal CA1 region,

where eCBs modulate both excitatory and inhibitory inputs[61,62], and recorded eCB2.0
signal in acute slices using 2P microscopy (**Fig. 4f,g**). Electrical stimuli from 5 pulses to
100 pulses at 20 Hz evoked eCB2.0 fluorescence increases (**Fig. 4g,h**), similar to the
results in DLS. Perfusion of 10 µM AEA to the hippocampal slice evoked a large signal
increase, which was blocked by further perfusion of 10 µM AM251 (**Fig. 4i**). In the presence
AM251, even 100 pulses no longer elicited eCB2.0 signal (**Fig. 4j**). These data together
demonstrate that eCB2.0 can be used to detect endogenously released eCBs in acute
brain slices with high sensitivity, specificity and spatiotemporal resolution.
229

Detection of eCBs in basolateral amygdala during foot shock in freely moving mice
Basolateral amygdala (BLA) is a key brain region mediating fear responses and processing
aversive memories[63]. Previous studies demonstrated that the CB1R is highly expressed in
the BLA, and the eCB system in BLA participates in stress expression[64-66]. To directly
record eCB dynamics in animals during an aversive stimulus, we co-expressed eCB2.0 (or
eCBmut) and mCherry in mouse BLA using AAVs and conducted local fiber photometry
recordings (**Fig. 5a,b**). A 2-s electrical foot-shock stimulus induced a time-locked eCB2.0
signal increase in the BLA (**Fig. 5c**), and the response was reproducible in 5 consecutive
trials (**Fig. 5d**). As negative controls, mCherry and eCBmut showed no fluorescence
change during the foot shock (**Fig. 5c,e**). The rise and decay kinetics of eCB2.0 signal
were ~1.0 s and ~6.3 s respectively (**Fig. 5f**). Taken together, these data demonstrate that
eCB2.0 can be used to measure eCB dynamics *in vivo* with high sensitivity and specificity
in freely moving animals.
243

Dual color imaging of eCB and calcium in mouse hippocampal CA1 during running
and seizures
Have shown that eCB2.0 enables the detection of electrically evoked eCB signals in mouse
hippocampal CA1 in acute slices, we then asked whether we could detect *in vivo* eCB
dynamics in CA1 during behavior. We injected AAVs carrying eCB2.0 (or eCBmut) and a
red calcium indicator jRGECO1a[67] in mouse CA1, and conducted head-fixed 2P dual-color
imaging through an implanted cannula above the hippocampus (**Fig. 6a**). Expression of
eCB2.0 and jRGECO1a was clearly observed in CA1 4-6 weeks after virus injection (**Fig.**
6b). We focused on the *stratum pyramidale* layer, which is composed of pyramidal neuron
somata and interneuron axons, including a class that densely express CB1R. When mice
spontaneously ran on a treadmill (**Fig. 6c**), we found rapid increases of both calcium and
eCB signals aligned to the start of running, and decreases of both signals when the running
stopped (**Fig. 6d,e**). In the control group, which expressed eCBmut and jRGECO1a,
calcium signals were intact while eCBmut showed no fluorescence change (**Fig. 6d,e**). The
calcium signal appeared earlier than the eCB signal, although both signals had similar
rising kinetics, while the decay phase of eCB signal was slower than that of the calcium
signal (**Fig. 6f**). These results demonstrate that under normal physiological conditions, the
eCB2.0 sensor enables detection of locomotion-induced eCB signals in mouse
hippocampus *in vivo*.
Epilepsy is a neurological disease characterized by excessive and synchronous
neuronal firing. eCBs are proposed to provide negative feedback during epilepsy to

attenuate the synaptic activity and protect the nervous system, which is exemplified by the
observation that animals with compromised eCB system all exhibit a pro-epileptic
phenotype[68]. To explore whether eCB2.0 could be used to study seizure-related eCB
signals *in vivo*, we used electrical kindling stimulation of the hippocampus contralateral to
the sensor expressing hemisphere to elicit brief self-terminating seizures (**Fig. 6g**). Strong
calcium and eCB signal increases were detected during electrical seizure activity (**Fig. 6h**).
Recent work has shown that seizures are often followed by a spreading calcium wave that
propagates across the cell layer[69]. Interestingly, we also found a propagating eCB wave
that closely followed the calcium wave (**Fig. 6h and Extended Data Movie 1**). In contrast,
eCBmut showed no response during and after seizures (**Fig. 6i**). The velocity and direction
of eCB waves were evident when we extracted the eCB2.0 signal from individual neurons
in the field of view (**Fig. 6j,k**). Notably, eCB waves and calcium waves varied across
experiment sessions and animals (**Fig. 6l**), but for each instance, the calcium and eCB
waves were similar, in agreement with the calcium- and activity-dependence of the eCB
signal. In summary, the running and seizure data demonstrate the ability of the eCB2.0
sensor to report eCB dynamics with high specificity and high spatiotemporal resolution in
both physiological and pathological conditions *in vivo*.
282
DISCUSSION
Here, we report the development and characterization of a genetically-encoded fluorescent
sensor eCB2.0 for detecting eCBs both *in vitro* and *in vivo*. With high sensitivity, selectivity
and kinetics, eCB2.0 enables the detection of endogenous eCB signals in cultured neurons
and acute brain slices. eCB2.0 also enables detection of eCB release in the amygdala of
freely behaving mice, running induced eCB signals in mouse hippocampus, as well as eCB
waves in mouse hippocampus under seizure conditions.
The on kinetics of eCB2.0 measured by local micropressure application at room
temperature is ~1.6 s, which is likely overestimated since the on kinetics of eCB2.0 signal
in DLS slices and in the BLA *in vivo* was within or about 1 s. The temporal resolution of
eCB2.0 is dramatically improved compared to microdialysis (minutes), and it may be further
optimized in the future to capture faster signals[70]. Currently, eCB2.0 detects both 2-AG and
AEA. Given 2-AG and AEA have different regulatory pathways, brain region and cell type
specificities, the development of new sensors with distinct eCB molecular specificity, as
well as different color spectra, would be also desirable for future studies.
eCB-mediated retrograde synaptic modulation was identified during the study of
depolarization induced suppression of inhibition or excitation (DSI or DSE) in the
hippocampus and cerebellum[25,26,28]. Those and subsequent experiments relied on the
electrophysiological recordings of synaptic transmission in combination with
pharmacological (such as activation and inhibition of cannabinoid receptors, inhibition of
production or degradation enzymes) or genetic (such as knockout of corresponding
receptors and enzymes) manipulations, thus lacking direct and physiologically relevant
eCB detection. Moreover, recording on the somata of neurons is unable to capture the
precise spatial distribution of eCBs. For example, DSI recorded from paired whole-cell
recordings in hippocampal slices indicate that the depolarization of one neuron could inhibit
GABAergic inputs to neurons ~20 μm or less away from it, suggesting the diffusion distance

of eCBs from a single neuron[25]. Similar results were obtained in cerebellar slices using two

separate stimulation electrodes to evoke eCB release from two dendritic regions of a single

Purkinje cell[56]. These data demonstrate that eCB signaling is relatively local and is likely

to be tightly controlled. However, the detailed spatial profile of eCB signal is still not known.

In addition, although the sampling rate of electrophysiological recording is generally high

(>kHz), the eCB signals revealed by the change of evoked postsynaptic currents (ePSCs)

have a sampling interval of ~2 s, which forms a temporal bottleneck. eCB2.0, therefore, for

the first time provides the opportunity to examine the eCB signal at high spatial (e.g.,

synaptic) and temporal (sub-second) resolution, similar to studies using other

neurotransmitter sensors[71,72]. In cultured neurons, we detected spontaneous eCB

transients with a diameter of ~11 μm, which is already smaller than the previous estimation

of the eCB diffusion distance. It will be interesting to test whether the local transient signals

originate from single spines.

We have demonstrated that eCB2.0 could be used in multiple preparations *in vitro* and

in vivo to report real-time neuronal eCB dynamics. Given the complexity of the nervous

system, future directions for research based on eCB2.0 applications may include the

identity of cell types that release eCBs, the mechanisms and temporal properties of eCB

release, characteristics of eCB diffusion, the duration of eCB signals, the nature of the cell

types and subcellular elements targeted by eCBs and the effects on them. Answering these

fundamental questions will significantly enrich our understanding of the mechanisms and

functions of eCB signaling in neural circuits.

Malfunction of eCB system is associated with multiple neurological disorders including

stress/anxiety, movement disorders, substance use disorders and epilepsy. The results of

eCB detection during foot shock, running and seizure in mice show clear examples of how

the eCB2.0 sensor could help to elucidate the fast eCB dynamics during both physiological

and pathological processes. The eCB2.0 sensor should be able to detect all CB1R agonists

(**Extended Data Fig. 3**) including Δ-9-tetrahydrocannabinol (Δ-9-THC) in the brain and

periphery following drug administration. This would allow investigators to track the time

course of Δ-9-THC actions and the impact of cannabis drugs on eCB signaling. Thus, eCB

sensors open a new era of endocannabinoid research aimed at understanding this system

at unprecedented, physiologically-relevant spatial and temporal scales.

METHODS

Molecular biology

DNA fragments were amplified by PCR with primers (TSINGKE Biological Technology) containing 25–30 bp overlaps. Plasmids were constructed by restriction enzyme cloning or Gibson Assembly. Plasmid sequences were verified by Sanger sequencing. For characterization of eCB2.0 and eCBmut in HEK293T cells, eCB2.0 and eCBmut genes were cloned into pDisplay vector with a IgK leader sequence before the sensor gene. An IRES-mCherry-CAAX cassette was inserted after the sensor gene for indicating cell membrane and calibrating the sensor fluorescence. For characterization of eCB2.0 in neurons, eCB2.0 gene was cloned into pAAV vector under control of a human synapsin promoter (pAAV-hSyn), PSD95-mScarlet and synaptophysin-mScarlet genes were cloned into pDest vetor under control of a CMV promoter. For the $G_{\beta\gamma}$ sensor assay, human CB1 was cloned into the pCI vector (Promega), eCB2.0 and eCBmut genes were cloned into the peGFP-C1 vector (Takara), replacing the eGFP open reading frame. For the Tango assay, CB1, eCB2.0 and eCBmut genes were cloned into pTango vector. pAAV-hsyn-eCBmut and pAAV-hsyn-R^{ncp}-iGluSnFR were also constructed for virus production.

AAV virus preparation

AAV2/9-hSyn-eCB2.0 (9.5×10^{13} viral genomes (vg)/mL), AAV2/9-hSyn-eCBmut (8.0×10^{13} vg/mL), AAV2/9-hSyn-R^{ncp}-iGluSniFR (6.2×10^{13} vg/mL, all packaged at Vigene Biosciences, China), AAV8-hSyn-mCherry (#114472, Addgene) and AAV1-Syn-NES-jRGECO1a-WPRE-SV40 (Penn Vector Core) were used to infect cultured neurons or *in vivo*.

Cell cultures

HEK293T cells were cultured at 37°C, 5% CO_2 in DMEM (Biological Industries) supplemented with 10% (v/v) fetal bovine serum (Gibco) and penicillin (100 unit/mL)-streptomycin (0.1 mg/mL) (Biological Industries). HEK293T cells were plated on 96-well plates or 12 mm glass coverslips in 24-well plates. Cells at 60–70% confluency were transfected with plasmids using polyethylenimine (PEI) (300 ng DNA/well for 96-well plates and 1 μg DNA/well for 24-well plates, DNA:PEI = 1:3) for 4–6 h before changing fresh culture medium. Imaging was performed 24–36 h after transfection. Rat cortical neurons were prepared from postnatal day 0 (P0) Sprague-Dawley rat. Briefly, rat cortical neurons were dissociated from rat brain cortex after dissection and digestion in 0.25% Trypsin-EDTA (Biological Industries), and plated in 12 mm glass coverslips in 24-well plates coated with poly-D-lysine (Sigma-Aldrich). Neurons were cultured at 37°C, 5% CO_2 in Neurobasal Medium (Gibco) supplemented with 2% B-27 Supplement (Gibco), 1% GlutaMAX (Gibco), and penicillin (100 unit/mL)-streptomycin (0.1 mg/mL) (Biological Industries). Cultured neurons were transfected at 7–9 day *in vitro* (DIV7–9) using calcium phosphate transfection method. Imaging was performed 48 h after transfection. Cultured neurons were infected by AAVs expressing eCB2.0, eCBmut and R^{ncp}-iGluSnFR at DIV3–5, and imaging was performed at DIV12–20. Calbryte590 (AAT Bioquest) was loaded into neurons for 1 h before imaging.

Animals

All experiment protocols were approved by the respective Laboratory Animal Care and Use Committees of Peking University, National Institute on Alcohol Abuse and Alcoholism, Cold Spring Harbor Laboratory (CSHL) and Stanford University, and studies were performed in accordance with the guidelines by the US National Institutes of Health. Postnatal day 0 (P0) Sprague-Dawley rats of both sexes (Beijing Vital River Laboratory) and P42–P150 C57BL/6J mice of both sexes (Beijing Vital River Laboratory and The Jackson Laboratory) were used in this study. Mice were housed under a normal 12 h light/dark cycle with food and water available *ad libitum*.

Confocal imaging in cultured cells

Before imaging, the culture medium for cells was replaced with Tyrode's solution consisting of (in mM): 150 NaCl, 4 KCl, 2 $MgCl_2$, 2 $CaCl_2$, 10 HEPES, and 10 glucose (pH 7.4). 0 mM Ca^{2+} Tyrode's solution contained 0 mM $CaCl^2$ and 2 mM EGTA. The Opera Phenix high-content screening system (PerkinElmer, USA) was used for imaging HEK293T cells in 96-well plates. It was equipped with a 60x/1.15 NA water-immersion objective, a 488 nm laser and a 561 nm laser. A 525/50 nm emission filter and a 600/30 nm emission filter were used to collect green and red fluorescence respectively. A Ti-E A1 confocal microscopy (Nikon, Japan) was used for imaging cultured cells in 12 mm coverslips. It was equipped with a 10x/0.45 NA objective, a 20x/0.75 NA objective, a 40x/1.35 NA oil-immersion objective, a 488 nm laser and a 561 nm laser. A 525/50 nm emission filter and a 595/50 nm emission filter were used to collect green and red fluorescence respectively. Drugs, including 2-AG (Tocris), AEA (Cayman), AM251 (Tocris), LPA (Tocris), S1P (Tocris), ACh (Solarbio), DA (Sigma-Aldrich), GABA (Tocris), Glu (Sigma-Aldrich), Gly, NE (Tocris), 5-HT (Tocris), His (Tocris), Epi (Sigma-Aldrich), Ado (Tocris), Tyr (Sigma-Aldrich), WIN55212-2 (Cayman), DO34 (MCE), JZL184 (Cayman), URB597 (Cayman) were applied by replacing drug-containing Tyrode's solution for 96-well plate imaging, or applied by bath application or by a custom made perfusion system for 12 mm coverslip imaging. The micropressure application of drugs was controlled by Pneumatic PicoPump PV800 (World Precision Instruments). Cultured neurons were field stimulated using parallel platinum electrodes at a distance of 1 cm controlled by a Grass S88 stimulator (Grass Instruments). The voltage was 80 V and the duration of each stimulation pulse was 1 ms.

BRET G$_{\beta\gamma}$ sensor assay

eCB2.0, eCBmut or CB1 genes were transfected using PEI into HEK293 tsA201 cells in 24 well plates in a 1:5 plasmid mass ratio with a single construct designed to separately express human GNAOa, human GNB1 (fused to amino acids 156-239 of Venus), human GNG2 (fused to amino acids 2-155 of Venus) and NanoLuc (Promega) fused to the amino terminal 112 amino acids of human phosducin circularly permutated at amino acids 54/55. The NanoLuc/phosducin fusion portion also contained a kRAS membrane targeting sequence on the carboxy terminal end. The G$_{\beta\gamma}$ sensor components were combined by either restriction enzyme cloning or InFusion (Takara) assembly. Templates for assembly were derived from human whole brain cDNA (Takara) for GNAOa and human retinal cDNA (Takara) for phosducin. Templates for hGNB1 and hGNG2 Venus fusions were a generous

gift from Nevin Lambert (Augusta University). Cells were harvested approximately 24 hours
post-transfection with 10mM EDTA in PBS (pH7.2), pelleted and resuspended in
Dulbecco's modified PBS (Life Technologies) without calcium or magnesium. Furimazine
(Promega) was added at a 1/100 dilution to 100 µl of cell suspension in a black 96 well
plate and BRET readings were taken in a Pherastar FS plate reader (Berthold) equipped
with a Venus BRET cube. Acceptor (Venus) and donor (NanoLuc) signals were monitored
at 535nm and 475nm respectively. Net BRET was calculated as the acceptor/donor ratio
of each sample minus the acceptor/donor ratio of a donor only sample. Readings were
taken before and 2-4 minutes after agonist 2-AG (Tocris, 20 µM final concentration)
application to activate CB1 or eCB sensors.
439

Tango assay
eCB2.0, eCBmut and CB1R genes fused with the tTA gene in pTango vectors were
transfected into the HTLA reporter cell line using the PEI in 6 well plates. The HLTA cell
line expresses a β-arrestin2-TEV fusion gene and a tTA-dependent luciferase reporter
gene. 24 h after transfection, cells in 6 well plates were collected after trypsin treatment
and plated in 96 well plates. CB1R agonist AEA was applied at concentrations ranging from
0.01 nM to 10 µM. The luciferase was expressed for 12 h before the luminescence
measurement. Bright-Glo (5 µM, Promega) was added and luminescence was measured
by the VICTOR X5 multi-label plate reader (PerkinElmer).
449

Acute brain slices
Photometry recording in the DLS in acute mouse brain slices
Adult (> 10 weeks) male C57BL/6J mice were anesthetized with isoflurane and injected
with AAV vectors into dorsal lateral striatum (300 nL, coordinates relative to bregma in mm:
A/P: + 0.75; M/L: ± 2.5; D/V: − 3.5) at a rate of 50 nL/min. Mice were given an injection of
ketoprofen (5 mg/kg, s.c.) and postoperative care was provided daily until mice regained
their preoperative weight. After a minimum of 4 weeks following AAV injection, mice were
deeply anesthetized with isoflurane, decapitated and the brains were extracted and placed
in ice cold sucrose cutting solution (in mM): 194 sucrose, 30 NaCl, 4.5 KCl, 26 $NaHCO_3$,
1.2 NaH_2PO_4, 10 D-glucose, 1 $MgCl_2$ saturated with 5% CO_2/ 95% O_2. Coronal brain slices
(250 µm) were prepared and slices were incubated at 32°C for ~60 min in artificial
cerebrospinal fluid (ACSF) (in mM): 124 NaCl, 4.5 KCl, 26 $NaHCO_3$, 1.2 NaH_2PO_4, 10 D-
glucose, 1 $MgCl_2$, 2 $CaCl_2$. After incubation at 32°C, slices were held at room temperature
until initiating an experiment. Photometry recordings were acquired using an Olympus
BX41 upright epifluorescence microscope equipped with a 40x/0.8 NA water-emersion
objective and a FITC filter set. Slices were superfused at ~2 mL/min with ACSF (29–31°C).
A twisted bipolar polyimide-coated stainless-steel stimulating electrode (~200 µm tip
separation) was placed in the DLS just medial to the corpus callosum and slightly below
the tissue surface in a region with visible eCB sensors fluorescence. GRAB$_{eCB}$ sensors
were excited using a 470 nm light emitting diode (LED) (ThorLabs, USA). Photons passing
through a 180 µm^2 aperture, positioned just lateral to the stimulating electrode, were
directed to a PMT (Model D-104, Photon Technology International, USA). The PMT output
was amplified (gain: 0.1 µA/V; time constant: 5 ms), filtered at 50 Hz and digitized at 250

Hz using a Digidata 1550B and Clampex software (Molecular Devices LLC, USA). For all

experiments, $GRAB_{eCB}$ measurements were acquired as discrete trials repeated every 3

minutes. For each trial, the light exposure period was 35–45 seconds to minimize $GRAB_{eCB}$

photobleaching, while capturing peak responses and the majority of the decay phase. To

evoke an eCB transient, a train of electrical pulses (1.0–1.5 mA, 200–500 µs) was delivered

5 s after initiating $GRAB_{eCB}$ excitation.

2-photon imaging in the hippocampus in acute mouse brain slices

Adult (6–8 weeks) C57BL/6J mice of both sexes were anesthetized with an intraperitoneal

injection of 2,2,2-tribromoethanol (Avertin, 500 mg/kg body weight, Sigma-Aldrich) and

injected with AAV vectors into hippocampal CA1 (400 nL, coordinates relative to bregma

in mm: A/P: − 1.8; M/L: ± 1.0; D/V: − 1.2) at a rate of 46 nL/min. After a minimum of 4 weeks

following AAV injection, mice were deeply anesthetized with an intraperitoneal injection of

2,2,2-tribromoethanol, decapitated and the brains were extracted and placed in ice cold

choline chloride cutting solution (in mM): 110 choline-Cl, 2.5 KCl, 0.5 $CaCl_2$, 7 $MgCl_2$, 1

NaH_2PO_4, 1.3 Na ascorbate, 0.6 Na pyruvate, 25 NaHCO3 and 25 glucose saturated with

5% CO_2/ 95% O_2. Coronal brain slices (300 µm) were prepared and slices were incubated

at 34°C for ~40 min in modified ACSF solution (in mM): 125 NaCl, 2.5 KCl, 2 $CaCl_2$, 1.3

$MgCl_2$, 1 NaH_2PO_4, 1.3 Na ascorbate, 0.6 Na pyruvate, 25 $NaHCO_3$ and 25 glucose

saturated with 5% CO_2/ 95% O_2. 2-photon imaging were performed under an FV1000MPE

2-photon microscope (Olympus) equipped with a 25x/1.05 NA water-immersion objective

and a mode–locked Mai Tai Ti: Sapphire laser (Spectra-Physics). Slices were superfused

at ~4 mL/min with modified ACSF (32-34°C). A 920 nm laser was used to excite eCB2.0

sensor, and fluorescence was collected using a 495–540 nm filter. For electrical stimulation,

a bipolar electrode (cat. number WE30031.0A3, MicroProbes for Life Science) was

positioned near the Stratum radiatum layer in CA1 using fluorescence guidance.

Fluorescence imaging and electrical stimulation were synchronized using an Arduino board

with custom-written programs. All images collected during electrical stimulation were

recorded at a frame rate of 0.3583 s/frame with 256×192 pixels per frame. The stimulation

voltage was 4–6 V, and the pulse duration was 1 ms. Drugs were applied to the imaging

chamber by perfusion at a flow rate at 4 mL/min.

Fiber photometry recording of eCB signals in the BLA during foot shock

Adult (10–12 weeks) C57 BL/6J mice of both sexes were injected with 300 nL of a 10:1

mixture of AAV-hSyn-eCB2.0 and AAV-hSyn-mCherry viruses, or a 10:1 mixture of AAV-

hSyn-eCBmut and AAV-hSyn-mCherry viruses, into the right basolateral amygdala at A/P

−1.78 mm, M/L −3.30 mm, D/V −4.53 mm (relative to Bregma). Virus injection was

performed using a glass pipette with a Picospritzer III microinjection system (Parker

Hannifin, USA). After injection, a 200-µm-diameter, 0.37 NA fiber (Inper, China) was

implanted into the same location, and secured with self-adhesive resin cement (3M). A

head bar was also mounted with resin cement onto the skull. At least 14 days were waited

before doing photometry recording. Photometry recording was taken using a commercial

photometry system (Neurophotometrics, USA). A patch cord (0.37 NA, Doric Lenses,

Canada) was attached to the photometry system, and to the fiber in the mouse brain. A

470 nm LED was used to excite eCB and eCBmut sensors, and a 560 nm LED was used

to excite mCherry. The average power levels of LED (measured at the output end of the
patch cord) were 160 µW for eCB/eCBmut sensors, and 25 µW for mCherry. The recording
frequency was 10 Hz. Photometry data were acquired with Bonsai 2.3.1 software. Mice
were free-moving in a shock box (Habitest, Coulbourn Instruments, USA) inside of a
sound-proof behavior box with light on. The FreezeFrame software was used to generate
triggers to the shock generator (Coulbourn Instruments) and Bonsai software. Five pulses
of 2-s electric shocks with an intensity of 0.7 mA were delivered to the shock box with 90–
120 s ITIs. After photometry recording, animals were perfused with phosphate-buffered
saline (PBS), and subsequently with 4% paraformaldehyde (PFA) in PBS. Brain tissues
were fixed in PFA solution overnight, and then were dehydrated with 30% sucrose in PBS
solution for 24 h. Brain slices were cut using a Leica SM 2010R microtome (Leica
Biosystems, USA). Floating brain slices were blocked (5% BSA, 0.1% Triton in PBS) at
room temperature for 2 h, and then stained with chicken anti-GFP (1:1000, Aves, #GFP-
1020) and rabbit anti-RFP (1:500, Rockland, #600-401-379) primary antibodies in antibody
solution (3% BSA, 0.1% Triton in PBS) at 4 °C for 24 h. Slices were next rinsed with PBS
15 min for 3 times, and stained with DAPI (5 µg/mL, Invitrogen, #D1306), Alexa Fluor 488
donkey anti-chicken (1:250, Jackson ImmunoResearch, #703-545-155) and Alexa Fluor
568 donkey anti-rabbit (1:250, Invitrogen, #A10042) in antibody solution at 4 °C for 24 h.
Confocal images were taken with a Zeiss LSM780 confocal microscopy (Zeiss, Germany).
536

2-photon imaging in living mice
Adult (P100− 150) C57BL/6J mice of both sexes were used. Mice were injected with a mix
of AAV1-Syn-NES-jRGECO1a-WPRE-SV40 and AAV9-hSyn-eCB2.0 (or eCBmut) viruses
(300-400 nL each, full titer) into the right CA1 (2.3 mm posterior, 1.5 mm lateral, 1.35 mm
ventral to bregma) using a Hamilton syringe. After recovery from virus injection, the cortex
above the injection site was aspirated and a stainless steel cannula with attached
coverglass was implanted over the hippocampus as previously described[73,74], followed by
a stainless steel headbar. A chronic bipolar wire electrode (tungsten, 0.002", 0.5 mm tip
separation, A-M systems) was implanted into the left ventral hippocampus (3.2 mm
posterior, 2.7 mm lateral, 4.0 mm ventral to bregma) as previously described[75]. Head-fixed
mice running on a linear treadmill with a cue-less belt (2 m) were imaged using a resonant
scanner 2-photon microscope (Neurolabware), equipped with a pulsed IR laser tuned at
1000 nm (Mai Tai, Spectra-Physics), GaAsP PMT detectors (H11706P-40, Hamamatsu),
and a 16x objective (0.8 NA WI, Nikon). 2-photon image acquisition and treadmill speed
monitoring were controlled by Scanbox (Neurolabware). Bipolar electrodes were recorded
using a differential amplifier (Model 1700, A-M Systems). Seizures were elicited by electric
stimulation above seizure threshold by 150 µA of current delivered in 1ms biphasic pulses
at 60Hz for 1 second, using a constant-current stimulator (A-M Systems model 2100).
Following *in vivo* recordings, mice were anaesthetized by isoflurane, prior to an
intraperitoneal injection of a mixture of ketamine (100 mg/kg) and Xylazine (10 mg/kg) in
saline. Animals were transcardially perfused with saline (0.9% NaCl for 1 minute) and then
with fixative solution (4% paraformaldehyde and 0.2% picric acid in 0.1 M phosphate buffer).
Perfused brains were then post-fixed in the same fixative solution for 24 h at 4 °C, prior to
slicing on a vibratome (VTS1200, Leica Biosystems). Sections were then washed and

mounted in Vectashield (Vector Laboratories). Confocal images were acquired on a Zeiss
LSM 710 imaging system using a 20x 0.8 NA objective.
563
Data processing
Confocal imaging
Data were collected and analyzed using Harmony high-content imaging and analysis
software (PerkinElmer) for 96-well plate imaging. Briefly, membrane regions were selected
as ROIs and green fluorescence of sensors was normalized by red fluorescence of
mCherry-CAAX (G/R). $\Delta F/F_0$ was calculated as $(G/R_{after\ durg} - G/R_{before\ drug}) / (G/R_{before\ drug})$.
For 12 mm coverslip imaging, data were collected using the NIS-Element software (Nikon)
and analyzed using the ImageJ software (NIH). $\Delta F/F_0$ was calculated as $(F_t - F_0)/F_0$, while
F_0 was quantified before drug application or stimulation. Data were plotted using OriginPro
2020 (OriginLab).
Slice photometry and 2-photon imaging
For slice photometry recording, eCB transients were calculated as $\Delta F/F_0$ by averaging the
PMT voltage (V) for a period of 1 s just prior to electrical stimulation (F_0) and then
calculating V/F_0-1 for each digitized data sample. The eCB decay phase was fitted with a
single exponential decay accounting for a sloping baseline. Rise $t_{1/2}$ was calculated in
Graphpad Prism (v8.3; San Diego, CA, USA) by fitting the rising phase of the eCB transient
with an asymmetrical logistics curve. Photometry sweeps were exported to Microsoft Excel
(v16.3; Redmond, WA, USA) to calculate normalized $\Delta F/F_0$ traces and peak $\Delta F/F_0$ values.
For slice 2-photon imaging, data were collected using FV10-ASW (Olympus) and analyzed
using ImageJ (NIH). $\Delta F/F_0$ was calculated as $(F_t - F_0)/F_0$, while F_0 was quantified before
stimulation. Data were plotted using OriginPro 2020 (OriginLab).
Fiber photometry in mice during foot shock
Off-line analysis of the photometry data was conducted with the Matlab software
(MathWorks). Data were plotted using OriginPro 2020 (OriginLab).
2-photon imaging in mice during locomotion and seizure
Imaging data were processed and analyzed using Python scripts. For the analysis of
single-cell responses, movies were initially motion corrected by rigid translation, followed
by non-rigid correction (*HiddenMarkov2D*) by the sima package[76]. Binary regions of
interest (ROIs) were selected in a semi-automated manner. For the initial automated
detection, movies were divided into segments of 100 frames each, the average intensity
projection of each segment was computed and the resulting resampled movie was used
for detection. In sessions with electric stimulation, only the baseline period before the
stimulation was used for segmentation. The *PlaneCA1PC* method of sima was run on the
inverted resampled movie, which resulted in detection of the hollow nuclei of cells. These
ROIs were filtered based on size, and binary dilation was performed to include the
cytoplasm around the nuclei. In a subsequent step, ROIs were detected in the non-inverted
resampled movie, filtered based on size and those that did not overlap with existing ROIs
were added to the set. ROIs outside the str. pyramidale were excluded. Next, the
fluorescence intensity traces were extracted for each ROI by averaging the included pixel
intensities within each frame. For analyzing run responses, only sessions with no electric
stimuli were included, and signals were pulled from the motion-corrected movies. These

raw traces were processed following standard steps for obtaining $\Delta F/F_0$ traces with a modified approach to determine the time-dependent baseline. A 3rd degree polynomial was fit on the trace after applying temporal smoothing, removing peaks (detected using continuous wavelet transform by scipy.signal), eliminating periods of running, and ignoring the beginning and end of the recording. The calculated polynomial was then used as a baseline. Z-scored traces were obtained after determining the standard deviation (SD) of each cell's baseline by excluding events exceeding 2 SD in two iterations.

For analyzing spreading activity, only sessions with electric stimulus that triggered an electrographic seizure and a spreading wave were included, and while the segmentation was performed based on the motion-corrected baseline segments of the recordings, signals were pulled from non-motion-corrected movies (as image-based motion correction was not feasible during seizures). $\Delta F/F_0$ traces were obtained using a constant baseline determined by averaging the pre-stimulus segments of the traces. For the analysis changes in average fluorescence intensity, a single large region of interest (ROI) was manually drawn to include the cell bodies in the pyramidal layer, and $\Delta F/F_0$ traces were pulled and processed as described above. Event-triggered averages were calculated after automatically detecting the frames with running onsets and stops by fixed criteria across all sessions. The average was computed in two steps (i.e. events were first averaged by cell, then cells averaged by virus type (eCB or eCBmut). Decay time constants were computed as the parameter of a 2nd degree polynomial fit after a log transform on the trace following the peak of the stop-triggered average trace. Rise times were determined between the frame where the start-triggered average signal first reached 90% of the range between baseline and peak, and the last frame before that where the signal was below 10% of the range. To determine the spreading speed and direction of waves, first the peak time of the wave was determined in each session by inspecting the average $\Delta F/F_0$ trace (including all cells). Then, the relative peak location (Δt) of the $\Delta F/F_0$ trace of each cell in the trace including ± 200 frames (12.8 s) around the wave peak was determined. Finally, two linear fits were computed using the x and y centroid coordinates of each ROI ($\Delta t \sim x$, $\Delta t \sim y$). The 2D speed was computed from the slopes of the two 1D fits. The direction was determined by computing the unity vector from the starting- to the end point of the fits between ± 3 s around the wave peak. The average speed was obtained by averaging the speed of individual sessions, while the average direction was obtained from the sum of the unity vectors of individual sessions. Data were plotted using Python and OriginPro 2020 (OriginLab).

Statistical analysis

Values with error bars indicate mean ± SEM. Group data were analyzed using the Student's t test or one-way ANOVA, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$; n.s., not significant, $p > 0.05$.

Data and software availability

Plasmids for expressing eCB2.0 and eCBmut used in this study were deposited at Addgene (https://www.addgene.org/Yulong_Li/).

ACKNOWLEDGMENTS
This work was supported by Beijing Municipal Science & Technology Commission
(Z181100001318002, Z181100001518004), National Natural Science Foundation of China
(31925017), NIH BRAIN Initiative (NS103558), Shenzhen-Hong Kong Institute of Brain
Science (NYKFKT2019013) and grants from the Peking-Tsinghua Center for Life Sciences
and the State Key Laboratory of Membrane Biology at Peking University School of Life
Sciences to Y.L.; a Division of Intramural Clinical and Biological Research of the NIAAA
grant (ZIA AA000416) to D.M.L; a NIH BRAIN Initiative (NS103558) grant to J.D.; NIH
grants (R01MH101214 and R01NS104944) to B.L.; an American Epilepsy Society
Postdoctoral Fellowship and a NIH grant (K99NS117795) to B.D.; a Canadian Institutes for
Health Research postdoctoral fellowship to J.S.F.; a NIH grant to I.S. (NS99457). We thank
Li lab members for helpful discussions. We thank Yi Rao for sharing the 2-photon
microscope and Xiaoguang Lei at PKU-CLS for providing support for the Opera Phenix
high-content screening system.
663

AUTHOR CONTRIBUTIONS

Y.L. conceived the project. A.D., K.H., H.L.P., R.C. and J.D. performed experiments related
to developing, optimizing, and characterizing the sensors in cultured HEK293T cells and
neurons. L.J.D. performed the surgery and photometry recording experiments related to
the validation of the sensor in DLS brain slices under the supervision of D.M.L.. A.D.
performed the surgery and 2-photon imaging in the hippocampal brain slices. E.A.
performed the surgery and 2-photon imaging in the striatal brain slices under the
supervision of J.D.. W. G. performed fiber photometry recordings in freely moving mice
during foot shock under the supervision of B.L.. B.D. and J.S.F. performed the *in vivo* 2-
photon imaging in the hippocampus in mice during running and seizure under the
supervision of I.S. All authors contributed to the data interpretation and analysis. A.D. and
Y.L. wrote the manuscript with input from other authors.
676

677 **COMPETING FINANCIAL INTERESTS**
678 Y. L. has filed patent applications whose value might be affected by this publication.
679

REFERENCES
1.   Zuardi, A.W. History of cannabis as a medicine: a review. *Braz J Psychiatry* **28**, 153-157
(2006).
2.   Piomelli, D. The molecular logic of endocannabinoid signalling. *Nat Rev Neurosci* **4**, 873-
884 (2003).
3.   Wilson, R.I. & Nicoll, R.A. Endocannabinoid signaling in the brain. *Science* **296**, 678-682
(2002).
4.   Kano, M., Ohno-Shosaku, T., Hashimotodani, Y., Uchigashima, M. & Watanabe, M.
Endocannabinoid-mediated control of synaptic transmission. *Physiological reviews* **89**, 309-380
(2009).
5.   Hebert-Chatelain, E.*, et al.* A cannabinoid link between mitochondria and memory. *Nature*
539, 555-559 (2016).
6.   Benard, G.*, et al.* Mitochondrial CB(1) receptors regulate neuronal energy metabolism. *Nat*

Neurosci **15**, 558-564 (2012).

7.   Jimenez-Blasco, D*., et al.* Glucose metabolism links astroglial mitochondria to cannabinoid effects. *Nature* **583**, 603-608 (2020).

8.   Stella, N. Cannabinoid signaling in glial cells. *Glia* **48**, 267-277 (2004).

9.   Navarrete, M., Diez, A. & Araque, A. Astrocytes in endocannabinoid signalling. *Philos Trans R Soc Lond B Biol Sci* **369**, 20130599 (2014).

10. Chevaleyre, V., Takahashi, K.A. & Castillo, P.E. Endocannabinoid-mediated synaptic plasticity in the CNS. *Annu Rev Neurosci* **29**, 37-76 (2006).

11. Oddi, S., Scipioni, L. & Maccarrone, M. Endocannabinoid system and adult neurogenesis: a focused review. *Curr Opin Pharmacol* **50**, 25-32 (2020).

12. Moreira, F.A. & Lutz, B. The endocannabinoid system: emotion, learning and addiction. *Addict Biol* **13**, 196-212 (2008).

13. Guindon, J. & Hohmann, A.G. The endocannabinoid system and pain. *CNS Neurol Disord Drug Targets* **8**, 403-421 (2009).

14. Kesner, A.J. & Lovinger, D.M. Cannabinoids, Endocannabinoids and Sleep. *Frontiers in molecular neuroscience* **13**, 125 (2020).

15. Silvestri, C. & Di Marzo, V. The endocannabinoid system in energy homeostasis and the etiopathology of metabolic disorders. *Cell Metab* **17**, 475-490 (2013).

16. Katona, I. & Freund, T.F. Endocannabinoid signaling as a synaptic circuit breaker in neurological disease. *Nat Med* **14**, 923-930 (2008).

17. Fernandez-Espejo, E., Viveros, M.P., Nunez, L., Ellenbroek, B.A. & Rodriguez de Fonseca, F. Role of cannabis and endocannabinoids in the genesis of schizophrenia. *Psychopharmacology (Berl)* **206**, 531-549 (2009).

18. Fraguas-Sanchez, A.I., Martin-Sabroso, C. & Torres-Suarez, A.I. Insights into the effects of the endocannabinoid system in cancer: a review. *Br J Pharmacol* **175**, 2566-2580 (2018).

19. Patel, S., Hill, M.N., Cheer, J.F., Wotjak, C.T. & Holmes, A. The endocannabinoid system as a target for novel anxiolytic drugs. *Neurosci Biobehav Rev* **76**, 56-66 (2017).

20. Ligresti, A., Petrosino, S. & Di Marzo, V. From endocannabinoid profiling to 'endocannabinoid therapeutics'. *Curr Opin Chem Biol* **13**, 321-331 (2009).

21. Sabatini, B.L. & Regehr, W.G. Timing of synaptic transmission. *Annu Rev Physiol* **61**, 521-542 (1999).

22. van den Pol, A.N. Neuropeptide transmission in brain circuits. *Neuron* **76**, 98-115 (2012).

23. Zoerner, A.A*., et al.* Quantification of endocannabinoids in biological systems by chromatography and mass spectrometry: a comprehensive review from an analytical and biological perspective. *Biochim Biophys Acta* **1811**, 706-723 (2011).

24. Marchioni, C*., et al.* Recent advances in LC-MS/MS methods to determine endocannabinoids in biological samples: Application in neurodegenerative diseases. *Anal Chim Acta* **1044**, 12-28 (2018).

25. Wilson, R.I. & Nicoll, R.A. Endogenous cannabinoids mediate retrograde signalling at hippocampal synapses. *Nature* **410**, 588-592 (2001).

26. Kreitzer, A.C. & Regehr, W.G. Retrograde inhibition of presynaptic calcium influx by endogenous cannabinoids at excitatory synapses onto Purkinje cells. *Neuron* **29**, 717-727 (2001).

27. Maejima, T., Hashimoto, K., Yoshida, T., Aiba, A. & Kano, M. Presynaptic inhibition caused

by retrograde signal from metabotropic glutamate to cannabinoid receptors. *Neuron* **31**, 463-
475 (2001).

28.   Ohno-Shosaku, T., Maejima, T. & Kano, M. Endogenous cannabinoids mediate retrograde
signals from depolarized postsynaptic neurons to presynaptic terminals. *Neuron* **29**, 729-738
(2001).

29.   Wiskerke, J.*, et al.* Characterization of the effects of reuptake and hydrolysis inhibition on
interstitial endocannabinoid levels in the brain: an in vivo microdialysis study. *ACS Chem*
Neurosci **3**, 407-417 (2012).

30.   Walker, J.M., Huang, S.M., Strangman, N.M., Tsou, K. & Sanudo-Pena, M.C. Pain
modulation by release of the endogenous cannabinoid anandamide. *Proc Natl Acad Sci U S A*
96, 12198-12203 (1999).

31.   Sun, F.*, et al.* A Genetically Encoded Fluorescent Sensor Enables Rapid and Specific
Detection of Dopamine in Flies, Fish, and Mice. *Cell* **174**, 481-496 e419 (2018).

32.   Patriarchi, T.*, et al.* Ultrafast neuronal imaging of dopamine dynamics with designed
genetically encoded sensors. *Science* **360**(2018).

33.   Jing, M.*, et al.* A genetically encoded fluorescent acetylcholine indicator for in vitro and in
vivo studies. *Nat Biotechnol* **36**, 726-737 (2018).

34.   Feng, J.*, et al.* A Genetically Encoded Fluorescent Sensor for Rapid and Specific In Vivo
Detection of Norepinephrine. *Neuron* **102**, 745-761 e748 (2019).

35.   Peng, W.*, et al.* Regulation of sleep homeostasis mediator adenosine by basal forebrain
glutamatergic neurons. *Science* **369**(2020).

36.   Patriarchi, T.*, et al.* An expanded palette of dopamine sensors for multiplex imaging in vivo.
Nat Methods (2020).

37.   Jing, M.*, et al.* An optimized acetylcholine sensor for monitoring in vivo cholinergic activity.
Nature Methods (2020).

38.   Sun, F.*, et al.* New and improved GRAB fluorescent sensors for monitoring dopaminergic
activity in vivo. 2020.2003.2028.013722 (2020).

39.   Wan, J.*, et al.* A genetically encoded GRAB sensor for measuring serotonin dynamics
in vivo. 2020.2002.2024.962282 (2020).

40.   Howlett, A.C.*, et al.* International Union of Pharmacology. XXVII. Classification of
cannabinoid receptors. *Pharmacol Rev* **54**, 161-202 (2002).

41.   Hua, T.*, et al.* Crystal structures of agonist-bound human cannabinoid receptor CB1.
Nature **547**, 468-471 (2017).

42.   Hua, T.*, et al.* Crystal Structure of the Human Cannabinoid Receptor CB1. *Cell* **167**, 750-
762 e714 (2016).

43.   Krishna Kumar, K.*, et al.* Structure of a Signaling Cannabinoid Receptor 1-G Protein
Complex. *Cell* **176**, 448-458 e412 (2019).

44.   Li, X.*, et al.* Crystal Structure of the Human Cannabinoid Receptor CB2. *Cell* **176**, 459-467
e413 (2019).

45.   Shao, Z.*, et al.* Structure of an allosteric modulator bound to the CB1 cannabinoid receptor.
Nat Chem Biol **15**, 1199-1205 (2019).

46.   Shao, Z.*, et al.* High-resolution crystal structure of the human CB1 cannabinoid receptor.
Nature (2016).

47.   Masuho, I.*, et al.* Distinct profiles of functional discrimination among G proteins determine

the actions of G protein-coupled receptors. *Sci Signal* **8**, ra123 (2015).

48. Hollins, B., Kuravi, S., Digby, G.J. & Lambert, N.A. The c-terminus of GRK3 indicates rapid

dissociation of G protein heterotrimers. *Cell Signal* **21**, 1015-1021 (2009).

49. Kroeze, W.K.*, et al.* PRESTO-Tango as an open-source resource for interrogation of the

druggable human GPCRome. *Nat Struct Mol Biol* **22**, 362-369 (2015).

50. Kim, S.H., Won, S.J., Mao, X.O., Jin, K. & Greenberg, D.A. Molecular mechanisms of

cannabinoid protection from neuronal excitotoxicity. *Mol Pharmacol* **69**, 691-696 (2006).

51. Wu, J.*, et al.* Genetically Encoded Glutamate Indicators with Altered Color and Topology.

ACS Chem Biol **13**, 1832-1837 (2018).

52. Alger, B.E. Retrograde signaling in the regulation of synaptic transmission: focus on

endocannabinoids. *Prog Neurobiol* **68**, 247-286 (2002).

53. Ogasawara, D.*, et al.* Rapid and profound rewiring of brain lipid signaling networks by

acute diacylglycerol lipase inhibition. *Proc Natl Acad Sci U S A* **113**, 26-33 (2016).

54. Long, J.Z.*, et al.* Selective blockade of 2-arachidonoylglycerol hydrolysis produces

cannabinoid behavioral effects. *Nat Chem Biol* **5**, 37-44 (2009).

55. Mor, M.*, et al.* Cyclohexylcarbamic acid 3'- or 4'-substituted biphenyl-3-yl esters as fatty

acid amide hydrolase inhibitors: synthesis, quantitative structure-activity relationships, and

molecular modeling studies. *J Med Chem* **47**, 4998-5008 (2004).

56. Brenowitz, S.D. & Regehr, W.G. Associative short-term synaptic plasticity mediated by

endocannabinoids. *Neuron* **45**, 419-431 (2005).

57. Soler-Llavina, G.J. & Sabatini, B.L. Synapse-specific plasticity and compartmentalized

signaling in cerebellar stellate cells. *Nat Neurosci* **9**, 798-806 (2006).

58. Lerner, T.N. & Kreitzer, A.C. RGS4 is required for dopaminergic control of striatal LTD and

susceptibility to parkinsonian motor deficits. *Neuron* **73**, 347-359 (2012).

59. Gerdeman, G.L., Ronesi, J. & Lovinger, D.M. Postsynaptic endocannabinoid release is

critical to long-term depression in the striatum. *Nat Neurosci* **5**, 446-451 (2002).

60. Kreitzer, A.C. & Malenka, R.C. Endocannabinoid-mediated rescue of striatal LTD and

motor deficits in Parkinson's disease models. *Nature* **445**, 643-647 (2007).

61. Yasuda, H., Huang, Y. & Tsumoto, T. Regulation of excitability and plasticity by

endocannabinoids and PKA in developing hippocampus. *Proc Natl Acad Sci U S A* **105**, 3106-

3111 (2008).

62. Edwards, D.A., Zhang, L. & Alger, B.E. Metaplastic control of the endocannabinoid system

at inhibitory synapses in hippocampus. *Proc Natl Acad Sci U S A* **105**, 8142-8147 (2008).

63. Li, B. Central amygdala cells for learning and expressing aversive emotional memories.

Curr Opin Behav Sci **26**, 40-45 (2019).

64. Katona, I.*, et al.* Distribution of CB1 cannabinoid receptors in the amygdala and their role

in the control of GABAergic transmission. *J Neurosci* **21**, 9506-9518 (2001).

65. Morena, M., Patel, S., Bains, J.S. & Hill, M.N. Neurobiological Interactions Between Stress

and the Endocannabinoid System. *Neuropsychopharmacology* **41**, 80-102 (2016).

66. Gunduz-Cinar, O., Hill, M.N., McEwen, B.S. & Holmes, A. Amygdala FAAH and

anandamide: mediating protection and recovery from stress. *Trends Pharmacol Sci* **34**, 637-

644 (2013).

67. Dana, H.*, et al.* Sensitive red protein calcium indicators for imaging neural activity. *Elife*

5(2016).

68.   Soltesz, I.*, et al.* Weeding out bad waves: towards selective cannabinoid circuit control in
epilepsy. *Nat Rev Neurosci* **16**, 264-277 (2015).

69.   Farrell, J.S.*, et al.* In vivo assessment of mechanisms underlying the neurovascular basis
of postictal amnesia. *Sci Rep* **10**, 14992 (2020).

70.   Heinbockel, T.*, et al.* Endocannabinoid signaling dynamics probed with optical tools. *J*
Neurosci **25**, 9449-9459 (2005).

71.   Ju, N.*, et al.* Spatiotemporal functional organization of excitatory synaptic inputs onto
macaque V1 neurons. *Nat Commun* **11**, 697 (2020).

72.   Sethuramanujam, S.*, et al.* Rapid 'multi-directed' cholinergic transmission at central
synapses. 2020.2004.2018.048330 (2020).

73.   Kaifosh, P., Lovett-Barron, M., Turi, G.F., Reardon, T.R. & Losonczy, A. Septo-hippocampal
GABAergic signaling across multiple modalities in awake mice. *Nat Neurosci* **16**, 1182-1184
(2013).

74.   Lovett-Barron, M.*, et al.* Dendritic inhibition in the hippocampus supports fear learning.
Science **343**, 857-863 (2014).

75.   Farrell, J.S.*, et al.* Postictal behavioural impairments are due to a severe prolonged
hypoperfusion/hypoxia event that is COX-2 dependent. *Elife* **5**(2016).

76.   Kaifosh, P., Zaremba, J.D., Danielson, N.B. & Losonczy, A. SIMA: Python software for
analysis of dynamic fluorescence imaging data. *Front Neuroinform* **8**, 80 (2014).

Fig. 1 | Development and characterization of GRAB$_{eCB}$ sensors in HEK293T cells

Fig. 2 | Characterization of GRAB$_{eCB}$ in primary cultured neurons

Fig. 3 | Imaging endogenously released eCBs in primary cultured neurons

Fig. 4 | Detection of eCBs in acute brain slices

Fig. 5 | Detection of foot shock-evoked eCB signals in the mouse basolateral amygdala

Fig. 6 | Imaging eCB dynamics in mouse hippocampus during running and seizure

a

Optimization on cpEGFP

eCB1.0 —(H18C, L83F, S127G, E136V)→ eCB1.5 —

eCBmut ←(F177^2.64A)— eCB2.0 ←(S383^7.39T)—

Optimization on CB1 ligand binding pocket

b

180°

F83, V136, N16, C18, I225, D199

F83, E208, V136, I225, G127

c

TM7, TM1, TM6, TM5, TM4, TM3, TM2

S383^7.39

AEA

V196^3.32

F177^2.64

d

EC_50

- ■ eCB2.0 — 10 μM
- ■ eCB1.5 — 8 μM
- ■ eCB1.0 — 3 μM
- ■ eCBmut

Norm. $\Delta F/F_0$

[2-AG] (LogM)

Extended Data Fig. 1 | Evolution of GRAB_eCB sensors during the screening

Extended Data Fig. 2 | Full amino acid sequences of eCB2.0 and eCBmut

a

Norm. $\Delta F/F_0$

1.5

0.5

EC$_{50}$: ~0.5 µM

-9 -8 -7 -6 -5 -4
[WIN55212-2] (LogM)

b

Norm. $\Delta F/F_0$

1.5

0.5

EC$_{50}$: ~0.02 µM

-10 -8 -6 -4
[CP55940] (LogM)

Extended Data Fig. 3 | eCB2.0 responses to synthetic CB1R agonists

Extended Data Fig. 4 | 2P imaging of eCB2.0 in acute mouse striatal slices

	FIGURE LEGENDS

Fig. 1 | Development and characterization of GRAB$_{eCB}$ sensors in HEK293T cells

a, Schematic diagrams depicting the design and principle of the GRAB$_{eCB}$ sensor.

b, Screening steps of GRAB$_{eCB}$ sensors and fluorescence responses to 10 µM 2-AG of

GRAB$_{eCB}$ candidates in HEK293T cells. eCBmut was generated by introducing F177$^{2.64}$A

to eCB2.0.

c, Expression and fluorescence response to 100 µM 2-AG and AEA of eCB2.0 in HEK293T

cells.

d, Dosage dependent responses of eCB2.0 to 2-AG and AEA in HEK293T cells. EC$_{50}$s of

2-AG and AEA were labelled. n = 3 wells.

e, Responses of eCB2.0 to eCBs and varied neurotransmitters and neuromodulators. Note

the responses to eCBs were blocked by CB1R inverse agonist AM251. LPA,

lysophosphatidic acid; S1P, sphingosine-1-phosphate; ACh, acetylcholine; DA, dopamine;

GABA, gamma-aminobutyric acid; Glu, glutamate; Gly, glycine; NE, norepinephrine; 5-HT,

5-hydroxytryptamine; His, histamine; Epi, epinephrine; Ado, adenosine; Tyr, tyramine. n =

3-4 wells.

f, Illustration of a local puff system with a glass pipette filled with 100 µM 2-AG or AM251

placed close to an eCB2.0-expressing cell. Black dash line indicates the scanning region

of interest.

g, Line scanning fluorescence responses (left) and summary of on/off time constants (right)

of eCB2.0 to 2-AG or AM251. n = 11 (or 4) cells for rise (or decay) kinetics measurement.

h, G protein coupling of CB1R and GRAB$_{eCB}$ sensors measured using a BRET G$_{\beta\gamma}$ sensor.

n = 3 repeats.

i, β-arrestin coupling of CB1R and GRAB$_{eCB}$ sensors measured using the Tango assay. n

= 3 wells.

Values with error bars indicate mean ± SEM. Student's t test performed; ***$p < 0.001$. Scale

bar in (c) and (h), 30 µm.

Fig. 2 | Characterization of GRAB$_{eCB}$ in primary cultured neurons

a, Localization of eCB2.0 in primary cultured rat cortical neurons. eCB2.0 displayed well in

neuronal membrane of axons (indicated by arrow-labelled synaptophysin-mScarlet),

dendrites and spines (indicated by arrowhead-labelled psd95-mScarlet).

b, Expression and fluorescence response to 100 μM 2-AG and AEA of eCB2.0 and eCBmut.

Insets on eCBmut images are contrast-enhanced to indicate the expression of the sensor.

c, Dose dependent responses of eCB2.0 and eCBmut in neurites to 2-AG and AEA. n = 3

cultures.

d, Quantification of eCB2.0 responses to 100 μM 2-AG and AEA in neuronal somata and

neurites. n = 3 cultures.

e, Images, trace and quantification of eCB2.0 fluorescence to long-term WIN55212-2

application in neurons. n = 3 cultures.

Values with error bars indicate mean ± SEM. Student's t test and one-way ANOVA test

performed; ***$p < 0.001$. Scale bar in (a), 30 μm (upper) and 15 μm (lower); in (b), 30 μm;

in (e), 100 μm.

Fig. 3 | Imaging endogenously released eCBs in primary cultured neurons

a, Expression of R^{ncp}-iGluSnFR and eCB2.0 in neurons, pseudocolor images, and fluorescence traces showing the responses of R-iGluSnFR and eCB2.0 to 100 electrical pulses. 100 white circles (12.4 μm in diameter) indicate ROIs for quantification.

b, Loading of Calbryte590 and expression of eCB2.0 in neurons, pseudocolor images, and fluorescence traces showing the responses of Calbryte590 and eCB2.0 to 100 electrical pulses. 100 white circles (12.4 μm in diameter) indicate ROIs for quantification.

c, Relationship between peak Calbryte590 signals and peak eCB2.0 signals, evoked by 1, 5, 10, 20 electrical pulses and 20 electrical pulses without extracellular Ca^{2+}. Data were normalized to signals evoked by 200 electrical pulses at 50 Hz. n = 4 cultures.

d, Diagram showing the eCB synthesis pathway and the pharmacological perturbation.

e, Representative traces showing the eCB2.0 responses to 20 electrical pulses stimuli before (1) and after DO34 incubation (2), as well as the response to WIN55212-2.

f, Quantification of peak eCB2.0 responses to 20 pulses stimuli before (Ctrl) and after 26 min DO34 incubation (DO), as well as the response to WIN55212-2. n = 3 cultures.

g, Diagram showing the eCB degradation pathway and the pharmacological perturbation.

h, Representative traces showing the eCB2.0 responses to 20 electrical pulses stimuli before (1) and after JZL184 or URB597 incubation (2), as well as after the AM251 incubation.

i, Quantification of decay time constant before (Ctrl) and after 68 min JZL184 (or URB597) incubation (JZL or URB). n = 3 cultures.

j, Pseudocolor images of spontaneous eCB transients, single pulse stimulus evoked eCB signals, and 10μM WIN55212-2 induced eCB2.0 signals.

k, Pseudocolor images of a representative eCB transient event.

l, Traces of the single pulse stimulus evoked eCB signal, spontaneous eCB transient and eCB2.0 signal in the presence of AM251 shown in **k**. Normalized traces with rise time constants are shown on the right.

m, Spatial profile of the eCB transient shown in l (left). Quantification of eCB spontaneous transients FWHM is shown on the right. n = 12 eCB transients.

n, Cumulative eCB transients during 19 min recording in the absence (left) or presence of AM251 (right). Pseudocolor images were calculated as maximum temporal projection subtracting average temporal projection over 19 min.

o, Quantification of eCB transient frequency before (Ctrl) and after AM251 incubation (AM). n = 5 (Ctrl) or 3 (AM) x 10 min.

Values with error bars indicate mean ± SEM. Student's t test performed; *p < 0.05, ***p < 0.001. Scale bar in (a) and (b), 200 μm; in (j) and (n), 100 μm; in (k), 10 μm.

Fig. 4 | Detection of eCBs in acute brain slices

a, Schematic diagrams depicting the viral injection and expression in DLS. Acute slices are

prepared for electrical stimulation and photometry recording. Dashed box corresponds to

the image in b.

b, The expression of eCB2.0 in DLS and cartoons showing the electrical stimulation and

photometry recording.

c, Representative traces showing eCB2.0 fluorescence ($\Delta F/F_0$) increase evoked by

electrical stimulation with different stimulation pulses and frequencies.

d, Quantification of peak responses, rise and decay kinetics of electrical stimulation-

evoked eCB2.0 signals. n = 6 slices.

e, Representative traces and quantification of peak responses showing eCB2.0

fluorescence ($\Delta F/F_0$) increase evoked by electrical stimulation in control group and in

AM251 pre-incubated group. n = 3-4 slices.

f, Schematic diagrams depicting the viral injection and expression in CA1, hippocampus.

Acute slices were prepared for stimulation and 2 photon imaging.

g, Expression of eCB2.0 in CA1 and placement of the electrode, and pseudocolor images

representing the electrical stimulation evoked eCB2.0 response.

h, Representative traces and quantification of peak responses showing eCB2.0

fluorescence ($\Delta F/F_0$) increase evoked by electrical stimulation with varied stimulation

frequencies. n = 5 slices.

i, eCB2.0 fluorescence signal ($\Delta F/F_0$) when bath applying AEA and AM251.

j, Traces showing eCB2.0 fluorescence ($\Delta F/F_0$) increase evoked by electrical stimulation

before and after applying AM251.

Values with error bars indicate mean ± SEM. Scale bar in (b), 1 mm; in (g), 100 µm.

Fig. 5 | Detection of foot shock-evoked eCB signals in the mouse basolateral
amygdala
a, Schematic diagrams depicting the viral expression, fiber photometry recording and
behavior paradigm of the foot shock experiment.
b, Expression of eCB2.0 (green) and mCherry (red) in the BLA and placement of the
recording; the nuclei were counterstained with DAPI (blue).
c, Representative single trial traces showing eCB2.0 and mCherry fluorescence ($\Delta F/F_0$)
during an electrical foot shock.
d, Pseudocolor fluorescence responses of eCB2.0 to the foot shock. Five consecutive trials
from one mouse were recorded and plotted relative to the onset of each stimulus.
e, Averaged traces showing eCB2.0 and mCherry (top) or eCBmut and mCherry (bottom)
fluorescence during an electrical foot shock (left). Group summary of peak responses is
shown on the right. n = 6 mice.
f, Quantification of rise and decay kinetics of foot shock-evoked eCB2.0 signals. n = 21 (or
18) trials from 6 animals for rise (or decay) kinetics quantification.
Values with error bars indicate mean ± SEM. Student's t test performed; ***$p < 0.001$. Scale
bar in (b), 300 μm.
969

Fig. 6 | Imaging eCB dynamics in mouse hippocampus during running and seizure

a, Schematic diagrams depicting the viral expression and cannula placement.

b, Histology showing the expression of eCB2.0 in CA1 (left) and *in vivo* 2-photon image of pyramidal layer (right) showing the expression of eCB2.0 (green) and jRGECO1a (red).

c, Schematic illustration of the experiment in which mice expressing eCB2.0 and jRGECO1a in CA1 are placed on a treadmill. The fluorescence is recorded using 2-photon microscope.

d, Averaged traces of calcium and eCB transients recorded from somata of individual neurons in the pyramidal layer (left). Plots show the event-triggered average upon the start and stop of spontaneous running episodes (dashed lines).

e, Group summary of peak responses in panel **d**. n = 8 mice for eCB2.0 group and 4 mice for eCBmut group.

f, Quantification data of rise and decay kinetics of jRGECO1a and eCB2.0 signals during running start and stop.

g, Schematic illustration of the electrode placement and experiment in which mice expressing eCB2.0 and jRGECO1a in CA1 are recorded when inducing a kindling seizure.

h, Example trace of LFP, with medio-lateral projections (line profile) of eCB2.0 and jRGECO1a fluorescence during electrically induced non-convulsive seizures and the following spreading wave. Dashed line indicates the stimulus onset.

i, Individual and averaged traces of eCB2.0 (or eCBmut) and jRGECO1a fluorescence signals during the seizure. Dashed line indicates the stimulus onset. Group summary of AUC is shown on the right. n = 8 mice for eCB2.0 group and 4 mice for eCBmut group.

j, Spreading of eCB2.0 signal after a seizure. ROIs representing individual neurons are pseudo-colored according to the peak time of eCB2.0 signal relative to the peak time of the average signal (left). The symbols stand for (clockwise): anterior, lateral, posterior, medial.

k, eCB2.0 traces of individual cells sampled systematically along a line fitted to the spreading wave (arrow). The dashed line shows the spreading of peak signals.

l, Velocity and direction of the spreading eCB and calcium waves. Colored lines show individual sessions. Black lines show the average. n = 7 sessions from 6 mice.

Values with error bars indicate mean ± SEM. Student's t test performed; *p < 0.05, ***p < 0.001, and n.s., not significant. Scale bar in (b, from left to right), 200, 50 and 50 µm.

Extended Data Fig. 1 | Evolution of GRAB$_{eCB}$ sensors during the screening

a, Schematic representation of the evolution of GRAB$_{eCB}$ sensors.

b, 8 sites selected for optimization on cpEGFP.

c, 3 sites selected for optimization on the GPCR ligand binding pocket.

d, Normalized dose-dependent responses of eCB1.0, eCB1.5, eCB2.0 and eCBmut to 2-AG in HEK293T cells. n = 3 wells.

Values with error bars indicate mean ± SEM.

Extended Data Fig. 2 | Full amino acid sequences of eCB2.0 and eCBmut

a, Schematic representation of the eCB2.0 structure.

b, Amino acids sequence of the eCB2.0. 177$^{2.64}$ (labelled by the gray box) is mutated to A in eCBmut.

Extended Data Fig. 3 | eCB2.0 responses to synthetic CB1R agonists

a, Normalized dose-dependent responses of eCB2.0 to WIN55212-2.

b, Normalized dose-dependent responses of eCB2.0 to CP55940.

Extended Data Fig. 4 | 2P imaging of eCB2.0 in acute mouse striatal slices

a, 2P images of eCB2.0 (indicated by arrows) expressed in mouse striatum in saline, 10 μM AEA and 10 μM AM251.

b, eCB2.0 fluorescence signal ($\Delta F/F_0$) when bath applying AEA and AM251.

Scale bar, 10 μm.

Extended Data Movie 1 | eCB and calcium signals in mouse hippocampal CA1 during kindling seizure

2P imaging of eCB and calcium signals using eCB2.0 and jRGECO1a in mouse hippocampal CA1 during kindling seizure. LFP was simultaneously recorded to indicate the electrical stimulation and seizure.

Scale bar, 50 μm for images and 1 V for LFP.