1

**Atypical molecular basis for drug resistance to mitochondrial function inhibitors**

**in *Plasmodium falciparum***

Heather J. Painter[1,a], Joanne M. Morrisey[2], Michael W. Mather[2], Lindsey M. Orchard[1], Cuyler

Luck[1], Martin J. Smilkstein[3], Michael K. Riscoe[3], Akhil B. Vaidya[2], Manuel Llinás[1]

[1] Department of Biochemistry and Molecular Biology, Huck Center for Malaria Research,

Pennsylvania State University, University Park, PA, USA

[2] Center for Molecular Parasitology, Department of Microbiology and Immunology, Drexel

University College of Medicine, Philadelphia, PA, USA

[4] Medical Research Service, Department of Veterans Affairs Medical Center, Portland, OR, USA

[a] Current affiliation: United States Food & Drug Administration, Center for Biologics &

Evaluation Research, Office of Vaccine Research & Review, Division of Parasites, Bacteria &

Allergenic Products, Silver Spring, MD, USA

[#] Corresponding authors: manuel@psu.edu and Heather.Painter@fda.hhs.gov

**Abstract:**

The continued emergence of drug-resistant *Plasmodium falciparum* parasites hinders global attempts to eradicate malaria, emphasizing the need to identify new antimalarial drugs. Attractive targets for chemotherapeutic intervention are the cytochrome (cyt) $bc_1$ complex, which is an essential component of the mitochondrial electron transport chain (mtETC) necessary for ubiquinone recycling and mitochondrially localized dihydroorotate dehydrogenase (DHODH) critical for *de novo* pyrimidine synthesis. Despite the essentiality of this complex, resistance to a novel acridone class of compounds targeting cyt $bc_1$ was readily attained, resulting in a parasite strain (SB1-A6) that was pan-resistant to both mtETC and DHODH inhibitors. Here we describe the molecular mechanism behind the resistance of the SB1-A6 parasite line which lacks the common cyt $bc_1$ point mutations characteristic of resistance to mtETC inhibitors. Using Illumina whole-genome sequencing, we have identified both a copy number variation (~2x) and a single-nucleotide polymorphism (C276F) associated with *pfdhodh* in SB1-A6. We have characterized the role of both genetic lesions by mimicking the copy number variation via episomal expression of *pfdhodh* and introducing the identified SNP using CRISPR/Cas9 and assessed their contributions to drug resistance. Although both of these genetic polymorphisms have been previously identified as contributing to both DSM-1 (1) and atovaquone resistance (2, 3), SB1-A6 represents a unique genotype in which both alterations are present in a single line, suggesting that the combination contributes to the pan-resistant phenotype. This novel mechanism of resistance to mtETC inhibition has critical implications for the development of future drugs targeting the $bc_1$ complex or *de novo* pyrimidine synthesis that could help guide future anti-malarial combination therapies and reduce the rapid development of drug resistance in the field.

## Introduction:

Infection by the human malaria parasite *Plasmodium falciparum* rapidly leads to the clinical symptoms associated with the 48-hour asexual replicative cycle of the parasite within the blood of the host (4). *P. falciparum* malaria continues to present an enormous global public health burden due to the lack of an effective long-term vaccine (5) and the emergence of resistance to front-line antimalarial chemotherapies (6). This underscores the continued need to maintain a robust drug development pipeline that produces novel and effective antimalarial compounds for future deployment.

To accelerate novel anti-malarial compound identification, independent laboratories and global consortia have rapidly identified numerous chemical entities that are efficacious against multiple *Plasmodium* species and developmental stages. Despite the diverse chemical scaffolds in the compounds tested, most effective candidates tend to have similar modes-of-action in the malaria parasite due to the paucity of molecular targets that are significantly divergent from the human host (7-9). As such, it is not surprising that a large and diverse set of compounds selectively target the highly divergent *Plasmodium* mitochondrion (7), which is essential for *de novo* pyrimidine synthesis during the blood stages of parasite development (10, 11).

Although the *Plasmodium* mitochondrion maintains the canonical tricarboxylic acid cycle pathway of central carbon metabolism (12-15), these enzymes are dispensable for blood-stage development (15). We have previously shown the main role of the parasite mitochondrion during the blood-stage is the provision of precursors for *de novo* pyrimidine synthesis (10), a process requiring a ubiquinone-dependent, mitochondrional dihydroorotate dehydrogenase (DHODH, PF3D7_0603300) and a functioning electron transport chain (ETC) to maintain turnover of

ubiquinol, regenerating ubiquinone. This sole essential function of the mitochondrion was

established by genetic supplementation with an alternative DHODH from *S. cerevisiae*

(yDHODH), which is localized to the cytoplasm and utilizes fumarate as an electron acceptor

rather than ubiquinone (10). Therefore, parasites expressing yDHODH from an episome,

bypassing the mitochondrially localized PfDHODH, are highly resistant to all chemical inhibitors

of the mtETC. Interestingly, the addition of the biguanide prodrug proguanil, combined with the

mtETC cytochrome *b* inhibitor atovaquone in the antimalarial drug Malarone, reversed the pan-

resistant phenotype of yDHODH expressing *P. falciparum* strain D10 (D10::yDHODH) (10).

Although the mode-of-action of atovaquone, as well as other mtETC inhibitors, is well-understood,

the molecular target of proguanil remains unknown.

Due to the efficacy of disrupting mitochondrial function in the malaria parasite as a strategy for

chemotherapeutic intervention, these pathways are the focus of comprehensive anti-mitochondrial

drug development programs, and significant efforts have been made to identify novel compounds

that target the mtETC both directly (cytochrome *b*) and indirectly (DHODH). To this end, there

are several antimalarials, some of which are in the advanced stages of clinical trials, whose

proposed modes-of action include interaction with the ubiquinone binding pocket(s) of DHODH

(DSM-1 and DSM265) (16-18) or cytochrome *b* (ELQ-300 and 6-NH$_2$Ac) (19, 20). Despite the

essentiality of both components of the mtETC, resistance to these compounds readily arose and

genotyping of resistant parasites revealed either amplification of or point mutations in DHODH,

as well as point mutations within cytochrome *b* (1, 2, 16, 21-23). However, parasites resistant to

6-NH$_2$Ac, an acridone compound thought to target the mtETC *bc$_1$* complex (20), did not exhibit

traditional *bc$_1$* resistance mutations (21). The acridone-resistant parasite line SB1-A6 was not only

resistant to 6-NH$_2$Ac but also exhibited pan-resistance to all mitochondrial inhibitors to which the

parent strain, D6, was susceptible (21).  Again, this resistance was abrogated by the addition of

proguanil (21), similar to *P. falciparum* parasites that express yDHODH, and confirming the target

of acridones to be the *P. falciparum* mtETC, although the provenance of the phenotypic similarities

in resistance to $bc_1$ complex inhibitors between SB1-A6 and the yDHODH transgenic parasite line

was unclear (10).

In this study, we present a clear genotype for the *P. falciparum* SB1-A6 acridone-resistant clonal

parasite strain and, through a combination of targeted and whole-cell methods, establish that the

mechanism of resistance to both cytochrome $bc_1$ and DHODH inhibitors results from the

contribution of multiple genetic polymorphisms.  We find that *P. falciparum* SB1-A6 accumulates

both a copy number variation and a specific mutation in PfDHODH, and both of these genetic

polymorphisms contribute to the pan-resistant phenotype.  This study uncovers a mechanism of

cross-resistance between PfDHODH and mtETC inhibitors and serves as a cautionary note to

future anti-malarial combination therapy formulations containing such drugs.

**Results:**

**Strain SB1-A6 displays pan-resistance to mtETC inhibitors.**

To confirm the resistance phenotype of *P. falciparum* SB1-A6 (21) and the line's similarity to *P.*

*falciparum* D10::yDHODH (10), we first tested the susceptibility of both strains to the classic

malaria parasite cytochrome $bc_1$ inhibitor, atovaquone, using a growth inhibition assay (Fig. 1A,

Table S1).  As expected, both SB1-A6 and D10::yDHODH displayed resistance to atovaquone,

while the parental strains (D6 and D10, respectively) were susceptible (Fig. 1A. Table S1), as

previously published (21).  It has been well established that expression of yDHODH in *P.*

*falciparum* results in resistance to the PfDHODH inhibitor DSM-1 (24, 25).  Therefore, we tested

the susceptibility of SB1-A6 to direct inhibition of PfDHODH with DSM-1 (17) and found that

this parasite line was resistant to both DHODH, as well as cytochrome $bc_1$ inhibitors (Figure 1B,

Table S1),  making it the first *P. falciparum* clone generated to exhibit such cross-resistance (21).

However, more recent studies aimed at generating resistance to atovaquone by increasing drug

concentration exposure step-wise have also resulted in cross-resistance to DSM-1 (2).  Resistance

to mtETC inhibitors due to an alternative source of pyrimidines is abolished by the addition of the

prodrug proguanil (10). Likewise, resistance to atovaquone inhibition of the mtETC in SB1-A6

was reversed upon the addition of 1.5µM proguanil (Fig. 1B, Table S1), suggesting that the

mechanism of pan-resistance to mitochondrial inhibitors is linked to the function of PfDHODH,

as was previously demonstrated for D10::yDHODH (10).

Drug resistance mechanisms in both prokaryotes and eukaryotes can involve "rewiring" to

overcome the inhibition of essential metabolic pathways by genetic variation, metabolic bypass,

or efflux mechanisms.  Metabolomics has become a useful tool for unraveling anti-parasite drug

mechanisms of action and resistance (8, 26, 27). To elucidate the mechanism(s) of resistance in

SB1-A6, we determined if any metabolic perturbations occurred in the SB1-A6 drug-resistant line

due to drug treatment that could account for cross-resistance to both PfDHODH and cytochrome

$bc_1$ inhibition, using whole-cell metabolomics analysis.  Trophozoite-stage D6 and SB1-A6 were

exposed to ~10x IC50 of either atovaquone (10nM) or DSM-1 (1.5µM) for 2 hours, followed by

whole-cell metabolite extraction, and quantitated by LC-MS/MS analysis (8).  Metabolomic

analysis showed that treatment of SB1-A6 with DSM-1 or atovaquone did not result in any

significant metabolic perturbation (Fig. 1C and 1D; Table S1), whereas treatment of the wild-type

D6 revealed an accumulation of *de novo* pyrimidine synthesis precursors compared to an untreated

control (Fig 1C and 1D). The accumulation of pyrimidine precursors in the parental line upon

treatment with either atovaquone or DSM-1 and lack of accumulation in the treated mutant line

(Fig. 1D), in neither a stage- nor time-dependent manner (Fig. S1) is consistent with the

involvement of DHODH in the mechanism of resistance.

**Genetic determinants of pan-resistance to mtETC and pyrimidine synthesis inhibition in**

**SB1-A6.**

Previous studies have demonstrated that genetic mechanisms of resistance resulting in the ability

of the parasite to overcome mtETC and *de novo* pyrimidine synthesis inhibition include mutations

in either *cytochrome b* (28, 29) or *pfdhodh* (1, 30) or copy number variation of *pfdhodh* (2, 3).

Analysis of the SB1-A6 *cytochrome b* sequence by Smilkstein *et al.* (21) did not reveal any

mutations. Therefore, to identify any genetic polymorphisms that could lead to a molecular

mechanism of resistance, we examined the genomes of both the D6 wild-type and SB1-A6 drug-

resistant parasites using Illumina-based next-generation DNA sequencing. Whole-genome

sequencing revealed a non-synonymous G → T point-mutation in the coding sequence of

PfDHODH, resulting in the amino acid change C276F (Fig. 2A, Table S2). Amino acid C276 is

localized adjacent to the flavin mononucleotide binding pocket (Fig. 2C), and the substitution to a

phenylalanine has been demonstrated to reduce the size of the binding pocket where DSM-1 binds

(1), resulting in a binding pocket structure resembling that of HsDHODH (30). Although this

mutation was not the only change found in SB1-A6 (Table S2), it was the only change in an

annotated gene known to be directly linked to pyrimidine synthesis and/or mitochondrial function.

In addition, we identified a ~20kb, 2-fold copy number variation (CNV) in the SB1-A6 genome,

including the coding sequence for PfDHODH, as well as six other protein-coding genes (Fig. 2B).

We further confirmed the presence of this copy number increase by DNA microarray comparative

genome hybridization (Table S3). This region has previously been shown to increase in copy number in *P. falciparum* selected for resistance to DSM-1 (3), resulting in cross-resistance to atovaquone (2). Although each of these genetic polymorphisms has individually been previously identified as contributing significantly to both DSM-1 (1) and atovaquone resistance (2, 3), SB1-A6 possesses a unique genotype in which both alterations are present in a single parasite line, suggesting that the combination contributes to the pan-resistant phenotype of this line. Intriguingly, the increased CNV of PfDHODH did not directly translate to higher protein levels (Fig. 2D), nor did it produce any obvious mislocalization (Fig. S2) of the protein in the parasite.

Due to previous observations that linked the PfDHODH C276F mutation and CNV to both DSM-1 and atovaquone resistance, respectively (2, 3), we examined the function of PfDHODH in mitochondria isolated from D6 and SB1-A6 to determine if these polymorphisms could account for the reduced sensitivity of SB1-A6 parasites to mitochondrial inhibition. Measurement of DHODH activity in isolated mitochondria from trophozoite stages of D6 and SB1-A6 revealed that the enzymes of both strains are equally susceptible to inhibition by DSM-1 (Fig. 2E), confirming the results from purified wild-type and C276F mutant PfDHODH activity assays (1). Interestingly, measurement of the specific activity of DHODH from SB1-A6 isolated mitochondria revealed a two-fold increase compared to D6 (Fig. 2F), possibly due to the increase in copy number of PfDHODH. Although the C276F mutation does not lead to a decrease in the ability of DSM-1 to inhibit PfDHODH, these data suggest that the SB1-A6 parasite line has gained the ability to withstand either mtETC or pyrimidine biosynthesis inhibition by increasing the overall activity of PfDHODH allowing sufficient levels of nucleotide synthesis for parasite survival.

**Genetic supplementation of DHODH increase copy number and CRISPR-Cas9 introduction of C276F point mutation**

Previous studies have established that *P. falciparum* grown in the presence of step-wise increasing

concentrations of DSM-1 leads to an increase in copy number of a chromosomal region that

includes PfDHODH leading to resistance to both DSM-1 (3) and atovaquone (2).  To test the

hypothesis that a copy number increase in PfDHODH also results in decreased sensitivity of SB1-

A6 parasites to inhibition of electron transport or *de novo* pyrimidine synthesis, we introduced

either wild-type or mutant (C276F) PfDHODH expressed from an episomal vector under the

control of a constitutive promoter in the drug-susceptible strain D6 (Fig. 3A).  After selection of a

parasite population containing the mutant or wild-type PfDHODH, we assessed the susceptibility

of these parasites to atovaquone or DSM-1 using a standard growth inhibition assay.   The

introduction of additional copies of PfDHODH resulted in decreased susceptibility to DSM-1 and

atovaquone (Fig. 3B).  Notably, the episomal expression of the mutant PfDHODH also results in

almost complete resistance to atovaquone (Fig. 3B), phenocopying the genotype of SB1-A6.

Episomal increase of wild-type PfDHODH results in a ten to 100-fold decrease in susceptibility to

DSM-1 (Fig. 3B), confirming the phenotype established by Guler *et al.* of DSM-1 and atovaquone

resistant parasites with an increased copy number of PfDHODH (2, 3).

Since the discovery that *P. falciparum* survival is dependent upon *de novo* pyrimidine synthesis

(10), PfDHODH has become a popular target for chemical intervention strategies (17, 18, 31, 32).

For example, discovery efforts have led to the development of triazolopyrimidine-based inhibitors

for PfDHODH, which are currently undergoing clinical trials for safety and efficacy (16, 33).  *In*

*vitro* selection of *P. falciparum* for resistance to compound DSM265 led to the identification of

various mutations within PfDHODH (1).  Of these mutations, C276F was determined to reduce

the binding capacity of chemical inhibitors using in an *in vitro* expression system (1), as well as

computational docking studies (30). To test the hypothesis that the same observed point mutation

in SB1-A6 PfDHODH contributes to *P. falciparum* resistance to both PfDHODH and cytochrome

$bc_1$ inhibitors, we utilized CRISPR/Cas9 (34) to genetically introduce the C276F mutations into

this gene in D6 and also repair it in SB1-A6 (Fig. 3C; Fig S3A and S3B). After confirming the

introduction or repair of the mutation at amino acid 276 (Fig. S3A and S3B), we then performed

drug susceptibility assays to quantitate the contribution of this mutation to the pan-resistant

phenotype (Table S1). While we were able to reverse this mutation to the wild-type background

in A6$^{F276C}$ decreasing the overall resistance to both atovaquone and DSM-1 (Fig. 3D, Table S1),

there remains an increased CNV of PfDHODH, which is likely contributing to the higher IC50

than demonstrated for D6 (Fig. 3D; Table S1). Conversely, susceptibility to atovaquone is only

decreased by 2-fold due to the introduction of C276F into strain D6 (Fig. 3D; Table S1).

Interestingly, these assays show that this mutation results in a 10-fold increase in the IC50 of DSM-

1 in D6$^{C276F}$ (Fig. 3D) and a 40-fold decrease in A6$^{F276C}$ (Fig. 3D). These results suggest that the

point mutation directly contributes to decreased efficacy of direct inhibition of PfDHODH and

enhances the ability of *P. falciparum* to overcome mtETC inhibition (Table S1). Taken together,

these data demonstrate the coordination between genetic polymorphisms of the PfDHODH SNP

and CNV in the SB1-A6 parasite line that mediates the pan-resistance phenotype.

**<u>Discussion:</u>**

In this study, we have demonstrated the coordination of two genetic polymorphisms that result in

resistance to compounds that target the mtETC. First, the copy number increase of DHODH results

in the reduced susceptibility to $bc_1$ complex and *de novo* pyrimidine synthesis inhibition.

However, a two-fold increase in DHODH, by itself, was not sufficient for complete resistance to

mtETC inhibitors, potentially due to a reliance on other genes within the CNV amplicon. Aside

from DHODH, this amplicon encompasses seven genes, a majority of which are unannotated,

except for PF3D7_0603200 (*bcs1*: mitochondrial chaperone, putative) and PF3D7_0602800 (*jmjc*:

jumanji-C demethylase domain containing), the gene products of which could contribute to the

drug-resistant phenotype. This is supported by previous studies that demonstrated that a similar

amplicon results in resistance to both atovaquone and DSM-1 (2, 3). Further work is necessary to

determine if the proteins encoded for on the amplicon are involved in reducing the parasite's

susceptibility to mtETC and *de novo* pyrimidine synthesis inhibition.

The implications of the SB1-A6 phenotype should be emphasized in considerations of the mtETC-

pyrimidine-biosynthesis axis as an anti-malarial drug target. While the adaptation that led to SB1-

A6 seems to have been a unique event, both the CNV and C276F point mutations have each been

derived independently from exposure to atovaquone and DSM-256, a next-generation PfDHODH

inhibitor currently in clinical trials. Therefore, the possibility that adaptations similar to those

found in SB1-A6 may arise during clinical trials or deployment of new $bc_1$ complex or DHODH

inhibitors deserves consideration. While concerns regarding atovaquone resistant mutations in the

*cytb* gene on mtDNA have been largely allayed due to lack of inheritance during transmission

within the mosquito vector (35), we have no reason to discount the transmissibility of nuclear-

encoded *pfdhodh* polymorphisms at present. Since these adaptations can be overcome by the

addition of proguanil, the continued inclusion of proguanil or an effective replacement should be

considered in future combinations. In addition to the question of transmissibility and choice of

partner drugs, the ultimate significance of the SB1-A6 phenotype to future anti-malaria drug

choices may also differ between treatment use and prophylaxis. During treatment parasite number

is high, increasing the risk for selection of both pre-existing and *de novo* resistant parasites,

whereas, in causal prophylactic use, in the absence of pre-existing resistance the risk would be

expected to be negligible. In either case, assessing the risk and monitoring for the emergence of
the SB1-A6 phenotype should be done.

This study presents the first identification of the acquisition of multiple genetic polymorphisms in
response to a $bc_1$ complex inhibitor that results in pan-resistance to both direct and indirect
inhibitors of DHODH. Fully elucidating the molecular mechanisms of this pan-resistance is
essential to inform future chemotherapeutic targeting strategies for malaria elimination. The
emergence of resistance to many of the drugs in the antimalarial arsenal has highlighted the
urgency of developing future medicines with modes of action distinct to those of the current
therapies. More recently, considerations have been made for combination therapies that target a
single essential metabolic pathway via different modes-of-action. In light of the potential
development of cross-resistance to a single inhibitor due to its combinatorial partner, the molecular
mechanisms of resistance for future therapeutic strategies should be thoroughly investigated prior
to approval for use in the clinical setting.

Recently, the essentiality of DHODH has been demonstrated in cell-cycle arrested cancer cells
without mitochondria wherein reactivation of growth occurs upon the acquisition of mtETC-linked
DHODH (36), suggesting that the lack of de novo pyrimidine synthesis is a major obstacle for in
vivo growth of respiration-compromised tumor cells. The combined detrimental effects of both
dysfunctional mitochondrial respiration and mtETC-linked de novo pyrimidine synthesis on cell
growth strengthen their relevance as drug targets against both parasite development and
metastatic cell growth. This further emphasizes the continued benefit of identifying novel
inhibitors of DHODH that could ultimately be appropriated for both parasite elimination and
perhaps extended to treatment targeting metastatic cells within the human host.

**Methods:**

***Parasite Strains and Culture:*** *Plasmodium falciparum* strains D6 and SB1-A6 were provided by

Michael Riscoe and Martin Smilkstein (21). All parasites obtained from other sources or generated

in this study were maintained under standard conditions (37) at 2% hematocrit with O+ human

erythrocytes in RPMI1640 (Thermo Fisher Scientific) containing hypoxanthine, $NaH_2CO_3$

(Sigma-Aldrich), HEPES (Sigma-Aldrich), glutamine (Sigma-Aldrich) and 2.5 g/L AlbuMAX II

(Thermo Fisher Scientific). Intraerythrocytic developing parasites were synchronized with 5%

sorbitol (Sigma-Aldrich) over three subsequent developmental cycles to synchronize parasites

within 4-6 hours, as previously described (38).

Parasite cultures were maintained Mycoplasma-free and were confirmed prior to metabolomic

analysis. Separate mycoplasma-free culture apparatus and solutions were used for all

metabolomics culturing to prevent contamination. Cultures were checked for mycoplasma

contamination weekly using IntronBio *e*-Myco™, Mycoplasma PCR Detection kit (Boca

Scientific; cat. no. 25235). If parasite cultures tested positive for Mycoplasma, cultures were

treated as per manufacturer's instructions with Mycoplasma Removal Agent (MP Biomedicals,

LLC.; cat. no. 30-500-44).

***Drug Inhibition Assays:*** All parasite growth inhibition assays were performed in 96-well plates

as described by Desjardin *et al.* (39) and adapted for SYBR Green I (40). Drugs were prepared in

DMSO as a 500× dilution series (typically a 2- or 3-fold dilution series was used, yielding a final

concentration in media ranging from 0.001−30 μM depending on the cell line) and were then

diluted 1:50 into media to yield a final DMSO concentration of 0.2%. Parasites were diluted to 1%

parasitemia and 1% hematocrit in each well of a 96-well plate then propagated for 48 h and growth

was assessed using the SYBR Green method.  SYBR green fluorescence was measured (ex./em.

485/535 nm) and data analysis was performed using GraphPad Prism 7.  For transgenic *P.*

*falciparum* strains generated in this study, parasites were propagated for three intraerythrocytic

cycles in media lacking drug before plating. All data were collected in triplicate and biological

duplicate.

***Mitochondrial Activity Assays:*** *P. falciparum* cultures were synchronized at least twice by

treatment with alanine (41), expanded and harvested at 8-15% parasitemia in the mid to late

trophozoite stages.  A hemozoin-depleted mitochondrial fraction was isolated from the parasites

using nitrogen cavitation, magnetic separation, and differential centrifugation, as previously

described (42).

DHODH activity assays were conducted essentially as previously described (18, 42) with minor

modifications, utilizing the ubiquinone analog decylubiquinone as the primary electron acceptor,

coupled to the redox dye 2,6-dichloroindophenol to take advantage of the latter's higher extinction

coefficient and absorbance in the visible range. Assays contained 1 mM L-dihydroorotate, 100 µM

decylubiquinone, 60 µM 2,6-dichloroindophenol, 50 mM sodium malonate, 2 mM potassium

cyanide, 100 mM potassium chloride, 0.05% Triton X-100, 10 % glycerol, and 0.1 M HEPES (pH

7.6) in a final volume of 1 ml. 5 ul aliquots of diluted inhibitor or vehicle (DMSO) was added, and

each assay was initiated by addition of a 7 µl aliquot of mitochondrial preparation (50-110 µg

protein) and recorded with a modified SLM-AMINCO DW2C dual-wavelength

spectrophotometer (On-Line Instrument Systems, Inc., Bogart, GA, USA) in dual mode (600 nm

– 528 nm) at 30°C.

***Inhibitors:*** The antimalaria compound atovaquone was a gift from Glaxo Wellcome, Research

Triangle Park, N.C. Proguanil was kindly provided by the Jacobus Pharmaceutical Company,

Princeton, NJ, USA. Both ELQ-300 and 6-NH2Ac were contributed to this study by Michael

Riscoe, Oregon Health Sciences University, Portland, OR, USA. DSM-1 (17) and DSM-265 (16)

were provided by Margaret Phillips, University of Texas Southwestern, Dallas, TX, USA.

Chloroquine (Cat# C6628) was purchased from Sigma-Aldrich, St. Louis, MO, USA.

***Whole-cell Metabolomics:*** Metabolite extractions, data collection, and analysis were performed

as described previously (8, 43, 44). Briefly, a ~20 µL pellet of cells was resuspended in 1 mL of

pre-chilled 90:10 methanol:water and placed at 4˚C. The internal standard $^{13}C_4$-$^{15}N_1$-aspartate was

spiked into the extraction methanol solution to control for sample preparation and handling.

Samples were vortexed, resuspended, and centrifuged for 10 min at 15000 RPM and 4˚C.

Supernatants were collected and stored at -80˚C or dried down immediately under nitrogen flow.

The dried metabolites were resuspended in HPLC-grade water (Sigma-Aldrich, CHROMASOLV)

to between $1x10^5$ and $1x10^6$ cells/µL, based on hemocytometer counts of purified parasites. All

samples were processed in technical triplicate with method blanks to reduce unwanted variation

and account for background signal, respectively. Samples were randomized and 10µL of

resuspended metabolite extract or method blank was injected for ultraHigh-Performance Liquid

Chromatography-Mass Spectrometry (HPLC-MS) analysis.

Extracts were analyzed using HPLC-MS on a Thermo Exactive Plus Orbitrap™. Metabolite

separation was performed with a C18 column (Phenomenex Hydro-RP; cat. No 00D-4387-B0)

using a 25 minute gradient of A: 3% aq. methanol/15mM acetic acid/10 mM tributylamine ion-

pairing agent and B: 100% MeOH (45). Metabolite detection was performed using a scan range

of 85-1000 m/z and a resolution of 140,000 @ m/z 200, in negative ion mode. The detection of

cellular metabolites was aided by the generation of a database from 292 pure metabolite standards

using the same instrument and method to determine detection capability, mass/charge ratio (m/z),

and retention time for each metabolite. Analysis of data was carried out as previously described

(8).

***Whole-genome Sequencing and Analysis:*** Genomic DNA was isolated, prepared from *P.*

*falciparum* parasite lines, and sequenced as previously described (46). Briefly, a total of 10 μg of

gDNA from each line was sheared to obtain a fragment size of ~200–400 bp. The resulting sheared

gDNA was size selected on a 2% (w/v) low-melting agarose gel and then purified and concentrated

using MinElute purification columns followed by the QIAquick PCR purification kit (QIAGEN).

Barcoded libraries for Illumina TruSeq single-end sequencing were then constructed from the size-

selected, sheared material and size selected using Agencourt AMPure XP magnetic beads

(Agencourt Biosciences, Beckman Coulter). The quality of the final sequencing libraries was

assessed using an Agilent 2100 Bioanalyzer (Agilent Technologies) and the concentration of each

library was quantified using a Quant-iT dsDNA Broad-Range Assay Kit (Invitrogen). The final

libraries were multiplexed and 20% (v/v) PhiX control DNA (Illumina, Catalog # FC-110-3001)

per lane and were sequenced using an Illumina HiSeq 2500 Rapid Run (150 bp) system (Illumina).

Sequencing outputs were uploaded into Galaxy (47), which is hosted locally at the Millennium

Science Complex at Pennsylvania State University.  Sequence reads were mapped to the *P.*

*falciparum* 3D7 reference genome v. 10.0 (http://plasmodb.org/common/downloads/release-

10.0/Pfalciparum/), using the Burrows-Wheeler alignment tool while selecting and filtering unique

reads for Map Quality >30 (48). Sequence variations were detected by Freebayes (version 0.9.0.a)

using stringent filtering parameters based on quality and read depth (49, 50).  Then SNPeff (version

3.3) was applied to annotate and determine the statistical significance of each variant (51). Genome

copy number variations were detected based upon local chromosomal read depth using CNVnator

(version 0.3) and annotated with Intansv (version 0.99.3) (52).

***DHODH Western Blot Analysis:*** Total protein extracts were run on a 4-12% Bis-Tris gel

(ThermoFisher) in MES buffer and then transferred to a nitrocellulose membrane. After blocking

with 5% milk in PBS-T for 30 minutes, membranes were incubated overnight with the primary

antibody in 5% milk. Primary antibodies used were: 1:300 dilution of mouse anti-PfDHODH #3

(a kind gift from Margeret Phillips, University of Texas Southwestern) or 1/5000 mouse anti-

*Pf*Aldolase-HRP (Abcam, Cat #ab38905). The next day, membranes were washed three times with

PBS-T and incubated for two hours with the appropriate secondary antibody, then washed again.

The secondary antibody that was used was 1/5,000 goat anti-mouse HRP conjugate (Pierce). ECL

reagent (Pierce) was used to detect bound antibody.

***DNA Constructs:*** To mutate DHODH using CRISPR-Cas9, complementary oligonucleotides

encoding the guide RNA were inserted into pDC2-U6A-hDHFR (kind gift of Marcus Lee) using

BbsI sites. A 515 bp homology region containing the mutation in PfDHODH (C276F) or the

wildtype sequence from either SB1-A6 or D6 was cloned into StrataClone (Agilent) and then

inserted using the NotI site of pDC2-U6A-hDHFR. The resulting plasmids are pDC2-C276F-

hDHFR and pDC2-F276C-hDHFR which were co-transfected with Cas9-BSD plasmid from Jose-

Juan Lopez-Rubio (34) into D6 and SB1-A6, respectively.

To introduce multiple copies of either the mutant or wild-type PfDHODH into D6 parasites, *Pf*

DHODH was amplified from SB1-A6 (mutant) or D6 (wild-type) genomic DNA including the

stop codon, cloned into StrataClone (Agilent), sequence-verified, and then inserted into pLN-

ENR-GFP (53) using the AvrII and BsiWI restriction sites.  The resulting plasmids, pLN-

A6DHODH was transfected into D6 parasites.

***Transfection of Plasmodium falciparum.***  Transfection of *P. falciparum* was performed as

previously described (54). Briefly, 5-7% ring stage parasite cultures were washed three times with

cytomix (55). The parasitized RBC pellet was resuspended to 50% hematocrit in cytomix.  The

plasmid, isolated using a Qiagen Maxi kit and stored in 50 μg aliquots in ethanol, was centrifuged

down and resuspended in 100 μL Cytomix.  The plasmid and 250 μL of the 50% parasitized RBC

suspension were combined and transferred to a 0.2 cm cuvette on ice.  Electroporations were

carried out using a BioRad GenePulser set at 0.31kV, 960uF.  The electroporated cells were

immediately transferred to a T-25 flask containing 0.2 ml uninfected 50% RBCs and 7 mL

medium.  To select for parasites containing plasmid, medium containing 5 nM WR99210 or 1.5

μg/mL Blasticidin was added at 48 hrs post-transfection.  Cultures were maintained under constant

5 nM WR92210 pressure, splitting weekly, until viable parasites were observed.  To confirm the

successful mutation of PfDHODH, genomic DNA was purified using the Qiagen DNeasy Blood

& Tissue kit and the region of interest was PCR amplified and Sanger sequenced. Positive

transfectants were cloned by limiting dilution to obtain a pure population of transgenic parasites.

**Author's contributions:**

H.P., M.M., J.M., S.C., and C.L. designed and performed experiments and analyzed data. H.P.

wrote the manuscript and generated figures. M.M., M.S., and M.R. provided reagents, critical

comments, and edits. M.L. and A.V. designed the experiments, provided critical comments, edits,

and oversight.

**Acknowledgments:** We would like to thank the many people who have contributed to this project

including Dr. Andrew D. Patterson and Dr. Philip B. Smith of the Penn State Huck Institutes of

the Life Sciences Metabolomics Core Facility and Dr. Simon Cobbold previously of the Llinás lab

at Princeton University, for analytical expertise, technical oversight; Dr. Craig Praul of the

Genomics Core Facility of the Penn State Huck Institutes of the Life Sciences; and the Genomics

Core Facility at Princeton University and its staff, especially Donna Storton and Jessica Buckles

Wiggins. We also thank Dr. Meg Phillips and Dr. Pradip Rathod for sharing reagents and critical

comments. Lastly, we thank the many past and present members of the Llinás lab who supported

this project by enhancing the evaluation of data and provided critical comments.

**Funding Information:**

This work was supported by generous funding awarded to M.L. from the Bill and Melinda Gates

Foundation Grand Challenges Grant (Phase II - OPP1119049), an NIH Director's New Innovators

Award (1DP2OD001315-01), the Center for Quantitative Biology (P50 GM071508), and startup

funding from the Pennsylvania State University. J.M.M., M.W.M. and A.B.V. were supported by

NIH grant R01 AI028398. M.K.R. is a recipient of a VA Research Career Scientist Award (14S-

RCS001) and his group receives financial support from the VA's Merit Review Program (i01

BX003312). Research reported in this publication was also supported by the U.S. National

Institutes of Health under award number AI100569 (M.K.R., A.B.V., J.M.M., and M.W.M.) and

by the U.S. Department of Defense Peer Reviewed Medical Research Program (Log # PR130649;

Contract # W81XWH-14-1-0447) (M.K.R.). The team also acknowledges valuable support

provided by the Medicines for Malaria Venture. This work was also supported by the Intramural

Research Program of the Center for Biologics Evaluation and Research, U.S. Food and Drug

Administration funding awarded to H.J.P.

**List of Supplemental Information:**

Supplemental Methods

Table S1: Drug susceptibility assays of various parasite lines

Fig S1: Effects of Inhibitors on pyrimidine synthesis precursors in both sensitive and resistant *P.*

*falciparum.*

Table S2: SNP analysis of the *P.f.* SB1-A6 genome

<u>Table S3:</u> Copy number analysis of the *P.f.* SB1-A6 genome

<u>Fig S2:</u> IFA of DHODH localization in D6 and SB1-A6

<u>Fig S3:</u> Sequence trace of C276F introduction or F276C repair

**<u>References:</u>**

1.    White J, Dhingra SK, Deng X, El Mazouni F, Lee MCS, Afanador GA, Lawong A, Tomchick DR, Ng CL, Bath J, Rathod PK, Fidock DA, Phillips MA. 2019. Identification and mechanistic understanding of dihydroorotate dehydrogenase point mutations in *Plasmodium falciparum* that confer in vitro resistance to the clinical candidate DSM265. ACS Infect Dis 5:90-101.

2.    Guler JL, White J, Phillips MA, Rathod PK. 2015. Atovaquone tolerance in *Plasmodium falciparum* parasites selected for high-level resistance to a dihydroorotate dehydrogenase inhibitor. Antimicrob Agents and Chemother 59:686-689.

3.    Guler JL, Freeman DL, Ahyong V, Patrapuvich R, White J, Gujjar R, Phillips MA, DeRisi J, Rathod PK. 2013. Asexual populations of the human malaria parasite, *Plasmodium falciparum,* use a two-step genomic strategy to acquire accurate, beneficial DNA amplifications. PLOS Pathogens 9:e1003375.

4.    Miller LH, Ackerman HC, Su XZ, Wellems TE. 2013. Malaria biology and disease pathogenesis: Insights for new treatments. Nat Med 19:156-67.

5.    RTS SCTP. 2015. Efficacy and safety of RTS,S/AS01 malaria vaccine with or without a booster dose in infants and children in Africa: final results of a phase 3, individually randomised, controlled trial. Lancet 386:31-45.

6.    Cui L, Lindner S, Miao J. 2015. Translational regulation during stage transitions in malaria parasites. Ann N Y Acad Sci 1342:1-9.

7.    Cowell AN, Istvan ES, Lukens AK, Gomez-Lorenzo MG, Vanaerschot M, Sakata-Kato T, Flannery EL, Magistrado P, Owen E, Abraham M, LaMonte G, Painter HJ, Williams RM, Franco V, Linares M, Arriaga I, Bopp S, Corey VC, Gnadig NF, Coburn-Flynn O, Reimer C, Gupta P, Murithi JM, Moura PA, Fuchs O, Sasaki E, Kim SW, Teng CH, Wang LT, Akidil A, Adjalley S, Willis PA, Siegel D, Tanaseichuk O, Zhong Y, Zhou Y, Llinas M, Ottilie S, Gamo FJ, Lee MCS, Goldberg DE, Fidock DA, Wirth DF, Winzeler EA. 2018. Mapping the malaria parasite druggable genome by using in vitro evolution and chemogenomics. Science 359:191-199.

8.    Allman EL, Painter HJ, Samra J, Carrasquilla M, Llinas M. 2016. Metabolomic profiling
of the malaria box reveals antimalarial target pathways. Antimicrob Agents and Chemother
doi:10.1128/AAC.01224-16.
9.    Cobbold SA, Chua HH, Nijagal B, Creek DJ, Ralph SA, McConville MJ. 2016. Metabolic
dysregulation induced in *Plasmodium falciparum* by dihydroartemisinin and other front-
line antimalarial drugs. J Infect Dis 213:276-86.
10.   Painter HJ, Morrisey JM, Mather MW, Vaidya AB. 2007. Specific role of mitochondrial
electron transport in blood-stage *Plasmodium falciparum*. Nature 446:88-91.
11.   Painter HJ, Morrisey JM, Vaidya AB. 2010. Mitochondrial electron transport inhibition
and viability of intraerythrocytic *Plasmodium falciparum*. Antimicrob Agents and
Chemother 54:5281-7.
12.   Vaidya AB, Painter HJ, Morrisey JM, Mather MW. 2008. The validity of mitochondrial
dehydrogenases as antimalarial drug targets. Trends in Parasitol 24:8-9.
13.   MacRae JI, Dixon MW, Dearnley MK, Chua HH, Chambers JM, Kenny S, Bottova I,
Tilley L, McConville MJ. 2013. Mitochondrial metabolism of sexual and asexual blood
stages of the malaria parasite *Plasmodium falciparum*. BMC Biology 11:67.
14.   Cobbold SA, Vaughan AM, Lewis IA, Painter HJ, Camargo N, Perlman DH, Fishbaugher
M, Healer J, Cowman AF, Kappe SH, Llinas M. 2013. Kinetic flux profiling elucidates
two independent acetyl-CoA biosynthetic pathways in *Plasmodium falciparum*. J Biol
Chem 288:36338-50.
15.   Ke H, Lewis IA, Morrisey JM, McLean KJ, Ganesan SM, Painter HJ, Mather MW, Jacobs-
Lorena M, Llinas M, Vaidya AB. 2015. Genetic investigation of tricarboxylic acid
metabolism during the *Plasmodium falciparum* life cycle. Cell Reports 11:164-74.
16.   Phillips MA, Lotharius J, Marsh K, White J, Dayan A, White KL, Njoroge JW, El Mazouni
F, Lao Y, Kokkonda S, Tomchick DR, Deng X, Laird T, Bhatia SN, March S, Ng CL,
Fidock DA, Wittlin S, Lafuente-Monasterio M, Benito FJ, Alonso LM, Martinez MS,
Jimenez-Diaz MB, Bazaga SF, Angulo-Barturen I, Haselden JN, Louttit J, Cui Y, Sridhar
A, Zeeman AM, Kocken C, Sauerwein R, Dechering K, Avery VM, Duffy S, Delves M,
Sinden R, Ruecker A, Wickham KS, Rochford R, Gahagen J, Iyer L, Riccio E, Mirsalis J,
Bathhurst I, Rueckle T, Ding X, Campo B, Leroy D, Rogers MJ, et al. 2015. A long-
duration dihydroorotate dehydrogenase inhibitor (DSM265) for prevention and treatment
of malaria. Science Trans Med 7:296ra111.
17.   Phillips MA, Gujjar R, Malmquist NA, White J, El Mazouni F, Baldwin J, Rathod PK.
2008. Triazolopyrimidine-based dihydroorotate dehydrogenase inhibitors with potent and
selective activity against the malaria parasite *Plasmodium falciparum*. J Med Chem.
18.   Baldwin J, Michnoff CH, Malmquist NA, White J, Roth MG, Rathod PK, Phillips MA.
2005. High-throughput screening for potent and selective inhibitors of *Plasmodium*
*falciparum* dihydroorotate dehydrogenase. J Biol Chem 280:21847-21853.
19.   Nilsen A, LaCrue AN, White KL, Forquer IP, Cross RM, Marfurt J, Mather MW, Delves
MJ, Shackleford DM, Saenz FE, Morrisey JM, Steuten J, Mutka T, Li Y, Wirjanata G,

Ryan E, Duffy S, Kelly JX, Sebayang BF, Zeeman AM, Noviyanti R, Sinden RE, Kocken CH, Price RN, Avery VM, Angulo-Barturen I, Jimenez-Diaz MB, Ferrer S, Herreros E, Sanz LM, Gamo FJ, Bathurst I, Burrows JN, Siegl P, Guy RK, Winter RW, Vaidya AB, Charman SA, Kyle DE, Manetsch R, Riscoe MK. 2013. Quinolone-3-diarylethers: a new class of antimalarial drug. Sci Transl Med 5:177ra37.

20. Winter RW, Kelly JX, Smilkstein MJ, Dodean R, Bagby GC, Rathbun RK, Levin JI, Hinrichs D, Riscoe MK. 2006. Evaluation and lead optimization of anti-malarial acridones. Exp Parasitol 114:47-56.

21. Smilkstein MJ, Forquer I, Kanazawa A, Kelly JX, Winter RW, Hinrichs DJ, Kramer DM, Riscoe MK. 2008. A drug-selected *Plasmodium falciparum* lacking the need for conventional electron transport. Molec Biochem Parasitol 159:64-68.

22. Stickles AM, De Almeida MJ, Morrisey JM, Sheridan KA, Forquer IP, Nilsen A, Winter RW, Burrows JN, Fidock DA, Vaidya AB, Riscoe MK. 2015. Subtle changes in endochin-like quinolone structure alter the site of inhibition within the cytochrome *bc 1* complex of *Plasmodium falciparum*. Antimicrob Agents Chemother 59:1977-1982.

23. Mandt REK, Lafuente-Monasterio MJ, Sakata-Kato T, Luth MR, Segura D, Pablos-Tanarro A, Viera S, Magan N, Ottilie S, Winzeler EA, Lukens AK, Gamo FJ, Wirth DF. 2019. In vitro selection predicts malaria parasite resistance to dihydroorotate dehydrogenase inhibitors in a mouse infection model. Sci Transl Med 11.

24. Ke H, Morrisey JM, Ganesan SM, Painter HJ, Mather MW, Vaidya AB. 2011. Variation among *Plasmodium falciparum* strains in their reliance on mitochondrial electron transport chain function. Eukaryotic Cell 10:1053-61.

25. Ganesan SM, Morrisey JM, Ke H, Painter HJ, Laroiya K, Phillips MA, Rathod PK, Mather MW, Vaidya AB. 2011. Yeast dihydroorotate dehydrogenase as a new selectable marker for *Plasmodium falciparum* transfection. Molec Biochem Parasitol 177:29-34.

26. Vincent IM, Barrett MP. 2015. Metabolomic-based strategies for anti-parasite drug discovery. J Biomol Screen 20:44-55.

27. Creek DJ, Barrett MP. 2014. Determination of antiprotozoal drug mechanisms by metabolomics approaches. Parasitol 141:83-92.

28. Ross LS, Gamo FJ, Lafuente-Monasterio MJ, Singh OM, Rowland P, Wiegand RC, Wirth DF. 2014. In vitro resistance selections for *Plasmodium falciparum* dihydroorotate dehydrogenase inhibitors give mutants with multiple point mutations in the drug-binding site and altered growth. J Biol Chem 289:17980-95.

29. Srivastava IK, Morrisey JM, Darrouzet E, Daldal F, Vaidya AB. 1999. Resistance mutations reveal the atovaquone-binding domain of cytochrome *b* in malaria parasites. Mol Microbiol 33:704-11.

30. Bedingfield PTP, Cowen D, Acklam P, Cunningham F, Parsons MR, McConkey GA, Fishwick CWG, Johnson AP. 2012. Factors Influencing the specificity of inhibitor binding to the human and malaria parasite dihydroorotate dehydrogenases. J Med Chem 55:5841-5850.

31.	Munier-Lehmann H, Vidalain PO, Tangy F, Janin YL. 2013. On dihydroorotate dehydrogenases and their inhibitors and uses. J Med Chem 56:3148-67.

32.	Patel V, Booker M, Kramer M, Ross L, Celatka CA, Kennedy LM, Dvorin JD, Duraisingh MT, Sliz P, Wirth DF, Clardy J. 2008. Identification and characterization of small molecule inhibitors of *Plasmodium falciparum* dihydroorotate dehydrogenase. J Biol Chem 283:35078-35085.

33.	Wells TNC, Van Huijsduijnen RH, Van Voorhis WC. 2015. Malaria medicines: a glass half full? Nat Rev Drug Discov 14:424-442.

34.	Ghorbal M, Gorman M, Macpherson CR, Martins RM, Scherf A, Lopez-Rubio J-J. 2014. Genome editing in the human malaria parasite *Plasmodium falciparum* using the CRISPR-Cas9 system. Nat Biotech 32:819-821.

35.	Goodman CD, Siregar JE, Mollard V, Vega-Rodriguez J, Syafruddin D, Matsuoka H, Matsuzaki M, Toyama T, Sturm A, Cozijnsen A, Jacobs-Lorena M, Kita K, Marzuki S, McFadden GI. 2016. Parasites resistant to the antimalarial atovaquone fail to transmit by mosquitoes. Science 352:349-53.

36.	Bajzikova M, Kovarova J, Coelho AR, Boukalova S, Oh S, Rohlenova K, Svec D, Hubackova S, Endaya B, Judasova K, Bezawork-Geleta A, Kluckova K, Chatre L, Zobalova R, Novakova A, Vanova K, Ezrova Z, Maghzal GJ, Magalhaes Novais S, Olsinova M, Krobova L, An YJ, Davidova E, Nahacka Z, Sobol M, Cunha-Oliveira T, Sandoval-Acuña C, Strnad H, Zhang T, Huynh T, Serafim TL, Hozak P, Sardao VA, Koopman WJH, Ricchetti M, Oliveira PJ, Kolar F, Kubista M, Truksa J, Dvorakova-Hortova K, Pacak K, Gurlich R, Stocker R, Zhou Y, Berridge MV, Park S, Dong L, Rohlena J, Neuzil J. 2019. Reactivation of dihydroorotate dehydrogenase-driven pyrimidine biosynthesis restores tumor growth of respiration-deficient cancer cells. Cell Metabol 29:399-416.e10.

37.	Trager W, Jensen JB. 1976. Human malaria parasites in continuous culture. Science 193:673-5.

38.	Lambros C, Vanderberg JP. 1979. Synchronization of *Plasmodium falciparum* erythrocytic stages in culture. J Parasitol 65:418-20.

39.	Desjardins RE, Canfield CJ, Haynes JD, Chulay JD. 1979. Quantitative assessment of antimalarial activity in vitro by a semiautomated microdilution technique. Antimicrob Agents Chemother 16:710-718.

40.	Smilkstein M, Sriwilaijaroen N, Kelly JX, Wilairat P, Riscoe M. 2004. Simple and inexpensive fluorescence-based technique for high-throughput antimalarial drug screening. Antimicrob Agents Chemother 48:1803-6.

41.	Haynes JD, Moch JK. 2002. Automated synchronization of *Plasmodium falciparum* parasites by culture in a temperature-cycling incubator. Methods Molec Med 72:489-497.

42.	Mather MW, Morrisey JM, Vaidya AB. 2010. Hemozoin-free *Plasmodium falciparum* mitochondria for physiological and drug susceptibility studies. Mol Biochem Parasitol 174:150-3.

43. Olszewski KL, Llinas M. 2013. Extraction of hydrophilic metabolites from *Plasmodium falciparum*-infected erythrocytes for metabolomic analysis. Methods Molec Biol 923:259-66.

44. Cobbold SA, Chua HH, Nijagal B, Creek DJ, Ralph SA, McConville MJ. 2015. Metabolic dysregulation induced in *Plasmodium falciparum* by dihydroartemisinin and other front line antimalarial drugs. J Infect Dis doi:10.1093/infdis/jiv372.

45. Lu W, Clasquin MF, Melamud E, Amador-Noguez D, Caudy AA, Rabinowitz JD. 2010. Metabolomic analysis via reversed-phase ion-pairing liquid chromatography coupled to a stand alone orbitrap mass spectrometer. Anal Chem 82:3212-21.

46. Pulcini S, Staines HM, Lee AH, Shafik SH, Bouyer G, Moore CM, Daley DA, Hoke MJ, Altenhofen LM, Painter HJ, Mu J, Ferguson DJ, Llinas M, Martin RE, Fidock DA, Cooper RA, Krishna S. 2015. Mutations in the *Plasmodium falciparum* chloroquine resistance transporter, PfCRT, enlarge the parasite's food vacuole and alter drug sensitivities. Sci Rep 5:14552.

47. Goecks J, Nekrutenko A, Taylor J, Galaxy T. 2010. Galaxy: a comprehensive approach for supporting accessible, reproducible, and transparent computational research in the life sciences. Genome Biol 11:R86.

48. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25:1754-60.

49. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, del Angel G, Rivas MA, Hanna M, McKenna A, Fennell TJ, Kernytsky AM, Sivachenko AY, Cibulskis K, Gabriel SB, Altshuler D, Daly MJ. 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. Nat Gen 43:491-8.

50. Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing. arXiv:12073907 [q-bioGN].

51. Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. Fly (Austin) 6:80-92.

52. Abyzov A, Urban AE, Snyder M, Gerstein M. 2011. CNVnator: an approach to discover, genotype, and characterize typical and atypical CNVs from family and population genome sequencing. Genome Res 21:974-84.

53. Nkrumah LJ, Muhle RA, Moura PA, Ghosh P, Hatfull GF, Jacobs WR, Jr., Fidock DA. 2006. Efficient site-specific integration in *Plasmodium falciparum* chromosomes mediated by mycobacteriophage Bxb1 integrase. Nat Methods 3:615-21.

54. Fidock DA, Wellems TE. 1997. Transformation with human dihydrofolate reductase renders malaria parasites insensitive to WR99210 but does not affect the intrinsic activity of proguanil. PNAS 94:10931-10936.

55.    van den Hoff MJ MA, and Lamers WH. 1992. Electroporation in 'intracellular' buffer
increases cell survival. Nucleic Acid Res 20:2902.

**Figure Legends:**

**Figure 1: Drug-resistant phenotyping of *P.f.* strain SB1-A6.** A) Early ring-stage *P. falciparum*

strain D6 (blue) and SB1-A6 (red) were exposed to titrating concentrations of atovaquone or DSM-

1. Their susceptibility was directly compared to a phenocopy strain, D10:yDHODH (grey, red

outline) and its parent, D10 (grey, blue outline). Parasite survival was determined using a

traditional 48-h SYBR-green growth inhibition assay and plotted as an average of three technical

replicates (±SD). B) Reversal of the SB1-A6 phenotype is demonstrated by exposure of ring-stage

parasites to titrating concentrations of atovaquone with (red circles, dashed line) or without the

addition of 1.0 μM proguanil (red circles, solid line). Parasite survival was determined as in (A).

C) Whole-cell metabolomic analysis was performed on trophozoite stages of both *P.f.* strain D6

and SB1-A6 after two hours of 10x $1C_{50}$ exposure to either atovaquone or DSM-1. Metabolic

changes are displayed as a MetaPrint profile (8) and highlighted are the increasing (blue outline)

metabolites in D6 compared to no change SB1-A6 (red outline). D) The increased metabolites

highlighted (C) in *P.f.* strain D6 (blue) are plotted as absolute signal determined by the average

total ion counts (biological duplicates in technical triplicate) for *N*-carbamoyl-L-aspartate and

dihydroorotate in the presence of atovaquone (10nM), DSM-1 (1.5 μM) or absence of drug relative

to *P.f.* strain SB1-A6 (red).

**Figure 2: Genetic and molecular differences in *P.f.* strain SB1-A6.** A) Whole-genome

sequencing revealed a point mutation in PF3D7_0603300 (*pfdhodh*) and this mutation was verified

by Sanger sequencing. A trace of the PfDHODH coding sequence is shown from nucleotide 818-

836 and the single nucleotide polymorphism change from the parental D6 to SB1-A6 at nucleotide

G827T (yellow highlight) which results in the amino acid change C276F. Additionally, WGS

revealed B) a 1.9x copy number increase of a ~20kb region (red highlight) on chromosome 6 which

includes PF3D7_0603300 (*pfdhodh*). Copy number increase was calculated and graphically

represented using Intansiv (v. 0.99.3).  C) Structural comparisons of wild-type (PDB 4RX0, light

grey, C276 highlighted in blue) and mutant (PDB 6E0B, dark grey, C276F highlighted in red)

PfDHODH bound with FMN, as well as the merged structure surrounding amino acid 276 (inset;

cysteine in blue and phenylalanine in red, represent the isosurface of each sidechain) were made

using PyMOL 2.3.4. D) Western blot analysis of the PfDHODH expression in both D6 and SB1-

A6 probed with anti-mouse PfDHODH antibody (1:300) compared to *Pf*Aldolase (anti-mouse

*Pf*Aldolase-HRP, 1:5000) as a loading control.  E) Percent inhibition of PfDHODH activity and

F) capture of total PfDHODH activity in isolated mitochondria from SB1-A6 (red) and D6 (blue)

in the presence of titrating concentrations of DSM-1.

**Figure 3: Phenotyping Individual Genetic Contributions to *P.f.* SB1-A6 Resistance.** A)

Strategy for phenotyping a mimic of the PfDHODH genotype in the wild-type *P.f.* D6 strain.

Episomal expression of an additional copy of wild-type PfDHODH (blue) aims to genocopy a

CNV and episomal expression of the mutant PfDHODH (red) genocopies a CNV of the mutation

to capture the contribution of these genotypes to the resistance phenotype. B) These transgenic

lines, D6::wtDHODH (blue triangle, dashed) and D6::mutDHODH (red open circle, dashed), were

assessed for their susceptibility to titrating concentrations of both atovaquone and DSM-1 in

comparison to the parental strains, A6 (red solid triangle) and D6 (blue solid square). Parasite

survival was determined using a traditional 48-h SYBR-green growth inhibition assay and plotted

as an average of three technical replicates (±SD). C) Graphic representation of the CRISPR/Cas9-

mediated mutational strategy for modification of PfDHODH in both the wild-type D6 and mutant

strain SB1-A6. A plasmid expressing Cas9 was cotransfected with a plasmid that contains the

DHODH repair template that encodes for C276 (wild-type) or F276 (mutant) that are provided to

introduce a single point mutation converting strain D6 to C276F and SB1-A6 to F276C,

respectively. The CRISPR/Cas9 mutated strains D6$^{C276F}$ (blue solid diamond, dashed) and SB1-

A6$^{F276C}$ (red solid triangle, dashed) were then assayed for their susceptibility to titrating

concentrations of both atovaquone and DSM-1 in comparison to the parental strains, A6 (red solid

triangle) and D6 (blue solid square). Parasite survival was determined using a traditional 48-h

SYBR-green growth inhibition assay and plotted as an average of three technical replicates (±SD).

**Figure 1**

## Figure 2

**Figure 3**