Diets affect the age-dependent decline of associative learning in *Caenorhabditis elegans*

Authors

Satoshi Higurashi[1, 2, *], Sachio Tsukada[1,2, *], Binta Maria Aleogho[2], Joo Hyun Park[2], Masaru

Tanaka[1,2], Shunji Nakano[3], Ikue Mori[3], and Kentaro Noma[2, #]

Affiliations

1. Milk Science Research Institute, Megmilk Snow Brand Co. Ltd., 1-1-2 Minamidai, Kawagoe, Saitama,

350-1165, Japan

2. Group of Nutritional Neuroscience, Neuroscience Institute, Graduate School of Science, Nagoya

University, Nagoya, 464-8602, Japan

3. Group of Molecular Neurobiology, Neuroscience Institute, Graduate School of Science, Nagoya

University, Nagoya, 464-8602, Japan

Current address: Group of Microbial Motility, Department of Biological Science, Division of Natural

Science, Graduate school of Science, Nagoya University, Nagoya, 464-8602, Japan

*These authors equally contributed to this work.

Correspondence to

Kentaro Noma

noma.kentaro.f1@f.mail.nagoya-u.ac.jp

Abstract

The causality and mechanism of dietary effects on brain aging are still unclear due to the long time scales

of aging. The nematode *Caenorhabditis elegans* has contributed to aging research because of its short

lifespan and easy genetic manipulation. When fed the standard laboratory diet, *Escherichia coli*, *C.*

elegans experiences an age-dependent decline in temperature-food associative learning, called

thermotaxis. To address if diet affects this decline, we screened 35 lactic acid bacteria as alternative diets

and found that animals maintained high thermotaxis ability when fed a clade of *Lactobacilli* enriched with

heterofermentative bacteria. Among them, *Lactobacillus reuteri* maintained the thermotaxis of aged

animals without affecting their lifespan and motility. The effect of *Lb. reuteri* depends on the DAF-16

transcription factor functioning in neurons. Furthermore, RNA sequencing analysis revealed that

differentially expressed genes between aged animals fed different bacteria were enriched with DAF-16

targets. Our results demonstrate that diet can impact brain aging in a *daf-16* dependent manner without

changing the lifespan.

Introduction

Human life expectancy has increased since the nineteenth century (Dong, Milholland, & Vijg, 2016), which has led to the social problem related to age-dependent cognitive dysfunction. Although human studies suggest that genetic background, diet, and lifestyle might affect brain aging, the causality and mechanism of how they affect brain aging remain unclear (Deary et al., 2009).

The nematode *Caenorhabditis elegans* (*C. elegans*) is ideal for addressing the mechanism of age-related phenotypes because of the two to three-week lifespan and the variety of available genetic tools. In *C. elegans*, the age-related phenotypes can be readily separable from the organismal lifespan by directly measuring the lifespan. In the past decades, studies using *C. elegans* have contributed to aging research by revealing the molecular mechanism of how dietary restriction, insulin-like signaling, and germline stem cells affect organismal lifespan (Mack, Heimbucher, & Murphy, 2018; Wolff & Dillin, 2006). Like mammals, *C. elegans* experiences age-dependent functional changes in the nervous system (Stein & Murphy, 2012). Aged animals are defective in locomotion (Hahm et al., 2015; Mulcahy, Holden-Dye, & O'Connor, 2013), mechanosensory response (Beck & Rankin, 1993), chemotaxis (Leinwand et al., 2015), thermotaxis (Huang et al., 2020; H. Murakami, Bessinger, Hellmann, & Murakami, 2005; S. Murakami & Murakami, 2005), and food-butanone associative learning (Kauffman, Ashraf, Corces-Zimmerman, Landis, & Murphy, 2010). Age-dependent memory decline in the food-butanone association is ameliorated in the mutant of *nkat-1* encoding kynurenic acid-synthesizing enzyme prevents (Vohra, Lemieux, Lin, & Ashrafi, 2018). Overactivation of Gα signaling in AWC sensory neurons also maintains the ability to form memory in aged animals in the food-butanone association (Arey, Stein, Kaletsky, Kauffman, & Murphy, 2018). These emerging evidence suggests that genetic manipulations can prevent age-dependent functional decline in the nervous system.

Compared to genetic manipulations, the modification of diet can be easily applicable to our daily lives. Studies in humans and mice imply that diets affect the cognitive decline in aged animals (Joseph, Cole, Head, & Ingram, 2009; Vauzour et al., 2017). Here, we use *C. elegans* to address the dietary effect

on the age-dependent behavioral decline and its underlying mechanism. In laboratories, *C. elegans* is

maintained monoxenically with a uracil auxotroph *Escherichia coli* (*E. coli*) strain, OP50, as the standard

diet (Brenner, 1974). On the other hand, *C. elegans* in natural habitat eats a wide variety of bacteria (Berg

et al., 2016; Dirksen et al., 2016; Johnke, Dirksen, & Schulenburg, 2020; Samuel, Rowedder, Braendle,

Felix, & Ruvkun, 2016; Zhang et al., 2017). These bacteria affect the physiology of *C. elegans*, such as

growth rate, reproduction, and sensory behavior (Dirksen et al., 2016; O'Donnell, Fox, Chao, Schroeder,

& Sengupta, 2020; Samuel et al., 2016). However, the effect of different bacteria on the behavioral

decline during aging is unexplored. Among the potential bacterial diet for *C. elegans* in natural habitat

(Berg et al., 2016; Dirksen et al., 2016; Samuel et al., 2016), we focused on Lactic Acid Bacteria (LAB),

which are the most common probiotics for humans (Hill et al., 2014). LAB, such as *Lactobacilli* (*Lb.*) and

Bifidobacteria (*B.*), are Gram-positive, non-spore-forming bacteria that produce lactic acid from

carbohydrates as the primary metabolic product. Depending on the species, LAB have various effects on

C. elegans physiology. *Lb. gasseri*, *B. longum*, and *B. infantis* extend lifespan in *C. elegans* (Komura,

Ikeda, Yasui, Saeki, & Nishikawa, 2013; Nakagawa et al., 2016; L. Zhao et al., 2017). On the other hand,

Lb. helveticus does not increase the lifespan (Nakagawa et al., 2016). Even in the same species, different

strains have different effects on lifespan, body size, and locomotion (Wang et al., 2020). In *C. elegans*,

LAB modulate evolutionarily conserved genetic pathways such as the insulin/insulin-like growth factor-1

(IGF-1) signaling (IIS) pathway (Grompone et al., 2012; Sugawara & Sakamoto, 2018), which consists of

the insulin receptor DAF-2, phosphoinositide 3 (PI3) kinase cascade, and the downstream transcription

factor DAF-16 (Kenyon, Chang, Gensch, Rudner, & Tabtiang, 1993; Lin, Dorman, Rodan, & Kenyon,

1997). DAF-16 is a sole *C. elegans* ortholog of mammalian FOXO transcription factor and is involved in

various biological processes (Stein & Murphy, 2012; Tissenbaum, 2018).

To comprehensively understand the effect of LAB, we screened 35 different LAB species,

including some subspecies. We examined the age-dependent functional decline of thermotaxis behavior,

which reflects associative learning between temperature and food (Hedgecock & Russell, 1975; Mori &

Ohshima, 1995). We demonstrate that *C. elegans* fed *Lactobacilli* in a clade maintained the thermotaxis

behavior when aged, while *E. coli*-fed animals did not. Among those *Lactobacilli*, *Lb. reuteri* maintained

the thermotaxis ability of aged animals without affecting the organismal lifespan or locomotion. The

effect of *Lb. reuteri* on the thermotaxis of aged animals depends on the DAF-16 transcription factor

functioning in neurons.

Results

C. elegans thermotaxis behavior declines with age

After being cultivated with food at a temperature within the physiological range (15~25 °C), *C. elegans*

migrates toward and stays at the past cultivation temperature (T_{cult}) on a linear thermal gradient without

food (Fig. 1A). This behavior is called thermotaxis (Hedgecock & Russell, 1975; Mori & Ohshima,

1995). To see the effect of aging on thermotaxis, we cultivated animals at 20 °C with an *E. coli* strain,

OP50 (hereafter, *E. coli*, unless otherwise noted), commonly used in laboratory conditions (Brenner,

1974). When the animals were placed at 17 °C on a temperature gradient without food, young adults (day

1 of adulthood, D1) migrated up the temperature gradient toward 20 °C (Fig. 1B). On the other hand,

aged animals (day 5 of adulthood, D5) remained around the spotted area and did not reach the area near

T_{cult} (Fig. 1B), as previously reported (Huang et al., 2020). The ability to perform the thermotaxis

behavior was quantified using the performance index, indicating the fraction of animals around T_{cult} (Fig.

1A). The performance index declined from D1 to D5 (Fig. 1C). To further accelerate aging (Fig. S1)

(Klass, 1977), we cultivated animals at 23 °C and placed them on a temperature gradient centered at

20 °C. In this condition, animals gradually lost the ability to move toward T_{cult} during aging (Fig. S2), and

the performance index declined from ~0.75 at D1 to ~0.25 at D5 (Fig. 1D). This age-dependent

thermotaxis decline was not specific to OP50-fed animals because animals fed another *E. coli* strain,

HT115, also showed a similar decline (Fig. S3).

　　　　The low thermotaxis performance of *E. coli*-fed aged animals appeared to be independent of

defects in motility or temperature sensation because D5 animals cultivated at 20 °C could migrate down

the thermal gradient relatively normally when the origin was at 23 °C (Figs. 1B and 1C). Consistent with

this notion, we did not observe the loss of AFD or AIY neurons at D5 (Fig. S4). Moreover, aged animals

could sense food normally based on the basal slowing response in the presence of food (Figs. S5A and

S5B) (Sawin, Ranganathan, & Horvitz, 2000). The food sensation of aged animals is also reported

to be normal, based on the attraction to *E. coli* (Cornils et al., 2016). In contrast to the basal

slowing response, we note that aged animals did not show a significantly enhanced slowing

response in the starved condition (Fig. S5B), implying that aged animals might not sense starvation

normally.

To address if the defects can be observed in another associative learning behavior, we tested the

salt avoidance behavior using two settings of assays (Saeki, Yamamoto, & Iino, 2001; Wicks, de Vries,

van Luenen, & Plasterk, 2000) (Figs. S6A and S6B). As previously reported, naïve D1 animals were

attracted by NaCl, while they avoided NaCl when cultivated with NaCl in the absence of food in both

settings (Saeki et al., 2001; Wicks et al., 2000) (Figs. S6C and S6D). In contrast to the thermotaxis

behavior, D5 animals showed normal salt-avoidance behaviors (Figs. S6C and S6D).

We thus concluded that age-dependent thermotaxis changes reflected the decline of specific

associative learning behavior and determined to use it to examine dietary effects.

Specific LAB prevent the age-dependent thermotaxis decline

To address if diets affect the age-dependent decline in thermotaxis, we fed animals with different lactic

acid bacteria (LAB) species instead of the regular *E. coli*. We selected 35 LAB, consisting of 17

Lactobacilli (Lb.), two *Pediococci (P.)*, two *Lactococci (Lc.)*, two *Streptococci (S.)*, five *Leuconostoc*

(Ls.), and seven *Bifidobacteria (B.)* (Table S1). To avoid developmental effects by feeding with LAB, we

cultivated animals with *E. coli* until D1 before switching to LAB (Fig. 2A). Animals were cultivated at

23 °C and spotted at the center of the 17-23 °C gradient (Fig. 2A). Five LAB did not support the survival

of animals during aging (Fig. 2B, NA). While eight LAB did not affect the thermotaxis performance

indices of the aged animals compared to *E. coli*, 22 LAB significantly increased them (Fig. 2B). Among

those, *P. pentosaceus*, *Lb. reuteri*, *Lb. rhamnosus*, and *Lb. plantarum* gave the highest performance

indices (Fig. 2B).

We first ruled out the possibility that aged animals fed the LAB were constitutively thermophilic,

irrespective of the T_{cult}. Thermophilicity is reported for mutants of genes such as *pkc-1/ttx-4* encoding

protein kinase C (Okochi, Kimura, Ohta, & Mori, 2005) and *tax-6* encoding calcineurin A subunit

(Kuhara, Inada, Katsura, & Mori, 2002). To distinguish between associative learning and thermophilicity,

we shifted the thermal gradient of assay plates from 17-23 °C to 20-26 °C for animals cultivated at 23 °C

(Fig. S7A). D5 *tax-6* mutants migrated toward a higher temperature than T_{cult} (Fig. S7B). On the other

hand, LAB-fed D5 animals stayed around the T_{cult} (Fig. S7B). We calculated the thermotaxis index,

instead of the performance index, to quantify animals' thermal preference (Ito, Inada, & Mori, 2006)

(Figs. S7A). Unlike thermophilic *tax-6* mutants, LAB-fed D5 animals showed thermotaxis indices

comparable to the D1 wild type (Fig. S7C), suggesting that LAB-fed D5 animals were not constitutively

thermophilic.

We next addressed if LAB-fed D5 animals can remember a new temperature by shifting the T_{cult}

from 23 °C to 17 °C one day before the thermotaxis assay. D1 animals could learn the new temperature

and shift their thermal preference from 23 °C (Fig. 3A) to the new T_{cult}, 17 °C (Fig. 3B). The thermotaxis

index was also shifted accordingly (Fig. 3C). LAB-fed D5 animals showed behavioral plasticity like D1

animals (Figs. 3B and 3C). This result suggests that LAB-fed aged animals retained an ability to learn a

new T_{cult}.

Different LAB show various effects on lifespan and locomotion

Some LAB extend the lifespan of *C. elegans* (Komura et al., 2013; Nakagawa et al., 2016; Wang et al.,

2020; Y. Zhao et al., 2013). Therefore, better thermotaxis performance of LAB-fed aged animals might

result from a systemic effect of prolonged organismal lifespan. To address this possibility, we measured

the lifespan of animals fed the four LAB: *P. pentosaceus*, *Lb. reuteri*, *Lb. rhamnosus*, and *Lb. plantarum.*

To avoid the growth of *E. coli* on LAB plates after transferring animals, we used peptone-free NGM

plates (T. Ikeda, Yasui, Hoshino, Arikawa, & Nishikawa, 2007; Lee, Kwon, & Lim, 2015). The lack of

peptone in the culture plates did not affect the dietary effects on the thermotaxis of aged animals (Fig.

S8). LAB had various effects on the lifespan of animals: *P. pentosaceus* prolonged the lifespan; *Lb.*

reuteri did not affect the lifespan; *Lb. rhamnosus* and *Lb. plantarum* shortened the lifespan (Fig. 4A).

We next examined the effect of LAB on the locomotion of aged animals using two assays:

thrashing assay (Miller et al., 1996) and motility assay. As previously reported (Glenn et al., 2004; Hahm

et al., 2015; Mulcahy et al., 2013), aged animals fed *E. coli* showed slight locomotion defects in both

assays (Figs. 4B and 4C). In the thrashing assay, *Lb. reuteri*- and *Lb. rhamnosus*-fed aged animals

showed better locomotion than *E. coli*-fed aged animals, while *P. pentosaseus* and *Lb. plantarum* did not

have effects (Fig. 4B). In the motility assay, we measured the distance animals migrate on a plate. *Lb.*

plantarum- and *Lb. rhamnosus*-fed aged animals showed reduced locomotion than *E. coli*-fed aged

animals, while *P. pentosaseus* and *Lb. reuteri* did not have effects (Fig. 4C). Thus, the four LAB selected

based on thermotaxis had different effects on the lifespan and locomotion, implying that the dietary effect

on thermotaxis is independent of lifespan and motility.

Bacteria affect the age-dependent thermotaxis decline as nutrition

How do different bacteria affect the thermotaxis of aged animals? *C. elegans* shows different preferences

in the bacterial diet (Shtonda & Avery, 2006). Our LAB screen used different diets during the

temperature-food association before the thermotaxis assays (Fig. 2A). It raises the possibility that the

different strengths of the association during learning caused the difference in thermotaxis of aged animals.

To address this issue, we switched the foods one day before the thermotaxis assay (Fig. 5A). We used *Lb.*

reuteri because it did not affect the lifespan (Fig. 4A). Aged animals whose diet was switched from *Lb.*

reuteri to *E. coli* showed the high thermotaxis performance, while aged animals with the opposite

condition did not (Fig. 5A). This result suggests that the dietary effects of thermotaxis on aged animals do

not reflect the strength of association.

To examine if LAB affects animals as live bacteria or serves as nutrition, we examined the effect

of bacteria heat-killed at 65 °C for one hour or at 100 °C for ten minutes (see Materials and Methods).

Like aged animals fed live bacteria, ones fed 65 °C-treated *E. coli* and *Lb. reuteri* showed low and high

performance indices in thermotaxis, respectively (Fig. 5B). Thus, the effect of both bacteria on

thermotaxis is independent of the condition of the bacteria being alive. On the other hand, animals fed

100 °C-treated *E. coli* showed higher performance than those fed live *E. coli* (Fig. 5B). This result

suggests that a component in *E. coli* resistant to 65 °C but sensitive to 100 °C facilitates thermotaxis

decline during aging. Moreover, animals fed crushed *E. coli* also showed higher thermotaxis performance,

implying that the said component might be diffused after crushing (Fig. 5B).

Next, we examined which bacteria, *E. coli* or *Lb. reuteri* has dominant effects on the thermotaxis

of aged animals. We mixed *E. coli* and *Lb. reuteri* and fed animals from D1. Both *E. coli* and *Lb. reuteri*

were ingested by *C. elegans* even when mixed, based on FITC labeling of bacteria (Fig. S9). In this

condition, *E. coli* had a dominant effect even when *Lb. reuteri* was mixed with twice as much as *E. coli*

(Fig. 5C). The dominant effect of *E. coli* seemed to require ingestion of bacteria because the exposure to

the *E. coli* odor did not affect the thermotaxis ability of *Lb. reuteri*-fed aged animals (Fig. 5D).

Collectively, we concluded that ingestion of bacterial nutrition during aging affects the

thermotaxis of aged animals.

The effects of LAB are associated with the phylogenetic tree

We explored the different features of bacteria which might affect *C. elegans* physiology. Gram-staining

showed that *E. coli* was Gram-negative, while LAB were Gram-positive as expected (Fig. S10). The

morphologies and sizes of the four select LAB were different (Fig. S10); these physical properties of

LAB may not explain the high performance indices of aged animals.

To examine the features of the genus of LAB, we analyzed a phylogenetic tree of 35 LAB strains

with the heatmap of the associated thermotaxis performance indices (Fig. 6). This analysis revealed that

LAB associated with high performance indices was significantly enriched in a specific clade, henceforth

referred to as Clade A (Fig. 6). Clade A, including *Lactobacilli* and *Pediococci,* was enriched in

obligatory and facultatively heterofermentative species except for *P. pentosaceus*, which was obligatory

homofermentative (Fig. 6). On the other hand, the *Lactobacilli*, associated with relatively low thermotaxis

indices (Clade B), are all homofermentative (Fig. 6). These results suggest that the nutritional feature of

LAB shared among the clades of *Lactobacilli* might affect *C. elegans*.

Neuronal *daf-16* is involved in the thermotaxis performance of *Lb. reuteri*-fed aged animals

Since the dietary effect on thermotaxis required more than one day to be manifested, bacterial diets seem

to alter the internal state of *C. elegans*. We addressed the molecular mechanism of how different diets

affect *C. elegans*. LAB can induce dietary restriction, which leads to a prolonged lifespan (Y. Zhao et al.,

2013). However, three out of four select LAB did not increase lifespan (Fig. 4A). *pha-4*, an ortholog of

the human FOXA2 transcription factor, is required for dietary restriction-induced longevity, and its

expression is increased by dietary restriction (Panowski, Wolff, Aguilaniu, Durieux, & Dillin, 2007). In

our condition, *pha-4* expression decreased in LAB-fed aged animals compared to *E. coli*-fed aged

animals, suggesting that LAB-fed animals were not under dietary restriction (Fig. S11A). To directly test

the effect of dietary restriction on the thermotaxis, we used *eat-2* mutants, which exhibit dietary

restriction by defective pharyngeal pumping (Lakowski & Hekimi, 1998). *eat-2* mutants did not increase

the performance index of *E. coli*-fed aged animals, although it may be due to the involvement of *eat-2* in

thermotaxis as shown in D1 animals (Fig. S9B). These results suggest that the high thermotaxis

performance of LAB-fed aged animals is not due to dietary restriction.

To find the gene involved in the dietary effect on aged animals, we tested mutants of three genes:

nkat-1 and *kmo-1* genes that encode enzymes in the kynurenic acid synthesizing pathway are known to be

involved in butanone-associated memory in aged animals (Vohra et al., 2018); *daf-16* that is an ortholog

of mammalian FOXO transcription factor involved in longevity (Kenyon et al., 1993) and LAB-

dependent lifespan extension (Grompone et al., 2012; Lee et al., 2015; Sugawara & Sakamoto, 2018).

Aged *nkat-1* and *kmo-1* mutants maintained thermotaxis ability like wild type when fed *Lb. reuteri*. On

the other hand, aged *daf-16* mutants showed significantly less ability to perform thermotaxis than their D1

counterpart (Fig. 7A); aged *daf-16* mutants fed *Lb. reuteri* distributed around a temperature slightly lower

than the T$_{cult}$ (Fig. 7B). This decreased thermotaxis ability in aged *daf-16* mutants fed *Lb. reuteri* was not

due to shortened lifespan because *daf-16* mutants had comparable lifespan to wild-type animals when fed

Lb. reuteri (Fig. 7C). DAF-16 is known to be activated in *daf-2* mutants. Therefore, we speculated that

daf-2 mutants might show high thermotaxis performance even in *E. coli*-fed aged animals. However, *daf-*

2 was required for nomal thermotaxis both young and aged animals (Fig. 7D).

daf-16 possesses several isoforms with different expression patterns and functions (Fig. 8A)

(Kwon, Narasimhan, Yen, & Tissenbaum, 2010). The b isoform is a neuronal isoform involved in AIY

development (Christensen, de la Torre-Ubieta, Bonni, & Colon-Ramos, 2011). The other isoforms are

more broadly expressed in most tissues, including neurons. We addressed which isoform is necessary for

the effect on thermotaxis of aged animals. *daf-16(mg54)* has a single nucleotide polymorphism (SNP)

which introduces an amber stop codon mutation and affects the exons of all *daf-16* isoforms except the b

isoform (Fig. 8A) (Ogg et al., 1997). To knock out only the b isoform, we generated *daf-16(knj36)* which

introduced 7-bp deletion in the first exon of the b isoform, located in the intron of the other isoforms (Fig.

8A). Both *daf-16(mg54)* and *daf-16(knj36)* did not affect the dietary effects on the thermotaxis of aged

animals (Fig. 8B). This result implies that the b isoform and other isoforms complement each other and

that having either one is sufficient to give the dietary effect. Consistently, the expression of *daf-16b* under

the control of its own promoter rescued the deletion mutants, *daf-16(mu86),* confirming the sufficiency of

the *daf-16 b* isoform.

We addressed in which tissue *daf-16* functions. Given that the expression of either b isoform or

non-b isoforms was sufficient to provide the dietary effects (Fig. 8B), *daf-16* might function in a tissue

where both b and non-b isoforms are expressed. We thus focused on pharyngeal muscles, body wall

muscles, and neurons (Nagashima, Iino, & Tomioka, 2019). Expression of *daf-16b* in pharyngeal muscles

or body wall muscles did not rescue *daf-16(mu86)* (Fig. 8C). On the other hand, the *daf-16 b* isoform

expressed in all neurons rescued the low performance of *Lb. reuteri*-fed aged *daf-16(mu86)* mutants (Fig.

8C). This result suggests that *daf-16* functioning in neurons is involved in the high thermotaxis

performance of *Lb. reuteri*-fed aged animals.

Diet and age affect DAF-16 target genes

To comprehensively understand the dietary effect on aging, we carried out RNA sequencing of eight

samples: D1, D5 fed *E. coli*, D5 fed homo-fermentative LAB, which gave low thermotaxis index (*Lb.*

gasseri and *Lb. delbrueckii*), and D5 fed heterofermentative LAB, which gave high thermotaxis index (*P.*

pentosaceus, *Lb. reuteri*, *Lb. rhamnosus*, and *Lb. plantarum*) (Fig. 9A). Principal component analysis of

transcriptome data revealed that the D5 fed LAB clustered together separately from D1 and D5 fed *E.*

coli, irrespective of the fermentation mode of LAB (Fig. 9B). The first principal component (PC1)

explained 76% of the entire variance and appeared to represent the difference in age irrespective of diets

(Fig. 9B, PC1). To characterize what genes contributed to PC1, we selected the top 5% genes positively

or negatively correlating to PC1 (666 genes, each, among 13331 genes in total, Fig. S12A) and performed

Gene Ontology (GO) analysis using Metascape (Y. Zhou et al., 2019). The genes correlating to PC1 were

enriched with the categories such as oogenesis (GO:0048477; $p=1.0 \times 10^{-10}$; Enrichment, 5.7) and

structural constituent of cuticle (GO:0042302; $p=1.0 \times 10^{-34}$; Enrichment, 8.3) (Fig. S12C). The second

principal component (PC2) explained 9% of the entire variance and appeared to represent the difference

between *E. coli* and LAB (Fig. 9B). GO analysis showed that the genes correlating to PC2 were enriched

with the categories such as glucuronosyltransferase activity (GO:0015020; $p=1.0 \times 10^{-10}$; Enrichment, 6.7)

and biological process involved in interspecies interaction between organisms (GO:0044419; $p=1.0 \times 10^{-34}$;

Enrichment, 5.3) (Figs. S12B and S12D). Among these categories, the neuropeptide signaling pathway

(GO:0007218; $p=5.1 \times 10-4$; Enrichment, 2.9) is particularly intriguing because it can work as signals

from the intestine to neurons.

The principal component analysis revealed the feature of gene expression of animals of different

ages and diets. However, it did not explain the difference in thermotaxis ability in (1) D5 animals fed *E.*

coli or homo fermentative LAB and (2) D5 animals fed hetero fermentative LAB. Therefore, we

investigated differentially expressed genes between these two groups (Fig. 9A). We found 65 genes

whose expression is >2 times higher in Group 1 (*E. coli* and Homofermentative LAB) than in Group 2

(Heterofermentative LAB). These genes included fungus-induced protein-related genes (*fipr-22*, *fipr-23*,

and *fipr-24*) (Fig. 9C) and were enriched with the category such as regulation of lipid localization

(GO:1905952; p=1.3×10^{-4}; Enrichment, 29.0) (Fig. 9D). On the other hand, we found 71 genes whose

expression is >2 times higher in Group 2 than in Group 1. These genes included lysozyme genes (*lys-5*

and *lys-6*) (Fig. 9C) and were enriched with the category such as regulation of defense response to other

organisms (GO:0098542; p=2.0×10^{-5}; Enrichment, 6.6) (Fig. 9D).

Since the high thermotaxis ability of *Lb. reuteri*-fed aged animals were *daf-16*-dependent, we

investigated the enrichment of *daf-16* targets in the whole body (1663 upregulated genes and 1733

downregulated genes (Tepper et al., 2013)). The genes highly expressed in aged animals fed *E. coli* or

homofermentative LAB were significantly enriched with upregulated genes by DAF-16 (Normalized

Enrichment Score (NES), -1.34; False Discovery Rate (FDR), 0.11; p<0.001) (Fig. 9E). On the other

hand, the genes highly expressed in heterofermentative LAB were significantly enriched with

downregulated genes by DAF-16 (NES, 1.53; FDR, 0.046; p<0.001) (Fig. 9E). This result implied that

DAF-16 might be activated in aged animals fed *E. coli* or homofermentative LAB. Interestingly, this is

the opposite of the prediction based on the *daf-16* dependency of the thermotaxis of *Lb. reuteri*-fed

animals. Namely, *daf-16* is predicted to be activated in heterofermentative *Lb. reuteri*-fed aged animals.

Therefore, *Lb. reuteri* might activate DAF-16 in a tissue and/or target-specific manner. Consistent with

this notion, the general activation of DAF-16 using *daf-2* mutants did not improve the thermotaxis ability

of aged animals (Fig. 7D).

Discussion

Addressing the causal relationship between diets and their effects on animals' physiology is challenging in human or mammalian models because microbiota in the gut and diet are complex. It is especially true in the context of aging because of their long lifespan. Using *C. elegans* as a model, we provide evidence of the dietary effect on the age-dependent behavioral decline discernible from the lifespan.

Aging and diet show various effects on different behaviors and organismal lifespan

This study demonstrated that diets affect the age-dependent decline of thermotaxis behavior. We ruled out the possibility that the high performance of LAB-fed aged animals was due to thermophilicity, stronger association to LAB, better motility, dietary restriction, or longer lifespan. Thus, aging and diets likely affect the thermosensory circuit. The primary thermosensory neuron AFD (Mori & Ohshima, 1995) cell-autonomously stores temperature memory (Kobayashi et al., 2016). Although the Ca^{2+} response in AFD is reported to be defective in aged animals (Huang et al., 2020), the temperature sensation itself does not seem to be abolished in aged animals because they could migrate down the gradient (Figs. 1B and 1C). AFD thermosensory neurons synapse onto and regulate AIY interneurons by switching excitatory and inhibitory signals in a context-dependent manner (Mori & Ohshima, 1995; Nakano et al., 2020; White, Southgate, Thomson, & Brenner, 1986). AIY neurons are reported to be a cite of action of *age-1* PI3 kinase, which is upstream of *daf-16* in isothermal tracking behavior (H. Murakami et al., 2005). Therefore, aging and diet might affect AIY interneuorns.

The degree of age-dependent decline seems to depend on behaviors. *E. coli*-fed animals experienced an age-dependent decline in thermotaxis but not in salt avoidance behavior. Despite the similar effects of *P. pentosaceus, Lb. reuteri, Lb plantarum,* and *Lb. rhamnosus* on thermotaxis in aged animals, these LAB showed various effects on locomotion. In thermotaxis, aged animals showed more severe defects in migrating up the thermal gradient than migrating down (Figs. 1B and 1C). Thermotaxis is achieved by multiple steps: sensing temperature, recognizing food, associating food and temperature,

memorizing T_{cult}, and migrating toward T_{cult} (Aoki & Mori, 2015; Goodman & Sengupta, 2018; Kimata,

Sasakura, Ohnishi, Nishio, & Mori, 2012). Thus, the different severities of thermotaxis decline between

migration up and down the gradient in aged animals might be attributed to the different neural circuits

responsible for those conditions, as previously reported (M. Ikeda et al., 2020).

Neuronal aging is discernible from an organismal lifespan. *nkat-1* mutants prevent age-dependent

memory decline in associative learning between food and butanone without changing lifespan (Vohra et

al., 2018). Similarly, we found that *Lb. reuteri* improved thermotaxis in aged animals without changing

their lifespan. More strikingly, *Lb plantarum* and *Lb. rhamnosus* shortened the lifespan while they had

beneficial effects on the thermotaxis of D5 adults. This different dietary condition will allow us to address

the mechanism underlying phenotypic variation in aged animals independent from organismal lifespan

and genetic perturbation.

Diets affect age-dependent thermotaxis decline as nutrients

Previous reports elucidated how bacterial diet affects *C. elegans* as nutritional components, gut

microbiota, and/or pathogen (Kumar et al., 2019; J. J. Zhou, Chun, & Liu, 2019) probably by changing *C.*

elegans metabolites (Gao et al., 2017; Reinke, Hu, Sykes, & Lemire, 2010) and gene expression

(MacNeil, Watson, Arda, Zhu, & Walhout, 2013).

Both live *E. coli* and LAB can colonize animals (Berg et al., 2016; Chelliah et al., 2018; Park et

al., 2018; Portal-Celhay, Bradley, & Blaser, 2012). Live bacteria are necessary for some physiological

roles; secreted enterobactin from live *E. coli* in the gut promotes *C. elegans* growth (Qi & Han, 2018);

live, but not dead, LAB reduces the susceptibility to pathogenic bacteria *Pseudomonas aeruginosa*. On

the other hand, live bacteria are unnecessary in different contexts; heat-killed *Lb. paracasei* and

Bifidobacterium longum extend *C. elegans* lifespan (Sugawara & Sakamoto, 2018; Wang et al., 2020). In

our thermotaxis assay on aged animals, *E. coli* and LAB killed by 65 °C treatment had similar effects to

live bacteria. This result implies that, instead of the action of live bacteria, bacterial nutrition might be

responsible for the effect on the thermotaxis of aged *C. elegans*. Feeding mixed bacteria suggests that *E.*

coli has a dominant effect over *Lb. reuteri* to reduce the thermotaxis ability of aged animals. *E. coli*

appeared to have components that are vulnerable at 100 °C and diffusible after crushing. Since the smell

of bacteria did not reverse the effect of diet, the ingestion of bacterial nutrition causes the dietary effects.

Metabolites in bacterial diet affect *C. elegans* physiology; some metabolites are beneficial, while others

are toxic (J. J. Zhou et al., 2019). Coenzyme Q in *E. coli* shortens the lifespan of *C. elegans* (Larsen &

Clarke, 2002). Bacterial nitric oxide and folate are also positive and negative regulators of *C. elegans*

lifespan, respectively (Gusarov et al., 2013; Virk et al., 2012). Vitamin 12 in *Comamonas aquatica*

accelerates development and reduces fertility without changing lifespan (Watson et al., 2014). Given that

different metabolites are produced by different LAB (Tomita, Saito, Nakamura, Sekiyama, & Kikuchi,

2017), these metabolites might be responsible for the different effects on the thermotaxis of aged *C.*

elegans.

Our results indicated that LAB associated with high performance indices of thermotaxis are

associated with a clade enriched in heterofermentative *Lactobacilli* and *Pediococci* (Clade A in Figure

5A). Heterofermentative LAB produce not only lactic acid and ATP but also several other end products

such as ethanol and CO_2 from glucose. On the other hand, homofermentative LAB converts glucose into

two molecules of lactic acid and ATP. Heterolactic fermentation itself does not explain the high

performance index in thermotaxis of aged animals because heterofermentative *Leuconostoc* and

Bifidobacteria species did not give the high performance indices. Metabolites other than lactic acid,

ethanol, and CO_2 also differ between hetero- and homofermentative *Lactobacilli* (Tomita et al., 2017).

Metabolites enriched in heterofermentative *Lactobacilli* include a neurotransmitter GABA and tyramine,

a substrate to synthesize neurotransmitter octopamine; metabolites enriched in homofermentative

Lactobacilli include 4-hydroxyphenyllactic acid (HPLA) and acetoin. We note that Tomita *et al*. reported

the metabolites in the media (Tomita et al., 2017) while we supply bacteria to animals after washing off

the bacterial media. Nonetheless, metabolites enriched in different *Lactobacilli* might affect the age-

dependent thermotaxis decline.

Bacterial diets modulate genetic pathways in *C. elegans* neurons

LAB can extend the lifespan of *C. elegans* either by dietary restriction-dependent (Y. Zhao et al., 2013) or

independent mechanisms (Komura et al., 2013; Nakagawa et al., 2016). The mechanism underlying the

dietary effect on the thermotaxis decline does not seem to depend on the activation of the dietary

restriction pathway. First, the expression of *pha-4* was low. Second, the lifespan of LAB-fed animals was

not necessarily prolonged. Third, *eat-2* mutants, which mimic dietary restriction, did not improve

thermotaxis in aged animals fed *E. coli.* Fourth, *kmo-1* and *nkat-1* genes involved in dietary restriction-

dependent beneficial effects on associative learning (Vohra, Lemieux, Lin, & Ashrafi, 2017) did not

affect the dietary effects on thermotaxis of aged animals. Fifth, no correlation was observed between the

thermotaxis ability of aged animals and body size or fat accumulation.

400   Different LAB activate distinct genetic pathways such as insulin and IGF-1 signaling (IIS)

pathway important for lifespan regulation and p38 mitogen-activated protein kinase (MAPK) pathway

important for innate immunity. *Lb. rhamnosus* and *B. longum* extend the lifespan of *C. elegans* by

modulating the IIS pathway consisting of DAF-2 and DAF-16 (Grompone et al., 2012; Sugawara &

Sakamoto, 2018). *B. infantis* extends the lifespan of *C. elegans* via the PMK-1 p38 MAPK pathway and a

downstream transcription factor SKN-1, an ortholog of mammalian Nrf, but not via DAF-16 (Komura et

al., 2013). The PMK-1 pathway is also activated by *Lb. acidophilus* and *Lactobacillus fermentum* (Y.

Kim & Mylonakis, 2012; Park et al., 2018). Animals fed a lactic acid bacterium, *Weissella*, show higher

expression of *daf-16*, *aak-2,* and *jnk-1,* and extend lifespan in these genes-dependent manners (Lee et al.,

2015). the *daf-16* pathway is also involved in thermotaxis (Kodama et al., 2006; H. Murakami et al.,

2005) and salt-avoidance behaviors (Tomioka et al., 2006). In our condition, *daf-16* did not strongly

affect thermotaxis at D1, while it was necessary for the maintenance of thermotaxis of *Lb. reuteri*-fed

aged animals. This result suggests that DAF-16 had a specific role in the *Lb reuteri*'s effects on aged

animals. Since *daf-16* functions in neurons (Fig. 8C) and has neuron-specific targets (Kaletsky et al.,

2016), differential expressions of these genes with different diets might affect thermotaxis behavior.

Indeed, our transcriptome analysis revealed that differentially expressed genes between animals fed *E.*

coli or homofermentative LAB and those fed heterofermentative LAB were enriched with DAF-16

targets. Interestingly, our transcriptome data suggested that DAF-16 was activated in animals fed *E. coli*

or homofermentative LAB although heterofermentative *Lb. reuteri*-fed aged animals showed *daf-16*

dependency on thermotaxis. The regulation of DAF-16 in specific neurons might be opposite from that in

the whole body. Single-cell transcriptome analysis for aged animals fed different diets will address this

question in the future (Cao et al., 2017).

Bacterial screen to address age-dependent phenotypes

Even with *C. elegans*, it is challenging to address age-dependent neuronal phenotypes because powerful

forward genetic screens are not readily applicable to aged animals. Our study showed that bacterial screen

could be useful for generating phenotypic diversity and addressing underlying molecular mechanisms in

aged animals. The bacterial screen has been applied to various *C. elegans* phenotypes. Watson *et al.*

carried out unbiased mutant screens of *Escherichia coli* and *Comamonas aquatica* to identify bacterial

genes that affect the "dietary sensor" in *C. elegans*, which increases the GFP intensity when fed

Comamonas; they found that mutations in genes involved in vitamin B12 biosynthesis/import increase *C.*

elegans dietary sensor activity (Watson et al., 2014). Zhou et al. screened 13 LAB and found that

Lactobacillus zeae protects *C. elegans* from enterotoxigenic *E. coli* (M. Zhou et al., 2014). Given that *C.*

elegans has its natural microbiota (Berg et al., 2016; Dirksen et al., 2016; Samuel et al., 2016; Zhang et

al., 2017), the nervous system of animals in a natural environment may be affected by complex bacteria.

Indeed, a recent study has revealed that tyramine produced from commensal bacteria affects *C. elegans*

avoidance behavior (O'Donnell et al., 2020). Hence, bacterial screens will provide a unique angle of

understanding for *C. elegans* research.

Materials and Methods

Worm maintenance and strains

C. elegans strains were maintained at 23 °C on Nematode Growth Medium (NGM) plates with *E. coli*, OP50, as previously reported (Brenner, 1974), except CB1370, which was maintained at 15 °C. N2 (Bristol) was used as the wild type. The following mutant strains were used for thermotaxis assays: CB1370 *daf-2(e1370ts)*, DA1116 *eat-2(ad1116)*, CF1038 *daf-16(mu86)*, IK0656 *tax-6(db60)*, NUJ69 *kmo-1(tm4529)*, NUJ71 *nkat-1(ok566)*. NUJ69 *kmo-1(tm4529)* is a one-time outcrossed FX04529 *kmo-1(tm4529)* strain. NUJ71 *nkat-1(ok566)* is a two-time outcrossed RB784 *nkat-1(ok566)* strain. The transgenic strains were generated as described below.

CRISPR knockout

To generate a *daf-16* allele that affects only the exon of the b isoform, we used the Co-CRISPR strategy (H. Kim et al., 2014). tracrRNA, *daf-16* crRNA (5'-GUCAUGCCAGAUGAAGAACA-3'), and *dpy-10* crRNA (5'-GCUACCAUAGGCACCACGAG-3') were annealed and injected into N2 animals with *eft-3*p::Cas9::NLS-3'UTR(*tbb-2*) plasmids. After picking dumpy and/or roller F_1 animals on individual plates, the mutation of the *daf-16* locus was detected by PCR and confirmed using Sanger sequencing. We obtained *knj36* which introduced 7-bp deletion in the first exon of the *daf-16* b isoform, corresponding to the intron of other isoforms. NUJ298 *daf-16(knj36) I* was used for thermotaxis assays.

Plasmids and single-copy insertion of transgenes

Single-copy insertions of transgenes were generated using Cas9-based homologous recombination following the established protocol (Andrusiak et al., 2019). *daf-16*bp::*daf-16*b (gDNA) was cloned into the pCR8 vector (Thermo Fisher Scientific) vector using Gibson Assembly (New England Biolabs) (pKEN891). pKEN891 was recombined with the pCZGY2729 repair template vector using the LR reaction (Gateway Technology, Thermo Fisher Scientific) to generate the pKEN1016 repair template

vector. pKEN1016 contains homology arms for *cxTi10882* site on chromosome IV and *rps-0*p::Hyg^R in

addition to the *daf-16*b promoter (4.9 kbp) and *daf-16*b gDNA. To swap promoters of pKEN1016, XhoI

and AgeI sites were introduced before and after the *daf-16*b promoter, respectively (pKEN973).

pKEN973 was digested with XhoI and AgeI, and the promoter was substituted to *myo-2*p (1.0 kbp,

pKEN976), *myo-3*p (2.4 kbp, pKEN975), or *rgef-1*p (3.5 kbp, pKEN974) using Gibson assembly.

We injected the repair template described above and *eft-3*p::Cas9 + *cxTi10882* sgRNA

(pCZGY2750) into N2 or CF1038 *daf-16(mu86)* with three red markers originally used for MosSCI

insertion (Frøkjær-Jensen et al., 2008): *rab-3*p::mCherry (pGH8), *myo-2*p::mCherry(pCFJ90), and *myo-*

*3*p::mCherry (pCFJ104). The animals with the single copy insertion were selected based on the

hygromycin-resistance (Hyg^R) and the absence of red fluorescence. The following strains were used for

the rescue experiments of thermotaxis assays: NUJ306 *daf-16(mu86) I; knjSi17[daf-16bp::daf-16b] IV*,

NUJ368 daf-16(mu86) I; knjSi19[myo-2p::daf-16b] IV, NUJ373 *daf-16(mu86) I; knjSi18[myo-3p::daf-*

16b] IV, NUJ372 *daf-16(mu86) I; knjSi22[rgef-1p::daf-16b] IV*.

Bacterial plates

E. coli, OP50 or HT115, was inoculated into Super Broth (32 g Bacto Tryptone (BD), 20 g Bacto Yeast

extract (BD), 5 g NaCl (Wako), 5 mL 1M NaOH in 1 L water) and cultured overnight at 37 °C. LAB

strains, provided by Megmilk Snow Brand company (Table S1), were inoculated into the liquid medium

from glycerol stocks and cultured in the conditions described in Supplementary Table 1. Bacterial cells

were collected by centrifugation at 7,000x g for 10 min at 4 °C. Cells were washed twice with sterile

0.9% NaCl solution. The washed bacteria were adjusted to a final concentration of 0.1 g/ml (wet weight)

in NG buffer (25 mM K-PO$_4$ (pH6), 50 mM NaCl, 1 mM CaCl$_2$, 1 mM MgSO$_4$). For heat killing, 0.1 g/ml

bacteria in tubes were incubated for 1 h in a 65 °C incubator or 10 min in boiled water. By this treatment,

bacterial colony-forming unit (cfu) became $<1.0\times10^2$ cfu/ml, which is at least 10^8 lower than live bacteria

(>1.5x10^{10}). For the mixed condition, bacteria were mixed to make the final concentration of 0.1 g/ml in

total before spread on NGM plates. To crush bacteria, bacterial suspension was vigorously vibrated with

glass beads at 4,200 rpm for 15 cycles of 30-sec ON and 30-sec OFF using a bead-based homogenizer

(MS-100R, Tomy). Two hundred microliters of the bacterial suspension were spread onto 60-mm NGM

plates and dried overnight. NGM plates with peptone were used except for thermotaxis to see the effect of

peptone and lifespan assays, in which NGM plates without peptone were used.

Preparation of aged animals fed different bacteria

For behavioral assays, synchronized eggs were prepared by bleaching gravid hermaphrodites using 0.5x

household bleach in 0.5 M NaOH and placed onto NGM plates with OP50. The eggs were cultivated at

23 °C for 72 hours to obtain day one adults (D1). For the thermotaxis of *daf-2* mutants, eggs of N2 and

CB1370 *daf-2(e1370)* were incubated at 15 °C for 96 hours and subsequently at 23 °C for 24 hours to

obtain D1. D1 animals were washed with NG buffer and transferred to NGM plates with OP50 or LAB

every day for thermotaxis of aged animals. To expose animals to the bacterial odor, NG buffer as control

or 0.1 g/ml bacterial solution was spotted on the lid of bacterial plates described above. Animals were

cultivated on the upside-down plates as described in Figure 5D. For the thrashing and locomotion assay,

animals were transferred individually by picking instead of washing.

Thermotaxis assay

Animals were cultivated at 23 °C (T$_{cult}$=23 °C) unless otherwise noted. Population thermotaxis assays

with a linear thermal gradient were performed as described (Ito et al., 2006). Fifty to 250 animals on

cultivation plates were washed with M9 and placed at the center of the assay plates without food and with

a temperature gradient of 17-23 or 20-26 °C. The temperature gradient was measured to be ~0.5 °C/cm.

After letting them move for 1 h, the animals were killed by chloroform. The number of adult animals in

each of eight sections along the temperature gradient (Fig. 1A) was scored under a stereomicroscope. The

fraction of animals in each section was plotted on histograms. The performance and thermotaxis indices

were calculated, as shown in Fig. 1A and Fig. 3C, respectively. For temperature shift assays, T_{cult} was

shifted 24 h prior to the assay.

Thrashing assay

A thrashing assay was performed, as previously described with a few modifications (Tsalik & Hobert,

2003). Animals were washed with NG buffer and transferred with a drop of NG buffer onto an NGM

plate without food using a capillary glass pipet. In liquid, animals show lateral swimming movements

(thrashes). We defined a single thrash as a complete movement through the midpoint and back and

counted the number of thrashes for 30 seconds.

Motility assay

Assay plates were prepared by placing circular filter paper with a one-inch hole on NGM plates with

OP50 or LAB and soaking the paper with ~100 µl of 100 mM $CuCl_2$. A single animal was transferred to

an assay plate with the cultured bacteria and left at 23 °C for three minutes. The images of the bacterial

lawn were captured by a digital camera (Fujifilm) through an eyepiece of a stereomicroscope, Stemi 508

(Zeiss). The trajectory of an animal on the lawn was traced using FIJI (Schindelin et al., 2012) and

measured as the distance of locomotion.

Lifespan assay

Animals were synchronized by bleaching gravid adults and grown with regular NGM plates with OP50

until D1. D1 animals were washed three times with M9 buffer and transferred to peptone-free NGM

plates supplemented with 50 mg/ml OP50 or LAB. Animals were transferred to new plates every day

until they became D4 and every other day afterward. Dead animals were defined as having no voluntary

movement after several touches on the head and tail and counted every day. Four independent sessions

with 25 animals per session were combined for each condition.

Phylogenetic tree

16S rRNA sequences of LAB were obtained from the Genome database of NCBI

(http://www.ncbi.nlm.nih.gov/genome/). The accession numbers are shown in Table S1. The phylogenetic

tree was inferred by the Neighbor-Joining method based on the 16S rRNA gene sequence of model LAB

strains. The evolutionary distances were computed using the Maximum Composite Likelihood method

conducted in MEGA X (Hall, 2013).

RNA sequencing

Non-gravid young adult animals were used to avoid the effect of eggs inside the body. D5 animals fed *E.*

coli or LAB (*Lb. gasseri*; *Lb. delbrueckii*, *P. pentosaceus*, *Lb. reuteri*, *Lb. rhamnosus*, and *Lb. plantarum)*

were prepared as described above. Total RNA was extracted from whole animals using RNAiso Plus

reagent (Takara) and sequenced using NovaSeq 6000 System by Macrogen Corp. Japan. We detected

13329 genes. The fragments per kilobase of exon per million mapped fragments (FPKM) of D1, D5 (*E.*

coli), and D5 (LAB, 6 in total) were used to perform the principal component analysis (PCA) based on the

variance-covariance matrix using R program. The eigenvalues of each transcript were ranked by the

percentage of explained variance for each principal component, and the top and bottom 5% (666 genes

each) were used to perform gene ontology (GO) analysis using Metascape (Y. Zhou et al., 2019).

To further analyze the difference between D5 animals fed *E. coli* or homofermentative LAB (*Lb.*

gasseri; *Lb. delbrueckii*) and D5 animals fed heterofermentative LAB (*P. pentosaceus*, *Lb. reuteri*, *Lb.*

rhamnosus, and *Lb. plantarum*), we compared the expression of genes between these two groups and

extracted differentially expressed genes as those with p-value<0.05 by Student's t-test. Those

differentially expressed genes were used to perform GO analysis using Metascape and compared with

DAF-16-regulated genes (Tepper et al., 2013). Gene enrichment was calculated using Gene Set

Enrichment Analysis (GSEA) (Subramanian et al., 2005).

Statistical analyses

Box-and-whisker plots represent medians as center lines; boxes as first and third quartiles; whiskers as

maximum and minimum values except for outliers, which are 1.5 times greater than the upper limit or 1.5

times smaller than the lower limit of the interquartile range; dots as outliers. In some figures, all the data

points were overlayed on the box-and-whisker plots. We used Student's t-test to compare two samples

and one-way or two-way ANOVA, followed by Dunnett's or Tukey-Kramer test to compare multiple

samples using R (R core team, https://www.R-project.org/, Vienna, Austria) or GraphPad Prism 7.0

(GraphPad Software, La Jolla, CA). In all figures, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$. $p > 0.05$ is

considered as not significant (ns).

Acknowledgments

Megmilk Snow Brand Company supported this work. We thank members of the Nutritional Neuroscience

laboratory and the Mori laboratory for their comments on the manuscript. Wei Huang and

Pauline Rouillard helped with basic experiments. *C. elegans* mutant strains were provided by

Caenorhabditis Genetics Center (CGC), funded by the NIH Office of Research Infrastructure Programs

(P40 OD010440), and Dr. Shoehei Mitani of the National Bioresource Project of Japan. Dr. Yishi Jin

provided pCZGY2729 and pCZGY2750 plasmids.

Competing interests

S.H. and S.T. are former employees of Megmilk Snow Brand Co., Ltd. M.T. is an employee of Megmilk

Snow Brand Co., Ltd. The other authors declare no competing interests.

References

Andrusiak, M. G., Sharifnia, P., Lyu, X., Wang, Z., Dickey, A. M., Wu, Z., . . . Jin, Y. (2019). Inhibition

of Axon Regeneration by Liquid-like TIAR-2 Granules. *Neuron, 104*(2), 290-304 e298.

doi:10.1016/j.neuron.2019.07.004

Aoki, I., & Mori, I. (2015). Molecular biology of thermosensory transduction in *C. elegans*. *Curr Opin*

Neurobiol, 34, 117-124. doi:S0959-4388(15)00066-5 [pii]

10.1016/j.conb.2015.03.011

Arey, R. N., Stein, G. M., Kaletsky, R., Kauffman, A., & Murphy, C. T. (2018). Activation of Galphaq

Signaling Enhances Memory Consolidation and Slows Cognitive Decline. *Neuron, 98*(3), 562-

574 e565. doi:10.1016/j.neuron.2018.03.039

Beck, C. D., & Rankin, C. H. (1993). Effects of aging on habituation in the nematode *Caenorhabditis*

elegans. *Behav Processes, 28*(3), 145-163. doi:10.1016/0376-6357(93)90088-9

Berg, M., Stenuit, B., Ho, J., Wang, A., Parke, C., Knight, M., . . . Shapira, M. (2016). Assembly of the

Caenorhabditis elegans gut microbiota from diverse soil microbial environments. *ISME J, 10*(8),

1998-2009. doi:10.1038/ismej.2015.253

Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics, 77*(1), 71-94.

Cao, J., Packer, J. S., Ramani, V., Cusanovich, D. A., Huynh, C., Daza, R., . . . Shendure, J. (2017).

Comprehensive single-cell transcriptional profiling of a multicellular organism. *Science,*

357(6352), 661-667. doi:10.1126/science.aam8940

Chelliah, R., Choi, J. G., Hwang, S. B., Park, B. J., Daliri, E. B., Kim, S. H., . . . Oh, D. H. (2018). In

vitro and in vivo defensive effect of probiotic LAB against *Pseudomonas aeruginosa* using

Caenorhabditis elegans model. *Virulence, 9*(1), 1489-1507. doi:10.1080/21505594.2018.1518088

Christensen, R., de la Torre-Ubieta, L., Bonni, A., & Colon-Ramos, D. A. (2011). A conserved

PTEN/FOXO pathway regulates neuronal morphology during *C. elegans* development.

Development, 138(23), 5257-5267. doi:10.1242/dev.069062

Cornils, A., Maurya, A. K., Tereshko, L., Kennedy, J., Brear, A. G., Prahlad, V., . . . Sengupta, P. (2016).

Structural and Functional Recovery of Sensory Cilia in *C. elegans* IFT Mutants upon Aging.

PLoS Genet, 12(12), e1006325. doi:10.1371/journal.pgen.1006325

Deary, I. J., Corley, J., Gow, A. J., Harris, S. E., Houlihan, L. M., Marioni, R. E., . . . Starr, J. M. (2009).

Age-associated cognitive decline. *Br Med Bull, 92*, 135-152. doi:10.1093/bmb/ldp033

Dirksen, P., Marsh, S. A., Braker, I., Heitland, N., Wagner, S., Nakad, R., . . . Schulenburg, H. (2016).

The native microbiome of the nematode *Caenorhabditis elegans*: gateway to a new host-

microbiome model. *BMC Biol, 14*, 38. doi:10.1186/s12915-016-0258-1

Dong, X., Milholland, B., & Vijg, J. (2016). Evidence for a limit to human lifespan. *Nature, 538*(7624),

257-259. doi:10.1038/nature19793

Frøkjær-Jensen, C., Davis, M. W., Hopkins, C. E., Newman, B. J., Thummel, J. M., Olesen, S. P., . . .

Jorgensen, E. M. (2008). Single-copy insertion of transgenes in *Caenorhabditis elegans*. *Nat*

Genet, 40(11), 1375-1383. doi:ng.248 [pii]

10.1038/ng.248

Gao, A. W., Chatzispyrou, I. A., Kamble, R., Liu, Y. J., Herzog, K., Smith, R. L., . . . Houtkooper, R. H.

(2017). A sensitive mass spectrometry platform identifies metabolic changes of life history traits

in *C. elegans*. *Sci Rep, 7*(1), 2408. doi:10.1038/s41598-017-02539-w

Glenn, C. F., Chow, D. K., David, L., Cooke, C. A., Gami, M. S., Iser, W. B., . . . Wolkow, C. A. (2004).

Behavioral deficits during early stages of aging in Caenorhabditis elegans result from locomotory

deficits possibly linked to muscle frailty. *J Gerontol A Biol Sci Med Sci, 59*(12), 1251-1260.

doi:10.1093/gerona/59.12.1251

Goodman, M. B., & Sengupta, P. (2018). The extraordinary AFD thermosensor of *C. elegans*. *Pflugers*

Arch, 470(5), 839-849. doi:10.1007/s00424-017-2089-5

Grompone, G., Martorell, P., Llopis, S., Gonzalez, N., Genoves, S., Mulet, A. P., . . . Ramon, D. (2012).

Anti-inflammatory *Lactobacillus rhamnosus* CNCM I-3690 strain protects against oxidative

stress and increases lifespan in *Caenorhabditis elegans*. *PLoS ONE, 7*(12), e52493.

doi:10.1371/journal.pone.0052493

Gusarov, I., Gautier, L., Smolentseva, O., Shamovsky, I., Eremina, S., Mironov, A., & Nudler, E. (2013).

Bacterial nitric oxide extends the lifespan of *C. elegans*. *Cell, 152*(4), 818-830.

doi:10.1016/j.cell.2012.12.043

Hahm, J. H., Kim, S., DiLoreto, R., Shi, C., Lee, S. J., Murphy, C. T., & Nam, H. G. (2015). *C. elegans*

maximum velocity correlates with healthspan and is maintained in worms with an insulin receptor

mutation. *Nat Commun, 6*, 8919. doi:10.1038/ncomms9919

Hall, B. G. (2013). Building phylogenetic trees from molecular data with MEGA. *Mol Biol Evol, 30*(5),

1229-1235. doi:10.1093/molbev/mst012

Hedgecock, E. M., & Russell, R. L. (1975). Normal and mutant thermotaxis in the nematode

Caenorhabditis elegans. *Proc Natl Acad Sci U S A, 72*(10), 4061-4065.

Hill, C., Guarner, F., Reid, G., Gibson, G. R., Merenstein, D. J., Pot, B., . . . Sanders, M. E. (2014).

Expert consensus document. The International Scientific Association for Probiotics and

Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev*

Gastroenterol Hepatol, 11(8), 506-514. doi:10.1038/nrgastro.2014.66

Hoogewijs, D., Houthoofd, K., Matthijssens, F., Vandesompele, J., & Vanfleteren, J. R. (2008). Selection

and validation of a set of reliable reference genes for quantitative sod gene expression analysis in

C. elegans. *BMC Mol Biol, 9*, 9. doi:10.1186/1471-2199-9-9

1471-2199-9-9 [pii]

Huang, T. T., Matsuyama, H. J., Tsukada, Y., Singhvi, A., Syu, R. T., Lu, Y., . . . Pan, C. L. (2020). Age-

dependent changes in response property and morphology of a thermosensory neuron and

thermotaxis behavior in *Caenorhabditis elegans*. *Aging Cell*. doi:10.1111/acel.13146

Ikeda, M., Nakano, S., Giles, A. C., Xu, L., Costa, W. S., Gottschalk, A., & Mori, I. (2020). Context-

dependent operation of neural circuits underlies a navigation behavior in *Caenorhabditis elegans*.

Proc Natl Acad Sci U S A, 117(11), 6178-6188. doi:10.1073/pnas.1918528117

Ikeda, T., Yasui, C., Hoshino, K., Arikawa, K., & Nishikawa, Y. (2007). Influence of lactic acid bacteria

on longevity of *Caenorhabditis elegans* and host defense against salmonella enterica serovar

enteritidis. *Appl Environ Microbiol, 73*(20), 6404-6409. doi:10.1128/AEM.00704-07

Ito, H., Inada, H., & Mori, I. (2006). Quantitative analysis of thermotaxis in the nematode *Caenorhabditis*

elegans. *J Neurosci Methods, 154*(1-2), 45-52. doi:10.1016/j.jneumeth.2005.11.011

Johnke, J., Dirksen, P., & Schulenburg, H. (2020). Community assembly of the native *C. elegans*

microbiome is influenced by time, substrate and individual bacterial taxa. *Environ Microbiol,*

22(4), 1265-1279. doi:10.1111/1462-2920.14932

Joseph, J., Cole, G., Head, E., & Ingram, D. (2009). Nutrition, brain aging, and neurodegeneration. *J*

Neurosci, 29(41), 12795-12801. doi:10.1523/JNEUROSCI.3520-09.2009

Kaletsky, R., Lakhina, V., Arey, R., Williams, A., Landis, J., Ashraf, J., & Murphy, C. T. (2016). The *C.*

elegans adult neuronal IIS/FOXO transcriptome reveals adult phenotype regulators. *Nature,*

529(7584), 92-96. doi:10.1038/nature16483

nature16483 [pii]

Kauffman, A. L., Ashraf, J. M., Corces-Zimmerman, M. R., Landis, J. N., & Murphy, C. T. (2010).

Insulin signaling and dietary restriction differentially influence the decline of learning and

memory with age. *PLoS Biol, 8*(5), e1000372. doi:10.1371/journal.pbio.1000372

Kenyon, C., Chang, J., Gensch, E., Rudner, A., & Tabtiang, R. (1993). A *C. elegans* mutant that lives

twice as long as wild type. *Nature, 366*(6454), 461-464. doi:10.1038/366461a0

Kim, H., Ishidate, T., Ghanta, K. S., Seth, M., Conte, D., Jr., Shirayama, M., & Mello, C. C. (2014). A co-

CRISPR strategy for efficient genome editing in *Caenorhabditis elegans*. *Genetics, 197*(4), 1069-

1080. doi:10.1534/genetics.114.166389

Kim, Y., & Mylonakis, E. (2012). *Caenorhabditis elegans* immune conditioning with the probiotic

bacterium *Lactobacillus acidophilus* strain NCFM enhances gram-positive immune responses.

Infect Immun, 80(7), 2500-2508. doi:10.1128/IAI.06350-11

Kimata, T., Sasakura, H., Ohnishi, N., Nishio, N., & Mori, I. (2012). Thermotaxis of *C. elegans* as a

model for temperature perception, neural information processing and neural plasticity. *Worm,*

1(1), 31-41. doi:10.4161/worm.19504

Klass, M. R. (1977). Aging in the nematode *Caenorhabditis elegans:* major biological and environmental

factors influencing life span. *Mech Ageing Dev, 6*(6), 413-429.

Kobayashi, K., Nakano, S., Amano, M., Tsuboi, D., Nishioka, T., Ikeda, S., . . . Mori, I. (2016). Single-

Cell Memory Regulates a Neural Circuit for Sensory Behavior. *Cell Rep, 14*(1), 11-21.

doi:10.1016/j.celrep.2015.11.064

S2211-1247(15)01420-5 [pii]

Kodama, E., Kuhara, A., Mohri-Shiomi, A., Kimura, K. D., Okumura, M., Tomioka, M., . . . Mori, I.

(2006). Insulin-like signaling and the neural circuit for integrative behavior in *C. elegans*. *Genes*

Dev, 20(21), 2955-2960. doi:10.1101/gad.1479906

Komura, T., Ikeda, T., Yasui, C., Saeki, S., & Nishikawa, Y. (2013). Mechanism underlying prolongevity

induced by bifidobacteria in *Caenorhabditis elegans*. *Biogerontology, 14*(1), 73-87.

doi:10.1007/s10522-012-9411-6

Kuhara, A., Inada, H., Katsura, I., & Mori, I. (2002). Negative regulation and gain control of sensory

neurons by the *C. elegans* calcineurin TAX-6. *Neuron, 33*(5), 751-763. doi:10.1016/s0896-

6273(02)00607-4

Kumar, A., Baruah, A., Tomioka, M., Iino, Y., Kalita, M. C., & Khan, M. (2019). *Caenorhabditis*

elegans: a model to understand host-microbe interactions. *Cell Mol Life Sci*. doi:10.1007/s00018-

019-03319-7

Kwon, E. S., Narasimhan, S. D., Yen, K., & Tissenbaum, H. A. (2010). A new DAF-16 isoform regulates

709   longevity. *Nature, 466*(7305), 498-502. doi:10.1038/nature09184

710   Lakowski, B., & Hekimi, S. (1998). The genetics of caloric restriction in *Caenorhabditis elegans*. *Proc*

711   *Natl Acad Sci U S A, 95*(22), 13091-13096.

712   Larsen, P. L., & Clarke, C. F. (2002). Extension of life-span in *Caenorhabditis elegans* by a diet lacking

713   coenzyme Q. *Science, 295*(5552), 120-123. doi:10.1126/science.1064653

714   Lee, J., Kwon, G., & Lim, Y. H. (2015). Elucidating the Mechanism of *Weissella*-dependent Lifespan

715   Extension in *Caenorhabditis elegans*. *Sci Rep, 5*, 17128. doi:10.1038/srep17128

716   Leinwand, S. G., Yang, C. J., Bazopoulou, D., Chronis, N., Srinivasan, J., & Chalasani, S. H. (2015).

717   Circuit mechanisms encoding odors and driving aging-associated behavioral declines in

718   *Caenorhabditis elegans*. *Elife, 4*, e10181. doi:10.7554/eLife.10181

719   Lin, K., Dorman, J. B., Rodan, A., & Kenyon, C. (1997). *daf-16*: An HNF-3/forkhead family member that

720   can function to double the life-span of *Caenorhabditis elegans*. *Science, 278*(5341), 1319-1322.

721   doi:10.1126/science.278.5341.1319

722   Mack, H. I. D., Heimbucher, T., & Murphy, C. T. (2018). The nematode *Caenorhabditis elegans* as a

723   model for aging research. *Drug Discov Today Dis Model., 27*, 3-13.

724   MacNeil, L. T., Watson, E., Arda, H. E., Zhu, L. J., & Walhout, A. J. (2013). Diet-induced developmental

725   acceleration independent of TOR and insulin in *C. elegans*. *Cell, 153*(1), 240-252.

726   doi:10.1016/j.cell.2013.02.049

727   Mann, F. G., Van Nostrand, E. L., Friedland, A. E., Liu, X., & Kim, S. K. (2016). Deactivation of the

728   GATA Transcription Factor ELT-2 Is a Major Driver of Normal Aging in *C. elegans*. *PLoS*

729   *Genet, 12*(4), e1005956. doi:10.1371/journal.pgen.1005956

730   Miller, K. G., Alfonso, A., Nguyen, M., Crowell, J. A., Johnson, C. D., & Rand, J. B. (1996). A genetic

731   selection for *Caenorhabditis elegans* synaptic transmission mutants. *Proc Natl Acad Sci U S A,*

732   *93*(22), 12593-12598.

733   Mori, I., & Ohshima, Y. (1995). Neural regulation of thermotaxis in *Caenorhabditis elegans*. *Nature,*

734   *376*(6538), 344-348.

735   Mulcahy, B., Holden-Dye, L., & O'Connor, V. (2013). Pharmacological assays reveal age-related changes

736   in synaptic transmission at the *Caenorhabditis elegans* neuromuscular junction that are modified

737   by reduced insulin signalling. *J Exp Biol, 216*(Pt 3), 492-501. doi:10.1242/jeb.068734

738   Murakami, H., Bessinger, K., Hellmann, J., & Murakami, S. (2005). Aging-dependent and -independent

739   modulation of associative learning behavior by insulin/insulin-like growth factor-1 signal in

740   *Caenorhabditis elegans*. *J Neurosci, 25*(47), 10894-10904. doi:10.1523/JNEUROSCI.3600-

741   04.2005

742   Murakami, S., & Murakami, H. (2005). The effects of aging and oxidative stress on learning behavior in

743   *C. elegans*. *Neurobiol Aging, 26*(6), 899-905. doi:10.1016/j.neurobiolaging.2004.08.007

744   Nagashima, T., Iino, Y., & Tomioka, M. (2019). DAF-16/FOXO promotes taste avoidance learning

745   independently of axonal insulin-like signaling. *PLoS Genet, 15*(7), e1008297.

746   doi:10.1371/journal.pgen.1008297

747   Nakagawa, H., Shiozaki, T., Kobatake, E., Hosoya, T., Moriya, T., Sakai, F., . . . Miyazaki, T. (2016).

748   Effects and mechanisms of prolongevity induced by *Lactobacillus gasseri* SBT2055 in

749   *Caenorhabditis elegans*. *Aging Cell, 15*(2), 227-236. doi:10.1111/acel.12431

750   Nakano, S., Ikeda, M., Tsukada, Y., Fei, X., Suzuki, T., Niino, Y., . . . Mori, I. (2020). Presynaptic MAST

751   kinase controls opposing postsynaptic responses to convey stimulus valence in *Caenorhabditis*

752   *elegans*. *Proc Natl Acad Sci U S A, 117*(3), 1638-1647. doi:10.1073/pnas.1909240117

753   O'Donnell, M. P., Fox, B. W., Chao, P. H., Schroeder, F. C., & Sengupta, P. (2020). A neurotransmitter

754   produced by gut bacteria modulates host sensory behaviour. *Nature*. doi:10.1038/s41586-020-

755   2395-5

Ogg, S., Paradis, S., Gottlieb, S., Patterson, G. I., Lee, L., Tissenbaum, H. A., & Ruvkun, G. (1997). The

Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in

C. elegans. *Nature, 389*(6654), 994-999. doi:10.1038/40194

Okochi, Y., Kimura, K. D., Ohta, A., & Mori, I. (2005). Diverse regulation of sensory signaling by *C.

elegans* nPKC-epsilon/eta TTX-4. *Embo J, 24*(12), 2127-2137. doi:10.1038/sj.emboj.7600697

Panowski, S. H., Wolff, S., Aguilaniu, H., Durieux, J., & Dillin, A. (2007). PHA-4/Foxa mediates diet-

restriction-induced longevity of *C. elegans*. *Nature, 447*(7144), 550-555.

doi:10.1038/nature05837

Park, M. R., Ryu, S., Maburutse, B. E., Oh, N. S., Kim, S. H., Oh, S., . . . Kim, Y. (2018). Probiotic

Lactobacillus fermentum strain JDFM216 stimulates the longevity and immune response of

Caenorhabditis elegans through a nuclear hormone receptor. *Sci Rep, 8*(1), 7441.

doi:10.1038/s41598-018-25333-8

Portal-Celhay, C., Bradley, E. R., & Blaser, M. J. (2012). Control of intestinal bacterial proliferation in

regulation of lifespan in *Caenorhabditis elegans*. *BMC Microbiol, 12*, 49. doi:10.1186/1471-

2180-12-49

Qi, B., & Han, M. (2018). Microbial Siderophore Enterobactin Promotes Mitochondrial Iron Uptake and

Development of the Host via Interaction with ATP Synthase. *Cell, 175*(2), 571-582 e511.

doi:10.1016/j.cell.2018.07.032

Reinke, S. N., Hu, X., Sykes, B. D., & Lemire, B. D. (2010). *Caenorhabditis elegans* diet significantly

affects metabolic profile, mitochondrial DNA levels, lifespan and brood size. *Mol Genet Metab,

100*(3), 274-282. doi:10.1016/j.ymgme.2010.03.013

Saeki, S., Yamamoto, M., & Iino, Y. (2001). Plasticity of chemotaxis revealed by paired presentation of a

chemoattractant and starvation in the nematode *Caenorhabditis elegans*. *J Exp Biol, 204*(Pt 10),

1757-1764.

Samuel, B. S., Rowedder, H., Braendle, C., Felix, M. A., & Ruvkun, G. (2016). *Caenorhabditis elegans*

responses to bacteria from its natural habitats. *Proc Natl Acad Sci U S A, 113*(27), E3941-3949.

doi:10.1073/pnas.1607183113

Sawin, E. R., Ranganathan, R., & Horvitz, H. R. (2000). *C. elegans* locomotory rate is modulated by the

environment through a dopaminergic pathway and by experience through a serotonergic pathway.

Neuron, 26(3), 619-631.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., . . . Cardona, A.

(2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods, 9*(7), 676-682.

doi:10.1038/nmeth.2019

nmeth.2019 [pii]

Shtonda, B. B., & Avery, L. (2006). Dietary choice behavior in *Caenorhabditis elegans*. *J Exp Biol,*

209(Pt 1), 89-102. doi:10.1242/jeb.01955

Stein, G. M., & Murphy, C. T. (2012). The Intersection of Aging, Longevity Pathways, and Learning and

Memory in *C. elegans*. *Front Genet, 3*, 259. doi:10.3389/fgene.2012.00259

Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., . . . Mesirov, J.

P. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-

wide expression profiles. *Proc Natl Acad Sci U S A, 102*(43), 15545-15550.

doi:10.1073/pnas.0506580102

Sugawara, T., & Sakamoto, K. (2018). Killed *Bifidobacterium longum* enhanced stress tolerance and

prolonged life span of *Caenorhabditis elegans* via DAF-16. *Br J Nutr, 120*(8), 872-880.

doi:10.1017/S0007114518001563

Tepper, R. G., Ashraf, J., Kaletsky, R., Kleemann, G., Murphy, C. T., & Bussemaker, H. J. (2013). PQM-

1 complements DAF-16 as a key transcriptional regulator of DAF-2-mediated development and

longevity. *Cell, 154*(3), 676-690. doi:10.1016/j.cell.2013.07.006

Tissenbaum, H. A. (2018). DAF-16: FOXO in the Context of *C. elegans*. *Curr Top Dev Biol, 127*, 1-21.

805   doi:10.1016/bs.ctdb.2017.11.007

806   Tomioka, M., Adachi, T., Suzuki, H., Kunitomo, H., Schafer, W. R., & Iino, Y. (2006). The insulin/PI 3-

807   kinase pathway regulates salt chemotaxis learning in *Caenorhabditis elegans*. *Neuron, 51*(5),

808   613-625. doi:S0896-6273(06)00589-7 [pii]

809   10.1016/j.neuron.2006.07.024

810   Tomita, S., Saito, K., Nakamura, T., Sekiyama, Y., & Kikuchi, J. (2017). Rapid discrimination of strain-

811   dependent fermentation characteristics among *Lactobacillus* strains by NMR-based metabolomics

812   of fermented vegetable juice. *PLoS ONE, 12*(7), e0182229. doi:10.1371/journal.pone.0182229

813   Tsalik, E. L., & Hobert, O. (2003). Functional mapping of neurons that control locomotory behavior in

814   *Caenorhabditis elegans*. *J Neurobiol, 56*(2), 178-197. doi:10.1002/neu.10245

815   Vauzour, D., Camprubi-Robles, M., Miquel-Kergoat, S., Andres-Lacueva, C., Banati, D., Barberger-

816   Gateau, P., . . . Ramirez, M. (2017). Nutrition for the ageing brain: Towards evidence for an

817   optimal diet. *Ageing Res Rev, 35*, 222-240. doi:10.1016/j.arr.2016.09.010

818   Virk, B., Correia, G., Dixon, D. P., Feyst, I., Jia, J., Oberleitner, N., . . . Weinkove, D. (2012). Excessive

819   folate synthesis limits lifespan in the *C. elegans*: *E. coli* aging model. *BMC Biol, 10*, 67.

820   doi:10.1186/1741-7007-10-67

821   Vohra, M., Lemieux, G. A., Lin, L., & Ashrafi, K. (2017). The beneficial effects of dietary restriction on

822   learning are distinct from its effects on longevity and mediated by depletion of a neuroinhibitory

823   metabolite. *PLoS Biol, 15*(8), e2002032. doi:10.1371/journal.pbio.2002032

824   Vohra, M., Lemieux, G. A., Lin, L., & Ashrafi, K. (2018). Kynurenic acid accumulation underlies

825   learning and memory impairment associated with aging. *Genes Dev, 32*(1), 14-19.

826   doi:10.1101/gad.307918.117

827   Wang, S., Ahmadi, S., Nagpal, R., Jain, S., Mishra, S. P., Kavanagh, K., . . . Yadav, H. (2020).

828   Lipoteichoic acid from the cell wall of a heat killed *Lactobacillus paracasei* D3-5 ameliorates

aging-related leaky gut, inflammation and improves physical and cognitive functions: from *C.*

elegans to mice. *Geroscience, 42*(1), 333-352. doi:10.1007/s11357-019-00137-4

Watson, E., MacNeil, L. T., Ritter, A. D., Yilmaz, L. S., Rosebrock, A. P., Caudy, A. A., & Walhout, A.

J. (2014). Interspecies systems biology uncovers metabolites affecting *C. elegans* gene expression

and life history traits. *Cell, 156*(4), 759-770. doi:10.1016/j.cell.2014.01.047

White, J. G., Southgate, E., Thomson, J. N., & Brenner, S. (1986). The structure of the nervous-system of

the nematode *Caenorhabditis elegans*. *Philosophical Transactions of the Royal Society of London

Series B-Biological Sciences, 314*(1165), 1-340.

Wicks, S. R., de Vries, C. J., van Luenen, H. G., & Plasterk, R. H. (2000). CHE-3, a cytosolic dynein

heavy chain, is required for sensory cilia structure and function in *Caenorhabditis elegans*. *Dev

Biol, 221*(2), 295-307. doi:10.1006/dbio.2000.9686

Wolff, S., & Dillin, A. (2006). The trifecta of aging in *Caenorhabditis elegans*. *Exp Gerontol, 41*(10),

894-903. doi:10.1016/j.exger.2006.06.054

Zhang, F., Berg, M., Dierking, K., Felix, M. A., Shapira, M., Samuel, B. S., & Schulenburg, H. (2017).

Caenorhabditis elegans as a Model for Microbiome Research. *Front Microbiol, 8*, 485.

doi:10.3389/fmicb.2017.00485

Zhao, L., Zhao, Y., Liu, R., Zheng, X., Zhang, M., Guo, H., . . . Ren, F. (2017). The Transcription Factor

DAF-16 is Essential for Increased Longevity in *C. elegans* Exposed to *Bifidobacterium longum

BB68. *Sci Rep, 7*(1), 7408. doi:10.1038/s41598-017-07974-3

Zhao, Y., Zhao, L., Zheng, X., Fu, T., Guo, H., & Ren, F. (2013). *Lactobacillus salivarius* strain FDB89

induced longevity in *Caenorhabditis elegans* by dietary restriction. *J Microbiol, 51*(2), 183-188.

doi:10.1007/s12275-013-2076-2

Zhou, J. J., Chun, L., & Liu, J. F. (2019). A Comprehensive Understanding of Dietary Effects on *C.*

elegans Physiology. *Curr Med Sci, 39*(5), 679-684. doi:10.1007/s11596-019-2091-6

Zhou, M., Yu, H., Yin, X., Sabour, P. M., Chen, W., & Gong, J. (2014). *Lactobacillus zeae* protects

Caenorhabditis elegans from enterotoxigenic *Escherichia coli*-caused death by inhibiting

enterotoxin gene expression of the pathogen. *PLoS ONE, 9*(2), e89004.

doi:10.1371/journal.pone.0089004

Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., . . . Chanda, S. K.

(2019). Metascape provides a biologist-oriented resource for the analysis of systems-level

datasets. *Nat Commun, 10*(1), 1523. doi:10.1038/s41467-019-09234-6

 Figure legends

Figure 1 Thermotaxis performance declines with age

(A) Schematic of thermotaxis assay. Animals were placed at light blue circles on a thermal gradient

without food. The pink rectangle indicates the sections around the T_{cult}. After one hour, the number of

animals in each section was counted to calculate the thermotaxis performance index using the indicated

formula. (B and C) Age-dependent changes in thermotaxis behavior. D1 and D5 animals were cultivated

with *E. coli* at 20 °C and placed at the center of a 14-20 or 20-26 °C gradient. (B) Distributions of animals

(pink rectangle: the sections around the T_{cult}) on the thermotaxis plates. (C) Box plots of thermotaxis

performance indices. The number of experiments is shown. Statistics: Student's t-test compared to D1

adults. **$p<0.01$, ***$p<0.001$. (D) Box plots summarizing thermotaxis performance indices of animals at

different ages. Animals were cultivated with *E. coli* at 23 °C and placed at the center of a 17-23 °C

gradient. The number of experiments is shown. Statistics: The mean indices marked with distinct

alphabets are significantly different ($p < 0.05$) according to One-way ANOVA followed by Tukey–

Kramer test.

Figure 2 LAB screen for thermotaxis in aged animals

(A) Schematic of the screening procedure. Animals were cultivated at 23 °C with *E. coli* until D1 and

transferred to *E. coli* or LAB plates every day until D5. At D5, animals were subjected to thermotaxis

assays with a thermal gradient of 17-23 °C. (B) Box plots comparing thermotaxis performance indices of

D5 animals fed LAB to those of D1 (pink dashed line) and D5 animals (light blue dashed line) fed *E. coli*.

"Not applicable" (NA) indicates that animals fed those LAB were not subjected to the assay because they

were sick or dead. Abbreviations: *B, Bifidobacterium; Lb, Lactobacillus; Lc: Lactococcus; Ls,*

Leuconostoc; P, Pediococcus; S, Streptococcus. The number of experiments is shown. Statistics: One-

way ANOVA followed by Dunnett's multiple comparison test compared to D5 adults fed *E. coli*,

***$p<0.001$; **$p<0.01$; *$p<0.05$.

Figure 3 LAB-fed aged animals learn a new T$_{cult}$

(A and B) The distribution of animals on thermotaxis plates. Pink rectangles indicate the sections around

the T$_{cult}$. (A) D1 or D5 animals fed indicated bacteria were cultivated at 23 °C and placed at the center of

a 17-23 °C gradient. (B) Temperature shift assay. T$_{cult}$ was shifted from 23 °C to 17 °C one day before the

assay. Animals were placed at the center of the 17-23 °C gradient. (C) Box plots summarizing

thermotaxis indices corresponding to (A) and (B). Thermotaxis indices were calculated to examine the

mean distribution of animals on thermotaxis plates using the indicated formula. The number of

experiments is shown. Statistics: Student's t-test for comparison between T$_{cult}$=23 °C and T$_{cult}$=23 °C →

17 °C, ***p<0.001.

Figure 4 LAB shows various effects on lifespan and locomotion

Animals were fed *E. coli* until D1 and indicated bacteria after D1. (A) Survival curves of animals fed

indicated LAB are shown with control animals fed *E. coli*. NGM plates without peptone were used to

avoid the undesired growth of *E. coli* on LAB plates. N=4 experiments with 25 animals/experiment (100

animals in total). Statistics: Log-rank test. p values are shown. (B and C) The number of thrashes in liquid

(B) and distance of migration in three minutes on plates with *E. coli* (C) were measured to examine the

locomotion of aged animals. The number of animals is shown in bars. Error bars: S.E.M. Statistics: One-

way ANOVA followed by Dunnett's multiple comparison test compared to D5 fed *E. coli*, p***<0.001;

ns, p>0.05.

Figure 5 Bacteria affect thermotaxis of aged animals as nutrition

Box plots show thermotaxis performance indices of animals fed indicated bacteria and cultivated at

23 °C. Aged animals were transferred every day to new plates from D1. The number of experiments is

shown. Statistics: The mean indices marked with distinct alphabets are significantly different ($p < 0.05$)

according to One-way ANOVA followed by Tukey–Kramer test. (A) The short-term effects of diet. The

diet was switched one day before the thermotaxis assay, as indicated in the schematic. (B) The effect of

heat treatment and crushing of bacteria. The bacteria were killed by incubating at 65 °C for 1 hour or

100 °C for 10 min. After heat treatment, bacteria were crushed using a bead-based homogenizer. (C) The

mixture of bacteria. Live *E. coli* and *Lb. reuteri* were mixed at a 1:1 or 1:2 ratio with the final

concentration of 0.1 g/ml and used as a diet. (D) The effect of bacterial odor. Animals were exposed to

the bacterial odor by putting the bacterial solution on the lid and cultivated, as shown in the schematic.

Ec: *E. coli*, Lr: *Lb. reuteri*.

Figure 6 *Lactobacilli* in a clade are associated with high thermotaxis performance of aged animals

Phylogenetic tree of LAB based on 16S rRNA is shown with fermentation mode, and heatmap of

performance indices of aged animals fed indicated LAB from D1. Bootstrap values are indicated at each

node on the phylogenetic tree. Fermentation modes were categorized based on previous studies (see Table

S1). The same data as Figure 2B were used for thermotaxis performance indices. NA in the performance

indices heatmap indicates that animals fed those LAB were not subjected to the thermotaxis assay

because they were sick or dead. Fermentation mode indicates obligatory hetero- (green), facultatively

hetero- (light green), and obligatory homofermentative (orange) LAB.

Figure 7 *daf-16* is involved in the effect of *Lb. reuteri* on thermotaxis in aged animals

(A) Box plots summarizing thermotaxis performance indices of animals with indicated genotypes.

Animals were cultivated at 23 °C with *E. coli* or *Lb. reuteri* from D1. The number of experiments is

shown. Statistics: One-way ANOVA followed by Dunnett's multiple comparison test compared to the D1

control for each condition, $p^{***}<0.001$; ns, $p>0.05$. (B) Distribution of animals of indicated conditions on

thermotaxis plates. Pink rectangles indicate the sections around the T_{cult}. (C) Survival curves of animals

with indicated genotypes fed *E. coli* or *Lb. reuteri* from D1 and cultivated at 23 °C. NGM plates without

peptone were used. N=4 experiments with 25 animals/experiment (100 animals in total). Statistics: Log-

rank test, p***<0.001; ns, p>0.05. (D) Box plots summarizing thermotaxis performance indices of

animals with indicated genotypes. Animals were cultivated at 15 °C for 96 hours to avoid dauer formation

of *daf-2(e1370)* and then incubated at 23 °C until D1 or D5 with *E. coli* or *Lb. reuteri*. The number of

experiments is shown. Statistics: The mean indices marked with distinct alphabets are significantly

different (p < 0.05) according to One-way ANOVA followed by Tukey–Kramer test.

Figure 8 *daf-16* b isoform functions in neurons to maintain high thermotaxis ability in aged animals

fed *Lb. reuteri*

(A) Schematic of *daf-16* locus with representative isoforms based on WormBase (https://wormbase.org).

Black boxes and black lines indicate exons and introns, respectively. Arrows indicate 3'UTR. Alleles

used in this study are shown in magenta. The promoter of the *daf-16* b isoform (4.9 kbp) is indicated in

light blue. (B and C) Box plots summarizing thermotaxis performance indices of animals with indicated

genotypes. Animals were cultivated at 23 °C with *E. coli* or *Lb. reuteri* from D1. The number of

experiments is shown. Statistics: The mean indices marked with distinct alphabets are significantly

different (p < 0.05) according to One-way ANOVA followed by Tukey–Kramer test.   (B) Analysis of

different alleles of *daf-16* indicated in (A). (C) Tissue-specific rescue. The single copy-insertions of *daf-*

*16*b fragment with introns under tissue-specific promoters were used to examine if it rescues *daf-*

16(mu86): *myo-2*p, pharynx; *myo-3*p, body-wall muscle; *rgef-1*p, pan-neuron.

Figure 9 Transcriptome analysis on the effect of aging and diet

(A) Schematic of transcriptome analysis. We carried out RNA sequencing of indicated samples (eight in

total). Principal component analysis was carried out using all samples. Differentially expressed genes

were analyzed using D5 samples to examine the difference between Group A and Group B. (B) A scatter

plots of the two variables projected on the first and second principal components. The percentage of

variance explained by each principal component is indicated in brackets. (C) Differentially expressed

genes (DEG) between Group A and Group B in (A). DEG were ranked by the ratio between Group A's

and B's average expression, and the top 10 genes are shown. Magenta and light blue indicate genes that

are upregulated and downregulated by DAF-16 (Tepper et al., 2013), respectively. (D) Gene ontology

analysis of differentially expressed genes between Group A and B. x-axis indicates $\log_{10}$[p-value of

enrichment]. The number of genes is indicated in brackets. (E) Venn diagram showing the overlap

between differentially expressed genes of our samples and DAF-16 targets (Tepper et al., 2013). The

number of genes is indicated in brackets.

Supplementary information

Supplementary Materials and Methods

AFD and AIY imaging

To examine if AFD and AIY experience cell death at D5, NUJ296 *knjIs15[gcy-8Mp::GCaMP6m + gcy-*

8Mp::tagRFP + ges-1p::tagRFP] and IK1144 *njIs26[AIYp::GCaMP3 + AIYp::tagRFP + ges-*

1p::tagRFP] were imaged, respectively. D1 and D5 animals were immobilized using 1 mM levamisole in

M9, mounted on an agarose pad, and imaged using an Axio Imager.A2 equipped with a Plan-Apochromat

63x/1.4 oil objective (Zeiss). TagRFP signals in the cell body of AFD or AIY were visualized by green

LED (555/30 nm) of Colibri 7 light source, and a quad-band path filter ser 90 HE LED (Zeiss) and used

to evaluate possible cell death. We did not observe any loss of cell bodies in young or aged animals.

Food recognition assay

The food recognition assay was performed as previously described with a few modifications (Sawin et al.,

2000). Assay plates with food were prepared by spreading OP50 onto NGM plates. For well-fed animals,

animals were washed twice in S basal buffer (Brenner, 1974) and transferred using a capillary glass

pipette into a drop of the buffer on an assay plate with or without food. Five minutes after transfer, the

number of body bends in 20 s intervals was counted. For starved animals, 5–15 animals were washed

twice in S basal buffer and incubated on NGM plates without food for 30 min. After transferring them on

assay plates with or without food, we measured the number of body bends.

Salt avoidance assay

For a gradient assay of salt chemotaxis (Saeki et al., 2001), a salt gradient was formed overnight by

placing an agar plug containing 100 mM of NaCl (5 mm diameter) 2 cm away from the edge of the 90

mm assay plate (2% Bacto Agar, 5 mM K-PO$_4$ (pH 6.0), 1 mM CaCl$_2$, 1 mM MgSO$_4$). D1, *E. coli*-fed D5,

and *Lb. reuteri*-fed D5 animals were divided into three groups. The first group of animals received no

conditioning (Naïve). For conditioning, animals were washed three times with chemotaxis buffer (5 mM

KPO$_4$ (pH 6.0), 1 mM CaCl$_2$, 1 mM MgSO$_4$), transferred to the same buffer with 20 mM NaCl (NaCl-

conditioned) or without NaCl (Mock-conditioned), and incubated at 25°C for 1 hr. These animals were

placed at the center of the assay plates and then incubated at 25°C for 30 min. The chemotaxis index was

calculated as $(N_{NaCl} - N_{control})/(N_{total} - N_{origin})$ as indicated in Figure S4B. One hundred to two hundred

animals were used in each assay.

For a quadrant assay of salt taxis (Wicks et al., 2000), we used a compartmentalized plate (Falcon

X, Becton Dickinson Labware) as an assay plate (2% Bacto Agar, 5 mM K-PO$_4$ (pH 6.0), 1 mM CaCl$_2$, 1

mM MgSO$_4$). The plates were freshly prepared on the day of the assay with two different agar solutions, 0

mM NaCl and 25 mM NaCl. D1, *E. coli*-fed D5, and *Lb. reuteri*-fed D5 animals were divided into three

groups. The first group of animals received no conditioning (Naïve). For conditioning, animals were

washed three times with chemotaxis buffer (5 mM K-PO$_4$ (pH 6.0), 1 mM CaCl$_2$, 1 mM MgSO$_4$),

transferred to the same buffer with 100 mM NaCl (NaCl-conditioned) or without NaCl (Mock-

conditioned), and incubated at room temperature for 15 min. These animals were placed at the center of

the assay plates and then incubated at room temperature for 10 min. The chemotaxis index was calculated

as $(N_{NaCl} - N_{control})/(N_{total} - N_{origin})$ as indicated in Figure S4B. One hundred to two hundred animals were

used in each assay.

1012

1013   **Gram staining**

1014   Bacteria are fixed with methanol and stained using Gram Color Kit (Muto Pure Chemicals Co., Ltd.,

1015   Tokyo, Japan). Stained bacteria are imaged using an Axio Imager.A2 equipped with a Plan-Apochromat

1016   63x/1.4 oil objective (Zeiss).

1017

 FITC-labeling of bacteria

To examine whether *C. elegans* ingests bacteria, we used fluorescently labeled bacteria. Bacterial cells

were incubated with Phosphate-Buffered Saline (PBS, Takara) (unlabeled) or 0.1 mg/ml FITC-I (Wako)

in PBS for one hour, washed with PBS three times, and resuspended with PBS at 0.1 g/ml. For the mixed

condition, an equal amount of unlabeled and FITC-labeled bacteria were mixed. Bacteria were spread on

NGM plates and dried. D1 adult animals were placed on the bacterial plate and incubated at 23 °C for 20

min. Excess amounts of fluorescent bacteria were removed by letting worms crawl on an NGM plate

without food for a few minutes. Animals were imaged after washing using an Axio Imager.A2 equipped

with a Plan-Apochromat 63x/1.4 oil objective (Zeiss).

1027

Quantitative RT-PCR

RNA was prepared as described in the RNA sequencing section. Two micrograms of total RNA were

reverse transcribed to cDNA with a mixture of random and oligo dT primers using ReverTra Ace qPCR

RT Master Mix with gDNA Remover (TOYOBO). The cDNA and gene-specific primers were used for

qPCR reaction with THUNDERBIRD SYBR qPCR Mix (TOYOBO), and the products were detected

using a LightCycler 96 System (Roche). The gene-specific primers were used for *pha-4*: KN1370, 5'-

GGTTGCCAGGTCCCCTGACA-3'; KN1371, 5'-GCCTACGGAGGTAGCATCCA-3'. *cdc-42* was used

as a reference because it is stable and unaltered during aging (Hoogewijs, Houthoofd, Matthijssens,

Vandesompele, & Vanfleteren, 2008; Mann, Van Nostrand, Friedland, Liu, & Kim, 2016) (KN1170, 5'-

CTGCTGGACAGGAAGATTACG-3'; KN1171, 5'-CTCGGACATTCTCGAATGAAG-3').

1038

 Supplementary Figure Legends

 Figure S1 Survival curve of animals cultivated at different temperature

1041 Survival curves of wild-type animals cultivated with *E. coli* from eggs at the indicated temperatures. N=4

1042 experiments with 25 animals/experiment (100 animals in total). Statistics: Log-rank test, p***<0.001.

1043

1044 **Figure S2 Distributions of animals on the temperature gradient at different age**

1045 The distributions of animals at D1 - D5 on the temperature gradient after thermotaxis assays. Animals

1046 were cultivated at 23 °C and placed at the center of the 17-23 °C gradient. Pink rectangles indicate the

1047 sections around the T_{cult}.

1048

1049 **Figure S3 HT115-fed animals showed thermotaxis decline**

1050 Box plots summarizing thermotaxis performance indices. Animals were cultivated with OP50 until D1

1051 and then with OP50 or HT115 until D5. The number of experiments is shown. Statistics: The mean

1052 indices marked with distinct alphabets are significantly different (p < 0.05) according to One-way

1053 ANOVA followed by Tukey–Kramer test.

1054

Figure S4 AFD and AIY are alive at D5

TagRFP of NUJ296 *knjIs15[gcy-8Mp::GCaMP6m + gcy-8Mp::tagRFP + ges-1p::tagRFP]* and IK1144

njIs26[AIYp::GCaMP3 + AIYp::tagRFP + ges-1p::tagRFP] were imaged to visualize AFD and AIY,

respectively. (A and C) Representative images of the head of C. elegans for tagRFP (top panels) or

tagRFP + Differential interference contrast (DIC) (bottom panels). Scale bar: 50 µm. Arrows indicate the

cell bodies of the neurons. Arrowheads in (A) indicate tagRFP expression in the intestine used as a co-

injection marker. (B and D) Bar graphs show the percentage of animals with AFD (B) and AIY (D) in D1

and D5, based on tagRFP signals in the cell body. The number indicates the number of animals with

visible neurons over the total number of animals examined.

1064

Figure S5 *E. coli*-fed aged animals sense food

(A and B) Food recognition assays of D1 in (A) and D5 adults in (B). Animals were pre-conditioned with

(well-fed) or without (food-deprived) *E. coli* and assayed on plates with or without *E. coli*. Animals'

locomotion was evaluated by body bends in 20 sec. The presence of food on the assay plate slows down

the locomotion of well-fed animals (basal slowing response). Pre-conditioning animals without food

enhanced the basal slowing response (enhanced slowing response). The number of animals examined is

shown. Error bars: S.E.M. Statistics: Two-way ANOVA with Turkey's multiple comparison test,

***p<0.001; **p<0.01; ns, p>0.05.

1073

Figure S6 Aged *E. coli*-fed animals perform salt avoidance behavior irrespective of diet

(A) Schematic of conditioning of animals. Animals were conditioned in the absence of food with (NaCl)

or without (Mock) NaCl. (B) Schematic of salt avoidance assays. Gradient and quadrant assays were used

to evaluate salt avoidance behavior. The chemotaxis index was calculated using the indicated formula

based on the number of animals in each area after the assay. (C and D) Box plots summarizing

chemotaxis indices of salt avoidance assays. The number of experiments is shown. One-way ANOVA

followed by Dunnett's multiple comparison test compared to Naïve condition in each condition,

***p<0.001; ns, p>0.05.

1082

1083

Figure S7 Animals fed heterofermentative LAB are not thermophilic

Animals were cultivated at 23 °C and placed at the center of a 20-26 °C gradient to examine the

thermophilicity. D5 animals with indicated genotypes were cultivated with *E. coli* or *Lb. reuteri* from D1.

tax-6(db60) was used as a thermophilic mutant. (A) Schematic of thermotaxis assay and formula to

calculate thermotaxis index. (B) Distribution of animals on the temperature gradient after thermotaxis

assays. Pink rectangles indicate the sections around the T_{cult}. (C) Box plots summarizing thermotaxis

indices of animals under indicated conditions. The number of experiments is shown. Statistics: The mean

indices marked with distinct alphabets are significantly different ($p < 0.05$) according to One-way

ANOVA followed by Tukey–Kramer test.

1093

1094   **Figure S8 Diets affect thermotaxis of aged animals on peptone-free plates**

1095   Box plots summarizing performance indices of animals cultivated on peptone-free NGM plates. D5

1096   animals were cultivated at 23 °C with *E. coli* or *Lb. reuteri* from D1. The number of experiments is

1097   shown. Statistics: One-way ANOVA followed by Dunnett's multiple comparison test compared to the D1

1098   control, p***<0.001; ns, p>0.05.

1099

1100   **Figure S9 Animals ingest *E. coli* and *Lb. reuteri***

1101   Representative images show fluorescently labeled bacteria in the intestine of *C. elegans*. Animals were

1102   cultivated with FICT-labeled *E. coli*, *Lb. reuteri*, or the mixture of fluorescently labeled and unlabeled

1103   bacteria. Both fluorescence and transmission light was simultaneously imaged to show that *C. elegans*

1104   ingested bacteria.

1105

1106   **Figure S10 Images of Gram-stained bacteria**

1107   Images of Gram-stained *E. coli* and select LAB. *E. coli* and LAB are Gram-negative and positive,

1108   respectively. Scale bar: 10 µm.

1109

Figure S11 LAB-fed bacteria are not dietary-restricted

(A) Expression of *pha-4* transcripts in aged animals fed indicated LAB relative to aged animals fed *E.*

coli. (B) Box plots summarizing thermotaxis indices of wild type and *eat-2* mutant animals cultivated at

23 °C with *E. coli*. The number of experiments is shown. Statistics: Student's t-test, ns, p>0.05.

Figure S12 Principal component analysis of the transcriptome data of the animals of different ages

and diets

Principal component analysis was carried out for the transcriptome data of D1, and D5 animals fed *E. coli*

or LAB (*Lb. gasseri*; *Lb. delbrueckii*, *P. pentosaceus*, *Lb. reuteri*, *Lb. rhamnosus*, and *Lb. plantarum*).

Loadings of genes were ranked. (A and B) Loadings of each gene to PC1 (A) and PC2. Magenta indicates

the top 5% of genes most correlated and anti-correlated to PC1 or PC2. the x-axis indicates genes, but the

orders are different between (A) and (B). (C and D) Gene ontology analysis of the top 5% of genes

correlated or anti-correlated with PC1 (C) or PC2 (D).

1122

1123 **Table S1 List of LAB strains**

1124

A   T$_{cult}$=20 ℃, 17 ℃ center

$$Performance\ index = \frac{n_j + n_{j+1}}{N}$$

N: total number of worms
n$_i$: number of worms in section i (i=1~8)
j, j+1: two sections near T$_{cult}$

B   T$_{cult}$=20 ℃

C   T$_{cult}$=20 ℃

D   T$_{cult}$=23 ℃, 20 ℃ center

A   T$_{cult}$=23 ℃, 20 ℃ center

B   T$_{cult}$=23 ℃→17 ℃, 20 ℃ center

C

$$Thermotaxis\ index = \sum_{i=1}^{8} i\,\frac{n_i}{N}$$

N: total number of worms
n_i: number of worms in section i (i=1~8)

A T_{cult}=23 ℃, 20 ℃ center

B T_{cult}=23 ℃, 20 ℃ center

C T_{cult}=23 ℃, 20 ℃ center

D T_{cult}=23 ℃, 20 ℃ center

A

B T_{cult}=23 ℃, 20 ℃ center

C T_{cult}=23 ℃, 20 ℃ center

■ **D1** ■ **D5 (*E.coli*)** ■ **D5 (*Lb. reuteri*)**

Performance index

A   **RNA sequencing**

B

C

Homo *Lactobacillus* , *E. coli* > Hetero *Lactobacillus*

Gene	Description	Average FPKM		Ratio	p value	DAF-16 target
		Hetero LAB	Homo LAB, *E. coli*	(Hetero/Homo)		
fipr-24	FIP (Fungus-Induced Protein) Related	0.93	7.92	0.1	0.006	
scl-12	SCP-Like extracellular protein	0.17	1.04	0.2	0.013	
fipr-23	FIP (Fungus-Induced Protein) Related	0.80	4.43	0.2	0.003	
fipr-22	FIP (Fungus-Induced Protein) Related	2.21	12.04	0.2	0.009	
F56D6.8	hypothetical protein	20.51	109.86	0.2	0.007	
C37A5.3	hypothetical protein	1.26	6.50	0.2	0.000	
drd-10	dietary restriction down regulated	0.82	4.20	0.2	0.038	
dct-8	DAF-16/FOXO Controlled, germline Tumor affecting	18.84	94.12	0.2	0.005	
T01C3.11	hypothetical protein	0.94	4.24	0.2	0.004	
F56D6.9	hypothetical protein	15.85	67.32	0.2	0.015	

Hetero *Lactobacillus* > Homo *Lactobacillus* , *E. coli*

Gene	Description	Average FPKM		Ratio	p value	DAF-16 target
		Hetero LAB	Homo LAB, *E. coli*	(Hetero/Homo)		
clec-125	C-type LECtin	0.46	0.07	6.1	0.024	
lys-6	LYSozyme	10.42	1.85	5.6	0.020	
arrd-6	ARRestin Domain protein	0.28	0.06	4.8	0.001	
lys-5	LYSozyme	12.58	2.68	4.7	0.028	
comt-2	Catechol-O-MethylTransferase family	3.15	0.74	4.2	0.027	
pho-13	intestinal acid PHOsphatase	4.10	1.05	3.9	0.000	
C06E1.7	Putative glycosyltransferase	0.42	0.11	3.9	0.048	
C54G6.3	hypothetical protein	0.23	0.06	3.7	0.046	
npax-2	N-terminal PAX (PAI domain only) protein	0.31	0.09	3.5	0.034	
F59A1.13	hypothetical protein	0.12	0.04	3.2	0.001	

D

Homo *Lb, E. coli* > Hetero *Lb* (65)

Hetero *Lb* > Homo *Lb, E. coli* (71)

E

T$_{cult}$=23 ℃, 20 ℃ center

T_{cult}=23 ℃, 20 ℃ center,

A

$$Thermotaxis\ index = \sum_{i=1}^{8} i\,\frac{n_i}{N}$$

N: total number of worms
n_i: number of worms in section i (i=1~8)

C T$_{cult}$=23 ℃, 23 ℃ center

B T$_{cult}$=23 ℃, 23 ℃ center

T$_{cult}$=23 ℃, 20 ℃ center

Peptone-free

E. coli NC

Lb. reuteri NC

E. coli FITC

Lb. reuteri FITC

E. coli FITC **+** *Lb. reuteri* NC

E. coli NC **+** *Lb. reuteri* FITC

A

B T_{cult}=23 ℃, 20 ℃ center

A

B

C

Positively correlated to PC1

Negatively correlated to PC1

D

Positively correlated to PC2

Negatively correlated to PC2