Alteration of gut microbiota with a broad-spectrum antibiotic does not impair

maternal care in the European earwig.

Sophie Van Meyel*, Séverine Devers, Simon Dupont, Franck Dedeine and Joël Meunier

Institut de Recherche sur la Biologie de l'Insecte, UMR 7261 CNRS – Université de Tours, Tours,

France

*Corresponding author: S. Van Meyel, sophie.vanmeyel@univ-tours.fr

ABSTRACT

The microbes residing within the gut of an animal host often increase their own fitness by

modifying their host's physiological, reproductive, and behavioural functions. Whereas recent

studies suggest that they may also shape host sociality and therefore have critical effects on

animal social evolution, the impact of the gut microbiota on maternal care remains unexplored.

This is surprising, as this behaviour is widespread among animals, often determines the fitness

of both juveniles and parents, and is essential in the evolution of complex animal societies.

Here, we tested whether life-long alterations of the gut microbiota with rifampicin - a broad-

spectrum antibiotic - impair pre- and post-hatching maternal care in the European earwig. Our

results first confirm that rifampicin altered the mothers' gut microbial communities and

indicate that the composition of the gut microbiota differs before and after egg care. Contrary

to our predictions, however, the rifampicin-induced alterations of the gut microbiota did not

modify pre- or post-hatching care. Independent of maternal care, rifampicin increased the

females' feces production and resulted in lighter eggs and juveniles. By contrast, rifampicin

altered none of the other 21 physiological, reproductive and longevity traits measured over the

300 days of a female's lifetime. Overall, these findings reveal that altering the gut microbiota

with a large spectrum antibiotic such as rifampicin does not necessarily affect host sociality.

They also emphasize that not all animals have evolved a co-dependence with their microbiota

and call for caution when generalizing the central role of gut microbes in a host biology.

Keywords: Antibiotic, Dermaptera, Insect, Microbiome, Parental care

1-INTRODUCTION

Almost all animals harbour a gut microbiota, i.e. a community of microorganisms residing within the gut of the host [1]. Some of these gut microbes have long been known for their pathogenic effects in the hosts [2] and others for the benefits they provide to the hosts in terms of nutritional mutualism [3]. Over the last decades, however, a growing number of studies has been revealing that the effects of gut microbes are much more diverse than previously thought and shape numerous physiological, reproductive, and behavioural functions of the host [4]. In the fruit fly *Drosophila melanogaster*, for instance, the gut microbiota is associated with hormone signalling, metabolism and ageing [5]. Gut microbes can also shape the hosts' immunocompetence and resistance against pesticides and toxic plant metabolites [6], such as in the mosquito *Anopheles stephensi* [7], the bean bug *Riptortus pedestri* [8] and the wasp *Nasonia vitripennis* [9]. Similarly, the gut microbiota is a key parameter in host reproduction and mating incompatibilities, as found in the fruit fly *Bactrocera minax* [10], the terrestrial isopod *Armadillidium vulgare* [11], and the parasitic wasp *Asobara tabida* [12]. Finally, gut microbes shape the expression of numerous host behaviours, such as offspring activity in the stinkbug *Megacopta punctissima* [13], and different tasks in honeybees [14].

Recent studies also suggest that gut microbiota can play a critical role in the sociality of their hosts by shaping the expression and nature of social interactions and/or by transforming mediators of social aggregation. For instance, family-living rats with a diet-altered gut microbiota exhibit deficient sociability and increased social avoidance [15,16]. Antibiotic-induced modifications of gut microbiota also alter the chemical signatures of social hosts and lead to higher levels of aggressiveness toward conspecifics in the leaf-cutting ant *Acromyrmex*

echinator [17] and the honeybee *Apis mellifera* [18]. Finally, alteration of the gut microbiota

reduces the production of aggregation pheromones in the swarm-living desert locust

Schistocerca gregaria [19] and diminishes the presence of aggregation pheromones in feces of

the gregarious German cockroach *Blattella germanica* [20].

Despite these causal and correlative links between the hosts' gut microbial communities

and sociality, the role of gut microbes on the expression of parental care remains

experimentally unexplored. This is surprising, because this form of social behaviour is present in

a large and taxonomically diverse number of animal species [21], has considerable effects on

the fitness of both juveniles and parents [22] and because shedding light on this link may

provide crucial information on the role of gut microbes in the early evolution of complex animal

societies [23]. On one hand, gut microbes could indirectly alter the investment in parental care,

because parents are expected to adjust their level of care to their own condition [24] and

altered gut microbial communities can lower these conditions in multiple ways (see above). On

the other hand, gut microbes could serve as a direct promoter of parental care because, by

enforcing the expression of care in adult hosts, parental gut microbes could maximize their

chances to reach novel hosts [25–30](but see [31]). The transfer of gut microbes through

parental care has been reported in several insect species, such as the stinkbug *Parastrachia*

japonensis [32], the Japanese common plataspid stinkbug *Megacopta punctatissima* [33], the

wood cockroach *Cryptocercus punctulatus* [34] and the wood-feeding termite *Reticulitermes*

grassei [35]. However, we still know very little about whether and how the gut microbiota

influences the investment in parental care.

In this study, we address this gap in knowledge by investigating whether gut microbiota

alteration with rifampicin - a broad-spectrum antibiotic - impaired the expression of pre- and

post-hatching maternal care in the European earwig *Forficula auricularia*. In this omnivorous

insect, mothers tend their first clutch of eggs over winter and their second and terminal clutch

(when present) during spring. During these periods, mothers stop their foraging activity and

provide egg care in the forms of protection against desiccation, predators and pathogens [36].

Egg care also mediates the transfer of microbes with antifungal properties to eggshell in the

maritime earwig *Anisolabis maritima* [37], a process that possibly occurs in the European

earwig [38]. Upon egg hatching, *F. auricularia* mothers continue tending their brood of newly

emerged juveniles (called nymphs) for two weeks, during which they provide care in the forms

of fierce protections against predators, grooming behaviours, and food provisioning through

regurgitation [39]. Pre-hatching care is necessary to ensure egg development and hatching [38],

whereas post-hatching care is facultative for the development and survival of nymphs [40].

Earwig females present important inter-individual variation in the expression of maternal care

within populations [41,42], and this variation is partly inherited from the parents [43] and partly

depends on environmental inputs, such as the social environment or food resources [44–46].

We altered the gut microbiota of *F. auricularia* females by feeding them with rifampicin

during their entire adult lifetime (about 14 months) and measured whether and how it affected

gut microbial communities, maternal care, and other life-history traits. Specifically, we first

determined how the antibiotherapy alters the diversity and structure of the gut bacterial

community of females at two periods of their life-cycle (just before the production and at the

hatching of their 1st clutch eggs) by sequencing 16S rRNA gene (V3-V4 region) amplicons. We

then tested the effects of rifampicin on the expression of four pre- and two post-hatching forms

of maternal care toward 1st clutch eggs and nymphs, respectively. Finally, to disentangle

whether the potential link between gut microbiota alteration and the level of maternal care is

direct and/or indirect, we investigated the effects of rifampicin on 24 other traits measured

throughout the mothers' lifetime and reflecting their general physiological state, investment in

future reproduction and longevity.

2-MATERIALS AND METHODS

2.1 Insect rearing and rifampicin treatment

The experiment involved a total of 296 *Forficula auricularia* L. (clade B [47]) males and females.

These were the first laboratory-born descendants of 74 females field-sampled in Pont-de-Ruan,

France, in 2017 and then maintained under standard laboratory conditions [42]. The entire

experiment consisted of feeding these 296 earwigs with either rifampicin or water from adult

emergence to death, and measuring the effects on mothers' behaviour, physiology,

reproduction and longevity (Figure S1). To obtain rifampicin- and control-treated mothers, we

first isolated two virgin males and two virgin females per family (n = 74 families) four days after

adult emergence, and then fed them for two weeks with green-coloured pollen pellets mixed

with either 10 µL of rifampicin (Sigma-Aldrich, PHR1806; 0.2 mg/ml) or 10 µL of water. During

these two weeks, each individual was isolated in a Petri Dish grounded with a humid filter paper

that was changed twice a week (together with the treated food) to limit the risk of self-

transplantation of gut microbes through the consumption of feces deposited on paper. The

production of green-coloured feces was always observed, confirming the consumption of

rifampicin-treated food. At the end of these two weeks, we used the 296 earwigs to set up 148

mating pairs composed of 1 virgin female and 1 virgin male from the same family and the same

treatment. There are only limited signs of inbreeding depression in this species [48] and sib-

mating allowed us reducing the risk of poor reproductive success due to possible inter-familial

cytoplasmic incompatibility between certain bacterial strains, as reported with *Wolbachia* and

Cardinium in several arthropod species [49,50]. Each of the resulting 148 mating pair received a

standard food source mostly composed of agar, carrots, pollen, and cat and bird dry food [42]

twice a week during two months. This food was mixed with either 10 µL of rifampicin (0.2

mg/ml) or 10 µL of water to follow-up on the previous treatments. After these two months,

females were isolated and maintained under winter conditions to mimic natural dispersal, allow

oviposition and subsequently measure four forms of egg care (details below) [42]. Mothers

were not provided with food and thus not treated with rifampicin from oviposition to egg

hatching, as mothers typically stop foraging during this period [40]. One day after egg hatching,

each family was maintained under spring conditions and fed with the standard food source

mixed with either 10 µL of rifampicin (0.2 mg/ml) or 10 µL of water according to the pre-

oviposition treatment. The food and treatment were renewed twice a week. We measured

three forms of maternal care towards juveniles during the following 14 days (details below),

which corresponds to the duration of family life in this species [42]. Nymphs were then

discarded from the experiment to allow newly isolated mothers to produce a 2nd clutch. These

mothers were then maintained under spring conditions and continued to receive the same

treatment (rifampicin or water) until they die. Except when stated otherwise, individuals were

always maintained in Petri dishes (diameter 9cm) lined with non-sterile moistened sand.

Rifampicin is considered one of the most potent and broad-spectrum antibiotics due to

its high-affinity binding to the RNAP β subunit, which causes the inhibition of the bacterial DNA-

dependent RNA polymerase RNAP by directly blocking the RNA elongation path [51]. It is also

commonly used to experimentally alter gut microbial communities in insects (e.g. [52–54]). The

high dose of rifampicin used in this study (about 10 times higher than the dose generally used

in smaller insect species [53,54]) was chosen to ensure gut microbial alteration and because it

did not trigger an excess of mortality in the German cockroach [52], an insect that is about the

size of the European earwig.

2.2 Effects of rifampicin on the gut microbiota

To test whether and how rifampicin treatment altered the earwigs' gut microbial communities,

we extracted the gut of 10 females per treatment (n total = 20) on the day we observed the

first oviposition of their 1[st] clutch (i.e. about 2 months after being fed with or without

rifampicin), and 10 rifampicin- and 8 water-treated females one day after their 1[st] clutch eggs

have hatched (i.e. about 1 month later; Figure S1). For gut extraction, we first anaesthetized

each female for 2 min at -20°C and then dissected them in a watch glass with sterilized double

Q water. All dissections and manipulations were conducted on a sterilized bench, under a

Bunsen burner's sterility area and using sterile material. Whole individual guts were extracted,

placed in 100 µl of T1 extraction buffer (Nucleo-Spin Tissue, Macherey-Nagel), and stored at

−80°C until DNA extraction. Two PCR amplifications were performed for each sample in a final

volume of 35 µl to amplify a 450-bp portion of the V3–V4 region of the 16S rRNA gene (forward

primer: 5ꞌ-CTT TCC CTA CAC GAC GCT CTT CCG ATC **TAC GGR AGG CAG CAG**-3ꞌ; reverse primer:

5⊠-GGA GTT CAG ACG TGT GCT CTT CCG ATC **TTA CCA GGG TAT CTA ATC**-3⊠; the Illumina

adapters and primers *per se* are indicated in non-bold and bold, respectively). 50 µl of PCR

product were then sent to the GeT-PlaGe genomic platform (GenoToul, Toulouse, France),

which performed library preparation and 2 × 250 paired-end Illumina Miseq sequencing

according to the manufacturer's instructions. DNA extractions protocols, sequencing process

and bioinformatic pipelines are detailed in the supplementary material.

2.3 Measurements of pre- and post-hatching maternal care

We first measured the effects of rifampicin on four classical forms of earwig maternal egg care:

egg grooming, egg defence, delay of maternal return and exploration rate while searching for

its eggs [55,56]. Egg grooming, which is used by earwig females to deposit chemical substances

on the eggs and to clean eggshell from dirt and fungi [38], was measured 15 days after egg

production. We first isolated each mother for 30 min, then returned them to their Petri dish

and gently deposited them at a distance of 5 cm from their clutch, and finally video-recorded

their behaviours for the subsequent 15 minutes (SONY© Handycam HDR-CX700 camera). The

resulting movies were analysed using the software BORIS v4.0.3 [57] and the total duration of

egg grooming was defined as the total amount of time each female spent on cleaning eggs with

their mandibles [38]. Clutch defence, which reflects the females' willingness to protect their

eggs from a predator attack [44], was measured 16 days after egg production. We standardly

poked each female on the pronotum with a glass capillary (one poke per second) and then

recorded the number of pokes required until the female moved more than one body length

away from the clutch. The delay of maternal return after clutch abandonment [56], which

represents the delay after which females return to their clutch after being chased away by a

simulated predator attack [44], was measured by recording the time the female took to return

to its clutch just after the end of the clutch defence measurement. Finally, the exploration rate,

which represents the level of exploration of a novel area by a mother looking for her eggs, was

measured 21 days after egg production. We removed each mother from its clutch of eggs,

subsequently deposited her at the centre of a square arena (W: 9 cm; L: 9 cm; H: 0.5 cm)

covered by a glass sheet, and then video-tracked its activity during 35 min. The video was done

under infra-red light, while the individual video tracking and calculation of exploration rate

were conducted using the software ToxTrac v2.83 [58].

We then measured the effects of rifampicin on two classical forms of post-hatching

maternal care: nymphs defence and exploration rate while searching for its nymphs [44,55].

These two forms of care were measured 10 and 12 days after egg hatching, respectively,

following the above-detailed protocols for egg defence and egg searching.

All the measurements of pre- and post-hatching maternal care were conducted in the

afternoon and under a red light as earwigs are nocturnal. These measurements were conducted

blindly regarding the treatments (rifampicin versus control). The number of replicates for each

of our measurements ranged from 9 to 59 per treatment (median = 36 per treatment; details in

Tables 1 and S1). The recorded range of values of maternal care is comparable to the range of

values obtained in previous studies conducted in other populations [41,42,44] and thus likely

reflects the natural variation in maternal care exhibited by earwig females.

2.4 Measurements of the 24 other life-history traits in mothers

We used standard protocols to test the effects of rifampicin on 7 proxies of female physiology,

as well as 16 proxies of female reproduction and female longevity [42,59]. Proxies of females'

physiology were the number of feces pellets produced per 24 hours (a number positively

associated with their digestive/foraging activity [60]) and the gain in fresh weight between two

life stages. Feces production was measured two months after the beginning of the treatments.

Females were isolated in a new Petri Dish for 24 hours, after which we counted the number of

feces pellets present on the ground. The weight gained by each female was measured between

the days of adult emergence and oviposition, and between the days of oviposition and egg

hatching. Proxies of female reproduction were measured in the 1^{st} and 2^{nd} clutches (if any) by

counting the number of eggs produced, the number of days between oviposition and egg

hatching (egg development time), and by measuring the mean egg weight at oviposition, the

egg hatching rate, and the mean offspring weight at egg hatching. We also counted the number

of days between adult emergence and oviposition (days until 1^{st} clutch oviposition), between

the females' isolation after family life and 2^{nd} clutch oviposition (days until 2^{nd} clutch

production), and between adult emergence and death (female longevity). We finally assessed

whether females produced a 2^{nd} clutch (yes = 1 or no = 0) and females' reproductive allocation

between the two clutches (i.e. the females' reproductive strategy [42]) defined as the number

of 2^{nd} clutch eggs divided by the total number of eggs produced by a female. Overall, weighing

was done to the nearest 0.01 mg using a microbalance (OHAUS© Discovery DV215CD). Sample

sizes are detailed in Tables 1 and S1.

2.5 Statistical analyses

Analyses of the α and β-diversity indices. The structure, composition and diversity of the

microbial communities were based on the 161 identified bacterial Operational Taxonomic Units

(OTUs) (see results) and analysed using PHYLOSEQ R package [61] implemented in the

FROGSSTAT Phyloseq tools [62]. Diversity within the gut microbial communities (alpha-

diversity) was assessed using two richness indices, which estimate the number of OTUs in the

microbiome with correction for subsampling (Chao1; ACE), and three metrics that aim to

measure diversity by accounting for evenness or homogeneity (Shannon, Simpson, Inverse

Simpson, and Fisher) [63]. Diversity between the gut microbial communities (beta-diversity)

was assessed using 4 measures of community similarity: 1) Jaccard indice, which does not

consider phylogeny of OTUs but takes into account their presence/absence; 2) Bray Curtis

dissimilarity, which does not consider the phylogeny but considers the number of reads

assigned to an OTU (i.e. its abundance); 3) UniFrac indice, which considers phylogeny but not

abundance; and finally 4) Weighed UniFrac indice, which considers both phylogeny and

abundance. The metrics were analysed individually using either a General Linear Model for α-

diversity, or a Permutational Multivariate Analysis of Variance Using Distance Matrices

(PERMANOVA) for β-diversity. In these models, the values (or distance matrix for β-diversity) of

each index were entered as a response variable, while the treatment (rifampicin or water), the

sampling stage of the female (before 1st oviposition or at 1st clutch egg hatching) and the

interaction between them were used as fixed factors. When required, a post-hoc analysis was

conducted by splitting the data set according to the sampling stage and then conducting

PERMANOVA on each of the two resulting subsets. To correct for multiple testing in these post-

hoc analyses, the significance level was adjusted to alpha = 0.0375 using the Mean False

Discovery Rate approach [64].

Analyses of the life-history traits. Although the presented experimental design was originally

paired, i.e. 2 females per family distributed among the two treatments, the 38 life-history traits

were often measured in only one of the pairs (see Tables 1 and S1). This was mostly due to time

constraints, and because some females died during the 18-months course of this experiment.

These overall led to critical reductions in the number of replicates that could be involved in a

paired statistical approach (details in Table S2). We, therefore, analysed the effects of

rifampicin on the 30 measurements using a series of 29 exact Mann Whitney U tests and 1

Pearson's Chi-squared test (for 2^{nd} clutch production), in which we compared the values of all

the available replicates fed with rifampicin to the values of all the available replicates fed with

water. Note that the results do not qualitatively change when we use paired analyses with the

associated smallest sample sizes (results presented in Table S2). To correct for the inflated risk

of Type I errors due to multiple testing, all p-values were adjusted using the False Discovery

Rate (FDR) method [64]. To confirm the robustness of non-significant results, we also calculated

the effect size *r* of each analysis and the number of replicates that would have been required to

detect a statistically significant effect with this effect size and a statistical power of 0.8. All

these analyses were conducted using the software R v4.0.2 (http://www.r-project.org) loaded

with the packages *exactRankTests, car, rcompanion* and *pwr*.

3-RESULTS

3.1 Description of the earwig gut microbiota

A total of 1 636 435 sequenced reads of the 16S rRNA V3-V4 region were obtained from the 38

female earwig gut samples. After sequence processing, this number went down to 1 130 241,

with 21069 to 35385 sequences per sample (median = 30595.5). The sequences were

aggregated and filtered in a total of 161 unique OTUs, which were resolved down to the family

or genus level to increase the confidence in the taxonomic assignation. All detailed information

on OTUs is given in Table S3. More than 99.90% of the sequences were assigned to four

bacterial phyla: Proteobacteria (65.94%), Firmicutes (21.12%), Bacteroidota (9.89%) and

Actinobacteriota (2.96%) (Figure 1). The remaining OTUs were assigned to Bdellovibrionota

(0.04%) and Patescibacteria (0.04%). The prevalence (i.e. frequency) of these 161 OTUs among

the 38 tested females ranged from 0.211 to 1.000 (Table S3). The vast majority of OTUs were

found in at least one female in each experimental modality (Table S3), indicating that our

rifampicin treatment did not eliminate specific OTUs.

3.2 Comparative analyses of the α and β diversity of the gut microbiota

The gut microbial α-diversity (i.e. species richness) decreased between oviposition and egg

hatching when this diversity was measured using Chao1 ($F_{1,34}$ = 21.63, P < 0.0001), ACE ($F_{1,34}$ =

24.46, P < 0.0001) and Fisher ($F_{1,34}$ = 20.85, P < 0.0001; Figure 2) indices. This decrease was,

however, not significant when α-diversity was measured using Shannon ($F_{1,34}$ = 3.18, P = 0.084;

Figure 2) and Simpson ($F_{1,34}$ = 1.60, P = 0.214) indices. Similarly, the α-diversity did not decrease

in the rifampicin treatment compared to the control (Chao1: $F_{1,34}$ = 0.72, P = 0.401; ACE: $F_{1,34}$ =

0.62, P =0.435; Fisher: $F_{1,34}$ = 0.59, P = 0.447; Shannon: $F_{1,34}$ = 1.67, P = 0.205; Simpson: $F_{1,34}$ =

0.55, P = 0.465; Figure 2), and it was independent of an interaction between female sampling

stage and rifampicin treatment (all P > 0.525).

The gut microbiota β-diversity (i.e. species composition) overall changed with female

sampling stage and rifampicin treatment. This was the case with the four measured indices of

β-diversity: Bray-Curtis (Stage: $F_{1,34}$ = 5.77, P < 0.0001; Rifampicin: $F_{1,34}$ = 4.23, P < 0.0001),

Jaccard (Stage: $F_{1,34}$ = 7.76, P < 0.0001; Rifampicin: $F_{1,34}$ = 2.37, P = 0.0036), unweighted UniFrac

(Stage: $F_{1,34}$ = 6.51, P < 0.0001; Rifampicin: $F_{1,34}$ = 3.39, P = 0.0006) and weighted UniFrac (Stage:

$F_{1,34}$ = 14.10, P < 0.0001; Rifampicin: $F_{1,34}$ = 6.42, P = 0.0006). In particular, females before

oviposition harboured less Actinobacteriota and Proteobacteria compared to females at egg

hatching, while rifampicin females overall harboured less Bacteroidota and more Firmicutes

compared to untreated females (Figure 1). The interaction between female sampling stage and

rifampicin had no effect on the β-diversity measured using all (all P > 0.117) but the weighted

UniFrac indices ($F_{1,34}$ = 2.94, P = 0.026). This interaction reflected an effect of rifampicin on the

β-diversity before oviposition ($F_{1,34}$ = 0.17, P = 0.018) but not at egg hatching ($F_{1,34}$ = 0.97, P =

0.356).

3.3 Rifampicin and maternal care

We did not detect any effect of rifampicin on the six measured forms of egg and nymph care

(Table 1). In particular, rifampicin- and control-fed mothers spent the same amount of time in

egg grooming (Figure 3A), showed the same levels of both egg and juvenile defences against a

simulated predator attack (Figures 3B and 3E), exhibited the same delay to return to their eggs

after egg defence (Figure 3C) and showed comparable exploration rates when searching for

their eggs and juveniles (Figures 3D and 3F).

3.4 Rifampicin and female's physiology, reproduction, and longevity

The consumption of rifampicin altered only 3 of the 24 measured proxies of female physiology,

reproduction, and longevity. In particular, females fed with rifampicin produced on average

twice as many feces pellets per 24h (Figure 4; Table 1), had newly hatched nymphs that were

15% lighter (W = 1244, P = 0.002, adjusted-P = 0.025; Table S1) and laid 2^{nd} clutch eggs that

were 7% lighter (W = 628, P = 0.002, adjusted-P = 0.025; Table S1) compared to control females.

By contrast, we did not detect any effect of rifampicin on the 21 other traits (Tables 1 and S1).

4-DISCUSSION

Whereas gut microbial communities shape the physiology, reproduction and behaviour of a

great diversity of hosts, their importance on parental care – a key behaviour in social evolution

[21–23] - remains poorly explored [30]. In this study, we addressed this gap in knowledge by

treating females of the European earwig with rifampicin and measuring the effects on gut

microbial communities, maternal care and female physiology, reproduction, and longevity. Our

results first reveal that rifampicin altered the composition of the gut microbial community of

earwig females and show that this modification diminishes during the period of egg care.

Contrary to our predictions, however, the rifampicin-induced alterations of gut microbial

communities did not impair the expression of pre- or post-hatching care: rifampicin-treated and

control mothers showed similar levels of egg grooming, clutch defences against a predator,

maternal return and clutch searching. Independent of maternal care, our results reveal that the

consumption of rifampicin increased the females' production of feces pellets, as well as lead to

the production of lighter nymphs and lighter 2nd clutch eggs. By contrast, rifampicin affected

none of the other 21 physiological, reproductive and longevity traits measured over the

females' lifetime.

Our experiment first demonstrates that the ingestion of rifampicin by earwig females

modified the composition (β-diversity) but not the richness (α-diversity) of bacterial OTUs

present in the gut. The fact that rifampicin only shapes indices of β-diversity controlling for

phylogeny suggests that OTUs' phylogeny is the most prominent difference between the

community structures present in treated versus non-treated individuals. Because weighted

uniFrac is significant and (unweighted) uniFrac is not, our results then indicate that this

phylogenetic difference is specific to clades that diverged in the more distant past compared to

recently evolved nodes, a pattern in line with broad-spectrum antibiotics acting on conserved

bacterial traits. Overall, our findings thus confirm that our treatment successfully altered gut

microbial communities in earwigs (just like in other animal species [52–54]). It also indicates

that both α- and β-diversity change from pre-oviposition to egg hatching. This stage-specific

pattern may result from the absence of food intake for about four weeks before gut sampling in

females at egg hatching compared to before oviposition [65], and/or from the different rearing

temperatures [66] and differences in female age [67] between the two life stages.

Although gut microbial communities shape the expression of host's sociality in

numerous vertebrate and arthropod species [15–17,19,20], our findings reveal that rifampicin-

induced alterations of this community did not affect the expression of pre- and post-hatching

maternal care in earwigs. One might have expected that gut microbes directly drive the

expression of parental care, as enforcing this social behaviour may allow (at least some)

symbionts to reach new hosts (i.e. offspring) that are typically young (thus offering long-lived

habitats), display poor immune defences (thus facilitating bacterial establishment and

development [68]) and harbour only a few resident microbes (thus limiting the risk of

competition within the microbiome [28]). However, our results are at odds with this prediction.

This may first suggest that earwig parental care is primarily shaped by microbes that are non-

sensitive to rifampicin or that non-sensitive microbes can take over this function (functional

redundancy). In insects, gut microbial communities do not only encompass a broad diversity of

bacteria (among which some are resistant to rifampicin) but also fungi, protists and other

microorganisms that could have key roles and functional redundancies in hosts biology [37,60].

Even if the previous experimental studies linking gut microbiota and host sociality focused on

bacteria [15–17,19,20], future studies will be required to confirm that no other members of the

gut microbiota shape parental care in our study species, and to explore causal links between

the presence of certain members of the microbiome and the level of maternal care expressed

by its host. A second potential explanation of our results is that microbial symbionts never

developed any specific capabilities to manipulate host sociality, either because adapted strain

never occurred within the microbial populations associated with these earwig species (or

populations), or because certain antagonistic interactions (e.g. competition) among the

members of the microbial community have prevented the emergence of host social

manipulation. Any symbiont species (or strain) investing its resources to manipulate host

behaviour could indeed be outcompeted within the microbiome by other species or variants

that, instead, direct their resources into growth, survival or directly transmission [31](but see

[30]). Finally, a third potential explanation is that the symbionts' capability to manipulate host

sociality may change during host social evolution and could thus have vanished in the European

earwig. The evolutionary drivers of family life are indeed known to change over time [23] and,

while gut microbes may have (at least partly) driven the ancestral evolutionary shift from

solitary to family living for the reasons detailed above, the resulting benefits of parental care

for the hosts could have consolidated the expression of care and thus reduced the capability of

symbionts to control host social behaviours and/or reduced the sensitivity of the hosts to this

manipulation. Based on this hypothesis, alterations in gut microbial communities should not

shape the expression of parental care once this behaviour is established. This process could

have limited the maintenance of symbiont control over parental care in earwigs, where

maternal care is well established. Notwithstanding the drivers of our results, our findings

provide first experimental evidence that broad alteration of the gut microbiota (with rifampicin)

does not directly or indirectly impair the expression of maternal care, and thus call for caution

when considering the role of gut microbiota in this important social behaviour.

Despite its apparent lack of effects on maternal care, rifampicin altered three maternal

traits related to physiology and reproduction. The first trait is the production of feces pellets,

which was twice as high in rifampicin compared to control females. This result was not

surprising, as the gut microbiota often plays a key role in nutrient extraction and digestion [69]

and its alteration by antibiotics typically disturbs the host's digestive efficiency and triggers an

overproduction of fecal material. The two other traits were the weights of the 2^{nd} clutch

juveniles and 2^{nd} clutch eggs, which were (slightly) lighter in rifampicin compared to control

females. Light eggs and newly hatched juveniles are often thought to reflect low offspring

quality in insects [70], and further studies are required to confirm this association in earwigs.

Rifampicin altered none of the 21 others physiological, reproductive and longevity traits

measured in earwig mothers. Whereas these findings contrast with a large body of literature

showing the broad impact of altered gut microbial communities on host biology [4], they are in

line with a few recent studies showing that antibiotic-induced alteration of gut microbial

communities does not affect the development and survival of three Lepidoptera species

(*Danaus chrysippus*, *Ariadne merione* and *Choristoneura fumiferana* [71–73]). Together with

these findings, our results thus provide support to the idea that essential microbial symbioses

are not universal across insect species [71,74]. In these lepidopterans, the lack of microbial

symbioses has been explained by the fact that they do not depend on specific gut bacteria to

derive critical nutrition from their dietary resources [73,75]. This might also be the case in the

European earwig because it is omnivorous [42] and thus a partnership with bacteria facilitating

the digestion of specific food sources might not have been required during species evolution.

Follow-up studies will investigate whether (and which part of) the earwigs gut microbiota is

transient.

To conclude, our study reveals that rifampicin consumption alters female gut microbial

communities in earwigs, but provides no evidence for a link between this alteration and the

expression of maternal care, as well as no evidence for a strong impact of this alteration on

earwig physiology, reproduction and survival. Overall, these findings provide support to a

recent proposal that microbial enforcement of host social interactions is unlikely to evolve [31]

and to the emerging idea that not all animals have evolved a co-dependence with their

microbiome [71,74]. Nevertheless, shedding light on whether and how a symbiotic community shape hosts biology is a difficult task, mostly due to the number of players possibly involved and the complexity of their potential interactions [72]. Hence, our findings call for follow-up studies testing whether and how other members (non-sensitive to rifampicin) of the gut microbial community could shape the expression of parental care in family-living animals and/or drive important fitness parameters of earwig biology.

5-ACKNOWLEDGEMENTS

We thank Jos Kramer, Maximilian Körner, as well as Nadia Aubin-Horth, Gabrielle Davinson, an anonymous reviewer and Trine Bilde (Peer Community in Evolutionary Biology referees and recommender, respectively) for their insightful comments on this manuscript. This research has been supported by a research grant from the French Ministry of Research (to S.V.M.) and a research grant from the Agence Nationale de la Recherche (ANR-20-CE02-0002 to J.M.).

6-DATA ACCESSIBILITY

The raw metabarcoding sequence data have been deposited in the NCBI Sequence Read Archive under the BioProject PRJNA646389 with BioSample accession numbers SAMN15547835 to SAMN15547872. The Dataset and R script used for analyses of life-history traits and behaviour as well as the detailed bioinformatics pipelines reported in this manuscript are available on Zenodo [76].

7-REFERENCES

1. Kolodny O, Callahan BJ, Douglas AE. 2020 The role of the microbiome in host evolution.

Philos. Trans. R. Soc. B Biol. Sci. **375**, 20190588. (doi:10.1098/rstb.2019.0588)

2. Engel P, Moran NA. 2013 The gut microbiota of insects - diversity in structure and

function. *FEMS Microbiol. Rev.* **37**, 699–735. (doi:10.1111/1574-6976.12025)

3. Cleveland LR. 1923 Symbiosis between termites and their intestinal protozoa. *Proc. Natl.*

Acad. Sci. U. S. A. **9**, 424–428. (doi:10.1073/pnas.0913714)

4. Moran NA, Ochman H, Hammer TJ. 2019 Evolutionary and ecological consequences of

gut microbial communities. *Annu. Rev. Ecol. Syst.* **50**, 20.1-20.25. (doi:10.1146/annurev-

ecolsys-110617- 062453-062453)

5. Broderick NA, Lemaitre B. 2012 Gut-associated microbes of Drosophila melanogaster.

Gut Microbes **3**. (doi:10.4161/gmic.19896)

6. Itoh H, Tago K, Hayatsu M, Kikuchi Y. 2018 Detoxifying symbiosis: Microbe-mediated

detoxification of phytotoxins and pesticides in insects. *Nat. Prod. Rep.* **35**, 434–454.

(doi:10.1039/c7np00051k)

7. Wei G, Lai Y, Wang G, Chen H, Li F, Wang S. 2017 Insect pathogenic fungus interacts with

the gut microbiota to accelerate mosquito mortality. *Proc. Natl. Acad. Sci. U. S. A.* **114**,

5994–5999. (doi:10.1073/pnas.1703546114)

8. Kikuchi Y, Hayatsu M, Hosokawa T, Nagayama A, Tago K, Fukatsu T. 2012 Symbiont-

mediated insecticide resistance. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 8618–22.

(doi:10.1073/pnas.1200231109)

9. Wang GH, Berdy BM, Velasquez O, Jovanovic N, Alkhalifa S, Minbiole KPC, Brucker RM.

2020 Changes in microbiome confer multigenerational host resistance after sub-toxic

pesticide exposure. *Cell Host Microbe* **27**, 213-224.e7. (doi:10.1016/j.chom.2020.01.009)

10.   Andongma AA, Wan L, Xue-ping D, Akami M, He J, Clarke AR, Chang-Ying N. 2018 The

impact of nutritional quality and gut bacteria on the fitness of *Bactrocera minax* (Diptera:

Tephritidae). *R. Soc. Open Sci.* **5**. (doi:10.1098/rsos.180237)

11.   Fortin M, Meunier J, Laverré T, Souty-Grosset C, Richard F-J. 2019 Joint effects of group

sex-ratio and *Wolbachia* infection on female reproductive success in the terrestrial

isopod *Armadillidium vulgare*. *BMC Evol. Biol.* **19**, 65. (doi:10.1186/s12862-019-1391-6)

12.   Dedeine F, Vavre F, Fleury F, Loppin B, Hochberg ME, Bouletreau M. 2001 Removing

symbiotic Wolbachia bacteria specifically inhibits oogenesis in a parasitic wasp. *Proc.*

Natl. Acad. Sci. U. S. A. **98**, 6247–6252. (doi:10.1073/pnas.101304298)

13.   Hosokawa T, Kikuchi Y, Shimada M, Fukatsu T. 2008 Symbiont acquisition alters

behaviour of stinkbug nymphs. *Biol. Lett.* **4**, 45–48. (doi:10.1098/rsbl.2007.0510)

14.   Jones JC, Fruciano C, Marchant J, Hildebrand F, Forslund S, Bork P, Engel P, Hughes WOH.

2018 The gut microbiome is associated with behavioural task in honey bees. *Insectes Soc.*

65, 419–429. (doi:10.1007/s00040-018-0624-9)

15.   Buffington SA, Di Prisco GV, Auchtung TA, Ajami NJ, Petrosino JF, Costa-Mattioli M. 2016

Microbial reconstitution reverses maternal diet-induced social and synaptic deficits in

offspring. *Cell* **165**, 1762–1775. (doi:10.1016/j.cell.2016.06.001)

16.   Desbonnet L, Clarke G, Shanahan F, Dinan TG, Cryan JF. 2014 Microbiota is essential for

social development in the mouse. *Mol. Psychiatry* **19**, 146–148.

(doi:10.1038/mp.2013.65)

17.   Teseo S, van Zweden JS, Pontieri L, Kooij PW, Sørensen SJ, Wenseleers T, Poulsen M,

Boomsma JJ, Sapountzis P. 2019 The scent of symbiosis: gut bacteria may affect social

interactions in leaf-cutting ants. *Anim. Behav.* **150**, 239–254.

(doi:10.1016/j.anbehav.2018.12.017)

18.   Vernier CL, Chin IM, Adu-Oppong B, Krupp JJ, Levine J, Dantas G, Ben-Shahar Y. 2020 The

gut microbiome defines social group membership in honey bee colonies. *Sci. Adv.* **6**,

eabd3431. (doi:10.1126/sciadv.abd3431)

19.   Dillon RJ, Vennard CT, Charnley AK. 2000 Exploitation of gut bacteria in the locust. *Nature*

403, 851.

20.   Wada-Katsumata A, Zurek L, Nalyanya G, Roelofs WL, Zhang A, Schal C. 2015 Gut bacteria

mediate aggregation in the German cockroach. *Proc. Natl. Acad. Sci. U. S. A.* **112**,

201504031. (doi:10.1073/pnas.1504031112)

21.   Royle NJ, Smiseth PT, Kölliker M. 2012 *The evolution of parental care*. Oxford Uni. Oxford:

Oxford University Press.

22.   Klug H, Bonsall MB. 2014 What are the benefits of parental care? The importance of

parental effects on developmental rate. *Ecol. Evol.* **4**, 2330–2351.

(doi:10.1002/ece3.1083)

23.   Kramer J, Meunier J. 2019 The other facets of family life and their role in the evolution of

animal sociality. *Biol. Rev.* **94**, 199–215. (doi:10.1111/brv.12443)

24.   Klug H, Alonso S, Bonsall MB, Alonzo SH, Bonsall MB. 2012 Theoretical foundations of

parental care. In *The evolution of parental care* (eds NJ Royle, PT Smiseth, M Kölliker), pp.

21–39. Oxford, UK: Oxford University Press.

25.   Lewin-Epstein O, Aharonov R, Hadany L. 2017 Microbes can help explain the evolution of

host altruism. *Nat. Commun.* **8**, 14040. (doi:10.1038/ncomms14040)

26.   Nalepa CA. 2020 Origin of mutualism between termites and flagellated gut protists:

transition from horizontal to vertical transmission. *Front. Ecol. Evol.* **8**, 1–15.

(doi:10.3389/fevo.2020.00014)

27.   Biedermann PHW, Rohlfs M. 2017 Evolutionary feedbacks between insect sociality and

microbial management. *Curr. Opin. Insect Sci.* **22**, 92–100.

(doi:10.1016/j.cois.2017.06.003)

28.   Figueiredo ART, Kramer J. 2020 Cooperation and conflict within the microbiota and their

effects on animal hosts. *Front. Ecol. Evol.* **8**, 1–15. (doi:10.3389/fevo.2020.00132)

29.   Stilling RM, Bordenstein SR, Dinan TG, Cryan JF. 2014 Friends with social benefits: host-

microbe interactions as a driver of brain evolution and development? *Front. Cell. Infect.*

Microbiol. **4**, 147. (doi:10.3389/fcimb.2014.00147)

30.   Gurevich Y, Lewin-Epstein O, Hadany L. 2020 The evolution of paternal care: a role for

microbes? *Philos. Trans. R. Soc. B Biol. Sci.* **375**, 20190599. (doi:10.1098/rstb.2019.0599)

31.   Johnson KV, Foster KR. 2018 Why does the microbiome affect behaviour? *Nat. Rev.*

Microbiol. **16**, 647–655. (doi:10.1038/s41579-018-0014-3)

32.   Hosokawa T, Hironaka M, Mukai H, Inadomi K, Suzuki N, Fukatsu T. 2012 Mothers never

miss the moment: a fine-tuned mechanism for vertical symbiont transmission in a

subsocial insect. *Anim. Behav.* **83**, 293–300. (doi:10.1016/j.anbehav.2011.11.006)

33.   Fukatsu T, Hosokawa T. 2002 Capsule-transmitted gut symbiotic bacterium of the

Japanese common plataspid stinkbug, *Megacopta punctatissima*. *Appl. Environ.*

Microbiol. **68**, 389–396. (doi:10.1128/AEM.68.1.389-396.2002)

34.   Nalepa CA. 1990 Early development of nymphs and establishment of hindgut symbiosis

in *Cryptocercus punctulatus* (Dictyoptera: Cryptocercidae). *Ann. Entomol. Soc. Am.* **83**,

786–789. (doi:10.1093/aesa/83.4.786)

35.   Michaud C, Hervé V, Dupont S, Dubreuil G, Bézier AM, Meunier J, Brune A, Dedeine F.

2020 Efficient but occasionally imperfect vertical transmission of gut mutualistic protists

in a wood-feeding termite. *Mol. Ecol.* **29**, 308–324. (doi:10.1111/mec.15322)

36.   van Meyel S, Devers S, Meunier J. 2019 Love them all: Mothers provide care to foreign

eggs in the European earwig Forficula auricularia. *Behav. Ecol.* **30**.

(doi:10.1093/beheco/arz012)

37.   Greer JA, Swei A, Vredenburg VT, Zink AG. 2020 Parental care alters the egg microbiome

of maritime earwigs. *Microb. Ecol.* (doi:10.1007/s00248-020-01558-x)

38.   Boos S, Meunier J, Pichon S, Kölliker M. 2014 Maternal care provides antifungal

protection to eggs in the European earwig. *Behav. Ecol.* **25**, 754–761.

(doi:10.1093/beheco/aru046)

39.   Mas F, Kölliker M. 2011 Differential effects of offspring condition-dependent signals on

maternal care regulation in the European earwig. *Behav. Ecol. Sociobiol.* **65**, 341–349.

(doi:10.1007/s00265-010-1051-8)

40.   Kölliker M. 2007 Benefits and costs of earwig (*Forficula auricularia*) family life. *Behav.*

Ecol. Sociobiol. **61**, 1489–1497. (doi:10.1007/s00265-007-0381-7)

41.   Ratz T, Kramer J, Veuille M, Meunier J. 2016 The population determines whether and

how life-history traits vary between reproductive events in an insect with maternal care.

Oecologia **182**, 443–452. (doi:10.1007/s00442-016-3685-3)

42.   Meunier J, Wong JWY, Gómez Y, Kuttler S, Röllin L, Stucki D, Kölliker M. 2012 One clutch

or two clutches? Fitness correlates of coexisting alternative female life-histories in the

European earwig. *Evol. Ecol.* **26**, 669–682. (doi:10.1007/s10682-011-9510-x)

43.   Meunier J, Kölliker M. 2012 Parental antagonism and parent-offspring co-adaptation

interact to shape family life. *Proc. R. Soc. London B Biol. Sci.* **279**, 3981–8.

(doi:10.1098/rspb.2012.1416)

44.   Thesing J, Kramer J, Koch LK, Meunier J. 2015 Short-term benefits, but transgenerational

costs of maternal loss in an insect with facultative maternal care. *Proc. R. Soc. London B*

Biol. Sci. **282**, 20151617. (doi:10.1098/rspb.2015.1617)

45.   Kramer J, Körner M, Diehl JM, Scheiner C, Yüksel-Dadak A, Christl T, Kohlmeier P,

Meunier J. 2017 When earwig mothers do not care to share: Parent-offspring

competition and the evolution of family life. *Funct. Ecol.* **31**, 2098–2107.

(doi:10.1111/1365-2435.12915)

46.   Kramer J, Thesing J, Meunier J. 2015 Negative association between parental care and

sibling cooperation in earwigs: a new perspective on the early evolution of family life? *J.*

Evol. Biol. **28**, 1299–1308. (doi:10.1111/jeb.12655)

47.   González-Miguéns R, Muñoz-Nozal E, Jiménez-Ruiz Y, Mas-Peinado P, Ghanavi HR,

García-París M. 2020 Speciation patterns in the *Forficula auricularia* species complex:

cryptic and not so cryptic taxa across the western Palaearctic region. *Zool. J. Linn. Soc.*

(doi:10.1093/zoolinnean/zlaa070)

48.   Meunier J, Kölliker M. 2013 Inbreeding depression in an insect with maternal care:

influences of family interactions, life stage and offspring sex. *J. Evol. Biol.* **26**, 2209–20.

(doi:10.1111/jeb.12217)

49.   Werren JH. 1997 Biology of Wolbachia. *Annu. Rev. Entomol.* **42**, 587–609.

(doi:10.1146/annurev.ento.42.1.587)

50.   Zchori-Fein E, Perlman SJ. 2004 Distribution of the bacterial symbiont Cardinium in

arthropods. *Mol. Ecol.* **13**, 2009–2016. (doi:10.1111/j.1365-294X.2004.02203.x)

51.   Campbell EA, Korzheva N, Mustaev A, Murakami K, Nair S, Goldfarb A, Darst SA. 2001

Structural mechanism for rifampicin inhibition of bacterial RNA polymerase. *Cell* **104**,

901–912. (doi:10.1016/S0092-8674(01)00286-0)

52.   Rosas T, García-Ferris C, Domínguez-Santos R, Llop P, Latorre A, Moya A. 2018 Rifampicin

treatment of *Blattella germanica* evidences a fecal transmission route of their gut

microbiota. *FEMS Microbiol. Ecol.* **94**, 1–12. (doi:10.1093/femsec/fiy002)

53.   Shan HW, Zhang CR, Yan TT, Tang HQ, Wang XW, Liu SS, Liu YQ. 2016 Temporal changes

of symbiont density and host fitness after rifampicin treatment in a whitefly of the

Bemisia tabaci species complex. *Insect Sci.* **23**, 200–214. (doi:10.1111/1744-7917.12276)

54.   Chouaia B *et al.* 2012 Delayed larval development in *Anopheles* mosquitoes deprived of

Asaia bacterial symbionts. *BMC Microbiol.* **12**, 1–8. (doi:10.1186/1471-2180-12-S1-S2)

55.   Diehl JM, Meunier J. 2018 Surrounding pathogens shape maternal egg care but not egg

production in the European earwig. *Behav. Ecol.* **29**, 128–136.

(doi:10.1093/beheco/arx140)

56.   Van Meyel S, Devers S, Meunier J. 2019 Love them all: Mothers provide care to foreign

eggs in the *European earwig Forficula auricularia. Behav. Ecol.* **30**, 756–762.

(doi:10.1093/beheco/arz012)

57.   Friard O, Gamba M. 2016 BORIS: a free, versatile open-source event-logging software for

video/audio coding and live observations. *Methods Ecol. Evol.* **7**, 1325–1330.

(doi:10.1111/2041-210X.12584)

58.   Rodriguez A, Zhang H, Klaminder J, Brodin T, Andersson PL, Andersson M. 2018 ToxTrac:

A fast and robust software for tracking organisms. *Methods Ecol. Evol.* **9**, 460–464.

(doi:10.1111/2041-210X.12874)

59.   Körner M, Diehl JM, Meunier J. 2016 Growing up with feces: benefits of allo-coprophagy

in families of the European earwig. *Behav. Ecol.* **27**, 1775–1781.

(doi:10.1093/beheco/arw113)

60.   Arcila F, Meunier J. 2020 Friend or foe? The apparent benefits of gregarine (Apicomplexa:

Sporozoa) infection in the European earwig. *Int. J. Parasitol.* **50**, 461–469.

(doi:10.1016/j.ijpara.2020.01.007)

61.   McMurdie PJ, Holmes S. 2013 Phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**.

(doi:10.1371/journal.pone.0061217)

62.   Bernard M, Pascal G, Mariadassou M, Cauquil L, Chaillou S. 2017 FROGS Training on

Galaxy⍰: statistics to explore metagenomics. *HAL* **02788316**, 116.

63.   Jost L. 2007 Partitioning diversity into independent Alpha and Beta components. *Ecology*

88, 2427–2439.

64.   Benjamini Y, Hochberg Y. 1995 Controlling the false discovery rate: a practical and

powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300.

(doi:10.2307/2346101)

598 65. Montoya-Ciriaco N, Gómez-Acata S, Muñoz-Arenas LC, Dendooven L, Estrada-Torres A,

599   Díaz De La Vega-Pérez AH, Navarro-Noya YE. 2020 Dietary effects on gut microbiota of

600   the mesquite lizard Sceloporus grammicus (Wiegmann, 1828) across different altitudes.

601   *Microbiome* **8**, 1–19. (doi:10.1186/s40168-020-0783-6)

602 66. Sepulveda J, Moeller AH. 2020 The effects of temperature on animal gut microbiomes.

603   *Front. Microbiol.* **11**, 1–9. (doi:10.3389/fmicb.2020.00384)

604 67. Anderson KE, Ricigliano VA, Mott BM, Copeland DC, Floyd AS, Maes P. 2018 The queen's

605   gut refines with age: longevity phenotypes in a social insect model. *Microbiome* **6**, 108.

606   (doi:10.1186/s40168-018-0489-1)

607 68. Vogelweith F, Körner M, Foitzik S, Meunier J. 2017 Age, pathogen exposure, but not

608   maternal care shape offspring immunity in an insect with facultative family life. *BMC*

609   *Evol. Biol.* **17**, 69. (doi:10.1186/s12862-017-0926-y)

610 69. Douglas AE. 2015 Multiorganismal insects: Diversity and function of resident

611   microorganisms. *Annu. Rev. Entomol.* **60**, 17–34. (doi:10.1146/annurev-ento-010814-

612   020822)

613 70. Berrigan D. 1991 The allometry of egg size and number in insects. *Oikos* **60**, 313–321.

614 71. Phalnikar K, Kunte K, Agashe D. 2019 Disrupting butterfly caterpillar microbiomes does

615   not impact their survival and development. *Proc. R. Soc. London B Biol. Sci.* **286**.

616   (doi:10.1098/rspb.2019.2438)

617 72. Ravenscraft A, Kish N, Peay K, Boggs C. 2019 No evidence that gut microbiota impose a

618   net cost on their butterfly host. *Mol. Ecol.* **28**, 2100–2117. (doi:10.1111/mec.15057)

619 73. Hammer TJ, Janzen DH, Hallwachs W, Jaffe SP, Fierer N. 2017 Caterpillars lack a resident

gut microbiome. *Proc. Natl. Acad. Sci. U. S. A.* , 1–6. (doi:10.1073/pnas.1707186114)

74.   Hammer TJ, Sanders JG, Fierer N. 2019 Not all animals need a microbiome. *FEMS*

Microbiol. Lett. **366**. (doi:10.1093/femsle/fnz117)

75.   Chaturvedi S, Rego A, Lucas LK, Gompert Z. 2017 Sources of variation in the gut microbial

community of *Lycaeides melissa* Caterpillars. *Sci. Rep.* **7**, 1–13. (doi:10.1038/s41598-017-

11781-1)

76.   Van Meyel S, Devers S, Dupont S, Dedeine F, Meunier J. 2020 Alteration of gut microbiota

with rifampicin does not impair maternal care in the European earwig [Data set]. Zenodo.

(doi.org/10.5281/zenodo.4506829)

Table 1. Effects of rifampicin on a representative selection of 16 of the 30 measured traits reflecting maternal care, physiology, reproduction, and longevity. The effects on the 30 traits are presented in table S1. P-values significant after correction for multiple comparisons (Adj-P) are in bold. Med = Median; 1Q and 3Q = first and third quartile, respectively. N = sample size. Expected N = number of replicates per treatment that would have been necessary to obtain a statistically significant difference with a power of 0.8.

	Water				Rifampicin				Statistics			Statistical powers	
	Med	1Q	3Q	N	Med	1Q	3Q	N	W	P	Adj-P	Effect size (r)	Expected N
MATERNAL CARE													
Egg grooming (sec)	379.3	259.2	492.3	26	359.5	264.5	450.2	22	288.5	0.963	0.963	-0.004	245274
Egg defense	12.0	7.0	27.0	55	14.0	8.0	26.8	56	1397.0	0.398	0.521	0.070	798
Delay maternal return (sec)	32.0	17.0	54.0	55	27.0	10.8	60.5	56	1677.5	0.417	0.521	-0.067	871
Egg searching (%)	68.4	59.7	76.3	27	69.5	50.8	81.9	24	339.5	0.775	0.802	-0.024	6810
Juveniles defense	6.0	3.0	13.5	35	5.0	3.0	8.8	30	592.5	0.377	0.521	-0.073	733
Nymph searching (%)	80.6	77.6	85.2	21	84.2	77.2	89.6	22	208.0	0.584	0.674	0.046	1852
FEMALE PHYSIOLOGY													
Feces production	6.50	4.00	11.00	36	13.00	10.00	14.00	36	303.0	**<0.001**	**<0.001**	0.321	35
Abs. weight gain during egg care (mg)	1.28	-1.28	4.82	52	2.32	0.19	4.48	59	1308.0	0.182	0.364	0.110	321
FEMALE REPRODUCTION & LONGEVITY													
No. eggs produced in the 1st clutch	55.00	48.50	60.00	59	53.00	43.00	58.75	62	2037.0	0.280	0.521	-0.089	492
Mean egg weight in the 1st clutch (mg)	0.62	0.58	0.67	59	0.59	0.57	0.64	62	2297.5	0.015	0.112	-0.200	95
Mean juvenile weight in the 1st clutch (mg)	1.75	1.53	1.85	43	1.52	1.39	1.74	42	1244.0	**0.002**	**0.020**	-0.247	61
No. eggs produced in the 2nd clutch	28.00	20.00	33.00	33	23.50	14.50	28.75	26	528.0	0.131	0.328	-0.124	252
Mean egg weight in the 2nd clutch	0.60	0.58	0.64	33	0.56	0.54	0.59	26	628.0	**0.002**	**0.020**	-0.250	60
Mean juvenile weight in the 2nd clutch (mg)	1.49	1.38	1.60	21	1.39	1.35	1.48	10	139.5	0.150	0.346	-0.120	270
Total No. nymphs produced	32.50	22.00	52.00	32	21.50	7.50	32.50	28	596.5	0.027	0.160	-0.181	117
Female longevity (days)	323.00	293.50	361.00	39	306.00	284.50	343.25	42	994.5	0.098	0.267	-0.136	209

6-FIGURES

Figure 1 – Gut microbial composition in females. Guts were sampled either before oviposition

or at the hatching of the 1st clutch in females treated either with water or rifampicin. The ID of

each female is provided on the x-axis. These results are presented at the phylum level for

clarity, whereas statistical analyses of gut microbial diversity were conducted using OTUs. More

details in table S3.

Figure 2 – Effects of rifampicin and female sampling stage on gut microbial α- and β-

diversities. Guts were sampled either before oviposition or at the hatching of the 1st clutch in

females treated either with water or rifampicin. (A, B) *Alpha*-diversity based on Shannon and

Fisher indices as representative of all the tested metrics. Box plots depict median (middle bar)

and interquartile range (box), with whiskers extending to 1.5 times the interquartile range and

dots/triangles representing values of each sample. (C, D) *Beta*-diversity based on Bray-Curtis

and weighed- uniFrac indices as representative of all the tested metrics. Illustrations report

multidimensional scaling (MDS) results, where dots show values and ellipses represent 95%

confidence intervals.

Figure 3 – Effect of rifampicin on maternal care. (A) duration of egg grooming, (B) egg defence

against a simulated predator, (C) delay of maternal return after egg defence, (D) egg searching,

(E) nymph defence against a simulated predator and (F) nymph searching. Box plots depict

median (middle bar) and interquartile range (light bar), with whiskers extending to 1.5 times

the interquartile range and dots representing experimental values. *ns* stands for P < 0.05.

Figure 4 – Effects of rifampicin on (A) females' feces production, (B) mean juveniles weight in

the 1st clutch and (C) mean egg weight in the 2nd clutch. Box plots depict median (middle bar)

and interquartile range (light bar), with whiskers extending to 1.5 times the interquartile range

and dots representing experimental values. ***$P < 0.001$ and *$P < 0.05$.