Alteration of gut microbiota with rifampicin does not impair maternal care in

the European earwig

Sophie Van Meyel*, Séverine Devers, Simon Dupont, Franck Dedeine and Joël Meunier

Institut de Recherche sur la Biologie de l'Insecte, UMR 7261 CNRS – Université de Tours, Tours,

France

*Corresponding author: S. Van Meyel, sophie.vanmeyel@univ-tours.fr

ABSTRACT

The microbes residing within the gut of an animal host often maximise their own fitness by

modifying their host's physiological, reproductive, and behavioural functions. Whereas recent

studies suggest that they may also shape host sociality and therefore have critical effects on

animal social evolution, the impact of the gut microbiota on maternal care remains unexplored.

This is surprising, as this social behaviour is widespread among animals, often determines the

fitness of both juveniles and parents, and is essential in the evolution of complex animal

societies. Here, we address this gap in knowledge by testing whether life-long alterations of the

gut microbiota with rifampicin - a broad-spectrum antibiotic - impair the expression of pre- and

post-hatching maternal care in the European earwig, an insect exhibiting extensive forms of

maternal care towards eggs and juveniles. Our results first confirm that rifampicin altered the

mothers' gut microbial communities and revealed that the composition of the gut microbiota

differs before and after egg care. Contrary to our predictions, however, the rifampicin-induced

alterations of the gut microbiota did not modify the expression of pre- or post-hatching care.

Independent of maternal care, rifampicin increased the females' feces production and resulted

in lighter eggs and juveniles. By contrast, rifampicin altered none of the other 23 physiological,

reproductive and longevity traits measured over the females' lifetime. Overall, these findings

reveal that altering the gut microbiota does not necessarily affect host sociality.   More

generally, our results emphasize that not all animals have evolved a co-dependence with their

microbiota.

Keywords: Antibiotic, Dermaptera, Insect, Microbiome, Parental care

1-INTRODUCTION

Almost all animals harbour a gut microbiota, i.e. a community of microorganisms residing within the gut of the host [1]. Some of these gut microbes have long been known for their pathogenic effects in the hosts [2] and others for the benefits they provide to the hosts in terms of nutritional mutualism [3]. Over the last decades, however, a growing number of works has been revealing that the effects of gut microbes are much more diverse than previously thought and shape numerous physiological, reproductive, and behavioural functions of the host [4]. In the fruit fly *Drosophila melanogaster*, for instance, the gut microbiota is associated with hormone signalling, metabolism and ageing [5]. Gut microbes can also shape the hosts' immunocompetence and resistance against pesticides and toxic plant metabolites [6], such as in the mosquito *Anopheles stephensi* [7], the bean bug *Riptortus pedestri* [8] and the wasp *Nasonia vitripennis* [9]. Similarly, the gut microbiota is a key parameter in host reproduction and mating incompatibilities, as found in the fruit fly *Bactrocera minax* [10], in the terrestrial isopod *Armadillidium vulgare* [11], and the parasitic wasp *Asobara tabida* [12]. Finally, gut microbes shape the expression of numerous host behaviours, such as nutritional preference and kin recognition in *D. melanogaster* [13,14], offspring activity in the stinkbug *Megacopta punctissima* [15], and different behavioural tasks in honeybees [16].

In addition to these multiple effects, recent studies suggest that gut microbiota also play a critical role in the sociality of their hosts by shaping the expression and nature of social interactions among group members. For instance, individuals with an altered gut microbiota exhibited deficient sociability and increased social avoidance in family-living rats [17,18], as well as showed higher levels of aggressiveness toward conspecifics in colonies of the leaf-cutting ant

Acromyrmex echinator [19]. Experimental alterations of gut microbial communities also

rendered hosts less attractive to conspecifics both in the gregarious German cockroach *Blattela*

germanica [20] and in the swarm-living desert locust *Schistocerca gregaria* [21]. Most of these

social alterations were reverted when individuals received transplants of their original gut

microbiota [17–20], suggesting that certain microbes and/or the gut community as a whole

were responsible for the sociality of their host and thus, more generally, supporting the

hypothesis that gut microbes could have a key role in the evolution of their hosts' social life.

Despite these apparent links between the hosts' gut microbial communities and their

social behaviours, the role of gut microbes on the expression of parental care remains

experimentally unexplored. This is surprising, because this form of social behaviour is present in

a large and taxonomically diverse number of animal species [22], has considerable effects on

the fitness of both juveniles and parents [23] and because shedding light on this link may

provide crucial information on the role of gut microbes in the early evolution of complex animal

societies [24]. On one hand, gut microbes may indirectly drive parental care, because parents

are expected to adjust their level of care to their own condition [25] and altered gut microbial

communities can lower these conditions in multiple ways (see above). On the other hand, gut

microbes could serve as a direct promoter of parental care because, by enforcing the

expression of care in adult hosts, parental gut microbes could maximize their chances to reach

novel hosts [26–31](but see [32]). The transfer of gut microbes through parental care has been

reported in several insect species, such as the stinkbug *Parastrachia japonensis* [33], the

Japanese common plataspid stinkbug *Megacopta punctatissima* [34] and the wood cockroach

Cryptocercus punctulatus [35,36]. However, whether the gut microbiota drives the expression

of parental care remains untested.

In this study, we address this gap in knowledge by investigating whether gut microbiota

alteration with rifampicin - a broad-spectrum antibiotic - impaired the expression of pre- and

post-hatching maternal care in the European earwig *Forficula auricularia* L. In this omnivorous

insect, mothers tend their clutch of eggs over winter. During this period, mothers stop their

foraging activity to provide extensive maternal egg care in the forms of protection against

desiccation, predators and pathogens [37,38]. Upon egg hatching, mothers continue tending

their brood of newly emerged juveniles (called nymphs) for two weeks, during which they

provide post-hatching care in the forms of fierce protections against predators, grooming

behaviours, and food provisioning through regurgitation [39,40]. Interestingly, pre-hatching

care allows mothers to transfer microbes exhibiting antifungal properties to their eggshell in

the maritime earwig *Anisolabis maritima* [41], a behaviour that could also occur in the

European earwig [42]. In *F. auricularia*, pre-hatching maternal care is necessary to ensure egg

development and hatching [42], whereas post-hatching maternal care is facultative for the

development and survival of nymphs [43]. Here, we altered the gut microbiota of *F. auricularia*

females by feeding them with rifampicin during their entire adult lifetime (about 14 months)

and measured whether and how this treatment affected gut microbial communities, maternal

care, and other life-history traits. Specifically, we first determined how the antibiotherapy alters

the diversity and the structure of the gut bacterial community of females at two periods of their

life-cycle (before oviposition and at egg hatching) by sequencing 16S rRNA gene (V3-V4 region)

amplicons. We then tested the effects of rifampicin on the expression of four pre- and two

post-hatching forms of maternal care. Finally, to disentangle whether the potential link

between gut microbiota alteration and the level of maternal care is direct and/or indirect, we

investigated the effects of rifampicin on 32 other traits measured throughout the females'

lifetime and reflecting their general physiological state, reproductive success and longevity, as

well as their juveniles' development, sex-ratio and survival.

2-MATERIALS AND METHODS

2.1 Insect rearing and rifampicin treatment

The experiment involved a total of 296 *Forficula auricularia* L. (clade B [44]) males and females.

These were the first laboratory-born descendants of 74 females field-sampled in Pont-de-Ruan,

France, in 2017 and then maintained under standard laboratory conditions [45]. For each of

these 74 families, 2 males and 2 females were isolated at adult emergence and immediately fed

with a standard food mixed with either 10 µL of rifampicin (1 male and 1 female per family;

Sigma-Aldrich, PHR1806; 0.2 mg/ml) or 10 µL of water (1 male and 1 female per family). The

standard food contained agar, carrots, pollen, and cat and bird dry food [45]. Two weeks later,

148 mating pairs were set up (1 female and 1 male from the same family and same treatment).

The use of sibling pairs allowed us limiting the risk of cytoplasmic incompatibility due to inter-

familial microbiome variability, and there are only limited signs of inbreeding depression in this

species [46]. They continued to receive the same rifampicin- and water-treatment for about

two months. At that time, females were isolated to mimic natural dispersal for oviposition [45].

From oviposition to egg hatching, four forms of egg care were measured (details below). During

that time, females were not provided with food and thus not treated with rifampicin, as

mothers typically stop foraging during the period of egg care [43]. One day after egg hatching,

each family (a mother with its newly hatched juveniles) was provided with either rifampicin or

water to follow up on the pre-oviposition treatment. Three forms of maternal care towards

juveniles were measured during the following 14 days (details below), which corresponds to the

natural duration of family life [45]. At the end of these 14 days, families were split to allow

newly isolated mothers to produce a 2^{nd} clutch and permit groups of nymphs to continue their

development. The mothers and groups of nymphs continued to receive the same treatment

(rifampicin or water) until the end of the experiment, i.e. until the mother died and nymphs

reached adulthood. Throughout the experiment, the treatments were renewed twice a week

(except during egg care). All isolated adults, groups of nymphs, and families were maintained in

Petri dishes (diameter 9 cm) lined with moistened sand. More details on the experimental

setup in the supplementary material.

Rifampicin is considered one of the most potent and broad-spectrum antibiotics due to

its high-affinity binding to the RNAP β subunit, which causes the inhibition of the bacterial DNA-

dependent RNA polymerase RNAP by directly blocking the RNA elongation path [47]. It is also

commonly used to experimentally alter gut microbial communities in insects (e.g. [48–50]). The

high dose of rifampicin used in this study (about 10 times higher than the dose generally used

in smaller insect species [49,50]) was chosen to ensure gut microbial alteration and because it

did not trigger an excess of mortality in the German cockroach [48], an insect that is about the

size of the European earwig.

2.2 Effects of rifampicin on the gut microbiota

To determine whether and how rifampicin treatment altered the earwigs' gut microbial

communities, we extracted the gut of 10 females per treatment (n total = 20) on the day we

observed the first oviposition (i.e. about 2 months after being fed with or without rifampicin),

and 10 rifampicin- and 8 water-treated females one day after egg hatching (i.e. about 1 month

later). For gut extraction, we first anaesthetized each female for 2 min at -20°C and then

dissected them in a watch glass with sterilized double Q water. All dissections and

manipulations were conducted on a sterilized bench, under a Bunsen burner's sterility area and

using sterile material. Whole individual guts were extracted, placed in 100 μl of T1 extraction

buffer (Nucleo-Spin Tissue, Macherey-Nagel), and stored at −80°C until DNA extraction.

Protocol for DNA extractions is detailed in the supplementary material. Two PCR amplifications

were performed for each sample in a final volume of 35 μl to amplify a 450-bp portion of the

V3–V4 region of the 16S rRNA gene (forward primer: 5–-CTT TCC CTA CAC GAC GCT CTT CCG

ATC T**AC GGR AGG CAG CAG**-3; reverse primer: 5–-GGA GTT CAG ACG TGT GCT CTT CCG ATC

TTA CCA GGG TAT CTA ATC-3; the Illumina adapters and primers *per se* appeared in non-bold

and bold, respectively). Fifty microliters of PCR product were then sent to the GeT-PlaGe

genomic platform (GenoToul, Toulouse, France), which performed library preparation and 2 ×

250 paired-end Illumina Miseq sequencing according to the manufacturer's instructions.

Protocols of the sequencing process and bioinformatic pipelines are detailed in the

supplementary material.

2.3 Measurements of pre- and post-hatching maternal care

We measured the effects of rifampicin on four classical forms of earwig maternal egg care: egg

grooming, egg defence, delay of maternal return and egg searching exploration rate [38,51].

Egg grooming, which is used by earwig females to deposit chemical substances on the eggs and

to clean eggshell from dirt and fungi [42], was measured 15 days after egg production. We first

isolated each mother for 30 min, then returned them to their Petri dish and gently deposited

them at a distance of 5 cm from their egg clutch, and finally recorded their behaviours for the

subsequent 15 minutes on camera (SONY© Handycam HDR-CX700 camera). The resulting

movies were analysed using the software BORIS v4.0.3 [52] and the total duration of egg

grooming was defined as the total amount of time each female spent on cleaning eggs with

their mandibles [42]. Clutch defence, which reflects the females' willingness to protect their

eggs from a predator attack [53], was measured 16 days after egg production. We standardly

poked each female on the pronotum with a glass capillary (one poke per second) and then

recorded the number of pokes required until the female moved more than one body length

away from the clutch. The delay of maternal return after clutch abandonment [38], which

represents the delay after which females return to their clutch after being chased away by a

simulated predator attack [53], was measured by recording the time the female took to return

to its clutch just after the end of the clutch defence measurement. Finally, the egg searching

exploration rate, which represents the level of exploration of a novel area by a mother looking

for her eggs, was measured 21 days after egg production. We removed each mother from its

clutch of eggs, subsequently deposited it at the centre of a square arena (W: 9 cm; L: 9 cm; H:

0.5 cm) covered by a glass sheet, and then video-tracked its activity during the following 35

179 min. The video was done under infra-red light, while the individual video tracking and

180 calculation of exploration rate were conducted using the software ToxTrac v2.83 [54].

181  We then measured the effects of rifampicin on two classical forms of post-hatching

182 maternal care: the defence of and search for juveniles [51,53]. These two forms of care were

183 measured 10 days and 12 days after egg hatching, respectively, following the above-detailed

184 protocols for egg defence and egg searching activity. All the measurements of pre- and post-

185 hatching maternal care were conducted in the afternoon and under a red light as earwigs are

186 nocturnal. These measurements were conducted blindly regarding the treatments (rifampicin

187 versus control). The number of replicates for each of our measurements ranged from 21 to 56

188 (details in Tables 1 and S1).

190 *2.4 Measurements of the 24 other life-history traits in mothers and offspring*

191 We tested the effects of rifampicin on 7 proxies of female physiology, 16 proxies of female

192 reproduction and on female longevity using standard protocols [45,55]. We measured the

193 females' physiology through their feces production (reflecting their digestive/foraging activity)

194 and weight gains between two life stages. Feces production was measured two months after

195 the beginning of the treatments. Females were isolated in a new Petri Dish for 24 hours, after

196 which we counted the number of feces pellets present on the ground. The weight gains of each

197 female were measured between the days of adult emergence and oviposition, and between the

198 days of oviposition and egg hatching. Reproductive success was measured in the 1st and 2nd

199 clutches (if any), by counting the number of eggs produced, the number of days between

200 oviposition and egg hatching (egg development time), and by measuring the mean egg weight

at oviposition, the egg hatching rate, and the mean offspring weight at egg hatching. We also

counted the number of days between adult emergence and oviposition (delay until oviposition),

between the females' isolation after family life and 2^{nd} clutch oviposition (delay until 2^{nd} clutch

production), and between adult emergence and death (female longevity). We finally assessed

the females' likelihood to produce a 2^{nd} clutch (1 or 0) and the females' reproductive allocation

between the two clutches (i.e. the females' reproductive strategy [45]). This allocation was

defined as the number of 2^{nd} clutch eggs divided by the total number of eggs produced by a

female.

Because mothers and groups of nymphs continued to receive their treatment after the

end of family life, we also tested the effects of rifampicin on juvenile developmental time

between each developmental instar, on the survival rates from egg hatching until both the end

of family life and adulthood, and on the sex ratio of the resulting adults. Juvenile

developmental time was defined as the differences between the moulting date in a given instar

(or hatching date) and the moulting date of its subsequent instar at the family-level, i.e.

focusing on the first individual moulting in the clutch. Every weighing was done to the nearest

0.01 mg using a microbalance (OHAUS© Discovery DV215CD). Sample sizes for each

measurement are detailed in Tables 1 and S1. More details on the methods are provided in the

supplementary material.

2.6 Statistical analyses

Analyses of the α and β-diversity indices. The structure, composition and diversity of the

microbial communities were analysed using PHYLOSEQ R package [56] implemented in the

FROGSSTAT Phyloseq tools [57]. Diversity within the gut microbial communities (alpha-

diversity) was assessed using two richness indices which estimate the number of species in the

microbiome with correction for subsampling (Chao1; ACE), and three metrics that aim to

measure diversity by accounting for evenness or homogeneity (Shannon, Simpson, Inverse

Simpson, and Fisher) [58]. Diversity between the gut microbial communities (beta-diversity)

was assessed using two non-phylogenetically informed (Bray Curtis dissimilarity; Jaccard indice)

and two phylogenetically informed (uniFrac; Weighed uniFrac) measures of community

similarity. The metrics were analysed individually using either a General Linear Model for α-

diversity, or a Permutational Multivariate Analysis of Variance Using Distance Matrices

(PERMANOVA) for β-diversity. In these models, the values (or distance matrix for β-diversity) of

each index were entered as a response variable, while the treatment (rifampicin or water), the

sampling stage of the female (before oviposition or at egg hatching) and the interaction

between them were used as fixed factors. When required, a post-hoc analysis was conducted

by splitting the data set according to the sampling stage and then conducting PERMANOVA on

each of the resulting subsets.

Analyses of the life-history traits. Although the presented experimental design was originally

paired, i.e. 2 females per family distributed among the two treatments, the 38 life-history traits

were often measured in only one of the paired individuals (detailed sample sizes in Tables 1 and

S1). This was mostly due to time constraints, and because some females died during the 18-

months course of this experiment. These overall led to critical reductions in the number of

replicates that could be involved in a paired statistical approach (details in Table S2). We,

therefore, analysed the effects of rifampicin on the 32 measurements using a series of 31 exact

Mann Whitney U tests and 1 Pearson's Chi-squared test (for the likelihood to produce a 2nd

clutch), in which we compared the values of all the available replicates fed with rifampicin to

the values of all the available replicates fed with water. Note that the results do not

qualitatively change when we use paired analyses with the associated smallest sample sizes

(results presented in Table S2). To correct for the inflated risk of Type I errors due to multiple

testing, all p-values were adjusted using the False Discovery Rate (FDR) method [59]. All these

analyses were conducted using the software R v4.0.2 (www.r-project.org).

3-RESULTS

3.1 Description of the earwig gut microbiota

A total of 1 636 435 sequenced reads of the 16S rRNA V3-V4 region were obtained from the 38

female earwig gut samples. After sequence processing, this number went down to 1 130 241,

with 21069 to 35385 sequences per sample (median = 30595.5). The sequences were

aggregated and filtered in a total of 161 unique OTUs, which were resolved down to the family

or genus level to increase the confidence in the taxonomic assignation. All detailed information

on OTUs is given in Table S3. More than 99.90% of the sequences were assigned to four

bacterial phyla: Proteobacteria (65.94%), Firmicutes (21.12%), Bacteroidota (9.89%) and

Actinobacteriota (2.96%) (Figure 1). The remaining OTUs were assigned to Bdellovibrionota

(0.04%) and Patescibacteria (0.04%).

3.2 Comparative analyses of the α and β diversity of the gut microbiota

The gut microbial α-diversity (i.e. species richness) decreased between oviposition and egg

hatching when this diversity was measured using Chao1 ($F_{1,34}$ = 21.63, P < 0.0001), ACE ($F_{1,34}$ =

24.46, P < 0.0001) and Fisher ($F_{1,34}$ = 20.85, P < 0.0001; Figure 2) indices. This decrease was,

however, not significant when α-diversity was measured using Shannon ($F_{1,34}$ = 3.18, P = 0.084;

Figure 2) and Simpson ($F_{1,34}$ = 1.60, P = 0.214) indices. The α-diversity was overall independent

of the rifampicin treatment (Chao1: $F_{1,34}$ = 0.72, P = 0.401; ACE: $F_{1,34}$ = 0.62, P =0.435; Fisher:

$F_{1,34}$ = 0.59, P = 0.447; Shannon: $F_{1,34}$ = 1.67, P = 0.205; Simpson: $F_{1,34}$ = 0.55, P = 0.465; Figure

2), and of an interaction between female sampling stage and rifampicin treatment (all P >

0.525).

The gut microbiota β-diversity (i.e. species composition) overall changed with female

sampling stage and rifampicin treatment. This was the case with the four measured indices of

β-diversity: Bray-Curtis (Stage: $F_{1,34}$ = 5.77, P < 0.0001; Rifampicin: $F_{1,34}$ = 4.23, P < 0.0001),

JACCARD (Stage: $F_{1,34}$ = 7.76, P < 0.0001; Rifampicin: $F_{1,34}$ = 2.37, P = 0.0036), unweighted

UniFrac (Stage: $F_{1,34}$ = 6.51, P < 0.0001; Rifampicin: $F_{1,34}$ = 3.39, P = 0.0006) and weighted

UniFrac (Stage: $F_{1,34}$ = 14.10, P < 0.0001; Rifampicin: $F_{1,34}$ = 6.42, P = 0.0006). In particular,

females before oviposition harboured less Actinobacteriota and Proteobacteria compared to

females at egg hatching, while rifampicin females overall harboured less Bacteroidota and more

Firmicutes compared to untreated females (Figure 1). Interestingly, the interaction between

female sampling stage and rifampicin had no effect on the β-diversity measured using all (all P >

0.117) but the weighted UniFrac indices ($F_{1,34}$ = 2.94, P = 0.026). This interaction reflected an

effect of rifampicin on the β-diversity before oviposition ($F_{1,34}$ = 0.17, P = 0.018) but not at egg

hatching ($F_{1,34}$ = 0.97, P = 0.356).

3.3 Rifampicin and maternal care

We did not detect any effect of rifampicin on the six measured forms of egg and nymph care

(Table 1). In particular, rifampicin- and control-fed mothers spent the same amount of time in

egg grooming (Figure 3A), showed the same levels of both egg and juvenile defences against a

simulated predator attack (Figures 3B and 3E), exhibited the same delay to return to their eggs

after egg defence (Figure 3C) and showed comparable exploration rates when looking for their

eggs or their juveniles (Figures 3D and 3F).

3.4 Rifampicin and female's physiology, reproduction, and longevity

The consumption of rifampicin altered 3 of the 23 measured proxies of female physiology,

reproduction, and longevity. In particular, females fed with rifampicin produced on average

twice as many feces pellets per 24h (Figure 4; Table 1), had newly hatched nymphs that were

15% lighter (W = 1244, P = 0.002, adjusted-P = 0.025; Table S1) and laid 2^{nd} clutch eggs that

were 7% lighter (W = 628, P = 0.002, adjusted-P = 0.025; Table S1) compared to control females.

By contrast, we did not detect any effect of rifampicin on the 20 other traits (Tables 1 and S1).

3.5 Rifampicin and nymphs' development, sex-ratio and survival

We did not detect any effect of rifampicin on the juveniles' developmental speed from hatching

to adult and at every step of their development, as well as the sex-ratio of the 1^{st} clutches

(Tables 1 and S1). Similarly, the survival rate of nymphs throughout family life (i.e. from egg

hatching until day fourteen) and from the end of family life until they reached adulthood were

not impacted by the consumption of rifampicin (Tables 1 and S1).

4-DISCUSSION

Whereas gut microbial communities shape the physiology, reproduction and behaviours of a

great diversity of hosts, their importance on parental care – a key behaviour in social evolution

[22–24] - remains poorly explored [31]. In this study, we addressed this gap in knowledge by

treating females of the European earwig with rifampicin and measuring the effects on gut

microbial communities, maternal care and female physiology, reproduction, and longevity. Our

results first reveal that rifampicin altered the composition of the gut microbial community of

earwig females and show that this modification diminishes during the period of egg care.

Contrary to our predictions, however, the rifampicin-induced alterations of gut microbial

communities did not impair the expression of pre- or post-hatching care: rifampicin-treated and

control mothers showed similar levels of egg grooming, clutch defences against a predator,

maternal return and clutch searching. Independent of maternal care, our results reveal that the

consumption of rifampicin increased the females' production of feces pellets, as well as lead to

the production of lighter nymphs and lighter 2^{nd} clutch eggs. By contrast, rifampicin affected

none of the other 23 physiological, reproductive and longevity traits measured over the

females' lifetime.

Our experiment first demonstrates that the ingestion of rifampicin by earwig females

induced stage-specific modifications in the species composition (β-diversity) of the gut

microbiota but did not shape its species richness (α-diversity). These findings confirm that the

earwig gut microbiota harbours both bacterial taxa (and/or genetic variants of certain taxa) that

are sensitive and taxa that are resistant to rifampicin, and thus that our treatment successfully

altered gut microbial communities (just like in other animal species [48–50]). Independent of

rifampicin, our data also reveal that both α- and β-diversity changed from pre-oviposition to

egg hatching. This stage-specific pattern may result from the absence of food intake for about

four weeks before gut sampling in females at egg hatching compared to before oviposition [60],

and/or from the different rearing temperatures [61] and differences in female age [62]

between the two life stages. Notwithstanding the drivers of this stage-specific effect, it is

important to note that all the tested females were re-treated with rifampicin (or water) after

egg hatching so that the resulting alteration of their gut microbiota reported at oviposition

likely operated during their entire lifetime and could thus have affected all the traits measured

after egg-hatching.

335   Although gut microbial communities shape the expression of hosts sociality in numerous

vertebrate and arthropod species [17–21], our findings reveal that rifampicin-induced

alterations of this community did not affect the expression of pre- and post-hatching maternal

care in earwigs. Gut microbes were expected to directly drive the expression of parental care,

because enforcing the expression of this social behaviour may allow symbionts to reach new

hosts (i.e. offspring) that are typically young (thus offering long-lived habitats), display poor

immune defences (thus facilitating bacterial establishment and development [63]) and harbour

only a few resident microbes (thus limiting the risk of competition within the microbiome [29]).

This absence of a link between rifampicin and maternal care may first suggest that earwig

parental care is shaped by microbes that are non-sensitive to rifampicin. In insects, gut

microbial communities do not only encompass a broad diversity of bacteria (among which some

are resistant to rifampicin) but also fungi, protists and other microorganisms that could have

key roles in hosts biology [41,64,65]. Even if the previous experimental works linking gut

microbiota and host sociality focused on bacteria [17–21], future studies will be required to

confirm that no other members of the gut microbiota shape parental care in our study species.

A second hypothesis is that microbial symbionts never developed any specific capabilities to

manipulate host sociality, either because adapted strain never occurred within the microbial

populations associated with these earwig species (or populations), or because certain

antagonistic interactions (e.g. competition) among the members of the microbial community

have prevented the emergence of host social manipulation. Any symbiont species (or strain)

investing its resources to manipulate host behaviour is indeed expected to be outcompeted

within the microbiome by other species or variants that, instead, direct their resources into

growth, survival or directly transmission [32](but see for the evolution of paternal care [31]).

Finally, a third hypothesis is that the symbionts' capability to manipulate host sociality may

have changed and/or disappeared during host social evolution and consequently vanished in

the European earwig. The evolutionary drivers of family life indeed change over time [24] and,

while gut microbes may have (at least partly) driven the ancestral evolutionary shift from

solitary to family living for the reasons detailed above, the resulting benefits of parental care

for the hosts could have consolidated the expression of care and thus reduced the capability of

symbionts to control host social behaviours and/or reduced the sensitivity of the hosts to this

manipulation. Based on this hypothesis, alterations in gut microbial communities should not

shape the expression of parental care once this behaviour is established. In earwigs, pre- and

post-hatching maternal care are well established (even if their levels of expression differ

between females and the associated benefits appear to be limited for juveniles [43,53]), which

may thus have limited the maintenance of symbiont control over parental care. Overall, our

findings provide the first experimental evidence that alteration of the gut microbiota (with

rifampicin) does not directly or indirectly impair the expression of maternal care and thus

emphasize the potentially limited role of the gut microbiota in this important social behaviour.

Despite its apparent lack of effects on maternal care, rifampicin altered three female

life-history traits related to physiology and reproduction. The first trait is the production of

feces pellets, which was twice as high in rifampicin-treated compared to control females. This

result was not surprising, as the gut microbiota often plays a key role in nutrient extraction and

digestion [66] and its alteration by antibiotics typically disturbs the host's digestive efficiency

and triggers an overproduction of fecal material. The two other traits were the weights of the

2^{nd} clutch juveniles and 2^{nd} clutch eggs, which were lighter in rifampicin compared to control

females. Light eggs and newly hatched juveniles are often thought to reflect low offspring

quality in insects [67]. In the present study, however, heavier eggs and newly hatched juveniles

did not translate into higher offspring survival and improved development compared to lighter

counterparts. On a proximate level, these findings suggest that rifampicin breaks the

association between offspring weight and quality, either due to alteration in gut microbial

communities and/or antibiotic toxicity. More generally, these findings stress that rifampicin

only has a limited impact on offspring fitness, as least under laboratory conditions.

Rifampicin altered none of the 23 others developmental, physiological, reproductive and

longevity traits measured in earwig mothers and offspring. Whereas these findings contrast

with a large body of literature showing the broad impact of altered gut microbial communities

on host biology [4], they are in line with a few recent studies showing that antibiotic-induced

alteration of gut microbial communities does not affect development and survival of the three

Lepidopteran *Danaus chrysippus*, *Ariadne merione* and *Choristoneura fumiferana* [68–70].

Together with these findings, our results thus provide support to the idea that essential

microbial symbioses are not universal across insect species [68,71]. In these three Lepidoptera

species, the lack of microbial symbioses has been explained by the fact that they do not depend

on specific gut bacteria to derive critical nutrition from their dietary resources [70,72,73]. This

might also be the case in the European earwig because it is omnivorous [45] and thus a

partnership with bacteria facilitating the digestion of specific food sources might not have been

required during species evolution. Future works are nevertheless required to test whether (or

which part of) its gut microbiota is transient.

To conclude, our study reveals that rifampicin consumption alters female gut microbial

communities in earwigs, but provides no evidence for a link between this alteration and the

expression of parental care, and no evidence for a strong impact of this alteration on earwig

physiology, reproduction and survival. Our study also shows that earwig females harbour

different gut microbial communities before and after the period of egg care, a result in line with

temporal variation in the microbial communities present on eggshells in the maritime earwig

[41]. Overall, these findings provide support to the recent proposal that microbial enforcement

of host social interactions is unlikely to evolve [32] and to the emerging idea that not all animals

have evolved a co-dependence with their microbiome [68,71]. Nevertheless, shedding light on

whether and how a symbiotic community shape hosts biology is a difficult task, mostly due to

the number of players possibly involved and the complexity of their potential interactions [69].

Hence, our findings pave the way for follow-up studies testing whether other (non-rifampicin

sensitive) members of the gut microbial community could shape the expression of parental care

in family-living animals and/or drive important fitness parameters of earwig biology.

5-ACKNOWLEDGEMENTS

415 We thank Jos Kramer and Maximilian Körner for their comments on this manuscript. This

416 research has been supported by a research grant from the French Ministry of Research (to

417 S.V.M.) and a research grant from the ANR (project *MicroSoc* to J.M.).

6-DATA ACCESSIBILITY

418 The raw metabarcoding sequence data have been deposited in the NCBI Sequence Read

419 Archive under the BioProject PRJNA646389 with BioSample accession numbers SAMN15547835

420 to SAMN15547872. The life-history traits and behaviour analyses reported in this manuscript

421 can be reproduced using the data and R script available in the public repository server Zenodo

422 [74].

7-REFERENCES

1. Kolodny O, Callahan BJ, Douglas AE. 2020 The role of the microbiome in host evolution.

Philos. Trans. R. Soc. B Biol. Sci. **375**, 20190588. (doi:10.1098/rstb.2019.0588)

2. Engel P, Moran NA. 2013 The gut microbiota of insects - diversity in structure and

function. *FEMS Microbiol. Rev.* **37**, 699–735. (doi:10.1111/1574-6976.12025)

3.   Cleveland LR. 1923 Symbiosis between termites and their intestinal protozoa. *Proc. Natl.*

Acad. Sci. U. S. A. **9**, 424–428. (doi:10.1073/pnas.0913714)

4.   Moran NA, Ochman H, Hammer TJ. 2019 Evolutionary and ecological consequences of

gut microbial communities. *Annu. Rev. Ecol. Syst.* **50**, 20.1-20.25. (doi:10.1146/annurev-

ecolsys-110617- 062453-062453)

5.   Broderick NA, Lemaitre B. 2012 Gut-associated microbes of Drosophila melanogaster.

Gut Microbes **3**. (doi:10.4161/gmic.19896)

6.   Itoh H, Tago K, Hayatsu M, Kikuchi Y. 2018 Detoxifying symbiosis: Microbe-mediated

detoxification of phytotoxins and pesticides in insects. *Nat. Prod. Rep.* **35**, 434–454.

(doi:10.1039/c7np00051k)

7.   Wei G, Lai Y, Wang G, Chen H, Li F, Wang S. 2017 Insect pathogenic fungus interacts with

the gut microbiota to accelerate mosquito mortality. *Proc. Natl. Acad. Sci. U. S. A.* **114**,

5994–5999. (doi:10.1073/pnas.1703546114)

8.   Kikuchi Y, Hayatsu M, Hosokawa T, Nagayama A, Tago K, Fukatsu T. 2012 Symbiont-

mediated insecticide resistance. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 8618–22.

(doi:10.1073/pnas.1200231109)

9.   Wang GH, Berdy BM, Velasquez O, Jovanovic N, Alkhalifa S, Minbiole KPC, Brucker RM.

2020 Changes in microbiome confer multigenerational host resistance after sub-toxic

pesticide exposure. *Cell Host Microbe* **27**, 213-224.e7. (doi:10.1016/j.chom.2020.01.009)

10. Andongma AA, Wan L, Xue-ping D, Akami M, He J, Clarke AR, Chang-Ying N. 2018 The

impact of nutritional quality and gut bacteria on the fitness of *Bactrocera minax* (Diptera:

Tephritidae). *R. Soc. Open Sci.* **5**. (doi:10.1098/rsos.180237)

11. Fortin M, Meunier J, Laverré T, Souty-Grosset C, Richard F-J. 2019 Joint effects of group

sex-ratio and *Wolbachia* infection on female reproductive success in the terrestrial

isopod *Armadillidium vulgare*. *BMC Evol. Biol.* **19**, 65. (doi:10.1186/s12862-019-1391-6)

12. Dedeine F, Vavre F, Fleury F, Loppin B, Hochberg ME, Bouletreau M. 2001 Removing

symbiotic Wolbachia bacteria specifically inhibits oogenesis in a parasitic wasp. *Proc.*

Natl. Acad. Sci. U. S. A. **98**, 6247–6252. (doi:10.1073/pnas.101304298)

13. Lizé A, McKay R, Lewis Z. 2014 Kin recognition in Drosophila: The importance of ecology

and gut microbiota. *ISME J.* **8**, 469–477. (doi:10.1038/ismej.2013.157)

14. Wong AC-N, Wang Q-P, Morimoto J, Senior AM, Lihoreau M, Neely GG, Simpson SJ,

Ponton F. 2017 Gut microbiota modifies olfactory-guided microbial preferences and

foraging decisions in Drosophila. *Curr. Biol.* **27**, 2397-2404.e4.

(doi:10.1016/j.cub.2017.07.022)

15. Hosokawa T, Kikuchi Y, Shimada M, Fukatsu T. 2008 Symbiont acquisition alters

behaviour of stinkbug nymphs. *Biol. Lett.* **4**, 45–48. (doi:10.1098/rsbl.2007.0510)

16. Jones JC, Fruciano C, Marchant J, Hildebrand F, Forslund S, Bork P, Engel P, Hughes WOH.

2018 The gut microbiome is associated with behavioural task in honey bees. *Insectes Soc.*

65, 419–429. (doi:10.1007/s00040-018-0624-9)

17. Buffington SA, Di Prisco GV, Auchtung TA, Ajami NJ, Petrosino JF, Costa-Mattioli M. 2016

Microbial reconstitution reverses maternal diet-induced social and synaptic deficits in

offspring. *Cell* **165**, 1762–1775. (doi:10.1016/j.cell.2016.06.001)

18. Desbonnet L, Clarke G, Shanahan F, Dinan TG, Cryan JF. 2014 Microbiota is essential for

social development in the mouse. *Mol. Psychiatry* **19**, 146–148.

(doi:10.1038/mp.2013.65)

19.   Teseo S, van Zweden JS, Pontieri L, Kooij PW, Sørensen SJ, Wenseleers T, Poulsen M,

Boomsma JJ, Sapountzis P. 2019 The scent of symbiosis: gut bacteria may affect social

interactions in leaf-cutting ants. *Anim. Behav.* **150**, 239–254.

(doi:10.1016/j.anbehav.2018.12.017)

20.   Wada-Katsumata A, Zurek L, Nalyanya G, Roelofs WL, Zhang A, Schal C. 2015 Gut bacteria

mediate aggregation in the German cockroach. *Proc. Natl. Acad. Sci. U. S. A.* **112**,

201504031. (doi:10.1073/pnas.1504031112)

21.   Dillon RJ, Vennard CT, Charnley AK. 2000 Exploitation of gut bacteria in the locust. *Nature*

403, 851.

22.   Royle NJ, Smiseth PT, Kölliker M. 2012 *The evolution of parental care*. Oxford Uni. Oxford:

Oxford University Press.

23.   Klug H, Bonsall MB. 2014 What are the benefits of parental care? The importance of

parental effects on developmental rate. *Ecol. Evol.* **4**, 2330–2351.

(doi:10.1002/ece3.1083)

24.   Kramer J, Meunier J. 2019 The other facets of family life and their role in the evolution of

animal sociality. *Biol. Rev.* **94**, 199–215. (doi:10.1111/brv.12443)

25.   Klug H, Alonso S, Bonsall MB, Alonzo SH, Bonsall MB. 2012 Theoretical foundations of

parental care. In *The evolution of parental care* (eds NJ Royle, PT Smiseth, M Kölliker), pp.

21–39. Oxford, UK: Oxford University Press.

26.   Lewin-Epstein O, Aharonov R, Hadany L. 2017 Microbes can help explain the evolution of

host altruism. *Nat. Commun.* **8**, 14040. (doi:10.1038/ncomms14040)

27.   Nalepa CA. 2020 Origin of mutualism between termites and flagellated gut protists:

transition from horizontal to vertical transmission. *Front. Ecol. Evol.* **8**, 1–15.

(doi:10.3389/fevo.2020.00014)

28.   Biedermann PHW, Rohlfs M. 2017 Evolutionary feedbacks between insect sociality and

microbial management. *Curr. Opin. Insect Sci.* **22**, 92–100.

(doi:10.1016/j.cois.2017.06.003)

29.   Figueiredo ART, Kramer J. 2020 Cooperation and Conflict Within the Microbiota and

Their Effects On Animal Hosts. *Front. Ecol. Evol.* **8**, 1–15. (doi:10.3389/fevo.2020.00132)

30.   Stilling RM, Bordenstein SR, Dinan TG, Cryan JF. 2014 Friends with social benefits: host-

microbe interactions as a driver of brain evolution and development? *Front. Cell. Infect.*

Microbiol. **4**, 147. (doi:10.3389/fcimb.2014.00147)

31.   Gurevich Y, Lewin-Epstein O, Hadany L. 2020 The evolution of paternal care: a role for

microbes? *Philos. Trans. R. Soc. B Biol. Sci.* **375**, 20190599. (doi:10.1098/rstb.2019.0599)

32.   Johnson KV, Foster KR. 2018 Why does the microbiome affect behaviour? *Nat. Rev.*

Microbiol. **16**, 647–655. (doi:10.1038/s41579-018-0014-3)

33.   Hosokawa T, Hironaka M, Mukai H, Inadomi K, Suzuki N, Fukatsu T. 2012 Mothers never

miss the moment: a fine-tuned mechanism for vertical symbiont transmission in a

subsocial insect. *Anim. Behav.* **83**, 293–300. (doi:10.1016/j.anbehav.2011.11.006)

34.   Fukatsu T, Hosokawa T. 2002 Capsule-transmitted gut symbiotic bacterium of the

Japanese common plataspid stinkbug, *Megacopta punctatissima*. *Appl. Environ.*

Microbiol. **68**, 389–396. (doi:10.1128/AEM.68.1.389-396.2002)

35.   Nalepa CA. 1984 Colony composition, protozoan transfer and some life history

characteristics of the woodroach *Cryptocercus punctulatus* Scudder (Dictyoptera:

Cryptocercidae). *Behav. Ecol. Sociobiol.* , 273–279.

36.   Nalepa CA. 1990 Early development of nymphs and establishment of hindgut symbiosis

in *Cryptocercus punctulatus* (Dictyoptera: Cryptocercidae). *Ann. Entomol. Soc. Am.* **83**,

786–789. (doi:10.1093/aesa/83.4.786)

37.   Lamb RJ. 1976 Parental behaviour in the Dermaptera with special reference to *Forficula*

auricularia. *Can. Entomol.* **108**, 609–619.

38.   Van Meyel S, Devers S, Meunier J. 2019 Love them all: Mothers provide care to foreign

eggs in the *European earwig Forficula auricularia*. *Behav. Ecol.* **30**, 756–762.

(doi:10.1093/beheco/arz012)

39.   Staerkle M, Kölliker M. 2008 Maternal food regurgitation to nymphs in earwigs (Forficula

auricularia). *Ethology* **114**, 844–850. (doi:10.1111/j.1439-0310.2008.01526.x)

40.   Mas F, Kölliker M. 2011 Differential effects of offspring condition-dependent signals on

maternal care regulation in the European earwig. *Behav. Ecol. Sociobiol.* **65**, 341–349.

(doi:10.1007/s00265-010-1051-8)

41.   Greer JA, Swei A, Vredenburg VT, Zink AG. 2020 Parental care alters the egg microbiome

of maritime earwigs. *Microb. Ecol.* (doi:10.1007/s00248-020-01558-x)

42.   Boos S, Meunier J, Pichon S, Kölliker M. 2014 Maternal care provides antifungal

protection to eggs in the European earwig. *Behav. Ecol.* **25**, 754–761.

(doi:10.1093/beheco/aru046)

43.   Kölliker M. 2007 Benefits and costs of earwig (Forficula auricularia) family life. *Behav.*

Ecol. Sociobiol. **61**, 1489–1497. (doi:10.1007/s00265-007-0381-7)

44.   González-Miguéns R, Muñoz-Nozal E, Jiménez-Ruiz Y, Mas-Peinado P, Ghanavi HR,

538   García-París M. 2020 Speciation patterns in the *Forficula auricularia* species complex:

539   cryptic and not so cryptic taxa across the western Palaearctic region. *Zool. J. Linn. Soc.*

540   (doi:10.1093/zoolinnean/zlaa070)

541   45.   Meunier J, Wong JWY, Gómez Y, Kuttler S, Röllin L, Stucki D, Kölliker M. 2012 One clutch

542   or two clutches? Fitness correlates of coexisting alternative female life-histories in the

543   European earwig. *Evol. Ecol.* **26**, 669–682. (doi:10.1007/s10682-011-9510-x)

544   46.   Meunier J, Kölliker M. 2013 Inbreeding depression in an insect with maternal care:

545   influences of family interactions, life stage and offspring sex. *J. Evol. Biol.* **26**, 2209–20.

546   (doi:10.1111/jeb.12217)

547   47.   Campbell EA, Korzheva N, Mustaev A, Murakami K, Nair S, Goldfarb A, Darst SA. 2001

548   Structural mechanism for rifampicin inhibition of bacterial RNA polymerase. *Cell* **104**,

549   901–912. (doi:10.1016/S0092-8674(01)00286-0)

550   48.   Rosas T, García-Ferris C, Domínguez-Santos R, Llop P, Latorre A, Moya A. 2018 Rifampicin

551   treatment of *Blattella germanica* evidences a fecal transmission route of their gut

552   microbiota. *FEMS Microbiol. Ecol.* **94**, 1–12. (doi:10.1093/femsec/fiy002)

553   49.   Shan HW, Zhang CR, Yan TT, Tang HQ, Wang XW, Liu SS, Liu YQ. 2016 Temporal changes

554   of symbiont density and host fitness after rifampicin treatment in a whitefly of the

555   *Bemisia tabaci* species complex. *Insect Sci.* **23**, 200–214. (doi:10.1111/1744-7917.12276)

556   50.   Chouaia B *et al.* 2012 Delayed larval development in *Anopheles* mosquitoes deprived of

557   *Asaia* bacterial symbionts. *BMC Microbiol.* **12**, 1–8. (doi:10.1186/1471-2180-12-S1-S2)

51.   Diehl JM, Meunier J. 2018 Surrounding pathogens shape maternal egg care but not egg

production in the European earwig. *Behav. Ecol.* **29**, 128–136.

(doi:10.1093/beheco/arx140)

52. Friard O, Gamba M. 2016 BORIS: a free, versatile open-source event-logging software for

video/audio coding and live observations. *Methods Ecol. Evol.* **7**, 1325–1330.

(doi:10.1111/2041-210X.12584)

53. Thesing J, Kramer J, Koch LK, Meunier J. 2015 Short-term benefits, but transgenerational

costs of maternal loss in an insect with facultative maternal care. *Proc. R. Soc. London B*

Biol. Sci. **282**, 20151617. (doi:10.1098/rspb.2015.1617)

54. Rodriguez A, Zhang H, Klaminder J, Brodin T, Andersson PL, Andersson M. 2018 ToxTrac:

A fast and robust software for tracking organisms. *Methods Ecol. Evol.* **9**, 460–464.

(doi:10.1111/2041-210X.12874)

55. Körner M, Diehl JM, Meunier J. 2016 Growing up with feces: benefits of allo-coprophagy

in families of the European earwig. *Behav. Ecol.* **27**, 1775–1781.

(doi:10.1093/beheco/arw113)

56. McMurdie PJ, Holmes S. 2013 Phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**.

(doi:10.1371/journal.pone.0061217)

57. Bernard M, Pascal G, Mariadassou M, Cauquil L, Chaillou S. 2017 FROGS Training on

Galaxy : statistics to explore metagenomics. *HAL* **02788316**, 116.

58. Jost L. 2007 Partitioning diversity into independent Alpha and Beta components. *Ecology*

88, 2427–2439.

59. Benjamini Y, Hochberg Y. 1995 Controlling the false discovery rate: a practical and

powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300.

(doi:10.2307/2346101)

60.   Montoya-Ciriaco N, Gómez-Acata S, Muñoz-Arenas LC, Dendooven L, Estrada-Torres A,

Díaz De La Vega-Pérez AH, Navarro-Noya YE. 2020 Dietary effects on gut microbiota of

the mesquite lizard Sceloporus grammicus (Wiegmann, 1828) across different altitudes.

Microbiome **8**, 1–19. (doi:10.1186/s40168-020-0783-6)

61.   Sepulveda J, Moeller AH. 2020 The effects of temperature on animal gut microbiomes.

Front. Microbiol. **11**, 1–9. (doi:10.3389/fmicb.2020.00384)

62.   Anderson KE, Ricigliano VA, Mott BM, Copeland DC, Floyd AS, Maes P. 2018 The queen's

gut refines with age: longevity phenotypes in a social insect model. *Microbiome* **6**, 108.

(doi:10.1186/s40168-018-0489-1)

63.   Vogelweith F, Körner M, Foitzik S, Meunier J. 2017 Age, pathogen exposure, but not

maternal care shape offspring immunity in an insect with facultative family life. *BMC*

Evol. Biol. **17**, 69. (doi:10.1186/s12862-017-0926-y)

64.   Urubschurov V, Janczyk P. 2011 Biodiversity of yeasts in the gastrointestinal ecosystem

with emphasis on its importance for the host. In *The Dynamical Processes of Biodiversity*

- Case Studies of Evolution and Spatial Distribution, pp. 277–302. (doi:10.5772/24108)

65.   Arcila F, Meunier J. 2020 Friend or foe? The apparent benefits of gregarine (Apicomplexa:

Sporozoa) infection in the European earwig. *Int. J. Parasitol.* **50**, 461–469.

(doi:10.1016/j.ijpara.2020.01.007)

66.   Douglas AE. 2015 Multiorganismal insects: Diversity and function of resident

microorganisms. *Annu. Rev. Entomol.* **60**, 17–34. (doi:10.1146/annurev-ento-010814-

020822)

67.   Berrigan D. 1991 The allometry of egg size and number in insects. *Oikos* **60**, 313–321.

68.   Phalnikar K, Kunte K, Agashe D. 2019 Disrupting butterfly caterpillar microbiomes does

not impact their survival and development. *Proc. R. Soc. London B Biol. Sci.* **286**.

(doi:10.1098/rspb.2019.2438)

69.   Ravenscraft A, Kish N, Peay K, Boggs C. 2019 No evidence that gut microbiota impose a

net cost on their butterfly host. *Mol. Ecol.* **28**, 2100–2117. (doi:10.1111/mec.15057)

70.   Hammer TJ, Janzen DH, Hallwachs W, Jaffe SP, Fierer N. 2017 Caterpillars lack a resident

gut microbiome. *Proc. Natl. Acad. Sci. U. S. A.* , 1–6. (doi:10.1073/pnas.1707186114)

71.   Hammer TJ, Sanders JG, Fierer N. 2019 Not all animals need a microbiome. *FEMS*

Microbiol. Lett. **366**. (doi:10.1093/femsle/fnz117)

72.   Phalnikar K, Kunte K, Agashe D. 2018 Dietary and developmental shifts in butterfly-

associated bacterial communities. *R. Soc. Open Sci.* **5**. (doi:10.1098/rsos.171559)

73.   Chaturvedi S, Rego A, Lucas LK, Gompert Z. 2017 Sources of Variation in the Gut

Microbial Community of Lycaeides melissa Caterpillars. *Sci. Rep.* **7**, 1–13.

(doi:10.1038/s41598-017-11781-1)

74.   Van Meyel S, Devers S, Dupont S, Dedeine F, Meunier J. 2020 Alteration of gut microbiota

with rifampicin does not impair maternal care in the European earwig [Data set]. Zenodo.

(doi.org/10.5281/zenodo.4072981)

Table 1. Effects of rifampicin on a representative selection of 15 of the 36 measured traits reflecting maternal care, female physiology, reproduction and longevity, as well as nymph development and survival. The effects of rifampicin on all 36 traits are presented in table S1. P-values significant after correction for multiple comparisons (Adj-P) are in bold. Med = Median; 1Q and 3Q = first and third quartile, respectively. N = sample size.

	Water				Rifampicin				Statistics		
	Med	1Q	3Q	N	Med	1Q	3Q	N	W	P	Adj-P
Maternal care											
Egg grooming (sec)	379.34	259.20	492.28	26	359.54	264.49	450.18	22	288.5	0.963	0.963
Egg defense	12.00	7.00	27.00	55	14.00	8.00	26.75	56	1397.0	0.398	0.582
Delay maternal return (sec)	32.00	17.00	54.00	55	27.00	10.75	60.50	56	1677.5	0.417	0.582
Egg searching (%)	68.37	59.72	76.28	27	69.55	50.78	81.88	24	339.5	0.775	0.807
Juveniles defense	6.00	3.00	13.50	35	5.00	3.00	8.75	30	592.5	0.377	0.582
Nymph searching (%)	80.61	77.55	85.20	21	84.16	77.17	89.55	22	208.0	0.584	0.716
Female physiology											
Feces production	6.50	4.00	11.00	36	13.00	10.00	14.00	36	303.0	**<0.001**	**<0.001**
Abs. weight gain during egg care (mg)	1.28	-1.28	4.82	52	2.32	0.19	4.48	59	1308.0	0.182	0.407
Female reproduction & longevity											
No. eggs produced in the 1st clutch	55.00	48.50	60.00	59	53.00	43.00	58.75	62	2037.0	0.280	0.560
No. eggs produced in the 2nd clutch	28.00	20.00	33.00	33	23.00	14.00	28.50	27	561.0	0.087	0.383
Total No. nymphs produced	32.50	22.00	52.00	32	21.50	7.50	32.50	28	596.5	0.027	0.203
Female longevity (days)	323.00	293.50	361.00	39	306.00	284.50	343.25	42	994.5	0.098	0.310
Nymph development and survival											
Dvptal time from hatching to adults	81.00	75.50	85.50	35	82.00	77.25	85.75	34	565.5	0.727	0.807
Survival rate from hatching to day 14	84.38	77.50	90.00	35	85.16	75.37	91.37	34	572.0	0.786	0.807
Survival rate from day 14 to adults	42.86	29.86	59.91	35	39.88	27.27	66.67	34	620.5	0.763	0.807

6-FIGURES

Figure 1 – Gut microbial composition in females. Guts were sampled either before oviposition

or at egg hatching in females treated either with water or rifampicin. The ID of each female is

provided on the x-axis. More details in table S3.

Figure 2 – Effects of rifampicin and female sampling stage on gut microbial α- and β-

diversities. Guts were sampled either before oviposition or at egg hatching in females treated

either with water or rifampicin. (A, B) *Alpha*-diversity based on Shannon and Fisher indices

(representative of all the tested metrics), respectively. Box plots depict median (middle bar)

and interquartile range (box), with whiskers extending to 1.5 times the interquartile range and

dots/triangles representing values of each sample. (C, D) *Beta*-diversity based on Bray-Curtis

and weighed- uniFrac indices (representative of all the tested metrics). Illustrations report

multidimensional scaling (MDS) results, where dots show values and ellipses represent 95%

confidence intervals.

638 **Figure 3 – Effect of rifampicin on maternal care.** (A) duration of egg grooming, (B) egg defence

639 against a simulated predator, (C) delay of maternal return after egg defence, (D) egg searching,

640 (E) nymph defence against a simulated predator and (F) nymph searching. Box plots depict

641 median (middle bar) and interquartile range (light bar), with whiskers extending to 1.5 times

642 the interquartile range and dots representing experimental values. *ns* stands for P < 0.05.

643   **Figure 4 – Effects of rifampicin on (A) females' feces production, (B) mean juveniles weight in**

644   **the first clutch and (C) mean egg weight in the 2nd clutch.** Box plots depict median (middle bar)

645   and interquartile range (light bar), with whiskers extending to 1.5 times the interquartile range

646   and dots representing experimental values. ***$P < 0.001$ and *$P < 0.05$.