The transcriptome of regenerating zebrafish scales identifies genes involved in human bone disease

Dylan J.M. Bergen[1,2]*, Qiao Tong[1], Ankit Shukla[3], Elis Newman[1], Jan Zethof[4], Mischa Lundberg[3,5], Rebecca Ryan[1], Scott E. Youlten[6,7], Erika Kague[1] Eleftheria Zeggini[8,9], Peter I. Croucher[6,7,10], Gert Flik[4], Rebecca J. Richardson[1], John P. Kemp[3,11,$], Chrissy L. Hammond[1,$] and Juriaan R. Metz[4,$*]

$Contributed equally to this work

*Corresponding authors

dylan.bergen@bristol.ac.uk

j.metz@fnwi.ru.nl

Affiliation:

School of Physiology, Pharmacology, and Neuroscience, Faculty of Life Sciences, University of Bristol, Bristol, United Kingdom

Musculoskeletal Research Unit, Translational Health Sciences, Bristol Medical School, Faculty of Health Sciences, University of Bristol, Bristol, United Kingdom

The University of Queensland Diamantina Institute, University of Queensland, Woolloongabba, Queensland, Australia

Department of Animal Ecology and Physiology, Institute for Water and Wetland Research, Faculty of Science, Radboud University Nijmegen, Netherlands

Transformational Bioinformatics, Commonwealth Scientific and Industrial Research Organisation, Sydney, New South Wales, Australia

Bone Biology, Garvan Institute of Medical Research, Sydney, New South Wales, Australia

St Vincent's Clinical School, Faculty of Medicine, UNSW Australia, Sydney New South Wales, Australia

Institute of Translational Genomics, Helmholtz Zentrum Munchen, Neuherberg, Germany

TUM School of Medicine, Technical University of Munich and Klinikum Rechts der Isar, Munich, Germany

School of Biotechnology and Biomolecular Sciences, UNSW Australia, Sydney, New South Wales, Australia

Medical Research Council Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, United Kingdom

Key words: bone, collagen, osteoblast, musculoskeletal disease, zebrafish, scales

Abstract
Zebrafish scales are mineralised plates that can regenerate involving *de novo* bone formation. This
presents an opportunity to uncover genes and pathways relevant to human musculoskeletal disease
relevant to impaired bone formation. To investigate this hypothesis, we defined transcriptomic
profiles of ontogenetic and regenerating scales, and identified 604 differentially expressed genes
(DEGs) that were enriched for extracellular matrix, ossification, and cell adhesion pathways. Next, we
showed that human orthologues of DEGs were 2.8 times more likely to cause human monogenic
skeletal diseases ($P<8\times10^{-11}$), and they showed enrichment for human orthologues associated with
polygenetic disease traits including stature, bone density and osteoarthritis ($P<0.005$). Finally,
zebrafish mutants of two human orthologues that were robustly associated with height and
osteoarthritis (*COL11A2*) or bone density only (*SPP1*) developed skeletal abnormalities consistent with
our genetic association studies. *Col11a2^{Y228X/Y228X}* mutants showed endoskeletal features consistent
with abnormal growth and osteoarthritis, whereas *spp1^{P160X/P160X}* mutants had elevated bone density
($P<0.05$). In summary, we show that transcriptomic studies of regenerating zebrafish scales have
potential to identify new genes and pathways relevant to human skeletal disease.

Introduction

The zebrafish skeleton, similar to higher vertebrates, provides structure, organ protection, endocrine and locomotive function. During development, bone is formed by two evolutionarily conserved processes via either endochondral or intramembranous (dermal) ossification. Endochondral bone formation occurs though progressive remodelling of a cartilage template into bone, while dermal or intramembranous bone is formed directly by mesenchymal condensation of osteoblasts. Due to mechanical loading induced microfractures, bone is a dynamic tissue that undergoes constant remodelling of the calcified extracellular matrix (ECM) via a tight coupling between the functional activity of the bone-building osteoblasts and bone-degrading osteoclasts, allowing a regenerative capacity of bone throughout life (Kenkre & Bassett, 2018). In common age-related conditions such as osteoporosis, this balance is dysregulated such that catabolism exceeds anabolism, making bone more fragile and susceptible to fracture. Current pharmacotherapies for osteoporosis are limited, with most targeting the catabolic activity of osteoclasts rather than on bone anabolism. However, human genetic studies of rare bone diseases have shed light on potential bone anabolic pathways leading to development of novel therapeutics (Pathak, Bravenboer et al., 2020). For example, linkage studies have identified the WNT pathway inhibitor Sclerostin (SOST) that leads to Sclerosteosis causing high bone mass (HBM) in humans and mice when mutated (Balemans, Ebeling et al., 2001, Li, Ominsky et al., 2008). The subsequent development of a humanised monoclonal antibody (Romozumab) against SOST now offers an osteo-anabolic therapeutic that reduces fracture risk for osteoporosis patients (Cosman, Crittenden et al., 2016). This demonstrates how understanding the genetic control of bone growth and osteoblast differentiation processes allow identification of novel therapeutic targets.

One potential route to identifying new pathways and targets is through the study of organisms and tissues capable of rapid regrowth of skeletal tissues, during which bone anabolism exceeds bone catabolism. While skeletal regenerative capacity in mammals is limited to bone remodelling and fracture healing, a number of other vertebrates species such as zebrafish are able to undergo epimorphic regeneration of many organs throughout life, including the bony structures of the fins and scales (Zhao, Qin et al., 2016). Zebrafish are increasingly used as a model for musculoskeletal (MSK) research, due to their genetic tractability, availability of transgenic reporter lines and the dynamic imaging opportunities, they offer as well as the similarities of their skeletal physiology to humans and other higher vertebrates (Lleras-Forero, Winkler et al., 2020). Zebrafish fin regeneration has been extensively studied and has revealed many of the mechanisms that underpin blastema

formation and skeletal dedifferentiation and re-differentiation during tissue regrowth (Sehring &

Weidinger, 2020).

Most teleost fishes have scales which function as a protective armour and a calcium reservoir (Sire,

Donoghue et al., 2009, Yasuo, 1980).The zebrafish elasmoid scales, that also undergo complete

epimorphic regeneration within days to weeks, have recently gained attention in studies aiming at

deeper understanding of mechanisms of bone growth, regeneration, and repair. In common with

intramembranous flat bones, the elasmoid scales are formed and mineralised directly by *de novo*

differentiated osteoblasts (Pasqualetti, Banfi et al., 2012, Sire, Allizard et al., 1997). Scales are

exoskeletal elements that have been strongly reduced or completely lost in terrestrial animals during

evolution but have been retained in bony fish. Long regarded to be odontogenic in origin, relatively

recent studies have provided new insights into the classification of skeletal structures and

evolutionary and developmental origins of exoskeletal teleost scales (Dhouailly, Godefroit et al.,

2019, Shimada, Kawanishi et al., 2013). Following removal of the ontogenetic scale from the dermal

socket initiating a wound healing inflammation phase, regeneration is initiated and a small

mineralised scale plate can be observed as early as two days post-harvest (dph) (Bereiter-Hahn &

Zylberberg, 1993, Richardson, Slanchev et al., 2013, Sire & Akimenko, 2004). Regenerating scales

have a high density of osteoblasts at the posterior edge of the plate forming the leading growth

plane whose dynamics can be tracked *in toto* (Cox, De Simone et al., 2018). These osteoblasts form a

(hyposquamal) monolayer and deposit hydroxyapatite into a type I collagen rich matrix that can be

resorbed by osteoclasts (de Vrieze, Sharif et al., 2011, Guellec & Zylberberg, 1998). The collagen

matrix of scales is organised in a plywood manner resembling lamellar bone in humans (Bergen,

Kague et al., 2019, Bigi, Burghammer et al., 2001, Giraud-Guille, 1988). Similar to human bones,

scale bone turnover responds to exposure of prednisolone, alendronate, chronic hyperglycaemia,

and fatty acids (e.g. OMEGA-6) highlighting evolutionarily conserved mechanisms of bone

metabolism (Carnovali, Luzi et al., 2016, Carnovali, Ottria et al., 2016, de Vrieze, van Kessel et al.,

2014, Pasqualetti, Congiu et al., 2015). Moreover, ontogenetic scales show a fracture healing

response with recruitment of *trapc* expressing cells to the fracture site (Kobayashi-Sun, Yamamori et

al., 2020). The possibility to *ex vivo* culture fish scales as small transparent bone units, in which vital

intercellular interactions (between osteoblasts and osteoclasts), as well as cell-matrix interactions

remain intact, has triggered their use in screenings for osteogenic compounds (de Vrieze et al., 2011,

de Vrieze, Zethof et al., 2015). Hence, these features therefore warrant a deeper analysis of genes,

pathways in the scale and a comparison with genes associated with human MSK disease.

As regeneration leads to rapid bone regrowth we hypothesised that (i) osteoanabolic genes can be

identified by contrasting the transcriptomic profiles of ontogenetic and regenerating zebrafish

scales, and (ii) that these identified "bone formation" genes were likely to be enriched for human

orthologues that influence bone growth, mineralisation and susceptibility to MSK disease. To test

our hypotheses, we identified differentially expressed genes (DEGs) that distinguished regenerating

scales from ontogenetic scales, identified biological pathways that DEGs were involved in, and

investigated whether DEGs were associated with human monogenic disease, and polygenetic

disease traits: height, bone mineral density and osteoarthritis susceptability. Lastly, we performed

functional studies of two DEGs that were robustly associated with one or more human MSK disease

traits and observed abnormal endoskeletal phenotypes consistent with the human genetic

association analyses.

Results

Defininig the transcriptome of ontogentic and regenerating scales by RNA-sequencing

We first confirmed that,regenerating scales show an increase in the number of *sp7* positive osteoblasts compared to original scales formed during development (ontogenetic) as previously reported (Cox et al., 2018, de Vrieze et al., 2014) (**figure 1A**). Alkaline phosphatase (ALP) staining showed that regenerating scales contained more ALP activity (**figure 1B**). To identify the biological pathways underpinning the regeneration process, we preformed RNA-sequencing (RNA-seq) on ontogenetic (original) and regenerating scales 9 days into their regeneration (**figure 1C**).

A total of 13,170 protein coding genes were consistently expressed (i.e. 51.4% of total protein coding genome) in both ontogentic and regenerating scales (**data file S1**). Principal component analysis (PCA) of the two groups of all genes expressed showed that ontogenetic and regenerating scale groups cluster together (**figure S1A**) and that there was a a high level of correlation (Pearson's correlation of 97.1%, **figure S1B**), confirming similar expression patterns within the PCA groups. Setting the arbitrary threshold at 1.25 $\log_2$ fold change and a false discovery rate (FDR) of <0.05 showed that 604 protein coding genes had substantial differences in gene expression (**data file S2**). We observed a skew towards upregulation of genes (*n = 514* compared to *n = 90* downregulated); consistent with regenerating tissues such as the caudal fin (Padhi, Joly et al., 2004, Schmidt, Geurtzen et al., 2019) (**figure 1D**). The osteoblast transcription factor *sp7(osxterix)* was substantially upregulated (2.39 $\log_2$ fold change) in regenerating scales (**figure 1D**), but 128 genes showed a larger up regulation than *sp7*. Among the top DEGs, we noticed a high number of collagen and cell adhesion related genes (**table 1**).Two genes were highly downregulated and a homology search revealed that both genes encoded different proteins [i.e. ENSDARG00000068621 (*si:ch211-181d7.3*) and ENSDARG00000088274 (*si:ch211-181d7.1*)] that were unique to fish(**figure S2**). Both genes have NOD-like receptor (NLR) domains that possess NACHT (NTPase) and leucine-rich repeat (LRR) protein domains important for innate immune system antigen interactions (Wu, Chen et al., 2018).

Osteoblast factors such as *sp7, entpd5a/b, postnb, smpd3, phospho1, plod1a, plod2,* and *plod3* were upregulated in 9 dph scales, as were genes related to ECM growth, collagens (predominantly fibrillar collagens) and collagen remodelling including *bmp1a*, *mmp9*, *spp1*, *ostn*, *col1a1a/b*, *col1a2*, *col5a1, col5a2a/b, col10a1a/b*, *bgna/b*, *col11a1a/b* and *col11a2* (Ricard-Blum, 2011). We also observed elevated expression of genes normally associated with endochondral ossification: *col11a1a/b* and *col11a2, ihha, ptch2, dlx5a,* and *mef2cb*, suggesting that some genes are common to both modes of ossification (Long & Ornitz, 2013). While osteoclast and monocyte related factors (e.g. *tnfrsf11a* (*RANK*), chloride channel *clc7n7, cathepsin-K* (*ctsk*), and *acp5a/b (trapc)*) were expressed in both

ontogenetic and regenerating scales, differential expression was not observed (**data file S1**). To

validate the RNA-seq dataset, quantitative real-time PCR (qRT-PCR) was performed on osteoblast

and osteoclast markers and showed concordant data between RNA-seq and qRT-PCR assays of all

selected amplicons (**figure 2A**). These results indicate osteoblast activity is elevated to a greater

degree than osteoclast activity. To visualise newly formed bone, *in vivo* calcein green staining was

performed (**figure 2B**). The calcein green signal was stronger in regenerating scales, and

predominantly seen in the posterior regions of the scale, consistent with elevated expression of

bone anabolic factors such as *sp7* and *entpd5a* (**figure 2A** and **2B**).

The collagen processing pathway is highly enriched and upregulated during scale regeneration

To identify biological pathways overrepresented among DEGs, we performed gene ontology (GO)

enrichment analysis. Several, bone specific GO terms were significantly overrepresented among

DEGS, such as 'regulation of bone biomineralization' (GO:0110149; 10.9-fold, FDR=0.018),

'ossification' (GO:0001503; 6.0-fold, FDR=0.0019), and 'skeletal system development' (GO:0001501;

3.6-fold, FDR=5.3×10^{-6}). Collagen related terms were also significantly enriched (>7.1 fold,

FDR=8.7×10^{-6}) (**figure 2C, data file S3**), including 'collagen trimer', 'collagen-containing extracellular

matrix' and 'fibrillar collagen trimer' GO terms (**figure 2D**). Additional bone matrix associated

processes such as 'calcium-ion binding', 'glycosaminoglycan binding' and. 'matrix metalloprotease

(MMP) activity' were identified (**figure S3 and S4**). Further, the 'integrin signalling pathway' (P00034;

3.5-fold, FDR=0.016) was the only enriched signalling pathway, which is known to bind collagen

during cell adhesion related processes (Zeltz & Gullberg, 2016) (**data file S4**). Together, this indicated

DEGs were enriched for biological pathways involved in bone formation and collagen-matrix

synthesis, consistent with increased osteoblast activity.

We next visualised protein-protein interactions (PPI) within this set using STRING Network analysis.

This broadly confirmed GO findings and showed that the DEGs PPI network has 6.6-fold higher

number of nodes (connections) compared to the whole reactome (P<1.0×10^{-16}). There were ten

clusters with distinct functions that were significantly enriched that were highly associated with

collagen-rich ECM and cell adhesion (**figure 3**). Additional terms not identified by GO included

'regulation of insulin-like growth factor (IGF) signalling' and 'signalling by hedgehog' (HH) (**figure 3**).

Note that STRING added 10 'high-scoring' interactors to the network that were not DEGs (**figure**

S5A). There were four clusters unlinked to the main network, related to cytoskeletal structure or cell

motility such as actin-myosin (cluster 1 and 4), cell motility proteins that regulate leading edge

cytoskeletal modelling (cluster 2), and enzyme function (cluster 3) (**figure 3** and **table S1**). Allowing

less stringent interaction evidence scores (≥ 0.4) showed that *sp7* and *secreted phospho protein 1*

(*spp1*, encoding Osteopontin (Opn)) factors have connections with both the collagen and bone factor

clusters, the latter showing many connections to adjacent osteogenic clusters that for example

contain *phospo1* and *ostn* (**figure S5B**). These findings were verified with PANTHER 'Reactome

pathways' analysis (**data file S5**). These GO and PPI analyses showed that the RNA-seq dataset is

specific for an osteogenic collagen rich regenerating tissue.

We next verified RNA expression of *col1a2,* osteoblast deposited collagen *col10a1a,* the collagen

nucleation factor proteoglycan *bgna,* and hedgehog signalling ligand *ihha* that represent some of

these clusters identified above. The expression of the gap junction gene *cx43* was also assessed as it

regulates fin regeneration growth, and within our PPI network it was connected to the IGF cluster

known to modulate bone growth (Bhattacharya, Hyland et al., 2020). These amplicons were

assessed in regenerating scales from an independent experiment and all showed similar trends

between RNA-seq and qRT-PCR, validating these findings (**figure 4A**). Note, a subset of genes

showed similar profiles between the RNA-seq and independent experiments (**figure S6**). Ontogenetic

scales showed weak *col1a1a* expression, predominantly located in the epidermis adjacent to *sp7*

positive cells at the posterior edge of the scale (**figure 4B**). In regenerating scales, *col1a1a* promoter

activity was elevated along with expanded *sp7* expression covering most of the bony unit. We

observed elevated activity at newly forming circuli associated with thickening of the calcified ECM

(**figure 2B** and **figure 4B**). We performed the same procedure in *col10a1a:Citrine* and

col2a1a:mChrerry double transgenic zebrafish where the *col2a1a* reporter functions as a negative

control as it was not a DEG. This showed that ontogenetic reporter expression of both transgenes

was absent. At 9 dph, *col10a1a* reporter expression was observed at the posterior edge of the scale

and interestingly also at the newly developing circuli (thicker ECM) located at the anterior region of

the scale (**figure 4C**). A mCherry signal was not detected in ontogenetic or regenerating scales. These

findings confirm that the RNA-seq dataset of regenerating scales contains gene networks with osteo-

active factors that actively assemble (nucleate) a calcified collagen-rich matrix.

Differentially expressed genes are enriched for human orthologues that cause monogenic skeletal

disease, and that associate with polygenic musculoskeletal traits and disease

Next, we used a Fisher's Exact test of independence, in conjunction with the ISDS Nosology and

Classification of Skeletal Disorders database (Mortier, Cohn et al., 2019) to show that human

orthologues of zebrafish DEGs were 2.8 times more likely to cause monogenic skeletal dysplasia in

humans than expected by chance (P=$8×10^{-11}$) (**figure 5A**). Specifically, human orthologues of 47 DEGs

resulted in one or more primary bone dysplasia in humans when mutated (**data file S6**). Subgroup

analysis revealed further evidence of of enrichment for genes that cause: 'Osteogenesis Imperfecta

and decreased bone density' (8-fold enrichment, P=7.9×10⁻¹⁰), 'abnormal bone mineralisation' (7-fold,

P=1.6×10⁻³), 'collagen type 11' (18-fold, P=3.1×10⁻³), 'metaphyseal dysplasia' (6.7-fold, P=7.9×10⁻³),

and several others (**figure 5B & data file S6**).

MAGMA competitive gene set analysis was used to investigate the relationship between genetic

variation surrounding human orthologues of 459 mappable zebrafish DEGs and quantitative

ultrasound derived heel bone mineral density (eBMD) measured in 378,484 white European adults

(**figure 5A**). Strong evidence of enrichment was observed, suggesting that human orthologues of

zebrafish DEGs were on average more strongly associated with the eBMD than all other human protein

coding genes in the genome (β=0.20, SE=0.079, P=0.005, **table 2**). In a sensitivity analysis, we adjusted

further for the set of orthologues that could not be mapped between fish and humans, and for the

set of orthologues that was not expressed in ontogenetic and/or regenerating scales. Adjustment for

these potential confounders reduced the magnitude of enrichment by a third, however the set of DEGs

remained enriched for BMD associated orthologues (β=*0.14, SE=0.080, P=0.04*). Post hoc permutation

analysis suggested that enrichment was not attributable to all DEGs, but rather that a subset of DEGs

accounted for most of the enrichment. To further investigate this finding, DEGs were stratified

according to whether they were up- or downregulated, and each set was re-analysed. Enrichment for

BMD associated orthologues was stronger (β=0.27, SE=0.084, P=0.0006) for the 402 upregulated DEGs

and no evidence of enrichment was observed for the 57 downregulated DEGs (β=–0.31, SE=0.23,

P=0.91) (**figure S7A**). Subsequent post hoc analysis suggested that enrichment for BMD associated

orthologues was attributable to at least ~40% of the upregulated DEGs. We repeated the analysis

using height and osteoarthritis (OA) and observed evidence that DEGs were enriched for both

outcomes (P<0.005, **figure 5A**, **table 2, figure S7B**). Notably, post hoc analysis revealed that the

enrichment for height and OA associated human orthologues was not attributable to all DEGs.

Stratified analysis revealed that upregulated DEGs were more strongly enriched for height associated

orthologues, whereas the enrichment for OA associated genes remained largely unchanged (**figure

S7B, table 2**).

Finally, one advantage of MAGMA is that it first performs gene-based tests of associations between

each human protein coding gene in the genome, and the outcome of interest (see supplementary

methods for more detail). Consequently, once enrichment is observed we can prioritise orthologues

of DEGs for functional validation by comparing their relative strength of association with BMD, height

and/or OA (**figure 5C** and **data file S7**). Using this approach, we identified human orthologues of *spp1*

and *col11a2* that were both differentially upregulated in regenerating fish scales, and showed

different patterns of association with all three traits. Specifically *SPP1* was robustly associated with BMD ($P=4\times10^{-15}$), but not height ($P=0.16$) or OA ($P=0.16$), whereas *COL11A2* was robustly associated with height ($P=4.1\times10^{-13}$) and OA ($P=3.6\times10^{-7}$), but did not meet the stringent gene-wide significance threshold of $P<2.63\times10^{-6}$ for BMD (i.e. $P=2.5\times10^{-3}$). Given these contrasting patterns of association, together with the availability of the corresponding mutant zebrafish lines, we chose to further investigate *spp1* and *col11a2* expression and establish whether mutant fish developed skeletal abnormalities that were consistent with the above mentioned human genetic associations.

Zebrafish mutants of differentially expressed genes show skeletal phenotypes consistent with their expression profile and with human genetic associations

Human orthologues of DEGS that had unique, or pleiotropic associations with eBMD, height and/or OA were identified (**figure 5C**), and qRT-PCR was used to confirm that the corresponding DEGs (including *spp1* and *col11a2*) were differentially expressed (**figure 6A**). We further assessed transgenic reporter expression of *spp1* (*spp1:mCherry*) to determine gene expression localisation. The expression pattern of *spp1:mCherry* was exclusively localised at the distal rim of the ontogenetic scale, similar to *sp7:GFP* positive osteoblast localisation (**figure 6B**). Interestingly, these *sp7* positive (sub)marginal cells at the rim are classed as more mesenchymal osteoblasts involved in *de novo* bone formation in ontogenetic scales (Iwasaki, Kuroda et al., 2018). In regenerating scales, *sp7* expression was seen more broadly but elevated at the posterior edge which was followed proximally by an increased *spp1* signal with reduced *sp7* expression in the same area (**figure 6B**). The width of *spp1* expression was increased compared to ontogenetic scales, implying an expansion of its activity during bone growth and mineralisation in the regenerating scale (**figure 6C** and **6D**).

We next measured axial skeleton length from 3D micro-CT renders of *col11a2*[Y228X/Y228X] and *spp1*[P160X/P160X] mutant zebrafish. This showed significant reduced length in *col11a2* mutants whilst *spp1* mutants exhibited a milder tendency (**figure 7A**). Additionally, we segmented the lower jaw (mandibular arch) and caudal vertebrae of these 3D micro-CT renders and measured several histomorphological parameters using element landmarks as set out in **figure S8A-B**. These measurements of the lower jaw demonstrated that *col11a2* fish have reduced width and length in the lower jaw whereas *spp1* fish did not show significant changes (**figure 7B**). When we calculated the ratio between length and width, both mutant lines showed a mild alteration in lower jaw element proportion (**figure 7C**). Interestingly, in contrast to wildtype and *spp1* mutants, the *col11a2* mutants exhibited altered joint shape consistent with the predicted OA phenotype in adulthood and as seen in larval jaws previously (**figure 7D**)(Lawrence, Kague et al., 2018). Measurements of the centrum, neural arch, and haemal arch of the vertebra did not show significant changes in either

mutant line (**figure 7E-G, figure S8C-G**) These histomorphological data show that indeed *col11a2* fish

showed bone growth defects while *spp1* mutants did not.

To test the prediction that *spp1* mutants would have abnormal BMD, we quantified mean volumetric

BMD from sites formed via different modes of ossification: the parietal plate of the cranium and the

first caudal vertebra as these are formed by intramembranous ossification and by ossification

directed by the notochord sheath, respectively. Consistent with our prediction, we observed an

increase in BMD for *spp1* mutants in both elements whereas no evidence of altered BMD was

observed for *col11a2* mutants (**figure 7H**). Interestingly, we observed a physical thickening of the

vertebral arches and anterior shaft of the vertebral centrum in *spp1* mutants consistent with the

relationship seen with the increased BMD (**figure 7G** and **figure S8E**). Taken together, our data

suggest that genes that are differentially expressed during scale regeneration play a role in wider

regulation of skeletal homeostasis at other skeletal sites in the zebrafish, and that mutants in these

genes have phenotypes that are largely consistent with those observed in humans.

Discussion

In this study we define, for the first time, the transcriptome of ontogenetic and regenerating zebrafish scales and show that DEGs are enriched for biological pathways involved in osteoblast mediated bone formation, many of which are conserved in humans. By integrating this information with large-scale human genetic association studies, we show that human orthologues of DEGs were likely to play a role in the pathogenesis of monogenic and polygenic human MSK disease. Our findings suggest that zebrafish scale regeneration has the potential to help us better understand biological function and pathways relevant to human skeletal health and disease.

Transcriptomic profiling revealed that regenerating scales predominantly upregulate gene expression of many osteoblast genes, correlating with the high abundance of metabolically active osteoblasts. The high expression of genes involved in ECM deposition, principally genes regulating collagen synthesis, processing, and deposition fits with the regeneration of the collagen-rich matrix of the bony scale. We demonstrated that while *col1a1a* is expressed throughout the regenerating scale it is largely excluded from the leading edge labelled by *sp7.* By contrast, *col10a1a* is strongly expressed in leading edge concomitant and newly forming lateral circuli that are associated with thickening of the calcified matrix in accordance with the role of type X collagen as an early marker of ossifying tissues in the zebrafish (Debiais-Thibaud, Simion et al., 2019, Eames, Amores et al., 2012, Hammond & Schulte-Merker, 2009). Amongst DEG collagens, *col11a2* was one of the highest DEGs in this profile. Interestingly, type XI collagen is more frequently associated with cartilage matrix formation; interacting with type II collagen to regulate spacing and nucleation of fibrils in the ECM (Gregory, Oxford et al., 2000, Li, Lacerda et al., 1995). As well as the collagens themselves, matrix processing genes coding for MMPs were also upregulated. MMPs are important to breakdown the dense type I collagen matrix and this process is crucial for tissue growth, shape, release and distribution of signalling molecules, and cell migration (Page-McCaw, Ewald et al., 2007). We have previously shown that *mmp9* is upregulated during scale regeneration and *mmp9* expression was observed adjacent to newly deposited matrix and TRAP positive cells (de Vrieze et al., 2011).

Moreover, we observed transcriptional upregulation of ECM proteoglycans that bind collagens, such as biglycan *(bgna/b*), which are important in regulating collagen nucleation and remodelling (Douglas, Heinemann et al., 2006). Interestingly, biglycan has been shown to modulate ECM accessibility and regulate diffusion of signalling molecules, such as Bmp4, regulating bone strength in mice (Chen, Fisher et al., 2004, Xu, Bianco et al., 1998). Related to the abundant presence of collagen related factors, cell adhesion genes, in particular integrin signalling, were enriched and connected to the collagen and MMP networks. Cell adhesion and the ECM are interlinked and both must be tightly regulated during rapid growth (Zeltz & Gullberg, 2016), such as osteogenic regeneration of the scale.

Cell adhesions are heavily modulated to allow cell rearrangements and tissue mechanics in

regenerative tissue expansion contexts. For example, in a mammalian epidermal wound healing

response it is crucial for epithelial regeneration (Mosaffa, Tetley et al., 2020).

Pathway analysis revealed signalling pathways related to ECM formation and expansion, such as

integrin, HH and IGF. The absence of enrichment (note these were also not under-represented) of

Wnt, Bmp, and Fgf pathway genes is likely to be associated with the timing of the transcriptomic

profile (9 dph). For example, it has been shown that Wnt and Fgf signalling are required for the

initiation of scale regeneration, with inhibiting Fgf or canonical Wnt signalling leading to a failure to

form scales (Aman, Fulbright et al., 2018). HH signalling controls morphogenesis of the scale over a

prolonged time period, where the HH ligand secreted from the epidermis regulates the rate of ECM

deposition by scale osteoblasts (Aman et al., 2018, Iwasaki et al., 2018).

Gene ontology analysis suggested that transcriptomic profiles of regenerating zebrafish scales were

enriched for biological pathways involved in bone formation and mineralisation. These pathways are

largely conserved in humans, prompting us to investigate whether DEGs were enriched for human

orthologues that cause human monogenic skeletal disorders, and that associate with polygenic MSK

traits and disease. Gene set analysis involving human genetic association studies complemented

findings from our gene ontology analysis, suggesting that DEGs were most strongly enriched for

human orthologues that cause monogenic disorders that were broadly characterised by abnormal or

decreased bone mineralisation, and separately, abnormal cartilage formation and bone growth.

Further analysis involving human polygenetic disease traits provided additional evidence that

upregulated (but not downregulated) DEGs, were enriched for human orthologues associated with

bone mineralisation as captured by heel bone ultrasound. Importantly, post hoc permutation analysis

revealed that although the set of upregulated DEGs was strongly enriched, it was likely that

enrichment was not attributable to all DEGS, but rather a nested subset (or subsets) of genes that

corresponded to one or more biological pathway(s) that regulates bone homeostasis in humans. A

possible explanation is that DEGs were identified using bulk RNA-sequencing, a method that captures

a heterogenous mixture of transcriptome signature profiles that define different cell types, states, and

their biological pathways, some of which may or may not be involved in bone mineralisation during

scale regeneration. Importantly, the small sample size of our transcriptomic study also limited our

ability to identify all DEGs and their associated biological pathways. However, as the scale contains a

heterogeneous cell population, this is the most likely cause of not identifying the nested subsets of

DEGs responsible for the enrichment. Hence, future studies may benefit by focusing on transcriptomic

profiling of separate cell populations from scales at different stages of regeneration to determine their

expression profiles to dissect the different pathways responsible for osteo-anabolic growth. This could

be achieved by, for example, using fluorescent automated cell sorting (FACS) to isolate various

populations using cell specific transgenic reporters (e.g. *sp7:GFP*) or even single cell RNA-seq

technologies to elucidate the transcriptional state of all different cell type populations in the

regenerating scale.

Given that DEGs appeared enriched for human orthologues associated with human MSK traits and

disease, we chose to perform detailed skeletal phenotyping on *spp1* and c*ol11a2* zebrafish mutants

to investigate whether they developed skeletal abnormalities that were consistent with their gene-

based associations with height, eBMD and/or OA. Both genes were upregulated DEGs in

regenerating scales and human orthologues of these genes showed contrasting patterns of

association with human disease traits (i.e. *SPP1* robustly associated with eBMD only*, and *COL11A2*

robustly associated with height and OA, and suggestively associated with *BMD).* Mutant fish

developed skeletal abnormalities largely consistent with those predicted by the human genetic

association studies. Notably, in humans, mutations in *COL11A2* lead to Stickler syndrome, a

condition associated with craniofacial dysplasias, and joint abnormalities that lead to premature OA

(Couchouron & Masson, 2011). No clear human phenotype for *SPP1* (osteopontin) has been defined

yet, but it plays a complex role in regulating repair and regeneration of multiple tissues in

vertebrates. We show that *spp1:mCherry* was up-regulated in the regenerating scale adjacent to pre-

osteoblastic sp7+ cells which is consistent with previous reports that expression of *spp1* is up-

regulated during bone remodelling in both zebrafish fins and mammals (Morinobu, Ishijima et al.,

2003, Sousa, Valerio et al., 2012, Terai, Takano-Yamamoto et al., 1999). Moreover, *spp1* mutants

have elevated BMD in the cranial and axial endoskeleton, in line with murine studies where loss-of

Spp1 leads to enhanced mineral content in some parts of trabecular bone (Boskey, Spevak et al.,

2002). Mouse studies have shown that *Spp1* determines mineralisation rate in the skeleton through

regulating *Phospho1* expression (Holm, Gleberzon et al., 2014, Yadav, Huesa et al., 2014). For

example, Osteopontin expression is detected during cranial suture closure in both mice and

zebrafish, suggesting a role in appropriate timing of pre-osteoblast differentiation (Kim, Lee et al.,

2003, Morinobu et al., 2003, Topczewska, Shoela et al., 2016). Outside of its function in skeletal

tissue, mammalian *Spp1* is functionally diverse, as it promotes angiogenesis and its expression is also

triggered during cutaneous wound healing to control the rate of repair (Dai, Peng et al., 2009, Mori,

Shaw et al., 2008). As *Spp1* is a component of the ECM and possesses integrin binding domains, it

can bind various integrins to modulate cell adhesion to the collagen matrix important during i.e.

tissue growth (Lamort, Giopanou et al., 2019). As we see high *spp1:mCherry* expression adjacent to

sp7+ pre-osteoblasts in the regenerating scale, it implies that *spp1* could play a role in differentiation

of these pre-osteoblasts and therefore timing of scale mineralisation and the association with

human eBMD is suggestive of a conserved function in control of BMD across species.

Since the prevalence of diseases with pronounced bone fragility phenotypes is increasing due to an

ageing population there is a need to discover and rapidly test new bone growth candidate genes that

could act as drug targets. These multi-factorial diseases have complex genetic and physiological

underpinnings that are not well understood, and as human bone samples are hard to obtain there is

a strong demand for models to study bone pathophysiology. The regenerating zebrafish scale could

therefore function as an additional model to study and discover osteo-anabolic factors relevant to

human skeletal diseases. The relative abundance of teleost scales (around 200 per fish) and their

amenability for imaging make them an ideal model for regenerative studies, and unlike caudal fins,

scales can be treated as independent bone units, each harbouring their native complex tissue

environment. Scales can be cultured *ex vivo* in a semi-high throughput multi-well format, offering an

avenue to test compounds that could complement established tissue culture and *in vivo*

pharmacology studies (Bergen et al., 2019, de Vrieze et al., 2015). By using a multi-disciplinary

approach we: i) described a transcriptomic profile of regenerating zebrafish scales, ii) showed that

DEGs are enriched for biological pathways involved in bone formation, and for genes associated with

MSK disease in human populations and, iii) studied mutant zebrafish for two DEGs and observed

skeletal abnormalities in both fish that were consistent with results from human genetic association

studies. In conclusion, we show that integrative analysis involving zebrafish transcriptomics and

human genetic association studies is feasable, and that future studies involving zebrafish scales have

potential to better our understanding of bone formation in general, and how defects in this complex

process contribute to musculoskeletal disease pathogenesis in humans.

Material and Methods

Zebrafish husbandry, mutant and transgenic lines

Zebrafish were maintained under normal husbandry conditions(Alestrom, D'Angelo et al., 2020). Experiments were locally ethically reviewed (by AWERB at University of Bristol (Bristol, UK) (UoB) and Radboud University (Nijmegen, NL) (RU) respectively) and performed under UK Home Office project licence 30/3408 (UoB) or RU-DEC2014-059 (RU).

Wildtype AB/TL (UoB) and AB (RU) strains were used. Mutant lines (AB/TL background, UoB) have been previously described; *col11a2^{sa18324}* carries a nonsense mutation causing a premature stop codon at tyrosine position 228 (ENSDART00000151138.3), henceforth called *col11a2^{Y228X}* (Lawrence et al., 2018) and *spp1^{CGAT327-330del}* carrying a deletion leading to a frameshift resulting in a premature stop codon nonsense mutation at proline position 160 (ENSDART00000101261.6), henceforth called *spp1^{P160X}* (Bevan, Lim et al., 2020). Transgenic lines are listed in the **Supplemental Methods**.

Elasmoid scale harvesting and imaging

Anaesthetised fish (0.05% (v/v) tricaine methanesulfonate (MS-222) (UoB), 0.1% (v/v) 2-phenoxyethanol (RU)) were put on a wet tissue containing system water and anaesthetic, and scales plucked under a microscope with a watchmaker's tweezers from the midline of the lateral flanks near the dorsal fin. Fluorescent microscope images of flanks (*in toto*) and single harvested scales (*in situ*) were acquired on a fluorescent stereomicroscope (Leica Microsystems, Germany), using 2x, 4x, and 8x magnification.

Adult fish were immersed in 40 µM Calcein (Sigma-Aldrich, cat# 154071-48-4) Danieau's buffer solution (pH 7.4) for two hours and washed in system water for at least 15 minutes prior to imaging.

Alkaline phosphatase staining

Scales were collected in ALP buffer (100 mM Tris-HCl (pH 9.5), 100 mM NaCl, and 50 mM $MgCl_2$) and stained in ALP buffer containing 2% (v/v) NBT/BCIP (Sigma, cat# 11681451001). After a brief wash in deionised water, scales were mounted on a microscope slide containing 10 % (w/v) Mowiol® 4-88 (Sigma-Aldrich cat# 9002-89-5) in 25% (v/v) glycerol solution. Images were taken on an upright microscope (Leica).

RNA isolation, RNA sequencing, transcriptomic mapping and analysis

Approximately 40 scales were collected from a standardized area on the left flank of 1-year old male zebrafish; the area that extends from just behind the operculum to the implant of the dorsal fin. The

area may included multiple rows of scales of similar size and shape. Total RNA was isolated from

ontogenetic and regenerating scales (n=3 fish per group, RU) by using Trizol (Invitrogen) for RNA-

sequencing (RNA-seq) and downstream qRT-PCR testing.

RNA-seq was performed by ZF-GENOMICS (Leiden, NL) and involved quality control of total RNA

libraries on 2100expert bioanalyzer (Agilent) that resulted in RIN scores of >9.2. Illumina RNAseq

library preparation involved standard 6 nucleotide adaptor ligation. Paired single read 1 x 50

nucleotide runs (10 million reads; 0.5 Gb per sample) were performed on an Illumina Hiseq2500

system.

Transcriptome Mapping and Differential Expression Analysis
Raw reads were mapped to the GRCz11 primary assembly (Ensembl version 99) using STAR (version

STAR_2.5.4b) software pipeline (Dobin, Davis et al., 2013). The read count table for all genes

mapped was obtained from the mapping step and filtered to leave out lowly expressed genes by

only keeping genes that had at least 5 mapped reads over all samples. The differential gene

expression analysis was performed using the R-package DESeq2 (version 1.28.1)(Love, Huber et al.,

2014) including the medians of ratio normalisation step to account for the bias in sequencing

depth/coverage and RNA composition of samples. For determining DEGs, we used a threshold of

1.25 $\log_2$ fold (2.4-fold) change and a false discovery rate (FDR) of <0.05. Further details are in the

Supplemental Methods.

For downstream analyses, genes were classed as 'expressed' in scales when all three ontogenetic

and regenerating scale samples produced a >5 normalised read count (background gene list).

Arbitrary threshold for differential expression was set at ±1.25 $\log_2$ fold and an adjusted p-value

(padj) of ≤0.05 (DEG list).

Gene ontology enrichment and STRING network analysis
DEG and background expression gene lists' gene symbols were uploaded to GOrilla (Eden, Navon et

al., 2009) using *Danio rerio* and 'two unranked lists of genes' as settings. The hierarchical images and

Microsoft Excel files were used for figure making. Ensembl IDs of DEGs were analysed with an

'Overrepresentation Test' (Released 20200407) using Fisher's exact test and False Discovery Rate

correction in PANTHER Gene Ontology (release 2020-06-01), and PANTHER 'Pathways' and

'Reactome pathways' (PANTHER version 15.0) software (Mi, Muruganujan et al., 2019).

For STRING (v11) gene network analysis (Szklarczyk, Gable et al., 2019), the DEG set (zebrafish gene

symbols) was uploaded and interaction score (high, ≥0.7 or medium ≥0.4), number of interactions

(one shell with max. 10 interactions), and active interaction sources (all were on) were set. More

details are in the **Supplemental Methods**.

Gene set enrichment analysis involving monogenic and polygenic skeletal traits and disease

In conjunction with ISDS Nosology and Classification of Skeletal Disorders database (Mortier et al.,

2019), evidence of enrichment for human monogenic skeletal disease causing genes either total or

within each individual nosology-defined skeletal disorder groups) was performed using Fisher's Exact

Test of Independence as described in detail in the **Supplemental Methods** and (Youlten, Kemp et al.,

2020). Enrichment for polygenetic traits and disease was investigated using MAGMA competitive gene

set analysis. In the first stage zebrafish – human orthologues of all protein coding genes were mapped.

In the second stage, the strength association between each human protein coding gene and eBMD

and height and any form of self-reported or hospital defined OA was evaluated separately using the

weighted average of 3 different gene-based tests of association. Gene based tests of association were

performed using high quality genome-wide imputed genetic data (~12 million SNPs, INFO > 0.9, MAF

> 0.05%) from 378,484 white European adults from the UK-Biobank Study with eBMD and height

measures. For the analysis involving OA, publically available genome-wide association summary

results statistics were used instead of genome-wide imputed genotyping data (Tachmazidou,

Hatzikotoulas et al., 2019). In the final stage, competitive gene set analysis was used to compare the

mean strength of association of human orthologues of zebrafish DEGs with each trait / disease, to the

mean strength of association all other genes, and evidence of enrichment was quantified using a one-

sided test of statistical significance.

Sensitivity analysis was performed correcting for the set of human genes that could not be mapped

between human and zebrafish, and the set of human orthologues that was not expressed in zebrafish.

Finally, post hoc permutation analysis was performed for analyses that were suggestive of enrichment.

A detailed description of the methods is presented in the **Supplemental Methods**.

Quantitative real-time PCR

500ng of total RNA was treated with DNase (1 unit) and reverse-transcribed (random hexamer

primers) with SuperScript II (Invitrogen 100 units). iQ SYBR Green Supermix (Biorad) containing 350

nM primer and cDNA was used for amplification (primers and PCR conditions in **Supplemental**

Methods). Relative expression was calculated based on a normalisation index of two reference

genes: *eef1a1l1* and *rpl13*. For comparison of qRT-PCR and RNA-seq expression of amplicons, the

average of ontogenetic normalised expression (qRT-PCR) or read count (RNA-seq) were taken and

every individual value was compared to the average ontogenetic expression (e.g. read count

regenerating scales individual 2 / average read count ontogenetic). The p-values presented in the

figures were derived from a two-tailed t-test (qRT-PCR) and p-adjusted (padj) from the DESEQ2

analysis (RNA-seq).

Micro-Computed Tomography and BMD calculations

MicroCT was performed as previously described (Kague et al., 2019), with BMD calculated as

previously described (Stevenson et al., 2017). Briefly, adult WT sibling, *col11a2^{Y228X}* and *spp1^{P160X}*

mutant zebrafish (1 year old) were fixed, dehydrated to 70% EtOH and scanned at 21 μm voxel size

(scan settings 130 kV, 150 μA, 0.5-s exposure, 3141 projections). Images were reconstructed using

NRecon software (Version 1.7.1.0), with dynamic ranges calibrated against a scan of hydroxyapatite

phantoms (0.25 g.cm^{-3} and 0.75 g.cm^{-3}) scanned with identical parameters. For BMD calculations

Aviso software (Avison2020.2; Thermo Fisher Scientific) was used to isolate pixel greyscale values for

the skull (parietal) and vertebrae (vertebrae 11-13) and calibrated against the greyscale values of

known hydroxyapatite density phantoms (0.25 g.cm^{-3} and 0.75 g.cm^{-3}) to estimate mean BMD values

and their standard deviations. Histomorphological assessment of segmented jaw and vertebrae (11-

13) elements were determined by measuring the width, length, and depth of the elements (N.B. left

and right side of jaw elements were considered as independent measurements) in AVIZO between

element landmarks as shown in **figure S7A** and **7B**.

Statistical testing

All graphs and data of downstream RNA-seq functional data were analysed in GraphPad Prism (v

8.4.3) and were subjected to variance comparisons (F test) to determine distribution of the data

prior to performing an unpaired t-test comparing ontogenetic against regenerating or wildtype

against mutant conditions. P-values in figures as follows: n.s. >0.05, * ≤0.05, ** ≤0.01, *** ≤0.001.

RNA-seq, GO, STRING pathway, and GSEA statistical analyses are described in their relevant sections.

Acknowledgements
We would like to thank our funding sources: D.B. EK and C.H. received Fellowship funding from
Versus Arthritis (22044 DB and 21937 CH and EK). J.M. and G.F. were supported by the subsidy
programme Smartmix (SSM06010) of the Dutch Ministries of Economic Affairs and Education,
Culture and, Science. J.P.K was funded by a University of Queensland (UQ) Development Fellowship
(UQFEL1718945), a National Health and Medical Research Council (Australia) Investigator grant
(GNT1177938) and project grant (GNT1158758). M.L. is supported by a UQ Research Training
Scholarship and the Commonwealth Scientific and Industrial Research Organisation Postgraduate
Top-Up Scholarship. R.R. and R.J.R. were supported by the BHF Oxbridge Centre of Regenerative
Medicine (RM/17/2/33380) and a BHF Intermediate Fellowship to R.J.R. (FS/15/2/31225). This
research has been conducted using the UK Biobank Resource (accession ID: 53641). Lastly, we would
like to thank Mathew Green and Pablo Peon Garcia for fish care at UoB, and Tom Spanings and
Antoon van der Horst for fish care at RU.

Author contributions

D.B., and J.M. conceptualised and initiated the project. D.B., Q.T., J.P.K., CH., and J.M. designed the
experiments. D.B., Q.T., A.S, J.Z., E.N., J.P.K., M.L., S.Y. E.K, C.H., and J.M. performed the experiments
and analysed data. E.Z, S.Y., J.P.K, R.R. and R.J.R. provided reagents or access to datasets. D.B., J.P.K.,
C.H., and J.M. supervised the project. D.B., G.F. J.P.K., CH., and J.M. acquired funding for the project.
D.B. prepared manuscript figures. J.P.K, E.Z., S.Y, P.I.C., R.R., G.F. and R.J.R. provided expertise. D.B.,
J.P.K., C.H., and J.M. co-wrote the first version of the manuscript. All authors revised the first version
and approved the contents of the final version of the manuscript.

Conflict of interest and ethics statement

566 All authors declare no conflict of interest. Human genetic and animal research that is presented in
567 this work has been conducted with ethical approval as described in Material and Methods section.

Data availability

RNA-sequencing FASTQ files are available in ENA (https://www.ebi.ac.uk/ena) under accession
number PRJEB39971.

Tables and Figures

Table 1: Top-40 smallest false discovery rate (FDR) differentially regulated transcripts (up (green)
and down (magenta) regulated in Log2 fold change) in regenerated scales (vs ontogenetic)
showing mean and standard deviation (Stdev) for each group (normalised read count).

Ensembl ID	Gene symbol	Log$_2$ Fold Change	FDR	Gene description
ENSDARG00000068621	*si:ch211-181d7.3*	*-6.767*	2.41E-98	NOD-like receptor (NLR) with NACHT and LRR domain
ENSDARG00000074132	*scpp7*	3.756	2.83E-70	secretory calcium-binding phosphoprotein 7
ENSDARG00000093659	*entpd5b*	3.560	2.36E-48	ectonucleoside triphosphate diphosphohydrolase 5b
ENSDARG00000025089	*loxl4*	3.167	1.41E-47	lysyl oxidase-like 4
ENSDARG00000054753	*col10a1a*	3.610	3.10E-43	collagen, type X, alpha 1a
ENSDARG00000052633	si:ch211-106n13.3	4.676	7.84E-39	si:ch211-106n13.3
ENSDARG00000039999	*slc13a5b*	4.351	1.54E-37	solute carrier family 13 member 5b
ENSDARG00000012422	*col11a2*	3.484	6.80E-37	collagen, type XI, alpha 2
ENSDARG00000069216	*c6ast1*	4.275	1.02E-33	six-cysteine containing astacin protease 1
ENSDARG00000098226	*smpd3*	3.226	1.22E-32	sphingomyelin phosphodiesterase 3
ENSDARG00000004627	*panx3*	2.897	1.27E-27	pannexin 3
ENSDARG00000104267	*postnb*	2.775	1.18E-26	periostin, osteoblast specific factor b
ENSDARG00000024847	*col5a2b*	2.783	2.75E-26	collagen, type V, alpha 2b
ENSDARG00000009014	*col11a1b*	3.298	2.77E-26	collagen, type XI, alpha 1b
ENSDARG00000020007	*col1a2*	2.951	5.95E-26	collagen, type I, alpha 2
ENSDARG00000105153	*ifitm5*	3.133	1.29E-25	interferon induced transmembrane protein 5
ENSDARG00000042186	*paplnb*	3.405	7.38E-25	papilin b, proteoglycan-like sulfated glycoprotein
ENSDARG00000095259	*tmem119a*	2.622	6.06E-24	transmembrane protein 119a
ENSDARG00000088274	si:ch211-181d7.1	-4.630	6.06E-24	si:ch211-181d7.1
ENSDARG00000059746	*plod1a*	2.209	7.53E-24	procollagen-lysine, 2-oxoglutarate 5-dioxygenase 1a
ENSDARG00000115341	*col11a1a*	3.421	9.48E-24	collagen, type XI, alpha 1a
ENSDARG00000035809	*col1a1b*	2.926	1.34E-23	collagen, type I, alpha 1b
ENSDARG00000091667	si:dkey-234i14.3	3.501	3.99E-23	si:dkey-234i14.3
ENSDARG00000079572	*plcd3b*	3.515	1.50E-22	phospholipase C, delta 3b
ENSDARG00000058270	si:ch211-114l13.7	2.465	1.79E-22	si:ch211-114l13.7
ENSDARG00000055175	*c1qtnf6a*	3.482	2.61E-22	C1q and TNF related 6a
ENSDARG00000037145	*slc8a4b*	3.076	4.38E-21	solute carrier family 8 member 4b
ENSDARG00000101535	*col10a1b*	3.032	6.06E-21	collagen, type X, alpha 1b
ENSDARG00000078866	*sdk1a*	1.876	8.41E-21	sidekick cell adhesion molecule 1a
ENSDARG00000041703	*rrbp1b*	2.696	1.33E-20	ribosome binding protein 1b
ENSDARG00000040985	*itgbl1*	3.531	9.38E-20	integrin, beta-like 1
ENSDARG00000019949	*serpinh1b*	3.135	1.18E-19	serpin peptidase inhibitor, clade H (heat shock protein 47), member 1b
ENSDARG00000026348	*csad*	2.915	3.69E-19	cysteine sulfinic acid decarboxylase
ENSDARG00000012593	*col5a1*	2.447	5.67E-19	procollagen, type V, alpha 1
ENSDARG00000022437	*cd81b*	2.749	5.85E-19	CD81 molecule b
ENSDARG00000021265	*mybpc2b*	6.520	1.04E-18	myosin binding protein C, fast type b
ENSDARG00000009418	*mef2cb*	2.062	1.19E-18	myocyte enhancer factor 2cb
ENSDARG00000017884	*bgna*	2.896	3.32E-18	biglycan a
ENSDARG00000062363	*phex*	2.034	5.54E-18	phosphate regulating endopeptidase homolog, X-linked
ENSDARG00000079486	*fam20a*	2.263	1.12E-17	FAM20A golgi associated secretory pathway pseudokinase

577 **Table 2: MAGMA competitive gene set enrichment analysis with p-value of >0.05 in green.**

Model	Trait/Disease	#Genes	BETA	BETA STD	SE	P-value
ALL DEGs						
Base	**eBMD**	459	0.20	0.03	0.079	4.76E-03
	Height	459	0.35	0.05	0.102	3.18E-04
	Osteoarthritis	459	0.13	0.02	0.047	3.02E-03
Sensitivity	**eBMD**	459	0.14	0.02	0.080	4.02E-02
	Height	459	0.24	0.04	0.103	9.96E-03
	Osteoarthritis	459	0.12	0.02	0.048	6.91E-03
Upregulated DEGs						
Base	**eBMD**	402	0.27	0.04	0.084	5.61E-04
	Height	402	0.39	0.06	0.109	1.45E-04
	Osteoarthritis	402	0.14	0.02	0.050	3.26E-03
Sensitivity	**eBMD**	402	0.21	0.03	0.085	6.57E-03
	Height	402	0.29	0.04	0.110	4.56E-03
	Osteoarthritis	402	0.12	0.02	0.051	7.05E-03
Downregulated DEGs						
Base	**eBMD**	57	-0.31	-0.02	0.227	9.13E-01
	Height	57	0.01	0.00	0.292	4.86E-01
	Osteoarthritis	57	0.07	0.00	0.131	2.89E-01
Sensitivity	**eBMD**	57	-0.38	-0.02	0.228	9.53E-01
	Height	57	-0.10	-0.01	0.292	6.38E-01
	Osteoarthritis	57	0.06	0.00	0.131	3.29E-01
Key: described in supplemental data file S7						

Bergen et al _ Figure 1

Figure 1: RNA-sequencing transcriptomics of ontogenetic and 9 days regenerated scales. A) *In toto* fluorescent stereomicroscope images of flanks from *sp7:GFP* transgenic fish showing ontogenetic (O) and regenerated scales (R) in an anterior (left) to posterior (right) direction). Note the auto-fluorescent pigment stripe (P) and the compass orientation is used to depict *in toto* scales in this paper with the distal (posterior) edge pointing caudally. **B)** *In situ* alkaline phosphatase staining of a pre-collection and 8 dph scale with insets (i) showing anterior part of the scale. Blue arrows indicate ALP+ cells which appear to be larger and more extended in ontogenetic than regenerating scales. Compass orientation is used to depict *in situ* scales in this paper as the anterior region was closest to the scale dermal socket (pre-harvest). **C)** Schematic drawing of the RNA-sequencing approach. **D)** Volcano plot showing relative fold change ($\log_2$) and false discovery rate ($-\log_{10}$ converted) of coding RNA sequences expressed in both ontogenetic and regenerating scales. Green (≥±1.25 fold change, ≥ 1.3 FDR), magenta (≥±1.25 fold change, <1.3 FDR), blue (<±1.25 fold change, ≥1.3 FDR), and black (not passing any threshold) coloured dots mark the different criteria.

Bergen et al _ Figure 2

Figure 2: Validation and gene ontology of RNA-sequencing dataset. A) Comparison of relative expression levels of quantitative real-time PCR (qRT-PCR) and transcriptomic analysis of selected amplicons. **B)** Representative stereomicroscope images of live calcein green staining labelling newly deposited calcium phosphate of ontogenetic and regenerating scales (n = 4 fish each condition). Blue arrow indicates elevated levels of calcein green labelling compared to surrounding signal whereas orange arrows indicate small puncta of enhanced signal. Insets show lateral circuli (i) and posterior epidermal (containing pigmentating melanocytes (M) region (ii). **C)** GOrilla Gene ontology (GO) analysis showing high enrichment (>5 fold) of collagen and ECM related terms. **D)** GOrilla cellular component hierarchical clustering. Scale bar: 100 µm.

Bergen et al _ Figure 3

Figure 3: STRING network analysis of DEGs. Only DEG proteins with 1 or more protein-protein interactions (PPI) (indicated as edges) were shown.

Figure 4: Validation of gene ontology findings. A) Relative expression levels of qRT-PCR and
transcriptomic analysis of amplicons found in the collagen, proteoglycan and hedgehog signalling
networks. **B)** Images of *in toto* and harvested scales (*in situ*) of *col1a1a:GFP* and *sp7:mCherry-NTR*
double transgenic fish (n = 4 each condition). Blue arrow indicates the posterior (distal) fringe with
high expression of mCherry. Orange arrow points at co-expression of GFP and mCherry. **C)** *In toto*
and *in situ* stereomicroscope images of *col10a1a:Citrine* and *col2a1a:mCherry* double transgenic
ontogenetic and regenerating scales (n = 4 fish each condition). Orange arrow indicate Citrine signal

at the posterior distal edge (inset i) while blue arrow points at a newly forming lateral circulus (inset
ii) of the scale. Scale bar: 100 μm.

Bergen et al _ Figure 5

Figure 5: Gene set enrichment analysis strategy showing gene set enrichment of zebrafish DEGs in monogenic skeletal disorders and complex musculoskeletal traits and disease. A) Flow diagram summarising gene set enrichment analysis approach showing examples of enriched monogenic nosology groups and polygenic traits using human orthologues of zebrafish DEGs. **B)** Graph showing p-value of monogenic skeletal dysplasia nosology groups found in this study. The fold enrichment and human disease genes and zebrafish DEGs intersect are listed as well. **C)** Venn diagram of gene-based tests of association post-hoc analysis output showing number of associated DEGs with a stringent gene-wide significance threshold of $P<2.63\times10^{-6}$. The diagrams show the unique or intersecting genes between estimated bone mineral density (eBMD), osteoarthritis (OA), and height UK-Biobank gene based tests of association. The boxes show selected examples of DEG in each category and bold gene symbols are validated in this study.

Bergen et al _ Figure 6

Figure 6: Validation of genes associated with polygenic traits in humans. A) Q-RTPCR and RNA-seq
expression levels (relative to average ontogenetic expression) of selected DEG found in each
category. All but one (*tmem97*) reached statistical threshold of p<0.05. **B)** Images of scales from *in
toto* and harvested scales of *sp7:GFP* and *spp1:mCherry* double transgenic fish (ontogenetic: n = 5,
regenerating: n = 4 fish). For overview and inset i, blue arrows indicate high GFP signal and orange
the mCherry signal at the posterior edge (inset i) while light blue arrow indicates an ontogenetic
scale next to a regenerating scale. Inset ii shows anterior region of the scale with opposite GFP and
mCherry signals (blue arrow) in ontogenetic and regenerating conditions. **C)** Quantification of width
of the mCherry signal at the posterior edge from harvested scales (between blue and orange arrows
in panel C). **D)** Width of mCherry signal normalised by scale length (anterior to posterior). Scale bar:
100 µm.

Figure 7: Histomorphology and bone mineral density measurements on 3D micro-CT images of *col11a2*[Y228X] and *spp1*[P160X] homozygous fish revealed altered bone structures. A) Axial skeleton length was reduced in *col11a2* mutants whereas *spp1* mutants showed a mild reduced tendency. B) Quantification of lower jaw size. B) Calculation of lower jaw element proportions. D) Ventral view of the segmented lower jaw images. Orange arrow indicates anterior mandibular arch joint and blue arrow shows mandibular arch – palatoquadrate (not visible) joint. E) Quantification of the vertebral arches' length. F) Total length of the anterior facet of the vertebral centrum. G) Lateral view of segmented images of the first caudal vertebra showing the anterior facet (orange arrow) and haemal arch (blue arrow). H) Mean BMD calculations of dermal and notochord sheath derived bone.

References

Alestrom P, D'Angelo L, Midtlyng PJ, Schorderet DF, Schulte-Merker S, Sohm F, Warner S (2020) Zebrafish: Housing and husbandry recommendations. *Lab Anim* 54: 213-224

Aman AJ, Fulbright AN, Parichy DM (2018) Wnt/β-catenin regulates an ancient signaling network during zebrafish scale development. *eLife* 7: e37001

Balemans W, Ebeling M, Patel N, Van Hul E, Olson P, Dioszegi M, Lacza C, Wuyts W, Van Den Ende J, Willems P, Paes-Alves AF, Hill S, Bueno M, Ramos FJ, Tacconi P, Dikkers FG, Stratakis C, Lindpaintner K, Vickery B, Foernzler D et al. (2001) Increased bone density in sclerosteosis is due to the deficiency of a novel secreted protein (SOST). *Human Molecular Genetics* 10: 537-544

Bereiter-Hahn J, Zylberberg L (1993) Regeneration of teleost fish scale. *Comparative Biochemistry and Physiology Part A: Physiology* 105: 625-641

Bergen DJM, Kague E, Hammond CL (2019) Zebrafish as an Emerging Model for Osteoporosis: A Primary Testing Platform for Screening New Osteo-Active Compounds. *Front Endocrinol (Lausanne)* 10: 6

Bevan L, Lim ZW, Venkatesh B, Riley PR, Martin P, Richardson RJ (2020) Specific macrophage populations promote both cardiac scar deposition and subsequent resolution in adult zebrafish. *Cardiovasc Res* 116: 1357-1371

Bhattacharya S, Hyland C, Falk MM, Iovine MK (2020) Connexin 43 gap junctional intercellular communication inhibits evx1 expression and joint formation in regenerating fins. *Development* 147: dev190512

Bigi A, Burghammer M, Falconi R, Koch MHJ, Panzavolta S, Riekel C (2001) Twisted Plywood Pattern of Collagen Fibrils in Teleost Scales: An X-ray Diffraction Investigation. *Journal of Structural Biology* 136: 137-143

Boskey AL, Spevak L, Paschalis E, Doty SB, McKee MD (2002) Osteopontin Deficiency Increases Mineral Content and Mineral Crystallinity in Mouse Bone. *Calcified Tissue International* 71: 145-154

Carnovali M, Luzi L, Banfi G, Mariotti M (2016) Chronic hyperglycemia affects bone metabolism in adult zebrafish scale model. *Endocrine* 54: 808-817

Carnovali M, Ottria R, Pasqualetti S, Banfi G, Ciuffreda P, Mariotti M (2016) Effects of bioactive fatty acid amide derivatives in zebrafish scale model of bone metabolism and disease. *Pharmacological Research* 104: 1-8

Chen X-D, Fisher LW, Robey PG, Young MF (2004) The small leucine-rich proteoglycan biglycan modulates BMP-4-induced osteoblast differentiation. *The FASEB Journal* 18: 948-958

Cosman F, Crittenden DB, Adachi JD, Binkley N, Czerwinski E, Ferrari S, Hofbauer LC, Lau E, Lewiecki EM, Miyauchi A, Zerbini CAF, Milmont CE, Chen L, Maddox J, Meisner PD, Libanati C, Grauer A (2016) Romosozumab Treatment in Postmenopausal Women with Osteoporosis. *New England Journal of Medicine* 375: 1532-1543

Couchouron T, Masson C (2011) Early-onset progressive osteoarthritis with hereditary progressive ophtalmopathy or Stickler syndrome. *Joint Bone Spine* 78: 45-49

Cox BD, De Simone A, Tornini VA, Singh SP, Di Talia S, Poss KD (2018) In Toto Imaging of Dynamic Osteoblast Behaviors in Regenerating Skeletal Bone. *Current Biology* 28: 3937-3947.e4

Dai J, Peng L, Fan K, Wang H, Wei R, Ji G, Cai J, Lu B, Li B, Zhang D, Kang Y, Tan M, Qian W, Guo Y (2009) Osteopontin induces angiogenesis through activation of PI3K/AKT and ERK1/2 in endothelial cells. *Oncogene* 28: 3412-3422

de Vrieze E, Sharif F, Metz JR, Flik G, Richardson MK (2011) Matrix metalloproteinases in osteoclasts of ontogenetic and regenerating zebrafish scales. *Bone* 48: 704-712

de Vrieze E, van Kessel MA, Peters HM, Spanings FA, Flik G, Metz JR (2014) Prednisolone induces osteoporosis-like phenotype in regenerating zebrafish scales. *Osteoporos Int* 25: 567-78

de Vrieze E, Zethof J, Schulte-Merker S, Flik G, Metz JR (2015) Identification of novel osteogenic compounds by an ex-vivo sp7:luciferase zebrafish scale assay. *Bone* 74: 106-113

Debiais-Thibaud M, Simion P, Ventéo S, Muñoz D, Marcellini S, Mazan S, Haitina T (2019) Skeletal
Mineralization in Association with Type X Collagen Expression Is an Ancestral Feature for Jawed
Vertebrates. *Molecular Biology and Evolution* 36: 2265-2276
Dhouailly D, Godefroit P, Martin T, Nonchev S, Caraguel F, Oftedal O (2019) Getting to the root of
scales, feather and hair: As deep as odontodes? *Experimental Dermatology* 28: 503-508
Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013)
STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29: 15-21
Douglas T, Heinemann S, Bierbaum S, Scharnweber D, Worch H (2006) Fibrillogenesis of collagen
types I, II, and III with small leucine-rich proteoglycans decorin and biglycan. *Biomacromolecules* 7:
2388-93
Eames BF, Amores A, Yan Y-L, Postlethwait JH (2012) Evolution of the osteoblast: skeletogenesis in
gar and zebrafish. *BMC Evolutionary Biology* 12: 27
Eden E, Navon R, Steinfeld I, Lipson D, Yakhini Z (2009) GOrilla: a tool for discovery and visualization
of enriched GO terms in ranked gene lists. *BMC Bioinformatics* 10: 48-48
Giraud-Guille MM (1988) Twisted plywood architecture of collagen fibrils in human compact bone
osteons. *Calcified Tissue International* 42: 167-180
Gregory KE, Oxford JT, Chen Y, Gambee JE, Gygi SP, Aebersold R, Neame PJ, Mechling DE, Bächinger
HP, Morris NP (2000) Structural Organization of Distinct Domains within the Non-collagenous N-
terminal Region of Collagen Type XI. *Journal of Biological Chemistry* 275: 11498-11506
Guellec DL, Zylberberg L (1998) Expression of Type I and Type V Collagen mRNAs in the Elasmoid
Scales of a Teleost Fish as Revealed by In Situ Hybridization. *Connective Tissue Research* 39: 257-267
Hammond CL, Schulte-Merker S (2009) Two populations of endochondral osteoblasts with
differential sensitivity to Hedgehog signalling. *Development* 136: 3991-4000
Holm E, Gleberzon Jared S, Liao Y, Sørensen Esben S, Beier F, Hunter Graeme K, Goldberg Harvey A
(2014) Osteopontin mediates mineralization and not osteogenic cell development in vitro.
Biochemical Journal 464: 355-364
Iwasaki M, Kuroda J, Kawakami K, Wada H (2018) Epidermal regulation of bone morphogenesis
through the development and regeneration of osteoblasts in the zebrafish scale. *Developmental
Biology* 437: 105-119
Kenkre JS, Bassett JHD (2018) The bone remodelling cycle. *Annals of Clinical Biochemistry* 55: 308-
327
Kim H-J, Lee M-H, Park H-S, Park M-H, Lee S-W, Kim S-Y, Choi J-Y, Shin H-I, Kim H-J, Ryoo H-M (2003)
Erk pathway and activator protein 1 play crucial roles in FGF2-stimulated premature cranial suture
closure. *Developmental Dynamics* 227: 335-346
Kobayashi-Sun J, Yamamori S, Kondo M, Kuroda J, Ikegame M, Suzuki N, Kitamura KI, Hattori A,
Yamaguchi M, Kobayashi I (2020) Uptake of osteoblast-derived extracellular vesicles promotes the
differentiation of osteoclasts in the zebrafish scale. *Commun Biol* 3: 190
Lamort A-S, Giopanou I, Psallidas I, Stathopoulos GT (2019) Osteopontin as a Link between
Inflammation and Cancer: The Thorax in the Spotlight. *Cells* 8: 815
Lawrence EA, Kague E, Aggleton JA, Harniman RL, Roddy KA, Hammond CL (2018) The mechanical
impact of col11a2 loss on joints; col11a2 mutant zebrafish show changes to joint development and
function, which leads to early-onset osteoarthritis. *Philosophical Transactions of the Royal Society B:
Biological Sciences* 373: 20170335
Li X, Ominsky MS, Niu Q-T, Sun N, Daugherty B, D'Agostin D, Kurahara C, Gao Y, Cao J, Gong J,
Asuncion F, Barrero M, Warmington K, Dwyer D, Stolina M, Morony S, Sarosi I, Kostenuik PJ, Lacey
DL, Simonet WS et al. (2008) Targeted Deletion of the Sclerostin Gene in Mice Results in Increased
Bone Formation and Bone Strength. *Journal of Bone and Mineral Research* 23: 860-869
Li Y, Lacerda DA, Warman ML, Beier DR, Yoshioka H, Ninomiya Y, Oxford JT, Morris NP,
Andrikopoulos K, Ramirez F, Wardell BB, Lifferth GD, Teuscher C, Woodward SR, Taylor BA,
Seegmiller RE, Olsen BR (1995) A fibrillar collagen gene, Col11a1, is essential for skeletal
morphogenesis. *Cell* 80: 423-430

Lleras-Forero L, Winkler C, Schulte-Merker S (2020) Zebrafish and medaka as models for biomedical
research of bone diseases. *Developmental Biology* 457: 191-205

Long F, Ornitz DM (2013) Development of the endochondral skeleton. *Cold Spring Harbor*
perspectives in biology 5: a008334

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biology* 15: 550

Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD (2019) PANTHER version 14: more genomes, a
new PANTHER GO-slim and improvements in enrichment analysis tools. *Nucleic Acids Research* 47:
D419-D426

Mori R, Shaw TJ, Martin P (2008) Molecular mechanisms linking wound inflammation and fibrosis:
knockdown of osteopontin leads to rapid repair and reduced scarring. *J Exp Med* 205: 43-51

Morinobu M, Ishijima M, Rittling SR, Tsuji K, Yamamoto H, Nifuji A, Denhardt DT, Noda M (2003)
Osteopontin Expression in Osteoblasts and Osteocytes During Bone Formation Under Mechanical
Stress in the Calvarial Suture In Vivo. *Journal of Bone and Mineral Research* 18: 1706-1715

Mortier GR, Cohn DH, Cormier-Daire V, Hall C, Krakow D, Mundlos S, Nishimura G, Robertson S,
Sangiorgi L, Savarirayan R, Sillence D, Superti-Furga A, Unger S, Warman ML (2019) Nosology and
classification of genetic skeletal disorders: 2019 revision. *American Journal of Medical Genetics Part*
A 179: 2393-2419

Mosaffa P, Tetley RJ, Rodríguez-Ferran A, Mao Y, Muñoz JJ (2020) Junctional and cytoplasmic
contributions in wound healing. *Journal of The Royal Society Interface* 17: 20200264

Padhi BK, Joly L, Tellis P, Smith A, Nanjappa P, Chevrette M, Ekker M, Akimenko M-A (2004) Screen
for genes differentially expressed during regeneration of the zebrafish caudal fin. *Developmental*
Dynamics 231: 527-541

Page-McCaw A, Ewald AJ, Werb Z (2007) Matrix metalloproteinases and the regulation of tissue
remodelling. *Nature Reviews Molecular Cell Biology* 8: 221-233

Pasqualetti S, Banfi G, Mariotti M (2012) The zebrafish scale as model to study the bone
mineralization process. *Journal of Molecular Histology* 43: 589-595

Pasqualetti S, Congiu T, Banfi G, Mariotti M (2015) Alendronate rescued osteoporotic phenotype in a
model of glucocorticoid-induced osteoporosis in adult zebrafish scale. *Int J Exp Pathol* 96: 11-20

Pathak JL, Bravenboer N, Klein-Nulend J (2020) The Osteocyte as the New Discovery of Therapeutic
Options in Rare Bone Diseases. *Frontiers in endocrinology* 11: 405-405

Ricard-Blum S (2011) The collagen family. *Cold Spring Harbor perspectives in biology* 3: a004978

Richardson R, Slanchev K, Kraus C, Knyphausen P, Eming S, Hammerschmidt M (2013) Adult
Zebrafish as a Model System for Cutaneous Wound-Healing Research. *Journal of Investigative*
Dermatology 133: 1655-1665

Schmidt JR, Geurtzen K, von Bergen M, Schubert K, Knopf F (2019) Glucocorticoid Treatment Leads
to Aberrant Ion and Macromolecular Transport in Regenerating Zebrafish Fins. *Frontiers in*
Endocrinology 10

Sehring IM, Weidinger G (2020) Recent advancements in understanding fin regeneration in
zebrafish. *WIREs Developmental Biology* 9: e367

Shimada A, Kawanishi T, Kaneko T, Yoshihara H, Yano T, Inohaya K, Kinoshita M, Kamei Y, Tamura K,
Takeda H (2013) Trunk exoskeleton in teleosts is mesodermal in origin. *Nature Communications* 4:
1639

Sire JY, Akimenko MA (2004) Scale development in fish: a review, with description of sonic hedgehog
(shh) expression in the zebrafish (Danio rerio). *Int J Dev Biol* 48: 233-47

Sire JY, Allizard F, Babiar O, Bourguignon J, Quilhac A (1997) Scale development in zebrafish (Danio
rerio). *Journal of anatomy* 190 (Pt 4): 545-561

Sire JY, Donoghue PC, Vickaryous MK (2009) Origin and evolution of the integumentary skeleton in
non-tetrapod vertebrates. *J Anat* 214: 409-40

Sousa S, Valerio F, Jacinto A (2012) A new zebrafish bone crush injury model. *Biology open* 1: 915-21

Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, Simonovic M, Doncheva NT,
Morris JH, Bork P, Jensen LJ, Mering Cv (2019) STRING v11: protein-protein association networks
with increased coverage, supporting functional discovery in genome-wide experimental datasets.
Nucleic acids research 47: D607-D613

Tachmazidou I, Hatzikotoulas K, Southam L, Esparza-Gordillo J, Haberland V, Zheng J, Johnson T,
Koprulu M, Zengini E, Steinberg J, Wilkinson JM, Bhatnagar S, Hoffman JD, Buchan N, Süveges D, arc
OC, Yerges-Armstrong L, Smith GD, Gaunt TR, Scott RA et al. (2019) Identification of new therapeutic
targets for osteoarthritis through genome-wide analyses of UK Biobank data. *Nature genetics* 51:
230-236

Terai K, Takano-Yamamoto T, Ohba Y, Hiura K, Sugimoto M, Sato M, Kawahata H, Inaguma N,
Kitamura Y, Nomura S (1999) Role of Osteopontin in Bone Remodeling Caused by Mechanical Stress.
Journal of Bone and Mineral Research 14: 839-849

Topczewska JM, Shoela RA, Tomaszewski JP, Mirmira RB, Gosain AK (2016) The Morphogenesis of
Cranial Sutures in Zebrafish. *PLoS One* 11: e0165775-e0165775

Wu XM, Chen WQ, Hu YW, Cao L, Nie P, Chang MX (2018) RIP2 Is a Critical Regulator for NLRs
Signaling and MHC Antigen Presentation but Not for MAPK and PI3K/Akt Pathways. *Frontiers in*
Immunology 9

Xu T, Bianco P, Fisher LW, Longenecker G, Smith E, Goldstein S, Bonadio J, Boskey A, Heegaard A-M,
Sommer B, Satomura K, Dominguez P, Zhao C, Kulkarni AB, Robey PG, Young MF (1998) Targeted
disruption of the biglycan gene leads to an osteoporosis-like phenotype in mice. *Nature Genetics* 20:
78-82

Yadav MC, Huesa C, Narisawa S, Hoylaerts MF, Moreau A, Farquharson C, Millán JL (2014) Ablation
of osteopontin improves the skeletal phenotype of phospho1(-/-) mice. *J Bone Miner Res* 29: 2369-
2381

Yasuo M (1980) The source of calcium in regenerating scales of the goldfish, Carassius auratus.
Comparative Biochemistry and Physiology Part A: Physiology 66: 521-524

Youlten SE, Kemp JP, Logan JG, Ghirardello EJ, Sergio CM, Dack MRG, Guilfoyle SE, Leitch VD,
Butterfield NC, Komla-Ebri D, Chai RC, Corr AP, Smith JT, Morris JA, McDonald MM, Quinn JMW,
McGlade AR, Bartonicek N, Jansson M, Hatzikotoulas K et al. (2020) Osteocyte Transcriptome
Mapping Identifies a Molecular Landscape Controlling Skeletal Homeostasis and Susceptibility to
Skeletal Disease. *bioRxiv*: 2020.04.20.051409

Zeltz C, Gullberg D (2016) The integrin–collagen connection – a glue for tissue repair? *Journal of Cell*
Science 129: 653

Zhao A, Qin H, Fu X (2016) What Determines the Regenerative Capacity in Animals? *BioScience* 66:
735-746