Rapid Cortical Adaptation and the Role of Thalamic Synchrony During Wakefulness

Nathaniel C. Wright[1], Peter Y. Borden[1], Yi Juin Liew[2], Michael F. Bolus[1], William M. Stoy[3], Craig R. Forest[3], Garrett B. Stanley[1]

[1]Wallace H Coulter Department of Biomedical Engineering, Georgia Institute of Technology and Emory University, Atlanta, GA, USA

[2]Department of Biomedical Engineering, Georgia Institute of Technology, Emory University, Atlanta, GA, USA and Beijing University, Beijing China

[3]Department of Mechanical Engineering, Georgia Institute of Technology, Atlanta, Georgia, USA

Abbreviated Title: Cortical adaptation during wakefulness

Pages/Figures: 48 text pages, 8 Figures

Word counts: Abstract: 248, Introduction: 650, Discussion: 1482

Correspondence: Garrett B Stanley

Coulter Department of Biomedical Engineering
Georgia Institute of Technology & Emory University
313 Ferst Drive
Atlanta GA 30332-0535
USA

Phone: 404-385-5037
Fax: 404-385-5044
Email: garrett.stanley@bme.gatech.edu

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgements: We thank Aurelie Pala, Audrey Sederberg, Adam Willats, Christian Waiblinger, and Elaida Dimwamwa for helpful comments on the data analysis and manuscript. We thank Aurelie Pala for assistance with spike-sorting, and Mahmood S. Hoseini for his assistance with the network model. **Funding**. This work was supported by NIH National Institute of Neurological Disorders and Stroke BRAIN Grant R01NS104928 (GBS), NIH National Institute of Mental Health BRAIN Grant U01MH106027 (GBS and CRF), NIH National Institute of Neurological Disorders and Stroke Pre-doctoral NRSA NS098691 (PYB), NIH National Eye Institute R01EY023173 (CRF), and National Science Foundation Graduate Research Fellowships (WMS, MFB).

Abstract

Rapid sensory adaptation is observed across all sensory systems, and strongly shapes sensory percepts in complex sensory environments. Yet despite its ubiquity and likely necessity for survival, the mechanistic basis is poorly understood. A wide range of primarily in-vitro and anesthetized studies have demonstrated the emergence of adaptation at the level of primary sensory cortex, with only modest signatures in earlier stages of processing. The nature of rapid adaptation and how it shapes sensory representations during wakefulness, and thus the potential role in perceptual adaptation, is underexplored, as are the mechanisms that underlie this phenomenon. To address these knowledge gaps, we recorded spiking activity in primary somatosensory cortex (S1) and the upstream ventral posteromedial (VPm) thalamic nucleus in the vibrissa pathway of awake male and female mice, and quantified responses to whisker stimuli delivered in isolation and embedded in an adapting sensory background. We found that cortical sensory responses were indeed adapted by persistent sensory stimulation; putative excitatory neurons were profoundly adapted, and inhibitory neurons only modestly so. Further optogenetic manipulation experiments and network modeling suggest this largely reflects adaptive changes in synchronous thalamic firing combined with robust engagement of feedforward inhibition, with little contribution from synaptic depression. Taken together, these results suggest that cortical adaptation in the regime explored here results from changes in the timing of thalamic input, and the way in which this differentially impacts cortical excitation and feedforward inhibition, pointing to a prominent role of thalamic gating in rapid adaptation of primary sensory cortex.

Significance Statement
Rapid adaptation of sensory activity strongly shapes representations of sensory inputs across all sensory pathways over the timescale of seconds, and has profound effects on sensory perception. Despite its ubiquity and theoretical role in the efficient encoding of complex sensory environments, the mechanistic basis is poorly understood, particularly during wakefulness. In this study in the vibrissa pathway of awake mice, we show that cortical representations of sensory inputs are strongly shaped by rapid adaptation, and that this is mediated primarily by adaptive gating of the thalamic inputs to primary sensory cortex and the differential way in which these inputs engage cortical sub-populations of neurons.

Introduction

Our experience of the world around us depends upon context. For instance, a noisy environment provides persistent sensory stimulation, which can adapt the representations and percepts of salient sensory features embedded within the environment. Rapid sensory adaptation describes such interactions between stimulus history and perception, spanning milliseconds to seconds (Whitmire and Stanley, 2016). A wealth of human psychophysical studies have documented perceptual adaptation in audition (Smith and Faulkner, 2006; Bestelmeyer et al., 2010; Erb et al., 2013), vision (Blakemore and Campbell, 1969; Blakemore and Nachmias, 1971; Anstis et al., 1998; Ghodrati et al., 2019), and somatosensation (Tannan et al., 2007), suggesting rapid sensory adaptation lends a vital flexibility to organisms tasked with surviving and thriving during rapid environmental changes.

Despite the ubiquity of rapid sensory adaptation across sensory pathways, and its likely relevance to function (Barlow, 1961), the neural basis has not been conclusively identified. A large body of (mostly in vitro and anesthetized) work has implicated the early sensory pathway; adaptation effects cascade toward and culminate in primary sensory cortex (Ganmor et al., 2010; Maravall et al., 2013; Lampl and Katz, 2017), and the nature of cortical adaptation is consistent with perceptual effects. In the vibrissa pathway of the anesthetized rodent for example, adaptation induced through whisker stimulation shapes the amplitude (Webber and Stanley, 2006, 2004; Boloori and Stanley, 2006; Maravall et al., 2007; Heiss et al., 2008; Boloori et al., 2010; Ganmor et al., 2010; Wang et al., 2010; Cohen-Kashi Malina et al., 2013; Ollerenshaw et al., 2014; Kheradpezhouh et al., 2017) and spatial extent (Ollerenshaw et al., 2014; Zheng et al., 2015) of responses to subsequent stimuli in primary somatosensory cortex (S1). Yet two important questions remain unanswered. First, what is the nature of cortical response adaptation during wakefulness, where baseline activity is elevated relative to the anesthetized state (Greenberg et al., 2008; Vizuete et al., 2012; Aasebø et al., 2017)? The thalamocortical pathway during wakefulness may be in a baseline "adapted" state that is relatively impervious to additional adaptation (Castro-Alamancos, 2004), and the neural basis for perceptual adaptation lies elsewhere, but this has not been rigorously tested. Second, if the circuit is subject to sensory adaptation during wakefulness, what are the underlying mechanisms? Previous anesthetized and in-vitro work implicates thalamocortical and/or intracortical synaptic depression (Castro-Alamancos and Oldford, 2002; Chung et al., 2002; Gabernet et al., 2005; Cruikshank et al., 2007, 2010; Cohen-Kashi Malina et al., 2013), which could explain why mean evoked rates are generally more adapted in cortex than in thalamus (Chung et al., 2002; Khatri et al., 2004; Gabernet et al., 2005). Yet adaptation also reduces thalamic synchrony (Wang et al., 2010; Ollerenshaw et al., 2014) and single-unit bursting (Whitmire et al., 2016) under anesthesia. Given the sensitivity of

114   cortex to the timing of thalamic inputs (Usrey et al., 2000; Swadlow and Gusev, 2001;
115   Bruno and Sakmann, 2006; Wang et al., 2010; Ollerenshaw et al., 2014), this suggests a
116   potential role for thalamic spike timing in cortical adaptation.
117
Here, we address these unknowns by recording from and modeling S1 of the awake,
head-fixed mouse during rapid sensory adaptation. We found that despite the relatively
high level of baseline activity typical of wakefulness, putative excitatory neurons in S1
were profoundly adapted by background sensory stimulation. In particular, mean evoked
firing rates, theoretical stimulus detectability, and synchronous cortical spiking were all
significantly reduced in the adapted state, consistent with previously-observed changes
in perceptual report (Ollerenshaw et al., 2014; Waiblinger et al., 2015). Several lines of
evidence – including the recording of the thalamic inputs under a range of optogenetic
controls and computational modeling – suggest this primarily reflected reduced
synchronous thalamic firing and robust thalamically-driven feedforward inhibition in the
adapted condition, with little contribution from thalamocortical and intracortical synaptic
depression. Taken together, these results establish the role of the thalamocortical circuit
in rapid adaptation during wakefulness, and implicate a more critical role of thalamic input
than previously thought.
132
133

Materials and Methods

All procedures were approved by the Institutional Animal Care and Use Committee at the Georgia Institute of Technology (Protocol Numbers A100223 and A100225), and were in agreement with guidelines established by the National Institutes of Health.

Surgery

Experiments were carried out using C57BL/6J, Ai32xNR133 (Bolus et al., 2020) (Ai32 crossed with the NR133 Cre-recombinase driver line (Gerfen et al., 2013)), and Ai32xPV-Cre mice of both sexes. All mouse lines were purchased from Jackson Laboratories. Mice were induced with 5% isoflurane in an induction chamber, then transferred to a heating pad on a stereotaxic instrument, and maintained at 1 - 2% isoflurane for the remainder of the surgical procedure. A custom stainless steel headplate was fixed to the exposed skull with Metabond dental cement (Parkell, Inc.), exposed bone and tissue were then sealed with Metabond and either super-glue (Loctite 404; Henkel) or Vetbond Tissue Adhesive (3M). Metabond was used to fashion a well surrounding the left hemisphere. The well was filled with Kwik-Cast (World Precision Instruments, Inc.) and covered with a thin layer of Metabond, and the mouse was returned to its home cage. Mice were given pre- (buprenorphine) and post-operative (ketoprofen) analgesic, and were allowed to recover for three days before additional handling.

Habituation

Three days after headplate implantation, mice were handled for at least 15 minutes, and then returned to their home cage. On subsequent days, mice were gradually habituated to head-fixation on a custom platform consisting of a tunnel with headpost clamps at one end. The first three daily habituation sessions lasted 15, 30, and 45 minutes respectively, but mice were returned to their home cage if they displayed signs of distress. We then gradually extended session durations until mice would tolerate at least 1.5 hours of fixation and whisker stimulation without signs of distress. Mice that did not meet these criteria were removed from the study, or used for anesthetized recordings (see below).

Awake electrophysiological recordings

We recorded from five Ai32xNR133, four C57BL/6J (wild-type) and one Ai32xPV-cre awake mice of both sexes (up to three awake sessions per mouse). We used intrinsic optical signal imaging acquired under anesthesia to identify at least one putative principal column in S1. On the morning of the first recording session for each animal, we

anesthetized the mouse as described above, and opened an approximately 500 micron
x 500 micron craniotomy centered over a putative cortical column. When acquiring
simultaneous VPm and S1 recordings, we opened a second craniotomy of similar size
over the stereotactic coordinates for VPm (1.8 mm lateral from midline by 1.8 mm caudal
from bregma) and slowly inserted either a single-channel tungsten electrode (2 MOhm,
FHC) or 32-channel silicon probe (NeuroNexus A1x32-Poly3-5mm-25s-177) to a depth
of approximately 3 mm. We adjusted the electrode depth while presenting continuous 10
Hz "sawtooth" stimulus trains to individual whiskers until we could identify a putative
principal barreloid (by observing broad-waveform units with robust, short-latency,
minimally-adapting sensory responses to stimulation of a single whisker, and at most
comparatively weak responses to stimulation of surrounding whiskers (Brecht and
Sakmann, 2002)), before slowly retracting the electrode/probe. We then covered
exposed brain tissue with agarose, filled the well with Kwik-Cast, and allowed the mouse
to recover in its home cage for at least two hours. After recovery, we head-fixed the
awake mouse, removed the Kwik-Cast, filled the well with either saline, mineral oil, or
agarose, and inserted an electrode/probe into each open craniotomy using a Luigs and
Neumann manipulator. For S1 recordings, we inserted a multi-channel silicon probe
(NeuroNexus) oriented 35 degrees from vertical. We used either a 32-channel linear
(A1x32-5mm-25-177-A32), 32-channel "Poly3" (A1x32-Poly3-5mm-25s-177), or 64-
channel, four-shank "Poly2" (A4x16-Poly2-5mm-23s-200-177) configuration probe. For
VPm recordings, we inserted either a tungsten optoelectrode (2 Megaohm, FHC, with
attached 200 micron optic fiber, Thorlabs), 32-channel silicon probe (A1x32-Poly3-5mm-
25s-177) or 32-channel silicon optoelectrode (A1x32-Poly3-5mm-25s-177-OA32LP, with
attached 105 micron optic fiber coupled to a 200 micron optic fiber, Thorlabs). Optic fibers
were coupled to a 470 nm LED (M470F3, Thorlabs). When the barreloid we functionally
identified during the anesthetized VPm mapping session was not topographically-aligned
with the targeted S1 column, we referenced the (Coronal) Allen Brain Atlas to adjust the
position of the VPm probe before descending. We inserted the probe(s) slowly to avoid
excessive tissue dimpling, and waited at least 30 minutes after probe insertion to begin
recording, to allow the tissue to settle. Continuous signals were acquired using a either
a Cerebus (Blackrock Microsystems) or Tucker Davis Technologies acquisition system.
Signals were amplified, filtered between 0.3 Hz and 7.5 kHz, and digitized at either 30
kHz or 24414.0625 Hz.

After the first recording session, we removed the probe(s), covered exposed tissue with
agarose, and sealed the well with Kwik-Cast and a thin layer of Metabond. We obtained
either two or three recording sessions (one per day) from each mouse using the original
craniotomy, but each time targeting a different cortical column and barreloid.

Anesthetized extracellular recordings

We recorded from three C57BL/6J (wild-type), four Ai32xPV-Cre, and three Ai32xNR133
mice of both sexes under isoflurane anesthesia. Mice were anesthetized and implanted
with headplates, and we opened a single craniotomy (either approximately 500 microns
x 500 microns, or 1 mm x 1 mm) over S1, as described above. In some cases, the
principal column was first identified using intrinsic optical signal imaging, as described
above. In other cases, we inserted a single tungsten electrode into the stereotactic
coordinates for the center of S1, and defined the putative principal whisker to be that
which evoked the largest LFP response. We then inserted a 4 x 16 silicon probe array
(A4x16-Poly2-5mm-23s-200-177) to a depth of 700 microns. We oriented the probe to
avoid blood vessels on the cortical surface. For a subset of these experiments, we
obtained additional sessions by repeating the stimulation protocol using the whisker that
evoked the maximum LFP response on a shank different from the first. For each such
session, we determined the putative principal column off-line using white-noise-evoked
spiking. For each shank, we summed single- and multi-unit (see below) spiking across
all trials for the 1 s window preceding feature onset. We divided the across-trial mean
white-noise-evoked response by the across-trial standard deviation of spontaneous
spiking, and the shank with the largest resulting value was determined to correspond to
the principal column.

Anesthetized intracellular recordings

We recorded ongoing, sensory-evoked, and light-evoked subthreshold activity from four
sensory- and light-responsive neurons in two mice using an Autopatcher system
(Kodandaramaiah et al., 2016), as described in detail previously (Stoy et al.,
2020). Briefly, we head-plated and identified the putative C2 column using intrinsic optical
signal imaging in two isoflurane-anesthetized Ai32xNR133 transgenic mice, and opened
a 1 mm x 1 mm craniotomy over the column, as described above. We then used an
Autopatcher 1500 (Neuromatic Devices) to provide pressure and measure pipette
resistance, and an algorithm based on these measurements to navigate around blood
vessels in an automated fashion while the pipette descended through cortical tissue.
Finally, we applied a recently-developed automated motion-compensation procedure
(Stoy et al., 2020) for synchronizing the motion of the pipette tip to that of the targeted
neuron prior to forming a seal. These experiments utilized Multiclamp 700B amplifiers
(Molecular Devices), and signals were digitized at 20 kHz (cDAQ-9174, National
Instruments), and were recorded in PClamp 10 in current-clamp mode.

Whisker stimulation

We used a precise, computer-controlled galvanometer (Cambridge Technologies) with attached tube to stimulate individual whiskers (Whitmire et al., 2016; Waiblinger et al., 2018; Sederberg et al., 2019; Liew et al., 2020). The galvanometer was controlled using either a custom Matlab GUI and Simulink Real-Time (MathWorks), or the Real-time eXperiment Interface application (http://rtxi.org/), sampling at 1 kHz. We inserted the whisker into the tube, which was positioned approximately 10 mm from the whisker pad. We delivered "sawtooth" stimulus features (exponential rise and decay waveforms lasting approximately 17 ms, with reported velocity defined by the average over the 8.5 ms rising phase (Wang et al., 2010)) either in isolation, or embedded in frozen sensory white noise (i.e., white noise waveforms that were identical across trials). To generate the white noise waveforms, the value at each time-step was drawn from a Gaussian distribution (with standard deviation of one degree), and the resulting signal was lowpass-filtered at 100 Hz (3rd-order Butterworth (Waiblinger et al., 2015)). The white noise waveform around the feature waveform was dampened with an inverted Gaussian, with standard deviation 25.5 ms.

The stimulus conditions were randomized across trials. The stimulus consisted of 1.5 s of either white noise ("adapted" trials) or no white noise ("control" trials), with the onset of the embedded feature at 1 s. The inter-trial interval was a random value (drawn from a uniform distribution) between 2 and 3 s. In a subset of experiments, optogenetic stimulation of either VPm or thalamocortical terminals was randomly interleaved (see below). We typically obtained at least 100 trials per stimulus condition.

Optogenetic stimulation

In a subset of acute anesthetized experiments in Ai32xNR133 transgenic mice, we stimulated thalamocortical terminals in S1 using blue (470 nm) light from an LED (ThorLabs), and recorded either spiking or subthreshold S1 responses. We positioned either a 200 or 400 micron optic fiber (ThorLabs) just above the exposed cortical surface, adjacent to the probe or patch pipette. Light pulses were either 10 ms or 15 ms in duration, and were delivered either in isolation or embedded in sensory white noise delivered to the whisker by the galvanometer. We titrated the light level at the beginning of each recording session to evoke cortical responses that were comparable in amplitude to those evoked by punctate whisker stimulation.

In a subset of awake experiments in Ai32xNR133 transgenic mice, we presented the above sensory stimulus protocol, in addition to a set of "LED" trials in which we delivered a step input of 470 nm light to VPm beginning 1 s before and ending 0.5 s after the delivery of an isolated sensory feature. The light was delivered via LED-coupled fiber attached to the electrode/probe (described above). We titrated the light level at the beginning of each recording session such that steady-state light-evoked firing rates in VPm (based on

threshold crossings of high-pass-filtered voltage recordings) approximately matched
those evoked by the white noise whisker stimulus.

Videography and whisker motion analysis

In five awake recording sessions, we also recorded whisker videography using a CCD
camera (DMK 21BU04.H USB 2.0 monochrome industrial camera, The Imaging Source,
LLC). The face was illuminated using IR LEDs (860 nm, DigiKey), and the face opposite
the galvanometer was imaged at 30 Hz. We used external triggers to acquire frames,
and recorded these triggers in Synapse to synchronize frames to all other recorded
signals.   We analyzed the videos using the python implementation of FaceMap
(www.github.com/MouseLand/FaceMap) (Stringer et al., 2019). For each video, we used
the FaceMap GUI to select two regions of interest (ROIs): one that captured most of the
whiskers, and a second of the nose. We then processed the videos for each ROI, which
involved first calculating the "motion energy" of each frame (i.e., the absolute difference
between the current and previous frame), and then computing the singular vectors of the
motion energy. For each video, we confirmed that the time series of the resulting first
SVDs for each ROI (which we call the "motion timeseries") were qualitatively consistent
with nose/whisker dynamics. We then calculated the relative whisker and nose motion
as a function of the sensory stimulus for each recording session. To do this, we first
calculated the across-time mean value of the motion timeseries in the 1 s preceding
sensory feature onset for each trial, which we call the "mean motion". We compared the
mean motion values for the two stimulus conditions ("spontaneous" and "white noise")
using the Wilcoxon signed-rank test. For visualization, we then calculated the across-trial
mean absolute motion value for the "spontaneous" and "white noise" conditions, and
normalized to the "spontaneous" value.

Spike-sorting

We sorted spikes off-line using KiloSort2 (https://github.com/MouseLand/Kilosort2) for
clustering, and phy (https://github.com/cortex-lab/phy) for manual curation of clusters.
During manual curation, clusters were either merged or separated based primarily on
waveform distributions across the probe and cross-correlogram structure. We discarded
as "noise" those clusters with across-instance mean waveform that did not resemble a
characteristic spike on any channel. All remaining clusters were labeled as either single-
or multi-units by downstream analysis (see below).

Units retained for analysis

We labeled each curated cluster as either a single- or multi-unit based on the signal-to-
noise ratio (SNR) and inter-spike interval (ISI) distribution. For each cluster and recorded
spike, we calculated the absolute voltage difference between the trough and subsequent
peak (VTP) on each channel. We defined the SNR to be the across-trial mean VTP
divided by the across-trial standard deviation, for the channel on which the mean VTP
was greatest. Additionally, we calculated "ISI violation percentage" for each cluster using
the autocorrelogram (ACG). We defined the violation percentage to be the percentage
of spikes in the 0 - 1 ms ACG bin. We then defined a well-isolated single-unit to be a
cluster with SNR greater than 4.0, VTP greater than 50 mV, and ISI violation percentage
below 1%. All other clusters were classified as multi-units. In our anesthetized recording
sessions, we used an alternate probe configuration, which was somewhat less well-suited
to obtaining well-isolated units. We therefore slightly relaxed our inter-spike interval
violation constraints (to 1.5%) for defining "well-isolated" units, to yield more RS and FS
cells from these datasets (see Methods). This did not qualitatively change our results.
For S1, we further segregated single-units into regular- and fast-spiking units based on
the mean waveform. Again using the channel on which the waveform was largest, we
calculated the time from trough to subsequent peak (TTP). We classified S1 units as
either fast-spiking (FS, putative inhibitory) or regular-spiking (RS, putative excitatory)
neurons based on waveform width (McCormick et al., 1985; Niell and Stryker, 2010; Guo
et al., 2017; Isett et al., 2018; Speed et al., 2019; Yu et al., 2019). Specifically, S1 units
with TTP < 0.4 ms were classified as FS, and all others as RS. Waveforms were in
general narrower for VPm units than for cortical units, consistent with previous work
(Barthó et al., 2014). We therefore classified VPm "RS" cells as those with TTP > 0.3 ms,
and excluded units with narrower waveforms, which likely originated from either the cell
bodies or axon terminals of neurons in reticular thalamus (Barthó et al., 2014). For
putative VPm units, we further required that the absolute peak of the PSTH of responses
to isolated punctate stimuli occur between 2 and 10 ms of stimulus onset. Finally, when
analyzing activity of single- and/or multi-units, we only included those units with at least
0.25 mean post-feature spikes per trial, and a significant change ($p < 0.05$, Wilcoxon
signed-rank test) in firing rate after stimulus onset on control (unadapted) trials, using the
weakest sawtooth stimulus delivered during that recording session and 50 ms (30 ms)
pre- and post-stimulus windows for S1 (VPm).
350

Experimental Design and Statistical Analyses

When comparing two sets of values (that were matched samples) across stimulus
conditions, we used the Wilcoxon signed-rank test (implemented in Python using the
wilcoxon function in the Scipy library), and Bonferroni-corrected the resulting p-values for
multiple comparisons where applicable. When comparing two independent samples

(e.g., normalized response rates for RS and FS cells), we used the Kruskal-Wallis test
(implemented in Python using the kruskal function in the Scipy library).

For any analysis resulting in a single value for a given recording session calculated using
all trials (e.g., mean synchronous spikes), we tested for significance of change across
conditions by re-sampling trials with replacement, re-calculating the final value for the re-
sampled pseudo-data, and calculating the 95% confidence intervals (Bonferroni-
corrected if necessary) of the resulting distribution of values.

The number of cells and animals used to calculate each reported result is included in the
text of the Results section and/or figure captions.

Analyses

All analyses were performed using custom scripts in Python 3.0. The details of each
analysis are presented below.

ROC analysis

We calculated the theoretical detectability of sensory features for each significantly-
responsive RS unit (see above) by applying ideal observer analysis (Wang et al., 2010;
Ollerenshaw et al., 2014) to the "population response" distributions for baseline and
feature-evoked activity. For this analysis, we excluded putative inhibitory neurons (FS
units), as we were interested in interpreting the loss of excitatory drive from cortical
neurons, likely to play a role in downstream signaling and thus relevant for percepts. For
each unit, we calculated the firing rate in a 50 ms baseline (pre-feature) and post-feature
window on each trial. For baseline activity, we used a window beginning 500 ms
preceding feature onset, as the white noise was largely dampened in the 50 ms window
immediately preceding the feature on adapted trials. We then calculated the across-trial
mean (μ) and standard deviation (σ) of the observed firing rates, and generated
"population" firing rate distributions, or samples drawn from gamma distributions
parametrized by the data. Specifically, we drew 1000 samples from a gamma distribution
$\Gamma(N * \alpha, \theta)$, where $\alpha = \mu^2/\sigma^2$, $\theta = \sigma^2/\mu$, and N is the assumed number of identical
neurons to which the ideal observer has access (Britten et al., 1992; Stüttgen and
Schwarz, 2008; Wang et al., 2010). We report results using N = 10, but results were
qualitatively similar for N = 1, 5, and 20. For each feature velocity, we calculated the true-
and false-positive rates for 30 evenly-spaced threshold values between 0 and the
maximum response amplitude. We then generated the receiver operator characteristic
(ROC) curves by plotting the set of true-positive values vs. the false-positive values. We
quantified the theoretical detectability as the area under the ROC curve (AUROC).

Because the sensory white noise was presented on adapted trials, it was possible that
the adaptive decrease in feature detectability was due in some degree to elevated, white-
noise-evoked baseline firing. To isolate the effects of feature response adaptation, we
calculated AUROC for a "hybrid" condition using the feature-evoked distributions from
adapted trials, and baseline distributions from control trials. Finally, we repeated this
analysis by systematically varying the width of the baseline and feature-evoked windows,
from 5 ms to 50 ms in 5 ms increments, yielding a time-resolved measure of feature
detectability in each condition.

Feature response latency and synchrony analysis

We sought to estimate the feature response latency of each S1 unit from the PSTH.
However, in contrast to the relatively well-populated grand PSTHs, the sparsely-
populated PSTHs of individual neurons confound latency calculations. As such, we
convolved the spike trains of each neuron with a Gaussian kernel (1 ms standard
deviation), yielding a convolved aggregate spike count time series, or a smoothed PSTH.
We defined the response latency (T_{onset}) for each stimulus condition to be the time at
which the smoothed PSTH exceeded a threshold of four standard deviations of the pre-
stimulus values (calculated from control trials). We further calculated the T_{onset} adaptation
index for each unit as the difference between adapted and control values divided by the
sum of the values. Our reported results were qualitatively robust to reasonable choices
of onset threshold (2.5, 3.5, 4.0, 5.0, and 6.0 standard deviations) and Gaussian kernel
standard deviation (0.5, 1.0, 2.0, and 3.0 ms, not shown).

We calculated the population synchrony of feature responses using the population grand
cross-correlogram (CCG) of single-unit spiking, separately for VPm and S1. First, we
treated each sensory-responsive unit as a "reference" unit, and calculated the spike times
of all other simultaneously-recorded units relative to those of the reference unit.
Specifically, for each trial and spike in a reference unit, we calculated the spike times of
all other simultaneously-recorded units relative to the reference spike time in a +/- 20 ms
window around the reference spike time, and appended these to a grand set of relative
spike times. We repeated this for all spikes, trials, and reference units. We then binned
the grand set of relative spike times (using 1 ms bins) to generate the grand CCG. We
generated a "randomized" grand CCG by producing a grand set of random relative spike
times (+/- 20 ms, drawn from a uniform distribution), equal in length to the actual grand
set, followed by binning. We then subtracted the randomized CCG from the actual CCG,
such that the value at each time lag indicated the number of events beyond that predicted
by a random, uncorrelated process. Finally, we divided the CCG by the number of
contributing pairs. We defined the mean synchronous spike count to be the sum of the
resulting CCG in a +/- 7.5 ms window (Wang et al., 2010). We calculated confidence

intervals by re-sampling the grand set of relative spike times with replacement, and re-
calculating the CCG and synchrony as described above. To include a pair of neurons in
this analysis, we required at least 20 feature-evoked spikes across all trials for both
neurons in both the control and adapted conditions. Our results were qualitatively robust
to different choices of minimum spike count and synchrony window (not shown). For each
feature level, we further calculated the grand CCG and synchrony for a "hybrid" condition,
in which each neuron's spike times from the control condition were subsampled to match
the rate of the adapted condition. For the minority of neurons with larger responses in
the adapted condition, we retained all spikes from the control condition.

S1 membrane potential analysis

We removed action potentials from intracellular voltage recordings by first identifying
spike times and interpolating between the values 2.5 ms before and 2.5 ms after the peak
of the action potential. To identify spike times in each recording, we first calculated the
first time derivative time of the membrane potential at each time step. Spike onsets were
defined to be positive crossings of five standard deviations of this time series. For each
onset, the spike peak time corresponded to the next time at which the derivative was less
than or equal to zero. After interpolation, we low-pass-filtered the resulting time series
(100 Hz, 3rd order Butterworth). For each trial, we calculated the amplitude and time-to-
peak of the subthreshold response to the (sawtooth sensory or optogenetic terminal light)
stimulus, in a 50 ms window following stimulus onset. We first subtracted the minimum
voltage value in this window on each trial, and defined the amplitude of the response to
be the maximum voltage value, and the time-to-peak was the time between stimulus
onset and this peak value. We then calculated the across trial median amplitude and
time-to-peak.

Thalamocortical network model

We constructed a simple model of the thalamocortical network using custom scripts
written in Python 3.6.10. All code is freely available upon request. We modeled a single
cortical barrel as a clustered network of excitatory and inhibitory single-compartment
leaky integrate-and-fire (LIF) neurons, subject to excitatory synaptic inputs from a
"barreloid" of VPm neurons, and well as excitatory non-thalamic inputs that were
independent across cortical neurons. For each of the "control" and "adapted" conditions,
we simulated 50 trials, each lasting 150 ms, with a time-step of 0.05 ms.

We modeled a single VPm barreloid as forty independent trains of tonic and burst spikes,
drawn from the empirical VPm PSTHs. The ongoing and evoked rates for each neuron
were set to the mean empirical values, and were then multiplied by a rate modulation

factor drawn from a skewed gamma distribution (with a shape value of 2.0, a scale value of 1.0, then re-scaled to have a mean value of 1.0), to mimic the broad firing rate distributions of VPm neurons previously reported (Pinto et al., 2000; Bruno and Simons, 2002; Wang et al., 2010; Whitmire et al., 2016). Bursts were modeled as pairs of spikes with 2.5 ms ISI.

Non-zero thalamocortical (TC) synaptic weights were drawn from a broad distribution (a rectified Gaussian distribution with mean 1.0 and standard deviation 2.0), to mimic the reported variability in TC synaptic strengths and/or efficacies (Bruno and Simons, 2002; Gabernet et al., 2005; Bruno and Sakmann, 2006; Cruikshank et al., 2007, 2010; Sermet et al., 2019). Mean initial TC synaptic strengths were the same for excitatory and inhibitory neurons (Sermet et al., 2019), but TC convergence was higher for inhibitory neurons (75% for inhibitory neurons, 50% for excitatory neurons). Finally, VPm neurons with the highest firing rates did not synapse onto excitatory neurons (Bruno and Simons, 2002); for each VPm synapse onto an excitatory neuron, if the rate modulation factor of the VPm neuron was greater than 1.5 standard deviations from 1.0, we set that connection to zero, and replaced it with a connection (of the same strength) to a VPm neuron with rate modulation factor below 1.5. In response to a spike in a given thalamic neuron, all TC synapses from that neuron instantly decayed (by a factor of 0.75), followed by exponential recovery (with time constant 25 ms). We assigned a minimum latency of 2 ms (3 ms) to each TC synapse onto an inhibitory (excitatory) neuron, to reflect the slightly faster arrival of thalamic inputs to L4 FS cells (Cruikshank et al., 2007; Kimura et al., 2010). Each TC synapse was assigned an additional temporal latency value between 0 and 1 ms, drawn from a uniform distribution, which helped to promote evoked cortical firing that was less time-locked across the network. Importantly, the higher TC convergence for inhibitory neurons, the rate-dependent TC connectivity described above, the shorter TC synaptic latencies onto inhibitory neurons, and intrinsic neuronal and intracortical connection parameters (see below) together supported strong feedforward inhibition in this model network, despite the equivalent mean TC synaptic weights onto the two neuron types.

Each thalamic spike resulted in a postsynaptic conductance waveform in each postsynaptic cortical neuron, with excitatory reversal potential 50 mV. Given a thalamic spike at time zero, we modeled the resulting postsynaptic conductance waveform $g(t)$ as a conductance amplitude $\bar{g}$ multiplied by a gating variable $G(t)$. The gating variable at a given TC synapse had the form of a difference of two exponentials, resulting in rise and subsequent decay in the synaptic conductance following a presynaptic spike at time zero:

$$g(t) = \bar{g}G(t)$$

$$G(t) = W \frac{\tau_m^{exc}}{\tau_d - \tau_r} \left[e^{-(t - \tau_l^{VPm})/\tau_d} - e^{-(t - \tau_l^{VPm})/\tau_r} \right]$$

where W is the weight of the TC synapse, τ_m^{exc} is the membrane time constant of excitatory
neurons, τ_d is the decay time constant, τ_r is the rise time constant, and τ_l^{VPm} is the latency
of the synapse. We selected rise and decay time constants of 0.2 and 1.0 ms,
respectively, for all TC synapses, yielding a relatively fast rise and slow decay. We
include τ_m^{exc} in the numerator of the leading factor so that the gating variable is unitless.
Note that this means the across-synapse mean integrated value of G(t) will be equal to
τ_m^{exc}, which should be taken into account when interpreting absolute values of TC synaptic
conductance amplitudes. We tuned the TC synaptic conductance amplitude (to 30 nS)
such that near-simultaneous firing of multiple thalamic neurons was required to evoke
action potentials in target neurons (Bruno and Sakmann, 2006), when intracortical
synaptic strengths were set to zero.

Each cortical network neuron also received random excitatory "nonthalamic" inputs, to
promote baseline network activity. For each neuron, spike times were drawn from a
homogeneous Poisson process, with rate 60 Hz. The synaptic weights for random
external inputs were drawn from a rectified, normalized Gaussian distribution (with mean
of 1.0, standard deviation of 2.0, then rectified, then divided by its mean), to support
variable ongoing firing rates across network neurons. Each random spike resulted in a
postsynaptic conductance waveform, with the same rise and decay time constants as for
thalamocortical inputs, and with conductance amplitude 60 nS.

We modeled a single cortical column as a network of 800 excitatory and 100 inhibitory
neurons, which approximates the relative numbers of these neuron subtypes across
layers in S1 (Lefort et al., 2009). Excitatory-to-excitatory connection probability was 4%,
excitatory-to-inhibitory was 20%, inhibitory-to-excitatory was 17.5%, and inhibitory-to-
inhibitory was 17.5%. While these numbers are approximately three times lower than
what has been reported within a column in S1 (Avermann et al., 2012), these lower
connection probabilities were necessary to maintain network stability, and the relative
values are in qualitative agreement with experimental measurements. We imposed
spatial clustering of connectivity via a "small-world" network connectivity (Litwin-Kumar
and Doiron, 2012; Wright et al., 2017a, 2017b; Hoseini et al., 2019), with 10% re-wiring
probability. First, all neurons were organized on a continuous ring. Then, outputs from a
given cortical excitatory network neuron were projected to 32 of its nearest excitatory
neighbors. For each of these outputs, the connection was reassigned to another
randomly-selected (not previously connected) excitatory neuron, with probability 10%. In
other words, most connections were nearest-neighbor, but some were random (and
possibly long-distance). Each nonzero connection weight was drawn from a heavy-tailed
beta distribution (with alpha value 0.11, beta value 1.0, and then re-scaled to have mean
value 1.0), to approximate the variable intracortical synaptic strengths previously reported

in cortex (Avermann et al., 2012; Cossell et al., 2015; Pala and Petersen, 2015). This
process was repeated for all neuron pair types, according to the connection probabilities
listed above. All intracortical synapses were assigned a latency of τ_l^{Cx} = 1.0 ms, and rise
and decay time constants of 0.2 and 1.0 ms, respectively, and the synaptic gating
variables evolved as described above for TC synapses.   Intracortical synaptic
conductance amplitudes were 27.5 nS for excitatory-to-excitatory connections, 30 nS for
excitatory-to-inhibitory connections, 40 nS for inhibitory-to-excitatory connections, and 30
nS for inhibitory-to-inhibitory connections.

For network LIF neurons, we selected intrinsic neuronal properties that were consistent
with previous modeling studies, and/or were motivated by previous experimental work.
We set the membrane time constant of excitatory neurons (inhibitory neurons) to be 30
ms (20 ms), reflecting the moderately higher input resistances of cortical excitatory
neurons reported in awake mouse (Gentet et al., 2010). Inhibitory neurons also had
shorter refractory periods (1 ms vs. 2 ms), which supported higher firing rates, as
observed previously (Bruno and Simons, 2002; Khatri et al., 2004; Gentet et al., 2010;
Taub et al., 2013). Excitatory neurons were subject to an inhibitory spike-rate adaptation
conductance, with a 1 ms rise time constant, and 30 ms decay time constant. We tuned
the time constants and amplitude of this conductance such that the ISI of spikes evoked
by tonic current injection gradually increased over the duration of the stimulus. This
conductance served to stabilize excitatory firing, and therefore the network as a whole.
For each neuron, the leak reversal potential E_{leak} was drawn from a uniform distribution
between -70 mV and -60 mV (which added variability in spike timing across the network),
and the leak conductance amplitude was 100 nS. All neurons had a spike threshold of -
45 mV, reset to -59 mV immediately after a spike, and were held at the reset value during
the refractory period.

For each condition, we simulated 50 trials, each lasting 150 ms (including a 50 ms "buffer
window" to allow the network to reach stead-state, a 50 ms "pre-stimulus" window, and a
50 ms "post-stimulus" window), with a time-step of 0.05 ms. At each time-step, the
membrane potential V of a given network neuron of type Y (where Y is either E or I,
representing an excitatory or inhibitory neuron) evolved according to the equations

575
$$V[t] = V[t-1] - \frac{dt}{\tau_m g_{leak}} \{g_{leak}(V[t-1] - E_{leak}) + I_{exc}[t] + I_{inh}[t]\}$$

577
$$I_{exc}[t] = (g_{YE}[t - \tau_l^{Cx}] + g_{VPm}[t - \tau_l^{VPm}] + g_{rand}[t-1])(V[t-1] - E_{exc})$$

579
$$I_{inh}[t] = g_{YI}[t - \tau_l^{Cx}](V[t-1] - E_{inh})$$

$$581 \qquad\qquad\qquad g_{YZ}[t] = \bar{g}_{YZ}S_{YZ}[t]$$

$$583 \qquad\qquad S_{YZ}[t] = S_{YZ}[t-1]\left(1 - \frac{dt}{\tau_d}\right) + \frac{dt}{\tau_d}X_{YZ}[t-1]$$

$$585 \qquad\qquad X_{YZ}[t] = X_{YZ}[t-1]\left(1 - \frac{dt}{\tau_r}\right) + \tau_m^{exc}\frac{dt}{\tau_r}W \cdot T[t - \tau_l^{Cx}]$$

where g_{YZ} is the summed synaptic conductance time series from all presynaptic network
neurons of type $Z = E|I$ onto this target neuron of type $Y = E|I$, g_{VPm} is the summed
synaptic conductance time series from all presynaptic VPm neurons, and g_{rand} is an
excitatory synaptic conductance resulting from Poisson process spike times that were
drawn independently for each neuron. The equations for S_{YZ} and X_{YZ} represent the
implementation of the gating variable *G(t)* described above, associated with conductance
of type *Z* onto neuron of type *Y*. *W* is the (input) synaptic weight matrix for presynaptic
neurons of type *Z* onto neurons of type *Y*, and *T* is the binary spike train sequence for all
presynaptic neurons of type *Z*. Excitatory neurons were also subject to an inhibitory
spike-rate-adaptation current, described above*.*

We quantified model responses by calculating the peaks of the grand PSTHs for
excitatory and inhibitory LIF neurons and divided "adapted" values by "control" values,
yielding the normalized adapted response. We generated grand cross-correlograms (as
described above) for 200 randomly-selected excitatory-excitatory and inhibitory-inhibitory
pairs, and for 100 VPm-VPm pairs.

We further employed alternate models to parse the roles played by synchronous thalamic
spikes and feedforward inhibition. For the "reduced synch" models, we maintained the
mean spike rates of the original model, but manually adjusted drawn VPm spike times to
reduce synchrony. Specifically, if a drawn VPm spike time was within +/- 5 ms of the
empirical PSTH peak time, we shifted the spike to a random later time with probability
P(shift), within approximately 20 ms of the PSTH peak. We repeated this process for
P(shift) = 0.1, 0.2, 0.3, and 0.4. For the "Identical TC Connectivity" model, excitatory and
inhibitory neurons had the same TC convergence values (50%), TC synaptic latencies
were identical for excitatory and inhibitory neurons,s and we did not require that VPm
neurons with the highest rates synapse exclusively onto inhibitory neurons.

Results

To investigate the adaptive effects of persistent sensory stimulation on S1 sensory responses during wakefulness, we presented precise deflections to a single whisker of the awake, head-fixed mouse using a computer controlled galvanometer, and recorded extracellular spiking activity in the corresponding principal column of S1, and/or principal barreloid of VPm (Fig. 1A, see Methods). We presented punctate "sawtooth" sensory features either in isolation or embedded in an adapting background stimulus (frozen sensory white noise, Fig. 1A). The punctate stimulus feature captures the basic nature of the high velocity "stick-slip" whisker motion events that occur as a result of whisker contacts with larger surface irregularities during active sensation (Ritt et al., 2008; Wolfe et al., 2008; Jadhav et al., 2009; Jadhav and Feldman, 2010). During whisker contacts with surfaces, these stick-slip events are embedded in patterns of smaller-amplitude, irregular deflections (Jadhav and Feldman, 2010), simplistically captured here utilizing low-amplitude, repeatable background white noise whisker stimulation, which has been shown to alter perceptual reports in behaving rats (Waiblinger et al., 2015, 2018) and adapt thalamic feature responses under anesthesia (Whitmire et al., 2016). We first characterized the effects of the background stimulus on baseline and feature-evoked cortical firing during wakefulness (Figs. 1 – 3), and then sought to identify the mechanisms underlying these effects through a battery of additional experiments (Figs. 4 – 7) and thalamocortical network modeling (Fig. 8).

S1 exhibits profound and differential sensory adaptation during wakefulness

We first characterized the effects of the background white noise adapting stimulus on baseline spiking activity in S1. We segregated well-isolated, sensory-responsive cortical units into regular-spiking (RS, putative excitatory) and fast-spiking (FS, putative inhibitory) neurons (Fig. 1B, see Methods). The sensory white noise evoked a sharp increase in mean RS and FS firing, such that mean rates were significantly elevated during the "early" white noise response window relative to spontaneous activity (Fig. 1C, D). Firing rates rapidly adapted, but remained significantly elevated above spontaneous levels in the "late" white noise response window (Fig. 1D). The effect was generally more pronounced for FS cells (Fig. 1C, D). Importantly, this elevation in baseline firing rates did not reflect white-noise-evoked whisking; not only were there distinct features within each PSTH – consistent with stimulus-locked firing and therefore a largely stationary whisker (Fig. 1C) – but we also confirmed using whisker videography in a subset of recording sessions that whisker (Fig. 1D, bottom) and nose (not shown) motion did not systematically change when the whisker white noise was applied (see Methods). As such, it is likely that the induced elevation in cortical firing rates was largely due to afferent sensory drive.

Next, we investigated the adaptive effects of the background stimulus on responses to
sensory features during wakefulness. We delivered a 300 deg/s feature to a single
whisker, either in isolation (the "control" condition), or embedded in background
stimulation (the "adapted" condition, Fig. 1A, see Methods), with the feature delivered 1
s after background stimulus onset. To avoid distortion of the feature waveform, the
background stimulus was dampened with an inverted Gaussian waveform in the
neighborhood of the feature (Waiblinger et al., 2015; Whitmire et al., 2016; see Methods).
This also allowed pre-feature RS and FS firing rates to return to near-baseline values at
feature onset (Fig. 1E). We delivered both moderate (300 deg/s) and relatively strong
(900 deg/s) features in a subset of experiments. In the control condition, features
evoked robust, short-latency spiking responses (Fig. 1E), consistent with previous work
in the anesthetized (Bruno and Simons, 2002; Pinto et al., 2003; Khatri et al., 2004; Wang
et al., 2010) and awake (Musall et al., 2014) rodent. We next asked whether the
background stimulus appreciably adapted feature responses, or whether the relatively
high baseline firing rates during wakefulness (Fig. 1D) resulted in a "pre-adapted" circuit
(Castro-Alamancos, 2004). We found that S1 feature responses were in fact substantially
muted when features were embedded in the background stimulus (Fig. 1E). To
characterize RS and FS feature responses and the effects of adaptation, we calculated
across-trial mean evoked rates using a 50 ms window following feature onset. For both
cell types and feature velocities, the peak (Fig. 1E) and mean (Fig. 1F) evoked rates were
reduced in the adapted condition. Interestingly, adaptation appeared to be more profound
for RS cells, in terms of proportional changes in sensory responses (Fig. 1E, F). To
further quantify the effects of adaptation on a cell-by-cell basis, and to capture cell-type-
specific adaptation, we calculated the normalized adapted response for each cell (i.e., the
across-trial mean adapted response rate divided by the mean control rate, see Methods).
For both feature velocities and cell types, population median normalized adapted
responses were less than one (capturing the general adaptive reduction in evoked rate),
and RS cells were indeed significantly more adapted than FS cells (Fig. 1G). This
differential effect was not specific to any cortical depth (Fig. 1H).

In summary, we observed profound rapid sensory adaptation in S1 of the awake mouse,
and cortical putative excitatory neurons were more adapted than putative inhibitory
neurons.

Adaptation reduces the theoretical detectability of punctate sensory stimuli

The adaptation of S1 feature responses should diminish the ability to distinguish the
sensory feature from the sensory background. To quantify the potential impact of
adaptation on feature detectability, we applied a signal detection theory framework to the
spiking activity of putative excitatory neurons, which are likely to play a role in downstream

signaling and thus are relevant for percepts. Qualitatively, theoretical feature detectability
is inversely related to the degree of overlap between the baseline and feature-evoked
firing rate distributions. For each S1 RS unit, we generated spike rate distributions using
50 ms windows of baseline and feature-evoked spiking, and drew samples from
"population" gamma distributions (Fig. 2A) parametrized by the empirical mean and
standard deviation ((Wang et al., 2010) see Methods). Consistent with the adaptive
decrease in single-neuron response rates, adaptation tended to move the feature-evoked
distribution toward the baseline distribution, increasing the degree of overlap (Fig. 2A,
right). We quantified the overlap by calculating the area under the receiver operator
characteristic curve (Wang et al., 2010; Whitmire et al., 2016) (AUROC, Fig. 2B), which
has a value of 1.0 for non-overlapping distributions, and 0.5 for complete
overlap. Adaptation significantly reduced the across-unit mean AUROC for both feature
velocities (Fig. 2C, bottom). Notably, repeating this analysis for a "hybrid" scenario using
the adapted feature-evoked distributions and control baseline distributions did not
significantly alter AUROC values relative to the true adapted condition (Fig. 2C, left),
demonstrating that the decrease in detectability reflected feature response adaptation,
rather than an elevation of baseline rates in response to the adapting stimulus.
710
The above analysis considers spiking in 50 ms pre- and post-feature windows, and the
results are consistent with the adaptation of mean feature-evoked RS rates. But it is also
apparent from visual inspection of the RS grand PSTHs that the time-course of S1 spiking
varied with feature velocity and sensory adaptation. To resolve the dependence of
feature detectability on time, we varied the width of the post-feature window from 5 ms to
50 ms in 5 ms increments (see Methods). Consistent with the RS grand PSTHs, we found
that in the control condition, AUROC values increased drastically in the first 15 ms of the
feature response, and reached peak values within approximately 25 ms (Fig. 2D). The
separation between control and adapted AUROC values largely occurred during this early
phase of the response (Fig. 2D). In other words, the adaptive loss of theoretical
detectability largely reflected a decrease in short-latency feature-evoked spiking.
722
Together, these results quantify the degree to which sensory adaptation reduces the
theoretical detectability of sensory features during wakefulness, and demonstrate that this
largely reflects adaptation of the early feature response. This is qualitatively consistent
with previous reports showing significant adaptation of theoretical feature detectability in
the anesthetized rat (Wang et al., 2010; Ollerenshaw et al., 2014; Zheng et al., 2015),
and perceptual threshold in awake rats (Ollerenshaw et al., 2014; Waiblinger et al., 2015).
729
730 **Adaptation strongly influences the timing of feature-evoked spiking in S1**
731

As highlighted above, adaptation altered S1 sensory response timing, a key component of cortical sensory representations that strongly influences how and what signals are transmitted to downstream brain structures. Specifically, in addition to the dramatic reduction in early feature-evoked spiking (particularly among RS cells), visual inspection of the grand PSTHs suggested adaptation increased the latency of feature responses. These phenomena may be relevant for interpreting the mechanistic basis of S1 adaptation, and the likely perceptual implications; both are consistent with weaker feedforward excitatory drive to S1, and could influence signaling to downstream targets. We therefore sought to quantify these adaptive effects, and began by calculating the response latency (T_{onset}) for each S1 unit (see Methods). We found that for both RS and FS cells, and for both stimulus strengths, adaptation increased T_{onset} values (Fig. 3A). To further quantify and compare adaptive changes in T_{onset} across cell types, we calculated the T_{onset} adaptation index (AI) for each cell (see Methods) and compared population medians for RS and FS cells. For both stimulus strengths, the adaptive increase in response onset times was significantly greater for RS cells (300 deg/s RS median T_{onset} AI = 0.21, FS median T_{onset} AI = 0.02, p = 1.21 x 10^{-4}, Kruskal-Wallis test; 900 deg/s RS median T_{onset} AI = 0.13, FS median T_{onset} AI = 0.00, p = 4.93 x 10^{-4}).

Next, we reasoned that the observed changes in S1 PSTH shapes might reflect a decrease in synchronous spiking among cortical neurons. In general, the degree of synchronous firing among a population of neurons is likely related to the effect on synaptic targets; stronger inhibitory synchrony will tend to silence postsynaptic neurons, and stronger excitatory synchrony will be more efficacious for driving postsynaptic neurons – possibly downstream from S1. Indeed, it has been shown that coordinated population firing is a better predictor than mean rates of stimulus identity (Jadhav et al., 2009; Safaai et al., 2013; Zuo et al., 2015) and behavioral stimulus discrimination performance (Safaai et al., 2013; Zuo et al., 2015) when rodents whisk across textured surfaces.

To calculate the prevalence of synchronous spikes in feature responses, we populated the grand cross-correlogram (CCG) for RS-RS and FS-FS pairs for each stimulus condition using spike trains from all pairs of simultaneously-recorded responsive cells, and defined synchronous spikes to be those within a +/- 7.5 ms window around zero lag (Fig. 3B, see Methods, Wang et al. 2010). We found that adaptation drastically reduced the amplitude and sharpness of the RS-RS CCGs (Fig. 3C, left), while more modestly reducing the grand CCG amplitude for FS-FS pairs (Fig. 3C, right). This represented a significant decrease in synchronous spike counts for both pair types and stimulus velocities (Fig. 3D). As reflected in the grand CCGs, the decrease in FS-FS synchrony was significant, but proportionally smaller than that of RS-RS synchrony. We next asked whether the adaptive decrease in synchronous spikes simply reflected the decrease in mean evoked rate, or possibly also other de-synchronizing mechanisms. For each

feature velocity and cell type, we simulated a "hybrid" condition in which we sub-sampled spikes from the control condition, matching the rate of the adapted condition (see Methods). We found that synchronous spike counts were significantly lower in the adapted than in the hybrid condition (Fig. 3D). Thus, though the relationship between mean rate and synchronous spike count is nonlinear, the adaptive decrease in synchronous spiking was greater than that predicted by the loss of feature-evoked spikes in individual cortical neurons alone. This non-trivial decrease in synchronous firing suggests the feedforward drive to S1 is not only less efficacious for driving spikes in the adapted condition, but may also be less synchronous across thalamorecipient neurons.

Thus, rapid sensory adaptation in S1 during wakefulness not only reduced mean evoked spike rates and theoretical feature detectability, but also disrupted timing through increased response latencies and reduced synchrony. In both aspects of timing, as with mean evoked rates, the adaptive effects on RS cells were more dramatic. The loss of synchronous excitatory firing has implications for the driving of targets downstream of S1, and ultimately for perception and behavior. The more modest decrease in synchronous inhibitory firing implies that inhibitory neurons were still relatively strongly driven in the adapted condition. Further, the synchronous inhibitory spiking that survives adaptation should provide robust feedforward inhibition to S1 excitatory neurons, which may in part explain the more profound adaptation of RS cells.

S1 sensory responses are also adapted under anesthesia, but RS cells are not differentially adapted

Having established the extent and characteristics of rapid sensory adaptation in S1 of the awake mouse, we next turned our attention to the underlying mechanisms, which may reflect a combination of feedforward, recurrent, and top-down sources. Our first step in doing so was to compare these results to those from the anesthetized mouse, for two reasons. First, while the background stimulus elevated S1 firing rates at least in part via feedforward sensory inputs (as evidenced by the degree of stimulus-locked firing), it is still possible that this adapting stimulus also evoked top-down modulation, e.g., through systematic changes in ongoing S1 state via arousal-related neuromodulation (Reimer et al., 2014, 2016; Mcginley et al., 2015; McGinley et al., 2015). In the anesthetized mouse, ongoing state changes caused by endogenous processes should be independent of the sensory white noise, and as such, white-noise-induced changes in mean feature representations should not reflect top-down mechanisms. Second, isoflurane anesthesia tends to silence the secondary posteromedial nucleus of thalamus (Suzuki and Larkum, 2020), lowers baseline cortical firing rates (Greenberg et al., 2008; Vizuete et al., 2012; Aasebø et al., 2017), and generally weakens cortical inhibition (Haider et al., 2013) and other intracortical interactions (Suzuki and Larkum, 2020). We therefore repeated our

812   experiments in a different set of mice lightly-anesthetized with isoflurane, using the
813   anesthesia to unmask the feedforward inputs from VPm to S1.
814
As expected, baseline firing rates under anesthesia were quite low compared to those
recorded during wakefulness (Fig. 4B, C). Despite these differences in baseline activity,
the anesthetized experiments recapitulated several key aspects of the awake data. First,
the background stimulus evoked a sharp increase in RS and FS firing in the "early" white
noise response window, with rapid adaptation toward "late" response rates that were
significantly elevated above spontaneous values (Fig. 4B, C), and with stimulus-locked
firing evident among FS neurons (Fig. 4B). Further, adaptation clearly decreased the
peak (Fig. 4D) and mean evoked firing rates of responses to 300 deg/s features (Fig. 4E),
and qualitatively shifted PSTHs to higher response latencies (Fig. 4D). Notably, however,
excitatory neurons were not differentially adapted under anesthesia (Fig. 4F). In light of
this, it is possible that the stronger cortical inhibition typical of wakefulness (Haider et al.,
2013) contributed to adaptation of RS cells in our awake recordings via robust
feedforward inhibition.
828
In summary, the elevation of (stimulus-locked) firing with presentation of background
sensory stimulation, and net adaptation of S1 feature responses was robust to
anesthesia, suggesting that feedforward mechanisms are sufficient to explain these
phenomena. In contrast, the differential adaptation of S1 excitatory neurons was
abolished by anesthesia, suggesting the set of mechanisms underlying this phenomenon
– possibly including strong feedforward inhibition – were not all active in the anesthetized
state.
836
837   **Adaptation primarily influences feature response timing in VPm thalamus**
838
Having established that several aspects of S1 sensory adaptation may reflect feedforward
mechanisms, we next sought to identify those mechanisms. Previous *in vitro* and
anesthetized work has demonstrated adaptation of evoked rate (Hartings et al., 2003;
Khatri et al., 2004; Gabernet et al., 2005; Ganmor et al., 2010; Wang et al., 2010;
Ollerenshaw et al., 2014; Whitmire et al., 2016; Liu et al., 2017), single-unit bursting
(Whitmire et al., 2016) and population synchrony (Wang et al., 2010; Ollerenshaw et al.,
2014) in VPm, suggesting S1 adaptation is inherited to some degree directly from VPm,
yet this has never been tested during wakefulness. We therefore next repeated our
experiments while recording spiking activity in VPm of the awake mouse (Fig. 5A, see
Methods), and asked how adaptation altered the rate and timing of thalamic sensory
responses.
850

We first parsed VPm spikes into putative burst and tonic spikes, and asked how
presentation of sensory white noise affected baseline tonic and burst firing. We defined
bursts according to classic criteria (Lu et al., 1992; Reinagel et al., 1999; Swadlow and
Gusev, 2001; Lesica et al., 2006; Whitmire et al., 2016): two or more sequential spikes
from a single unit preceded by at least 100 ms of quiescence, and with at most 4 ms
between spikes (Fig. 5A, bottom). These criteria are consistent with the timing of burst
spikes resulting from de-inactivation of T-type calcium channels after prolonged
hyperpolarization (McCormick, 1992; Llinás and Steriade, 2006), though we did not
confirm this directly via intracellular recordings. As in S1, we found that the background
stimulus evoked stimulus-locked firing in VPm (Fig. 5B). Tonic firing rates increased
sharply at white noise onset, and rapidly adapted to "late" white noise response rates that
were significantly elevated above spontaneous levels (Fig. 5B, C). Burst rates were very
low during spontaneous activity (Fig. 5B, C), increased sharply at white noise onset,
before decreasing below spontaneous levels in the "late" response window (Fig. 5B, C) –
consistent with inactivation of T-type calcium channels. We next characterized VPm
responses to sensory features, and the effects of the background stimulus on these
responses.   We found that background stimulation reduced feature-evoked tonic spiking
(Fig. 5D, E), but this adaptive effect was modest in comparison to the adaptation of S1
RS cells. Adaptation had a more striking effect, however, on burst firing in individual
neurons. Burst firing has been shown to provide potent synaptic drive to S1 (Sherman,
2001; Swadlow and Gusev, 2001; Swadlow, 2002), and may therefore be critical for
shaping cortical sensory responses. Yet few studies have explored single-unit VPm
sensory responses during wakefulness, when VPm is likely to be on average relatively
depolarized, and T-type calcium channels inactivated. We did, in fact, observe feature-
evoked bursting, consistent with previous recordings in awake rats (Whitmire et al., 2016),
though burst spikes constituted a minority of total feature-evoked spikes in the control
condition (Fig. 5E). Adaptation profoundly reduced bursting, with feature-evoked burst
spikes nearly abolished in the adapted condition (Fig. 5D, E). Finally, we asked how
adaptation altered synchronous thalamic firing, shown previously to be crucial for driving
downstream cortical targets (Bruno and Sakmann, 2006; Wang et al., 2010; Bruno, 2011).
We thus generated grand CCGs and counted synchronous spikes for each stimulus
condition, as done previously for S1 cells. We found that adaptation significantly and
substantially decreased synchronous VPm spike counts for both feature velocities (Fig.
5F, G). Interestingly, adapted synchronous spike counts were not significantly different
from those of the hypothetical "hybrid condition" (Fig. 5G). Thus, unlike S1, VPm was
apparently not subject to mechanisms that actively de-synchronized firing across thalamic
neurons beyond the degree predicted by loss of rate alone. As such, the loss of
synchrony in this regime of adaptation may be attributable to weakened synaptic drive
from trigeminal neurons (Castro-Alamancos, 2002; Timofeeva and Lavalle, 2003;
Ganmor et al., 2010). This is in contrast to previous studies in anesthetized (Wang et al.,

2010) and awake (Ollerenshaw et al., 2014) rats, which demonstrated that sequences of
high-velocity stimulus features resulted in a gradual loss of synchrony greater than that
predicted by adaptation of mean rate alone. This discrepancy may owe to the relatively
weak adapting stimulus we employ here; stronger adapting stimuli may engage a broader
range of mechanisms that serve to desynchronize VPm firing, possibly including more
profound adaptation of trigeminal neurons, and weakened synaptic interactions between
VPm and TRN.

In summary, rapid sensory adaptation modestly reduced feature-evoked tonic spiking in
VPm, and substantially reduced bursting and synchronous spiking. Given the sensitivity
of S1 to bursting and synchronous firing in VPm (Usrey et al., 2000; Sherman, 2001;
Swadlow and Gusev, 2001; Swadlow, 2002; Bruno and Sakmann, 2006; Wang et al.,
2010; Ollerenshaw et al., 2014), this qualitatively predicts that cortical responses will be
smaller in amplitude and less synchronous across neurons, as observed in experiment.
Further, the more substantial loss of synchronous firing in S1 than in VPm demonstrates
an apparent enhancement of adaptation-induced desynchronization from thalamus to
cortex.

Thalamocortical and intracortical synaptic depression contribute little to S1
sensory adaptation

While the thalamic adaptation we observed likely played a key role in S1 adaptation, what
was the additional contribution from thalamocortical and/or intracortical synaptic
depression? Previous anesthetized work has demonstrated greater adaptation of mean
evoked rates in S1 than in VPm (Chung et al., 2002; Khatri et al., 2004; Gabernet et al.,
2005), and this has been interpreted to reflect thalamocortical synaptic depression.
However, this may also reflect the sensitivity of cortex to bursting and synchronous
thalamic firing, which we found to be profoundly adapted by persistent sensory
stimulation. We sought to disentangle the relative contributions of thalamic response
adaptation and thalamocortical and/or intracortical synaptic depression by performing two
complementary optogenetic experiments: one to isolate the contribution from
thalamocortical synaptic depression, and one to isolate the contribution from intracortical
mechanisms. In these experiments, we achieved targeted manipulation of either VPm
neurons or thalamocortical synapses – without directly manipulating whisker-responsive
POm or TRN neurons – by using a transgenic mouse (Ai32xNR133, see Methods)
expressing Channelrhodopsin in VPm/VPl cell bodies, axons, and thalamocortical axon
terminals.

In the first experiment, we sought to isolate the contribution of thalamocortical synaptic
depression to S1 response adaptation. We did this by presenting a background
optogenetic stimulus that bypassed the early sensory pathway upstream of VPm, and

elevated baseline VPm rates without elevating S1 rates. We inserted an optoelectrode
into VPm and a linear silicon probe into the topographically-aligned column of S1 in the
awake mouse (Fig. 6A, see Methods), and we randomly interleaved "white noise" and
"LED" trials. On "LED" trials, we elevated baseline VPm rates by substituting the adapting
sensory stimulus with a step input of blue light to thalamus (Fig. 6B, right). We titrated
the light power such that mean baseline thalamic rates were comparable to "white noise"
trials (Fig. 6B, 6C, top). Importantly, while the LED significantly increased VPm firing
rates (Fig. 6B, C top, spont vs. LED), it did not evoke even a transient increase in S1 firing
rates (Fig. 6B, C, bottom), likely because thalamic spiking was not sufficiently
synchronous to effectively drive cortical targets (Bruno and Sakmann, 2006). As such,
this manipulation did not engage activity-dependent intracortical adaptation mechanisms.
942
We next inspected the effects of our sensory and optogenetic manipulations on VPm and
S1 sensory feature responses. We were interested in the presence or absence of gross
adaptive effects, and so we grouped together tonic and burst spikes in VPm, and RS and
FS cells in S1 for this analysis. Because sensory adaptation was qualitatively similar for
300 deg/s and 900 deg/s features, we presented only the 300 deg/s sensory feature in
these experiments. On LED trials, we maintained a constant light level during
presentation of the sensory feature, to avoid transient VPm responses to reduction in light
power. As shown above, background sensory stimulation adapted feature responses in
VPm and S1 (Fig. 6D). We next inspected the effects of optogenetically-elevated
baseline VPm rates on feature responses. If the "artificial" elevation of baseline VPm rate
adapted TC synapses prior to delivery of the sensory feature, we would anticipate
adapted S1 feature responses on LED trials, despite the non-adapted VPm feature
response. On the contrary, we observed no significant differences in S1 feature response
rates between the control and LED conditions (Fig. 6D, bottom). Importantly, LED
presentation did not significantly enhance synchronous spike counts in the VPm feature
response relative to control trials (Fig. 6E), thus ruling out the possible confound of
enhanced feature-evoked thalamic synchrony masking the effects of TC synaptic
depression. As such, the results of this first optogenetic experiment suggest that S1
response adaptation did not largely reflect TC synaptic depression, but rather a
combination of VPm response adaptation and intracortical mechanisms.
963
We next performed a second, complementary optogenetic experiment to determine the
relative contribution from intracortical mechanisms engaged by the elevation of pre-
feature cortical firing rates (e.g., intracortical synaptic depression, a build-up of
intracortical inhibition, etc.). We positioned an LED-coupled optic fiber above the cortical
surface and recorded extracellular spiking activity with a silicon probe array in S1 (Fig.
7A). We performed these experiments in lightly-anesthetized mice – in which we also
observed profound S1 adaptation (Fig. 4) – which allowed additional recording time to

titrate light levels (see Methods). On each trial, we presented either a 300 deg/s sensory feature to the principal whisker (as described above), or a "light feature", which was a brief step input of light to stimulate TC terminals in the principal column, with sensory and light features randomly interleaved. In both cases, we presented the features either in isolation (control trials) or embedded in a background sensory stimulus (adapted trials). We recorded from a pool of 208 RS, FS, and multi-units across these experiments. Of these, 121 responded significantly to the sensory but not the light feature, 0 responded only to the light feature, 71 responded to both, and 16 did not respond to either (Fig. 7B, inset). For the following analyses, we consider only the units that responded significantly to both feature types (see Methods).

Consistent with the above results for awake and anesthetized mice, the S1 sensory feature response grand PSTH for pooled RS, FS, and multi-units exhibited profound sensory adaptation (Fig. 7B, left). If this largely reflected intracortical mechanisms engaged prior to feature delivery, we would expect that light feature responses to be similarly adapted by the background sensory stimulus, as the light feature precluded VPm response adaptation. Instead, there was comparatively little evidence of adaptation in the S1 grand PSTHs for light feature responses (Fig. 7B, right). This result also bore out in mean evoked rates: mean rates for sensory feature responses were profoundly adapted (Fig. 7C, left), while light feature responses were only slightly adapted (Fig. 7C, right), and the population median normalized adapted response was near 1 for light feature responses, but significantly smaller for sensory feature responses (Fig. 7D).

Importantly, this did not appear to simply reflect an overwhelmingly strong LED stimulus, or light-evoked TC synaptic activity that was unnaturally synchronous across synapses; not only were evoked rates generally lower for the light feature than for the sensory feature across all neurons (Fig. 7B, C), but in exploring a variety of LED stimulus amplitudes and durations across experiments, we found that both relatively large- and small-amplitude light-evoked responses were at most only modestly adapted, and that the degree to which sensory white noise adapted light-evoked responses was unrelated to the amplitude of the control response for a given unit (Fig. 7E). Further, given that we observed no units that were responsive to the light but not the sensory feature (Fig. 7B, inset), it is also unlikely that the light feature responses were driven primarily by TC synapses and/or cortical neurons that were not responsive to sensory stimulation (and therefore could not be adapted by sensory white noise).

We confirmed these observations of spiking activity by obtaining in vivo patch clamp recordings from neurons in S1 of the lightly-anesthetized mouse (Fig. 7F, left, see Methods). We recorded from four neurons that responded to both sensory and light features. While sensory- and light-evoked amplitudes varied across neurons (Fig. 7G,

top), light feature responses were comparatively impervious to sensory adaptation. Specifically, for sensory feature responses, the across-trial median amplitude significantly decreased for three cells, and the time to response peak significantly increased for all cells, (Fig. 7G, left, see Methods), consistent with the extracellular recordings (Fig. 7B, C). In contrast, light feature response amplitudes were unchanged and the time to peak response did not increase for any of these cells (Fig. 7G, right).

These two complementary optogenetic experiments together suggest that intracortical mechanisms engaged during pre-feature activity – including TC synaptic depression – contributed little to S1 response adaptation, which in this regime of adaptation appear to primarily reflect the profound loss of synchronous feature-evoked VPm spiking.

A model network identifies synchronous VPm spikes and robust feedforward inhibition as key mechanisms underlying S1 response adaptation

We observed that S1 feature response rates were more profoundly adapted than VPm feature response rates, and we have proposed that this reflected the sensitivity of S1 to the adaptive loss of synchronous VPm spikes. Further, we hypothesized that feedforward cortical inhibition that was relatively robust to sensory adaptation helped to dampen feature-evoked S1 excitatory firing in the adapted condition. In other words, inhibition increased the sensitivity of cortical excitatory neurons to adaptation of synchronous VPm spiking. To test the reasonableness of these assertions, and to identify the mechanistic basis for robust feedforward inhibition, we implemented a model thalamocortical network, and assessed its ability to reproduce profound, cell-type-specific adaptation.

We modeled a single S1 barrel as a clustered network of 800 excitatory and 100 inhibitory leaky integrate-and-fire neurons, subject to excitatory inputs from a model "VPm barreloid" (Fig. 8A). The barreloid was modeled as 40 independent trains of tonic and burst spikes, with spike times drawn from the empirical VPm PSTHs. We selected cortical network and intrinsic neuronal parameters that mimicked measurements from previous studies, and then adjusted parameters slightly to ensure stable ongoing and evoked network activity (see Methods). Importantly, we implemented differential thalamocortical (TC) connectivity, which we hypothesized might contribute to the robustness of inhibitory firing to VPm adaptation. This differential connectivity consisted of three key components motivated by previous experimental work. First, inhibitory neurons had higher "TC convergence" (or proportion of VPm neurons that synapse onto each cortical neuron) than excitatory neurons (0.75 vs. 0.5, (Bruno and Simons, 2002)). Second, the baseline and evoked firing rate of each VPm neuron was drawn from a skewed distribution, and VPm neurons with the highest rates synapsed exclusively onto inhibitory neurons ((Bruno and Simons, 2002), see Methods). Finally, TC synaptic latencies were on average 1 ms

shorter for inhibitory neurons (Cruikshank et al., 2007; Kimura et al., 2010). With this
architecture in place, we modeled 50 trials from each stimulus condition of interest.

We fine-tuned the model parameters to give qualitatively realistic peak rates for excitatory
neurons when VPm spike times were drawn from the control (unadapted) empirical
PSTHs for the 300 deg/s and 900 deg/s sensory features (Fig. 8B). Thus, the excitatory
population was appropriately tuned to the rate and synchrony of thalamic firing. Next, we
repeated the simulations using VPm spikes drawn from the adapted PSTHs (Fig. 8B, top,
filled PSTH), and found that cortical network excitatory neurons were in fact profoundly
adapted, despite the only modest reduction in mean VPm rates (Fig. 8B, center). Further,
the excitatory population was more strongly adapted than the inhibitory population (Fig.
8B, C). Finally, as observed in experiment, synchronous spike counts were significantly
reduced in the adapted condition for excitatory-excitatory (Fig. 8D, left) and inhibitory-
inhibitory (Fig. 8D, right) pairs (see Methods), with a more drastic reduction for excitatory
neurons. Thus, the mechanisms incorporated in this simple model were sufficient to
qualitatively reproduce our key experimental results.

In these simulations, both the mean VPm rate (Fig. 8B, top) and VPm synchronous
spiking (Fig. 8E, top) were reduced in the adapted condition. We next assessed the
degree to which the loss in synchronous VPm spikes alone could explain cortical
adaptation. To do this, we repeated the simulations while "manually" manipulating VPm
spike times. Specifically, we first drew VPm spike times from the control PSTHs, and for
each spike that occurred within 5 ms of the PSTH peak time, we shifted the spike to a
random higher latency (within approximately 20 ms of the peak) with 30% probability (see
Methods). This had the effect of maintaining the mean evoked VPm rate, while reducing
the number of near-coincident pairs of VPm spikes in the early response (Fig. 8E, bottom,
"reduced synch" condition). We found that this change alone – which only modestly
affected the resulting VPm grand PSTH (Fig. 8F, top) – was sufficient to profoundly adapt
mean excitatory evoked rates (Fig. 8F). Further, excitatory and inhibitory responses
decreased monotonically with percentage of shifted VPm spikes, with excitatory cells
more sensitive to this manipulation (Fig. 8G). In other words, a loss of synchronous VPm
spiking alone was sufficient to reproduce profound and differential cortical response
adaptation in this network.

Finally, we asked whether robust feedforward inhibition – mediated in part by differential
TC connectivity – contributed to the adaptation of network excitatory neurons. We
modified the network slightly by setting identical TC convergence and TC synaptic latency
values for excitatory and inhibitory neurons and eliminating rate-dependent TC
connectivity (Fig. 8H). Inhibitory and excitatory neurons therefore had identical mean
thalamocortical connection properties, though differences in intrinsic neuronal properties
and dense excitatory-to-inhibitory connectivity still allowed for higher mean firing rates in
the inhibitory population (see Methods). We then slightly reduced the mean TC synaptic

weight to yield reasonable excitatory responses in the control condition (see Methods),
before inspecting the responses to adapted VPm inputs. For this network, evoked rates
(Fig. 8I, top, J) and synchronous spike counts (Fig. 8K, left) for excitatory neurons were
only modestly adapted compared to the model with differential TC connectivity, and the
degree of adaptation more closely matched that of the inhibitory population (Fig. 8J, K,
right). In other words, the excitatory population was less sensitive to VPm adaptation
when differential TC connectivity was removed. We used an additional set of models to
further assess the relative importance of each component of the differential TC
connectivity in the original model. While each component contributed, we found that the
degree of excitatory adaptation was most sensitive to differences in TC synaptic latencies
(not shown). Importantly, imposing identical TC connectivity did not entirely remove
differential adaptation, as the intrinsic inhibitory neuronal properties and dense
intracortical connectivity also helped make them comparatively robust to losses in
synchronous spiking in both thalamic and cortical neurons excitatory neurons (not
shown). This model thus demonstrates the role of robust feedforward inhibition – due in
part to differential TC connectivity – in shaping the adaptation of cortical excitatory
neurons.

Taken together, these modeling results support our hypotheses that the profound
adaptation of cortical RS cells during wakefulness represents a loss of synchronous
feature-evoked thalamic spikes, in conjunction with strong feedforward inhibition that is
comparatively robust to this decrease in feedforward thalamic drive.

1111

DISCUSSION

To determine the nature of rapid cortical sensory adaptation during wakefulness, we
recorded single-unit activity in S1 of the awake, head-fixed mouse while presenting
punctate sensory features either in isolation, or embedded in a persistent background
stimulus. To elucidate the mechanistic basis of cortical adaptation, we recorded from
putative excitatory and inhibitory cortical neurons and the lemniscal inputs to S1, while
employing a battery of additional manipulations across the thalamocortical circuit. This
approach, in conjunction with a thalamocortical network model, allowed us to infer the
contributions from thalamic adaptation, thalamocortical synaptic depression, and
intracortical mechanisms.

Previous in vitro and anesthetized work has clearly demonstrated profound rapid sensory
adaptation in sensory cortex (Lesica et al., 2007; Heiss et al., 2008; Ganmor et al., 2010;
Wang et al., 2010; Cohen-Kashi Malina et al., 2013; Ollerenshaw et al., 2014; Zheng et
al., 2015; Kheradpezhouh et al., 2017), which is thought to represent the net effects on
the circuit of elevated firing rates. Yet because baseline cortical firing rates are elevated
during wakefulness compared to the anesthetized state (Greenberg et al., 2008; Vizuete

et al., 2012; Aasebø et al., 2017), it remains an open question how much room is left for
fine-tuning by the sensory environment (Castro-Alamancos, 2004). Here, we
demonstrate that cortical sensory responses can indeed be profoundly adapted during
wakefulness. The adaptive decrease in theoretical feature detectability and synchronous
firing of putative cortical excitatory neurons suggest that downstream targets of S1 will be
substantially less driven in the adapted state, which predicts a decrease in perceived
feature intensity and a loss of detectability. These observations are consistent with
previous behavioral work in rats, which demonstrated changes in perceptual reporting
following repetitive whisker stimulation (Ollerenshaw et al., 2014; Waiblinger et al., 2015).
In other words, this study supports previous in vitro and anesthetized work suggesting
cortical response adaptation could underlie perceptual adaptation.

Our observations are also consistent with elements of two previous studies in awake rats,
which demonstrated adaptation of S1 spiking (Musall et al., 2014) and LFP deflections
(Castro-Alamancos, 2004) during presentation of feature sequences across a range of
animal states. Interestingly, Castro-Alamancos (2004) showed that the extent of
adaptation varied across behavioral states, from profound during quiescence to nearly
absent in the very early stages of training in an active avoidance task (i.e., likely during
states of hyper-arousal). Musall et al. (2014), on the other hand, showed that the degree
of cortical response adaptation was largely unchanged by task engagement. This
suggests that sensory adaptation is likely relevant across much of the continuum of
behavioral states, and that brain state gates adaptation, but future work is needed to more
fully address the relationship between behavioral state and adaptation. Active sensation
is a particularly important and complex scenario: whisker self-motion may invoke sensory
adaptation via bottom-up inputs due to re-afference (Yu et al., 2016) and low-amplitude
whisker deflections resulting from surface contacts, but is also associated with top-down
modulation of VPm and S1 state (Poulet et al., 2012; Poulet and Crochet, 2019). While
our study provides insight into how isolated bottom-up inputs may adapt sensory
representations and percepts, it is not clear how whisking-related state modulation will
gate this process.

We next sought to identify the mechanistic basis for S1 response adaptation. One body
of literature implicates thalamocortical synaptic depression (Castro-Alamancos and
Oldford, 2002; Chung et al., 2002; Gabernet et al., 2005; Cruikshank et al., 2010), while
our previous work points to adaptation of thalamic spike timing (Wang et al., 2010;
Ollerenshaw et al., 2014; Whitmire et al., 2016), but both viewpoints have originated
largely from in vitro or anesthetized preparations. Our results suggest cortical response
adaptation is largely due to thalamic response adaptation in the regime we explored here.
First, adaptation profoundly reduced single-unit bursting and the rate of feature-evoked
synchronous spikes in VPm, likely due to a combination of persistent VPm depolarization
(Whitmire et al., 2016) and adapted inputs from trigeminal neurons (Castro-Alamancos,

2002; Timofeeva and Lavalle, 2003; Ganmor et al., 2010), which predicts attenuated
cortical firing (Swadlow and Gusev, 2001; Bruno and Sakmann, 2006; Wang et al., 2010;
Ollerenshaw et al., 2014). Further, optogenetic elevation of baseline VPm rates did not
adapt S1 feature responses, and background sensory stimulation had little effect on S1
responses to direct TC terminal stimulation. Finally, our modeling demonstrated that
modest reductions in synchronous VPm spiking alone predicted profound cortical
adaptation. Taken together, these results therefore demonstrate for the first time in the
awake animal the sensitivity of cortex to thalamic spike timing in the context of sensory
adaptation, and suggest that synaptic depression contributes little to the observed S1
feature response attenuation.

This apparent lack of TC synaptic depression appears to contradict the results of previous
anesthetized and in vitro studies (Castro-Alamancos and Oldford, 2002; Chung et al.,
2002; Gabernet et al., 2005; Cruikshank et al., 2010). We believe that this reflects a
difference in the strength of adapting stimuli. Specifically, these previous studies used
adapting sequences of high-velocity, punctate sensory stimuli and/or electrical
stimulation. Here, we employed a relatively low-amplitude white noise adapting stimulus,
to emulate small-amplitude whisker micromotions that occur during whisking across
textured surfaces (Jadhav and Feldman, 2010). In terms of total power (Zheng et al.,
2015), this white noise stimulus is many times weaker than the adapting stimuli used in
many previous studies (not shown). This likely adapts the TC circuit to a lesser degree
than punctate stimulus trains, which may explain the apparent contradiction with previous
work. This may also explain why we did not observe more pronounced adaptation of FS
units, which has been shown to reflect stronger adaptation of TC synapses onto inhibitory
neurons (Gabernet et al., 2005). It is possible that stronger adapting stimuli will generally
engage a broader range of adaptive mechanisms than we observed here, with
thalamocortical/intracortical synaptic depression playing more prominent roles in regimes
of more profound adaptation. It is also unclear whether sensory white noise presented
during wakefulness can alter aspects of cortical sensory coding not explored here (such
as input-output relationships (Maravall et al., 2007) and response tuning curves (Wang et
al., 2010)), and whether adaptation of thalamic synchrony is sufficient to explain such
changes. Future work probing a range of adapting stimulus strengths and effects can
investigate such questions.

While thalamic feature response adaptation appeared necessary for S1 adaptation, it did
not explain the differential adaptation of RS and FS cells during wakefulness. In other
words, RS and FS cells provided different read-outs of feature-evoked thalamic spiking.
In contrast, RS feature responses were no more adapted than FS rates under isoflurane
anesthesia, which has been shown to disproportionately weaken cortical inhibition
(Haider et al., 2013; Taub et al., 2013). This suggested to us that feedforward inhibition
contributed to the adaptation of RS cells during wakefulness. We explored this possibility

with a network model, in which we implemented cell-type-specific TC connectivity
motivated by previous experimental work (Bruno and Simons, 2002). We found that S1
response adaptation did largely reflect a loss of synchronous VPm spikes, but that the
profound and differential adaptation of excitatory neurons also required several additional
mechanisms, including cell-type-specific TC connectivity. Taken together, these
experimental and modeling results suggest a thalamocortical circuit basis for the
observed S1 adaptation, involving a profound loss of synchronous feedforward excitation,
and a comparatively modest decrease in dampening feedforward inhibition.

This adaptive shift in the feedforward E/I balance toward inhibition has implications for
cortical function and perception beyond attenuation of response amplitudes and
perceived stimulus intensities. For example, previous experimental work has
demonstrated that the relative strength and/or timing of cortical excitation and inhibition
contributes to the direction-selectivity (Wilent and Contreras, 2005) and receptive field
properties (Kyriazi and Simons, 1993; Bruno and Simons, 2002; Ramirez et al., 2014) of
excitatory neurons, maintains relatively low excitatory firing rates during bouts of whisking
(Yu et al., 2016; Gutnisky et al., 2017), shapes the "window of integration" during which
excitatory neurons integrate excitatory synaptic inputs and depolarize toward threshold
(Gabernet et al., 2005; Wilent and Contreras, 2005), and generally serves to "dampen"
thalamic-evoked spiking in the excitatory subnetwork (Pinto et al., 2003). Thalamocortical
adaptation exists on a continuum (Wang et al., 2010; Zheng et al., 2015), and more
moderate levels of adaptation than we imposed here may result in moderately attenuated
excitatory firing that is sharpened in space and time by comparatively non-adapted
inhibition, resulting in more faithful spatiotemporal cortical representations of complex
sensory stimuli. Future experiments exploring regimes of weaker adaptation and using
more complex single- and multi-whisker stimulation can explore these possibilities.

In summary, these results demonstrate the effects of rapid sensory adaptation on the
early sensory pathway during wakefulness, culminating in profound adaptation of primary
sensory cortex, which likely underlies perceptual adaptation on this timescale. Further,
they highlight the relative importance of thalamic gating in establishing cortical adaptation,
through population timing control of thalamic drive and the differential engagement of the
inhibitory cortical sub-population.

1237

1238

References

Aasebø IEJJ, Lepperød ME, Stavrinou M, Nøkkevangen S, Einevoll G, Hafting T, Fyhn M (2017) Temporal Processing in the Visual Cortex of the Awake and Anesthetized Rat. Eneuro 4:ENEURO.0059-17.2017.

Anstis S, Verstraten FAJ, Mather G (1998) The motion aftereffect. Trends Cogn Sci 2:111–117.

Avermann M, Tomm C, Mateo C, Gerstner W, Petersen CCH (2012) Microcircuits of excitatory and inhibitory neurons in layer 2/3 of mouse barrel cortex. J Neurophysiol 107:3116–3134.

Barlow HB (1961) Possible Principles Underlying the Transformations of Sensory Messages. In: Sensory Communication (Rosenblith WA, ed), pp 217–234. The MIT Press.

Barthó P, Slézia A, Mátyás F, Faradzs-Zade L, Ulbert I, Harris KD, Acsády L (2014) Ongoing network state controls the length of sleep spindles via inhibitory activity. Neuron 82:1367–1379.

Bestelmeyer PEG, Rouger J, DeBruine LM, Belin P (2010) Auditory adaptation in vocal affect perception. Cognition 117:217–223.

Blakemore C, Campbell FW (1969) On the existence of neurones in the human visual system selectively sensitive to the orientation and size of retinal images. J Physiol 203:237–260.

Blakemore C, Nachmias J (1971) The orientation specificity of two visual after-effects. J Physiol 213:157–174.

Boloori A-R, Jenks RA, Desbordes G, Stanley GB (2010) Encoding and Decoding Cortical Representations of Tactile Features in the Vibrissa System. J Neurosci 30:9990–10005.

Boloori A-R, Stanley GB (2006) The Dynamics of Spatiotemporal Response Integration in the Somatosensory Cortex of the Vibrissa System. J Neurosci 26:3767–3782.

Bolus MF, Willats AA, Rozell CJ, Stanley GB (2020) State-space optimal feedback control of optogenetically driven neural activity. J Neural Eng:0–31.

Brecht M, Sakmann B (2002) Whisker maps of neuronal subclasses of the rat ventral posterior medial thalamus, identified by whole-cell voltage recording and morphological reconstruction. J Physiol:495–515.

Britten K, Shadlen M, Newsome W, Movshon J (1992) The analysis of visual motion: a comparison of neuronal and psychophysical performance. J Neurosci 12:4745–4765.

Bruno RM (2011) Synchrony in sensation. Curr Opin Neurobiol 21:701–708.

Bruno RM, Sakmann B (2006) Cortex is driven by weak but synchronously active thalamocortical synapses. Science 312:1622–1627.

Bruno RM, Simons DJ (2002) Feedforward mechanisms of excitatory and inhibitory
cortical receptive fields. J Neurosci 22:10966–10975.

Castro-Alamancos MA (2002) Properties of Primary Sensory (Lemniscal) Synapses in
the Ventrobasal Thalamus and the Relay of High-Frequency Sensory Inputs. J
Neurophysiol 87:946–953.

Castro-Alamancos MA (2004) Absence of Rapid Sensory Adaptation in Neocortex
during Information Processing States. Neuron 41:455–464.

Castro-Alamancos MA, Oldford E (2002) Cortical sensory suppression during arousal is
due to the activity-dependent depression of thalamocortical synapses. J Physiol
541:319–331.

Chung S, Li X, Nelson SB (2002) Short-term depression at thalamocortical synapses
contributes to rapid adaptation of cortical sensory responses in vivo. Neuron
34:437–446.

Cohen-Kashi Malina K, Jubran M, Katz Y, Lampl I, Ebsch C, Rosenbaum R (2013)
Imbalance between Excitation and Inhibition in the Somatosensory Cortex
Produces Postadaptation Facilitation Graham LJ, ed. J Neurosci 33:8463–8471.

Cossell L, Iacaruso MF, Muir DR, Houlton R, Sader EN, Ko H, Hofer SB, Mrsic-Flogel
TD (2015) Functional organization of excitatory synaptic strength in primary visual
cortex. Nature 518:399–403.

Cruikshank SJ, Lewis TJ, Connors BW (2007) Synaptic basis for intense
thalamocortical activation of feedforward inhibitory cells in neocortex. Nat Neurosci
10:462–468.

Cruikshank SJ, Urabe H, Nurmikko A V., Connors BW (2010) Pathway-Specific
Feedforward Circuits between Thalamus and Neocortex Revealed by Selective
Optical Stimulation of Axons. Neuron 65:230–245.

Erb J, Henry MJ, Eisner F, Obleser J (2013) The brain dynamics of rapid perceptual
adaptation to adverse listening conditions. J Neurosci 33:10688–10697.

Gabernet L, Jadhav SP, Feldman DE, Carandini M, Scanziani M (2005) Somatosensory
integration controlled by dynamic thalamocortical feed-forward inhibition. Neuron
48:315–327.

Ganmor E, Katz Y, Lampl I (2010) Intensity-Dependent Adaptation of Cortical and
Thalamic Neurons Is Controlled by Brainstem Circuits of the Sensory Pathway.
Neuron 66:273–286.

Gentet LJ, Avermann M, Matyas F, Staiger JF, Petersen CCHH (2010) Membrane
potential dynamics of GABAergic neurons in the barrel cortex of behaving mice.
Neuron 65:422–435.

Gerfen CR, Paletzki R, Heintz N (2013) GENSAT BAC cre-recombinase driver lines to
study the functional organization of cerebral cortical and basal ganglia circuits.
Neuron 80:1368–1383.

Ghodrati M, Zavitz E, Rosa MGP, Price NSC (2019) Contrast and luminance adaptation alter neuronal coding and perception of stimulus orientation. Nat Commun 10.

Greenberg DS, Houweling AR, Kerr JND (2008) Population imaging of ongoing neuronal activity in the visual cortex of awake rats. Nat Neurosci 11:749–751.

Guo W, Clause AR, Barth-Maron A, Polley DB (2017) A Corticothalamic Circuit for Dynamic Switching between Feature Detection and Discrimination. Neuron 95:180-194.e5.

Gutnisky DA, Yu J, Hires SA, To MS, Bale MR, Svoboda K, Golomb D (2017) Mechanisms underlying a thalamocortical transformation during active tactile sensation.

Haider B, Häusser M, Carandini M (2013) Inhibition dominates sensory responses in the awake cortex. Nature 493:97–100.

Hartings JA, Temereanca S, Simons DJ (2003) Processing of Periodic Whisker Deflections by Neurons in the Ventroposterior Medial and Thalamic Reticular Nuclei. J Neurophysiol 90:3087–3094.

Heiss JE, Katz Y, Ganmor E, Lampl I (2008) Shift in the Balance between Excitation and Inhibition during Sensory Adaptation of S1 Neurons. J Neurosci 28:13320–13330.

Hoseini MS, Wright NC, Xia J, Clawson W, Shew W, Wessel R (2019) Dynamics and sources of response variability and its coordination in visual cortex. Vis Neurosci 36:E013.

Isett BR, Feasel SH, Lane MA, Feldman DE (2018) Slip-Based Coding of Local Shape and Texture in Mouse S1. Neuron 97:418-433.e5.

Jadhav SP, Feldman DE (2010) Texture coding in the whisker system. Curr Opin Neurobiol 20:313–318.

Jadhav SP, Wolfe J, Feldman DE (2009) Sparse temporal coding of elementary tactile features during active whisker sensation. 12.

Khatri V, Hartings JA, Simons DJ (2004) Adaptation in thalamic barreloid and cortical barrel neurons to periodic whisker deflections varying in frequency and velocity. J Neurophysiol 92:3244–3254.

Kheradpezhouh E, Adibi M, Arabzadeh E (2017) Response dynamics of rat barrel cortex neurons to repeated sensory stimulation. Sci Rep 7:1–10.

Kimura F, Itami C, Ikezoe K, Tamura H, Fujita I, Yanagawa Y, Obata K, Ohshima M (2010) Fast activation of feedforward inhibitory neurons from thalamic input and its relevance to the regulation of spike sequences in the barrel cortex. J Physiol 588:2769–2787.

Kodandaramaiah SB, Holst GL, Wickersham IR, Singer AC, Franzesi GT, McKinnon ML, Forest CR, Boyden ES (2016) Assembly and operation of the autopatcher for

automated intracellular neural recording in vivo. Nat Protoc 11:634–654.

Kyriazi H, Simons D (1993) Thalamocortical response transformations in simulated
whisker barrels. J Neurosci 13:1601–1615.

Lampl I, Katz Y (2017) Neuronal adaptation in the somatosensory system of rodents.
Neuroscience 343:66–76.

Lefort S, Tomm C, Floyd Sarria JC, Petersen CCH (2009) The Excitatory Neuronal
Network of the C2 Barrel Column in Mouse Primary Somatosensory Cortex.
Neuron 61:301–316.

Lesica NA, Jin J, Weng C, Yeh C-I, Butts DA, Stanley GB, Alonso J-M (2007)
Adaptation to Stimulus Contrast and Correlations during Natural Visual Stimulation.
Neuron 55:479–491.

Lesica NA, Weng C, Jin J, Yeh C-II, Alonso J-MM, Stanley GB (2006) Dynamic
Encoding of Natural Luminance Sequences by LGN Bursts Gallant J, ed. PLoS Biol
4:e209.

Liew JY, Pala A, Whitmire CJ, Stoy WM, Forest CR, Stanley GB (2020) Inferring
Monosynaptic Connectivity Across Brain Structures In-Vivo.

Litwin-Kumar A, Doiron B (2012) Slow dynamics and high variability in balanced cortical
networks with clustered connections. Nat Neurosci 15:1498–1505.

Liu C, Foffani G, Scaglione A, Aguilar J, Moxon KA (2017) Adaptation of thalamic
neurons provides information about the spatiotemporal context of stimulus history.
J Neurosci:0637–17.

Llinás RR, Steriade M (2006) Bursting of thalamic neurons and states of vigilance. J
Neurophysiol 95:3297–3308.

Lu SM, Guido W, Sherman SM (1992) Effects of membrane voltage on receptive field
properties of lateral geniculate neurons in the cat: contributions of the low-threshold
Ca2+ conductance. J Neurophysiol 68:2185–2198.

Maravall M, Alenda A, Bale MR, Petersen RS (2013) Transformation of Adaptation and
Gain Rescaling along the Whisker Sensory Pathway Solomon SG, ed. PLoS One
8:e82418.

Maravall M, Petersen RS, Fairhall AL, Arabzadeh E, Diamond ME (2007) Shifts in
Coding Properties and Maintenance of Information Transmission during Adaptation
in Barrel Cortex Ebner FF, ed. PLoS Biol 5:e19.

McCormick D (1992) Cellular mechanisms underlying cholinergic and noradrenergic
modulation of neuronal firing mode in the cat and guinea pig dorsal lateral
geniculate nucleus. J Neurosci 12:278–289.

McCormick DA, Connors BW, Lighthall JW, Prince DA (1985) Comparative
electrophysiology of pyramidal and sparsely spiny stellate neurons of the
neocortex. J Neurophysiol 54:782–806.

McGinley MJ, David S V., McCormick DA (2015) Cortical Membrane Potential Signature of Optimal States for Sensory Signal Detection. Neuron 87:179–192.

Mcginley MJ, Vinck M, Reimer J, Batista-brito R, Zagha E, Cadwell CR, Tolias AS, Cardin JA, Mccormick DA (2015) Waking State: Rapid Variations Modulate Neural and Behavioral Responses. Neuron 87:1143–1161.

Musall S et al. (2014) Tactile frequency discrimination is enhanced by circumventing neocortical adaptation. Nat Neurosci 17:1567–1573.

Niell CM, Stryker MP (2010) Modulation of visual responses by behavioral state in mouse visual cortex. Neuron 65:472–479.

Ollerenshaw DR, Zheng HJV, Millard DC, Wang Q, Stanley GB (2014) The Adaptive Trade-Off between Detection and Discrimination in Cortical Representations and Behavior. Neuron 81:1152–1164.

Pala A, Petersen CCH (2015) InVivo Measurement of Cell-Type-Specific Synaptic Connectivity and Synaptic Transmission in Layer 2/3 Mouse Barrel Cortex. Neuron 85:68–75.

Pinto DJ, Hartings JA, Brumberg JC, Simons DJ (2003) Cortical damping: Analysis of thalamocortical response transformations in rodent barrel cortex. Cereb Cortex 13:33–44.

Pinto DJD, Brumberg JCJ, Simons DJDJ (2000) Circuit dynamics and coding strategies in rodent somatosensory cortex. J Neurophysiol 83:1158–1166.

Poulet JFA, Crochet S (2019) The cortical states of wakefulness. Front Syst Neurosci 12:1–18.

Poulet JFAA, Fernandez LMJJ, Crochet S, Petersen CCHH (2012) Thalamic control of cortical states. Nat Neurosci 15:370–372.

Ramirez A, Pnevmatikakis EA, Merel J, Paninski L, Miller KD, Bruno RM (2014) Spatiotemporal receptive fields of barrel cortex revealed by reverse correlation of synaptic input. Nat Neurosci 17:866–875.

Reimer J, Froudarakis E, Cadwell CR, Yatsenko D, Denfield GH, Tolias AS (2014) Pupil Fluctuations Track Fast Switching of Cortical States during Quiet Wakefulness. Neuron 84:355–362.

Reimer J, Mcginley MJ, Liu Y, Rodenkirch C, Wang Q, McCormick DA, Tolias AS (2016) Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex. Nat Commun 7:1–7.

Reinagel P, Godwin D, Sherman SM, Koch C (1999) Encoding of Visual Information by LGN Bursts. J Neurophysiol 81:2558–2569.

Ritt JT, Andermann ML, Moore CI (2008) Embodied Information Processing: Vibrissa Mechanics and Texture Features Shape Micromotions in Actively Sensing Rats. Neuron 57:599–613.

Safaai H, von Heimendahl M, Sorando JM, Diamond ME, Maravall M (2013)
Coordinated Population Activity Underlying Texture Discrimination in Rat Barrel
Cortex. J Neurosci 33:5843–5855.

Sederberg AJ, Pala A, Zheng HJ V., He BJ, Stanley GB (2019) State-aware detection of
sensory stimuli in the cortex of the awake mouse Macke JH, ed. PLOS Comput Biol
15:e1006716.

Sermet BS, Truschow P, Feyerabend M, Mayrhofer JM, Oram TB, Yizhar O, Staiger JF,
Petersen CCH (2019) Pathway-, layer-and cell-type-specific thalamic input to
mouse barrel cortex. Elife 8:1–28.

Sherman SM (2001) A wake-up call from the thalamus. Nat Neurosci 4:344–346.

Smith MW, Faulkner A (2006) Perceptual adaptation by normally hearing listeners to a
simulated "hole" in hearing. J Acoust Soc Am 120:4019–4030.

Speed A, Del Rosario J, Burgess CP, Haider B (2019) Cortical State Fluctuations
across Layers of V1 during Visual Spatial Perception. Cell Rep 26:2868-2874.e3.

Stoy WM, Yang B, Kight A, Wright NC, Borden PY, Stanley GB, Forest CR (2020)
Compensation of physiological motion enables high-yield whole-cell recording in
vivo. J Neurosci Methods:109008.

Stringer C, Pachitariu M, Steinmetz N, Reddy CB, Carandini M, Harris KD (2019)
Spontaneous behaviors drive multidimensional, brainwide activity. Science (80-)
364:eaav7893.

Stüttgen MC, Schwarz C (2008) Psychophysical and neurometric detection
performance under stimulus uncertainty. Nat Neurosci 11:1091–1099.

Suzuki M, Larkum ME (2020) General Anesthesia Decouples Cortical Pyramidal
Neurons. Cell 180:666-676.e13.

Swadlow HA (2002) Thalamocortical control of feed-forward inhibition in awake
somatosensory "barrel" cortex. Philos Trans R Soc B Biol Sci 357:1717–1727.

Swadlow HA, Gusev AG (2001) The impact of "bursting" thalamic impulses at a
neocortical synapse. Nat Neurosci 4:402–408.

Tannan V, Simons S, Dennis RG, Tommerdahl M (2007) Effects of adaptation on the
capacity to differentiate simultaneously delivered dual-site vibrotactile stimuli. Brain
Res 1186:164–170.

Taub AH, Katz Y, Lampl I (2013) Cortical Balance of Excitation and Inhibition Is
Regulated by the Rate of Synaptic Activity. J Neurosci 33:14359–14368.

Timofeeva E, Lavalle P (2003) The Relay of High-Frequency Sensory Signals in the
Whisker-to-Barreloid Pathway. J Neurosci 23:6778–6787.

Usrey WM, Alonso J-M, Reid RC (2000) Synaptic Interactions between Thalamic Inputs
to Simple Cells in Cat Visual Cortex. J Neurosci 20:5461–5467.

Vizuete JA, Pillay S, Diba K, Ropella KM, Hudetz AG (2012) Monosynaptic functional
connectivity in cerebral cortex during wakefulness and under graded levels of
anesthesia. Front Integr Neurosci 6:1–11.

Waiblinger C, Brugger D, Whitmire CJ, Stanley GB, Schwarz C (2015) Support for the
slip hypothesis from whisker-related tactile perception of rats in a noisy
environment. Front Integr Neurosci 9:e53.

Waiblinger C, Whitmire CJ, Sederberg A, Stanley GB, Schwarz C (2018) Primary
Tactile Thalamus Spiking Reflects Cognitive Signals. J Neurosci 38:4870–4885.

Wang Q, Webber RM, Stanley GB (2010) Thalamic synchrony and the adaptive gating
of information flow to cortex. Nat Neurosci 13:1534–1541.

Webber RM, Stanley GB (2004) Nonlinear Encoding of Tactile Patterns in the Barrel
Cortex. J Neurophysiol 91:2010–2022.

Webber RM, Stanley GB (2006) Transient and Steady-State Dynamics of Cortical
Adaptation. J Neurophysiol 95:2923–2932.

Whitmire CJ, Stanley GB (2016) Rapid Sensory Adaptation Redux: A Circuit
Perspective. Neuron 92:298–315.

Whitmire CJ, Waiblinger C, Schwarz C, Stanley GB (2016) Information Coding through
Adaptive Gating of Synchronized Thalamic Bursting. Cell Rep 14:795–807.

Wilent WB, Contreras D (2005) Dynamics of excitation and inhibition underlying
stimulus selectivity in rat somatosensory cortex. Nat Neurosci 8:1364–1370.

Wolfe J, Hill DN, Pahlavan S, Drew PJ, Kleinfeld D, Feldman DE (2008) Texture coding
in the rat whisker system: Slip-stick versus differential resonance. PLoS Biol
6:1661–1677.

Wright NC, Hoseini MS, Wessel R (2017a) Adaptation modulates correlated
subthreshold response variability in visual cortex. J Neurophysiol 118:1257–1269.

Wright NC, Hoseini MS, Yasar TB, Wessel R (2017b) Coupling of synaptic inputs to
local cortical activity differs among neurons and adapts after stimulus onset. J
Neurophysiol 118:3345–3359.

Yu J, Gutnisky DA, Hires SA, Svoboda K (2016) Layer 4 fast-spiking interneurons filter
thalamocortical signals during active somatosensation. Nat Neurosci 19:1647–
1657.

Yu J, Hu H, Agmon A, Svoboda K (2019) Recruitment of GABAergic Interneurons in the
Barrel Cortex during Active Tactile Behavior. Neuron 104:412-427.e4.

Zheng HJ V., Wang Q, Stanley GB (2015) Adaptive shaping of cortical response
selectivity in the vibrissa pathway. J Neurophysiol 113:3850–3865.

Zuo Y, Safaai H, Notaro G, Mazzoni A, Panzeri S, Diamond ME (2015) Complementary
Contributions of Spike Timing and Spike Rate to Perceptual Decisions in Rat S1
and S2 Cortex. Curr Biol 25:357–363.

1505

1506 ***Figure Captions***

1507

Figure 1. S1 exhibits sensory adaptation during wakefulness, and regular-spiking
neurons are more profoundly adapted than fast-spiking neurons. A. Experimental
setup. We recorded in S1 of the awake, head-fixed mouse while presenting precise
single-whisker stimulation. "Sawtooth" punctate sensory features were delivered either
in isolation (control condition) or embedded in sensory white noise (adapted condition).
B. Top: Grand mean +/- SEM waveforms for all well-isolated fast-spiking (FS, blue, N =
95) and regular-spiking (RS, red, N = 119) significantly responsive single-units recorded
in S1 of awake mice (see Methods). Bottom: distribution of mean waveform widths (time
from trough to subsequent peak) for all units, with color denoting RS and FS designation.
C. Grand PSTHs for regular-spiking (RS, putative excitatory, top) and fast-spiking (FS,
putative inhibitory, bottom) from a subset of all recording sessions (for which we used the
same white noise stimulus). **D.** Top: grand-average mean (+/- SEM) rates for
spontaneous activity (i.e., no sensory stimulation) and "early" (0 – 200 ms) and "late" (500
– 700 ms) windows following onset of sensory white noise (with *** indicating $p < 0.0005$,
Wilcoxon signed-rank test). RS: spontaneous rate = 6.12 +/- 0.60 Hz; white noise early
rate = 9.75 +/- 0.80 Hz; white noise late rate = 7.77 +/- 0.76 Hz, mean +/- SEM. Spont
vs. white noise early: W = 274, $p = 4.59 \times 10^{-10}$; spont vs. white noise late: W = 799.5, p
$= 5.92 \times 10^{-4}$; white noise early vs. white noise late: W = 425, $p = 9.95 \times 10^{-9}$, Wilcoxon
signed-rank test, N = 119 units from 19 recording sessions, FS: spontaneous rate = 12.59
+/- 1.29 Hz, white noise early rate = 31.73 +/- 3.29 Hz; white noise late rate = 24.04 +/-
2.6 Hz, mean +/- SEM. Spont vs. white noise early: W = 4.0, $p = 9.13 \times 10^{-15}$; spont vs.
white noise late: W = 237.5, $p = 3.34 \times 10^{-11}$; white noise early vs. white noise late: W =
220.0, $p = 1.88 \times 10^{-11}$, Wilcoxon signed-rank test, N = 95 units from 19 sessions. Bottom:
relative whisker motion in the absence ("spont") and presence ("white noise") of white
noise whisker stimulation, from five recording sessions with simultaneous
electrophysiology and whisker videography (see Methods). **E.** Grand PSTHs for all
responsive RS (top, N = 119) and FS (bottom, N = 95) units, for two punctate stimulus

velocities. **F.** Across-neuron mean (+/- SEM) firing rates for all responsive neurons, for 300 deg/s (left) and 900 deg/s (right) punctate stimuli (*: $0.01 \leq p < 0.05$; ***: $p < 0.001$, Wilcoxon singed-rank test). RS 300 deg/s mean +/- SEM control: 17.08 +/- 1.02, adapted: 11.36 +/- 1.01, 33.5% decrease, W = 523.0, p = 1.03×10^{-15}, Wilcoxon signed-rank test, N = 119 units from 19 recording sessions; 900 deg/s control: 17.19 +/- 1.25 Hz, adapted: 9.92 +/- 1.08 Hz, 42.3% decrease, W = 43.0, p = 1.47×10^{-8}, N = 49 units from 8 sessions; FS: 300 deg/s control: 41.72 +/- 3.85 Hz, adapted: 34.1 +/- 3.39 Hz, 18.3% decrease, W = 785.5, p = 2.90×10^{-8}, N = 95 units from 19 sessions; 900 deg/s control: 42.88 +/- 5.61 Hz, adapted: 40.03 +/- 6.43 Hz, 6.6% decrease, W = 156.5, p = 0.19, N = 29 units from 8 sessions. **G.** Population median (+/- SEM) normalized adapted responses for all responsive RS (red) and FS (blue) neurons (see Methods; ***: $p < 0.001$, Kruskal-Wallis test). RS 300 deg/s median normed adapted response = 0.61, FS median normed adapted response = 0.77, H = 16.94, p = 3.85×10^{-5}, Kruskal-Wallis test; 900 deg/s RS median normed adapted response = 0.49, FS median normed adapted response = 0.87, H = 14.91, p = 1.13×10^{-4}. **H.** Population median normalized adapted responses (+/- SEM) by binned cortical depth (see Methods).

Figure 2. Adaptation reduces the theoretical detectability of punctate sensory stimuli. A. Baseline and feature-evoked population spike rate distributions for one example RS unit (see Methods). For this unit (qualitatively representative of the average), adaptation decreased the mean population response rate, increasing the overlap of the baseline and feature-evoked distributions (right). **B.** Receiver operator characteristic (ROC) curves for example unit in (B), for the control (black line) and adapted (gray line) conditions, and associated area under ROC curve (AUROC) values. **C.** Grand average mean (+/- SEM) theoretical detectability (AUROC) vs. stimulus condition for all significantly responsive RS units (***: $p < 0.0005$, Wilcoxon signed-rank test). 300 deg/s control: mean +/- SEM AUROC = 0.81 +/- 0.02, adapted: 0.66 +/- 0.02, hybrid: 0.66 +/- 0.02, control vs. adapted: W = 1936.0, p = 2.09×10^{-11}; control vs. hybrid: W = 1154.0, p = 1.91×10^{-16}; adapted vs. hybrid: W = 5243.0, p = 0.81, Wilcoxon signed-rank test, N = 119 units from 19 recording sessions; 900 deg/s control: mean +/- SEM AUROC = 0.89

1565 +/- 0.02, adapted: 0.73 +/- 0.03, hybrid: 0.71 +/- 0.03; control vs. adapted: W = 166.0, p

1566 = 3.18 x 10^{-6}; control vs. hybrid: W = 106, p = 1.78 x 10^{-7}; adapted vs. hybrid: W = 528.0,

1567 p = 0.21, N = 49 units from 8 recording sessions.

1568

Figure 3.   Adaptation increases response latency and reduces pairwise

synchronous spiking in S1. A. Grand mean onset times for RS (red) and FS (blue)

neurons, for control and adapted responses to 300 deg/s stimulus (***: p < 0.001; single

gray bar: p ≥ 0.05, Wilcoxon signed-rank test, control vs. adapted). 300 deg/s RS mean

+/ SEM control: T_{onset} = 8.81 +/- 0.53 ms, adapted: 15.31 +/- 0.98 ms, 73.9% increase, W

= 397.0, p = 3.11 x 10^{-11}, Wilcoxon signed-rank test, N = 119 units; FS control: T_{onset} =

6.69 +/- 0.37 ms, adapted: 8.95 +/- 0.7 ms, 33.9% increase, W = 245.0, p = 8.11 x 10^{-7},

N = 95 units; 900 deg/s RS control: T_{onset} = 7.89 +/- 0.49 ms, adapted: 11.36 +/- 1.07 ms,

43.9% increase, W = 90.0, p = 1.70 x 10^{-5}, N = 49 units; FS control: T_{onset} = 5.67 +/- 0.29

ms, adapted: 6.27 +/- 0.63 ms, 10.5% increase, W = 20.5, p = 0.27, N = 29 units.   **B.**

Illustration of synchronous spike-count calculation. The grand cross-correlogram (CCG)

was constructed using all valid pairs of simultaneously-recorded S1 units, then scaled by

the number of contributing pairs, and shuffle-corrected (see Methods). The synchronous

spike count was the number of spikes in a +/- 7.5 ms window around zero lag. **C.** Grand

RS-RS (left) and FS-FS (right) CCGs for responses to 300 deg/s stimulus, for the control

(dark lines) and adapted (light lines) conditions.   Bands indicate 97.5% confidence

intervals (from re-sampling spikes with replacement, see Methods). **D.** Synchronous AP

counts for control, adapted, and "hybrid" conditions (see Methods), calculated from grand

CCGs. (Error bars indicate 97.5% confidence intervals; *: p < 0.025, re-sampling spikes

with replacement, see Methods.) RS-RS pairs: 300 deg/s mean +/- 97.5% confidence

interval control: synch AP count = 54.15 +/- 3.58 spikes/pair; adapted: 13.69 +/- 2.05

spikes/pair; hybrid: 16.61 +/- 1.67 spikes/pair; N = 189 valid pairs from 17 sessions; 900

deg/s control: 141.93 +/- 5.42 spikes/pair; adapted: 21.18 +/- 2.8 spikes/pair; hybrid:

34.87 +/- 3.84 spikes/pair; N = 55 pairs from 6 sessions. FS-FS pairs: 300 deg/s control:

191.03 +/- 7.64 spikes/pair; adapted: 106.54 +/- 7.49 spikes/pair; hybrid: 126.33 +/- 6.69

spikes/pair; N = 118 pairs from 17 sessions; 900 deg/s control: 558.27 +/- 26.03

spikes/pair; adapted: 257.15 +/- 19.67 spikes/pair; hybrid: 423.61 +/- 17.97 spikes/pair;

N = 33 pairs from 6 sessions. See Methods for definition of "valid pairs".

1597

Figure 4. S1 sensory responses are also adapted under anesthesia, but RS cells

are not differentially adapted. A. Experimental setup. S1 sensory responses were

recorded in mice lightly-anesthetized with isoflurane (see Methods). **B.** Summed spiking

activity of regular-spiking (RS, putative excitatory) and fast-spiking (FS, putative

inhibitory) from one example recording session. Each row indicates spike times of all

simultaneously-recorded RS (red) and FS (blue) neurons on a single trial. **C.** Grand-

average mean (+/- SEM) rates for spontaneous activity (i.e., no sensory stimulation) and

"early" (0 – 200 ms) and "late" (500 – 700 ms) windows following onset of sensory white

noise (with *** indicating $p < 0.0005$, Wilcoxon signed-rank test). RS: spontaneous rate

= 2.34 +/- 0.38 Hz; white noise early rate = 4.96 +/- 0.50 Hz; white noise late rate = 3.33

+/- 0.45 Hz, mean +/- SEM. Spont vs. white noise early: $W = 55$, $p = 1.13 \times 10^{-7}$; spont

vs. white noise late: $W = 191$, $p = 3.89 \times 10^{-4}$; white noise early vs. white noise late: $W =$

214, $p = 3.61 \times 10^{-4}$, Wilcoxon signed-rank test, N = 46 units from 14 recording sessions,

FS: spontaneous rate = 0.85 +/- 0.28 Hz, white noise early rate = 8.97 +/- 0.85 Hz; white

noise late rate = 5.52 +/- 1.13 Hz, mean +/- SEM. Spont vs. white noise early: $W = 0$, p

= 5.95×10^{-5}; spont vs. white noise late: $W = 6$, $p = 1.41 \times 10^{-4}$; white noise early vs. white

noise late: $W = 20$, $p = 9.02 \times 10^{-4}$, Wilcoxon signed-rank test, N = 21 units from 14

sessions. **D.** Sensory response grand PSTHs for all responsive RS (left, N = 46) and FS

(right, N = 21) units recorded under anesthesia, for 300 deg/s punctate stimulus velocity.

E. Grand mean (+/- SEM) rates for cells contributing to PSTHs in (D). RS mean +/- SEM

control: 12.27 +/- 0.88 Hz, adapted: 7.75 +/- 1.01, 36.8% decrease, $W = 184$, $p = 9.82 \times$

10^{-5}; FS control: 18.61 +/- 2.19 Hz, adapted: 11.94 +/- 2.61, 35.8% decrease, $W = 21$, p

= 0.001. **F.** Population median (+/- SEM) normalized adapted responses for all

responsive RS (red) and FS (blue) neurons (see Methods). RS 300 deg/s median normed

adapted response = 0.54, FS median normed adapted response = 0.57, $H = 0.01$, $p =$

0.91, Kruskal-Wallis test.

1624

Figure 5. Adaptation reduces tonic and burst firing rates, and synchronous spike counts, in VPm sensory responses. A. Top: experimental setup. We recorded extracellular spiking in VPm of the awake mouse, primarily using high-density silicon probes (see Methods). Bottom: criteria for classification of putative tonic (black) and burst (red) VPm spikes. **B.** Grand PSTHs for putative tonic (black) and burst (red) VPm spikes from a subset of all recording sessions. **C.** Grand-average mean (+/- SEM) rates for spontaneous activity (i.e., no sensory stimulation) and "early" (0 – 200 ms) and "late" (500 – 700 ms) windows following onset of sensory white noise (with *** indicating p < 0.0005, Wilcoxon signed-rank test). Tonic: spontaneous rate = 10.28 +/- 1.31 Hz; white noise early rate = 16.69 +/- 2.77 Hz; white noise late rate = 14.84 +/- 2.39 Hz, mean +/- SEM. Spont vs. white noise early: W = 42, p = 8.91 x 10^{-5}; spont vs. white noise late: W = 75, p = 1.12 x 10^{-3}; white noise early vs. white noise late: W = 135, p = 0.045, Wilcoxon signed-rank test; burst: spontaneous rate = 0.36 +/- 0.07 Hz, white noise early rate = 0.85 +/- 0.2 Hz; white noise late rate = 0.26 +/- 0.08 Hz, mean +/- SEM. Spont vs. white noise early: W = 65.5, p = 3.0 x 10^{-3}; spont vs. white noise late: W = 51, p = 4.65 x 10^{-3}; white noise early vs. white noise late: W = 37, p = 4.35 x 10^{-4}, Wilcoxon signed-rank test, N = 30 units from 9 sessions. **D.** Grand PSTHs for tonic (black) and burst (red) spikes from all putative VPm neurons, for 300 deg/s punctate stimulus. Note the presence of both a short-latency primary peak, and a shorter, secondary peak in tonic firing rates (Fig. 5E). This secondary peak in the grand PSTH resulted from a subset of neurons with both early and late responses – often within individual trials – and was likely evoked by the return of the whisker to resting position in the second half of the sawtooth waveform. **E.** Across-neuron mean (+/- SEM) firing rates for all putative VPm neurons (**: 0.001 ≤ p < 0.01; ***: p < 0.001, Wilcoxon singed-rank test). Tonic 300 deg/s mean +/- SEM rate control: 30.2 +/- 3.41 Hz, adapted: 24.32 +/- 2.92 Hz, 19.5% decrease, W = 83, p = 0.002, Wilcoxon signed-rank test, N = 30 units; 900 control: 29.98 +/- 3.47 Hz, adapted: 26.87 +/- 3.18 Hz, W = 44.5, p = 0.22, N = 16 units; burst: 300 deg/s control: 3.79 +/- 1.08, adapted: 1.37 +/- 0.73 Hz, 64.0% decrease, W = 47, p = 6.21 x 10^{-4}; 900 deg/s control: 5.75 +/- 3.13 Hz, adapted: 2.02 +/- 0.88, 64.9% decrease, W = 5, p = 4.6 x 10^{-3}. **F.** Grand shuffle-corrected cross-correlograms for all simultaneously-recorded putative VPm neurons, for the control (black) and adapted (gray) conditions. Bands indicate 97.5% confidence intervals (from

re-sampling spikes with replacement, see Methods). **G.** Synchronous AP counts for

control, adapted, and "hybrid" conditions (see Methods), calculated from grand CCGs.

(Error bars indicate 97.5% confidence intervals; *: p < 0.025, re-sampling spikes with

replacement, see Methods.) 300 deg/s mean +/- 97.5% confidence interval control: synch

AP count = 46.69 +/- 8.12 spikes/pair; adapted: 17.29 +/- 4.0 spikes/pair; hybrid: 20.4 +/-

4.25 spikes/pair; N = 48 valid pairs from 7 sessions; 900 deg/s control: 51.38 +/- 8.22

spikes/pair; adapted: 30.81 +/- 6.3 spikes/pair; hybrid: 31.3 +/- 5.35 spikes/pair; N = 37

pairs from 3 sessions. (See Methods for definition of "valid pairs".)

Figure 6. Optogenetic elevation of baseline VPm firing rate does not adapt S1

sensory responses. A. Experimental setup. We recorded extracellular spiking activity

from topographically-aligned VPm barreloids and S1 barrels in awake, head-fixed,

transgenic mice expressing Channelrhodopsin in VPm/VPI neurons (see Methods). **B.**

Summed spiking activity of all well-isolated, responsive putative VPm (top) and S1

(bottom) units from one example recording session. Each row indicates spike times of all

such simultaneously-recorded units in that brain region on a single trial. **C.** Grand mean

(+/- SEM) firing rates for spontaneous activity ("spont"), and during presentation of the

adapting sensory stimulus ("white noise") and optogenetic depolarization of VPm ("LED"),

for VPm (top) and S1 (bottom; *: $0.005 \leq p < 0.025$; **: $5 \times 10^{-4} \leq p < 5 \times 10^{-3}$; ***: $p < 5 \times$

10^{-4}). VPm mean +/- SEM rate spontaneous: 7.68 +/- 1.1 Hz, white noise: 11.05 +/- 1.99

Hz, LED: 14.27 +/- 2.81 Hz; spont vs. white noise, W = 37, p = 0.011, Wilcoxon signed-

rank test; spont vs. LED: W = 37, p = 0.011; white noise vs. LED: W = 102, p = 0.91, N =

20 units from 6 sessions; S1 spontaneous: 8.87 +/- 0.82 Hz, white noise: 16.32 +/- 2.05

Hz, LED: 10.03 +/- 1.05 Hz; spont vs. white noise: W = 445, p = 5.05×10^{-10}; spont vs.

LED: W = 1568.5, p = 0.11; white noise vs. LED: W = 1013, p = 8.42×10^{-5}, N = 88 units

from six sessions. **D.** Left: grand PSTHs for each stimulus condition, for VPm (top) and

S1 (bottom). Right: Grand mean (+/- SEM) firing rates for each condition. Asterisks as

in (C). VPm mean +/- SEM rate control: 29.12 +/- 4.82 Hz, adapted: 19.07 +/- 3.1 Hz,

LED: 36.44 +/- 5.35 Hz; control vs. adapted: W = 26, p = 0.003; adapted vs. LED: W =

11, p = 4.5×10^{-4}; control vs. LED: W = 55, p = 0.06. S1 control: 27.33 +/- 2.61 Hz,

adapted: 20.91 +/- 2.47 Hz, LED: 27.82 +/- 2.79 Hz; control vs. adapted: W = 502.5, p =

1.39×10^{-9}; adapted vs. LED: W = 424, p = 2.86×10^{-10}; control vs. LED: W = 1837, p =

0.89, Wilcoxon signed-rank test. **E.** Top: Grand VPm CCGs for each stimulus condition.

Bands indicate 99.9% confidence intervals (re-sampling spikes with replacement (see

Methods). Bottom: synchronous spike counts calculated from CCGs in (E, Top), for each

stimulus condition. Error bars indicate 99.9% confidence intervals (re-sampling spikes

with replacement (see Methods). Mean +/- 99.9% confidence interval synch AP count

control: 47.14 +/- 11.0 spikes/pair, LED: 51.83 +/- 12.94 spikes/pair, adapted: 20.08 +/-

8.53 spikes/pair, p ≥ 0.05 control vs. LED, p < 0.001 control vs. adapted, re-sampling

spikes with replacement, N = 36 pairs from 19 units in five sessions.

1695

Figure 7. S1 responses to direct optogenetic stimulation of TC terminals are not

adapted by sensory white noise. **A.** Experimental setup for extracellular recordings.

S1 spiking activity was recorded in mice lightly-anesthetized with isoflurane (see

Methods). These transgenic mice expressed Channelrhodopsin in VPm cell bodies,

axons, and thalamocortical (TC) axon terminals. An optic fiber positioned above the

cortical surface was used to deliver punctate optogenetic stimulation to TC terminals on

"light feature" stimulation trials (see Methods). **B.** Grand PSTHs (using all sensory-

responsive single- and multi-units, see Methods) for sensory feature (black, left) and light

feature (blue, right) trials, for control (empty histogram) and adapted (filled histogram)

conditions. Inset: number of recorded units that were significantly responsive to the

sensory feature only ("sens only"), the light feature only, both the sensory and light

features ("sens & light"), and not responsive to either ("NR"). **C.** Across-unit mean (+/-

SEM) punctate-stimulus-evoked firing rates vs stimulus condition for all responsive single-

and multi-units, for sensory feature (black, left) and light feature (blue, right) stimuli. (***:

p < 0.001, Wilcoxon singed-rank test). Sensory feature mean +/- SEM control: 18.38 +/-

1.27 Hz, adapted: 6.59 +/- 0.65 Hz, W = 39, p = 1.39×10^{-12} N = 71 units, Wilcoxon signed-

rank test. Light feature mean +/- SEM control: 11.7 +/- 0.95 Hz, adapted: 11.38 +/- 0.99

Hz, W = 987, p = 0.19. **D.** Distributions of normalized adapted responses for all valid

units (see Methods). Triangles at top denote population median values (***: p < 0.001,

Wilcoxon signed-rank test). Population median normalized adapted response = -0.29

light feature trials, 0.96 sensory feature trials, W = 199, p = 3.12×10^{-11}, Wilcoxon signed-

rank test. **E.** Normalized adapted responses to the light feature vs. across-trial mean light-evoked rate in the control condition for all 71 units that responded to both sensory and light features (with r- and p-values from Pearson correlation test). **F.** Left: experimental setup for in vivo S1 patch clamp recordings in lightly-anesthetized transgenic mice. Right: Across-trial median membrane potential responses to sensory features (black traces) and light features (blue traces), for one example S1 neuron. **G.** Properties of subthreshold responses to sensory features (left) and light features (right): subthreshold response amplitude (top), and time from stimulus onset to peak subthreshold response (T_{peak}, center) for each of the four recorded cells. Dark lines connecting pairs of data points indicate significant difference across stimulus conditions ($p < 0.05$, Wilcoxon signed-rank test), and light lines indicate non-significance.

Figure 8. A thalamocortical network model identifies synchronous VPm spikes and feedforward inhibition as key mechanisms underlying response adaptation. A. Model schematic (see Methods). **B.** Grand mean PSTHs for VPm spike times used to stimulate the network (top) and network excitatory and inhibitory neurons, for the control (empty PSTHs) and adapted (filled PSTHs) conditions. **C.** Normalized adapted responses for both simulated stimulus velocities (see Methods). For each condition, the response is defined to be the peak of the grand PSTH, and the normalized adapted response is the adapted value divided by the control value. Error bars indicate 95% confidence intervals from re-sampling neurons with replacement. **D.** Grand exc-exc (left) and inh-inh (right) CCGs for 200 randomly-selected pairs of network neurons, for the control (dark line) and adapted (lighter line) conditions. CCGs normalized to max value in control condition for visualization purposes. **E.** Top: grand CCGs for VPm inputs to model in the control (dark line) and adapted (lighter line) conditions (corresponding to PSTHs in B, top). Bottom: grand CCGs for VPm inputs in the control (dark line) and "reduced synch" (lighter line) conditions, where the "reduced synch" condition results from manual changes to drawn VPm spike times (see Methods). **F.** Same as in B, but for control and "reduced synch" simulations. **G.** Top: grand PSTHs for various choices of percent shifted VPm spikes. Bottom: normalized adapted excitatory and inhibitory

responses vs. percent of shifted VPm spikes. **H.** Model schematic for "identical TC connectivity" network (see Results and Methods). **I.** Grand excitatory (top) and inhibitory (bottom) PSTHs for "identical TC connectivity" network. **J, K.** Same as in (H, I), but for "identical TC connectivity" network.

1747

1748

1749

1750

1751

1752

1753

1754

1755

1756

1757

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A Ai32 x nsfm-Cre

anesthetized

light feature

silicon probe

S1

sensory feature

adapting sensory inputs

VPm

ChR2

sensory
light

sensory feature   light feature

ctrl   500 ms

adpt

B S1 grand PSTH (N = 71 SU + MU)

sensory feature   light feature

sens only
121

NR
16

light only
0

sens & light
71

10 Hz
10 ms

ctrl
adpt

ctrl
adpt

sensory feature onset

light feature onset

C sensory feature   light feature

mean rate (Hz)

***

ctrl   adpt   ctrl   adpt

D

probability

***

sens
light

0   1
normed adpt resp

E

normed adpt resp

r = 0.01
p = 0.94

10   20   40
mean ctrl rate (Hz)

F *in vivo patch clamp*

S1

5 mV
15 ms

ctrl
adpt

ctrl
adpt

G sensory   light

amp (mV)

T_peak (ms)

ctrl   adpt   ctrl   adpt

Figure 8