TITLE

Malaria parasite fitness in healthy human red cells is driven by common host variation

AUTHORS

Emily R Ebel[1,2], Frans A Kuypers[3], Carrie Lin[2], Dmitri A Petrov[1*], Elizabeth S Egan[2,4*]

[1]Department of Biology, Stanford University, Stanford, CA 94305

[2]Department of Pediatrics, Stanford University School of Medicine, Stanford, CA 94305

[3]Children's Hospital Oakland Research Institute, Oakland, CA, 94609

[4]Department of Microbiology & Immunology, Stanford University School of Medicine,

Stanford, CA 94305

*corresponding authors

SUMMARY

The replication of *Plasmodium falciparum* malaria parasites within red blood cells

(RBCs) causes severe disease in humans, especially in Africa. The influence of host

erythrocyte variation on parasite replication is largely uncharted, aside from a handful of

deleterious alleles like sickle cell. Here, we integrated analyses of exome sequencing, RBC

phenotyping, and parasite fitness assays on blood donated by 122 individuals, most with

African ancestry. In donors lacking alleles for hemoglobinopathies or G6PD deficiency, RBC

phenotypes including size, deformability, and hydration status explained 21-38% of the

variation in parasite growth rate. Furthermore, non-pathogenic polymorphisms in 14 RBC

proteins including *SPTA1*, *PIEZO1,* and *ATP2B4* explained 45-70% of parasite growth

variation. Interestingly, we observed little evidence for divergent selection on this variation

between Africans and Europeans. These findings suggest a model in which widespread,

non-pathogenic variation in a moderate number of genes strongly modulates *P. falciparum*

fitness in RBCs.

INTRODUCTION

Malaria caused by the replication of *Plasmodium falciparum* parasites in red blood

cells (RBCs) kills hundreds of thousands of children each year[1]. In each 48-hour cycle of

blood stage malaria, parasites must deform RBC membranes to invade them[2,3]; consume

hemoglobin and tolerate the resulting oxidative stress[4]; multiply to displace half the RBC

volume[5]; and remodel the RBC membrane to avoid immune detection[6]. Consequently,

genetic disorders that alter aspects of RBC biology are well known to influence malaria

susceptibility[7]. For example, sickle cell trait impairs parasite growth by altering

hemoglobin polymerization at low oxygen tension[8,9], while deficiency of the G6PD enzyme

involved in oxidative stress tolerance is thought to make parasitized RBCs more susceptible

to breakdown[10]. Aside from these diseases, however, the genetic basis of malaria

susceptibility remains mostly unknown. Several large genome-wide association studies

(GWAS) have recently made progress on this question by identifying a few dozen loci that

collectively explain up to 11% of the heritability of the risk of severe versus uncomplicated

malaria[11–16]. Eleven of these loci are in or near genes expressed predominantly in RBCs,

and one new GWAS variant has since been shown to regulate expression of the *ATP2B4*

calcium channel[17] and to be associated with RBC dehydration[18]. However, most other RBC

loci that have been identified by GWAS are not novel, and a functional link between *ATP2B4*

and *P. falciparum* replication has yet to be demonstrated. More discoveries about the

influence of RBC variation on malaria susceptibility are not expected from similar

GWAS[19,20], in part because severe malaria is a complex phenotype that depends on a

combination of factors from RBCs, the vascular endothelium, the immune system, and the

environment[21,22]. Alternate approaches are therefore needed to discover more genetic

variation that impacts the replication of malaria parasites in human RBCs.

RBC phenotypes like mean cell volume (MCV), hemoglobin content (HGB/MCH), and

antigenic blood type vary widely within and between human populations[23–25]. 40-90% of

the variation in common RBC indices is heritable[26–28], and GWAS have demonstrated that

these phenotypes have polygenic bases[29–33]. Average hemoglobin levels, hematocrit, and

especially RBC membrane fragility have also been shown to differ between African and

European populations, likely for genetic reasons[34–38]. Although African populations are

highly genetically diverse[39], some of these differences between Africans and Europeans are

explained by RBC disease alleles[24,35,37] that have been widely selected across Africa for

their protective effects on malaria. It remains untested whether other extensive phenotypic

and genetic diversity in RBCs, aside from these disease alleles, also influences malaria

susceptibility and has been similarly shaped by malaria selection.

Here, we approach these questions by performing exome sequencing and extensive
RBC phenotyping on blood samples from a diverse human cohort of 122 individuals. We
show that *P. falciparum* fitness varies widely among donor cells *in vitro*, with the
distributions of parasite phenotypes in 'healthy' RBCs overlapping those from RBCs
carrying classic disease alleles. We apply Lasso variable selection to identify a small set of
alleles and phenotypes that strongly predict parasite fitness outside of the context of RBC
disease, highlighting RBC membrane and dehydration genes and phenotypes as key to
modulating *P. falciparum* fitness. However, we find no evidence that non-pathogenic alleles
or phenotypes that confer parasite protection are associated with African ancestry.
Overall, these findings advance our understanding of the origin and function of common
RBC variation and suggest new targets for therapeutic intervention for malaria.
74
RESULTS
<u>Many healthy blood donors with African ancestry carry alleles for RBC disease</u>
We collected blood samples from 121 donors with no known history of blood
disorders, most of whom self-identified as having recent African ancestry (Figure 1A). As a
positive control, we also sampled a patient with hereditary elliptocytosis (HE), a polygenic
condition characterized by extremely fragile RBC membranes that strongly inhibit *P.*
falciparum growth[40–43]. We performed whole-exome sequencing, both to check for the
presence of known RBC disease alleles and to confirm the population genetic ancestry of
our donors. A principal component analysis of more than 35,000 unlinked, exomic SNPs
showed that most donors fell along a continuum from African to European ancestry, as
defined by data from the 1000 Genomes Project (Figure 1A). We found that 16% of the
healthy donors carried pathogenic hemoglobin alleles (Figure 1B), including 5
heterozygotes for hemoglobin S (HbAS), 4 heterozygotes for hemoglobin C (HbAC), and 11
individuals with one or two copies of an *HBA2* deletion causing α-thalassemia[44]. We also
scored eight polymorphisms in *G6PD* that have been functionally associated with various
degrees of G6PD deficiency[45,46] and found that 32% of the study population carried at least
one, including 12 of the 20 donors with hemoglobinopathies. Among those with wild-type

Figure 1. Overview of study design and blood donors. (a) PCA of genetic variation across 35,759 unlinked exome SNPs. Samples collected in this study are plotted on coordinate space derived from 1000 Genomes reference populations. Points with white borders represent the six related individuals in this study. **(b)** Exome sequencing revealed that 37% of donors carried alleles for RBC disorders linked to *P. falciparum* resistance. Individuals with >1 disease allele were assigned to the category of the most severe condition. **non-carrier:** Donor without any of the following alleles or conditions. **G6PD⁻low:** Mild to medium G6PD deficiency (<42% loss of function). **G6PD⁻high:** Severe G6PD deficiency (>60% loss of function). **-α/αα:** heterozygous HBA2 deletion, or alpha thalassemia minima. **-α/-α:** homozygous HBA2 deletion, or alpha

102   thalassemia trait. **HbAC:** heterozygous HBB:E7K, or hemoglobin C trait. **HbAS:** heterozygous HBB:E7V, or
103   sickle cell trait. **HE:** hereditary elliptocytosis. **(c)** Parasite replication rate in donor RBCs was measured via
104   flow cytometry at three timepoints over 72 hours. 3D7 is a lab-adapted strain used in routine culture.
105   Th.026.09 is a recent clinical isolate from Senegal. **(d)** RBC phenotypes were measured using complete blood
106   counts with RBC indices, osmotic fragility tests, and ektacytometry.
107

hemoglobin, we identified one individual with polymorphisms associated with severe

G6PD deficiency (>60% loss of function) and 23 with polymorphisms associated with mild

to medium deficiency (<42% loss of function). To our knowledge, we detected no alleles

linked to other monogenic RBC disorders, including β-thalassemia or xerocytosis[47,48]. We

therefore classified the remaining 73 donors as 'non-carriers' of known disease alleles for

the purposes of this work.

P. falciparum replication rates vary widely among non-carrier RBCs

To determine the variation in _P. falciparum_ fitness among samples with different

genotypes, we performed invasion and growth assays with two strains of parasite: a

laboratory adapted strain (3D7) and a recent clinical isolate from Senegal (Th.026.09).

We observed a wide range of _P. falciparum_ growth rates among RBC samples, especially

among non-carriers that lacked known disease alleles (Figure 2A-C). Each strain's growth

rate is defined here as parasite multiplication over a full 48-hour cycle in donor RBCs

(Figure 1C), with the mean value for non-carriers set to 100% after normalization. Briefly,

we used a repeated control RBC sample (Figure 2, gray circles) and other batch-specific

factors to correct for variation in parasite growth across multiple experiments (see

Methods). Among non-carriers, growth rates ranged from 64-136% for 3D7 (SD = 17.7%)

and 76-128% for Th.026.09 (SD = 10.7%) (Figure 2A-B). Per-sample growth rates were

strongly correlated between the two strains (Figure 2C, $p < 3\times10^{-16}$) and positively

correlated when measured in different weeks (Figure S1A, $\rho=0.34$), demonstrating that

these data capture real variation among donor RBCs. Furthermore, as expected[42,49–51], we

detected reductions in mean growth rate for both strains in RBCs carrying known disease

alleles. These included severe G6PD deficiency (3D7, 60% of the non-carrier average,

p<0.014; Th.026.09, 64%, p<0.014), α-thalassemia trait (3D7, 75%, p=0.027; Th.026.09,

81%, p = 0.077), HbAS (3D7, 16%, $p=1.05\times10^{-7}$; Th.026.09, 19%, $p =1.2\times10^{-4}$), and HE

Figure 2. *P. falciparum* **replication rate varies widely among donor RBCs. (a-b)** Growth of *P. falciparum* strain 3D7 **(a)** or clinical isolate Th026.09 **(b)** over a full 48-hour cycle in donor RBCs (hours 24-72 of assay). Growth is presented relative to the average non-carrier rate after correction for batch effects, including comparison to a repeated RBC control (Methods). Repeated measurements of the same RBC control over the course of the experiment are shown as open gray circles. **(c)** Correlation in per-sample growth rates between the two *P. falciparum* strains. **(d-e)** Invasion efficiency of *P. falciparum* strain 3D7 **(d)** or clinical isolate Th026.09 **(e)** after a 24-hour incubation with donor RBCs, normalized in the same way as growth. Repeated measurements of the RBC control over time are shown as gray circles. **(f)** Correlation in per-sample invasion success for the two *P. falciparum* strains. Colors in **(c)** and **(f)** correspond to the classifications in other panels; R^2 and *p*-values are derived from OLS regression. *p<0.1; **p<0.05; ***p<0.01.

(3D7, 18%, p<0.014; Th.026.09, 33%, p<0.014). Notably, the wide distribution of growth

rates for non-carrier RBCs had considerable overlap with the growth rates in carrier RBCs.

Only the HbAS and HE samples fell entirely outside the non-carrier range. This observation

implies the existence of previously unknown RBC variation that impacts *P. falciparum*

growth, which may have effect sizes comparable to the milder known disease alleles.

We observed a similarly wide range in the efficiency of *P. falciparum* invasion into

donor RBCs (Figure 2D-F). Invasion is defined here as the fold-change in parasitemia over

the first 24 hours of the assay, when parasites previously maintained in standard culture

conditions egressed and invaded new donor RBCs (Figure 1C). Among non-carriers,

invasion rates ranged from 66-173% for 3D7 (SD=15.8%) and 41-193% for Th.026.09

(SD=28.9%) (Figure 2D-E). Compared to growth rates, no disease alleles conferred

protection against invasion that was extreme enough to fall outside the broad non-carrier

range. In 3D7 only, HbAC was associated with a 9% decrease in invasion (p=0.019) and

mild G6PD deficiency was associated with a 7% increase (p=0.072). Only HE had a

significant effect on the invasion efficiency of both strains (3D7, 67% of the non-carrier

average, p<0.014; Th.026.09, 57%, p<0.014). There was little correlation between the

invasion efficiencies of the two parasite strains in individual donor samples (Figure 2F),

potentially reflecting strain-specific differences in the pathways used for invasion[52].

However, as we also observed greater variability between repeated samples for invasion

than for growth (Figure S1), the invasion results may be influenced by greater

experimental noise.

<u>RBC phenotypes vary widely among non-carriers</u>

To assess phenotypic variation across donor RBCs, we measured 19 common

indices of RBC size and hemoglobin content from complete blood counts using an Advia

hematology analyzer (Figure 3A-D; Figure S2). Mean cellular volume (MCV) and

hemoglobin mass (MCH) are closely related traits, which can be represented together as

cellular hemoglobin concentration (CHCM) or the fraction of RBCs with 'normal'

hemoglobin and volume indices (M5). As expected, each known disease allele was

associated with a distinct set of RBC abnormalities. These included elevated CHCM for

HbAC (p=0.033), consistent with dehydration, and very low MCV (p=8.7×10^{-5}) and MCH

(p=4.6×10^{-7}) for α-thalassemia trait ($\alpha\alpha$/--), consistent with microcytic anemia[44]. RBCs

from the HE patient also had very low MCV and MCH (both p<0.014), reflecting the

membrane breakage and volume loss characteristic of this disease. However, for all of these

phenotypic measures, we also observed broad distributions in non-carriers that

Figure 3. Red cell phenotypes that are abnormal in carriers also vary widely among non-carriers.
MCV: mean RBC volume. **MCH:** mean cellular hemoglobin. **CHCM:** cellular hemoglobin concentration. **M5:**
fraction of RBCs with 'normal' indices of hemoglobin concentration (28-41 g/dL) and volume (60-120 fL).
Fragility: Tendency to lyse when osmotically overhydrated, defined as [NaCl] causing 50% lysis. **DI$_{max}$:**
Maximum membrane deformability under physiological salt conditions. **O$_{hyper}$:** Tendency to resist osmotic
dehydration and loss of deformability, defined as the hypertonic [NaCl] at which deformability is half
maximum. *p<0.1; **p<0.05; ***p<0.01; tests and samples as in **Figure 2.**

overlapped the distributions of most carriers (Figure 3A-D; Figure S2). Notably, the

breadth of the non-carrier distribution for each phenotype was large (e.g., 24 fL range for

MCV) compared to the average difference between Africans and Caucasians (e.g., 3-5

fL[24,35]). This wide diversity and substantial overlap between non-carrier and carrier traits

suggest that 'healthy' RBCs exist on the same phenotypic continuum as RBCs carrying

known disease alleles.

We observed similar patterns of variation in RBC membrane fragility (Figure 3E-F)

and membrane deformability (Figure 3G-H), as measured with osmotic fragility tests and

osmotic gradient ektacytometry (see Methods). Both sets of curves represent RBC

tolerance to osmotic stress, which can result in swelling and lysis (fragility, Figure 3E-F) or

dehydration and decreased deformability (O_{hyper}, Figure 3G-H). Specific

hemoglobinopathies were associated with moderate to strong reductions in fragility ($\alpha\alpha/\alpha$-

p=0.039; $\alpha\alpha$/-- p=0.003; HbAS p=0.014), deformability (DI_{max}; HbAC p=0.007; HbAS

p=4.4x10^{-6}), and/or resistance to loss of deformability when dehydrated (O_{hyper}; HbAC p =

0.019; HbAS p=0.041). HE cells were both extremely fragile (p<0.014) and extremely non-

deformable (p<0.014). In non-carriers, the distributions for all membrane measures were

wide, continuous, and overlapped the distribution of most carriers (Figure 3E-H; Figure

S3). Overall, these data demonstrate that multiple phenotypic alterations associated with

RBC disease alleles are also present in non-carrier RBCs.

<u>Non-carrier variation in RBC phenotypes predicts *P. falciparum* replication rate</u>

To identify sets of phenotypes associated with *P. falciparum* replication in non-

carrier RBCs, we used a machine learning method called Lasso (least absolute shrinkage

and selection operator) that performs regularization and variable selection[53] (see

Methods). Briefly, Lasso shrinks the regression coefficients for some possible predictors (in

this case, phenotypes) to zero, in order to obtain a subset of predictors that minimizes

prediction error. This method is well-suited for datasets in which possible predictors are

correlated, as are standard measurements of RBC size, hemoglobin, and membrane

dynamics. For each trait in each predictive subset selected by Lasso, we then applied

univariate OLS regression to more accurately estimate the sign of its effect on all measured

components of parasite fitness. The highest-confidence results are summarized in

Figure 4. RBC phenotypes predict *P. falciparum* fitness in non-carriers. **(a)** Phenotypes shown were selected by Lasso in at least one of four models (columns) with at least 60% cross-validation (CV) support, as indicated by blue shading. **(+/-)** shows the direction of effect if the phenotype was significantly correlated ($p<0.1$) with the parasite fitness component in a separate, univariate linear model. **(b)** Lasso model predictions of parasite growth as a function of RBC phenotypes are plotted against measured growth values. Each point represents one non-carrier. The solid line shows a perfect 1:1 fit; R^2 indicates the proportion of variance explained by Lasso predictions. **(c)** For 1000 permutations of the growth data, the histograms show the proportion of variance explained by the Lasso modeling procedure. Dashed lines indicate R^2 for the actual data. *p*-values indicate the fraction of permutations that explain at least as much variance as the real data. **(d,e)** As in **(b,c)**, but for *P. falciparum* invasion. **O_{hyper}:** Tendency to resist osmotic dehydration and loss of deformability, defined as the hypertonic [NaCl] at which deformability is half maximum (mM). **MCV:** mean RBC volume (fL). **DI_{max}:** Maximum membrane deformability under physiological salt conditions. **CHCM:** cellular hemoglobin concentration (g/dL). **Fragility:** Tendency to lyse when osmotically overhydrated, defined as [NaCl] causing 50% lysis (mM). **M5:** fraction of RBCs with 'normal' indices of hemoglobin concentration (28-41 g/dL) and volume (60-120 fL) (%). **MCH:** mean corpuscular hemoglobin (pg/RBC). **MPV:** mean platelet volume (fL). **RBC**: red cell number (x10^6/uL). **PLT:** platelet number (x10^3/uL).). **HCT:** hematocrit (%).

Figure 4A, with complete details provided in Table S2.

P. falciparum fitness in non-carrier RBCs was strongly predicted by variation in

several RBC traits related to deformability, dehydration, volume, and hemoglobin content.

Among 24 tested phenotypes, the most important trait was the ektacytometry parameter

O_{hyper}, which represents a cell's tendency to retain deformability in the face of dehydration

(Figure 3G). In univariate models, non-carrier RBCs with the largest O_{hyper} values supported

1.2-1.5X faster parasite growth (3D7 1.47X, p=0.003; Th.026.09 1.24X, p=0.005) and 1.4-

2.0X more effective invasion (3D7 1.36X, p=0.018; Th.026.09 1.94X, p=0.006) than RBCs

with the smallest O_{hyper} values (Table S2; Figure S5A). Consistent with this result, *P.*

falciparum replication was relatively inhibited in RBCs that were more dehydrated at

baseline, as indicated by higher CHCM (3D7 invasion p=0.007; 3D7 growth p=0.042;

Th.026.09 invasion p=0.006; Th.026.09 growth p=0.36). Parasites also grew faster in RBCs

with larger mean volume, or MCV (3D7 p=0.001; Th.026.09, p=0.001); a greater mass of

hemoglobin per cell, or MCH (3D7 p=0.073; Th.026.09 p=0.019); and more deformable

membranes, measured by DI_{max} (3D7 p=0.0006; Th.026.09 p=0.006). Some measures of

parasite fitness were also reduced in RBCs with more fragile membranes (3D7 invasion

p=0.65; 3D7 growth p=0.003; Th.026.09 invasion p=0.067; Th.026.09 growth p=0.14).

Additional phenotypes related to platelets and RBC density were selected for some models,

but the direction of their effects was unclear since they were not significant when

considered individually. These results indicate that common, non-pathogenic variation in

RBC size, membrane dynamics, and other correlated traits have meaningful effects on *P.*

falciparum replication rate in RBCs.

Together, 13 phenotype measurements explained approximately 40% of the

variation in 3D7 growth and invasion in non-carriers (Figure 4B,D). To test for overfitting,

we applied the same model-fitting procedure many times to randomly shuffled parasite

data, which usually failed to produce predictive models (i.e., R^2=0, Figure 4C,E). These

permutation tests found that real data explained, on average, 6.5X the variance in 3D7

growth and 6.3X the variance in 3D7 invasion than expected by chance (both p=0.01,

Figure 4C,E). For Th.026.09, the phenotypes selected by Lasso explained 17-21% of the

total variation in parasite fitness (Figure 4B), which was marginally better than expected

by chance for growth (3.7X, p=0.07, Figure 4C) and marginally worse than expected for

invasion (3.0X, p=0.13, Figure 4E). The weaker result for Th.026.09 may be explained by its

smaller dynamic growth range (Figure 2A-B), perhaps because clinical isolates are

relatively poorly adapted to laboratory conditions. Overall, these results demonstrate that

multiple, variable phenotypes impact *P. falciparum* susceptibility in healthy RBCs. Non-

carrier cells that are less hospitable to parasites share specific traits with RBCs that carry

disease alleles, including smaller size, decreased deformability, and an increased tendency

to lose deformability when dehydrating.

A small set of RBC genotypes predicts *P. falciparum* replication in non-carriers

Next, we tested whether non-carrier genotypes derived from exome sequencing

could improve our predictions of *P. falciparum* replication rate. To limit false positives and

increase statistical power[54,55], we focused on 23 candidate RBC proteins that have been

previously associated with *P. falciparum* resistance in the literature (Table S3). These

proteins contained 106 unlinked genetic variants (pairwise $r^2 < 0.1$) among the 73 non-

carriers. Along with these polymorphisms and phenotypes described above, we also

treated the top 10 principal components (PCs) of 1000 Genomes population structure as

possible predictors of *P. falciparum* fitness.

The addition of genetic factors substantially improved all four models of parasite

fitness. Across strains, 73-82% of the parasite variation in non-carriers was explained by a

total of 47 genetic and phenotypic variables selected by Lasso (Figure 5A; Table S4). For

3D7 and Th.026.09 respectively, polymorphisms in 14 of 23 malaria-related genes

explained 71% and 41% of the growth variation in non-carriers (Figure 5B). These models

were significant by permutation (both p=0.011; Figure S6AB), indicating that the results

were unlikely to be due to overfitting. Nonetheless, to confirm that the genetic signal was

specific to our set of 23 malaria-related genes, we used polymorphisms from 1000 sets of

random genes from the RBC proteome (Table S6) to build 1000 alternative models (see

Methods). On average, variants from random RBC genes explained only 17.9% of the

observed variation in *P. falciparum* growth for 3D7 (p=0.002) and 16.4% for Th.026.09

(p=0.047) (Figure 5B). In contrast to growth, the genotype/phenotype models of *P.*

falciparum invasion (Figure 5C) were only marginally significant by permutation (3D7

p=0.09, Th.026.09 p=0.11, Figure S6C-D) and the genetic signal was not specific to the set of

23 malaria-related genes (3D7 p=0.26, Th.026.09 p=0.52, Figure 5D). Nonetheless, 4 of the

8 variants selected by Lasso for 3D7 invasion were also selected for 3D7 growth. Together,

these results indicate that *P. falciparum* growth rate in RBCs is strongly determined by host

genetic variation concentrated in a small number of genes.

Figure 5. RBC genotypes substantially improve predictions of *P. falciparum* fitness in non-carriers. (a) Lasso model predictions of parasite growth as a function of RBC phenotypes, PCs of population structure, and variants in 23 malaria-related proteins (Table S3) are plotted against measured growth values. Each point represents one non-carrier. The solid line shows a perfect 1:1 fit; R^2 indicates the proportion of variance explained by Lasso-selected predictors. **(b)** For 1000 random sets of 23 genes drawn from the RBC proteome, the histograms show the variance explained *by variants within those genes* after applying the same Lasso modeling procedure as (a). Dashed lines indicate the variance explained by variants within the 23 malaria-related genes. *p*-values indicate the fraction of random gene sets that explain at least as much variance as the real data. **(c,d)** As in (a,b), but for *P. falciparum* invasion. **(e)** Variants and PCs counted in each cell were selected in the designated Lasso model with at least 70% cross-validation (CV) support. Several tested genes had 0 associated variants: *CD55*, *EPB41*, *HP*, *HBA1/2, GYPA, GYPC, GYPB,* and *HBB.* Details on lower-confidence variants are available in Table S4.

Of the top 31 variants associated by Lasso with *P. falciparum* growth, seventeen (55%) were synonymous (Table S4). This fraction did not significantly differ from the input set of 106 variants (p=0.64, 2-tailed z-score). Although 15/17 of these synonymous variants have no known, direct function, 7/15 are in linkage disequilibrium ($r^2 > 0.1$) with other variants that have been associated with RBC phenotypes by GWAS (Table S4). Similarly, 6 of the top 14 non-synonymous polymorphisms associated with *P. falciparum* growth have been previously linked to RBC or other traits (Table S4). This overlap with previous studies suggests that the variants associated with *P. falciparum* growth in this study are likely to tag functional genetic loci.

Nearly all of the top 31 growth-associated polymorphisms occurred in either (1) ion

channel proteins, which regulate RBC hydration; (2) components of the flexible RBC

membrane backbone; or (3) red cell plasma membrane proteins, including known invasion

receptors (Figure 5E). In the first category, the highly polymorphic ion channel *PIEZO1*

contained six polymorphisms associated with small (-3%) to moderate (-32%) reductions

of *P. falciparum* growth rate (Table S4). The microsatellite variant *PIEZO1*-E756del, which

has been a focus of several recent studies[56–59], predicted a moderate reduction in Th.026.09

growth (-8.2%, p=0.011) but was not related to RBC dehydration in these data (Table S5).

We also detected one growth-associated variant in *ATP2B4,* which encodes the primary

RBC calcium channel PMCA4b (-13% 3D7, p=0.006). This variant tags an *ATP2B4* haplotype

implicated by GWAS in protection from severe malaria[16,17,60] and many RBC

phenotypes[18,31,60,61] including RDW, which was replicated in our data set (Table S4). *SPTA1*

and *SPTB,* which encode the flexible spectrin backbone of RBCs, also contained several

variants associated with the growth of at least one *P. falciparum* strain, as did the structural

linker proteins ANK1, SLC4A1, and EPB42 (Figure 5E; Table S4). Finally, we identified one

polymorphism in ABCB6, two in CD44, and two in basigin (BSG) that were strongly

associated with *P. falciparum* growth (Figure 5E; Table S4). These plasma membrane

proteins have previously been implicated in *P. falciparum* invasion by *in vitro* genetic

deficiency studies[62–64]. Notably, two of the natural polymorphisms identified here are

synonymous quantitative trait loci (QTL) for CD44 splicing (rs35356320) and BSG

expression (rs4682)[65]. We also detected one variant in the RBC iron exporter *FPN* and one

in *G6PD* with small effects on Th.026.09 growth (2.4%, p=0.015 and 3.4%, p=0.010

respectively; Table S4). No associated variants were detected in the other nine tested genes

(Table S2), including three hemoglobin proteins, three glycophorins, *CD55*, *EPB41*, and *HP*.

Together, these data demonstrate that *P. falciparum* growth rate in non-carrier RBCs is

strongly determined by dozens of host genetic variants, which together shape the

phenotypic distribution of red cell susceptibility.

In addition to these specific genetic variants, PCs of population structure explained

22-26% of the variation in non-carrier invasion and 10-12% of the variation in growth.

Interestingly, neither PC1 nor PC2—which distinguish Africans from other populations

(Figure 1A)—were related to parasite fitness. Instead, this signal of correlated variation

was partially driven by a six-member family with unique ancestry in our sample (Figure

1A, white borders) and extreme parasite fitness values (Figure S6). Although the present

study is limited by sample size, the association between global genetic PCs and *P.*

falciparum fitness suggests that additional functional variants remain to be discovered in

many populations.

<u>African ancestry does not predict *P. falciparum* resistance in red cells</u>

Based on evidence from balanced disease alleles like HbAS, it has been suggested

that anti-malarial selection has shaped polygenic red cell phenotypes in African

populations as a whole[37,56,66]. We tested this hypothesis by examining the correlation

between African ancestry and *P. falciparum* replication rate in RBCs from non-carriers

(Figure 6A-D). We found no evidence that these traits were related, apart from an weakly

positive relationship between African ancestry and the invasion rate of Th.026.09, a clinical

Senegalese strain (p=0.01, R^2=0.09, Figure 6D). To understand this result, we next

examined how key RBC phenotypes identified in this study (Figure 4A) vary with African

ancestry. We found that greater African ancestry predicts reduced osmotic fragility (p=

9.03 x 10^{-6}), increased hypotonic deformability (DI_{min}, p=0.001), lower cellular hemoglobin

concentration (CHCM, reflecting reduced dehydration, p=0.007), and a reduced fraction of

RBCs with normal volume and hemoglobin content (M5, p=0.033). All of these traits

actually predict *greater* red cell susceptibility to *P. falciparum* (Figure 4A), although

together they explain less than 9% of the non-carrier variation in parasite fitness. The

remaining key phenotypes (Figure 4A) do not vary with African ancestry, which may

explain why African ancestry itself is mostly unassociated with *P. falciparum* fitness in non-

carrier RBCs (Fig 6A-D).

Next, we used allele frequency data from the global gnomAD collection[67] to test

whether polymorphisms associated with *P. falciparum* fitness in RBCs (Figure 5E) vary

between African and European populations. Geographical differences in malaria selection

are hypothesized to have increased the frequency of hundreds or thousands of

undiscovered anti-malarial alleles in Africa[22,68], as has been shown for several variants

causing common RBC disorders (Figure 1B)[7]. Here, aside from well-known disease

variants, we find no evidence that African populations as a whole are enriched for RBC

Figure 6. No evidence of widespread selection in Africa for slower *P. falciparum* replication or protective alleles in non-carriers. (a-d) Parasite growth and invasion measurements plotted against the exome-wide fraction of African ancestry, determined by comparison to 1000 Genomes reference populations. Adjusted-R^2 and *p*-values are shown for OLS regression. **(e)** Blue points represent non-carrier alleles selected by Lasso that also have significant effects on parasite growth ($p<0.1$) in univariate linear models (Table S4). The maximum estimated effect across two *P. falciparum* strains is shown. Red and purple points represent HbAS and the HBA2 deletion in an additive model, respectively. F_{ST} was calculated between pan-African and pan-European samples in gnomAD (Methods). Other disease alleles are not shown because they do not function as single loci (G6PD⁻, HE) or do not affect growth (HbAC).

polymorphisms that impair *P. falciparum* growth in at least one strain (Figure 6E). Of 25 alleles with detectable protective effects, 52% are in fact more common in Europeans than Africans (p=0.95, binomial test). Besides HbAS, the polymorphism with the largest effect is a synonymous variant in PIEZO1 (-31.9%, p=0.001), which is found at <4% frequency in both Africans and Europeans. A few other protective polymorphisms, including two in *PIEZO1* and two in *SPTB*, are nearly fixed in both populations. We do observe one F_{ST} outlier variant, a synonymous allele at SPTA1 L958, which is substantially more common in Africa (70% frequency) than in Europe (25%) and has a mild protective effect against *P. falciparum* growth (-6.5%, p=0.027, Figure 6E). Given its high frequency, as well as its

direct association with a number of RBC phenotypes in previous GWAS (Table S4), this

synonymous SPTA1 allele could be a promising candidate for investigating anti-malarial

selection in Africa. However, the total set of natural RBC variants that limit *P. falciparum*

growth *in vitro* do not appear to support that general hypothesis.

DISCUSSION

Healthy red blood cells (RBCs) harbor extensive phenotypic and genetic variation,

both within and between human populations. In this work, we demonstrate that this

variation modulates a wide range of RBC susceptibility to *P. falciparum* parasites. Our

findings add to a growing understanding of the genetic and phenotypic basis of RBC

resistance to *P. falciparum*, especially for RBCs that lack population-specific disease alleles.

In addition to suggesting new targets for future malaria interventions, these findings

challenge assumptions about the role of malaria selection in shaping human RBC diversity.

Since exponential replication of *P. falciparum* is a significant driver of malaria

disease progression[69], the ample variation in this trai*t* that we observed *in vitro* could be

relevant for clinical outcomes in endemic regions. Growth inhibition from HbAS, for

example, reduces the risk of death from malaria by reducing parasite density in the

blood[70,71]. While HbAS has an extreme effect size, the 3-fold range of parasite replication

rates we found among non-carrier RBCs shares substantial overlap with RBCs carrying

other protective variants. Although the physiologically complex basis of severe malaria[72]

makes it difficult to estimate the precise contribution of RBC factors to severe malaria risk,

the genotypes and phenotypes we have associated with *P. falciparum* fitness may be

promising targets for future therapeutic interventions.

We have shown here that widespread, 'normal' variation in RBC traits like volume,

hydration status, and maximum deformability is associated with *P. falciparum* fitness in

non-carrier RBCs. Reassuringly, these phenotypes are present to a stronger degree in RBCs

that do carry disease alleles[73–75]. They are also consistent with published reports of

extreme RBC traits that limit parasite replication. For example, Tiffert *et al.*[76] used an

experimental gradient of ion concentrations to show that *P. falciparum* growth is strongly

reduced in chemically dehydrated RBCs. Here, we found that naturally occurring, mildly

dehydrated RBCs also supported less efficient parasite growth and invasion. Similarly, the

rare Dantu variant of the glycophorin A/B receptors has been associated with increased

RBC membrane tension, decreased *P. falciparum* invasion efficiency, and decreased risk of

severe malaria[11,77,78]. Our data on RBC membrane deformability and fragility support the

same link between more flexible RBC membranes and greater *P. falciparum* success.

Moreover, they expand upon previous studies by demonstrating for the first time that

common, healthy phenotypic variation in RBCs contributes meaningfully to *P. falciparum*

fitness.

In our linear models of *P. falciparum* growth, phenotypic variation was strongly

outperformed by genetic variation in a small number of RBC proteins. This result implies

the existence of additional RBC phenotypes that we did not measure (or did not measure

with sufficient accuracy), but which have a genetic basis in a small number of genes

important to *P. falciparum*. Approximately half of the polymorphisms we identified are

non-synonymous, and may therefore exert direct effects on phenotypes like RBC

membrane structure or ion transport. The abundances of RBC gene transcripts and

proteins are other clear candidates for future phenotype studies, given that the other half

of associated polymorphisms were synonymous. A recent metanalysis showed that 'silent'

and 'coding' SNPs are equally likely to be associated with human disease, and moreover

have similar effect sizes[83]. Although synonymous SNPs may sometimes be linked to coding

variants, some have been shown to directly impact mRNA expression, protein translation

speed, and protein folding by altering transcription levels, codon usage, and mRNA

stability[84]. Synonymous SNPs that impact splicing, like rs35356320 in CD44, may also

impact protein structure. Other conceivable RBC phenotypes, such as the dynamics of

membrane modification during *P. falciparum* development, may only become evident in

more detailed time course experiments. The true number of RBC phenotypes important to

P. falciparum may be effectively infinite[85], making it very useful in practice that static

genetic variation is so strongly predictive of parasite growth.

Overall, just 31 polymorphisms in 14 RBC genes explained the majority of the

variation in *P. falciparum* growth among non-carriers. One reason that our explanatory

power is so high is our focus on *P. falciparum* replication in RBCs, which is a relatively

simple component of a larger, complex disease. This focus allowed us to use controlled, *in*

vitro experiments to detect genetic variants with small to moderate effects. Another

important explanation is our use of Lasso variable selection on just 106 polymorphisms in

genes with strong existing links to malaria. This approach obviated the need to meet an

exome-wide significance threshold, while still allowing for the discovery of novel,

putatively functional alleles in well-known disease genes. Furthermore, we confirmed that

this set of genes was enriched for signal compared to random sets of genes drawn from the

exome-wide background. It is important to note that genetic linkage complicates the

identification of the exact functional polymorphisms in any population sample[87], and we

cannot rule out that some variants identified here are merely linked to the true functional

variants. Indeed, about half of the variants we identified occur in linkage blocks containing

other SNPs associated with RBC traits in GWAS studies. In any case, we present significant

evidence that loci containing 14 RBC genes are strongly enriched for polymorphisms with

significant impacts on *P. falciparum* growth.

One of the unique aspects of our study is the participation of individuals with a

range of African ancestry, defined by similarity to donors from five 1000 Genomes

reference populations. We found that African ancestry was associated with RBC

phenotypes that *improved* parasite fitness, particularly for the clinical Senegalese strain

Th.026.09. In the future, it would be very interesting to test for local parasite adaptation to

human RBCs using *P. falciparum* strains and RBC samples from around the globe. We also

found that the polymorphisms associated with *P. falciparum* growth by Lasso are not

enriched in African populations included in the gnomAD database of human variation. This

is not consistent with African-specific selection on these smaller-effect variants, despite the

famous examples provided by RBC disease alleles. This unexpected result for 'healthy'

variation has at least three possible explanations. First, relatively small-effect alleles may

not have had sufficient time to increase in frequency since *P. falciparum* began expanding

in humans some 5,000-10,000 years ago[88,89]. Second, the complexity of severe malaria

could mean that the variants discovered here do not ultimately impact disease outcome,

especially relative to known disease variants. Third, human adaptation may be too local to

detect with coarse-grain sampling of Sub-Saharan African genetic diversity (e.g [90]). Our

data do suggest, however, that few RBC alleles remain to be discovered that are both

widespread in Africa and have large effects on *P. falciparum* proliferation in RBCs (e.g. [56]).

More broadly, these data show that it is inaccurate to make assumptions about RBC

susceptibility to *P. falciparum* based on a person's race or continental ancestry. These kinds

of hypotheses[37,56,66,68] are based on well-known examples of balanced disease alleles, but

our data suggest that these examples are exceptions rather than the rule. It is important to

recognize that, at least in biology classes, the use of racially-based genetic examples

increases the belief that genes encode absolute, functional differences between races[91,92].

This perspective neglects the fact that >90% of genetic differences among humans occur

within populations, rather than across them[93]. In this study, global population structure

captured by PCs showed that non-African variation has its own distinct impacts on *P.*

falciparum replication in RBCs. These data are an important reminder that most human

genetic diversity is a result of demography, not population-specific selection[94,95].

In conclusion, this study demonstrates that substantial phenotypic and genetic

diversity in healthy human RBCs impacts the replication of malaria parasites. Whether or

not this diversity is shaped by malaria selection, a better understanding of how *P.*

falciparum biology is impacted by natural RBC variation can help lead to new therapies for

one of humanity's most important infectious diseases.

METHODS

Sample collection and preparation

Subjects with no known history of RBC disorders were recruited to donate blood at

the Stanford Clinical and Translational Research Unit. Written informed consent was

obtained from each subject and/or their parent as part of a protocol approved by the

Stanford University Institutional Review Board (#40479). To help control for weekly batch

effects, Subject 1111 donated fresh blood for parasite assays each week. 11 other subjects

donated blood on at least two different weeks. Whole blood samples from a hereditary

elliptocytosis patient were obtained from Dr. Bertil Glader under a separate approved

protocol (#14004) that permitted sample sharing among researchers. All samples were de-

identified upon collection by labeling with a random four-digit number. Two samples were

eventually removed from analysis based on a failed sequencing library (6449KD) and

history of stem cell transplant (8715).

Whole blood was drawn into CPDA tubes and spun down within 36 hours to

separate serum, buffy coat, and red blood cells. Red blood cells were washed and stored in

RPMI-1640 medium (Sigma) supplemented with 25 mM HEPES, 50 mg/L hypoxanthine,

and 2.42 mM sodium bicarbonate at 4°C. Buffy coat was transferred directly to cryotubes

and stored at -80°C.

Exome sequencing and genotype calling

Genomic DNA was isolated from frozen buffy coats using a DNeasy Blood and Tissue

Kit (Qiagen). Libraries were prepared using a KAPA Hyperplus kit (Roche) and hybridized

to human exome probes using the SeqCap EZ Prime Exome kit (Roche). The resulting

exome libraries were sequenced with paired-end 150 bp Illumina reads on the HiSeq or

NextSeq platforms at Admera Health (South Plainfield, NJ).

Reads were aligned to the hg38 human reference genome using bwa mem[96],

yielding an average coverage of 42X across targeted exome regions (excluding sample

6449KD). Variants were called using GATK best practices[97] and hard filtered with the

following parameters: QD < 2.0, FS > 60.0, ReadPosRankSum < -2.5, SOR > 2.5, MQ < 55.0,

MQRankSum < -1.0, DP < 500. To minimize the effects of sequencing errors, variants not

present in 1000 Genomes, dbSNP_138, or the Mills indel collection[98] were discarded.

Variants were also discarded if their frequencies in our sample did not fall between

gnomAD African or European frequencies[67] and significantly differed from both groups'

frequencies by Fisher's Exact Test (p≤0.05).

PIEZO1 E756del was genotyped via PCR and Sanger sequencing according to a

previously published protocol[57]. To call deletion variants that cause α-thalassemia in the

paralogous genes HBA2 and HBA1, we extracted reads from each .bam file that lacked any

mismatches or soft-clipping and had MAPQ ≥ 13 (i.e., <5% chance of mapping error).

Coverage with these well-mapped reads was calculated over the 73 and 81 bp of unique

sequence in HBA2 and HBA1, respectively, and normalized to each sample's exome-wide

coverage. To determine which samples has unusually low coverage, we formed an *ad hoc*

reference panel of seven donors who were unlikely to carry deletion alleles based on their

normal MCH, MCV, and HGB and >96% exome-wide European ancestry[99]. We called

heterozygous HBA2 deletions when normalized coverage across three unique regions of

the HBA2 gene was below the minimum reference value. Similarly, we called homozygous

HBA2 deletions when normalized coverage across three unique regions of the HBA2 gene

was less than half of the minimum reference value. This approach resulted in an estimated

HBA2 copy number of 2.0 in the reference panel, 0.95 in eight putative heterozygotes and

0.12 in four putative homozygotes. The same method produced no evidence of HBA1

deletion in any sample.

<u>Variant classification and linkage pruning</u>

Exonic variants in RefSeq genes were identified using ANNOVAR[100]. Variants were

classified into three categories: those within 23 malaria-related genes (Table S2); those

within 887 other red blood cell proteins (Table S5) derived with a medium-confidence

filter from the Red Blood Cell Collection database (rbcc.hegelab.org); and those within any

other gene. Singleton variants were removed to avoid excessive false positives.

Linkage between all pairs of bi-allelic, exonic variants in our 121 genotyped samples

was calculated using the --geno-r2 and --interchrom-geno-r2 functions in vcftools [101].

Variants in RBC genes that shared $r^2 > 0.1$ with any variant in the 23-gene set were

removed. Within the 23-gene set and RBC-gene set separately, non-carrier variants were

ranked by the *p*-values of their OLS regression with all four parasite measures. Then, one

variant was removed from each pair with $r^2 > 0.1$, prioritizing retention in the following

order: greater significance across models; non-synonymous protein change; higher

frequency in our sample; and finally by random sampling. We report results from additive

genetic models (genotypes coded 0/1/2), which performed as well or better than recessive

(0/0/2) and dominant (0/2/2) models.

<u>Population analysis</u>

The population ancestry of our donors was assessed by comparison with African,

European, East Asian, and South Asian reference populations from the 1000 Genomes

Project[102]. Briefly, variants called from an hg38 alignment of the 1000 Genomes data [103]

were filtered for concordance with the variants genotyped in this study. The --indep-

pairwise command in PLINK[104] was used to prune SNPs with $r^2 > 0.1$ with any other SNP in

a 50-SNP sliding window, producing 35,759 unlinked variants. These variants were

analyzed in both PLINK --pca and in ADMIXTURE[105] with K=4 for the 121 genotyped

individuals in this study, alongside 2,458 individuals from 1000 Genomes. Pan-African and

pan-European allele frequencies were obtained from gnomAD v3[67]. F_{ST} for specific alleles

was calculated as $(H_T-H_S)/H_T$ and then polarized, such that positive values indicate variants

more common in Africa.

P. falciparum culture and assays

P. falciparum strain 3D7 is a laboratory-adapted strain that was obtained from the

Walter and Eliza Hall Institute (Melbourne, Australia) and routinely cultured in human

erythrocytes obtained from the Stanford Blood Center. Th.026.09 is a clinical strain isolated

from a patient in Senegal in 2009 that was then adapted to short-term culture[106] and kindly

provided by Daouda Ndiaye and Sarah Volkman. 3D7 was maintained at 2% hematocrit in

RPMI-1640 supplemented with 25 mM HEPES, 50 mg/L hypoxanthine, 2.42 mM sodium

bicarbonate and 4.31 mg/ml Albumax (Invitrogen) at 37°C in 5% CO2 and 1% O2.

Th.026.09 was maintained in the same media, with the exception that half the Albumax was

replaced with heat-inactivated human AB serum.

Parasite growth and invasion assays were performed using schizont-stage parasites

isolated from routine culture using a MACS magnet (Miltenyi). Parasites were added at

~0.5% initial parasitemia to fresh erythrocytes suspended at 1% hematocrit in complete

RPMI, as above. Parasites were cultured in each erythrocyte sample for three to five days in

triplicate 100 uL wells. Parasitemia was determined on day 0, day 1 (24 hours), day 3 (72

hours), and in some cases day 5 (120 hours) by staining with SYBR Green 1 nucleic acid

stain (Invitrogen, ThermoFisher Scientific, Eugene, OR, USA) at 1:2000 dilution in

PBS/0.3% BSA for 20 minutes, followed by flow cytometry analysis on a MACSQuant flow

cytometer (Miltenyi). Raw invasion rate was defined as the day 1 parasitemia divided by

the day 0 parasitemia; raw growth rate was defined as the day 3 (or day 5) parasitemia

divided by the day 1 (or day 3) parasitemia. Day 0 parasitemia was not measured in weeks

1-3, so invasion rate estimates are absent for these samples. The parasite assays failed for

both strains in week 9 and for Th.026.09 in week 10, and so were repeated in weeks 10 and

11 with RBCs that had been stored for 1 or 2 weeks.

To correct for batch effects, such as week-to-week variation in *P. falciparum*
replication, we extracted the residuals from a linear regression of the raw parasite values
against up to four significantly related batch variables: (1) the raw values for control donor
1111 each week; (2) the parasitemia measured at the previous time point; (3) the age in
weeks of the RBCs being measured; and (4) the experimenter performing the assays.
Notably, there was no additional effect of 'Week' or the length of the experiment (i.e., 3 or 5
days) once the previous variables were regressed out. To convert these residuals (mean 0)
to relative percents, we first transformed them with a linear model parameterized by data
from control donor 1111 and the most extreme donors (HbAS for growth; G6PD$^-$$_{high}$ and HE
for invasion), normalized by the 1111 rates from the appropriate week. These values were
finally arithmetically adjusted so that the mean invasion and growth values for non-
carriers was 100%.

<u>Red cell phenotyping and normalization</u>
Complete blood count (CBC) data for RBCs, reticulocytes, and platelets were
obtained with an Advia 120 hematology analyzer (Siemens, Laguna Hills, CA) at the Red
Cell Laboratory at Children's Hospital Oakland Research Institute. These data were: RBC,
HGB, HCT, MCV, MCH, MCHC, CHCM, RDW, HDW, PLT, MPV, Reticulocyte number and
percentage, and the fraction of RBCs in each of nine cells of the RBC matrix (M1-M9, with
MCH cutoffs of 28 and 41 and MCV cutoffs of 60 and 120). Systematic biases were evident
for some measures in certain weeks, but data from control donor 1111 were not available
for all weeks. Therefore, CBC data were normalized such that the median value for non-
carrier samples was equal across weeks.
Osmotic fragility tests were performed by incubating 20 μL of washed erythrocytes
for 5 minutes in 500 μL solutions of NaCl in 14 concentrations: 7.17, 6.14, 5.73, 5.32, 4.91,
4.50, 4.30, 4.09, 3.89, 3.68, 3.27, 3.07, 2.66, and 2.46 g/L. Tubes were spun for 5 minutes at
1000g and 100 μL of supernatant was transferred to a 96-well plate in duplicate.
Hemoglobin concentration was determined by adding 100 μL of Drabkin's reagent (Ricca
Chemical) to each well and measuring absorbance at OD$_{540nm}$ with a Synergy H1 plate
reader (Biotek). Relative lysis was determined by normalizing to the maximum hemoglobin

concentration in the 14-tube series for each sample. After outlier points were manually

removed, sigmoidal osmotic fragility curves were estimated under a self-starting logistic

model in the nls package in R. Curves were summarized by the relative tonicity at which

50% lysis occurred and normalized within weekly batches, such that this value was equal

for control sample 1111 across weeks.

670   Osmotic gradient ektacytometry[74,107] was performed at the Red Cell Laboratory at

Children's Hospital Oakland Research Institute (CHORI). Red cell deformability estimates

across a range of NaCl concentrations were fitted to a 20-parameter polynomial model to

generate a smooth curve, which was manually verified to closely fit the data. Each curve

was summarized with three standard points (Figure S3A)[74], which were normalized such

that the median x- and y-values of the three points was equal for non-carrier samples

across weeks.

<u>Statistical analysis</u>

679   Student's t-test was used to compare trait values between non-carriers and each

680   other group. Where n=1 (i.e., for G6PD$^-$$_{high}$ and HE), p-values were defined as the percentile

681   of the non-carrier distribution. Significance in these tests was defined as $p<0.1$. OLS linear

682   regression was performed with the lm function in R for all comparisons of two continuous

683   variables, unless otherwise specified. Adjusted R^2 values are reported.

684   Lasso regression[53,108] was performed with the cv.glmnet function in R. To account

for moderate uncertainty in the choice of lambda (due to random division of data for cross-

validations), each model was run 1000 times. For each predictor, we report the fraction of

runs in which it was selected (e.g Figure 4A; Table S1; Table S3) .The R^2 values reported

between actual and predicted parasite measures are medians from 1000 runs. We noted

that effect size estimates for each predictor varied considerably across runs; therefore, we

used univariate OLS regression to separately estimate the effect size of each predictor

selected at least once in at least one model.

692   The significance of each Lasso result was assessed by permutation. Parasite values

were randomly shuffled 1000 times, preserving correlations among potential predictors.

The median models for 6 cross-validations of each of the 1,000 permuted data sets were

compared to the median model from the real data. To compare the 23 malaria-related

genes to other sets of genes, a gene-specific R^2 for the target set was calculated using OLS

with only specific variants (no phenotypes or PCs) selected in at least 70% of cross-

validations. 1000 random sets of 23 genes were then generated from the RBC proteome

(Table S5) and analyzed identically. *P*-values were calculated as the percentile of this

permuted distribution in which the malaria-related gene set fell.

ACKNOWLEDGEMENTS

We gratefully acknowledge the invaluable participation of all 123 volunteer blood donors.

Nick Bondy, Sandra Larkin, Brian Fleischer, Ashley Dunn, Talal Seddik, Trung Pham, David

Vu, and Spectrum Child Health provided crucial assistance in donor coordination and

sample processing. For quantitative advice, we thank Grant Kinsler and the Stanford

Statistics Consulting Group. This work was primarily supported by a Pilot Early Career

award from the Stanford Maternal Child Health Research Institute and a Gabilan Faculty

Award from the Stanford University School of Medicine Office of Faculty Development and

Diversity. E.R.E. received additional support from the Stanford Center for Computational,

Evolutionary, and Human Genomics. D.A.P. was funded through an NIH MIRA award

5R35GM118165-05. E.S.E. was funded by the Doris Duke Charitable Foundation (no.

2016098) and the NIH (DP2HL13718601). Local blood samples were drawn at the

Stanford Clinical and Translational Research Unit, which is supported by CTSA Grant UL1

TR001085.

REFERENCES

1.   WHO. *Malaria eradication: benefits, future scenarios and feasibility. Executive*

summary, WHO Strategic Advisory Group on Malaria Eradication.

https://www.who.int/publications/i/item/who-cds-gmp-2019-10 (2019).

2.   Kariuki, S. N. *et al.* Red blood cell tension protects against severe malaria in the Dantu

blood group. *Nature* 1–5 (2020) doi:10.1038/s41586-020-2726-6.

3.   Koch, M. *et al.* Plasmodium falciparum erythrocyte-binding antigen 175 triggers a

biophysical change in the red blood cell that facilitates invasion. *Proc. Natl. Acad. Sci.*

U. S. A. **114**, 4225–4230 (2017).

4.   Francis, S. E., Sullivan, D. J. & Goldberg, and D. E. HEMOGLOBIN METABOLISM IN

THE MALARIA PARASITE *PLASMODIUM FALCIPARUM*. *Annu. Rev. Microbiol.* **51**, 97–

123 (1997).

5.   Hanssen, E. *et al.* Soft X-ray microscopy analysis of cell volume and hemoglobin

content in erythrocytes infected with asexual and sexual stages of Plasmodium

falciparum. *J. Struct. Biol.* **177**, 224–232 (2012).

6.   Zhang, Y. *et al.* Multiple stiffening effects of nanoscale knobs on human red blood

cells infected with Plasmodium falciparum malaria parasite. *Proc. Natl. Acad. Sci. U. S.*

A. **112**, 6068–6073 (2015).

7.   Kwiatkowski, D. P. How malaria has affected the human genome and what human

genetics can teach us about malaria. *American Journal of Human Genetics* vol. 77 171–

192 (2005).

8.   Pasvol, G., Weatherall, D. J. & Wilson, R. J. M. Cellular mechanism for the protective

effect of haemoglobin S against P. falciparum malaria [22]. *Nature* vol. 274 701–703

(1978).

9.   Archer, N. M. *et al.* Resistance to Plasmodium falciparum in sickle cell trait

erythrocytes is driven by oxygen-dependent growth inhibition. *Proc. Natl. Acad. Sci.*

U. S. A. **115**, 7350–7355 (2018).

10.   Ruwende, C. & Hill, A. Glucose-6-phosphate dehydrogenase deficiency and malaria.

Journal of Molecular Medicine vol. 76 581–588 (1998).

11.   Leffler, E. M. *et al.* Resistance to malaria through structural variation of red blood cell

invasion receptors. *Science (80-.).* **356**, 1140–1152 (2017).

12.   MGE Network. Insights into malaria susceptibility using genome-wide data on 17,000

individuals from Africa, Asia and Oceania. *Nat. Commun.* **10**, 5732 (2019).

13.   Ndila, C. M. *et al.* Human candidate gene polymorphisms and risk of severe malaria in

children in Kilifi, Kenya: a case-control association study. *Lancet Haematol.* **5**, e333–

e345 (2018).

14.   Band, G. *et al.* A novel locus of resistance to severe malaria in a region of ancient

balancing selection. *Nature* **526**, 253–257 (2015).

15.   Rockett, K. A. *et al.* Reappraisal of known malaria resistance loci in a large

multicenter study. *Nat. Genet.* **46**, 1197–1204 (2014).

16.   Timmann, C. *et al.* Genome-wide association study indicates two novel resistance loci

for severe malaria. *Nature* **489**, 443–446 (2012).

17.   Zámbó, B. *et al.* Decreased calcium pump expression in human erythrocytes is

connected to a minor haplotype in the ATP2B4 gene. *Cell Calcium* **65**, 73–79 (2017).

18.   Li, J. *et al.* GWAS of blood cell traits identifies novel associated loci and epistatic

interactions in Caucasian and African-American children. *Hum. Mol. Genet.* (2013)

doi:10.1093/hmg/dds534.

19.   Boyle, E. A., Li, Y. I. & Pritchard, J. K. An Expanded View of Complex Traits: From

Polygenic to Omnigenic. *Cell* vol. 169 1177–1186 (2017).

20.   MGE Network. Insights into malaria susceptibility using genome-wide data on 17,000

individuals from Africa, Asia and Oceania. *Nat. Commun.* **10**, 5732 (2019).

21.   De Mendonça, V. R. R., Goncalves, M. S. & Barral-Netto, M. The Host Genetic Diversity

in Malaria Infection. *J. Trop. Med.* **2012**, 17 (2012).

22.   Mackinnon, M. J., Mwangi, T. W., Snow, R. W., Marsh, K. & Williams, T. N. Heritability

of malaria in Africa. *PLoS Med.* **2**, 1253–1259 (2005).

23.   Canela-Xandri, O., Rawlik, K. & Tenesa, A. An atlas of genetic associations in UK

Biobank. *Nat. Genet.* **50**, 1593–1599 (2018).

24.   Lo, K. S. *et al.* Genetic association analysis highlights new loci that modulate

hematological trait variation in Caucasians and African Americans. *Hum. Genet.* **129**,

307–317 (2011).

25.   Cooling, L. Blood groups in infection and host susceptibility. *Clin. Microbiol. Rev.* **28**,

801–870 (2015).

26.   Whitfield, J. B., Martin, N. G. & Rao, D. C. Genetic and environmental influences on the

size and number of cells in the blood. *Genet. Epidemiol.* **2**, 133–144 (1985).

27.   Evans, D. M., Frazer, I. H. & Martin, N. G. Genetic and environmental causes of

variation in basal levels of blood cells. *Twin Res.* **2**, 250–257 (1999).

28.   Pilia, G. *et al.* Heritability of Cardiovascular and Personality Traits in 6,148

Sardinians. *PLoS Genet.* **2**, e132 (2006).

29.   Chen, M.-H. *et al.* Trans-ethnic and Ancestry-Specific Blood-Cell Genetics in 746,667

Individuals from 5 Global Populations. *Cell* **182**, 1198-1213.e14 (2020).

30.   Vuckovic, D. *et al.* The Polygenic and Monogenic Basis of Blood Traits and Diseases.

Cell **182**, 1214-1231.e11 (2020).

31.   van der Harst, P. *et al.* Seventy-five genetic loci influencing the human red blood cell. *Nature* **492**, 369–75 (2012).

32.   Chami, N. *et al.* Exome Genotyping Identifies Pleiotropic Variants Associated with Red Blood Cell Traits. *Am. J. Hum. Genet.* **99**, 8–21 (2016).

33.   Astle, W. J. *et al.* The Allelic Landscape of Human Blood Cell Trait Variation and Links to Common Complex Disease. *Cell* **167**, 1415-1429.e19 (2016).

34.   Perry, G. S., Byers, T., Yip, R. & Margen, S. Iron nutrition does not account for the hemoglobin differences between blacks and whites. *J. Nutr.* **122**, 1417–1424 (1992).

35.   Beutler, E. & West, C. Hematologic differences between African-Americans and whites: The roles of iron deficiency and α-thalassemia on hemoglobin levels and mean corpuscular volume. *Blood* **106**, 740–745 (2005).

36.   Garn, S. M. Lower Hematocrit Levels in Blacks Are Not Due to Diet or Socioeconomic Factors. *Pediatrics* **67**, (1981).

37.   Kanias, T. *et al.* Ethnicity, sex, and age are determinants of red blood cell storage and stress hemolysis: Results of the REDS-III RBC-Omics study. *Blood Adv.* **1**, 1132–1141 (2017).

38.   Page, G. *et al.* GWAS of Osmotic Hemolysis in 12,352 Healthy Blood Donors Identifies Red Cell Genetic Variants Associated with Steady State Hemolysis in Patients with Sickle Cell Disease. *Blood* **130**, 1117–1117 (2017).

39.   Campbell, M. C. & Tishkoff, S. A. African Genetic Diversity: Implications for Human Demographic History, Modern Human Origins, and Complex Disease Mapping. *Annu. Rev. Genomics Hum. Genet.* **9**, 403–433 (2008).

40.   Gallagher, P. G. Abnormalities of the erythrocyte membrane. *Pediatric Clinics of North America* vol. 60 1349–1362 (2013).

41.   Schulman, S. *et al.* Growth of Plasmodium falciparum in human erythrocytes containing abnormal membrane proteins. *Proc. Natl. Acad. Sci. U. S. A.* **87**, 7339–7343 (1990).

42.   Facer, C. A. *Erythrocytes carrying mutations in spectrin and protein 4.1 show differing sensitivities to invasion by Plasmodium falciparum. Parasitol Res* vol. 81 (1995).

43.   Dhermy, D., Schrével, J. & Lecomte, M.-C. Spectrin-based skeleton in red blood cells and malaria. *Curr. Opin. Hematol.* **14**, 198–202 (2007).

44.   Galanello, R. & Cao, A. Alpha-thalassemia. *Genetics in Medicine* vol. 13 83–88 (2011).

45.   Yoshida, A., Beutler, E. & Motulsky, A. G. Human glucose-6-phosphate dehydrogenase

822   variants. *Bull. World Health Organ.* **45**, 243–253 (1971).

823   46.   Clarke, G. M. *et al.* Characterisation of the opposing effects of G6PD deficiency on

824   cerebral malaria and severe malarial anaemia. *Elife* **6**, (2017).

825   47.   Cao, A. & Galanello, R. Beta-thalassemia. *Genetics in Medicine* vol. 12 61–76 (2010).

826   48.   Glogowska, E. *et al.* Novel mechanisms of PIEZO1 dysfunction in hereditary

827   xerocytosis. *Blood* **130**, 1845–1856 (2017).

828   49.   Greene, L. S. G6PD deficiency as protection againstfalciparum malaria: An

829   epidemiologic critique of population and experimental studies. *Am. J. Phys. Anthropol.*

830   **36**, 153–178 (1993).

831   50.   Ifediba, T., Stern, A., Ibrahim, A. & Rieder, R. Plasmodium falciparum in vitro:

832   diminished growth in hemoglobin H disease erythrocytes. *Blood* **65**, 452–455 (1985).

833   51.   Friedman, M. J. Erythrocytic mechanism of sickle cell resistance to malaria. *Proc. Natl.*

834   *Acad. Sci. U. S. A.* **75**, 1994–1997 (1978).

835   52.   Wright, G. J. & Rayner, J. C. Plasmodium falciparum Erythrocyte Invasion: Combining

836   Function with Immune Evasion. *PLoS Pathog.* **10**, (2014).

837   53.   Tibshirani, R. & Tibshirani, R. Regression Shrinkage and Selection Via the Lasso. *J. R.*

838   *Stat. Soc. Ser. B* **58**, 267--288 (1994).

839   54.   Ioannidis, J. P. A. Why Most Published Research Findings Are False. *PLoS Med.* **2**, e124

840   (2005).

841   55.   Flynn, C. J., Hurvich, C. M. & Simonoff, J. S. On the sensitivity of the Lasso to the

842   number of predictor variables. *Stat. Sci.* **32**, 88–105 (2017).

843   56.   Ma, S. *et al.* Common PIEZO1 Allele in African Populations Causes RBC Dehydration

844   and Attenuates Plasmodium Infection. *Cell* **173**, 443-455.e12 (2018).

845   57.   Nguetse, C. N. *et al.* A common polymorphism in the mechanosensitive ion channel

846   PIEZO1 is associated with protection from severe malaria in humans. *Proc. Natl.*

847   *Acad. Sci.* (2020) doi:10.1073/pnas.1919843117/-/DCSupplemental.

848   58.   Rooks, H. *et al.* A gain of function variant in PIEZO1 (E756del) and sickle cell disease.

849   *Haematologica* vol. 104 e91–e93 (2019).

59.   Ilboudo, Y. *et al.* A common functional PIEZO1 deletion allele associates with red

blood cell density in sickle cell disease patients. *American Journal of Hematology* vol.
93 E362–E365 (2018).

60.   Lessard, S. *et al.* An erythroid-specific ATP2B4 enhancer mediates red blood cell
hydration and malaria susceptibility. *J. Clin. Invest.* **127**, 3065–3074 (2017).

61.   Lin, S.-H., Brown, D. & Machiela, M. LDtrait: an online tool for identifying published
phenotype associations in linkage disequilibrium. *Cancer Res.* canres.0985.2020
(2020) doi:10.1158/0008-5472.CAN-20-0985.

62.   Crosnier, C. *et al.* Basigin is a receptor essential for erythrocyte invasion by
Plasmodium falciparum. *Nature* **480**, 534–537 (2011).

63.   Egan, E. S. *et al.* Erythrocytes lacking the Langereis blood group protein ABCB6 are
resistant to the malaria parasite Plasmodium falciparum. *Commun. Biol.* **1**, 1–9
(2018).

64.   Egan, E. S. *et al.* A forward genetic screen identifies erythrocyte CD55 as essential for
Plasmodium falciparum invasion. *Science (80-.).* **348**, 711–714 (2015).

65.   GTEx Consortium. Genetic effects on gene expression across human tissues. *Nature*
550, 204–213 (2017).

66.   Goheen, M. M. *et al.* Anemia Offers Stronger Protection Than Sickle Cell Trait Against
the Erythrocytic Stage of Falciparum Malaria and This Protection Is Reversed by Iron
Supplementation. *EBioMedicine* **14**, 123–130 (2016).

67.   Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation
in 141,456 humans. *Nature* **581**, 434–443 (2020).

68.   Williams, T. N. Human red blood cell polymorphisms and malaria. *Current Opinion in*
Microbiology vol. 9 388–394 (2006).

69.   Bejon, P. *et al.* Defining Childhood Severe Falciparum Malaria for Intervention
Studies. *PLoS Med.* **4**, e251 (2007).

70.   Allison, A. C. Protection afforded by sickle-cell trait against subtertian malarial
infection. *Br. Med. J.* **1**, 290–294 (1954).

71.   Luzzatto, L. Sickle cell anaemia and malaria. *Mediterranean Journal of Hematology*
and Infectious Diseases vol. 4 2012065 (2012).

72.   Okwa, O. O. *Malaria Parasites*. *Malaria Parasites* (InTech, 2012). doi:10.5772/1477.

73.   Pengon, J., Svasti, S., Kamchonwongpaisan, S. & Vattanaviboon, P. Hematological

parameters and red blood cell morphological abnormality of Glucose-6-Phosphate

dehydrogenase deficiency co-inherited with thalassemia. *Hematol. Oncol. Stem Cell*

Ther. **11**, 18–24 (2018).

74.   Clark, M., Mohandas, N. & Shohet, S. Osmotic gradient ektacytometry: comprehensive

characterization of red cell volume and surface maintenance. *Blood* **61**, 899–910

(1983).

75.   Mockenhaupt, F. P. *et al.* Anaemia in pregnant Ghanaian women: importance of

malaria, iron deficiency, and haemoglobinopathies. *Trans. R. Soc. Trop. Med. Hyg.* **94**,

477–483 (2000).

76.   Tiffert, T. *et al.* The hydration state of human red blood cells and their susceptibility

to invasion by Plasmodium falciparum. *Blood* **105**, 4853–4860 (2005).

77.   Kariuki, S. N. *et al.* Red blood cell tension controls Plasmodium falciparum invasion

and protects against severe malaria in the Dantu blood group. *bioRxiv* 475442 (2018)

doi:10.1101/475442.

78.   Field, S. P., Hempelmann, E., Mendelow, B. V. & Fleming, A. F. Glycophorin variants

and Plasmodium falciparum: protective effect of the Dantu phenotype in vitro. *Hum.*

Genet. **93**, 148–150 (1994).

79.   Bei, A. K. *et al.* A flow cytometry-based assay for measuring invasion of red blood

cells by *Plasmodium falciparum*. *Am. J. Hematol.* **85**, 234–237 (2010).

80.   Awandare, G. A., Nyarko, P. B., Aniweh, Y., Ayivor-Djanie, R. & Stoute, J. A.

Plasmodium falciparum strains spontaneously switch invasion phenotype in

suspension culture. *Sci. Rep.* **8**, 1–10 (2018).

81.   Dolan, S. A., Miller, L. H. & Wellems, T. E. Evidence for a switching mechanism in the

invasion of erythrocytes by Plasmodium falciparum. *J. Clin. Invest.* **86**, 618–624

(1990).

82.   Boyle, M. J. *et al.* Isolation of viable Plasmodium falciparum merozoites to define

erythrocyte invasion events and advance vaccine and drug development. *Proc. Natl.*

Acad. Sci. U. S. A. **107**, 14378–14383 (2010).

83.   Chen, R., Davydov, E. V., Sirsota, M. & Butte, A. J. Non-synonymous and synonymous

coding SNPS show similar likelihood and effect size of human disease association.

PLoS One **5**, (2010).

84.   Sauna, Z. E. & Kimchi-Sarfaty, C. Understanding the contribution of synonymous

mutations to human disease. *Nature Reviews Genetics* vol. 12 683–691 (2011).

85.   Kinsler, G., Geiler-Samerotte, K. & Petrov, D. A genotype-phenotype-fitness map

reveals local modularity and global pleiotropy of adaptation. *bioRxiv*

2020.06.25.172197 (2020) doi:10.1101/2020.06.25.172197.

86.   Gusev, A. *et al.* Partitioning heritability of regulatory and cell-type-specific variants

across 11 common diseases. *Am. J. Hum. Genet.* **95**, 535–552 (2014).

87.   Sohail, M. *et al.* Polygenic adaptation on height is overestimated due to uncorrected

stratification in genome-wide association studies. *Elife* **8**, (2019).

88.   Otto, T. D. *et al.* Genomes of all known members of a Plasmodium subgenus reveal

paths to virulent human malaria. *Nat. Microbiol.* **3**, 687–697 (2018).

89.   Sundararaman, S. A. *et al.* Genomes of cryptic chimpanzee Plasmodium species reveal

key evolutionary events leading to human malaria. *Nat. Commun.* **7**, (2016).

90.   Pankratov, V. *et al.* Differences in local population history at the finest level: the case

of the Estonian population. *Eur. J. Hum. Genet.* 1–12 (2020) doi:10.1038/s41431-020-

0699-4.

91.   Donovan, B. M. Learned inequality: Racial labels in the biology curriculum can affect

the development of racial prejudice. *J. Res. Sci. Teach.* **54**, 379–411 (2017).

92.   Sparks, R. A., Baldwin, K. E. & Darner, R. Using Culturally Relevant Pedagogy to

Reconsider the Genetics Canon †. *J. Microbiol. Biol. Educ.* **21**, (2020).

93.   Rosenberg, N. A. A Population-Genetic Perspective on the Similarities and Differences

among Worldwide Human Populations. *Hum. Biol.* **83**, 659–684 (2011).

94.   Coop, G. *et al.* The Role of Geography in Human Adaptation. *PLoS Genet.* **5**, e1000500

(2009).

95.   Hofer, T., Ray, N., Wegmann, D. & Excoffier, L. Large allele frequency differences

between human continental groups are more likely to have occurred by drift during

range expansions than by selection. *Ann. Hum. Genet.* **73**, 95–108 (2009).

96.   Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. *arXiv* (2013).

97.   Van der Auwera, G. A. *et al.* From fastQ data to high-confidence variant calls: The

genome analysis toolkit best practices pipeline. *Curr. Protoc. Bioinforma.* **11**, 11.10.1

(2013).

98.   Mills, R. E. *et al.* An initial map of insertion and deletion (INDEL) variation in the
human genome. *Genome Res.* **16**, 1182–1190 (2006).

99.   Weatherall, D. J. Phenotype-genotype relationships in monogenic disease: Lessons
from the thalassaemias. *Nature Reviews Genetics* vol. 2 245–255 (2001).

100.   Wang, K., Li, M. & Hakonarson, H. ANNOVAR: Functional annotation of genetic
variants from high-throughput sequencing data. *Nucleic Acids Res.* **38**, (2010).

101.   Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
(2011).

102.   The 1000 Genomes Project Consortium. A global reference for human genetic
variation. *Nature* vol. 526 68–74 (2015).

103.   Lowy-Gallego, E. *et al.* Variant calling on the GRCh38 assembly with the data from
phase three of the 1000 Genomes Project. *Wellcome Open Res.* **4**, 50 (2019).

104.   Purcell, S. *et al.* PLINK: A tool set for whole-genome association and population-
based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

105.   Alexander, D. H. & Lange, K. Enhancements to the ADMIXTURE algorithm for
individual ancestry estimation. *BMC Bioinformatics* **12**, 246 (2011).

106.   Park, D. J. *et al.* Sequence-based association and selection scans identify drug
resistance loci in the Plasmodium falciparum malaria parasite. *Proc. Natl. Acad. Sci. U.*
S. A. **109**, 13052–13057 (2012).

107.   Kuypers, F. A., Scott, M. D., Schott, M. A., Lubin, B. & Chiu, D. T. Y. Use of ektacytometry
to determine red cell susceptibility to oxidative stress. *J. Lab. Clin. Med.* **116**, 527–534
(1990).

108.   Chatterjee, S. Assumptionless consistency of the Lasso. *arXiv* (2013).

968