TITLE

Common host variation drives malaria parasite fitness in healthy human red cells

AUTHORS

Emily R Ebel[1,2], Frans A Kuypers[3], Carrie Lin[2], Dmitri A Petrov[1*], Elizabeth S Egan[2,4*]

[1]Department of Biology, Stanford University, Stanford, CA 94305

[2]Department of Pediatrics, Stanford University School of Medicine, Stanford, CA 94305

[3]Children's Hospital Oakland Research Institute, Oakland, CA, 94609

[4]Department of Microbiology & Immunology, Stanford University School of Medicine, Stanford, CA 94305

*co-corresponding authors: eegan@stanford.edu; dpetrov@stanford.edu

SUMMARY

The replication of *Plasmodium falciparum* parasites within red blood cells (RBCs) causes severe disease in humans, especially in Africa. The influence of host RBC variation on parasite replication is largely uncharted, aside from a handful of deleterious alleles like sickle cell. Here, we integrated analyses of exome sequencing, RBC phenotyping, and parasite fitness assays on blood from 122 individuals, most with African ancestry. In donors lacking alleles for hemoglobinopathies or G6PD deficiency, RBC phenotypes including size, deformability, and hydration status explained 21-38% of the variation in parasite growth rate. With the addition of non-pathogenic polymorphisms in 14 RBC proteins including *SPTA1*, *PIEZO1,* and *ATP2B4,* our model explained 73-82% of the variation in parasite growth rate*.* Interestingly, we observed little evidence for divergent selection on this variation between Africans and Europeans. These findings suggest a model in which widespread, non-pathogenic variation in a moderate number of genes strongly modulates *P. falciparum* fitness in RBCs.

KEY WORDS: malaria, *Plasmodium falciparum*, red blood cells, parasite fitness, natural variation

INTRODUCTION

Malaria caused by the replication of *Plasmodium falciparum* parasites in red blood cells (RBCs) kills hundreds of thousands of children each year (WHO, 2019). In each 48-hour cycle of blood stage malaria, parasites must deform RBC membranes to invade them(Koch *et al.*, 2017; Kariuki *et al.*, 2020); consume hemoglobin and tolerate the resulting oxidative stress (Francis, Sullivan and Goldberg, 1997); multiply to displace half the RBC volume (Hanssen *et al.*, 2012); and remodel the RBC membrane to avoid immune detection (Zhang *et al.*, 2015). Consequently, genetic disorders that alter aspects of RBC biology are well known to influence malaria susceptibility (Kwiatkowski, 2005). For example, sickle cell trait impairs parasite growth by altering hemoglobin polymerization at low oxygen tension (Pasvol, Weatherall and Wilson, 1978; Archer *et al.*, 2018), while deficiency of the G6PD enzyme involved in oxidative stress tolerance is thought to make parasitized RBCs more susceptible to breakdown (Ruwende and Hill, 1998). Aside from these diseases, however, the genetic basis of malaria susceptibility remains mostly unknown. Several large genome-wide association studies (GWAS) have recently made progress on this question by identifying a few dozen loci that collectively explain up to 11% of the heritability of the risk of severe versus uncomplicated malaria (Timmann *et al.*, 2012; Rockett *et al.*, 2014; Band *et al.*, 2015; Leffler *et al.*, 2017; Ndila *et al.*, 2018; MGE Network, 2019a). Eleven of these loci are in or near genes expressed predominantly in RBCs, and one new GWAS variant has since been shown to regulate expression of the *ATP2B4* calcium channel (Zámbó *et al.*, 2017) and to be associated with RBC dehydration (Li *et al.*, 2013). However, most other RBC loci that have been identified by GWAS are not novel, and a functional link between *ATP2B4* and *P. falciparum* replication has yet to be demonstrated. More discoveries about the influence of RBC variation on malaria susceptibility are not expected from similar GWAS (Boyle, Li and Pritchard, 2017; MGE Network, 2019b), in part because severe malaria is a complex phenotype that depends on a combination of factors from RBCs, the vascular endothelium, the immune system, and the environment (Mackinnon *et al.*, 2005; De Mendonça, Goncalves and Barral-Netto, 2012). Alternate approaches are therefore needed to discover more genetic variation that impacts the replication of malaria parasites in human RBCs.

RBC phenotypes like mean cell volume (MCV), hemoglobin content (HGB/MCH), and

antigenic blood type vary widely within and between human populations (Lo *et al.*, 2011;

Cooling, 2015; Canela-Xandri, Rawlik and Tenesa, 2018). 40-90% of the variation in

common RBC indices is heritable (Whitfield, Martin and Rao, 1985; Evans, Frazer and

Martin, 1999; Pilia *et al.*, 2006), and GWAS have demonstrated that these phenotypes have

polygenic bases (van der Harst *et al.*, 2012; Astle *et al.*, 2016; Chami *et al.*, 2016; Chen *et al.*,

2020; Vuckovic *et al.*, 2020). Average hemoglobin levels, hematocrit, and especially RBC

membrane fragility have also been shown to differ between African and European

populations, likely for genetic reasons (Garn, 1981; Perry *et al.*, 1992; Beutler and West,

2005; Kanias *et al.*, 2017; Page *et al.*, 2017). Although African populations are highly

genetically diverse(Campbell and Tishkoff, 2008), some of these differences between

Africans and Europeans are explained by RBC disease alleles (Beutler and West, 2005; Lo *et*

al., 2011; Kanias *et al.*, 2017) that have been widely selected across Africa for their

protective effects on malaria. It remains untested whether other extensive phenotypic and

genetic diversity in RBCs, aside from these disease alleles, also influences malaria

susceptibility and has been similarly shaped by malaria selection.

Here, we approach these questions by performing exome sequencing and extensive

RBC phenotyping on blood samples from a diverse human cohort of 122 individuals. We

show that *P. falciparum* fitness varies widely among donor cells *in vitro*, with the

distributions of parasite phenotypes in 'healthy' RBCs overlapping those from RBCs

carrying classic disease alleles. We apply Lasso variable selection to identify a small set of

alleles and phenotypes that strongly predict parasite fitness outside of the context of RBC

disease, highlighting RBC membrane and dehydration genes and phenotypes as key to

modulating *P. falciparum* fitness. However, we find no evidence that non-pathogenic alleles

or phenotypes that confer parasite protection are associated with African ancestry.

Overall, these findings advance our understanding of the origin and function of common

RBC variation and suggest new targets for therapeutic intervention for malaria.

RESULTS

Many healthy blood donors with African ancestry carry alleles for RBC disease

We collected blood samples from 121 donors with no known history of blood

Figure 1. Overview of study design and blood donors. (a) PCA of genetic variation across 35,759 unlinked
exome SNPs. Samples collected in this study are plotted on coordinate space derived from 1000 Genomes
reference populations. Points with white borders represent the six related individuals in this study. **(b)**
Exome sequencing revealed that 37% of donors carried alleles for RBC disorders linked to *P. falciparum*
resistance. Individuals with >1 disease allele were assigned to the category of the most severe condition.
non-carrier: Donor without any of the following alleles or conditions. **G6PD⁻low:** Mild to medium G6PD
deficiency (<42% loss of function). **G6PD⁻high:** Severe G6PD deficiency (>60% loss of function). **-α/αα:**
heterozygous HBA2 deletion, or alpha thalassemia minima. **-α/-α:** homozygous HBA2 deletion, or alpha

103 thalassemia trait. **HbAC**: heterozygous HBB:E7K, or hemoglobin C trait. **HbAS:** heterozygous HBB:E7V, or
104 sickle cell trait. **HE:** hereditary elliptocytosis. **(c)** Parasite replication rate in donor RBCs was measured via
105 flow cytometry at three timepoints over 72 hours. 3D7 is a lab-adapted strain used in routine culture.
106 Th.026.09 is a recent clinical isolate from Senegal. **(d)** RBC phenotypes were measured using complete blood
107 counts with RBC indices, osmotic fragility tests, and ektacytometry.
108

disorders, most of whom self-identified as having recent African ancestry (Figure 1A). As a

positive control, we also sampled a patient with hereditary elliptocytosis (HE), a polygenic

condition characterized by extremely fragile RBC membranes that strongly inhibit *P.*

falciparum growth (Schulman *et al.*, 1990; Facer, 1995; Dhermy, Schrével and Lecomte,

2007; Gallagher, 2013). We performed whole-exome sequencing, both to check for the

presence of known RBC disease alleles and to confirm the population genetic ancestry of

our donors. A principal component analysis of more than 35,000 unlinked, exomic SNPs

showed that most donors fell along a continuum from African to European ancestry, as

defined by data from the 1000 Genomes Project (Figure 1A). We found that 16% of the

healthy donors carried pathogenic hemoglobin alleles (Figure 1B), including 5

heterozygotes for hemoglobin S (HbAS), 4 heterozygotes for hemoglobin C (HbAC), and 11

individuals with one or two copies of an *HBA2* deletion causing α-thalassemia (Galanello

and Cao, 2011). We also scored eight polymorphisms in *G6PD* that have been functionally

associated with various degrees of G6PD deficiency (Yoshida, Beutler and Motulsky, 1971;

Clarke *et al.*, 2017) and found that 32% of the study population carried at least one,

including 12 of the 20 donors with hemoglobinopathies. Among those with wild-type

hemoglobin, we identified one individual with polymorphisms associated with severe

G6PD deficiency (>60% loss of function) and 23 with polymorphisms associated with mild

to medium deficiency (<42% loss of function). To our knowledge, we detected no alleles

linked to other monogenic RBC disorders, including β-thalassemia or xerocytosis (Cao and

Galanello, 2010; Glogowska *et al.*, 2017). We therefore classified the remaining 73 donors

as 'non-carriers' of known disease alleles for the purposes of this work.

***P. falciparum* replication rates vary widely among non-carrier RBCs**

To determine the variation in *P. falciparum* fitness among samples with different

genotypes, we performed invasion and growth assays with two strains of parasite: a

laboratory adapted strain (3D7) and a recent clinical isolate from Senegal (Th.026.09).

We observed a wide range of *P. falciparum* growth rates among RBC samples, especially

among non-carriers that lacked known disease alleles (Figure 2A-C). Each strain's growth

rate is defined here as parasite multiplication over a full 48-hour cycle in donor RBCs

(Figure 1C), with the mean value for non-carriers set to 100% after normalization. Briefly,

we used a repeated control RBC sample (Figure 2, gray circles) and other batch-specific

factors to correct for variation in parasite growth across multiple experiments (see

Methods). Among non-carriers, growth rates ranged from 64-136% for 3D7 (SD = 17.7%)

and 76-128% for Th.026.09 (SD = 10.7%) (Figure 2A-B). Per-sample growth rates were

strongly correlated between the two strains (Figure 2C, $p < 3 \times 10^{-16}$) and positively

correlated when measured in different weeks (Figure S1A, $\rho = 0.34$), demonstrating that

these data capture real variation among donor RBCs. Furthermore, as expected (Friedman,

1978; Ifediba *et al.*, 1985; Greene, 1993; Facer, 1995), we detected reductions in mean

growth rate for both strains in RBCs carrying known disease alleles. These included severe

G6PD deficiency (3D7, 60% of the non-carrier average, p<0.014; Th.026.09, 64%,

p<0.014), α-thalassemia trait (3D7, 75%, p=0.027; Th.026.09, 81%, p = 0.077), HbAS (3D7,

16%, $p=1.05 \times 10^{-7}$; Th.026.09, 19%, $p =1.2 \times 10^{-4}$), and HE (3D7, 18%, p<0.014; Th.026.09,

33%, p<0.014). Notably, the wide distribution of growth rates for non-carrier RBCs had

considerable overlap with the growth rates in carrier RBCs. Only the HbAS and HE samples

fell entirely outside the non-carrier range. This observation implies the existence of

previously unknown RBC variation that impacts *P. falciparum* growth, which may have

effect sizes comparable to the milder known disease alleles.

We observed a similarly wide range in the efficiency of *P. falciparum* invasion into

donor RBCs (Figure 2D-F). Invasion is defined here as the fold-change in parasitemia over

the first 24 hours of the assay, when parasites previously maintained in standard culture

conditions egressed and invaded new donor RBCs (Figure 1C). Among non-carriers,

invasion rates ranged from 66-173% for 3D7 (SD=15.8%) and 41-193% for Th.026.09

(SD=28.9%) (Figure 2D-E). Compared to growth rates, no disease alleles conferred

protection against invasion that was extreme enough to fall outside the broad non-carrier

range. In 3D7 only, HbAC was associated with a 9% decrease in invasion (p=0.019) and

Figure 2. ***P. falciparum*** **replication rate varies widely among donor RBCs. (a-b)** Growth of *P. falciparum* strain 3D7 **(a)** or clinical isolate Th026.09 **(b)** over a full 48-hour cycle in donor RBCs (hours 24-72 of assay). Growth is presented relative to the average non-carrier rate after correction for batch effects, including comparison to a repeated RBC control (Methods). Repeated measurements of the same RBC control over the course of the experiment are shown as open gray circles. **(c)** Correlation in per-sample growth rates between the two *P. falciparum* strains. **(d-e)** Invasion efficiency of *P. falciparum* strain 3D7 **(d)** or clinical isolate Th026.09 **(e)** after a 24-hour incubation with donor RBCs, normalized in the same way as growth. Repeated measurements of the RBC control over time are shown as gray circles. **(f)** Correlation in per-sample invasion success for the two *P. falciparum* strains. Colors in **(c)** and **(f)** correspond to the classifications in other panels; R^2 and *p*-values are derived from OLS regression. *p<0.1; **p<0.05; ***p<0.01.

mild G6PD deficiency was associated with a 7% increase (p=0.072)*.* Only HE had a

significant effect on the invasion efficiency of both strains (3D7, 67% of the non-carrier

average, p<0.014; Th.026.09, 57%, p<0.014). There was little correlation between the

invasion efficiencies of the two parasite strains in individual donor samples (Figure 2F),

potentially reflecting strain-specific differences in the pathways used for invasion (Wright

and Rayner, 2014). However, as we also observed greater variability between repeated

samples for invasion than for growth (Figure S1), the invasion results may be influenced by

greater experimental noise.

RBC phenotypes vary widely among non-carriers

To assess phenotypic variation across donor RBCs, we measured 19 common

indices of RBC size and hemoglobin content from complete blood counts using an Advia

hematology analyzer (Figure 3A-D; Figure S2). Mean cellular volume (MCV) and

hemoglobin mass (MCH) are closely related traits, which can be represented together as

cellular hemoglobin concentration (CHCM) or the fraction of RBCs with 'normal'

hemoglobin and volume indices (M5). As expected, each known disease allele was

associated with a distinct set of RBC abnormalities. These included elevated CHCM for

HbAC (p=0.033), consistent with dehydration, and very low MCV ($p=8.7 \times 10^{-5}$) and MCH

($p=4.6 \times 10^{-7}$) for α-thalassemia trait ($\alpha\alpha$/--), consistent with microcytic anemia (Galanello

and Cao, 2011). RBCs from the HE patient also had very low MCV and MCH (both p<0.014),

reflecting the membrane breakage and volume loss characteristic of this disease. However,

for all of these phenotypic measures, we also observed broad distributions in non-carriers

that overlapped the distributions of most carriers (Figure 3A-D; Figure S2). Notably, the

breadth of the non-carrier distribution for each phenotype was large (e.g., 24 fL range for

MCV) compared to the average difference between Africans and Caucasians (e.g., 3-5

fL(Beutler and West, 2005; Lo *et al.*, 2011)). This wide diversity and substantial overlap

between non-carrier and carrier traits suggest that 'healthy' RBCs exist on the same

phenotypic continuum as RBCs carrying known disease alleles.

We observed similar patterns of variation in RBC membrane fragility (Figure 3E-F)

and membrane deformability (Figure 3G-H), as measured with osmotic fragility tests and

osmotic gradient ektacytometry (see Methods). Both sets of curves represent RBC

tolerance to osmotic stress, which can result in swelling and lysis (fragility, Figure 3E-F) or

dehydration and decreased deformability (O_{hyper}, Figure 3G-H). Specific

hemoglobinopathies were associated with moderate to strong reductions in fragility ($\alpha\alpha$/α-

p=0.039; $\alpha\alpha$/-- p=0.003; HbAS p=0.014), deformability (DI_{max}; HbAC p=0.007; HbAS

$p=4.4 \times 10^{-6}$), and/or resistance to loss of deformability when dehydrated (O_{hyper}; HbAC

Figure 3. Red cell phenotypes that are abnormal in carriers also vary widely among non-carriers.
MCV: mean RBC volume. **MCH:** mean cellular hemoglobin. **CHCM:** cellular hemoglobin concentration. **M5:**
fraction of RBCs with 'normal' indices of hemoglobin concentration (28-41 g/dL) and volume (60-120 fL).
Fragility: Tendency to lyse when osmotically overhydrated, defined as [NaCl] causing 50% lysis. **DI_max:**
Maximum membrane deformability under physiological salt conditions. **O_hyper:** Tendency to resist osmotic
dehydration and loss of deformability, defined as the hypertonic [NaCl] at which deformability is half
maximum. *p<0.1; **p<0.05; ***p<0.01; tests and samples as in **Figure 2.**

p=0.019; HbAS p=0.041). HE cells were both extremely fragile (p<0.014) and extremely

non-deformable (p<0.014). In non-carriers, the distributions for all membrane measures

were wide, continuous, and overlapped the distribution of most carriers (Figure 3E-H;

Figure S3). Overall, these data demonstrate that multiple phenotypic alterations associated

with RBC disease alleles are also present in non-carrier RBCs.

Non-carrier variation in RBC phenotypes predicts *P. falciparum* replication rate

To identify sets of phenotypes associated with *P. falciparum* replication in non-

carrier RBCs, we used a machine learning method called Lasso (least absolute shrinkage

and selection operator) that performs regularization and variable selection (Tibshirani and

Tibshirani, 1994) (see Methods). Briefly, Lasso shrinks the regression coefficients for some

possible predictors (in this case, phenotypes) to zero, in order to obtain a subset of

predictors that minimizes prediction error. This method is well-suited for datasets in which

possible predictors are correlated, as are standard measurements of RBC size, hemoglobin,

and membrane dynamics. For each trait in each predictive subset selected by Lasso, we

then applied univariate OLS regression to more accurately estimate the sign of its effect on

all measured components of parasite fitness. The highest-confidence results are

summarized in Figure 4A, with complete details provided in Table S2.

P. falciparum fitness in non-carrier RBCs was strongly predicted by variation in

several RBC traits related to deformability, dehydration, volume, and hemoglobin content.

Among 24 tested phenotypes, the most important trait was the ektacytometry parameter

O_{hyper}, which represents a cell's tendency to retain deformability in the face of dehydration

(Figure 3G). In univariate models, non-carrier RBCs with the largest O_{hyper} values supported

1.2-1.5X faster parasite growth (3D7 1.47X, p=0.003; Th.026.09 1.24X, p=0.005) and 1.4-

2.0X more effective invasion (3D7 1.36X, p=0.018; Th.026.09 1.94X, p=0.006) than RBCs

with the smallest O_{hyper} values (Table S2; Figure S5A). Consistent with this result, *P.*

falciparum replication was relatively inhibited in RBCs that were more dehydrated at

baseline, as indicated by higher CHCM (3D7 invasion p=0.007; 3D7 growth p=0.042;

Th.026.09 invasion p=0.006; Th.026.09 growth p=0.36). Parasites also grew faster in RBCs

with larger mean volume, or MCV (3D7 p=0.001; Th.026.09, p=0.001); a greater mass of

hemoglobin per cell, or MCH (3D7 p=0.073; Th.026.09 p=0.019); and more deformable

Figure 4. RBC phenotypes predict *P. falciparum* fitness in non-carriers. **(a)** Phenotypes shown were
selected by Lasso in at least one of four models (columns) with at least 60% cross-validation (CV) support, as
indicated by blue shading. **(+/-)** shows the direction of effect if the phenotype was significantly correlated
($p<0.1$) with the parasite fitness component in a separate, univariate linear model. **(b)** Lasso model
predictions of parasite growth as a function of RBC phenotypes are plotted against measured growth values.
Each point represents one non-carrier. The solid line shows a perfect 1:1 fit; R^2 indicates the proportion of
variance explained by Lasso predictions. **(c)** For 1000 permutations of the growth data, the histograms show
the proportion of variance explained by the Lasso modeling procedure. Dashed lines indicate R^2 for the actual
data. p-values indicate the fraction of permutations that explain at least as much variance as the real data.
(d,e) As in **(b,c)**, but for *P. falciparum* invasion. **O_{hyper}:** Tendency to resist osmotic dehydration and loss of
deformability, defined as the hypertonic [NaCl] at which deformability is half maximum (mM). **MCV:** mean
RBC volume (fL). **DI_{max}:** Maximum membrane deformability under physiological salt conditions. **CHCM:**
cellular hemoglobin concentration (g/dL). **Fragility:** Tendency to lyse when osmotically overhydrated,
defined as [NaCl] causing 50% lysis (mM). **M5:** fraction of RBCs with 'normal' indices of hemoglobin
concentration (28-41 g/dL) and volume (60-120 fL) (%). **MCH:** mean corpuscular hemoglobin (pg/RBC).
MPV: mean platelet volume (fL). **RBC**: red cell number (x10^6/uL). **PLT:** platelet number (x10^3/uL).). **HCT:**
hematocrit (%).

membranes, measured by DI_{max} (3D7 p=0.0006; Th.026.09 p=0.006). Some measures of

parasite fitness were also reduced in RBCs with more fragile membranes (3D7 invasion

p=0.65; 3D7 growth p=0.003; Th.026.09 invasion p=0.067; Th.026.09 growth p=0.14).

Additional phenotypes related to platelets and RBC density were selected for some models,

but the direction of their effects was unclear since they were not significant when

considered individually. These results indicate that common, non-pathogenic variation in

RBC size, membrane dynamics, and other correlated traits have meaningful effects on *P.*

falciparum replication rate in RBCs.

Together, 13 phenotype measurements explained approximately 40% of the

variation in 3D7 growth and invasion in non-carriers (Figure 4B,D). To test for overfitting,

we applied the same model-fitting procedure many times to randomly shuffled parasite

data, which usually failed to produce predictive models (i.e., R^2=0, Figure 4C,E). These

permutation tests found that real data explained, on average, 6.5X the variance in 3D7

growth and 6.3X the variance in 3D7 invasion than expected by chance (both p=0.01,

Figure 4C,E). For Th.026.09, the phenotypes selected by Lasso explained 17-21% of the

total variation in parasite fitness (Figure 4B), which was marginally better than expected

by chance for growth (3.7X, p=0.07, Figure 4C) and marginally worse than expected for

invasion (3.0X, p=0.13, Figure 4E). The weaker result for Th.026.09 may be explained by its

smaller dynamic growth range (Figure 2A-B), perhaps because clinical isolates are

relatively poorly adapted to laboratory conditions. Overall, these results demonstrate that

multiple, variable phenotypes impact *P. falciparum* susceptibility in healthy RBCs. Non-

carrier cells that are less hospitable to parasites share specific traits with RBCs that carry

disease alleles, including smaller size, decreased deformability, and an increased tendency

to lose deformability when dehydrating.

A small set of RBC genotypes predicts *P. falciparum* replication in non-carriers

Next, we tested whether non-carrier genotypes derived from exome sequencing could

improve our predictions of *P. falciparum* replication rate. To limit false positives and

increase statistical power (Ioannidis, 2005; Flynn, Hurvich and Simonoff, 2017), we

focused on 23 candidate RBC proteins that have been previously associated with *P.*

falciparum resistance in the literature (Table S3). These proteins contained 106 unlinked

genetic variants (pairwise r^2 < 0.1) among the 73 non-carriers. Along with these

polymorphisms and phenotypes described above, we also treated the top 10 principal

components (PCs) of 1000 Genomes population structure as possible predictors of *P.*

falciparum fitness.

The addition of genetic factors substantially improved all four models of parasite

fitness. Across strains, 73-82% of the parasite variation in non-carriers was explained by a

	3D7 growth	Th.026 growth	3D7 invasion	Th.026 invasion
Gene	associated variants			
Flexible membrane backbone				
SPTA1	3	3	0	0
SPTB	4	1	0	0
ANK1	0	2	1	0
SLC4A1	1	2	0	0
EPB42	1	0	0	0
Ion channels (hydration)				
PIEZO1	5	2	1	1
ATP2B4	1	0	0	0
Plasma membrane proteins				
CD44	2	2	0	0
CR1	1	1	1	1
BSG	1	2	0	0
ABCB6	1	0	0	0
GYPB	1	0	0	0
Other				
FPN	0	1	0	0
G6PD	0	1	0	0
	associated 1000G PCs			
of top 10	1	1	1	1

Figure 5. RBC genotypes substantially improve predictions of *P. falciparum* fitness in non-carriers. (a) Lasso model predictions of parasite growth as a function of RBC phenotypes, PCs of population structure, and variants in 23 malaria-related proteins (Table S3) are plotted against measured growth values. Each point represents one non-carrier. The solid line shows a perfect 1:1 fit; R^2 indicates the proportion of variance explained by Lasso-selected predictors. **(b)** For 1000 random sets of 23 genes drawn from the RBC proteome, the histograms show the variance explained *by variants within those genes* after applying the same Lasso modeling procedure as (a). Dashed lines indicate the variance explained by variants within the 23 malaria-related genes. *p*-values indicate the fraction of random gene sets that explain at least as much variance as the real data. **(c,d)** As in (a,b), but for *P. falciparum* invasion. **(e)** Variants and PCs counted in each cell were selected in the designated Lasso model with at least 70% cross-validation (CV) support. Several tested genes had 0 associated variants: *CD55*, *EPB41*, *HP*, *HBA1/2*, *GYPA*, *GYPC*, *GYPB*, and *HBB.* Details on lower-confidence variants are available in Table S4.

total of 47 genetic and phenotypic variables selected by Lasso (Figure 5A; Table S4). For 3D7 and Th.026.09 respectively, polymorphisms in 14 of 23 malaria-related genes explained 71% and 41% of the growth variation in non-carriers (Figure 5B). These models were significant by permutation (both p=0.011; Figure S6AB), indicating that the results were unlikely to be due to overfitting. Nonetheless, to confirm that the genetic signal was specific to our set of 23 malaria-related genes, we used polymorphisms from 1000 sets of random genes from the RBC proteome (Table S6) to build 1000 alternative models (see Methods). On average, variants from random RBC genes explained only 17.9% of the

observed variation in *P. falciparum* growth for 3D7 (p=0.002) and 16.4% for Th.026.09

(p=0.047) (Figure 5B). In contrast to growth, the genotype/phenotype models of *P.*

falciparum invasion (Figure 5C) were only marginally significant by permutation (3D7

p=0.09, Th.026.09 p=0.11, Figure S6C-D) and the genetic signal was not specific to the set of

23 malaria-related genes (3D7 p=0.26, Th.026.09 p=0.52, Figure 5D). Nonetheless, 4 of the

8 variants selected by Lasso for 3D7 invasion were also selected for 3D7 growth. Together,

these results indicate that *P. falciparum* growth rate in RBCs is strongly determined by host

genetic variation concentrated in a small number of genes.

Of the top 31 variants associated by Lasso with *P. falciparum* growth, seventeen

(55%) were synonymous (Table S4). This fraction did not significantly differ from the input

set of 106 variants (p=0.64, 2-tailed z-score). Although 15/17 of these synonymous

variants have no known, direct function, 7/15 are in linkage disequilibrium ($r^2 > 0.1$) with

other variants that have been associated with RBC phenotypes by GWAS (Table S4).

Similarly, 6 of the top 14 non-synonymous polymorphisms associated with *P. falciparum*

growth have been previously linked to RBC or other traits (Table S4). This overlap with

previous studies suggests that the variants associated with *P. falciparum* growth in this

study are likely to tag functional genetic loci.

Nearly all of the top 31 growth-associated polymorphisms occurred in either (1) ion

channel proteins, which regulate RBC hydration; (2) components of the flexible RBC

membrane backbone; or (3) red cell plasma membrane proteins, including known invasion

receptors (Figure 5E). In the first category, the highly polymorphic ion channel *PIEZO1*

contained six polymorphisms associated with small (-3%) to moderate (-32%) reductions

of *P. falciparum* growth rate (Table S4). The microsatellite variant *PIEZO1*-E756del, which

has been a focus of several recent studies (Ilboudo *et al.*, 2018; Ma *et al.*, 2018; Rooks *et al.*,

2019; Nguetse *et al.*, 2020), predicted a moderate reduction in Th.026.09 growth (-8.2%,

p=0.011) but was not related to RBC dehydration in these data (Table S5). We also detected

one growth-associated variant in *ATP2B4,* which encodes the primary RBC calcium channel

PMCA4b (-13% 3D7, p=0.006). This variant tags an *ATP2B4* haplotype implicated by GWAS

in protection from severe malaria (Timmann *et al.*, 2012; Lessard *et al.*, 2017; Zámbó *et al.*,

2017) and many RBC phenotypes (van der Harst *et al.*, 2012; Li *et al.*, 2013; Lessard *et al.*,

2017; Lin, Brown and Machiela, 2020) including RDW, which was replicated in our data set

(Table S4). *SPTA1* and *SPTB,* which encode the flexible spectrin backbone of RBCs, also

contained several variants associated with the growth of at least one *P. falciparum* strain,

as did the structural linker proteins ANK1, SLC4A1, and EPB42 (Figure 5E; Table S4).

Finally, we identified one polymorphism in ABCB6, two in CD44, and two in basigin (BSG)

that were strongly associated with *P. falciparum* growth (Figure 5E; Table S4). These

plasma membrane proteins have previously been implicated in *P. falciparum* invasion by *in*

vitro genetic deficiency studies (Crosnier *et al.*, 2011; Egan *et al.*, 2015, 2018). Notably, two

of the natural polymorphisms identified here are synonymous quantitative trait loci (QTL)

for CD44 splicing (rs35356320) and BSG expression (rs4682) (GTEx Consortium, 2017).

We also detected one variant in the RBC iron exporter *FPN* and one in *G6PD* with small

effects on Th.026.09 growth (2.4%, p=0.015 and 3.4%, p=0.010 respectively; Table S4). No

associated variants were detected in the other nine tested genes (Table S2), including three

hemoglobin proteins, three glycophorins, *CD55*, *EPB41*, and *HP*. Together, these data

demonstrate that *P. falciparum* growth rate in non-carrier RBCs is strongly determined by

dozens of host genetic variants, which together shape the phenotypic distribution of red

cell susceptibility.

In addition to these specific genetic variants, PCs of population structure explained

22-26% of the variation in non-carrier invasion and 10-12% of the variation in growth.

Interestingly, neither PC1 nor PC2—which distinguish Africans from other populations

(Figure 1A)—were related to parasite fitness. Instead, this signal of correlated variation

was partially driven by a six-member family with unique ancestry in our sample (Figure

1A, white borders) and extreme parasite fitness values (Figure S6). Although the present

study is limited by sample size, the association between global genetic PCs and *P.*

falciparum fitness suggests that additional functional variants remain to be discovered in

many populations.

African ancestry does not predict *P. falciparum* resistance in red cells

396   Based on evidence from balanced disease alleles like HbAS, it has been suggested

397 that anti-malarial selection has shaped polygenic red cell phenotypes in African

398 populations as a whole (Goheen *et al.*, 2016; Kanias *et al.*, 2017; Ma *et al.*, 2018). We tested

399 this hypothesis by examining the correlation between African ancestry and *P. falciparum*

400
Figure 6. No evidence of widespread selection in Africa for slower *P. falciparum* replication or
protective alleles in non-carriers. (a-d) Parasite growth and invasion measurements plotted against the
exome-wide fraction of African ancestry, determined by comparison to 1000 Genomes reference populations.
Adjusted-R^2 and *p*-values are shown for OLS regression. **(e)** Blue points represent non-carrier alleles selected
by Lasso that also have significant effects on parasite growth (*p*<0.1) in univariate linear models (Table S4).
The maximum estimated effect across two *P. falciparum* strains is shown. Red and purple points represent
HbAS and the HBA2 deletion in an additive model, respectively. F_{ST} was calculated between pan-African and
pan-European samples in gnomAD (Methods). Other disease alleles are not shown because they do not
function as single loci (G6PD⁻, HE) or do not affect growth (HbAC).

replication rate in RBCs from non-carriers (Figure 6A-D). We found no evidence that these

traits were related, apart from an weakly positive relationship between African ancestry

and the invasion rate of Th.026.09, a clinical Senegalese strain (p=0.01, R²=0.09, Figure

6D). To understand this result, we next examined how key RBC phenotypes identified in

this study (Figure 4A) vary with African ancestry. We found that greater African ancestry

predicts reduced osmotic fragility (p= 9.03 x 10⁻⁶), increased hypotonic deformability

(DI$_{min}$, p=0.001), lower cellular hemoglobin concentration (CHCM, reflecting reduced

dehydration, p=0.007), and a reduced fraction of RBCs with normal volume and

hemoglobin content (M5, p=0.033). All of these traits actually predict *greater* red cell

susceptibility to *P. falciparum* (Figure 4A), although together they explain less than 9% of

the non-carrier variation in parasite fitness. The remaining key phenotypes (Figure 4A) do

not vary with African ancestry, which may explain why African ancestry itself is mostly

unassociated with *P. falciparum* fitness in non-carrier RBCs (Figure 6A-D).

Next, we used allele frequency data from the global gnomAD collection (Karczewski

et al., 2020) to test whether polymorphisms associated with *P. falciparum* fitness in RBCs

(Table S4) vary between African and European populations. Geographical differences in

malaria selection are hypothesized to have increased the frequency of hundreds or

thousands of undiscovered anti-malarial alleles in Africa (Mackinnon *et al.*, 2005; Williams,

2006), as has been shown for several variants causing common RBC disorders

(Kwiatkowski, 2005). Here, aside from well-known disease variants, we find no evidence

that African populations as a whole are enriched for RBC polymorphisms that

impair *P. falciparum* growth in at least one strain (Figure 6E). Of 25 alleles with detectable

protective effects, 52% are in fact more common in Europeans than Africans (p=0.95,

binomial test). Besides HbAS, the polymorphism with the largest effect is a synonymous

variant in PIEZO1 (-31.9%, p=0.001), which is found at <4% frequency in both Africans and

Europeans. A few other protective polymorphisms, including two in *PIEZO1* and two in

SPTB, are nearly fixed in both populations. We do observe one F_{ST} outlier variant, a

synonymous allele at SPTA1 L958, which is substantially more common in Africa (70%

frequency) than in Europe (25%) and has a mild protective effect against *P. falciparum*

growth (-6.5%, p=0.027, Figure 6E). Given its high frequency, as well as its direct

association with a number of RBC phenotypes in previous GWAS (Table S4), this

synonymous SPTA1 allele could be a promising candidate for investigating anti-malarial

selection in Africa. However, the total set of natural RBC variants that limit *P. falciparum*

growth *in vitro* do not appear to support the hypothesis of widespread divergent selection

related to malaria between Africans and Europeans.

DISCUSSION

Healthy red blood cells (RBCs) harbor extensive phenotypic and genetic variation,

both within and between human populations. In this work, we demonstrate that this

variation modulates a wide range of RBC susceptibility to *P. falciparum* parasites. Our

findings add to a growing understanding of the genetic and phenotypic basis of RBC

resistance to *P. falciparum*, especially for RBCs that lack population-specific disease alleles.

In addition to suggesting new targets for future malaria interventions, these findings

challenge assumptions about the role of malaria selection in shaping human RBC diversity.

Since exponential replication of *P. falciparum* is a significant driver of malaria

disease progression (Bejon *et al.*, 2007), the ample variation that we observed in this trait

in vitro could be relevant for clinical outcomes in endemic regions. Growth inhibition from

HbAS, for example, reduces the risk of death from malaria by reducing parasite density in

the blood (Allison, 1954; Luzzatto, 2012). While HbAS has an extreme effect size, the 3-fold

range of parasite replication rates we found among non-carrier RBCs shares substantial

overlap with RBCs carrying other protective variants. Although the physiologically complex

basis of severe malaria (Okwa, 2012) makes it difficult to estimate the precise contribution

of RBC factors to severe malaria risk, the genotypes and phenotypes we have associated

with *P. falciparum* fitness may be promising targets for future therapeutic interventions.

We have shown here that widespread, 'normal' variation in RBC traits like volume,

hydration status, and maximum deformability is associated with *P. falciparum* fitness in

non-carrier RBCs. Reassuringly, these phenotypes are present to a stronger degree in RBCs

that do carry disease alleles (Clark, Mohandas and Shohet, 1983; Mockenhaupt *et al.*, 2000;

Pengon *et al.*, 2018). They are also consistent with published reports of extreme RBC traits

that limit parasite replication. For example, (Tiffert *et al.*, 2005) used an experimental

gradient of ion concentrations to show that *P. falciparum* growth is strongly reduced in

chemically dehydrated RBCs. Here, we found that naturally occurring, mildly dehydrated

RBCs also supported less efficient parasite growth and invasion. Similarly, the rare Dantu

variant of the glycophorin A/B receptors has been associated with increased RBC

membrane tension, decreased *P. falciparum* invasion efficiency, and decreased risk of

severe malaria (Field *et al.*, 1994; Leffler *et al.*, 2017; Kariuki *et al.*, 2018). Our data on RBC

membrane deformability and fragility support the same link between more flexible RBC

membranes and greater *P. falciparum* success. Moreover, they expand upon previous

studies by demonstrating for the first time that common, healthy phenotypic variation in

RBCs contributes meaningfully to *P. falciparum* fitness.

In our linear models of *P. falciparum* growth, phenotypic variation was strongly

outperformed by genetic variation in a small number of RBC proteins. This result implies

the existence of additional RBC phenotypes that we did not measure (or did not measure

with sufficient accuracy), but which have a genetic basis in a small number of genes

important to *P. falciparum*. Approximately half of the polymorphisms we identified are

non-synonymous, and may therefore exert direct effects on phenotypes like RBC

membrane structure or ion transport. The abundances of RBC gene transcripts and

proteins are other clear candidates for future phenotype studies, given that the other half

of associated polymorphisms were synonymous. A recent metanalysis showed that 'silent'

and 'coding' SNPs are equally likely to be associated with human disease, and moreover

have similar effect sizes (Chen *et al.*, 2010). Although synonymous SNPs may sometimes be

linked to coding variants, some have been shown to directly impact mRNA expression,

protein translation speed, and protein folding by altering transcription levels, codon usage,

and mRNA stability (Sauna and Kimchi-Sarfaty, 2011). Synonymous SNPs that impact

splicing, like rs35356320 in CD44, may also impact protein structure. Other conceivable

RBC phenotypes, such as the dynamics of membrane modification during *P. falciparum*

development, may only become evident in more detailed time course experiments. The

true number of RBC phenotypes important to *P. falciparum* may be effectively infinite

(Kinsler, Geiler-Samerotte and Petrov, 2020), making it very useful in practice that static

genetic variation is so strongly predictive of parasite growth.

Overall, just 31 polymorphisms in 14 RBC genes explained the majority of the

variation in *P. falciparum* growth among non-carriers. One reason that our explanatory

power is so high is our focus on *P. falciparum* replication in RBCs, which is a relatively

simple component of a larger, complex disease. This focus allowed us to use controlled, *in*

vitro experiments to detect genetic variants with small to moderate effects. Another

important explanation is our use of Lasso variable selection on just 106 polymorphisms in

genes with strong existing links to malaria. This approach obviated the need to meet an

exome-wide significance threshold, while still allowing for the discovery of novel,

putatively functional alleles in well-known disease genes. Furthermore, we confirmed that

this set of genes was enriched for signal compared to random sets of genes drawn from the

exome-wide background. It is important to note that genetic linkage complicates the

identification of the exact functional polymorphisms in any population sample (Sohail *et al.*,

2019), and we cannot rule out that some variants identified here are merely linked to the

true functional variants. Indeed, about half of the variants we identified occur in linkage

blocks containing other SNPs associated with RBC traits in GWAS studies. In any case, we

present significant evidence that loci containing 14 RBC genes are strongly enriched for

polymorphisms with significant impacts on *P. falciparum* growth.

One of the unique aspects of our study is the participation of individuals with a

range of African ancestry, defined by similarity to donors from five 1000 Genomes

reference populations. We found that African ancestry was associated with RBC

phenotypes that *improved* parasite fitness, particularly for the clinical Senegalese strain

Th.026.09. In the future, it would be very interesting to test for local parasite adaptation to

human RBCs using *P. falciparum* strains and RBC samples from around the globe. We also

found that the polymorphisms associated with *P. falciparum* growth by Lasso are not

enriched in African populations included in the gnomAD database of human variation. This

is not consistent with African-specific selection on these smaller-effect variants, despite the

famous examples provided by RBC disease alleles. This unexpected result for 'healthy'

variation has at least three possible explanations. First, relatively small-effect alleles may

not have had sufficient time to increase in frequency since *P. falciparum* began expanding

in humans some 5,000-10,000 years ago (Sundararaman *et al.*, 2016; Otto *et al.*, 2018).

Second, the complexity of severe malaria could mean that the variants discovered here do

not ultimately impact disease outcome, especially relative to known disease variants.

Third, human adaptation may be too local to detect with coarse-grain sampling of Sub-

Saharan African genetic diversity (e.g (Pankratov *et al.*, 2020)). Our data do suggest,

however, that few RBC alleles remain to be discovered that are both widespread in Africa

and have large effects on *P. falciparum* proliferation in RBCs (e.g. (Ma *et al.*, 2018)).

More broadly, these data show that it is inaccurate to make assumptions about RBC

susceptibility to *P. falciparum* based on a person's race or continental ancestry. These kinds

of hypotheses (Williams, 2006; Goheen *et al.*, 2016; Kanias *et al.*, 2017; Ma *et al.*, 2018) are

based on well-known examples of balanced disease alleles, but our data suggest that these

examples are exceptions rather than the rule. It is important to recognize that, at least in

biology classes, the use of racially-based genetic examples increases the belief that genes

encode absolute, functional differences between races (Donovan, 2017; Sparks, Baldwin

and Darner, 2020). This perspective neglects the fact that >90% of genetic differences

among humans occur within populations, rather than across them (Rosenberg, 2011). In

this study, global population structure captured by PCs showed that non-African variation

has its own distinct impacts on *P. falciparum* replication in RBCs. These data are an

important reminder that most human genetic diversity is a result of demography, not

population-specific selection (Coop *et al.*, 2009; Hofer *et al.*, 2009).

In conclusion, this study demonstrates that substantial phenotypic and genetic

diversity in healthy human RBCs impacts the replication of malaria parasites. Whether or

not this diversity is shaped by malaria selection, a better understanding of how *P.*

falciparum biology is impacted by natural RBC variation can help lead to new therapies for

one of humanity's most important infectious diseases.

ACKNOWLEDGMENTS

We gratefully acknowledge the invaluable participation of all volunteer blood donors. Nick

Bondy, Sandra Larkin, Brian Fleischer, Ashley Dunn, Talal Seddik, Trung Pham, David Vu,

and Spectrum Child Health provided crucial assistance in donor coordination and sample

processing. *P. falciparum* strain Th.026.09 was kindly provided by Daouda Ndiaye and

Sarah Volkman. For quantitative advice, we thank Grant Kinsler and the Stanford Statistics

Consulting Group. This work was primarily supported by a Pilot Early Career award from

the Stanford Maternal Child Health Research Institute and a Gabilan Faculty Award from

the Stanford University School of Medicine Office of Faculty Development and Diversity.

E.R.E. received additional support from the Stanford Center for Computational,

Evolutionary, and Human Genomics. D.A.P. was funded through an NIH MIRA award

5R35GM118165-05. E.S.E. was funded by the NIH (DP2HL13718601), and is a Tashia and

John Morgridge Endowed Faculty Scholar in Pediatric Translational Medicine through the

Stanford Maternal Child Health Research Institute. Local blood samples were drawn at the

Stanford Clinical and Translational Research Unit, which is supported by CTSA Grant UL1

TR001085.

AUTHOR CONTRIBUTIONS

Conceptualization, E.R.E., D.A.P., and E.S.E; Methodology, E.R.E., D.A.P., F.A.K., and E.S.E.;

Investigation, E.R.E., F.A.K., C.L., and E.S.E., Formal Analysis, E.R.E.; Visualization, E.R.E.,

Writing- Original Draft, E.R.E.; Writing- Review & Editing, E.R.E., F.A.K., D.A.P., and E.S.E.;

Funding Acquisition, E.R.E. and E.S.E.; Resources, F.A.K., D.A.P., and E.S.E; Supervision,

D.A.P. and E.S.E.

DECLARATION OF INTERESTS

The authors declare no competing interests.

METHODS

Sample collection and preparation

Subjects with no known history of RBC disorders were recruited to donate blood at

the Stanford Clinical and Translational Research Unit. Written informed consent was

obtained from each subject and/or their parent as part of a protocol approved by the

Stanford University Institutional Review Board (#40479). To help control for weekly batch

effects, Subject 1111 donated fresh blood for parasite assays each week. 11 other subjects

donated blood on at least two different weeks. Whole blood samples from a hereditary

elliptocytosis patient were obtained from Dr. Bertil Glader under a separate approved

protocol (#14004) that permitted sample sharing among researchers. All samples were de-

identified upon collection by labeling with a random four-digit number. Two samples were

eventually removed from analysis based on a failed sequencing library (6449KD) and

history of stem cell transplant (8715).

Whole blood was drawn into CPDA tubes and spun down within 36 hours to

separate serum, buffy coat, and red blood cells. Red blood cells were washed and stored in

RPMI-1640 medium (Sigma) supplemented with 25 mM HEPES, 50 mg/L hypoxanthine,

and 2.42 mM sodium bicarbonate at 4°C. Buffy coat was transferred directly to cryotubes

and stored at -80°C.

Exome sequencing and genotype calling

Genomic DNA was isolated from frozen buffy coats using a DNeasy Blood and Tissue

Kit (Qiagen). Libraries were prepared using a KAPA Hyperplus kit (Roche) and hybridized

to human exome probes using the SeqCap EZ Prime Exome kit (Roche). The resulting

exome libraries were sequenced with paired-end 150 bp Illumina reads on the HiSeq or

NextSeq platforms at Admera Health (South Plainfield, NJ).

Reads were aligned to the hg38 human reference genome using bwa mem(Li, 2013),

yielding an average coverage of 42X across targeted exome regions (excluding sample

6449KD). Variants were called using GATK best practices(Van der Auwera *et al.*, 2013) and

hard filtered with the following parameters: QD < 2.0, FS > 60.0, ReadPosRankSum < -2.5,

SOR > 2.5, MQ < 55.0, MQRankSum < -1.0, DP < 500. To minimize the effects of sequencing

errors, variants not present in 1000 Genomes, dbSNP_138, or the Mills indel

collection(Mills *et al.*, 2006) were discarded. Variants were also discarded if their

frequencies in our sample did not fall between gnomAD African or European

frequencies(Karczewski *et al.*, 2020) and significantly differed from both groups'

frequencies by Fisher's Exact Test (p≤0.05).

PIEZO1 E756del was genotyped via PCR and Sanger sequencing according to a

previously published protocol(Nguetse *et al.*, 2020). To call deletion variants that cause α-

thalassemia in the paralogous genes HBA2 and HBA1, we extracted reads from each .bam

file that lacked any mismatches or soft-clipping and had MAPQ ≥ 13 (i.e., <5% chance of

mapping error). Coverage with these well-mapped reads was calculated over the 73 and 81

bp of unique sequence in HBA2 and HBA1, respectively, and normalized to each sample's

exome-wide coverage. To determine which samples has unusually low coverage, we

formed an *ad hoc* reference panel of seven donors who were unlikely to carry deletion

alleles based on their normal MCH, MCV, and HGB and >96% exome-wide European

ancestry(Weatherall, 2001). We called heterozygous HBA2 deletions when normalized

coverage across three unique regions of the HBA2 gene was below the minimum reference

value. Similarly, we called homozygous HBA2 deletions when normalized coverage across

three unique regions of the HBA2 gene was less than half of the minimum reference value.

This approach resulted in an estimated HBA2 copy number of 2.0 in the reference panel,

0.95 in eight putative heterozygotes and 0.12 in four putative homozygotes. The same

method produced no evidence of HBA1 deletion in any sample.

Variant classification and linkage pruning

Exonic variants in RefSeq genes were identified using ANNOVAR(Wang, Li and
Hakonarson, 2010). Variants were classified into three categories: those within 23 malaria-
related genes (Table S2); those within 887 other red blood cell proteins (Table S5) derived
with a medium-confidence filter from the Red Blood Cell Collection database
(rbcc.hegelab.org); and those within any other gene. Singleton variants were removed to
avoid excessive false positives.

Linkage between all pairs of bi-allelic, exonic variants in our 121 genotyped samples
was calculated using the --geno-r2 and --interchrom-geno-r2 functions in vcftools (Danecek
et al., 2011). Variants in RBC genes that shared $r^2 > 0.1$ with any variant in the 23-gene set
were removed. Within the 23-gene set and RBC-gene set separately, non-carrier variants
were ranked by the *p*-values of their OLS regression with all four parasite measures. Then,
one variant was removed from each pair with $r^2 > 0.1$, prioritizing retention in the
following order: greater significance across models; non-synonymous protein change;
higher frequency in our sample; and finally by random sampling. We report results from
additive genetic models (genotypes coded 0/1/2), which performed as well or better than
recessive (0/0/2) and dominant (0/2/2) models.

Population analysis

The population ancestry of our donors was assessed by comparison with African,
European, East Asian, and South Asian reference populations from the 1000 Genomes
Project(The 1000 Genomes Project Consortium, 2015). Briefly, variants called from an
hg38 alignment of the 1000 Genomes data (Lowy-Gallego *et al.*, 2019) were filtered for
concordance with the variants genotyped in this study. The --indep-pairwise command in
PLINK(Purcell *et al.*, 2007) was used to prune SNPs with $r^2 > 0.1$ with any other SNP in a
50-SNP sliding window, producing 35,759 unlinked variants. These variants were analyzed
in both PLINK --pca and in ADMIXTURE(Alexander and Lange, 2011) with K=4 for the 121
genotyped individuals in this study, alongside 2,458 individuals from 1000 Genomes. Pan-
African and pan-European allele frequencies were obtained from gnomAD v3 (Karczewski
et al., 2020). F_{ST} for specific alleles was calculated as $(H_T-H_S)/H_T$ and then polarized, such
that positive values indicate variants more common in Africa.

P. falciparum culture and assays

P. falciparum strain 3D7 is a laboratory-adapted strain that was obtained from the Walter and Eliza Hall Institute (Melbourne, Australia) and routinely cultured in human erythrocytes obtained from the Stanford Blood Center. Th.026.09 is a clinical strain isolated from a patient in Senegal in 2009 that was then adapted to short-term culture (Park *et al.*, 2012) and kindly provided by Daouda Ndiaye and Sarah Volkman. 3D7 was maintained at 2% hematocrit in RPMI-1640 supplemented with 25 mM HEPES, 50 mg/L hypoxanthine, 2.42 mM sodium bicarbonate and 4.31 mg/ml Albumax (Invitrogen) at 37°C in 5% CO_2 and 1% O_2. Th.026.09 was maintained in the same media, with the exception that half the Albumax was replaced with heat-inactivated human AB serum.

Parasite growth and invasion assays were performed using schizont-stage parasites isolated from routine culture using a MACS magnet (Miltenyi). Parasites were added at ~0.5% initial parasitemia to fresh erythrocytes suspended at 1% hematocrit in complete RPMI, as above. Parasites were cultured in each erythrocyte sample for three to five days in triplicate 100 uL wells. Parasitemia was determined on day 0, day 1 (24 hours), day 3 (72 hours), and in some cases day 5 (120 hours) by staining with SYBR Green 1 nucleic acid stain (Invitrogen, ThermoFisher Scientific, Eugene, OR, USA) at 1:2000 dilution in PBS/0.3% BSA for 20 minutes, followed by flow cytometry analysis on a MACSQuant flow cytometer (Miltenyi). Raw invasion rate was defined as the day 1 parasitemia divided by the day 0 parasitemia; raw growth rate was defined as the day 3 (or day 5) parasitemia divided by the day 1 (or day 3) parasitemia. Day 0 parasitemia was not measured in weeks 1-3, so invasion rate estimates are absent for these samples. The parasite assays failed for both strains in week 9 and for Th.026.09 in week 10, and so were repeated in weeks 10 and 11 with RBCs that had been stored for 1 or 2 weeks.

To correct for batch effects, such as week-to-week variation in *P. falciparum* replication, we extracted the residuals from a linear regression of the raw parasite values against up to four significantly related batch variables: (1) the raw values for control donor 1111 each week; (2) the parasitemia measured at the previous time point; (3) the age in weeks of the RBCs being measured; and (4) the experimenter performing the assays. Notably, there was no additional effect of 'Week' or the length of the experiment (i.e., 3 or 5

days) once the previous variables were regressed out. To convert these residuals (mean 0)

to relative percents, we first transformed them with a linear model parameterized by data

from control donor 1111 and the most extreme donors (HbAS for growth; G6PD$^-_{high}$ and HE

for invasion), normalized by the 1111 rates from the appropriate week. These values were

finally arithmetically adjusted so that the mean invasion and growth values for non-

carriers was 100%.

Red cell phenotyping and normalization

Complete blood count (CBC) data for RBCs, reticulocytes, and platelets were

obtained with an Advia 120 hematology analyzer (Siemens, Laguna Hills, CA) at the Red

Cell Laboratory at Children's Hospital Oakland Research Institute. These data were: RBC,

HGB, HCT, MCV, MCH, MCHC, CHCM, RDW, HDW, PLT, MPV, Reticulocyte number and

percentage, and the fraction of RBCs in each of nine cells of the RBC matrix (M1-M9, with

MCH cutoffs of 28 and 41 and MCV cutoffs of 60 and 120). Systematic biases were evident

for some measures in certain weeks, but data from control donor 1111 were not available

for all weeks. Therefore, CBC data were normalized such that the median value for non-

carrier samples was equal across weeks.

Osmotic fragility tests were performed by incubating 20 μL of washed erythrocytes

for 5 minutes in 500 μL solutions of NaCl in 14 concentrations: 7.17, 6.14, 5.73, 5.32, 4.91,

4.50, 4.30, 4.09, 3.89, 3.68, 3.27, 3.07, 2.66, and 2.46 g/L. Tubes were spun for 5 minutes at

1000g and 100 μL of supernatant was transferred to a 96-well plate in duplicate.

Hemoglobin concentration was determined by adding 100 μL of Drabkin's reagent (Ricca

Chemical) to each well and measuring absorbance at OD_{540nm} with a Synergy H1 plate

reader (Biotek). Relative lysis was determined by normalizing to the maximum hemoglobin

concentration in the 14-tube series for each sample. After outlier points were manually

removed, sigmoidal osmotic fragility curves were estimated under a self-starting logistic

model in the nls package in R. Curves were summarized by the relative tonicity at which

50% lysis occurred and normalized within weekly batches, such that this value was equal

for control sample 1111 across weeks.

Osmotic gradient ektacytometry (Clark, Mohandas and Shohet, 1983; Kuypers *et al.*,

1990) was performed at the Red Cell Laboratory at Children's Hospital Oakland Research

Institute (CHORI). Red cell deformability estimates across a range of NaCl concentrations

were fitted to a 20-parameter polynomial model to generate a smooth curve, which was

manually verified to closely fit the data. Each curve was summarized with three standard

points (Figure S3A)(Clark, Mohandas and Shohet, 1983), which were normalized such that

the median x- and y-values of the three points was equal for non-carrier samples across

weeks.

Statistical analysis

Student's t-test was used to compare trait values between non-carriers and each

other group. Where n=1 (i.e., for G6PD$^-$$_{high}$ and HE), *p*-values were defined as the percentile

of the non-carrier distribution. Significance in these tests was defined as *p*<0.1. OLS linear

regression was performed with the lm function in R for all comparisons of two continuous

variables, unless otherwise specified. Adjusted R^2 values are reported.

Lasso regression (Tibshirani and Tibshirani, 1994; Chatterjee, 2013) was performed

with the cv.glmnet function in R. To account for moderate uncertainty in the choice of

lambda (due to random division of data for cross-validations), each model was run 1000

times. For each predictor, we report the fraction of runs in which it was selected (e.g Figure

4A; Table S1; Table S3) .The R^2 values reported between actual and predicted parasite

measures are medians from 1000 runs. We noted that effect size estimates for each

predictor varied considerably across runs; therefore, we used univariate OLS regression to

separately estimate the effect size of each predictor selected at least once in at least one

model.

The significance of each Lasso result was assessed by permutation. Parasite values

were randomly shuffled 1000 times, preserving correlations among potential predictors.

The median models for 6 cross-validations of each of the 1,000 permuted data sets were

compared to the median model from the real data. To compare the 23 malaria-related

genes to other sets of genes, a gene-specific R^2 for the target set was calculated using OLS

with only specific variants (no phenotypes or PCs) selected in at least 70% of cross-

validations. 1000 random sets of 23 genes were then generated from the RBC proteome

(Table S5) and analyzed identically. *P*-values were calculated as the percentile of this

permuted distribution in which the malaria-related gene set fell.

LEAD CONTACT: Further information and requests for resources and reagents should be

directed to and will be fulfilled by the Lead Contact, Elizabeth Egan (eegan@stanford.edu).

MATERIALS AVAILABILITY

This study did not generate any new unique reagents.

DATA and CODE AVAILABILITY

The human sequence datasets generated during this study will be available at dbGAP (in

progress). Phenotype data and code generated during this study is available at

https://github.com/emily-ebel/RBC.

REFERENCES

Alexander, D. H. and Lange, K. (2011) 'Enhancements to the ADMIXTURE algorithm for

individual ancestry estimation', *BMC Bioinformatics*. BioMed Central, 12(1), p. 246. doi:

10.1186/1471-2105-12-246.

Allison, A. C. (1954) 'Protection afforded by sickle-cell trait against subtertian malarial

infection', *British Medical Journal*. BMJ Publishing Group, 1(4857), pp. 290–294. doi:

10.1136/bmj.1.4857.290.

Archer, N. M. *et al.* (2018) 'Resistance to Plasmodium falciparum in sickle cell trait

erythrocytes is driven by oxygen-dependent growth inhibition', *Proceedings of the National*

Academy of Sciences of the United States of America. National Academy of Sciences, 115(28),

pp. 7350–7355. doi: 10.1073/pnas.1804388115.

Astle, W. J. *et al.* (2016) 'The Allelic Landscape of Human Blood Cell Trait Variation and

Links to Common Complex Disease', *Cell*. Cell Press, 167(5), pp. 1415-1429.e19. doi:

10.1016/j.cell.2016.10.042.

Van der Auwera, G. A. *et al.* (2013) 'From fastQ data to high-confidence variant calls: The

genome analysis toolkit best practices pipeline', *Current Protocols in Bioinformatics*. John

Wiley and Sons Inc., 11(SUPL.43), p. 11.10.1. doi: 10.1002/0471250953.bi1110s43.

Band, G. *et al.* (2015) 'A novel locus of resistance to severe malaria in a region of ancient

balancing selection', *Nature.* Nature Publishing Group, 526(7572), pp. 253–257. doi:

10.1038/nature15390.

Bejon, P. *et al.* (2007) 'Defining Childhood Severe Falciparum Malaria for Intervention

Studies', *PLoS Medicine.* Edited by G. Pasvol. Public Library of Science, 4(8), p. e251. doi:

10.1371/journal.pmed.0040251.

Beutler, E. and West, C. (2005) 'Hematologic differences between African-Americans and

whites: The roles of iron deficiency and α-thalassemia on hemoglobin levels and mean

corpuscular volume', *Blood.* American Society of Hematology, 106(2), pp. 740–745. doi:

10.1182/blood-2005-02-0713.

Boyle, E. A., Li, Y. I. and Pritchard, J. K. (2017) 'An Expanded View of Complex Traits: From

Polygenic to Omnigenic', *Cell.* Cell Press, pp. 1177–1186. doi: 10.1016/j.cell.2017.05.038.

Campbell, M. C. and Tishkoff, S. A. (2008) 'African Genetic Diversity: Implications for

Human Demographic History, Modern Human Origins, and Complex Disease Mapping',

Annual Review of Genomics and Human Genetics. Annual Reviews , 9(1), pp. 403–433. doi:

10.1146/annurev.genom.9.081307.164258.

Canela-Xandri, O., Rawlik, K. and Tenesa, A. (2018) 'An atlas of genetic associations in UK

Biobank', *Nature Genetics.* Nature Publishing Group, 50(11), pp. 1593–1599. doi:

10.1038/s41588-018-0248-z.

Cao, A. and Galanello, R. (2010) 'Beta-thalassemia', *Genetics in Medicine.* Nature Publishing

Group, pp. 61–76. doi: 10.1097/GIM.0b013e3181cd68ed.

Chami, N. *et al.* (2016) 'Exome Genotyping Identifies Pleiotropic Variants Associated with

Red Blood Cell Traits', *American Journal of Human Genetics.* Cell Press, 99(1), pp. 8–21. doi:

10.1016/j.ajhg.2016.05.007.

Chatterjee, S. (2013) 'Assumptionless consistency of the Lasso', *arXiv.* Available at:

http://arxiv.org/abs/1303.5817 (Accessed: 10 March 2020).

Chen, M.-H. *et al.* (2020) 'Trans-ethnic and Ancestry-Specific Blood-Cell Genetics in 746,667

Individuals from 5 Global Populations', *Cell.* Cell Press, 182(5), pp. 1198-1213.e14. doi:

10.1016/j.cell.2020.06.045.

Chen, R. *et al.* (2010) 'Non-synonymous and synonymous coding SNPS show similar

likelihood and effect size of human disease association', *PLoS ONE*. PLoS One, 5(10). doi:

10.1371/journal.pone.0013574.

Clark, M., Mohandas, N. and Shohet, S. (1983) 'Osmotic gradient ektacytometry:

comprehensive characterization of red cell volume and surface maintenance', *Blood*, 61(5),

pp. 899–910. doi: 10.1182/blood.V61.5.899.899.

Clarke, G. M. *et al.* (2017) 'Characterisation of the opposing effects of G6PD deficiency on

cerebral malaria and severe malarial anaemia', *eLife*. eLife Sciences Publications Ltd, 6. doi:

10.7554/eLife.15085.

Cooling, L. (2015) 'Blood groups in infection and host susceptibility', *Clinical Microbiology*

Reviews. American Society for Microbiology, 28(3), pp. 801–870. doi: 10.1128/CMR.00109-

14.

Coop, G. *et al.* (2009) 'The Role of Geography in Human Adaptation', *PLoS Genetics*. Edited

by M. H. Schierup. Public Library of Science, 5(6), p. e1000500. doi:

10.1371/journal.pgen.1000500.

Crosnier, C. *et al.* (2011) 'Basigin is a receptor essential for erythrocyte invasion by

Plasmodium falciparum', *Nature*. NIH Public Access, 480(7378), pp. 534–537. doi:

10.1038/nature10606.

Danecek, P. *et al.* (2011) 'The variant call format and VCFtools', *Bioinformatics*, 27(15), pp.

2156–2158. doi: 10.1093/bioinformatics/btr330.

Dhermy, D., Schrével, J. and Lecomte, M.-C. (2007) 'Spectrin-based skeleton in red blood

cells and malaria', *Current Opinion in Hematology*, 14(3), pp. 198–202. doi:

10.1097/MOH.0b013e3280d21afd.

Donovan, B. M. (2017) 'Learned inequality: Racial labels in the biology curriculum can

affect the development of racial prejudice', *Journal of Research in Science Teaching*. John

Wiley and Sons Inc., 54(3), pp. 379–411. doi: 10.1002/tea.21370.

Egan, E. S. *et al.* (2015) 'A forward genetic screen identifies erythrocyte CD55 as essential

for Plasmodium falciparum invasion', *Science*. American Association for the Advancement

of Science, 348(6235), pp. 711–714. doi: 10.1126/science.aaa3526.

Egan, E. S. *et al.* (2018) 'Erythrocytes lacking the Langereis blood group protein ABCB6 are

resistant to the malaria parasite Plasmodium falciparum', *Communications Biology*. Nature

Research, 1(1), pp. 1–9. doi: 10.1038/s42003-018-0046-2.

Evans, D. M., Frazer, I. H. and Martin, N. G. (1999) 'Genetic and environmental causes of

variation in basal levels of blood cells', *Twin Research (1999)*. Twin Res, 2(4), pp. 250–257.

doi: 10.1375/136905299320565735.

Facer, C. A. (1995) *Erythrocytes carrying mutations in spectrin and protein 4.1 show differing*

sensitivities to invasion by Plasmodium falciparum, *Parasitol Res*. Springer Verlag.

Field, S. P. *et al.* (1994) 'Glycophorin variants and Plasmodium falciparum: protective effect

of the Dantu phenotype in vitro', *Human Genetics*. Springer-Verlag, 93(2), pp. 148–150. doi:

10.1007/BF00210600.

Flynn, C. J., Hurvich, C. M. and Simonoff, J. S. (2017) 'On the sensitivity of the Lasso to the

number of predictor variables', *Statistical Science*. Institute of Mathematical Statistics,

32(1), pp. 88–105. doi: 10.1214/16-STS586.

Francis, S. E., Sullivan, D. J. and Goldberg, and D. E. (1997) 'HEMOGLOBIN METABOLISM IN

THE MALARIA PARASITE *PLASMODIUM FALCIPARUM*', *Annual Review of Microbiology*.

Annual Reviews 4139 El Camino Way, P.O. Box 10139, Palo Alto, CA 94303-0139, USA ,

51(1), pp. 97–123. doi: 10.1146/annurev.micro.51.1.97.

Friedman, M. J. (1978) 'Erythrocytic mechanism of sickle cell resistance to malaria',

Proceedings of the National Academy of Sciences of the United States of America. National

Academy of Sciences, 75(4), pp. 1994–1997. doi: 10.1073/pnas.75.4.1994.

Galanello, R. and Cao, A. (2011) 'Alpha-thalassemia', *Genetics in Medicine*. Genet Med, pp.

83–88. doi: 10.1097/GIM.0b013e3181fcb468.

Gallagher, P. G. (2013) 'Abnormalities of the erythrocyte membrane', *Pediatric Clinics of*

North America. NIH Public Access, pp. 1349–1362. doi: 10.1016/j.pcl.2013.09.001.

Garn, S. M. (1981) 'Lower Hematocrit Levels in Blacks Are Not Due to Diet or

Socioeconomic Factors', *Pediatrics*, 67(4).

Glogowska, E. *et al.* (2017) 'Novel mechanisms of PIEZO1 dysfunction in hereditary

xerocytosis', *Blood*. American Society of Hematology, 130(16), pp. 1845–1856. doi:

10.1182/blood-2017-05-786004.

Goheen, M. M. *et al.* (2016) 'Anemia Offers Stronger Protection Than Sickle Cell Trait

Against the Erythrocytic Stage of Falciparum Malaria and This Protection Is Reversed by

Iron Supplementation', *EBioMedicine*. Elsevier B.V., 14, pp. 123–130. doi:

10.1016/j.ebiom.2016.11.011.

Greene, L. S. (1993) 'G6PD deficiency as protection againstfalciparum malaria: An

epidemiologic critique of population and experimental studies', *American Journal of*

Physical Anthropology, 36(S17), pp. 153–178. doi: 10.1002/ajpa.1330360609.

GTEx Consortium (2017) 'Genetic effects on gene expression across human tissues', *Nature*.

Nature Publishing Group, 550(7675), pp. 204–213. doi: 10.1038/nature24277.

Hanssen, E. *et al.* (2012) 'Soft X-ray microscopy analysis of cell volume and hemoglobin

content in erythrocytes infected with asexual and sexual stages of Plasmodium falciparum',

Journal of Structural Biology. Academic Press, 177(2), pp. 224–232. doi:

10.1016/j.jsb.2011.09.003.

van der Harst, P. *et al.* (2012) 'Seventy-five genetic loci influencing the human red blood

cell.', *Nature*. NIH Public Access, 492(7429), pp. 369–75. doi: 10.1038/nature11677.

Hofer, T. *et al.* (2009) 'Large allele frequency differences between human continental

groups are more likely to have occurred by drift during range expansions than by

selection', *Annals of Human Genetics*. Blackwell Publishing Ltd, 73(1), pp. 95–108. doi:

10.1111/j.1469-1809.2008.00489.x.

Ifediba, T. *et al.* (1985) 'Plasmodium falciparum in vitro: diminished growth in hemoglobin

H disease erythrocytes', *Blood*. American Society of Hematology, 65(2), pp. 452–455. doi:

10.1182/blood.V65.2.452.452.

Ilboudo, Y. *et al.* (2018) 'A common functional PIEZO1 deletion allele associates with red

blood cell density in sickle cell disease patients', *American Journal of Hematology*. Wiley-

Liss Inc., pp. E362–E365. doi: 10.1002/ajh.25245.

Ioannidis, J. P. A. (2005) 'Why Most Published Research Findings Are False', *PLoS Medicine*.

Springer International Publishing, 2(8), p. e124. doi: 10.1371/journal.pmed.0020124.

Kanias, T. *et al.* (2017) 'Ethnicity, sex, and age are determinants of red blood cell storage

and stress hemolysis: Results of the REDS-III RBC-Omics study', *Blood Advances*. American

Society of Hematology, 1(15), pp. 1132–1141. doi: 10.1182/bloodadvances.2017004820.

Karczewski, K. J. *et al.* (2020) 'The mutational constraint spectrum quantified from

variation in 141,456 humans', *Nature*. Nature Research, 581(7809), pp. 434–443. doi:

10.1038/s41586-020-2308-7.

Kariuki, S. N. *et al.* (2018) 'Red blood cell tension controls Plasmodium falciparum invasion

and protects against severe malaria in the Dantu blood group', *bioRxiv*. Cold Spring Harbor

Laboratory, (November), p. 475442. doi: 10.1101/475442.

Kariuki, S. N. *et al.* (2020) 'Red blood cell tension protects against severe malaria in the

Dantu blood group', *Nature*. Nature Publishing Group, pp. 1–5. doi: 10.1038/s41586-020-

2726-6.

Kinsler, G., Geiler-Samerotte, K. and Petrov, D. (2020) 'A genotype-phenotype-fitness map

reveals local modularity and global pleiotropy of adaptation', *bioRxiv*. Cold Spring Harbor

Laboratory, p. 2020.06.25.172197. doi: 10.1101/2020.06.25.172197.

Koch, M. *et al.* (2017) 'Plasmodium falciparum erythrocyte-binding antigen 175 triggers a

biophysical change in the red blood cell that facilitates invasion', *Proceedings of the*

National Academy of Sciences of the United States of America. National Academy of Sciences,

114(16), pp. 4225–4230. doi: 10.1073/pnas.1620843114.

Kuypers, F. A. *et al.* (1990) 'Use of ektacytometry to determine red cell susceptibility to

oxidative stress', *The Journal of Laboratory and Clinical Medicine*, 116(4), pp. 527–534.

Kwiatkowski, D. P. (2005) 'How malaria has affected the human genome and what human

genetics can teach us about malaria', *American Journal of Human Genetics*. University of

Chicago Press, pp. 171–192. doi: 10.1086/432519.

Leffler, E. M. *et al.* (2017) 'Resistance to malaria through structural variation of red blood

cell invasion receptors', *Science*. American Association for the Advancement of Science,

356(6343), pp. 1140–1152. doi: 10.1126/science.aam6393.

Lessard, S. *et al.* (2017) 'An erythroid-specific ATP2B4 enhancer mediates red blood cell

hydration and malaria susceptibility', *Journal of Clinical Investigation*. American Society for

Clinical Investigation, 127(8), pp. 3065–3074. doi: 10.1172/JCI94378.

Li, H. (2013) 'Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM', *arXiv*. Available at: http://arxiv.org/abs/1303.3997 (Accessed: 22 February 2020).

Li, J. *et al.* (2013) 'GWAS of blood cell traits identifies novel associated loci and epistatic

interactions in Caucasian and African-American children', *Human Molecular Genetics*. doi:

10.1093/hmg/dds534.

Lin, S.-H., Brown, D. and Machiela, M. (2020) 'LDtrait: an online tool for identifying

published phenotype associations in linkage disequilibrium', *Cancer Research*. American

Association for Cancer Research, p. canres.0985.2020. doi: 10.1158/0008-5472.CAN-20-

0985.

Lo, K. S. *et al.* (2011) 'Genetic association analysis highlights new loci that modulate

hematological trait variation in Caucasians and African Americans', *Human Genetics*.

Springer, 129(3), pp. 307–317. doi: 10.1007/s00439-010-0925-1.

Lowy-Gallego, E. *et al.* (2019) 'Variant calling on the GRCh38 assembly with the data from

phase three of the 1000 Genomes Project', *Wellcome Open Research*. F1000 Research Ltd, 4,

p. 50. doi: 10.12688/wellcomeopenres.15126.2.

Luzzatto, L. (2012) 'Sickle cell anaemia and malaria', *Mediterranean Journal of Hematology*

and Infectious Diseases. Catholic University in Rome, p. 2012065. doi:

10.4084/MJHID.2012.065.

Ma, S. *et al.* (2018) 'Common PIEZO1 Allele in African Populations Causes RBC Dehydration

and Attenuates Plasmodium Infection', *Cell*. Cell Press, 173(2), pp. 443-455.e12. doi:

10.1016/j.cell.2018.02.047.

Mackinnon, M. J. *et al.* (2005) 'Heritability of malaria in Africa', *PLoS Medicine*. Public

Library of Science, 2(12), pp. 1253–1259. doi: 10.1371/journal.pmed.0020340.

De Mendonça, V. R. R., Goncalves, M. S. and Barral-Netto, M. (2012) 'The Host Genetic

Diversity in Malaria Infection', *Journal of Tropical Medicine*. Hindawi Publishing

Corporation, 2012, p. 17. doi: 10.1155/2012/940616.

MGE Network (2019a) 'Insights into malaria susceptibility using genome-wide data on

17,000 individuals from Africa, Asia and Oceania', *Nature Communications*. NLM (Medline),

10(1), p. 5732. doi: 10.1038/s41467-019-13480-z.

MGE Network (2019b) 'Insights into malaria susceptibility using genome-wide data on

17,000 individuals from Africa, Asia and Oceania', *Nature communications*. NLM (Medline),

10(1), p. 5732. doi: 10.1038/s41467-019-13480-z.

Mills, R. E. *et al.* (2006) 'An initial map of insertion and deletion (INDEL) variation in the

human genome', *Genome Research*. Cold Spring Harbor Laboratory Press, 16(9), pp. 1182–

1190. doi: 10.1101/gr.4565806.

Mockenhaupt, F. P. *et al.* (2000) 'Anaemia in pregnant Ghanaian women: importance of

malaria, iron deficiency, and haemoglobinopathies', *Transactions of the Royal Society of*

Tropical Medicine and Hygiene. Royal Society of Tropical Medicine and Hygiene, 94(5), pp.

477–483. doi: 10.1016/S0035-9203(00)90057-9.

Ndila, C. M. *et al.* (2018) 'Human candidate gene polymorphisms and risk of severe malaria

in children in Kilifi, Kenya: a case-control association study', *The Lancet Haematology*.

Elsevier Ltd, 5(8), pp. e333–e345. doi: 10.1016/S2352-3026(18)30107-8.

Nguetse, C. N. *et al.* (2020) 'A common polymorphism in the mechanosensitive ion channel

PIEZO1 is associated with protection from severe malaria in humans', *Proceedings of the*

National Academy of Sciences. doi: 10.1073/pnas.1919843117/-/DCSupplemental.

Okwa, O. O. (2012) *Malaria Parasites*, *Malaria Parasites*. InTech. doi: 10.5772/1477.

Otto, T. D. *et al.* (2018) 'Genomes of all known members of a Plasmodium subgenus reveal

paths to virulent human malaria', *Nature Microbiology*. Nature Publishing Group, 3(6), pp.

687–697. doi: 10.1038/s41564-018-0162-2.

Page, G. *et al.* (2017) 'GWAS of Osmotic Hemolysis in 12,352 Healthy Blood Donors

Identifies Red Cell Genetic Variants Associated with Steady State Hemolysis in Patients

with Sickle Cell Disease', *Blood*. American Society of Hematology, 130(Supplement 1), pp.

1117–1117. doi: 10.1182/BLOOD.V130.SUPPL_1.1117.1117.

Pankratov, V. *et al.* (2020) 'Differences in local population history at the finest level: the

case of the Estonian population', *European Journal of Human Genetics*. Springer Science and

Business Media LLC, pp. 1–12. doi: 10.1038/s41431-020-0699-4.

Park, D. J. *et al.* (2012) 'Sequence-based association and selection scans identify drug

resistance loci in the Plasmodium falciparum malaria parasite', *Proceedings of the National*

Academy of Sciences of the United States of America. National Academy of Sciences, 109(32),

pp. 13052–13057. doi: 10.1073/pnas.1210585109.

Pasvol, G., Weatherall, D. J. and Wilson, R. J. M. (1978) 'Cellular mechanism for the

protective effect of haemoglobin S against P. falciparum malaria [22]', *Nature*. Nature, pp.

701–703. doi: 10.1038/274701a0.

Pengon, J. *et al.* (2018) 'Hematological parameters and red blood cell morphological

abnormality of Glucose-6-Phosphate dehydrogenase deficiency co-inherited with

thalassemia', *Hematology/ Oncology and Stem Cell Therapy*. King Faisal Specialist Hospital

and Research Centre, 11(1), pp. 18–24. doi: 10.1016/j.hemonc.2017.05.029.

Perry, G. S. *et al.* (1992) 'Iron nutrition does not account for the hemoglobin differences

between blacks and whites', *Journal of Nutrition*, 122(7), pp. 1417–1424. doi:

10.1093/jn/122.7.1417.

Pilia, G. *et al.* (2006) 'Heritability of Cardiovascular and Personality Traits in 6,148

Sardinians', *PLoS Genetics*. Edited by W. N. Frankel. Public Library of Science, 2(8), p. e132.

doi: 10.1371/journal.pgen.0020132.

Purcell, S. *et al.* (2007) 'PLINK: A tool set for whole-genome association and population-

based linkage analyses', *American Journal of Human Genetics*. Cell Press, 81(3), pp. 559–

575. doi: 10.1086/519795.

Rockett, K. A. *et al.* (2014) 'Reappraisal of known malaria resistance loci in a large

multicenter study', *Nature Genetics*. Nature Publishing Group, 46(11), pp. 1197–1204. doi:

10.1038/ng.3107.

Rooks, H. *et al.* (2019) 'A gain of function variant in PIEZO1 (E756del) and sickle cell

disease', *Haematologica*. Ferrata Storti Foundation, pp. e91–e93. doi:

10.3324/haematol.2018.202697.

Rosenberg, N. A. (2011) 'A Population-Genetic Perspective on the Similarities and

Differences among Worldwide Human Populations', *Human Biology*. Wayne State

University Press, 83(6), pp. 659–684. doi: 10.3378/027.083.0601.

Ruwende, C. and Hill, A. (1998) 'Glucose-6-phosphate dehydrogenase deficiency and

malaria', *Journal of Molecular Medicine*. J Mol Med (Berl), pp. 581–588. doi:

10.1007/s001090050253.

Sauna, Z. E. and Kimchi-Sarfaty, C. (2011) 'Understanding the contribution of synonymous

mutations to human disease', *Nature Reviews Genetics*. Nature Publishing Group, pp. 683–

691. doi: 10.1038/nrg3051.

Schulman, S. *et al.* (1990) 'Growth of Plasmodium falciparum in human erythrocytes

containing abnormal membrane proteins', *Proceedings of the National Academy of Sciences*

of the United States of America. National Academy of Sciences, 87(18), pp. 7339–7343. doi:

10.1073/pnas.87.18.7339.

Sohail, M. *et al.* (2019) 'Polygenic adaptation on height is overestimated due to uncorrected

stratification in genome-wide association studies', *eLife*. eLife Sciences Publications Ltd, 8.

doi: 10.7554/eLife.39702.

Sparks, R. A., Baldwin, K. E. and Darner, R. (2020) 'Using Culturally Relevant Pedagogy to

Reconsider the Genetics Canon †', *Journal of Microbiology & Biology Education*. American

Society for Microbiology, 21(1). doi: 10.1128/jmbe.v21i1.1901.

Sundararaman, S. A. *et al.* (2016) 'Genomes of cryptic chimpanzee Plasmodium species

reveal key evolutionary events leading to human malaria', *Nature Communications*. Nature

Publishing Group, 7. doi: 10.1038/ncomms11078.

The 1000 Genomes Project Consortium (2015) 'A global reference for human genetic

variation', *Nature*. Nature Publishing Group, pp. 68–74. doi: 10.1038/nature15393.

Tibshirani, R. and Tibshirani, R. (1994) 'Regression Shrinkage and Selection Via the Lasso',

JOURNAL OF THE ROYAL STATISTICAL SOCIETY, SERIES B, 58, pp. 267--288. Available at:

http://citeseer.ist.psu.edu/viewdoc/summary?doi=10.1.1.35.7574 (Accessed: 22 February

2020).

Tiffert, T. *et al.* (2005) 'The hydration state of human red blood cells and their susceptibility

to invasion by Plasmodium falciparum', *Blood*. The American Society of Hematology,

105(12), pp. 4853–4860. doi: 10.1182/blood-2004-12-4948.

Timmann, C. *et al.* (2012) 'Genome-wide association study indicates two novel resistance

loci for severe malaria', *Nature*, 489(7416), pp. 443–446. doi: 10.1038/nature11334.

Vuckovic, D. *et al.* (2020) 'The Polygenic and Monogenic Basis of Blood Traits and Diseases',

Cell. Cell Press, 182(5), pp. 1214-1231.e11. doi: 10.1016/j.cell.2020.08.008.

Wang, K., Li, M. and Hakonarson, H. (2010) 'ANNOVAR: Functional annotation of genetic

variants from high-throughput sequencing data', *Nucleic Acids Research*, 38(16). doi:

10.1093/nar/gkq603.

Weatherall, D. J. (2001) 'Phenotype-genotype relationships in monogenic disease: Lessons

from the thalassaemias', *Nature Reviews Genetics*. Nature Publishing Group, pp. 245–255.

doi: 10.1038/35066048.

Whitfield, J. B., Martin, N. G. and Rao, D. C. (1985) 'Genetic and environmental influences on

the size and number of cells in the blood', *Genetic Epidemiology*, 2(2), pp. 133–144. doi:

10.1002/gepi.1370020204.

WHO (2019) *Malaria eradication: benefits, future scenarios and feasibility. Executive*

summary, WHO Strategic Advisory Group on Malaria Eradication. Available at:

https://www.who.int/publications/i/item/who-cds-gmp-2019-10 (Accessed: 7 August

2020).

Williams, T. N. (2006) 'Human red blood cell polymorphisms and malaria', *Current Opinion*

in Microbiology. Elsevier Current Trends, pp. 388–394. doi: 10.1016/j.mib.2006.06.009.

Wright, G. J. and Rayner, J. C. (2014) 'Plasmodium falciparum Erythrocyte Invasion:

Combining Function with Immune Evasion', *PLoS Pathogens*. Public Library of Science,

10(3). doi: 10.1371/journal.ppat.1003943.

Yoshida, A., Beutler, E. and Motulsky, A. G. (1971) 'Human glucose-6-phosphate

dehydrogenase variants.', *Bulletin of the World Health Organization*. World Health

Organization, 45(2), pp. 243–253.

Zámbó, B. *et al.* (2017) 'Decreased calcium pump expression in human erythrocytes is

connected to a minor haplotype in the ATP2B4 gene', *Cell Calcium*. Elsevier Ltd, 65, pp. 73–

79. doi: 10.1016/j.ceca.2017.02.001.

Zhang, Y. *et al.* (2015) 'Multiple stiffening effects of nanoscale knobs on human red blood

cells infected with Plasmodium falciparum malaria parasite', *Proceedings of the National*

Academy of Sciences of the United States of America. National Academy of Sciences, 112(19),

pp. 6068–6073. doi: 10.1073/pnas.1505584112.

1081