1   **Common physiological processes control mercury reduction during**

2   **photosynthesis and fermentation**

4   Running title: Isotopes unify bacterial controls on Hg reduction
5

Daniel S. Grégoire [a*#], Sarah E. Janssen [b], Noémie C. Lavoie [a], Michael T. Tate [b], & Alexandre J.

Poulain [a]

a - Biology Department, University of Ottawa, 30 Marie Curie, Ottawa, ON, K1N 6N5, Canada.

b - Upper Midwest Water Science Center, USGS, Middleton, Wisconsin 53562, USA.

Corresponding author: Dr. Daniel S. Grégoire, daniel.gregoire@uwaterloo.ca

* Current affiliation: Biology Department, University of Waterloo, 200 University Ave W,

Waterloo, ON, N2L 3G1

Word count for abstract: 248

Word count for text (excluding abstract and importance sections): 4968

ABSTRACT

Mercury (Hg) is a global pollutant and potent neurotoxin that bioaccumulates in food webs as monomethylmercury (MeHg). The production of MeHg is driven by anaerobic and Hg redox cycling pathways such as Hg reduction, which control the availability of Hg to methylators. Anaerobes play an important role in Hg reduction in methylation hotspots, yet their contributions remain underappreciated due to how challenging these pathways are to study in the absence of dedicated genetic targets and low levels of Hg^0 in anoxic environments. In this study we used Hg stable isotope fractionation to explore Hg reduction during anoxygenic photosynthesis and fermentation in the model anaerobe *Heliobacterium modesticaldum* Ice1. We show that cells preferentially reduce lighter Hg isotopes in both metabolisms leading to mass-dependent fractionation, but mass-independent fractionation commonly induced by UV-visible light is absent. We show that isotope fractionation is affected by the interplay between pathways controlling Hg recruitment, accessibility, and availability alongside metabolic redox reactions. The combined contributions of these processes lead to isotopic enrichment during anoxygenic photosynthesis that is in between the values reported for anaerobic respiratory microbial Hg reduction and abiotic photoreduction. Isotope enrichment during fermentation is closer to what has been observed in aerobic bacteria that reduce Hg through dedicated detoxification pathways. These results demonstrate that common controls exist at the atomic level for Hg reduction during photosynthesis and fermentation in *H. modesticaldum*. Our work suggests that similar controls likely underpin diverse microbe-mediated Hg transformations that affect Hg's fate in oxic and anoxic habitats.

IMPORTANCE

Anaerobic and photosynthetic bacteria that reduce mercury affect mercury delivery to microbes in methylation sites that drive bioaccumulation in food webs. Anaerobic mercury reduction pathways remain underappreciated in the current view of the global mercury cycle because they are challenging to study, bearing no dedicated genetic targets to establish physiological mechanisms. In this study we used stable isotopes to show that common physiological processes control mercury reduction during photosynthesis and fermentation in the model anaerobe *Heliobacterium modesticaldum* Ice1. The sensitivity of isotope analyses highlighted the subtle contribution of mercury uptake towards the isotope signature associated with anaerobic mercury reduction. When considered alongside the isotope signatures associated with microbial pathways for which genetic determinants have been identified, our findings underscore the narrow range of isotope enrichment that is characteristic of microbial mercury transformations. This suggests that there exist common atomic-level controls for biological mercury transformations across a broad range of geochemical conditions.

INTRODUCTION

Mercury (Hg) is a global pollutant and potent neurotoxin that bioaccumulates in aquatic and terrestrial food webs as monomethylmercury (MeHg) (1). Anaerobic microbes with the *hgcAB* gene cluster, which encodes for the metabolic machinery responsible for Hg methylation (2), contribute to MeHg production in habitats such as aquatic sediments, water columns, and flooded soils (3-6). Hg methylation is thought to occur intracellularly and as such, it is ultimately controlled by the bioavailability of Hg to methylators in anoxic habitats (7).

Hg redox cycling plays a key role in determining the inorganic Hg substrates available to methylators. Its role is two-fold: anaerobic Hg^0 oxidation can supply dissolved Hg^{II} required to generate MeHg and evasion of Hg^0 due to reduction can limit MeHg production by removing inorganic Hg substrates. Although it is well established that abiotic photochemical Hg reduction dominates Hg redox cycling in oxic surface systems where light is present (8-16), Hg reduction pathways in anoxic zones where Hg methylation occurs have mostly been characterized under laboratory conditions and remain challenging to assess in the field.

Anoxic Hg reduction can occur via abiotic redox reactions with dissolved organic carbon (17) and magnetite (18, 19), but also through biotic pathways mediated by chemotrophic and phototrophic microorganisms (20-24). Unlike many aerobes that reduce Hg using dedicated enzymatic machinery encoded by the *mer* operon (25, 26), Hg reduction in anaerobes occurs through cometabolic pathways tied to anaerobic respiration, fermentation, and anoxygenic photosynthesis (20-24). In contrast to *mer*-mediated Hg reduction and demethylation, dedicated genetic determinants have yet to be identified for anaerobic Hg reduction, making these pathways challenging to study from a mechanistic standpoint. Furthermore, while some studies have shown that Hg^0 can significantly contribute to Hg speciation in anoxic habitats (27, 28), it is

possible that low levels of Hg^0 are maintained by an active cycle of reduction and oxidation (29).

In the absence of genetic targets and measurable Hg^0 accumulation, the contributions of

anaerobic Hg^0 production to global Hg cycling remain cryptic and difficult to assess.

The use of stable Hg isotope fractionation is a powerful tool for providing

biogeochemical proxies for different abiotic and biotic Hg transformations (30, 31). All abiotic

and biotic Hg transformation pathways studied to date demonstrate kinetic mass dependent

fractionation (MDF) in which the product pool becomes isotopically lighter (enrichment of light

isotopes) as the reaction proceeds (32). Some pathways, such as abiotic photochemical reduction

and demethylation, also demonstrate mass independent fractionation (MIF), which is defined as

the fractionation that occurs outside of the predicted MDF pattern (30). MIF is most commonly

observed due to the magnetic isotope effect for odd-Hg isotopes ^{199}Hg and ^{201}Hg in the presence

of sunlight (33) and the extent to which MIF occurs is controlled in part by water chemistry and

the presence of photosynthetic organisms (39).

The combination of MDF and MIF tracers, and the rate of isotopic enrichment, referred

to as fractionation factor, can be used to outline the underlying components of different Hg

microbe-mediated Hg transformation pathways. Over the last decade, Hg isotope fractionation

has been studied in microbes capable of *mer*-mediated reduction (34) and MeHg demethylation

(35); anaerobic non-*mer*-mediated Hg reduction (36, 37); Hg methylation (37-40); and oxygenic

phototrophic MeHg degradation and Hg reduction (41). These studies have yielded valuable

insights and mechanistic details that explain how microbes interact with Hg at the atomic level

making them a promising option for studying mechanisms for anaerobic Hg reduction.

Our objective in this study was to characterize Hg isotope fractionation patterns during

anoxygenic phototrophic and fermentative Hg^{II} reduction, for which there exists no data. By

providing the first isotopic characterization of these pathways we aim to better understand the

underlying physiological pathways involved in anaerobic Hg reduction and their role in

controlling Hg availability to methylators in anoxic habitats. In this work, we test whether Hg

reduction by the model anaerobe *Heliobacterium modesticaldum* Ice1 during anoxygenic

photosynthesis and fermentation leads to distinct isotope enrichment signatures, which would

suggest the different metabolic Hg reduction pathways are controlled by separate processes. We

also compare our results to previously published isotope signatures for biotic and abiotic Hg

transformations to critically evaluate whether common controls exist for microbe-mediated Hg

transformations. Finally, we discuss whether current analytical capacity for isotope fractionation

can be used to track the contributions of such pathways in the environment.

METHODS

Cell growth conditions and bioreactor setup

All experiments with live cells employed the strain *Heliobacterium modesticaldum* Ice1 grown

in PYE medium. Phototrophic cultures were grown at a constant visible light intensity of 80

μmol photon m^{-2} s^{-1} with a peak irradiance in the near-red to red spectrum (600 nm to 700 nm) at

50°C whereas chemotrophic cultures were grown in the dark at 50°C in line with our previous

work (23). *H. modesticaldum* Ice1 was obtained from the DSMZ culture collection (catalogue

number DSM-792). The bioreactor methodology employed in these experiments was similar to

the one used in previous work (23) with the following modifications: the bioreactor was

periodically supplied with sterile nitrogen gas that passed through an activated carbon filter and a

0.2 µm pore size air filter at a flow rate of 0.25 L min^{-1} and two gold traps were connected in

series at the reactor outlet to capture Hg^0 (**Fig S1**).

The bioreactor was kept in an incubator set to 50°C. Phototrophic experiments were

performed using a 60 W incandescent light bulb with a visible light intensity being maintained at

20 µmol photon m^{-2}s^{-1} and mirroring the irradiance of growth conditions. Chemotrophic

experiments were performed in the dark. Background Hg^0 was purged and removed from

reactors prior to phototrophically or chemotrophically-grown *H. modesticaldum* cells being

added as a 10 % (v/v) inoculum wherein an additional subsample was withdrawn to verify initial

cell density via O.D. 600 nm. For sterile treatments, no cells were added to the bioreactor and the

volume of growth medium was adjusted to ensure experiments were carried out at a working

volume of 500 mL. In all experiments, NIST 3133, a standard with known Hg isotopic

composition, was added to the bioreactor to approximately10 nM Hg final concentration.

Additional details on bioreactor setup are provided in **Supporting Methods**.

Hg^0 measurements

The bioreactor was bubbled periodically rather than continuously in contrast to previous work

because continuous bubbling led to fouling of gold traps and poor total Hg (THg) and isotope

recoveries. Subsamples were collected at designated timepoints by connecting dual gold traps to

the reactor, purging the reactor for 30 min to collect Hg^0. In the final 5 mins the reactor liquid

was sampled for Hg^{II} concentration, isotopic composition, and cell growth after which gas flow

was stopped. All gold traps were capped to prevent Hg^0 loss and aqueous samples were

conserved with 1% (v/v) trace element grade HCl at 4°C in the dark. Reactor washes were

performed following previously established protocols (23).

Total Hg analyses and sample preparation for stable isotope measurements

All samples for aqueous Hg analyses were oxidized with 10 % (v/v) bromine chloride (BrCl) in line with previous work (40) and THg concentrations were determined by stannous chloride reduction coupled to cold vapor atomic fluorescence spectrometry (42). During THg analysis, duplicates and matrix standard spikes were analyzed every 10 samples; analyses showed less than 10 % relative percent difference between duplicates and spike recoveries of 90-110%. The detection limit was 0.2 pM.

Total Hg mass balances were calculated in line with previous work (23, 24) with corrections to account for the mass of Hg removed for each subsample. For Hg^0 collected onto gold traps, a thermal desorption system was used to liberate Hg^0 over a 40 min desorption cycle which was then captured in 40% HNO_3:BrCl (3:1) oxidizing solution. An aliquot of the trapping solution was subsequently analyzed for THg concentration following the protocol established in previous work (43). Desorption efficiency during pre-concentration was tested by trapping a NIST 3133 standard with each batch of samples, and recoveries for standard trapping were 101.27 ± 4.26 (n = 15). THg recoveries from bioreactor experiments were 94.90 ± 9.50 % (n=7) (**Fig S2**).

Hg stable isotope analyses

Using the previously determined THg concentration, an aliquot of the trapping solution was diluted with ultra-high purity water to an acid content < 10% H^+ and a THg concentration of 3.75 to 5.00 nM (44). A Neptune Plus multicollector-inductively coupled plasma mass spectrometer (MC-ICP-MS) was used for Hg stable isotope ratio measurements. A concentration and matrix-matched NIST 3133 Hg was used for standard sample bracketing (45). To achieve detection at

low Hg concentrations, Hg was reduced with stannous chloride in-line then introduced

continuously into a custom designed gas liquid separator along with a thallium standard (NIST

SRM 997, 195 nM in 3% HCl) for mass bias correction (44, 46). MC-ICP-MS instrument

conditions and analytical expectations derived from previously published work and quality

control metrics for Hg stable isotope analysis are provided in **Supporting Methods, Table S1**

and **Table S2**.

Isotope calculations

Delta calculations followed the conventions set forth by others (45). This convention calls for

mass dependent fractionation (MDF) to be expressed in terms of δ^{xxx}Hg where graphically

δ^{202}Hg is used to signify MDF. δ^{xxx}Hg is calculated as:

$$\delta^{xxx}\text{Hg}(‰) = [(^{xxx}\text{Hg}/^{198}\text{Hg}_{sample})/(^{xxx}\text{Hg}/^{198}\text{Hg}_{\text{NIST-3133}}) - 1] \times 1000. \quad (\textbf{Eq 1})$$

XXX is used to signify the isotope of interest. Hg also undergoes MIF of both even and odd

isotopes. Here, odd-MIF is described by Δ^{199}Hg and even-MIF by Δ^{200}Hg. MIF is calculated as:

$$\Delta^{xxx}\text{Hg} \approx \delta^{xxx}\text{Hg} - (\delta^{202}\text{Hg} * \beta). \quad (\textbf{Eq 2})$$

β denotes the mass dependent scaling constant, which is determined by the laws of mass

dependence (45). Isotope enrichment effects based on the ratio of products to reactants (i.e. $\varepsilon_{p/r}$,

herein referred to as ε_{AP} for anoxygenic phototrophic reduction and ε_{FM} for fermentative

reduction) were calculated using Rayleigh fractionation models to account for the Hg^0 that was

removed following methods from previous work on microbial Hg reduction (34).

This method involved fitting a linear regression (shown in **Fig S3, Fig S4,** and **Table S3**) based

on the change in relative isotope ratios, $\ln(R/R_0)$, as a function of $\ln(f_r)$ where:

196             $R = (\delta^{202}Hg/1000) + 1$ for Hg^{II} (reactant) or Hg^0 (product) at a given time point and

197                     $R_0 = (\delta^{202}Hg/1000) + 1$ at the beginning of the experiment (**Eq 3**);

198     $f_r = Hg^{II}$ in the reactor at a given time / Hg^{II} in the reactor at the start of the experiment (**Eq 4**);

199   and the slope of the linear regression, which is equivalent to ε_{AP} or ε_{FM}:

200                     $$\varepsilon = \ln(R/R_0)/\ln(f_r) \ (\textbf{Eq 5}).$$

We chose to use the total Hg analyses from the bioreactor to establish f_r as this is a true

representation of the instantaneous isotope fractionation rather than a time integrated sample

following the methods outlined in previous work (34). Note that in our slope calculations we do

not have data for $^{202}Hg^0$ when $f_r = 1$ (the 0h sampling point) because no isotope fractionation was

taking place prior to the start of Hg reduction (**Table S2** and **Table S3**). Additional calculations

for isotope mass balances were carried out to test for deviation from the Rayleigh fraction model

and can be found in **Table S4.** ε values for MDF were compiled from previous studies on pure

microbial cultures and abiotic Hg transformations for comparison purposes. Currently, there is

no standard guideline for how to calculate ε and studies vary widely in how they present isotope

enrichment data and their associated variability. We have provided details on how we normalized

ε values from different studies in **Table S5**. In this work we only compared enrichment factors

from studies that used MC-ICP-MS techniques that were in line with our analyses, however we

have included all of the enrichment factors for microbe-driven Hg transformations published to

date in **Table S5**.

RESULTS AND DISCUSSION

Hg^0 production during anoxygenic photosynthesis and fermentation

In our experiments Hg^0 production relied on the presence of live cells for both chemotrophic and phototrophic growth conditions. Cumulative Hg^0 production was an order of magnitude higher for live cell treatments compared to abiotic controls (>0.60 vs <0.05 nmol) (**Fig 1**). Chemotrophic cells produced double the amount of Hg^0 compared to phototrophic cells (1.33 to 1.64 nmol vs 0.60 to 0.81 nmol, respectively) despite having been supplied with slightly less Hg^{II} (**Fig 1B, E**). In relative terms, phototrophic cells reduced 12 to 15% of the initial Hg^{II} supplied whereas chemotrophic cells reduced 31 to 40% (**Fig S2**).

Relative Hg^0 production in these experiments was lower, yet consistent with our previous work, which showed chemotrophically-grown cells reduced 15% more Hg^{II} than phototrophically-grown cells, reducing between 60 to 75% of the initial Hg^{II} supplied (23). The relative amount of Hg^0 production did not increase with higher Hg^{II} exposure (10 nM) versus previous experiments (250 pM) (23). It is unlikely that the lower reduction rate observed in the experiments presented here was due to the toxicity of 10 nM Hg supplied, given that *H. modesticaldum* grows well under the same conditions at 50 times higher Hg concentrations (23). Instead, we suspect that the lack of constant bubbling of the reactor may be responsible for the observed results. It is possible that the accumulation of CO_2 in the bioreactor following organic carbon oxidation may compete with Hg^{II} as an electron sink, decreasing the amount of Hg^0 produced. Inorganic carbon is required for anaplerotic reactions that fulfill the biosynthetic needs of *H. modesticaldum* and CO_2 could have competed with Hg^{II} for reducing power, as previously shown (47). These results could also be due to Hg^0 oxidation allowed by the increased residence

239　time of Hg⁰ in the reactor afforded by the intermittent sparging, as has been observed in other

240　anaerobes (48, 49).

Figure 1: Total Hg and cumulative Hg^0 production by *H. modesticaldum* grown phototrophically and chemotrophically. A, D THg for live cells and sterile controls without cells. **B, E** Cumulative Hg^0 production for live cells and sterile controls without cells. **C, F** Microbial growth as measured by O.D. 600 nm for live cells and sterile controls without cells. 10 nM Hg was supplied in all experiments from the same stock (NIST 3133).

Despite carefully controlling the growth conditions in our experiments, chemotrophic

cells from replicate #1 exhibited a lag phase of 12 hours (**Fig 1F**). Although this lag did not

affect final cell density, the slow-growing culture from replicate #1 initiated Hg^0 production

earlier than the culture in replicate #2 (**Fig 1E, F**). These results may be attributable to cells in

replicate #1 maintaining a higher Hg to cell ratio in the first 12 hours of the experiment, which

suggests that Hg is being reduced intracellularly in line with our previous work on Heliobacteria

(23, 50). Indeed, even at low initial densities, cells can maintain µM levels of intracellular

reducing power, which could be used to reduce nM levels of Hg^{II} (24).

Mass dependent fractionation during photosynthetic and fermentative growth

Hg reduction in phototrophically grown *H. modesticaldum* resulted in consistent positive MDF

whereas fermentatively grown cultures showed positive MDF with variability in fractionation

patterns across replicates (**Fig 2**). Abiotic controls showed minimal fractionation for both MDF

and MIF, confirming that Hg reduction and subsequent isotopic fractionation were

predominantly driven by cellular processes (**Fig 2**). Live cell experiments were devoid of MIF

(**Table S1** and **Table S2**).

For phototrophically-grown cultures, the reactant pool $\delta^{202}Hg^{II}$ increased steadily over time

suggesting cells preferentially reduced lighter Hg^{II} (**Fig 2A**). A mirror trend was observed for

$\delta^{202}Hg^0$, which was depleted in $^{202}Hg^0$ at the beginning of the experiment, became enriched with

heavier isotopes as the reaction proceeded, and eventually approached the initial isotopic

composition of the NIST 3133 standard (0‰) (**Fig 2B**).

Figure 2: Mass dependent fractionation of ^{202}Hg in Hg^{II} (A, C) and Hg^0 (B, D) during phototrophic and chemotrophic growth of *H. modesticaldum* and no cell controls. δ^{202}Hg values for Hg^{II} and Hg^0 are plotted with respect to the fraction of remaining total inorganic Hg^{II} in the bioreactor. Insufficient Hg^0 was recovered from the no cell controls and thus no results are displayed. 10 nM Hg was used in all experiments and Hg was supplied from the same stock (NIST 3133).

Curiously, the $\delta^{202}Hg^0$ for f_r = 0.94 at 3h in the two phototrophic live cell replicates showed

that the product pool was initially enriched with the $^{202}Hg^0$ isotope (**Fig 2B**). A similar pattern

has been observed in iron reducing bacteria that preferentially reduced heavier Fe^{III} (51) and the

model anaerobic Hg methylator and reducer *Geobacter sulfurreducens* PCA (37, 40). The most

recent work on *G. sulfurreducens* has shown that uptake of Hg selects for lighter Hg isotopes but

cells can also access an isotopically heavier pool of Hg^{II} from the dissolved phase (37). It is

possible that a similar process is occurring for phototrophically-grown *H. modesticaldum*

wherein cells access an isotopically heavier pool of Hg during equilibrium binding of Hg^{II} to the

outside of the cell, followed by an alternative uptake process that selects for lighter pools of Hg.

Based on these results, we included all time points in isotopic enrichment calculations for Hg^{II}

but omitted the 3h time point in calculations for Hg^0 given that other cellular fractionation

processes could be occurring (**Fig S3, Fig S4,** and **Table S3**).

Positive MDF was also observed for chemotrophic cultures but fractionation patterns differed

strongly between replicates. Replicate #1 showed an initial depletion in reactant pool $\delta^{202}Hg^{II}$ and

became progressively enriched with $^{202}Hg^{II}$ relative to the NIST 3133 suggesting lighter isotopes

were still preferentially reduced (**Fig 2C**). Note that the second $\delta^{202}Hg^{II}$ data point available for

replicate #1 occurs at f_r=0.72 due to the rapid rate of Hg^{II} reduction leading to substantially less

Hg remaining in the reactor (**Fig 1E** and **Fig 2C**). Results for replicate #2 further suggested that

lighter Hg^{II} isotopes were preferentially reduced with $\delta^{202}Hg^{II}$ slightly increasing at the

beginning of the experiment before rising to a maximum of 0.72 ‰ between f_r = 0.85 at 12h and

0.53 at 24h (**Fig 2C**).

The trends for $\delta^{202}Hg^0$ in chemotrophic cultures mirrored what was observed for $\delta^{202}Hg^{II}$.

$\delta^{202}Hg^0$ from replicate #1 showed that the product pool was initially enriched with heavy $^{202}Hg^0$,

further suggesting cells may have accessed a readily bioavailable pool of heavy Hg^{II}, before

undergoing a pronounced depletion and progressive enrichment with heavy isotopes similar to

phototrophic cultures (**Fig 2B, D**). In contrast, the $\delta^{202}Hg^0$ values for cells in replicate #2 suggest

that the product pool was depleted in $^{202}Hg^0$ before undergoing enrichment for heavier Hg^0

isotopes reaching a maximum of 1.73 ‰ (**Fig 2D**).

The enrichment in $^{202}Hg^0$ exceeded the initial composition of the NIST 3133 standard (0 ‰)

suggesting fractionation deviated from the predicted Rayleigh fractionation for Hg reduction

(**Table S4**). Recent work with *G. sulfurreducens* has shown that subcellular partitioning

processes contribute to Hg isotope fractionation and cannot be detected at the level of THg pool

(37). In our work, Hg reduction provided such a strong isotopic shift that we could detect

fractionation in the THg pool, however we cannot discount the contributions of subcellular

partitioning processes, which are smaller in magnitude (<0.10 ‰). Such processes could be

driving the depletion of $^{202}Hg^{II}$ early on for chemotrophic cells in replicate #1 where low cell

densities may have amplified uptake and adsorption-driven fractionation (**Fig 1F**). It is more

challenging to discuss contributions of such processes for replicate #2, where the low cumulative

Hg^0 production at $f_r = 0.94$ and 0.88 (3h and 6h time points, respectively) precluded our ability to

measure Hg^0 isotope fractionation (**Fig 1E** and **Table S2**). Despite this limitation, the high

$\delta^{202}Hg^0$ value obtained for the final timepoint for replicate #2 reinforces that additional processes

are contributing to fractionation under chemotrophic conditions that warrant further exploration.

Our experimental results suggest that Hg^0 oxidation is not a major contributor at earlier

stages of reduction (i.e., 3-12 hours) but may be contributing to isotope fractionation at later time

points associated with longer Hg residence times in the reactor. Under phototrophic conditions,

the measured values of $\delta^{202}Hg^{II}$ at 48h were isotopically heavier than predicted values, which

suggests Hg^0 oxidation may be occurring (**Table S4**). Under fermentative conditions the

potential contribution of Hg oxidation is unclear, since a consistent enrichment was not observed

between replicates (**Table S4**). The variability in isotope results for later time points in both

replicates suggests processes other than reduction may be occurring.

Although Hg^0 oxidation has never been observed in anoxygenic phototrophs, it has been

observed in other model chemotrophic anaerobes (49). Recent work showed that thiol-bearing

molecules preferentially oxidize heavy $^{202}Hg^0$, leading to negative MDF, in addition to a MIF

signal (52). Although we did not observe any negative MDF or MIF in our experiments (**Table**

S1 and **Table S2**), we acknowledge that such processes have the potential to contribute to net

fractionation alongside internal partitioning related to uptake or adsorption processes. These

results are an important first step to resolving the physiological processes that drive these Hg

transformations in *H. modesticaldum* and we plan to carry out additional experiments to assess

the contributions of uptake, adsorption, and redox transformations to the net Hg isotope

fractionation observed in the future.

Comparing enrichment factors for abiotic and biotic Hg transformations

The average isotopic enrichment factor calculated using the ratio of product to reactant ($\varepsilon_{p/r}$) ± 2

standard deviations (2SD) for MDF in the reactant pool Hg^{II} during anoxygenic photosynthesis

(ε_{AP}) was -0.80 ± 0.56 ‰ whereas the average isotopic enrichment factor obtained during

fermentation (ε_{FM}) was -1.13 ± 0.50 ‰ (**Fig 3, Fig S3, Table S3,** and **Table S5**).

Figure 3: Compilation of isotopic enrichment factors (ε) for abiotic and biotic Hg transformations. Hg transformations are denoted by different shapes and specific metabolic and chemical pathways have been colour coded. Error bars denote 2 standard deviations for studies where this data was available. In the event that data was presented as a standard error, standard deviation was calculated based on the sample size indicated in each study. Abbreviations on the y-axis denote the following: UV (ultraviolet light), DOC (dissolved organic carbon), AP (anoxygenic photosynthesis), ΔhgcAB (no *hgcAB* gene cluster), and FM (fermentation). The grey-shading in this figure represents light and oxygen. Lighter regions represent habitats where light and oxygen are present and darker regions represent where they are not. The values in this figure were obtained from the following studies: abiotic dark oxidation (52); abiotic photoreduction (32, 41); aerobic MerA reduction (34, 36); anaerobic respiratory methylation (40); anaerobic respiratory reduction (36, 37); anoxygenic photosynthetic reduction (this study); fermentative reduction (this study); oxygenic photosynthetic reduction (41). The raw data in this figure are in **Table S5**.

The enrichment factor could not be calculated using the product Hg^0 pool for fermentative

cultures and overestimated the value in phototrophic cultures (-2.86 ± 1.32 ‰) (**Fig S4** and

Table S3). Previous work has shown that using the product pool can overestimate the

fractionation factor due to difficulty obtaining analytical replicates of gaseous Hg^0 and the

complexity of quantitatively collecting this pool (53). As such, we carried out comparisons for

enrichment factors using the reactant Hg^{II} pool. Ideally, we would have been able to obtain more

replicates of our experiments but were unable to do so due the challenges stated previously. It is

important to note that low numbers of replicates are routinely discussed in isotope geochemistry

and the variability from our results is in line with previous work (**Fig 3** and **Table S5**).

Our initial motivation for this study was to test if phototrophic and fermentative Hg

reduction led to markedly different isotope enrichment factors and establish whether different

underlying processes supported different metabolic Hg reduction pathways. When comparing ε

values, which are used as an index to distinguish between different isotope fractionation

processes, we report a difference of 0.33 ‰ between ε_{AP} and ε_{FM} (**Fig 3** and **Table S5**). This

small difference in enrichment factors supports that common processes are controlling Hg

reduction during anoxygenic photosynthesis and fermentation in *H. modesticaldum*.

In our previous work with *H. modesticaldum* we demonstrated that inhibiting pyruvate

ferredoxin oxidoreductase, the enzyme responsible for the production of reduced ferredoxin,

greatly hampered Hg^{II} reduction during anoxygenic photosynthesis and fermentation (23). It is

possible that the similarity between ε_{AP} and ε_{FM} is the result of reactions originally involving the

same redox cofactors (e.g. ferredoxins) but with Hg^{II} being reduced by different electron donors

specific to each metabolism. Such reactions could involve direct electron transfer from

ferredoxin to Hg^{II} or from enzymes directly or indirectly relying on ferredoxin (23, 54). It is also

possible that the net isotopic enrichment stems from a combination of multiple processes

involved in Hg transport (e.g. adsorption, uptake, and efflux) that would occur regardless of the

intracellular transformations in question. Our results suggest that the relative contributions of

these processes may be more pronounced during fermentation compared to anoxygenic

photosynthesis although more controlled experiments are required to constrain the variability in

the isotope fractionation observed.

At the broader scale, the isotopic enrichment observed for both metabolisms tested in *H.*

modesticaldum falls between the values observed for abiotic and microbial Hg reduction

pathways (**Fig 3** and **Table S5**). The isotopic enrichment observed for phototrophic *H.*

modesticaldum is slightly lower than photoreduction in the presence of dissolved organic carbon

(-0.80 ± 0.56 ‰ vs -0.60 ± 0.20 ‰, respectively) (32) but higher than photoreduction in the

presence of marine exudates and UVB light (-0.80 ± 0.56 ‰ vs -1.47 ‰, respectively) (41) (**Fig**

3 and **Table S5**). When comparing Hg reduction in the anoxygenic phototroph *H. modesticaldum*

to the oxygenic phototrophic green alga *Isochrysis galbana*, the isotopic enrichment observed is

also slightly lower (-0.80 ± 0.56 ‰ vs -0.14 to -0.60 ‰, respectively) (41) (**Fig 3** and **Table S5**).

Though this similarity suggests anoxygenic and oxygenic photosynthetic Hg reduction share a

common physiological pathway, the presence of MIF in *I. galbana* due to free radical generation

within the cell rules out this possibility (41). (**Table S1**). Thus, Hg reduction during anoxygenic

photosynthetic Hg reduction could be distinguished from other photo-induced reduction

pathways due to the lack of MIF tracer.

When compared with previous work on microbial Hg reduction in pure cultures, the isotopic

enrichment observed for phototrophically-grown *H. modesticaldum* was higher than what has

previously been reported for aerobic *mer*-mediated reduction and anaerobic respiratory Hg

reduction (-0.80 ± 0.56 ‰ vs -1.80 to -1.20 ‰, respectively) (34, 36, 37) (**Fig 3** and **Table S5**).

Interestingly, the isotopic enrichment observed in fermentatively-grown *H. modesticaldum* (-1.13

± 0.50 ‰) was closer to aerobic *mer*-mediated reduction (-1.40 to -1.20 ‰) (34) than to

anaerobic respiratory Hg reduction in *Shewanella oneidensis* MR-1 and a modified strain of *G.*

sulfurreducens PCA incapable of Hg methylation (-1.80 to -1.60 ‰) (**Fig 3** and **Table S5**) (36,

37). An additional line of evidence supporting that Hg^0 oxidation is unlikely to be contributing to

the isotope signature observed in our system are the ε values for thiol mediated Hg^0 oxidation in

the dark. These values are considerably larger than what we observed for *H. modesticaldum*

(1.51 ± 0.20 ‰ for cysteine and 1.47 ± 0.24 ‰ for glutathione) (**Fig 3** and **Table S5**)**.**

These comparisons illustrate that isotopic enrichment for microbial Hg reduction

pathways that only display MDF fall within a narrow range (i.e. -0.80 to -1.80 ‰) (**Fig 3** and

Table S5). We find this to be a strikingly small difference given the ecological diversity of the

model organisms and variety of growth conditions used to study microbial Hg reduction. The

similar ε values reported for fermentation in this study and the *mer* operon are also noteworthy as

they suggest nearly identical isotope enrichment can occur in aerobes harbouring dedicated

enzymatic machinery to reduce Hg and anaerobes where *mer*-based strategies are largely absent.

Our study also shows that similar isotopic enrichment can be observed for different

anaerobic Hg transformations. The anaerobic Hg methylating chemotrophs *G. sulfurreducens*

PCA and *Desulfovibrio desulfuricans* ND132 displayed ε signatures (i.e. -0.92 and -1.1‰,

respectively) that fall within the range of ε values we report in our study (i.e. -0.80 to -1.13 ‰)

(**Fig 3** and **Table S5**) (40). The similarity in enrichment signatures suggests that common

processes may affect the delivery of inorganic Hg substrates to intracellular sites independently

of the Hg transformations in question. Although it is outside of the scope of the current study,

identifying the nature of these processes may provide insights into similar controls that exist for

Hg uptake and transformation in anaerobes.

Our work showcases the strengths but also the challenges of using isotope fractionation-

based approaches to decipher cryptic Hg cycling pathways. To illustrate the challenges of using

isotopes to track Hg cycling pathways in the environment, we visualized our comparison of ε

values in the context of a vertically redox stratified habitat (e.g. a water column) (**Fig 3**).

Through Hg stable isotope fractionation, we can distinguish between abiotic and biotic redox

reaction pathways that control Hg's availability in methylation hotspots, such as the limit of the

photic zone, on the bases of MIF signatures associated with photochemical reactions. In contrast,

teasing apart biological pathways that contribute to the net isotope signature of Hg^{II} in the

environment is more challenging because competing MDF processes occur in both dark and

photo-mediated pathways where Hg^{II} is a reactant (e.g. reduction and methylation) and a product

(e.g. demethylation and oxidation). This overlap makes it difficult to translate $\delta^{202}Hg$ values

observed under controlled conditions that are sensitive to changes in cellular physiology to

environments where pools of bioavailable Hg are not homogenous due to variable mixing.

CONCLUSION

In this study we have provided the first evidence that common physiological processes drive

MDF during Hg reduction associated with anoxygenic photosynthesis and fermentation in the

model anaerobe *H. modesticaldum*. We show that isotopic enrichment during anoxygenic

phototrophic Hg reduction leads to an intermediate fractionation process that lies between

photochemical reactions and dark microbial Hg reduction. We also show that fermentative Hg

reduction leads to isotopic enrichment that is similar to aerobic *mer*-mediated reduction.

Although current analytical capacities to measure stable isotope fractionation make

applying such approaches to the environment challenging, this is a burgeoning field that has

advanced considerably in terms of sensitivity and the different Hg species that can be targeted in

the last 10 years. Isotope fractionation approaches have proved extremely valuable for teasing

apart the complex interwoven processes that drive microbe-mediated Hg transformations and as

seen in our study, highlighting common processes that control Hg bioavailability to microbes.

In future work, it will be important to isolate the different steps involved in Hg uptake

and transformation by rigorously controlling Hg availability and cellular physiology. Leveraging

molecular tools alongside isotopes will be a useful strategy to address these knowledge gaps and

the recent availability of genetically tractable *H. modesticaldum* deletion mutants offers a

promising means to do so (55). In that regard, gene deletion approaches targeting redox-active

enzymes could be used to characterize the physiological components contributing directly to Hg^{II}

reduction, whereas targeting homologues for proteins involved in Hg uptake in other model

anaerobes could be the key to identifying entry routes for Hg into *H. modesticaldum* (56, 57).

In addition to gene deletions, whole-genome, transcriptome, and proteome approaches

could be combined with isotopic analyses in pure cultures to hone in on potential biomarkers for

Hg transformation pathways that remain poorly understood. Such biomarkers could be used to

examine publicly available genomic datasets to assess the distribution of anaerobic Hg reducers

in modern day environments and rock records, the latter serving as an archive for Hg isotope

signatures and the microbial community involved in historical Hg cycling (31, 58). These ancient

environments were likely dominated by anoxygenic phototrophs that were important primary

producers before the rise of oxygen on Earth. By using isotopes to create a window into the past,

we can compare the selective forces that gave rise to fortuitous Hg reduction pathways tied to

anaerobic metabolisms and dedicated detoxification strategies such as the *mer* operon.

ACKNOWLEDGMENTS

482 Our work was funded by NSERC Discovery and Accelerator grants, CFI funding to AJP and

483 NSERC graduate scholarship to DSG. Instrumentation and lab operations for Hg isotope

484 analyses were supported by the USGS Toxic Substances Hydrology Program. Any use of trade,

485 firm, or product names is for descriptive purposes only and does not imply endorsement by the

486 U.S. Government.

AUTHOR CONTRIBUTIONS

489 DSG, AJP, SEJ and MTT designed all experiments; DSG and NCL carried out bioreactor

490 experiments and data analyses; SEJ and MTT carried out total Hg and stable isotope analyses;

491 DSG, AJP, SEJ and MTT wrote the manuscript.

REFERENCES

REFERENCES

1.   Selin NE. 2009. Global biogeochemical cycling of mercury: A review. Annual Review of

Environment and Resources 34:43-63.

2.   Parks JM, Johs A, Podar M, Bridou R, Hurt Jr RA, Smith SD, Tomanicek SJ, Qian Y,

Brown SD, Brandt CC, Palumbo AV, Smith JC, Wall JD, Elias DA, Liang L. 2013. The

genetic basis for bacterial mercury methylation. Science 339:1332-1335.

3.   Fleming EJ, Mack EE, Green PG, Nelson DC. 2006. Mercury methylation from

unexpected sources: Molybdate-inhibited freshwater sediments and an iron-reducing

bacterium. Applied and Environmental Microbiology 72:457-464.

4.   Han S, Narasingarao P, Obraztsova A, Gieskes J, Hartmann AC, Tebo BM, Allen EE,

Deheyn DD. 2010. Mercury speciation in marine sediments under sulfate-limited

conditions. Environmental Science and Technology 44:3752-3757.

5.   Podar M, Gilmour CC, Brandt CC, Soren A, Brown SD, Crable BR, Palumbo AV,

Somenahally AC, Elias DA. 2015. Global prevalence and distribution of genes and

microorganisms involved in mercury methylation. Science Advances 1:1-13.

6.   Schartup AT, Balcom PH, Soerensen AL, Gosnell KJ, Calder RS, Mason RP, Sunderland

EM. 2015. Freshwater discharges drive high levels of methylmercury in Arctic marine

biota. Proceedings of the National Academy of Sciences of the United States of America

112:11789-94.

7.   Hsu-Kim H, Kucharzyk KH, Zhang T, Deshusses MA. 2013. Mechanisms regulating

mercury bioavailability for methylating microorganisms in the aquatic environment: a

critical review. Environmental Science and Technology 47:2441-56.

8.   Gustin MS, Biester H, Kim CS. 2002. Investigation of the light-enhanced emission of

mercury from naturally enriched substrates. Atmospheric Environment 36:3241-3254.

9.   Amyot M, Lean D, Mierle G. 1997. Photochemical formation of volatile mercury in high

Arctic lakes. Environmental Toxicology and Chemistry 16:2054-2063.

10.   Whalin L, Kim EH, Mason R. 2007. Factors influencing the oxidation, reduction,

methylation and demethylation of mercury species in coastal waters. Marine Chemistry

107:278-294.

11.   Poulain AJ, Amyot M, Findlay D, Findlay S, Telor S, Barkay T, Hintelmann H, Amyot

M, Findlay D. 2011. Biological and photochemical production of dissolved gaseous

mercury in a boreal lake. Limnology 49:2265-2275.

12.   Carpi A, Lindberg SE. 1997. Sunlight-mediated emission of elemental mercury from soil

amended with municipal sewage sludge. Environmental Science and Technology

31:2085-2091.

13.   Vandal GM, Fitzgerald WF, Rolfhus KR, Lamborg CH. 1995. Modeling the elemental

mercury cycle in Pallette Lake, Wisconsin, USA. Water, air, and soil pollution 80:529-

538.

14.   Zhang H, Lindberg SE. 2001. Sunlight and iron(III)-induced photochemical production

of dissolved gaseous mercury in freshwater. Environmental Science and Technology

35:928-935.

15.   Zhang H. 2006. Photochemical redox reactions of mercury, p 37-79. *In* Atwood DA (ed),

Recent Developments in Mercury Science, vol 120.

16.   O'Driscoll NJ, Lean DRS, Loseto LL, Carignan R, Siciliano SD. 2004. Effect of

dissolved organic carbon on the photoproduction of dissolved gaseous mercury in lakes:

Potential impacts of forestry. Environmental Science and Technology 38:2664-2672.

17.   Gu B, Bian Y, Miller CL, Dong W, Jiang X, Liang L. 2011. Mercury reduction and

complexation by natural organic matter in anoxic environments. Proceedings of the

National Academy of Sciences of the United States of America 108:1479-1483.

18.   Wiatrowski HA, Das S, Kukkadapu R, Ilton ES, Barkay T, Yee N. 2009. Reduction of

Hg(II) to Hg(0) by magnetite. Environmental Science and Technology 43:5307-5313.

19.   Bone SE, Bargar JR, Sposito G. 2014. Mackinawite (FeS) reduces mercury(II) under

sulfidic conditions. Environmental Science and Technology 48:10681-10689.

20.   Wiatrowski HA, Ward PM, Barkay T. 2006. Novel reduction of mercury(II) by mercury-

sensitive dissimilatory metal reducing bacteria. Environmental Science and Technology

40:6690-6696.

21.   Lu X, Liu Y, Johs A, Zhao L, Wang T, Yang Z, Lin H, Elias DA, Pierce EM, Liang L,

Barkay T, Gu B. 2016. Anaerobic mercury methylation and demethylation by *Geobacter*

bemidjiensis Bem. Environmental Science and Technology 50:4366-4373.

22.   Lin H, Morrell-Falvey JL, Rao B, Liang L, Gu B. 2014. Coupled mercury-cell sorption,

reduction, and oxidation on methylmercury production by *Geobacter sulfurreducens*

PCA. Environmental Science and Technology 48:11969-11976.

23.   Grégoire DS, Lavoie NC, Poulain AJ. 2018. Heliobacteria reveal fermentation as a key

pathway for mercury reduction in anoxic environments. Environmental Science and

Technology 52:4145-4153.

24.   Grégoire DS, Poulain AJ. 2016. A physiological role for Hg(II) during phototrophic

growth. Nature Geoscience 9:121-125.

25.   Barkay T, Kritee K, Boyd E, Geesey G. 2010. A thermophilic bacterial origin and

subsequent constraints by redox, light and salinity on the evolution of the microbial

mercuric reductase. Environmental Microbiology 12:2904-2917.

26.   Grégoire DS, Poulain AJ. 2014. A little bit of light goes a long way: the role of

phototrophs on mercury cycling. Metallomics 6:396-407.

27.   Poulin BA, Aiken GR, Nagy KL, Monceau A, Krabbenhoft DP, Ryan JN. 2016. Mercury

transformation and release differs with depth and time in a contaminated riparian soil

during simulated flooding. Geochimica Et Cosmochimica Acta 176:118-138.

28.   Bouffard A, Amyot M. 2009. Importance of elemental mercury in lake sediments.

Chemosphere 74:1098-1103.

29.   Grégoire DS, Poulain AJ. 2018. Shining light on recent advances in microbial mercury

cycling. FACETS 3:858-879.

30.   Blum JD, Sherman LS, Johnson MW. 2014. Mercury isotopes in earth and environmental

sciences. Annual Review of Earth and Planetary Sciences 42:249-269.

31.   Zheng W, Gilleaudeau GJ, Kah LC, Anbar AD. 2018. Mercury isotope signatures record

photic zone euxinia in the Mesoproterozoic ocean. Proceedings of the National Academy

of Sciences 115:10594-10599.

32.   Bergquist BA, Blum JD. 2007. Mass-dependent and -independent fractionation of Hg

isotopes by photoreduction in aquatic systems. Science 318:417-420.

33.   Bergquist BA, Blum JD. 2009. The odds and evens of mercury isotopes: Applications of

mass-dependent and mass-independent isotope fractionation. Elements 5:353-357.

34. Kritee K, Blum JD, Johnson MW, Bergquist BA, Barkay T. 2007. Mercury stable isotope

fractionation during reduction of Hg(II) to Hg(0) by mercury resistant microorganisms.

Environmental Science and Technology 41:1889-1895.

35. Kritee K, Barkay T, Blum JD. 2009. Mass dependent stable isotope fractionation of

mercury during mer mediated microbial degradation of monomethylmercury.

Geochimica et Cosmochimica Acta 73:1285-1296.

36. Kritee K, Blum JD, Barkay T. 2008. Mercury stable isotope fractionation during

reduction of Hg(II) by different microbial pathways. Environmental Science and

Technology 42:9171-9177.

37. Wang Y, Janssen SE, Schaefer JK, Yee N, Reinfelder JR. 2020. Tracing the uptake of

Hg(II) in an iron-reducing bacterium using mercury stable isotopes. Environmental

Science & Technology Letters 7:573-578.

38. Rodríguez-González P, Epov VN, Bridou R, Tessier E, Guyoneaud R, Monperrus M,

Amouroux D. 2009. Species-specific stable isotope fractionation of mercury during

Hg(II) methylation by an anaerobic bacteria (*Desulfobulbus propionicus*) under dark

conditions. Environmental Science and Technology 43:9183-9188.

39. Perrot V, Bridou R, Pedrero Z, Guyoneaud R, Monperrus M, Amouroux D. 2015.

Identical Hg isotope mass dependent fractionation signature during methylation by

sulfate-reducing bacteria in sulfate and sulfate-free environment. Environmental Science

and Technology 49:1365-1373.

40. Janssen SE, Schaefer JK, Barkay T, Reinfelder JR. 2016. Fractionation of mercury stable

isotopes during microbial methylmercury production by iron- and sulfate-reducing

bacteria. Environmental Science and Technology 50:8077-8083.

41.   Kritee K, Motta LC, Blum JD, Tsui MT-K, Reinfelder JR. 2018. Photomicrobial visible

light-induced magnetic mass independent fractionation of mercury in a marine microalga.

ACS Earth and Space Chemistry 2:432-440.

42.   EPA U. 2002. Method 1631, Revision E: Mercury in water by oxidation, purge and trap,

and cold vapor atomic fluorescence spectrometry. US Environmental Protection Agency

Washington, DC.

43.   Janssen SE, Lepak RF, Tate MT, Ogorek JM, DeWild JF, Babiarz CL, Hurley JP,

Krabbenhoft DP. 2019. Rapid pre-concentration of mercury in solids and water for

isotopic analysis. Analytica Chimica Acta 1054:95-103.

44.   Yin R, Krabbenhoft DP, Bergquist BA, Zheng W, Lepak RF, Hurley JP. 2016. Effects of

mercury and thallium concentrations on high precision determination of mercury isotopic

composition by Neptune Plus multiple collector inductively coupled plasma mass

spectrometry. Journal of Analytical Atomic Spectrometry 31:2060-2068.

45.   Blum JD, Bergquist BA. 2007. Reporting of variations in the natural isotopic

composition of mercury. Analytical and Bioanalytical Chemistry 388:353-359.

46.   Lepak RF, Yin R, Krabbenhoft DP, Ogorek JM, DeWild JF, Holsen TM, Hurley JP.

2015. Use of stable isotope signatures to determine mercury sources in the Great Lakes.

Environmental Science & Technology Letters 2:335-341.

47.   Tang KH, Feng XY, Zhuang WQ, Alvarez-Cohen L, Blankenship RE, Tang YJ. 2010.

Carbon flow of Heliobacteria is related more to Clostridia than to the Green Sulfur

Bacteria. Journal of Biological Chemistry 285:35104-35112.

48.   Colombo MJ, Ha J, Reinfelder JR, Barkay T, Yee N. 2013. Anaerobic oxidation of Hg(0)

626   and methylmercury formation by *Desulfovibrio desulfuricans* ND132. Geochimica et

627   Cosmochimica Acta 112:166-177.

628   49.   Colombo MJ, Ha J, Reinfelder JR, Barkay T, Yee N. 2014. Oxidation of Hg(0) to Hg(II)

629   by diverse anaerobic bacteria. Chemical Geology 363:334-340.

630   50.   Lavoie NC, Grégoire DS, Stenzler BR, Poulain AJ. 2020. Reduced sulphur sources

631   favour Hg(II) reduction during anoxygenic photosynthesis by Heliobacteria. Geobiology

632   18:70-79.

633   51.   Icopini GA, Anbar AD, Ruebush SS, Tien M, Brantley SL. 2004. Iron isotope

634   fractionation during microbial reduction of iron: The importance of adsorption. Geology

635   32:205-208.

636   52.   Zheng W, Demers JD, Lu X, Bergquist BA, Anbar AD, Blum JD, Gu B. 2019. Mercury

637   stable isotope fractionation during abiotic dark oxidation in the presence of thiols and

638   natural organic matter. Environmental Science and Technology 53:1853-1862.

639   53.   Kritee K, Blum JD, Reinfelder JR, Barkay T. 2013. Microbial stable isotope fractionation

640   of mercury: A synthesis of present understanding and future directions. Chemical

641   Geology 336:13-25.

642   54.   Tang KH, Yue H, Blankenship RE. 2010. Energy metabolism of *Heliobacterium*

643   *modesticaldum* during phototrophic and chemotrophic growth. Bmc Microbiology 10:150

644   55.   Baker PL, Orf GS, Kevershan K, Pyne ME, Bicer T, Redding KE. 2019. Using the

645   endogenous CRISPR-Cas system of *Heliobacterium modesticaldum* to delete the

646   photochemical reaction center core subunit gene. Applied and Environmental

647   Microbiology 85:e01644-19.

56.   Stenzler B, Hinz A, Ruuskanen M, Poulain AJ. 2017. Ionic strength differentially affects

the bioavailability of neutral and negatively charged inorganic Hg complexes.

Environmental Science and Technology 51:9653-9662.

57.   Schaefer JK, Szczuka A, Morel FM. 2014. Effect of divalent metals on Hg(II) uptake and

methylation by bacteria. Environmental Science and Technology 48:3007-13.

58.   Ruuskanen MO, Aris-Brosou S, Poulain AJ. 2020. Swift evolutionary response of

microbes to a rise in anthropogenic mercury in the Northern Hemisphere. The ISME

Journal 14:788-800.

 FIGURE CAPTIONS

Figure 1: Total Hg and cumulative Hg^0 production by *H. modesticaldum* grown

phototrophically and chemotrophically. A, D THg for live cells and sterile controls without

cells. **B, E** Cumulative Hg^0 production for live cells and sterile controls without cells. **C, F**

Microbial growth as measured by O.D. 600 nm for live cells and sterile controls without cells. 10

nM Hg was supplied in all experiments from the same stock (NIST 3133).

Figure 2: Mass dependent fractionation of ^{202}Hg in Hg^{II} (A, C) and Hg^0 (B, D) during

phototrophic and chemotrophic growth of *H. modesticaldum* and no cell controls. δ^{202}Hg

values for Hg^{II} and Hg^0 are plotted with respect to the fraction of remaining total inorganic Hg^{II}

in the bioreactor. Insufficient Hg^0 was recovered from the no cell controls and thus no results are

displayed. 10 nM Hg was used in all experiments and Hg was supplied from the same stock

(NIST 3133).

Figure 3: Compilation of isotopic enrichment factors (ε) for abiotic and biotic Hg

transformations. Hg transformations are denoted by different shapes and specific metabolic and

chemical pathways have been colour coded. Error bars denote 2 standard deviations for studies

where this data was available. In the event that data was presented as a standard error, standard

deviation was calculated based on the sample size indicated in each study. Abbreviations on the

y-axis denote the following: UV (ultraviolet light), DOC (dissolved organic carbon), AP

(anoxygenic photosynthesis), ΔhgcAB (no *hgcAB* gene cluster), and FM (fermentation). The

grey-shading in this figure represents light and oxygen. Lighter regions represent habitats where

light and oxygen are present and darker regions represent where they are not. The values in this

figure were obtained from the following studies: abiotic dark oxidation (52); abiotic

photoreduction (32, 41); aerobic MerA reduction (34, 36); anaerobic respiratory methylation

(40); anaerobic respiratory reduction (36, 37); anoxygenic photosynthetic reduction (this study);

fermentative reduction (this study); oxygenic photosynthetic reduction (41). The raw data in this

figure are in **Table S5**.

SUPPORTING INFORMATION CAPTIONS

Figure S1: Schematic of the bioreactor setup used to measure Hg stable isotope fractionation for

Hg^{II} and Hg^0 during the anaerobic reduction of Hg^{II}.

Figure S2: **Total Hg mass balance for isotope fractionation experiments under**

phototrophic and chemotrophic growth conditions. The average recovery across all

experiments was 94.40 ± 9.50 % (n=7) and is shown by the dashed red line.

Figure S3: Mass dependent fractionation of $^{202}Hg^{II}$ in phototrophically (A, B) and

chemotrophically (C, D) grown *H. modesticaldum*. $\delta^{202}Hg$ values for Hg^{II} are plotted as open

system Rayleigh fractionation models with $\ln(R/R_0)$ vs $\ln(f_r)$. The amount of Hg^{II} remaining in

the reactor was used to calculate f_r (see Methods). The 0h time point for chemotrophic live cells

replicate #1 was omitted from the regression fitted to the data for Hg^{II} (C) as this suggested an

alternative process was contributing to isotopic fractionation earlier on in the experiment.

Figure S4: Mass dependent fractionation of $^{202}Hg^0$ in phototrophically (A, B) and

chemotrophically (C, D) grown *H. modesticaldum*. $\delta^{202}Hg$ values for Hg^0 are plotted as open

system Rayleigh fractionation models with $\ln(R/R_0)$ vs $\ln(f_r)$. The amount of Hg^{II} remaining in

the reactor was used to calculate f_r (see Methods). The 3h time point for phototrophic cells in

replicates #1 and 2 were omitted from the regression fitted to the data for Hg^0 (A, B) as this

suggested an alternative process was contributing to isotopic fractionation earlier on in the

experiment. No significant regressions were obtained for chemotrophic cells and as such no

models are presented.

Table S1: Total Hg concentrations and Hg isotope ratios for the reactant pool Hg^{II} in

phototrophic and chemotrophic growth conditions experiments. Each line corresponds to an

individual sample and technical duplicates were taken for each time point. Std represents the

standard deviation from analytical replicates on the MC-ICP-MS. Abbreviations denote Metabo.

= Metabolism; Photo. = Phototrophy; Chemo. = Chemotrophy; Treat. = Treatment; Hg conc. =

Hg concentration, rep = replicate; Avg = average; Std = Standard deviation; ctl = control, MED

= medium. $\delta^{202}Hg$ values in **bold** denote those that deviate substantially from the composition of

the NIST 3133 standard in medium spiked controls.

Table S2: Total Hg concentrations and Hg isotope ratios for the Hg^0 pool in phototrophic and

chemotrophic growth conditions experiments. Each line corresponds to the Hg present on two

gold traps used to sample each time point in parallel. Std represents the standard deviation from

analytical replicates on the MC-ICP-MS. Abbreviations denote the following: Metabo. =

Metabolism; Photo. = Phototrophy; Chemo. = Chemotrophy; Treat. = Treatment; Hg conc. = Hg

concentration, rep = replicate; Avg = average; Std = Standard deviation.

Table S3: Summary of regression models for mass dependent fractionation of $^{202}Hg^{II}$ and $^{202}Hg^0$

in phototrophically and chemotrophically grown *H. modesticaldum* cultures. P-values for

regressions that displayed a significant (p<0.05) fit are shown in **bold.**

Table S4: Comparison of measured and calculated isotope fractionation values for the reactant

pool of Hg^{II} using a mass balance approach. Measured values of Hg concentration and $\delta^{202}Hg$ are

represented as an average of replicates from experiments presented in Tables S1 and S3.

Predicted values for media were obtained using the equations outlined for mass balance

calculations. Differences between the measured and predicted values that are substantially

greater than the analytical variability of measurements (2SD = 0.08) are shown in **bold**.

Abbreviations denote Photo. = phototrophic, Chemo. = chemotrophic, rep = replicates, Avg. =

average, Pred. = predicted.

Table S5: Comparison of enrichment factors and measures of variability used in this study and

the original cited work. Values in bold represent calculation ratios that differ between the

original article cited and the current study. Abbreviations: SD = standard deviation, SE =

standard error, p = product, r = reactant, Ox. = oxygenic, Anox. = anoxygenic, PS =

photosynthesis.