Developing Human Pluripotent Stem Cell-Based Cerebral Organoids with a Controllable Microglia Ratio for Modeling Brain Development and Pathology

Ranjie Xu[1, *], Andrew J. Boreland[1,5], Xiaoxi Li[2], Caroline Erickson[3], Mengmeng Jin[1], Colm Atkins[1], Zhiping Pang[4], Brian P. Daniels[1], Peng Jiang[1, *]

[1]Department of Cell Biology and Neuroscience, Rutgers University, Piscataway, NJ 08854, USA.
[2]Department of Immunology, Nanjing Medical University, Nanjing, China.
[3]Summer Undergraduate Research Program in Neuroscience (NeuroSURP), Rutgers Robert Wood Johnson Medical School, Piscataway, NJ 08854, USA
[4]Department of Neuroscience and Cell Biology and Child Health Institute of New Jersey, Rutgers Robert Wood Johnson Medical School, New Brunswick, NJ 08901, USA
[5]Graduate Program in Molecular Biosciences, Rutgers University, Piscataway, NJ 08854, USA.

*Correspondence: peng.jiang@rutgers.edu and ranjie.xu@rutgers.edu

Abstract

Microglia, as brain-resident macrophages, play critical roles in brain development, homeostasis, and disease. Microglia in animal models cannot accurately model the properties of human microglia due to notable transcriptomic and functional differences between human and other animal microglia. Efficient generation of microglia from human pluripotent stem cells (hPSCs) provides unprecedented opportunities to study the function and behavior of human microglia. Particularly, incorporating hPSCs-derived microglia into brain organoids facilitates their development in a 3-dimensional context, mimicking the brain environment. However, an optimized method that integrates an appropriate amount of microglia into brain organoids at a proper time point, resembling in vivo brain development, is still lacking. Here, we report the development of a new brain region-specific, microglia-containing organoid model by co-culturing hPSCs-derived primitive neural progenitor cells (pNPCs) and primitive macrophage progenitors (PMPs). In these organoids, hPSCs-derived pNPCs and PMPs interact with each other and develop into functional neurons, astroglia, and microglia, respectively. Importantly, the numbers of human microglia in the organoids can be controlled, resulting in a cell type ratio similar to that seen in the human brain. Using super-resolution microscopy, we demonstrate that these human microglia are able to phagocytize neural progenitor cells (NPCs) and dead cells, as well as to prune synapses at different developmental stages of the organoids. Furthermore, these human microglia respond to Zika virus infection of the organoids, as indicated by amoeboid-like morphology, increased expression of gene transcripts encoding inflammatory cytokines, and excessive pruning of synaptic materials. Together, our findings establish a new microglia-containing brain organoid model that will serve to study human microglial function in a variety of neurological disorders.

Introduction

Microglia, as the innate immune cells of the central nervous system (CNS), play critical roles in neurodevelopment and tissue homeostasis. During brain development, microglia phagocytize apoptotic cells, prune redundant synapses, and regulate neurogenesis and axonal growth (Casano and Peri, 2015; Wolf et al., 2017). Under homeostatic conditions, microglia survey the CNS environment, detect and respond to protein aggregates, pathogens, and other damage that may endanger the CNS and compromise tissue homeostasis (Bahmad et al., 2018; Casano and Peri, 2015). Increasing evidence has also demonstrated that microglial dysfunction is implicated in a broad range of neurological disorders, including schizophrenia, Alzheimer's disease, and traumatic brain injury (Wolf et al., 2017). Moreover, Genome-wide association studies show that many neurological disease risk genes are highly, and sometimes exclusively, expressed in microglia (Gosselin et al., 2017). Thus, there is a compelling incentive to understand the role of microglia in brain development and disease.

Most of our knowledge of microglia has relied on animal microglia, such as those derived from rodent or zebrafish (Pocock and Piers, 2018). However, human microglia are distinct from microglia in other animals, including differential expression of the complement cascade, phagocytic functions, and expression of susceptibility genes to neurodegeneration (Galatro et al., 2017; Geirsdottir et al., 2020; Gosselin et al., 2017). Because of these innate species differences, there is an urgent need to develop more representative strategies for investigating the unique functions of human microglia in health and disease. Recent advances in stem cell technologies have enabled the generation of human microglia from human pluripotent stem cells (hPSCs), greatly facilitating the study of the pathophysiology of human microglia (Douvaras et al., 2017; Haenseler et al., 2017; McQuade et al., 2018; Muffat et al., 2016).

Notably, previous studies demonstrated that the transcriptomic profile and morphology of hPSC-derived microglia cultured under 2-dimensinoal (2D) conditions are different than microglia in the *in vivo* brain. This is in part due to limited cell-cell and cell-matrix interactions in the 2D cultures(Gosselin et al., 2017; Jiang et al., 2020). Brain organoids derived from hPSCs in a 3D context contain diverse neural cell types, recapitulate embryonic brain architecture, and provide unprecedented opportunities to study human brain development and the pathogenesis of neurodevelopmental disorders (Di Lullo and Kriegstein, 2017; Qian et al., 2020). Thus, organoids may enhance human microglial survival and functional development, since human microglia in organoids likely are exposed to brain region specific signaling cues and interact with other cell types, including neurons and astrocytes. Previous studies have demonstrated the feasibility of generating microglia-containing organoids (Abud et al., 2017; Ormel et al., 2018). However, these models have significant limitations. Abud et al. reported integration of mature microglia into cerebral organoids that had been cultured for a long time period (Abud et al., 2017). During early embryonic development, PMPs originating from the yolk sac migrate to the CNS through the circulation (Ginhoux et al., 2010; Ginhoux and Guilliams, 2016). Those PMPs and NPCs interact with each other and undergo differentiation and maturation together *in vivo* (Li and Barres, 2018). As such, the model reported by Abud et al. is limited in examining the interactions between microglia and NPCs or developing neurons because microglia are introduced into the organoids at a stage when neurons have already developed. The timing of integrating microglia into organoids may impact neuronal population and maturation (Del Dosso et al., 2020). Recently, in order to promote mesoderm differentiation in brain organoids, Ormel et al. developed methods to generate organoids without using dual SMAD inhibition to induce neural differentiation (Ormel et al., 2018). Microglia innately develop in these cerebral organoids, simultaneously with neural differentiation. However, due to the absence of SMAD inhibition of direct cell differentiation, the heterogeneity of cell types in those organoids is high and uncontrollable.

In this study, we hypothesize that by co-culturing hPSCs-derived primitive neural progenitor cells (pNPCs) and PMPs, we can generate developmentally appropriate and brain region-specific microglia-containing brain organoids. The fact that the starting numbers of PMPs and pNPCs can be readily and precisely controlled further reduces the variability between individual organoids. Importantly, the numbers of human microglia in the organoids can be controlled, resulting in a cell type ratio similar to that seen in

the brain. Moreover, we demonstrate that in our microglia-containing organoids, human microglia exhibit phagocytic activity and synaptic pruning function. We also show that human microglia are dynamic in response to Zika virus (ZIKV) infection and find that ZIKV causes inflammation and excessive elimination of synaptic components by microglia.

Results

Generation of brain organoid with a controllable microglial ratio.
During embryonic CNS development, yolk sac-derived myeloid progenitors migrate into the developing neural tube, where both NPCs and myeloid progenitors interact, proliferate, and mature into neurons, astrocytes, oligodendrocytes, and microglia (Colonna and Butovsky, 2017). To mimic this early neurodevelopmental process, we designed a strategy to induce NPCs and PMPs from hPSCs first, and then co-culture NPCs and PMPs in 3D to generate microglia-containing brain organoids (Figure 1A). To induce neural differentiation, we used three hPSC lines, including two healthy individual-derived human induced pluripotent stem cells (hiPSCs) and one human embryonic stem cell (hESC), with dual inhibition of SMAD signaling(Chambers et al., 2009). At day 7, we plated the embryoid bodies (EB) on dishes, and on day 14, we manually separated neural rosettes from surrounding cells and further expanded as pNPCs, as described in our previous studies (Chen et al., 2016; Xu et al., 2019). The identity of hPSCs-derived pNPCs was confirmed by staining with Nestin, a marker for NPCs (Wiese et al., 2004) (Figure 1B). To induce microglia differentiation and obtain PMPs, we followed a published protocol (Haenseler et al., 2017; Xu et al., 2020). At day 5, yolk sac embryoid bodies (YS-EB) were plated on dishes, and PMPs were released from attached YS-EB into the supernatant starting around day 20; PMPs were continuously produced for more than three months, with a cumulative yield about 40-fold higher than the number of the input hPSCs. The identity of hPSCs-derived PMPs was confirmed by staining with CD235, a marker for YS primitive hematopoietic progenitors (Claes et al., 2018), and CD43, a marker for hematopoietic progenitor-like cells (Claes et al., 2018). Moreover, the PMPs were highly proliferative, as indicated by Ki67 staining (Figure 1C). Next, we cultured pNPCs and PMPs together and supplemented with mitogen FGF-2 for another two weeks (up to day 35), which is termed as the proliferation stage. At day 35, we transitioned organoids to neural differentiation medium, which contained growth factors, such as BDNF, GDNF to promote neural maturation, and IL-34, GM-CSF, which are important for microglia survival and maturation (Haenseler et al., 2017). We termed this stage as the neural differentiation stage. Organoids can be maintained in this stage for a long period for further characterization.

PMPs develop into microglia in brain organoids.
Microglia account for around 5-10% of total CNS cells (Frost and Schafer, 2016). To mimic the percentage of microglia in the CNS, we tested two different starting ratios of pNPCs and PMPs: 9:1 or 7: 3. We found that starting with 7000 pNPCs and 3000 PMPs generated organoids with around 8% microglia in the total cell population (Figure S1A and S1B). Also, organoids gradually increased in size during culture (Figure S1C). To confirm whether PMPs mature into microglia in the brain organoid, we examined the identity of microglia at different stages. At the proliferation stage (day 35), around 13% (13.7% ± 4.6) of cells in the organoid expressed PU.1, a transcription factor that is necessary for microglial differentiation (McKercher et al., 1996), and the majority of cells in the organoid were proliferative, as indicated by Ki67 staining (Figures 2A and 2G). Among all the cells, about 5% (4.6% ± 1.5) of cells expressed both PU.1 and Ki67, suggesting PMPs started to differentiate into microglia and were proliferative at this stage (Figure 2A). Next, we examined the identity of microglia at neural differentiation stage by using a panel of microglia markers. At day 45, organoids co-expressed Iba1, a canonical macrophage/microglia marker (Butovsky and Weiner, 2018), and CD45, which is expressed by all nucleated hematopoietic cells (Ford et al., 1995) (Figure 2B). At day 50, appropriate microglia morphology became increasingly evident. About 8% (8.0 % ± 1.6) of cells expressed Iba1, and most of these Iba1-expressing microglia showed ramified morphology (Figures 2C and 2H). Besides, some of the cells in day 55 organoids also co-expressed CD11b and CD45, as well as Iba1 and PU.1 (Figures 2D and 2E). We also found cells that expressed TMEM119 (Figure 2F), a marker that distinguishes microglia from other macrophages (Satoh et al., 2016). Taken together, these results demonstrate that PMPs differentiate into microglia in our brain organoids.

Neural maturation in microglia-containing brain organoid.

We then examined neural maturation in microglia-containing brain organoids. At day 40, ventricular zone-like regions were seen in organoids, which contained Ki67$^+$ and SOX2$^+$ progenitors, as well as TUJ-1$^+$ immature neurons (Figure 3A). At day 55, organoids had a large number of immature neurons, as indicated by the expression of Doublecortin (Dcx), a marker of immature neurons (Gleeson et al., 1999) (Figure 3B). As shown in Figure 2C and 2D, mature NeuN+ neurons and S100β$^+$ astrocytes were detected in day 55 and 75 organoids. To further confirm whether neurons were functionally mature in the microglia-containing organoids, we performed whole-cell patch-clamp recordings of neurons from acutely sectioned organoid slices (Figure 3E). Neurons recorded were simultaneously loaded with Biotin Ethylenediamine HBR for morphological representation and displayed complex arborization, long-distance projections, and spine-like structures (Figure 3F). We found that human neurons cultured in the 3D microglia-containing organoids were functionally mature, as indicated by these neurons robustly expressing voltage-gated K$^+$ and Na$^+$ currents (Figure 3G), firing spontaneous action potentials (Figure 3H) and repetitive evoked action potentials upon step-wise current injections (Figure 3I). More importantly, neurons in the organoids (90 days in culture) exhibit spontaneous post-synaptic currents (PSCs) (Figure 3J). Taken together, these results show that pNPCs differentiate into functional neurons and astrocytes in the presence of microglia in our organoids.

Microglia phagocytize NPCs and remove apoptotic cells in brain-region specific organoids.
As the resident macrophages of the CNS, microglia play vital roles in neurodevelopment and tissue homeostasis mediated in part by their phagocytic functions. Microglia phagocytize NPCs to control their relative abundance during development, clear extracellular debris and remove dead cells to maintain CNS homeostasis (Wolf et al., 2017). To investigate whether microglia have phagocytic functions in organoids, we examined the expression of CD68, a marker for the phagolysosome (Weinhard et al., 2018). We found the vast majority of microglia expressed CD68 (Figures 4A and 4B, and S2A), suggesting their active phagocytic functions. Next, we patterned the organoids into brain region-specific organoids, including the dorsal forebrain, as indicated by dorsal forebrain marker PAX6 (Sussel et al., 1999), or ventral forebrain, as indicated by ventral forebrain marker NKX2.1 (Sussel et al., 1999). As shown in Figures 4C and 4D, microglia phagocytized PAX6$^+$ and NKX2.1$^+$ NPCs on day 45. Additionally, CD68$^+$ microglia phagocytized developing neurons, as indicated by the intermediate progenitor stage marker Tbr2 (Sessa et al., 2010) (Figures 4E, S2B). To confirm whether microglia can remove apoptotic cells, we co-stained CD68 and activated caspase-3, an apoptotic cell marker, and found that microglia phagocytized the apoptotic cells (Figures 4F, S2C, S2D). Occasionally, we also found that microglia responded and aggregated around foreign bodies in the organoid caused by embedded fragments from a glass pipette tip (Figure S2E).

Microglia prune synapses in brain organoids.
As shown in Figure 5A, Iba1$^+$ microglia resided closely to MAP2$^+$ neuronal dendrites. In addition, we observed that synaptic puncta, labeled by PSD95, a post-synaptic marker, and synapsin I, a pre-synaptic marker, were distributed along the MAP2$^+$ dendrites (Figure 5B). Thus, we hypothesize that these human microglia are able to prune synapses in our organoids. We triple-stained Iba1, CD68, and PSD95 and employed a super-resolution imaging technique to visualize synapse engulfment by microglia. As shown in Figure 5C, PSD95$^+$ puncta were localized within CD68$^+$ phagolysosomes in Iba1$^+$ microglia, indicating that microglia phagocytized synaptic proteins in the day 55 organoid. Similarly, Synapsin I$^+$ puncta were found within CD68$^+$ phagolysosomes (Figure S3A). Next, we triple-stained CD68 with PSD95 and Synapsin I. The 3D reconstruction images clearly showed that PSD95$^+$ and synapsin I$^+$ puncta were colocalized within CD68$^+$ microglia, further indicating synaptic pruning function of the microglia in organoids (Figure 5D).

Microglia respond to Zika virus (ZIKV) infection in brain organoid.
We further explored whether our organoid model can be employed to study microglial functions under diseased or pathological conditions. Brain organoids have been used to model ZIKV infection before

(Dang et al., 2016; Garcez et al., 2016; Qian et al., 2016). However, microglia were missing in the previous organoid models and how human microglia are involved in ZIKV-related neuropathology is largely unknown. To address this question, we infected our microglia-containing organoids with two different stains of ZIKV, MR766 and Fortaleza. Consistent with previous studies, ZIKV-MR776 exhibited greater virulence than Fortaleza (Esser-Nobis et al., 2019), as indicated by the significant morphological abnormalities and cell detachment in the organoid (Figure 6A, S3B). Thus, we chose to use the MR766 strain for the following experiments. We measured viral replication using a standard plaque assay, which confirmed productive infection of the organoid (Figure 6B). Furthermore, flavivirus specific envelope (E)-protein, was found to be highly expressed in ZIKV-exposed organoids but not in mock-infected organoids, further indicating the success of ZIKV infection (Figure 6C). Interestingly, microglial morphology dramatically changed after virus exposure and became more amoeboid-like, suggesting microglia activation (Figure 6D). We then examined the expression of inflammatory cytokines IL-6, IL-1β, and TNF-α by qRT-PCR. Expression of these inflammatory cytokines was significantly increased in the ZIKV-exposed organoids, consistent with previous studies (Figure 6E). Type I interferon signaling is involved in response to viral and bacterial infections (Ivashkiv and Donlin, 2014). We examined the mRNA expression of INFAR1 and INFAR2 and found that both INFAR1 and INFAR2 mRNA expression increased after the ZIKV exposure (Figure 6F). These data confirm successful infection of the organoid, as well as the establishment of a robust innate immune response.

Abnormalities in synaptic pruning have been reported in many neurological disorders, brain injury, and viral infection (Garber et al., 2019; Gunner et al., 2019; Hong et al., 2016; Sellgren et al., 2019). To examine whether ZIKV exposure alters human microglial synaptic pruning function, we infected day 75 organoids, in which synapses have formed. As shown in Figure 6G and 6H, compared with mock-infected control, ZIKV exposure significantly increased PSD95[+] engulfment in CD68[+] microglial phagolysosome compartments, indicating increased microglial synaptic elimination following ZIKV infection. We further examined two receptors, complement receptor 3 (CR3), and the fractalkine receptor (CX3CR1), which mediate two major molecular pathways modulating microglia synaptic pruning in development and disease (Bialas and Stevens, 2013; Gunner et al., 2019; Hong et al., 2016). The qRT-PCR results showed that the expression of CR3, but not CX3CR1, increased after ZIKV exposure, suggesting CR3 might be involved in ZIKV-induced synaptic removal (Figure 6I). Taken together, these results show that microglia are able to respond to ZIKV exposure and exhibit increased synaptic elimination in organoids. Thus, this microglia-containing brain organoid model may provide a new platform to study human microglia function in ZIKV infection that was ignored previously due to the lack of this crucial cell type in the organoids.

Discussion

In this study, we develop a microglia-containing brain organoid model by co-culturing hPSC-derived pNPCs and PMPs at the onset of 3D organoid formation. In this organoid model, PMPs mature into functional microglia, as indicated by their ability to phagocytize NPCs and apoptotic cells as well as prune synapses. Meanwhile, NPCs differentiate into mature neurons that show electrophysiological activities in addition to their complex morphology. Importantly, using this organoid model, we find that human microglia excessively prune synapses under ZIKV infection.

Different from previous microglia-containing brain organoid models (Abud et al., 2017; Ormel et al., 2018), our model has several unique advantages. First, the ratio of microglia can be adjusted by controlling the starting number of pNPCs and PMPs. In the CNS, microglia account for around 5-10% of total CNS cells (Frost and Schafer, 2016). In our study, using a ratio of 7:3 for NPCs to PMPs, we generated organoids with around 8% microglia of the total cell population. Thus, this method can generate organoids with a consistent and physiologically relevant percentage of microglia. Second, since pNPCs are highly responsive to instructive neural patterning cues *in vitro* (Elkabetz et al., 2008; Li et al., 2011), these organoids can be assigned brain region identities at early stages of organoid culture by giving different patterning morphogens. This enables us to study microglia function in a brain region-dependent manner. In this study, we patterned organoid into dorsal or ventral forebrain identity and showed that microglia phagocytized dorsal or ventral forebrain NPCs in the corresponding organoids (Figures 4C and 4D). Third, co-culture of fate-committed ectoderm-derived pNPCs and mesoderm-derived PMPs at the onset of organoid formation better mimics early neurodevelopmental processes observed in *in vivo* brain development. This allows NPCs and PMPs to interact and mature reciprocally and significantly reduces possible contamination by other non-neural endoderm- or mesoderm-derived cells.

Our new microglia-containing brain organoid model provides unique opportunities to investigate human microglial function in development and disease, particularly with regard to microglial synaptic pruning function. During CNS development, microglia-mediated synaptic pruning plays a critical role in removing unwanted synapses and shaping neural circuits (Bialas and Stevens, 2013; Hong et al., 2016; Sakai, 2020). A previous study showed that human microglia engulfed post-synaptic materials in organoids, suggesting human microglia might prune synapses in organoids (Ormel et al., 2018). In this study, using super-resolution imaging techniques, we clearly demonstrated that synapses labeled by both pre-synaptic and post-synaptic markers were engulfed in microglia lysosomal compartments in the organoids (Figure 5D). Different from previous studies which reported either pre- or post-synaptic materials were engulfed by microglia(Ormel et al., 2018; Weinhard et al., 2018), our super-resolution imaging results reveal that human microglia appear to phagocytize the whole synapses. Moreover, we show that in addition to synaptic pruning, microglia phagocytize NPCs and remove dead cells in the organoid (Figures 4F), recapitulating microglial functions observed in the developing brain(Cunningham et al., 2013; Takahashi et al., 2005). Therefore, our microglia-containing organoid model enables us to study human microglial function in brain development, as well as neurodevelopmental disorders.

Previous studies using brain organoid models have shown that organoids are useful for studying ZIKV mediated pathology in humans. It was previously shown that ZIKV infection causes a reduction in human NPC pools and limits their proliferation in organoids (Dang et al., 2016; Qian et al., 2016). However, these organoids may not be able to fully model the disease mechanisms. For example, some receptors that are crucial for ZIKV infection are expressed by human astrocytes and microglia, but not human NPCs (Di Lullo and Kriegstein, 2017; Meertens et al., 2017; Qian et al., 2016). Thus, it is of particular importance to employ organoids containing diverse types of neural cells for studying ZIKV infection. Using our microglia-containing brain organoid model, we demonstrate that in response to ZIKV infection, human microglia target mature neurons and excessively remove synaptic materials. In addition to infecting the fetal brain, increasing evidence shows that ZIKV is also associated with a broad range of neurological disorders in adults (da Silva et al., 2017; Figueiredo et al., 2019). Notably, ZIKV can replicate in ex vivo adult human brain tissue(Figueiredo et al., 2019). In our study, after ZIKV infection, we found

that microglia adopted a more activated amoeboid-like morphology and expressed elevated levels of
inflammatory cytokines. Interestingly, human microglia excessively removed synaptic materials in our
model, consistent with previous findings in mice that mouse microglia engulf hippocampal presynaptic
terminals following ZIKV acute infection(Figueiredo et al., 2019; Garber et al., 2019). In addition, we also
examined two receptors, CX3CR1 and C3R, which are important for microglia-mediated synaptic pruning
during development and disease(Bialas and Stevens, 2013; Gunner et al., 2019; Hong et al., 2016).
Consistent with the previous findings that C3 expression was increased in the microglia after ZIKA
infection in the mouse brain(Figueiredo et al., 2019), we found increased expression of C3R after ZIKV
infection in the human brain organoid. Furthermore, we also found that expression of INFAR1/2, which
play critical roles in antiviral immune responses(Teijaro, 2016), were also upregulated in the brain
organoids. Thus, this microglia-containing organoid can be employed to study interactions between
human microglia and neurons in response to ZIKV infection and subsequent antiviral immune responses.
Increasing evidence indicates that adaptive immune cells play essential roles in brain
development and disease (Pasciuto et al., 2020; Prinz and Priller, 2017). Future studies incorporating
adaptive immune cells in brain organoid cultures will further present new opportunities for studying
interactions between human CNS and the adaptive immune system in development and neurological
disorders, including virus infection-associated brain dysfunctions such as ZIKV and SARS-CoV2
infections (Ellul et al., 2020; Iadecola et al., 2020). Overall, our new microglia-containing organoid model
will provide unique opportunities to investigate human microglial function in development and disease,
as well as provide a platform for drug screening.

Acknowledgements
This work was in part supported by grants from the NIH (R21HD091512 and R01NS102382 to P.J.). A.J.B is supported by the National Institute of General Medicine Sciences (NIGMS) NIH T32 GM008339.

Author Contributions
P.J. and R.X. designed experiments and interpreted data; R.X. carried out most of experiments with technical assistance from X.L., C.E., and M.J.; A.J.B. and Z.P. performed patch-clamp recordings. C.A, and B.P.D. performed ZIKV infection experiments. Z. P. and B.P.D. provided critical suggestions. P.J. and R.X wrote the manuscript with input from all co-authors.

Competing Financial Interests
The authors declare no competing financial interests.

Methods
Generation, culture, and quality control of hPSC lines.
Two healthy control hiPSC lines (female and male) and the female H9 ESC line were used in this study.
The hiPSC lines were generated from healthy adult-derived fibroblasts using the "Yamanaka"
reprogramming factors in our previous study (Chen et al., 2014). The hPSC line has been fully
characterized, including karyotyping, teratoma assay, Pluritest (www.PluriTest.org), DNA fingerprinting
STR (short tandem repeat) analysis and gene expression profiling(Xu et al., 2019; Xu et al., 2020). The
hPSCs, from passage number 15–30, were cultured on dishes coated with hESC-qualified Matrigel
(Corning) in mTeSR1 medium (STEM CELL Technologies) under a feeder-free condition and were
passaged once per week with ReLeSR medium (STEM CELL Technologies). All hPSC studies were
approved by the Embryonic Stem Cell Research Oversight committee at Rutgers University.
12
pNPC generation and culture.
Human pNPCs were generated from hPSCs as previously reported(Chen et al., 2016; Xu et al., 2019).
Briefly, to induce neural differentiation, embryoid bodies (EB) were treated with noggin (50 ng/ml,
Peprotech) and SB431542 (5 mM, Stemgent) for dual inhibition of SMAD signaling, in neural induction
medium consisting of DMEM/F12 (HyClone) and 1x N2 (Thermo Fisher Scientific) for 1 week. EBs were
then plated on dishes coated with growth factor-reduced Matrigel (BD Biosciences) in medium composed
of DMEM/F12, 1x N2, and laminin (1 mg/ml; Corning). pNPCs emerged in the form of neural rosettes for
another 7 days, after which, rosettes were manually isolated from surrounding cells and expanded for
another 7 days in pNPC medium, which is composed of a 1:1 mixture of Neurobasal (Thermo Fisher
Scientific) and DMEM/F12, supplemented with 1x N2, 1x B27-RA (Thermo Fisher Scientific), FGF2 (20
ng/ml, Peprotech), CHIR99021 (3 mM, Biogems), human leukemia inhibitory factor (hLIF, 10 ng/ml,
Millipore), SB431542 (2 mM), and ROCK inhibitor Y-27632 (10 mM, Tocris). Next, the expanded pNPCs
were dissociated into single cells using TrypLE Express (Thermo Fisher Scientific) for organoid
generation or long-term storage in liquid nitrogen.
27
PMP generation and culture.
The PMPs were derived from hPSCs, as previously reported(Haenseler et al., 2017; Xu et al., 2020), and
produced in a Myb-independent manner to recapitulate primitive hematopoiesis(Buchrieser et al., 2017;
Haenseler et al., 2017). Briefly, yolk sac embryoid bodies (YS-EBs) were generated by treating EBs with
bone morphogenetic protein 4 (BMP4, 50 ng/ml, Peprotech; to induce mesoderm), stem cell factor (SCF,
20 ng/ml, Peprotech), and vascular endothelial growth factor (VEGF, 50 ng/ml, Peprotech) in mTeSR1
medium (STEM CELL Technologies). Then, the YS-EBs were plated into dishes and cultured in PMP
medium, which is composed of X-VIVO 15 medium (Lonza), supplemented with macrophage colony-
stimulating factor (M-CSF,100 ng/ml, Peprotech) and interleukin-3 (IL-3, 25 ng/ml, Peprotech) to promote
myeloid differentiation. After 2–3 weeks of plating, human PMPs begin to emerge into the supernatant
and are continuously produced for around 3 months. The cumulative yield of PMPs can reach around 40-
fold of input hPSCs.
40
Brain organoid generation and culture.
To generate brain organoids, 10,000 cells were used per organoid similar to our previous studies (Kim et
al., 2019; Xu et al., 2019). In this study, 7,000 pNPCs plus 3,000 PMPs were co-cultured, supplemented
with ROCK inhibitor Y-27632 (10 mM), in ultra-low-attachment 96-well plates to form uniform organoids
in medium composed of a 1:1 mixture of PMP medium and NPC medium. NPC medium is composed of
a 1:1 mixture of Neurobasal (Thermo Fisher Scientific) and DMEM/F12, supplemented with 1x N2, 1x
B27-RA (Thermo Fisher Scientific), FGF2 (20 ng/ml, Peprotech) for three days (day 24). Then, organoids
were transferred to ultra-low-attachment 6-well plates and cultured with 1:1 mixture of PMP medium and
NPC medium for another 11 days (day 35). To pattern these organoids to the fate of dorsal or ventral
forebrain, we respectively culture the organoids without morphogen or treated organoids with sonic
hedgehog (SHH) (50 ng/mL, Peprotech) and purmorphamine (1 mM, Cayman Chem), an agonist of sonic

hedgehog signal pathway from day 21 to day 35(Figure 1A). The medium was replenished every day,
and the cell culture plates were kept on the orbital shaker with a speed of 80 rpm/min starting from day
27. Maximum 8 organoids were cultured in each well of the ultra-low-attachment 6-well plate. To promote
neural and microglial differentiation, organoids were cultured in differentiation medium, which contained
a 1:1 mixture of Neurobasal and DMEM/F12, supplemented with 1x N2 (Thermo Fisher Scientific), BDNF
(20 ng/ml, Peprotech), GDNF (20 ng/ml, Peprotech), dibutyryl-cyclic AMP (1mM, Sigma), ascorbic acid
(200 nM, Sigma), IL-34 (100 ng/ml, Peprotech), Granulocyte-macrophage colony-stimulating factor (GM-
CSF, 10 ng/ml, Peprotech) from day 35 onwards. Medium was replenished every other day. To further
promote neural maturation, organoids were cultured in media comprised of 1:1 mixture of the
aforementioned differentiation media and BrainPhys neuronal medium (STEMCELL Technologies).
These organoids were used for functional patch-clamp recordings.
12
Immunostaining and cell counting.
Immunostaining and cell counting were performed similarly to our previous report(Xu et al., 2019). Briefly,
organoids were fixed with 4% paraformaldehyde (PFA) for 1h at room temperature and then immersed
in 25% sucrose at 4 °C overnight for cryoprotection. Organoids were then embedded in OCT medium,
snap frozen in a dry-ice chilled methanol bath and sectioned in a cryostat (Leica) at 18-µm thickness
(unless otherwise specified). For immunofluorescence staining, the sections were washed with PBS
before permeabilization/blocking with 5% goat or donkey serum in PBS with 0.2% Triton X-100 at room
temperature (RT) for 1 h. The sections were incubated with primary antibodies diluted in the same
blocking solution at 4 °C overnight. All antibodies used are listed in Supplementary Table 1. Then, the
sections were washed with PBS and incubated with secondary antibodies for 1 h at RT. Finally, the slides
were washed with PBS and mounted with anti-fade Fluoromount-G medium containing 1, 40,6-diami-
dino-2-phenylindole dihydrochloride (DAPI) (Southern Biotechnology). A Zeiss 710 confocal microscope
was used for capturing images. Zeiss Airyscan super-resolution microscope at 63X with 0.2 mm z-steps
was used to acquire super-resolution images for visualizing synaptic puncta pruning. Imaris software
(Bitplane 9.5) was used to process super-resolution images and generate 3D-surface rendered images
as previously reported(Xu et al., 2020). To visualize the engulfed puncta within microglia, the mask
function of Imaris was used to subtract any fluorescence that was outside of the microglia, as previously
reported(Schafer et al., 2014). The cells were counted with ImageJ software, and at least three fields of
each organoid were chosen randomly for cell counting. To quantify PSD95 within the lysosome, the
volume of PSD95$^+$ synaptic puncta and CD68$^+$ phagosome was calculated by Imaris, and all data were
then normalized to the control, as previously reported(Gunner et al., 2019; Schafer et al., 2014). For the
visualization of neurons filled with Neurobiotin (Biotin Ethylenediamine HBR) during patch-clamp
recording, slices were immediately fixed for 30 min at RT in 4% PFA. Slices were washed with PBS,
permeabilized for 45 min in the same blocking solution above, and incubated with Alexa Fluor 546
streptavidin conjugate in blocking solution for 1 hr at RT. Finally, slices were washed with PBS and
mounted in Fluoromount-G medium for subsequent confocal imaging.
39
RNA isolation and qPCR
Total RNA was extracted from organoids using Trizol (Invitrogen), as previously reported(Xu et al., 2014).
Complementary DNA was prepared using a Superscript III First-Strand kit (Invitrogen). qPCR was
performed with Taqman primers (Life Technologies) on an ABI 7500 Real-Time PCR System. All primers
used are listed in Supplementary Table 2. All experimental samples were analyzed and normalized with
the expression level of housekeeping gene glyceraldehyde- 3-phosphate dehydrogenase (GAPDH).
Relative quantification was performed by applying the 2-DDCt method(Xu et al., 2019).
47
48 **Whole-cell patch-clamp recording in organoid slices**
49 Organoids were embedded in low melting temperature agarose (SeaPlaqueTM Low Melt Agarose) and
50 sectioned into 200-300 µm thick sections in ice-cold oxygenated (95% O_2/5% CO_2) ACSF containing the
51 following (in mM): 125 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 25 $NaHCO_3$, 2.5 glucose, 50 sucrose, 0.625 $CaCl_2$,

and 1.2 $MgCl_2$. The osmolarity of the cutting solution was 320-330 mOsm. The slices were then transferred to a recovery bath for 45 min at 32°C in oxygenated (95% O_2/5% CO_2) ACSF containing the following (in mM): 125 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 25 $NaHCO_3$, 2.5 glucose, 22.5 sucrose, 2.5 $CaCl_2$, and 1.2 $MgCl_2$. The osmolarity of the external solution was 305-315 mOsm. Electrophysiological recordings were performed in the same solution with a perfusion rate of 2 ml/min at room temperature. Both cutting and patching external ACSF solutions had a pH 7.2-7.4 when saturated with 95% O_2/5% CO_2. Neurons were identified by differential interference contrast (DIC) optics under an upright microscope (bx51, Olympus) with a 40x water-immersion objective. Patch pipettes were pulled from borosilicate glass with a resistance of 5-8 mΩ. For current-clamp and voltage-clamp recordings, a K-gluconate internal solution was used consisting of (in mM): 126 K-gluconate, 4 KCl, 10 HEPES, 0.05 EGTA, 4 ATP-magnesium, 0.3 GTP-sodium, and 10 phosphocreatine. The pH was adjusted to 7.2 with KOH and the osmolarity adjusted to 280-290 mOsm. Neurons were held at -70mV for voltage-clamp recordings. In combination with the electrophysiological recordings, we added 0.5% Biotin Ethylenediamine HBR (Biotum) to the internal solution in order to label patched neurons. Whole-cell patch-clamp recordings were performed using an Axon 700B amplifier. Data were filtered at 2 kHz, digitized at 10 kHz, and collected using Clampex 10.6 (Molecular Devices). Electrophysiological data were analyzed offline using Clampfit 10.6 software (Molecular Devices).

Zika virus infection.
Day 75 old organoids were infected with ZIKV. Two different strains were tested initially, including MR766 (Uganda) and Fortaleza (Brazil),, and MR766 was used for most experiments. Briefly, organoids were infected with ZIKA or mock, with a concentration of 10^4 PFU/ml in 6 well plates on the orbital shaker. After three days of infection, organoids were replenished with fresh medium without ZIKV and cultured for another four days. At day three or day seven post-infection, organoids were collected for immunohistochemistry or RNA extraction, and the supernatant was collected for plaque assay. Briefly, Vero CCL-81 cells (ATCC) were plated at 2E5 cells/well in 12-well plates. Supernatant from infected organoids was serially diluted 10-fold, and 100μL transferred to each well. Plates were rocked every 15 minutes for 1 hour before a semi-solid overly containing 0.5% Methylcellulose in viral growth media (VWR) was applied. 4 days post infection, the overlay was removed, and cells fixed in a solution of 10% neutral buffered formalin with 0.25% crystal violet for plaque counting.

Statistics and reproducibility. All data represent mean ± s.e.m. When only two independent groups were compared, significance was determined by two-tailed unpaired t test with Welch's correction. A p-value <0.05 was considered significant. The analyses were done in GraphPad Prism v.5. All experiments were independently performed at least three times with similar results.

References:

Abud, E.M., Ramirez, R.N., Martinez, E.S., Healy, L.M., Nguyen, C.H.H., Newman, S.A., Yeromin, A.V., Scarfone, V.M., Marsh, S.E., Fimbres, C.*, et al.* (2017). iPSC-Derived Human Microglia-like Cells to Study Neurological Diseases. Neuron *94*, 278-293 e279.

Bahmad, H.F., Cheaito, K., Chalhoub, R.M., Hadadeh, O., Monzer, A., Ballout, F., El-Hajj, A., Mukherji, D., Liu, Y.-N., Daoud, G.*, et al.* (2018). Sphere-Formation Assay: Three-Dimensional in vitro Culturing of Prostate Cancer Stem/Progenitor Sphere-Forming Cells. Frontiers in Oncology *8*, 347.

Bialas, A.R., and Stevens, B. (2013). TGF-beta signaling regulates neuronal C1q expression and developmental synaptic refinement. Nat Neurosci *16*, 1773-1782.

Buchrieser, J., James, W., and Moore, M.D. (2017). Human Induced Pluripotent Stem Cell-Derived Macrophages Share Ontogeny with MYB-Independent Tissue-Resident Macrophages. Stem Cell Reports *8*, 334-345.

Butovsky, O., and Weiner, H.L. (2018). Microglial signatures and their role in health and disease. Nat Rev Neurosci *19*, 622-635.

Casano, A.M., and Peri, F. (2015). Microglia: multitasking specialists of the brain. Dev Cell *32*, 469-477.

Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., and Studer, L. (2009). Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. Nat Biotechnol *27*, 275-280.

Chen, C., Jiang, P., Xue, H., Peterson, S.E., Tran, H.T., McCann, A.E., Parast, M.M., Li, S., Pleasure, D.E., Laurent, L.C.*, et al.* (2014). Role of astroglia in Down's syndrome revealed by patient-derived human-induced pluripotent stem cells. Nat Commun *5*, 4430.

Chen, C., Kim, W.Y., and Jiang, P. (2016). Humanized neuronal chimeric mouse brain generated by neonatally engrafted human iPSC-derived primitive neural progenitor cells. JCI Insight *1*, e88632.

Claes, C., Van den Daele, J., and Verfaillie, C.M. (2018). Generating tissue-resident macrophages from pluripotent stem cells: Lessons learned from microglia. Cell Immunol *330*, 60-67.

Colonna, M., and Butovsky, O. (2017). Microglia Function in the Central Nervous System During Health and Neurodegeneration. Annu Rev Immunol *35*, 441-468.

Cunningham, C.L., Martinez-Cerdeno, V., and Noctor, S.C. (2013). Microglia regulate the number of neural precursor cells in the developing cerebral cortex. J Neurosci *33*, 4216-4233.

da Silva, I.R.F., Frontera, J.A., Bispo de Filippis, A.M., Nascimento, O., and Group, R.-G.-Z.R. (2017). Neurologic Complications Associated With the Zika Virus in Brazilian Adults. JAMA Neurol *74*, 1190-1198.

Dang, J., Tiwari, S.K., Lichinchi, G., Qin, Y., Patil, V.S., Eroshkin, A.M., and Rana, T.M. (2016). Zika Virus Depletes Neural Progenitors in Human Cerebral Organoids through Activation of the Innate Immune Receptor TLR3. Cell Stem Cell *19*, 258-265.

Del Dosso, A., Urenda, J.P., Nguyen, T., and Quadrato, G. (2020). Upgrading the Physiological Relevance of Human Brain Organoids. Neuron *107*, 1014-1028.

Di Lullo, E., and Kriegstein, A.R. (2017). The use of brain organoids to investigate neural development and disease. Nat Rev Neurosci *18*, 573-584.

Douvaras, P., Sun, B., Wang, M., Kruglikov, I., Lallos, G., Zimmer, M., Terrenoire, C., Zhang, B., Gandy, S., Schadt, E.*, et al.* (2017). Directed Differentiation of Human Pluripotent Stem Cells to Microglia. Stem cell reports *8*, 1516-1524.

Elkabetz, Y., Panagiotakos, G., Al Shamy, G., Socci, N.D., Tabar, V., and Studer, L. (2008). Human ES cell-derived neural rosettes reveal a functionally distinct early neural stem cell stage. Genes Dev *22*, 152-165.

Ellul, M.A., Benjamin, L., Singh, B., Lant, S., Michael, B.D., Easton, A., Kneen, R., Defres, S., Sejvar, J., and Solomon, T. (2020). Neurological associations of COVID-19. Lancet Neurol *19*, 767-783.

Esser-Nobis, K., Aarreberg, L.D., Roby, J.A., Fairgrieve, M.R., Green, R., and Gale, M., Jr. (2019). Comparative Analysis of African and Asian Lineage-Derived Zika Virus Strains Reveals Differences in Activation of and Sensitivity to Antiviral Innate Immunity. J Virol *93*.

Figueiredo, C.P., Barros-Aragao, F.G.Q., Neris, R.L.S., Frost, P.S., Soares, C., Souza, I.N.O., Zeidler, J.D., Zamberlan, D.C., de Sousa, V.L., Souza, A.S.*, et al.* (2019). Zika virus replicates in adult human brain tissue and impairs synapses and memory in mice. Nat Commun *10*, 3890.

Ford, A.L., Goodsall, A.L., Hickey, W.F., and Sedgwick, J.D. (1995). Normal adult ramified microglia separated from other central nervous system macrophages by flow cytometric sorting. Phenotypic differences defined and direct ex vivo antigen presentation to myelin basic protein-reactive CD4+ T cells compared. J Immunol *154*, 4309-4321.

Frost, J.L., and Schafer, D.P. (2016). Microglia: Architects of the Developing Nervous System. Trends Cell Biol *26*, 587-597.

Galatro, T.F., Holtman, I.R., Lerario, A.M., Vainchtein, I.D., Brouwer, N., Sola, P.R., Veras, M.M., Pereira, T.F., Leite, R.E.P., Moller, T.*, et al.* (2017). Transcriptomic analysis of purified human cortical microglia reveals age-associated changes. Nat Neurosci *20*, 1162-1171.

Garber, C., Soung, A., Vollmer, L.L., Kanmogne, M., Last, A., Brown, J., and Klein, R.S. (2019). T cells promote microglia-mediated synaptic elimination and cognitive dysfunction during recovery from neuropathogenic flaviviruses. Nat Neurosci *22*, 1276-1288.

Garcez, P.P., Loiola, E.C., Madeiro da Costa, R., Higa, L.M., Trindade, P., Delvecchio, R., Nascimento, J.M., Brindeiro, R., Tanuri, A., and Rehen, S.K. (2016). Zika virus impairs growth in human neurospheres and brain organoids. Science *352*, 816-818.

Geirsdottir, L., David, E., Keren-Shaul, H., Weiner, A., Bohlen, S.C., Neuber, J., Balic, A., Giladi, A., Sheban, F., Dutertre, C.A.*, et al.* (2020). Cross-Species Single-Cell Analysis Reveals Divergence of the Primate Microglia Program. Cell *181*, 746.

Ginhoux, F., Greter, M., Leboeuf, M., Nandi, S., See, P., Gokhan, S., Mehler, M.F., Conway, S.J., Ng, L.G., Stanley, E.R.*, et al.* (2010). Fate mapping analysis reveals that adult microglia derive from primitive macrophages. Science *330*, 841-845.

Ginhoux, F., and Guilliams, M. (2016). Tissue-Resident Macrophage Ontogeny and Homeostasis. Immunity *44*, 439-449.

Gleeson, J.G., Lin, P.T., Flanagan, L.A., and Walsh, C.A. (1999). Doublecortin Is a Microtubule-Associated Protein and Is Expressed Widely by Migrating Neurons. Neuron *23*, 257-271.

Gosselin, D., Skola, D., Coufal, N.G., Holtman, I.R., Schlachetzki, J.C.M., Sajti, E., Jaeger, B.N., O'Connor, C., Fitzpatrick, C., Pasillas, M.P.*, et al.* (2017). An environment-dependent transcriptional network specifies human microglia identity. Science *356*.

Gunner, G., Cheadle, L., Johnson, K.M., Ayata, P., Badimon, A., Mondo, E., Nagy, M.A., Liu, L., Bemiller, S.M., Kim, K.W.*, et al.* (2019). Sensory lesioning induces microglial synapse elimination via ADAM10 and fractalkine signaling. Nat Neurosci *22*, 1075-1088.

Haenseler, W., Sansom, S.N., Buchrieser, J., Newey, S.E., Moore, C.S., Nicholls, F.J., Chintawar, S., Schnell, C., Antel, J.P., Allen, N.D.*, et al.* (2017). A Highly Efficient Human Pluripotent Stem Cell Microglia Model Displays a Neuronal-Co-culture-Specific Expression Profile and Inflammatory Response. Stem Cell Reports *8*, 1727-1742.

Hong, S., Beja-Glasser, V.F., Nfonoyim, B.M., Frouin, A., Li, S., Ramakrishnan, S., Merry, K.M., Shi, Q., Rosenthal, A., Barres, B.A.*, et al.* (2016). Complement and microglia mediate early synapse loss in Alzheimer mouse models. Science *352*, 712-716.

Iadecola, C., Anrather, J., and Kamel, H. (2020). Effects of COVID-19 on the Nervous System. Cell.

Ivashkiv, L.B., and Donlin, L.T. (2014). Regulation of type I interferon responses. Nat Rev Immunol
14, 36-49.

Jiang, P., Turkalj, L., and Xu, R. (2020). High-Fidelity Modeling of Human Microglia with Pluripotent
Stem Cells. Cell Stem Cell *26*, 629-631.

Kim, H., Xu, R., Padmashri, R., Dunaevsky, A., Liu, Y., Dreyfus, C.F., and Jiang, P. (2019). Pluripotent
Stem Cell-Derived Cerebral Organoids Reveal Human Oligodendrogenesis with Dorsal and Ventral
Origins. Stem cell reports *12*, 890-905.

Li, Q., and Barres, B.A. (2018). Microglia and macrophages in brain homeostasis and disease. Nat
Rev Immunol *18*, 225-242.

Li, W., Sun, W., Zhang, Y., Wei, W., Ambasudhan, R., Xia, P., Talantova, M., Lin, T., Kim, J., Wang, X.*, et*
al. (2011). Rapid induction and long-term self-renewal of primitive neural precursors from human
embryonic stem cells by small molecule inhibitors. Proc Natl Acad Sci U S A *108*, 8299-8304.

McKercher, S.R., Torbett, B.E., Anderson, K.L., Henkel, G.W., Vestal, D.J., Baribault, H., Klemsz, M.,
Feeney, A.J., Wu, G.E., Paige, C.J.*, et al.* (1996). Targeted disruption of the PU.1 gene results in multiple
hematopoietic abnormalities. EMBO J *15*, 5647-5658.

McQuade, A., Coburn, M., Tu, C.H., Hasselmann, J., Davtyan, H., and Blurton-Jones, M. (2018).
Development and validation of a simplified method to generate human microglia from pluripotent
stem cells. Mol Neurodegener *13*, 67.

Meertens, L., Labeau, A., Dejarnac, O., Cipriani, S., Sinigaglia, L., Bonnet-Madin, L., Le Charpentier, T.,
Hafirassou, M.L., Zamborlini, A., Cao-Lormeau, V.M.*, et al.* (2017). Axl Mediates ZIKA Virus Entry in
Human Glial Cells and Modulates Innate Immune Responses. Cell Rep *18*, 324-333.

Muffat, J., Li, Y., Yuan, B., Mitalipova, M., Omer, A., Corcoran, S., Bakiasi, G., Tsai, L.H., Aubourg, P.,
Ransohoff, R.M.*, et al.* (2016). Efficient derivation of microglia-like cells from human pluripotent
stem cells. Nat Med *22*, 1358-1367.

Ormel, P.R., Vieira de Sa, R., van Bodegraven, E.J., Karst, H., Harschnitz, O., Sneeboer, M.A.M.,
Johansen, L.E., van Dijk, R.E., Scheefhals, N., Berdenis van Berlekom, A.*, et al.* (2018). Microglia
innately develop within cerebral organoids. Nat Commun *9*, 4167.

Pasciuto, E., Burton, O.T., Roca, C.P., Lagou, V., Rajan, W.D., Theys, T., Mancuso, R., Tito, R.Y., Kouser,
L., Callaerts-Vegh, Z.*, et al.* (2020). Microglia Require CD4 T Cells to Complete the Fetal-to-Adult
Transition. Cell *182*, 625-640 e624.

Pocock, J.M., and Piers, T.M. (2018). Modelling microglial function with induced pluripotent stem
cells: an update. Nat Rev Neurosci *19*, 445-452.

Prinz, M., and Priller, J. (2017). The role of peripheral immune cells in the CNS in steady state and
disease. Nat Neurosci *20*, 136-144.

Qian, X., Nguyen, H.N., Song, M.M., Hadiono, C., Ogden, S.C., Hammack, C., Yao, B., Hamersky, G.R.,
Jacob, F., Zhong, C.*, et al.* (2016). Brain-Region-Specific Organoids Using Mini-bioreactors for
Modeling ZIKV Exposure. Cell *165*, 1238-1254.

Qian, X., Su, Y., Adam, C.D., Deutschmann, A.U., Pather, S.R., Goldberg, E.M., Su, K., Li, S., Lu, L., Jacob,
F.*, et al.* (2020). Sliced Human Cortical Organoids for Modeling Distinct Cortical Layer Formation.
Cell Stem Cell *26*, 766-781 e769.

Sakai, J. (2020). Core Concept: How synaptic pruning shapes neural wiring during development and,
possibly, in disease. Proc Natl Acad Sci U S A *117*, 16096-16099.

Satoh, J.-i., Kino, Y., Asahina, N., Takitani, M., Miyoshi, J., Ishida, T., and Saito, Y. (2016). TMEM119
marks a subset of microglia in the human brain. Neuropathology *36*, 39-49.

Schafer, D.P., Lehrman, E.K., Heller, C.T., and Stevens, B. (2014). An engulfment assay: a protocol to assess interactions between CNS phagocytes and neurons. J Vis Exp.

Sellgren, C.M., Gracias, J., Watmuff, B., Biag, J.D., Thanos, J.M., Whittredge, P.B., Fu, T., Worringer, K., Brown, H.E., Wang, J., *et al.* (2019). Increased synapse elimination by microglia in schizophrenia patient-derived models of synaptic pruning. Nat Neurosci *22*, 374-385.

Sessa, A., Mao, C.A., Colasante, G., Nini, A., Klein, W.H., and Broccoli, V. (2010). Tbr2-positive intermediate (basal) neuronal progenitors safeguard cerebral cortex expansion by controlling amplification of pallial glutamatergic neurons and attraction of subpallial GABAergic interneurons. Genes Dev *24*, 1816-1826.

Sussel, L., Marin, O., Kimura, S., and Rubenstein, J.L. (1999). Loss of Nkx2.1 homeobox gene function results in a ventral to dorsal molecular respecification within the basal telencephalon: evidence for a transformation of the pallidum into the striatum. Development *126*, 3359-3370.

Takahashi, K., Rochford, C.D., and Neumann, H. (2005). Clearance of apoptotic neurons without inflammation by microglial triggering receptor expressed on myeloid cells-2. J Exp Med *201*, 647-657.

Teijaro, J.R. (2016). Pleiotropic Roles of Type 1 Interferons in Antiviral Immune Responses. Adv Immunol *132*, 135-158.

Weinhard, L., di Bartolomei, G., Bolasco, G., Machado, P., Schieber, N.L., Neniskyte, U., Exiga, M., Vadisiute, A., Raggioli, A., Schertel, A., *et al.* (2018). Microglia remodel synapses by presynaptic trogocytosis and spine head filopodia induction. Nat Commun *9*, 1228.

Wiese, C., Rolletschek, A., Kania, G., Blyszczuk, P., Tarasov, K.V., Tarasova, Y., Wersto, R.P., Boheler, K.R., and Wobus, A.M. (2004). Nestin expression – a property of multi-lineage progenitor cells? Cellular and Molecular Life Sciences CMLS *61*, 2510-2522.

Wolf, S.A., Boddeke, H.W., and Kettenmann, H. (2017). Microglia in Physiology and Disease. Annu Rev Physiol *79*, 619-643.

Xu, R., Brawner, A.T., Li, S., Liu, J.J., Kim, H., Xue, H., Pang, Z.P., Kim, W.Y., Hart, R.P., Liu, Y., *et al.* (2019). OLIG2 Drives Abnormal Neurodevelopmental Phenotypes in Human iPSC-Based Organoid and Chimeric Mouse Models of Down Syndrome. Cell Stem Cell *24*, 908-926 e908.

Xu, R., Hu, Q., Ma, Q., Liu, C., and Wang, G. (2014). The protease Omi regulates mitochondrial biogenesis through the GSK3beta/PGC-1alpha pathway. Cell Death Dis *5*, e1373.

Xu, R., Li, X., Boreland, A.J., Posyton, A., Kwan, K., Hart, R.P., and Jiang, P. (2020). Human iPSC-derived mature microglia retain their identity and functionally integrate in the chimeric mouse brain. Nat Commun *11*, 1577.

Figure legends:

Figure 1. Generation of microglia-containing brain organoids.
(A) A schematic procedure for deriving microglia-containing brain organoids by co-culture of primitive neural progenitor cells (pNPCs) and primitive macrophage progenitors (PMPs) under 3-dimensional (3D) conditions. Scale bars: 100 µm or 10 µm in the original or enlarged images, respectively.
(B) Representatives of Nestin$^+$ pNPC. Scale bars: 20 µm in the original or enlarged images.
(C) Representatives of CD235$^+$, CD43$^+$, and Ki67$^+$ PMPs. Scale bars: 20 µm in the original or enlarged images.

Figure 2. Differentiation of PMPs into microglia in brain organoids.
(A) Representatives of PU.1$^+$/Ki67$^+$ cells in day 35 brain organoids. Scale bars: 20 µm.
(B) Representatives of Iba1$^+$/CD45$^+$ cells in day 45 brain organoids. Scale bars: 20 µm.
(C) Representatives of Iba1$^+$ cells in day 50 brain organoids. Scale bars: 50 µm or 10 µm in the original or enlarged images, respectively.
(D) Representatives of CD11b$^+$/CD45$^+$ cells in day 55 brain organoids. Scale bars: 50 µm or 20 µm in the original or enlarged images, respectively.
(E) Representatives of Iba1$^+$/PU.1$^+$ cells in day 55 brain organoids. Scale bars: 50 µm or 20 µm in the original or enlarged images, respectively.
(F) Representatives of TMEM119$^+$ cells in day 55 brain organoids. Scale bars: 20 µm.
(G) Quantification of pooled data from two hiPSC lines and one hESC line showing the percentage of PU.1$^+$, and PU.1$^+$/Ki67$^+$ cells in day 35 organoids (n = 3). Three independent experiments were performed and for each experiment, 4-6 organoids from each hPSC line were used. Data are presented as mean ± s.e.m.
(H) Quantification of pooled data from hiPSC and hESC lines showing the percentage of Iba1$^+$ cells in day 50 organoids (n = 3). Three independent experiments were performed, and for each experiment, 4-6 organoids from each hPSC line were used. Data are presented as mean ± s.e.m.

Figure 3. Neural maturation in microglia-containing brain organoid.
(A) Representatives of Ki67$^+$, SOX2$^+$, and Tuj1$^+$ cells in day 40 microglia-containing brain organoids. Scale bars: 50 µm or 20 µm as indicated.
(B) Representatives of DCX$^+$ cells in day 45 microglia-containing brain organoid. Scale bars: 20 µm.
(C) Representatives of NeuN$^+$ and S100β$^+$ cells in day 55 or 75 brain organoids with or without microglia. Scale bars: 20 µm.
(D) Quantification of pooled data from two hiPSC lines and one hESC line showing the diameter of day 35, 55 or 75 brain organoids with or without microglia. (n = 3), Three independent experiments were performed and for each experiment, 4-6 organoids from each hPSC line were used. Student's *t* test. *NS* represents no significance. Data are presented as mean ± s.e.m.
(E) A representative image showing a patched neuron in acute organoid slice.
(F) Sample images of a neuron filled with Biotin Ethylenediamine HBR from a day 92 microglia-containing organoid after whole-cell recording. Complex arborization (F1) and spine-like structures (F2, F3) were identified. The images are inverted to a white background to help visualize neuron morphology.
(G) Representative traces of whole-cell currents recorded from neurons in acutely sectioned organoid (90 day old in culture) slices. Insert: fast activation/inactivation voltage dependent sodium currents.
(H) Representative traces of spontaneous action potentials recorded from neurons in organoid (90 day old in culture) slices.
(I) Example of repetitive action potentials evoked by stepwise currents injections.
(J) Representative traces of spontaneous postsynaptic currents (PSCs) recorded from neurons in sliced collected from a day 92 organoid.

Figure 4. Phagocytic function of microglia in brain-region specific organoids.

(A) Representatives of Iba1$^+$/CD68$^+$ cells in day 50 brain organoids. Scale bars: 20 μm.
(B) Quantification of pooled data from two hiPSC lines and one hESC line showing the percentage of
CD68$^+$ cells in total Iba1$^+$ cells in day 50 organoids (n = 3, three lines each time, 4-6 organoids each line
each time). Data are presented as mean ± s.e.m.
(C) Representative images showing CD68$^+$ microglia phagocytosing PAX6$^+$ dorsal forebrain NPCs in day
46 dorsal forebrain organoids (enlarged images). Scale bars: 50 μm or 10 μm in the original or enlarged
images, respectively.
(D) Representative images showing CD68$^+$ microglia phagocytosing NKX2.1$^+$ ventral forebrain NPCs in
day 46 ventral forebrain organoids (enlarged images). Scale bars: 10 μm or 5 μm in the original or
enlarged images, respectively.
(E) Representative images showing CD68$^+$ microglia phagocytosing Tbr2$^+$ intermediate progenitor cells
in day 56 organoids (enlarged images). Scale bars: 10 μm or 5 μm in the original or enlarged images,
respectively.
(F) Representative images showing CD68$^+$ microglia phagocytosing activated caspase3$^+$ apoptotic cells
in day 56 organoids (enlarged images). Scale bars: 5 μm.
16
Figure 5. Microglia prune synapses in microglia-containing brain organoids.
(A) Representative images showing Iba1$^+$ microglia interact with MAP2$^+$ neuronal processes in day 55
brain organoids. Scale bars: 50 μm or 20 μm in the original or enlarged images, respectively.
(B) Representative images showing PSD95$^+$ and synapsin 1$^+$ synaptic puncta along MAP2$^+$ processes in
day 55 brain organoids. Scale bars: 20 μm or 5 μm in the original or enlarged images, respectively.
(C) Representative images showing colocalization of Iba1$^+$, CD68$^+$, and PSD95$^+$ staining in day 55
organoids. Left, a raw fluorescent super-resolution image. Right, a 3D-surface rendered image from the
same raw image. Scale bars, 3 μm.
(D) Representative 3D reconstruction of super-resolution images showing CD68$^+$ microglial engulfment
of synapsin I and PSD95 in day 55 organoids. Scale bars, 3 and 1 μm in the original or enlarged images,
respectively.
28
Figure 6. Responses of microglia to ZIKV infection in brain organoids.
(A) Representative phase-contrast images showing the morphology of day 75 microglia-containing
organoid after three and seven days of ZIKV or mock infection. Scale bars: 200 μm.
(B) Quantitation of infectious ZIKV particles in the culture medium by plaque assay.
(C) Representative images showing E protein expression in day 75 organoids after three days of ZIKV
or mock infection. Scale bars: 200 μm.
(D) Representatives of Iba1$^+$/CD68$^+$ microglia in day 75 organoids after three days of ZIKV or mock
infection. Scale bars: 50 μm or 20 μm in the original or enlarged images, respectively.
(E) qPCR analysis of IL-6, IL-β, and TNF-α mRNA expression in day 75 organoid after three days of ZIKV
or mock infection.
(F) qPCR analysis of IFNAR1 and IFNAR2 mRNA expression in day 75 organoid after three days of ZIKV
or mock infection.
(G) Representative 3D reconstruction of super-resolution images showing PSD95$^+$ puncta within CD68$^+$
microglial lysosomes in day 75 organoids after three days of ZIKV or mock infection. Scale bars, 5 and 2
μm in the original or enlarged images, respectively.
(H) Quantification of PSD95$^+$ puncta engulfment in CD68$^+$ microglial lysosomes in day 75 organoids after
three days of ZIKV or mock infection. Notably, larger volumes of PSD95$^+$ puncta were seen inside
microglia in response to ZIKV, as compared to mock infection group. (n = 3, three independent times
infections, each time, 9-12 organoids). Data normalized to engulfment in the mock control. Student's t
test. ** represents ***$P < 0.001$. Data are presented as mean ± s.e.m.
(I) qPCR analysis of CX3CR1 and CR3 mRNA expression in day 75 organoid after three days of ZIKV or
mock infection.
51

Supplementary Figure 1. Characterization of microglia-containing organoid.
(A) Representatives of CD11b$^+$ cells in day 50 brain organoids with different starting numbers of PMPs and pNPCs. Scale bars: 20 µm.
(B) Quantification of pooled data from an hiPSC and an hESC line showing the percentage of CD11b$^+$ cells in day 50 organoids with different starting numbers of PMPs and pNPCs (n = 3). Three independent experiments were performed, and for each experiment, 4-6 organoids from each hPSC line were used. Data are presented as mean ± s.e.m.
(C) Quantification of organoid size at different stages. Scale bars: 200 µm. Data are presented as mean ± s.e.m.

Supplementary Figure 2. Phagocytic function of microglia in brain organoids.
(A) Representative of CD68$^+$ cells in day 45 organoids. Scale bars: 50 µm.
(B) Representative images showing CD68$^+$ microglia phagocytosing Tbr2$^+$ intermediate progenitor cells in day 55 organoids. The enlarged images. Scale bars: 20 µm or 5 µm in the original or enlarged images, respectively.
(C and D) Representative images showing CD68$^+$ microglia phagocytosing activated-caspase-3$^+$ cell in day 55 organoids. Scale bars: 20 µm.
(E) Representative images showing Iba1$^+$ and PU.1$^+$ microglia accumulate around foreign body glass fragments from pipette tip in day 55 organoids. Scale bars: 20 µm.

Supplementary Figure 3. Microglia prune pre-synaptic component and respond to ZIKV--Fortaleza infection. (A) Representative 3D reconstruction of super-resolution images showing synapsin I$^+$ puncta within CD68$^+$ microglial lysosomes in day 55 organoids. Scale bars, 3 µm.
(B) Representative phase-contrast images showing the morphology of day 75 organoid at 3- and 7-day day post Fortaleza ZIKV-Fortaleza infection. Scale bars: 500 µm.

Supplementary Table 1. A list of antibodies used.

Antibodies	Vendor/Catalog #.	Type	Dilution
Caspase 3 activated	Sigma / C8487	Rabbit IgG	1:1000
CD235	Thermo Fisher Scientific / PA5-27154	Rabbit IgG	1:100
CD43	Thermo Fisher Scientific / 14-0439-82	Mouse IgG	1:25
CD45	Thermo Fisher Scientific / 14-9457-82	Mouse IgG	1:40
CD68	Invitrogen / MA5-13324	Mouse IgG	1:100
Dcx	Cell Signaling / 4604s	Rabbit IgG	1:500
TuJ-1	Millipore / MAB1637	Mouse IgG	1:200
flavivirus E-protein	Millipore / MAB10216	Mouse IgG	1:500
Ki67	Thermo Fisher Scientific / SP6	Rabbit IgG	1:200
MAP2	Millipore / AB3418	Mouse IgG	1:500
NeuN	Millipore / ABN78	Rabbit IgG	1:500
Nestin	Santa Cruz / SC-21249	Goat IgG	1:100
NKX2.1(TTF1)	Abcam / ab76013	Rabbit IgG	1:200
PAX6	GeneTex / GTX11324	Rabbit IgG	1:400
PSD95	Invitrogen 51-6900	Rabbit IgG	1:100
PSD95	Abcam / ab12093	Goat IgG	1:400
S100B	Sigma / S2532	Mouse IgG	1:1000
SOX2	Millipore / AB5603	Rabbit IgG	1:100
Synapsin I	Millipore / AB1543P	Rabbit IgG	1:400
Tbr2	Abcam / AB23345	Rabbit IgG	1:100
TMEM119	Thermo Fisher Scientific / PA-62505	Rabbit IgG	1:400

Supplementary Table 2. Taqman Primers.

Gene	Gene expression assay catalog number
IL-6	Hs00174131_m1
IL-β	Hs01555410_m1
TNF-α	Hs00174128_m1
IFNAR1	Hs01066118_m1
INFAR2	Hs01022060_m1
CX3CR1	Hs01922583_m1
C3R	Hs00355885_m1
GAPDH	Hs02758991_g1