Learning when effort matters: Neural dynamics underlying updating and adaptation to changes in

performance efficacy

Ivan Grahek[1*], Romy Frömer[1], Mahalia Prater Fahey[1], & Amitai Shenhav[1]

1. *Department of Cognitive, Linguistic, & Psychological Sciences and Carney Institute for*

   Brain Science, Brown University, Providence, RI 02912, USA

* Corresponding author. E-mail address: ivan_grahek@brown.edu

Conflicts of interest: The authors have no conflicts of interest to declare.

Abstract

To determine how much cognitive control to invest in a task, people need to consider whether exerting control *matters* for obtaining rewards. In particular, they need to account for the efficacy of their performance – the degree to which rewards are determined by performance or by independent factors. Yet it remains unclear how people learn about their *performance efficacy* in an environment. Here we combined computational modeling with measures of task performance and EEG, to provide a mechanistic account of how people (a) learn and update efficacy expectations in a changing environment, and (b) proactively adjust control allocation based on current efficacy expectations. Across two studies subjects performed an incentivized cognitive control task while their performance efficacy (the likelihood that rewards are performance-contingent or random) varied over time. We show that people update their efficacy beliefs based on prediction errors – leveraging similar neural and computational substrates as those that underpin reward learning – and adjust how much control they allocate according to these beliefs. Using computational modeling, we show that these control adjustments reflect changes in information processing, rather than the speed-accuracy tradeoff. These findings demonstrate the neurocomputational mechanism through which people learn how worthwhile their cognitive control is.

Keywords: Motivation; Cognitive control; Learning; Performance efficacy; Reward rate; EEG

Cognitive control is critical for achieving most goals, but it is effortful (Botvinick and

Cohen 2014; Shenhav et al. 2017). To decide how to invest control into a task (e.g., writing an

essay for a competition), a person must weigh these effort costs against the potential benefits of a

given type and amount of control (Manohar et al. 2015; Verguts et al. 2015; Kool and Botvinick

2018). One aspect of these benefits is the significance of the expected outcomes, both positive

(e.g., a monetary prize, social acclaim) and negative (e.g., missed revenue, social derision)

(Atkinson 1966; Leng et al., 2021). An equally important aspect of the expected benefits of

control is the extent to which control *matters* for achieving good outcomes and avoiding bad

ones (Frömer et al., 2021a; Shenhav et al., 2021). This can in turn be decomposed into the extent

to which higher levels of control translate into better performance (e.g., whether writing a good

essay will require substantial or only minimal control resources; *control efficacy*) and the extent

to which better performance translates into better outcomes (e.g., whether prizes are determined

by the strength of an essay or by arbitrary or even biased metrics unrelated to essay-writing

performance; *performance efficacy*). Whereas studies have increasingly characterized the ways

in which control allocation is influenced by expected outcomes (e.g., Parro et al., 2018; Leng et

al., 2021) and the expected efficacy of control (e.g., as a function of task difficulty; Krebs et al.,

2012; Vassena et al., 2014; Chiu and Egner 2019), much less is known about how people

estimate and adjust to the perceived efficacy of their performance in a given environment.

We recently showed that when participants are explicitly instructed about how

efficacious their performance will be on an upcoming trial, they exhibit behavioral and neural

responses consistent with increased control (Frömer et al., 2021a). We had participants perform a

standard cognitive control task (Stroop) for potential monetary rewards, and we varied whether

obtaining those rewards was contingent on performing well on the task (high performance

efficacy) or whether those rewards were determined at random (low efficacy). We showed that

people allocate more control when they expect to have high compared to low efficacy, reflected

in higher amplitudes of an EEG index of proactive control (the contingent negative variation

[CNV]) and in improved behavioral performance. These results demonstrate that participants

leverage expectations about the extent to which their performance matters when deciding how

much cognitive effort to invest in a task. However, in this work, participants were explicitly cued

with the level of performance efficacy to expect on a given trial, and those predictive cues

retained the same meaning across the session. Thus, how it is that people learn these efficacy

expectations in environments where contingencies are not instructed, and how they dynamically

update their expectations as contingencies change, remains unanswered.

Outside of the domain of cognitive control, a relevant line of work has examined how

people learn about the factors that determine future outcomes when selecting between potential

courses of action. In particular, work in this area has shown that people are able to learn about

and update their expectations of the likelihood that a given action will generate a given outcome

(action-outcome contingency; Dickinson and Balleine 1995; Moscarello and Hartley 2017; Ly et

al. 2019). People preferentially, and more vigorously, select actions that reliably lead to desired

outcomes (i.e., the more contingent those outcomes are on the action in question; Liljeholm et al.

2011; Manohar et al. 2017), and work in both animals (Balleine & O'Doherty, 2010) and humans

(Norton & Liljeholm, 2020; Dorfman et al., 2021; Ligneul et al., 2022; Morris et al., 2022) has

helped to characterize the neural systems that support this process of learning and action

selection. However, given the focus on discrete actions and their immediate relationship with

outcomes, research into these action-outcome contingencies is unable to capture key aspects that

are unique to selection of control states. Most notably, cognitive control signals (e.g., attention to

one or more features of the environment) are multidimensional, not immediately observable by

either the participant or experimenter, and their relationship with potential outcomes is

intermediated by the many-to-many relationship between control states and task performance

(Ritz et al., 2022). While there has been research into how people learn to adjust cognitive

control signals based on changes in their task environment, here again work has focused on how

people adapt to changes in outcomes (e.g., Otto and Daw 2019; Bustamante et al. 2021) and

changes in the relationship between control and performance (with increasing task difficulty;

Bugg et al. 2011; Nigbur et al. 2015; Bejjani et al. 2018; Jiang et al. 2020). The mechanisms by

which people learn about the relationship between performance and outcomes (performance

efficacy), and how they adjust their control allocation accordingly, remain largely unexplored.

Here, we seek to fill this gap by studying the mechanisms through which expectations of

performance efficacy are formed, updated, and used to guide control allocation. To do so, we

extend our previous approach (Frömer et al., 2021a) – which studied how behavioral and neural

correlates of control allocation vary when performance efficacy is explicitly cued – to examine

how participants learn and adapt their control under conditions where efficacy was un-cued

(having to instead be learned from feedback) and gradually varied across a wide range of

potential efficacy values over the course of the session. We use computational reinforcement

learning models to show that expected efficacy can be learned from feedback through iterative

updating based on weighted prediction errors (Sutton and Barto 2018), and model-based single-

trial EEG analyses to show that these efficacy prediction errors modulate a canonical neural

marker of reward-based learning and behavioral adjustment (Fischer and Ullsperger 2013). We

further provide evidence that efficacy estimates learned in this way are used to guide the

allocation of control. In our EEG study and a second behavioral study, participants tended to

perform better when efficacy was higher. We also provide evidence that a neural marker of

control allocation (Schevernels et al. 2014a) tends to increase with increasing model-based

efficacy estimates. Using a drift diffusion model (Ratcliff & McKoon, 2008; Wiecki, Sofer, &

Frank, 2013), in Study 2 we further show that the performance improvements related to

increased performance-efficacy reflect facilitation of task-related information processing

(reflected in increased drift rates), rather than changes in the speed-accuracy tradeoff (i.e.,

thresholds). Taken together, these results show that efficacy estimates can be learned and

updated based on feedback, leveraging general cognitive and neural mechanisms of predictive

inference.

80   **Materials and Methods**

Study 1

Participants

We recruited forty-one participants with normal or corrected-to-normal vision from the

Brown University subject pool. One participant was excluded due to technical issues. The final

data set included 40 participants (24 females, 16 males; median age = 19). Participants gave

informed consent and were compensated with course credits or a fixed payoff of $20. In

addition, they received up to $5 bonus that depended on their task performance ($3.25 on

average). The research protocol was approved by Brown University's Institutional Review

Board.

Experimental design

In the main task, taking approximately 45 minutes, participants performed 288 Stroop

trials (Figure 1A). Each trial started with the presentation of a cue (grey circle) that remained on

the screen throughout the trial. After a period of 1500 ms, a Stroop stimulus was superimposed

until a response was made or 1000 ms elapsed, at which time it was sequentially replaced with

two types of feedback presented for 1000 ms each. Each trial onset was preceded by a fixation

cross (randomly jittered between 1000 and 1500ms). Participants responded to the ink color of

the Stroop stimulus (red, green, blue, or yellow) by pressing one of four keyboard keys (D, F, J,

and K). Stimuli were either color words same as the ink color (congruent, n = 108) or different

(incongruent, n = 108), or a string of letters "XXXXX" (neutral, n= 72). Feedback informed

them whether they obtained a reward (reward, "$50c" or no reward, "$0c") and whether the

reward they received depended on their performance (performance-based feedback, a button

graphic), or not (random feedback, a dice graphic). In order to earn rewards in the performance-

based case, participants had to be both accurate and respond within an individually calibrated

response deadline (see details below). The order of the two types of feedback was pseudo-

randomized with half of the trials showing reward feedback first and the other half efficacy

feedback. Every 2-4 trials the feedback was followed by a probe of efficacy ("How much do you

think your rewards currently depend on your performance?") or reward rate ("How often do you

think you are currently being rewarded?") to which participants responded on a visual analog

scale ranging from 0 to 100. The number and timing of the probes was randomized per subject

resulting in a median of 45 efficacy probes (SD=3.38) and 47 reward probes (SD=2.81).

Efficacy (performance-based or random rewards) on each trial was sampled from a

binomial distribution with probabilities ranging between 0.1 and 0.9 that drifted over the course

of the experiment and were predetermined (Figure 1B). In order to ensure that the performance-

based and random trials did not differ in reward rate, reward feedback for the random trials was

sampled from the moving window of the reward feedback of the previous 10 performance-based

trials. At the beginning of the experiment a window with 8 rewards and 2 no rewards was used to

reflect the pre-calibrated reward rate (details below), and this moving window was then updated

after every trial. Thus, reward rate was not experimentally manipulated in the experiment and

remained constant. We confirmed that reward rate was matched across performance-based and

random trials by comparing reward probability between these trial types ($b = 0.01$; 95% CrI [-

0.07, 0.10]; $p_{b > 0} = 0.38$).

Prior to the main task, participants performed several practice phases of the Stroop task

(approximately 15 minutes). First, they practiced the mappings between colors and keyboard

keys (80 trials). Then they completed a short practice of the Stroop task with written feedback

("correct" or "incorrect") on each trial (30 trials). Participants then completed 100 more of such

trials during which we individually calibrated the reaction time deadline such that participants

yielded approximately 80% reward rate. The reaction time calibration started from a fixed

deadline of 750ms and increased or decreased this threshold in order to ensure that participants

earn rewards on 80% of trials (i.e., that they are both accurate and below the deadline). The

deadline obtained in this way (M = 796ms; SD = 73ms) was used in the main experiment and

was not further adjusted. In the final practice phase participants were introduced to the two types

of feedback which they would see in the main experiment (30 trials).

The experimental task was implemented in Psychophysics Toolbox (Brainard 1997; Pelli

1997; Kleiner et al. 2007) for Matlab (MathWorks Inc.) and presented on a 23 inch screen with a

1920 x 1080 resolution. All of the stimuli were presented centrally while the participants were

seated 80 cm away from the screen.

Psychophysiological recording and preprocessing

EEG data were recorded at a sampling rate of 500 Hz from 64 Ag/AgCl electrodes

mounted in an electrode cap (ECI Inc.), referenced against Cz, using Brain Vision Recorder

(Brain Products, München, Germany). Vertical and horizontal ocular activity was recorded from

below both eyes (IO1, IO2) and the outer canthi (LO1, LO2), respectively. Impedances were

kept below 10 kΩ. Offline, data were processed using custom made Matlab scripts (Frömer et al.

2018) employing EEGlab functions (Delorme and Makeig 2004). Data were re-referenced to

average reference, ocular artifacts were corrected using brain electric source analyses (Ille et al.

2002) based on separately recorded prototypical eye movements. The cleaned continuous EEG

was then low pass filtered at 40 Hz and segmented into epochs around cue onset (-200 to 1500

ms), stimulus onset, and both efficacy and reward feedback (-200 to 800 ms). Baselines were

corrected to the average of each 200 ms pre-stimulus interval. Segments containing artifacts,

values exceeding $\pm$ 150 µV or gradients larger than 50 µV, were excluded from further analyses.

Single trial ERPs were then exported for further analyses in R (R Core Team 2017). The

late CNV was quantified between 1000 and 1500 ms post neutral cue onset (Schevernels et al.

2014b; Frömer et al. 2016; Frömer et al. 2021a) as the average activity over 9 fronto-central

electrodes (Fz, F1, F2, FCz, FC1, FC2, Cz, C1, and C2). The P3b was quantified between 350

and 500 ms (Fischer and Ullsperger 2013) for both reward and efficacy feedback and calculated

as the averaged activity over 9 centro-parietal electrodes (Pz, P1, P2, POz, PO1, PO2, CPz, CP1,

and CP2).

Learning models and statistical analyses

Learning models. Participants provided their subjective estimates of efficacy and reward every

4-8 trials (a total of 45 estimates), and we sought to fit a learning model to these estimates to be

able to predict trial-by-trial adjustments in performance and neural markers of learning and

cognitive control allocation. In order to obtain trial-by-trial estimates of efficacy and reward rate,

we fitted two temporal difference learning models (Gläscher et al., 2010; Sutton & Barto, 2018)

to the continuous subjective estimates of efficacy and reward rate (Rutledge et al., 2014; Eldar et

al., 2016; Nagase et al., 2018). The first model ("1 learning rate efficacy model") assumed that

the estimate of efficacy for the next trial (E_{t+1}) depended on the current efficacy estimate (E_t)

and the prediction error (δ_t) weighted by a constant learning rate (α):

167 $$E_{t+1} = E_t + \alpha * \delta_t$$

Where $0 \leq \alpha \leq 1$, and the prediction error is calculated as the difference between the contingency

feedback on the current trial (e_t) and the efficacy estimate on that trial: $\delta_t = e_t - E_t$. The model

started from an initial value (free parameter) and updated the model-based efficacy estimate

based on the binary efficacy feedback on each trial. For example, assuming a learning rate of 0.5

and the initial value of 0.5, the model would update the initial estimate following efficacy

feedback signaling "performance-based" ($e_t = 1$) to 0.75. If on the next trial contingency

feedback was "random" ($e_{t+1} = 0$) the model-based efficacy estimate would drop to 0.6. The

model was fitted separately to the subjective estimates of efficacy with only the learning rate as a

free parameter. The second model ("2 learning rates efficacy model") was the same as the first

model, but it included two learning rates: one learning rate for learning from the "performance-

based" feedback, and another for learning from the "random" feedback. Finally, as a baseline, we

also included the "intercept model" which did not update the efficacy estimate throughout the

experiment, but just assumed that the estimate took one constant value. Importantly, the same

models were fitted to obtain the model-based estimates of reward on each trial ("1 learning rate

reward model" and the "2 learning rate reward model"). These models were fitted using trial-by-

trial reward feedback and the subjective estimates of reward. The models were fit hierarchically

to the data using maximum likelihood estimation (using mfit (https://github.com/sjgershm/mfit).

To calculate the likelihood of each data point, model-based estimates (0-1 range) were compared

to the subjective efficacy estimates (range normalized to 0-1 range for each participant).

Likelihood was evaluated on trials which included a subjective estimate, as the likelihood that

the difference between the model-based and the empirical estimate comes from a Gaussian

distribution centered on 0 with a variance which was fitted as a free parameter each subject. This

variance parameter served as the noise in the estimates. Likelihoods were log transformed,

summed, and then maximized using the fmincon function in MATLAB.

We performed a parameter recovery study to show that the most complex model (the 2

learning rates model) can be successfully recovered. We simulated a dataset with the same

number of trials and subjective efficacy or reward probes as in the actual experiment. We used

the efficacy drifts presented to the actual subjects (half of the simulated subjects saw one drift,

and half its inverse), and we used the reward feedback sequences of two actual subjects from our

experiment. We simulated 200 agents which learned both efficacy and reward with the noise

parameter fixed to 0.2, intercept fixed to 0.5, and the positive and negative learning rates

sampled from a uniform distribution ranging from 0.001 to 0.5. These parameters were matched

based on the range of values obtained from the empirical fits to our data. As shown in Figure S1

we were able to very reliably recover the simulated parameters for both efficacy and reward rate

learning.

Statistical analyses. The efficacy and reward rate estimates obtained through fitting the learning

model were then used to analyze the behavioral and EEG data. To this end, we fitted Bayesian

multilevel regressions to predict subjective estimates of efficacy and reward rates, reaction times,

accuracy, as well as the CNV and P3b amplitudes. Subjective estimates of efficacy and reward

rate were regressed onto efficacy or reward feedback. Reaction times and accuracies were

regressed onto trial-by-trial model-based estimates of efficacy and reward rate, as well as trial-

by-trial CNV amplitude, while controlling for congruency. The P3b amplitudes were analyzed in

two ways: with trial-by-trial model-based estimates of efficacy and reward rate and current

feedback as predictors, and with model-based prediction errors and learning rates for each

feedback type. CNV amplitudes were regressed onto trial-by-trial model-based estimates of

efficacy. All of the fitted models controlled for the influence of the reward rate estimates.

Parallel analyses were done to predict the P3b in response to reward feedback, while controlling

for the efficacy estimates.

The regression models were fitted in R with the *brms* package (Bürkner 2016) which

relies on the probabilistic programming language *Stan* (Carpenter et al. 2016) to implement

Markov Chain Monte Carlo (MCMC) algorithms and estimate posterior distributions of model

parameters. The analyses were done based on the recommendations for Bayesian multilevel

modeling using *brms* (Bürkner 2016; Bürkner 2017; Nalborczyk and Bürkner 2019). The fitted

models included constant and varying effects (also known as fixed and random) with weakly

informative priors (except for the behavioral and CNV analyses, see below for details) for the

intercept and the slopes of fixed effects and the likelihoods appropriate for the modeled data (Ex-

Gaussian for reaction times, Bernoulli for accuracy, and Gaussian for the subjective estimates

and the EEG amplitudes). The fitted models included all of the fixed effects as varying effects.

All of the continuous predictors in the model were centered and the categorical predictors were

contrast coded. Four MCMC simulations ("chains"; 20,000 iterations; 19,000 warmup) were run

to estimate the parameters of each of the fitted models. The convergence of the models was

confirmed by examining trace plots, autocorrelation, and variance between chains (Gelman and

Rubin 1992). After convergence was confirmed, we analyzed the posterior distributions of the

parameters of interest and summarized them by reporting the means of the distribution for the

given parameter (*b*) and the 95% credible intervals (95% CrI) of the posterior distributions of

that model. We report the proportion of the posterior samples on the relevant side of 0 (e.g., $p_{b<0}$

= 0.9), which represents the probability that the estimate is below or above 0. We also report

Bayes factors (BF) calculated using the Savage-Dickey method (Wagenmakers et al. 2010). We

report the BFs in support of the alternative hypothesis against the null (BF_{10}), except for the

analyses of accurate RT, accuracies, and CNV amplitude in which we have informative priors

based on our previous study (Frömer et al. 2021a), and in which case we support the evidence in

favor of the null (BF_{01}).

To compare the positive and negative learning rates we fitted a model in which the

learning rates were predicted by the learning rate type (Kruschke 2013). In this model we used

Gaussian distributions (mean, standard deviation) as priors (intercept: (0.5,0.5); slopes: (0,0.5)).

We fitted two separate models to predict the subjective estimates of efficacy and reward

rate based on previous feedbacks. At each timepoint the estimates were predicted by the current,

and previous 4 feedbacks. The feedback on each of the trials (performance-based vs. random or

reward vs. no reward) was entered as a constant effect and the models also included the intercept

as a varying effect. As the subjective estimates could vary between 0 and 1, we used Gaussian

distributions (intercept: (0.5,0.2); slopes: (0,0.2)) as priors.

For predicting the P3b amplitude in response to the onset of the efficacy feedback, we

fitted two models. First, we fitted a model which included the model-based estimate of efficacy

(prior to observing the current feedback), the actual feedback, and the interaction between the

expected efficacy and the observed efficacy feedback. Additionally, we controlled for the reward

rate estimate. Second, we fitted separate models which included the model-based prediction

errors, the influence of the between-subject learning rates, and their interaction with the

prediction errors, while controlling for the estimate of the reward rate. In this analysis the

learning rates (one for feedback type for each subject) were mean-centered within subjects and

thus any effect of the learning rates is driven by the difference between the random and the

performance-based learning rate. For these models we selected wide Gaussian priors (intercept:

(5,3); slopes (0,3)). The same logic in building models was applied for the analyses of the reward

feedback. In these analyses we focused on the reward feedback processing and how it interacted

with the model-based estimates of reward rates, while controlling for the model-based estimates

of efficacy. We analyzed only the trials with correct responses for both the efficacy and the

reward feedback analyses.

To test the influence of efficacy on the late CNV, we fitted a model which predicted the

CNV based on the model-based efficacy estimates, while controlling for the effect of the reward

rate estimates. Drawing on the results of our previous study (Frömer et al. 2021a), this model

included Gaussian priors for the intercept (-0.16, 1.59) and the efficacy (-0.30, 0.73) and reward

(0,0.73) slopes.

For predicting reaction times and accuracy we fitted models which included congruency

(Facilitation: difference between neutral and congruent trials; Interference: difference between

incongruent and neutral trials) and the model-based efficacy estimates, while controlling for the

reward rate estimates. We used Gaussian distributions as informative priors based on our

previous study (Frömer et al. 2021a), for both the reaction times (intercept: (624, 58.69);

facilitation (15.54, 21.93), interference (61.12, 37.49); efficacy (-10.26, 19.51); reward (0,

19.51)) and accuracy[1] analyses (intercept: (2.11, 0.81); facilitation (-0.45, 0.64), interference (-

0.53, 0.81); efficacy (0.09, 0.32); reward (0, 0.32)).

To investigate how the late CNV influences the behavior, we fitted two models in which

we predicted the reaction times and accuracy based on the CNV amplitude. The prior

distributions for these models were weakly informative Gaussian distributions for predicting

both the reaction times (intercept: (650, 200); slope: (0, 50)) and accuracy (intercept: (0.7, 0.2);

slope: (0, 0.2)).

To visualize the topographies of the relevant ERP effects, we fitted the relevant models to

all 64 channels and then plotted the posterior estimates of the effects of interest at each electrode

(cf. Frömer et al., 2021b).

Study 2

Participants

We recruited eighty-seven participants residing in the United States from Prolific – an

online platform for data collection. Participants had normal or corrected-to-normal vision and

gave informed consents. They were compensated with a fixed payoff of $8 per hour (median

completion time of 74 minutes) plus a monetary bonus based on points earned during the task

($1 on average). The research protocol was approved by Brown University's Institutional

Review Board.

We a priori excluded participants who did not pass attention checks (N=8) or who took

substantially longer than the average participant to complete the study (N=2 participants who

took over 130 minutes), suggesting that they did not sustain attention to the experiment over that

---

[1] Note that the prior distributions are set in log-odds.

time. We fit our learning models to data from the remaining 77 subjects, and then excluded

participants whose performance suggested inattention to the overall task (based on accuracies

less than 70% across all trials – including the trials in which performance efficacy was low, N=6)

or inattention to the task feedbacks and efficacy probes (based on low learning rates (N=19), and

one subject with no variance in responses to reward probes). To identify participants with

exceedingly low learning rates, we submitted all positive and negative efficacy learning rates to

unsupervised Gaussian mixture models (as implemented in the Mclust package; (Scrucca et al.,

2016) to determine the best fitting number and shape of clusters (model comparison via BIC).

This procedure identified four clusters of subjects with different overall learning rates (Figure

S2B-C) and we excluded subjects from the first cluster as they all had very low learning rates

relative to the other participants (both learning rates < 0.03). The subjects excluded based on low

learning rates were most likely not paying attention to efficacy feedback, or were always giving

very similar responses to the efficacy probes (Figure S3). The final sample included 51

participants (31 females, 20 males; median age = 29).

Experimental design

In order to better understand the computational mechanisms that lead to improved

behavioral performance in high efficacy states (Study 1), we wanted to fit a Drift Diffusion

model (DDM; Ratcliff & McKoon, 2008) to our behavioral data. If people allocate more

attention when they think they have high performance efficacy, this should be observed as an

increase in the drift rate (speed of evidence accumulation). However, Study 1 included a tight

respond deadline for earning a reward, making it more challenging to fit the DDM. To avoid this

issue, in Study 2 we adjusted the task to a free response paradigm which allowed us to

investigate drift rate and threshold adjustments (cf. Leng et al., 2021). We used this design to test

the hypothesis that higher efficacy estimates should predict increased drift rates.

Instead of single trials, participants now completed 288 intervals during which they could

respond to as many trials (congruent and incongruent, removing the neutral condition) as they

wished within a fixed time window (randomly selected between 2000, 3000, or 4000ms). Apart

from this, the structure of the task remained the same: participants saw a fixation cross (1000,

1500, or 2000ms), then completed as many trials as they wished during a fixed interval, followed

by the feedback (1500 ms) on how many points they earned and whether this was based on their

performance or awarded to them based on random chance. Note that participants now received

continuous reward feedback (10 points per correct response instead of the binary reward-no

reward in Study 1). For example, if participants completed 4 trials correctly and 2 trials

incorrectly within a performance-based interval they would receive 40 points and see feedback

informing them that the points were based on their performance. To determine the number of

points on random intervals, the same yoking procedure as in Study 1 was employed ensuring that

the amount of reward was matched between performance-based and random intervals (reward

amounts on random intervals were sampled from the moving average window of the past 10

performance-based intervals). We confirmed that the yoking procedure was successful by

comparing the reward amounts on the two interval types ($b = 0.00$; 95% CrI [-0.00, 0.02]; $p_{b > 0} =$

0.16). As in Study 1, participants were probed every 2-4 intervals to estimate either how much

they thought their rewards depended on their performance, or how often they were rewarded. We

again implemented an efficacy drift (modified, but comparable to the drift in Study 1; Figure

S2A), now across the 288 intervals of the task.

We gamified the task in order to make it more appealing for the participants. Instead of

the Stroop task, we used a picture-word interference task in which four grey-scaled images of

fruit (apple, pear, lemon, and peach) were overlaid by those fruit words. Participants responded

to the image, while ignoring the word, by pressing one of the 4 corresponding keys. They first

practiced this task, and then were introduced with a cover story telling them that they are in the

garden and need to water the fruit in little patches by pressing the correct keys. They were

instructed that they will be moving through a garden and that in some patches watering will

directly translate into how many points they will be earning, while in the others that will not be

the case (the efficacy drift). The experiment was implemented in Psiturk (Gureckis et al., 2016)

and the participants performed the task on their own computers and were required to have a

keyboard.

Learning models, Drift Diffusion model, and statistical analyses

Learning models. We fitted the same set of learning models as in Study 1, performed model

comparison, and got the interval-by-interval model-based estimates of performance efficacy and

reward. Note that in this version of the task participants earned points in each interval, unlike the

binary rewards (reward vs. no reward) in Study 1. This meant that the reward learning model

learned reward magnitudes rather than reward probabilities. However, model fitting and the

further analyses were the same as in Study 1.

Statistical analyses. We fitted the same Bayesian multilevel models as in Study 1 to predict the

influence of previous efficacy feedbacks on the subjective efficacy estimates, as well as the

influence of the model-based efficacy estimates on reaction times and accuracies. For the

analyses of the subjective efficacy estimates we used the same priors as in Study 1. For the

analyses of the reaction times and accuracies, we used the posterior distributions obtained in

Study 1 as the informative priors for the congruency, efficacy, and reward effects. The reaction

times and accuracy models also controlled for the effects of the average congruency level in the

interval and the interval length.

Drift Diffusion model. This model decomposes participant's behavior into drift rate (the speed

of evidence accumulation) and response threshold (the level of caution), allowing us to

investigate which of these two components is affected by the efficacy estimates. We fitted the

model using Bayesian hierarchical estimation as implemented in the HDDM package (Wiecki,

Sofer, & Frank, 2013). The fitted model included the main effects of efficacy and reward rate

estimates onto both drift rate and threshold, and included the effect of congruency on the drift

rate. The responses were coded as correct or incorrect, and trials with reaction times below

250ms were excluded. All of the effects were allowed to vary across subjects, and we ran five

MCMC chains (12,000 iterations; 10,000 warmup). We confirmed convergence by examining

trace plots and variance between chains.

377   **Results**

378   **Study 1**

379   To investigate how efficacy estimates are learned, and how they affect control allocation,

in Study 1 we recorded EEG while 40 participants performed a modified version of the Stroop

task (Figure 1A). Across trials we varied whether reward outcomes ($0.50 vs. $0.00) were

determined by a participant's performance on a given trial (responding accurately and below a

pre-determined response time criterion; *performance-based* trials) or whether those outcomes

were determined independent of performance (based on a weighted coin flip; *random* trials).

Over the course of the session, we gradually varied the likelihood that a given trial would be

performance-based or random such that, at some points in the experiment, most trials were

performance-based (high efficacy level), and at other points most trials had random outcomes

(low efficacy level) (Figure 1B). Importantly, unlike in our previous study (Frömer et al., 2021a),

participants were not told whether a given trial would be performance-based (maximal efficacy)

or random (minimal efficacy), but instead had to estimate their current efficacy level based on

recent trial feedback, which indicated both the reward outcome ($0.50 vs. $0.00) and how that

outcome was determined (performance-based [button symbol] vs. random [dice symbol]). We

held reward rate constant across both feedback types by yoking reward rate on random trials to

the reward rate on performance-based trials, and counter-balanced the time-course of the gradual

change in efficacy (see Methods for details). To capture changes in efficacy expectations over

the course of the session, we probed participants every 4-8 trials (averaging 44.3 probes per

participant) to report their current estimates of efficacy. These efficacy probes were interleaved

with probes asking participants to estimate the current reward rate, serving as foils and for

control analyses.

Figure 1. Manipulating expected efficacy and assessing learning in Study 1. A. Trial Schematic. On each trial participants saw a cue (gray circle), predicting the onset of a Stroop stimulus (target), and were then sequentially presented with reward and efficacy feedback. On half of the trials, efficacy feedback was presented first, and on the other half reward feedback was presented first. Every 2-4 trials participants were subsequently asked to estimate their current efficacy level ("How much do you think your rewards currently depend on your performance?") or reward rate ("How often do you think you are currently being rewarded?"). **B. Efficacy manipulation.** We let the probability of performance-based vs. random feedback continuously drift over the course of the experiment (inversed for one half of the sample). Arrows mark time points with low and high efficacy, respectively. When efficacy was low, rewards were more likely to be random, whereas when efficacy was high, rewards were more likely to be performance-based.

Participants dynamically update efficacy expectations based on feedback

To determine whether and how participants learned their current efficacy levels, we first

analyzed the influence of previous efficacy feedback (whether outcomes on a given previous trial

had been performance-based or random) on one's current subjective estimates of efficacy. We

found very strong evidence that participants adjusted their subjective efficacy upward or

downward depending on whether the most recent trial was Performance-Based or Random ($b =$

0.14; 95% CrI [0.12, 0.16]; $p_{b < 0} = 0$; $BF_{10} > 100$), and that this remained true (but to diminishing

degrees) up to five trials back (all $p_{b < 0} < 0.01$; Figure 2A; Table S1). This effect of feedback was

present only for the efficacy feedback, while reward feedback did not predict the subjective

estimates of efficacy (Figure 2A). These results suggest that participants were dynamically

updating their efficacy estimates based on efficacy feedback.

Figure 2. Efficacy learning is captured by a reinforcement learning model with separate learning rates for performance-based and random feedback in Study 1. A. Efficacy estimates track recent efficacy feedback. Regression weights for the influence of the current (t) and previous contingent vs. random feedback, as well as reward feedback, on the subjective efficacy estimate. Error bars represent 50% and 95% highest density intervals. **B. A separate learning rate model captures efficacy learning best.** BICs of fitted intercept only and one learning rate model relative to two learning rate models are plotted for each participant and favor the two learning rate model. **C. Learning rate biases vary between participants.** Positive and negative learning rate estimates are plotted for all participants. Points below the diagonal indicate higher learning rates for performance-based compared to random feedback, and points above the opposite. **D. Example learning trajectories.** Subjective and model-based efficacy estimates, and a running average of the previous 5 efficacy feedbacks, for a participant with a higher learning rate for the contingent compared to random feedback (upper) and a participant with the reverse bias (lower).

This pattern of learning was accounted for by a standard reinforcement learning (RL)

algorithm, the delta-rule, according to which efficacy estimates are updated in proportion to the

prediction error between the expected efficacy and the efficacy feedback on a given trial (i.e.,

whether a given outcome was determined by performance or not) (Sutton and Barto 2018).

Interestingly, consistent with studies of reward learning (Niv et al. 2012; Collins and Frank 2014;

Lefebvre et al. 2017; Chambon et al. 2020; Garrett and Daw 2020), we found that the RL model

that best accounted for our data allowed efficacy estimates to be updated differently from trials

that were more efficacious than expected (Performance-Based feedback) than from trials that

were less efficacious than expected (Random feedback), by having separate *learning rates*

scaling prediction errors in the two cases. Even when accounting for model complexity, this Two

Learning Rate Efficacy model outperformed a One Learning Rate Efficacy model as well as a

baseline model that fits a single constant efficacy estimate and no learning (Intercept Model)

(Figure 2B). In addition, we were able to successfully recover the parameters of this model from

a simulated dataset matched to our data (see Methods and Figure S1). We found that the two

learning rates for this best-fit model varied across the group (Figure 2E), but did not find that one

was consistently higher than the other ($b = 0.02$; 95% CrI [-0.04, 0.08]; $p_{b < 0} = 0.260$;

$BF_{01}=13.55$; Figure S4), despite most participants (80%) tending to learn more from

Performance-Based than Random trials. Finally, model-based efficacy estimates were strongly

related to the raw subjective estimates on trials on which participants reported efficacy ($b = 0.77$;

95% CrI [0.62, 0.91]; $p_{b < 0} < 0.001$; $BF_{01} > 100$; $R^2=0.50$), demonstrating that the model

successfully captured the raw estimates. Taken together, these results show that participants

dynamically updated their expected efficacy based on trial-by-trial feedback, and that they did so

differentially based on whether the trial was more or less efficacious than expected. The fitted

models further enable us to generate trial-by-trial estimates of expected efficacy and efficacy

prediction errors, which we use in model-based analyses of behavior and neural activity below.

Note that in all the following analyses we control for reward estimates obtained from models fit

to reward feedback (for details see below).

The feedback-related P3b indexes updating of efficacy expectations

440   To investigate the neural mechanism underlying feedback-based learning of efficacy, we

probed the centroparietal P3b ERP component (Figure 3A), an established neural correlate of

prediction-error based learning (Fischer and Ullsperger 2013; Nassar et al. 2019). If the P3b

indexes feedback-based updating of *efficacy* predictions, as it does for reward predictions, we

would expect this ERP to demonstrate several key markers of such a learning signal. First, we

would expect the P3b to reflect the extent to which efficacy feedback (Performance-Based vs.

Random) deviates from the current level of expected efficacy. In other words, the P3b should

track the magnitude of the *unsigned efficacy prediction error (PE)* on a given trial. We tested

this by examining how the amplitude of the P3b to a given type of efficacy feedback varied with

model-based estimates of the participant's efficacy expectation on that trial, while holding the

expected reward rate constant (see the Methods and Materials sections for the details of the

experimental design and the statistical models). If the P3b signaled efficacy PE then its

amplitude should scale inversely with the expected probability of a given type of feedback (i.e.,

how *unexpected* that feedback is), thus correlating *negatively* with expected efficacy on trials

providing Performance-Based feedback and correlating *positively* with expected efficacy on

trials providing Random feedback. In addition to overall higher P3b to performance-based

compared to random feedback ($b = 0.86$; 95% CrI [0.42, 1.31]; $p_{b<0} = 0$; $BF_{10} = 30.86$; Figure

2B), we found exactly this predicted interaction between feedback type and expected efficacy (b

= -2.40; 95% CrI [-4.07, -0.74]; $p_{b>0} = 0$; $BF_{10} = 24.40$; Figure 2B; Figure S5A; Table S2), with

the P3b amplitude decreasing with model-based estimates of expected efficacy on Performance-

Based trials (b = -1.49; 95% CrI [-2.72, -0.27]; $p_{b>0} = 0.03$; $BF_{10} = 1.52$) and increasing

numerically, but not robustly with expected efficacy on Random trials ($b = 0.91$; 95% CrI [-0.34,

2.21]; $p_{b<0} = 0.12$; $BF_{10} = 0.44$). Accordingly, when we regressed P3b amplitude on our model-

based estimates of trial-to-trial efficacy PE, we found a positive relationship ($b = 1.25$; 95% CrI

[0.35, 2.15]; $p_{b<0} = 0.01$; $BF_{10} = 5.84$; Figure 3C-left; Table S2).

Figure 3. P3b reflects dynamically changing efficacy estimates during processing of efficacy feedback in Study 1. A. ERP average for the P3b locked to the onset of the efficacy feedback separately for performance-based and random feedback. The grey area shows the time window used for quantifying the P3b. **B.** LMM predicted P3b amplitudes are plotted for performance-based and random feedback as a function of efficacy estimates. The topography shows the interaction of efficacy estimate with efficacy feedback in the P3b time window. **C.** Predicted (centered) effects of unsigned prediction errors (left) and model-based learning rates (right) on the P3b. Shaded error bars represent 95% confidence intervals. Topographies display fixed-effects estimates.

In addition to tracking efficacy PEs, the second key prediction for the P3b if it indexes

efficacy learning, is that it should track the extent to which PEs are used to update estimates of

efficacy (i.e., the learning rate). In the current study, we found that participants differed in their

learning rates for the two forms of efficacy feedback (Performance-Based vs. Random),

providing us with a unique opportunity to perform a within-subject test of whether P3b tracked

differences in learning rate across these two conditions. Specifically, we could test whether a

given subject's P3b was greater in the feedback condition for which they demonstrated a higher

learning rate. However, we have not found conclusive evidence for the increase in P3b for the

within-subject feedback condition with the higher learning rate ($b = 2.00$; 95% CrI [-2.21, 6.04];

$p_{b<0} = 0.17$; BF = 1.08; Figure 3C-right). While, theoretically, prediction error and learning rate

would interact in predicting the P3b amplitude, we did not observe such an interaction here. This

finding is in line with previous work on reward processing (Fischer and Ullsperger 2013), which

has found additive effects of prediction errors and learning rate on P3b. While we found the

effect of prediction errors, our learning rate effect was going in the expected direction, but was

not conclusive.

The CNV indexes control allocation based on updated expectations of efficacy

Thus far, our findings suggest that participants dynamically updated expectations of their

performance efficacy based on feedback, and that the P3b played a role in prediction error-based

updating of these expectations. Next, we tested the prediction that learned efficacy estimates

determine the expected benefits of control, and thus influence how much control is allocated

(Shenhav et al. 2013). We have previously shown that people exert more control when expecting

their performance to be more rather than less efficacious on the upcoming trial (Frömer et al.,

2021a). This was reflected in better behavioral performance and higher amplitudes of the

contingent negative variation (CNV; Figure 4B-left) - a slow negative fronto-central waveform

preceding target onset, which is increasingly negative as the amount of control allocated in

preparation for the task increases (Grent-'t-Jong and Woldorff 2007; Schevernels et al. 2014a).

Here we used the same marker of control, but, unlike in our previous study, efficacy expectations

were (a) learned rather than explicitly instructed; (b) varied over time rather than having a fixed

association with a cue; and (c) varied continuously across the range of efficacy probabilities

rather than alternating between maximal and minimal efficacy. We were therefore interested in

testing whether these dynamically varying expectations of efficacy, as estimated by our model,

would still exert the same influence on behavior and neural activity.

Consistent with our predictions and our previous findings, participants tended to perform

better when they expected performance to be more efficacious, responding faster on correct trials

with increasing model-based estimates of efficacy (Figure 4A; Figure S6A; Table S3). This

finding replicates the performance effects we observed using instructed cues, albeit with only

modest evidence (b = -16.00; 95% CrI [-34.91, 2.96]; $p_{b > 0}$ = 0.05; BF_{01} = 1.68). Like in our

previous studies, faster responding was not explained by a change in speed-accuracy trade-off, as

accuracy did not decrease with increasing efficacy (b = 0.12; 95% CrI [-0.20, 0.44]; $p_{b > 0}$ = 0.23;

BF_{01} = 1.99; Figure S6A). These analyses controlled for the standard behavioral effects related to

Stroop congruency (i.e., slower and less accurate responding for incongruent relative to

congruent trials; Figure S7), as well as for the reward rate estimates.

If the CNV provides an index of control allocation based on current incentive

expectations, it should both reflect one's latest efficacy estimate and should predict performance

on the upcoming trial. Our findings support both predictions. Regressing single-trial CNV

amplitude onto model-based efficacy estimates, and controlling for expectations of reward

(discussed below), we found that the CNV amplitudes had a positive relationship[2] with the

current efficacy expectations (b = -0.35; 95% CrI [-0.85, 0.16]; $p_{b>0}$ = 0.09; BF_{10} = 2.73; Figure

4B-right; Figure S5B; Table S4). However, this effect was weaker than in the previous

experiment with cued efficacy levels, which is to be expected given that in this experiment

participants had to learn their efficacy levels. As with the behavioral finding above, this result

provides evidence consistent with our previous CNV finding using instructed cues. We further

replicate our earlier finding (Frömer et al., 2021a) that larger CNV amplitude in turn predicted

better performance on the upcoming trial, with participants responding faster (b = 11.41; 95%

CrI [8.09, 14.72]; $p_{b<0}$ = 0; BF_{10} > 100; Figure 4C; Table S5) and more accurately (b = -0.07;

95% CrI [-0.12, -0.01]; $p_{b>0}$ = 0.01; BF_{10} = 3.14; Figure 4D; Table S5) as CNV increased.

---

[2] Note that the CNV is a negative component, thus higher CNV amplitudes (i.e., more control allocation) are more negative.

Figure 4. Efficacy estimates influence allocation of control and behavior in Study 1. A. Higher efficacy predicts faster accurate responses. **B. CNV increases with higher efficacy.** Left: Grand-average ERP for high and low efficacy estimates (median split used for plotting). The shaded area marks the time window used for CNV quantification. Time 0 corresponds to the onset of the neutral cue. Right: Predicted CNV amplitudes as a function of efficacy estimates. The topography shows the fixed effect of the efficacy estimate from the fitted linear model. **C. -D. Larger CNV amplitude predicts better performance.** Predicted accurate RT (**C.**) and accuracy (**D.**) as a function of efficacy estimates. Shaded error bars indicate 95% confidence intervals.

Parallel learning of efficacy and reward rate

We held the amount of reward at stake constant over the course of the experiment, but the

frequency of reward (reward rate) varied over the course of the session based on a participant's

recent performance, and participants were sporadically probed for estimates of their reward rate

(interleaved with trials that were followed by efficacy probes). Our efficacy manipulation

explicitly controlled for this variability by yoking Random-Outcome feedback to a participant's

recent reward rate (see methods for details). However, this additional source of variability also

provided an opportunity to examine additional mechanisms of learning and adaptation in our

task. As in the case of efficacy estimates, reward rate estimates were robustly predicted by

reward feedback on previous trials (Table S1; Figure S8A), and this reward rate learning process

was well captured by a two learning rate reward rate model (Garrett and Daw 2020; S8B-C),

with the model-based estimates successfully predicting the reported subjective estimates ($b =$

0.82; 95% CrI[$0.60, 1.02$]; $p_{b<0} < 0.001$; $BF_{01} > 100$; $R^2=0.58$; Figure S8B-C). Updates to these

reward rate estimates were reflected in P3b modulations of (unsigned) reward prediction errors

and associated learning rates (Fischer and Ullsperger 2013; Figure S9; Table S6). This pattern of

results provides additional evidence that efficacy learning involves similar neural and

computational mechanisms as reward-based learning.

Study 2

Learned efficacy modulates control over information processing

Our findings suggest that people rely on domain-general mechanisms to learn about their

performance efficacy in a given environment, and use these learned estimates of efficacy to

optimize performance on their task. Specifically, in Study 1 we found that higher levels of

learned efficacy are associated with faster responses, albeit with modest evidence ($b = $ -16.00;

95% CrI [-34.91, 2.96]; $p_{b > 0} = 0.05$; $BF_{01} = 1.68$). We also found that this speeding occurred on

correct but not incorrect trials, suggesting that these performance adjustments reflected

attentional control rather than adjustments to speed-accuracy tradeoffs. However, these findings

remain only suggestive in the absence of a formal model, and the presence of a stringent

response deadline in this study (individually calibrated for each subject during the practice phase

to ensure the reward rate of 80%; M = 796ms; SD = 73ms) presented an obstacle to fitting our

behavioral data to such a model without additional assumptions (e.g., regarding the form of a

collapsing threshold).

To provide further support for our proposal that learned efficacy influences control over

information processing, we ran an additional behavioral study (Study 2). Participants in this

study (N = 51) performed a web-based version of the task in Study 1, with the biggest

modification being that the Stroop trials (now using picture-word rather than color-word

interference) were performed over the course of short self-paced time intervals rather than trial

by trial as in Study 1. Specifically, participants were given limited time windows (2-4s) to

complete as many Stroop trials as they wanted to and were rewarded at the end of each interval

(cf. Leng et al., 2021). When rewards were performance-based, participants received a number of

points exactly proportional to the number of correct responses they gave during that window.

When rewards were random, the number of points was unrelated to performance on that interval

but (as in Study 1) was yoked to their performance in previous performance-contingent intervals.

Following our approach in Study 1, we varied the likelihood of a given interval being

performance-based or random over the course of the session (Figure S2A), and sporadically

probed participants for their subjective estimates of expected efficacy and reward rate. While in

most respects very similar to the paradigm in Study 1, one noteworthy feature of this self-paced

design is that it resulted in a much less stringent deadline for responding, thus producing reaction

time patterns more typical of free-response paradigms for which the traditional (fixed-threshold)

DDM was designed. Note that because this was an online sample we also employed additional

cutoff criteria to exclude inattentive participants, as detailed in the Methods section and Figures

S2B-C and S3.

Replicating the learning patterns observed in Study 1, subjective estimates of efficacy in

Study 2 reflected a running average over recent efficacy feedback ($b = 0.18$; 95% CrI [0.16,

0.20]; $p_{b < 0} = 0$; $BF_{10} > 100$). This effect was again weighted towards the most recent feedback

but still present up to five intervals back (all $p_{b < 0} < 0.01$; Figure 5A, Figure S10, and Table S7).

As in Study 1, this learning pattern was best captured by an RL algorithm with two learning rates

(Figure 5B), though positive and negative efficacy learning rates did not significantly differ from

one another on average ($b = 0.03$; 95% CrI [-0.03, 0.08]; $p_{b < 0} = 0.260$; $BF_{01}=11.74$; Figure 5C).

As in Study 1, the model-based efficacy estimates successfully predicted the raw subjective

estimates on intervals on which participants reported their efficacy beliefs ($b = 0.86$; 95% CrI

[0.81, 0.90]; $p_{b < 0} < 0.001$; $BF_{01} > 100$; $R^2 = 0.60$), and the same was true for the model-based

reward estimates predicting the subjective reward estimates ($b = 1.00$; 95% CrI[0.95, 1.04]; $p_{b < 0}$

< 0.001; $BF_{01} > 100$; $R^2=0.66$).

Critically, we once again found that higher model-based estimates of efficacy predicted

better performance on the upcoming interval. Participants responded faster ($b = -10.25$; 95% CrI

[-19.86, -0.20]; $p_{b > 0} = 0.02$; $BF_{01} = 2.33$) and more accurately ($b = 0.23$; 95% CrI [0.03, 0.43];

$p_{b > 0} = 0.01$; $BF_{01} > 100$) with increasing model-based efficacy estimates (Figure 6B, Figure

S6B, and Table S8). To formally test whether these behavioral patterns reflected adjustments in

information processing (i.e., the rate of evidence accumulation once the stimuli appeared) or

instead reflected adjustments in speed-accuracy tradeoffs (i.e., the threshold for responding), we

fit these data with the hierarchical drift diffusion model (HDDM; Wiecki, Sofer, & Frank, 2013).

We tested whether model-based efficacy estimates predicted trial-by-changes in drift rate,

threshold, or both, while controlling for influences of expected reward rate on those same DDM

parameters. We found that higher levels of expected efficacy were associated with higher drift

rates ($b = 0.07$; 95% CrI [0.14, 0.43]; $p_{b < 0} = 0.00$) but were uncorrelated with threshold levels (b

$= -0.00$; 95% CrI [-0.04, 0.04]; $p_{b < 0} = 0.74$). Expected reward rate was not correlated with either

drift rate or threshold (Table S9). These results suggest that participants responded to changes in

performance efficacy by adjusting their attention to the task, rather than simply adjusting their

response threshold (i.e., becoming more or less impulsive).

Figure 5. Efficacy is learned in the same way in a modified task and influences a behavioral marker of control allocation. A. Regression weights for the influence of previous feedback type (efficacy and reward) on the subjective efficacy estimate. Error bars represent 50% and 95% highest density intervals. **B. Two learning rate model captures efficacy learning best.** Differences in BICs between the two learning rate model, and the intercept-only and one learning rate models. **C. Efficacy learning rates.** Positive and negative efficacy learning rates for each participant. **D. Higher model-based efficacy estimates predict better behavioral performance.** Higher efficacy estimates reduce reaction times (left) and improve accuracy (right). **E. Higher model-based efficacy estimates predict increased allocation of attention.** Parameter estimates from the drift diffusion model. Higher efficacy estimates increase drift rates, but not response caution (thresholds).

Discussion

To evaluate the expected benefits of investing cognitive control into a task, people need

to consider whether this control is necessary and/or sufficient for achieving their desired

outcome (i.e., whether these efforts are worthwhile). A critical determinant of the

worthwhileness of control is performance efficacy, the extent to which performance on a control-

demanding task matters for outcome attainment versus those outcomes being determined by

performance-unrelated factors. However, the mechanisms through which people estimate the

efficacy of their performance based on previous experience are largely unknown. Here, we

identified the computational and neural mechanism through which efficacy is learned and used to

guide the allocation of cognitive control. Across two experiments, we found that participants

dynamically updated expectations of the efficacy of their task performance (i.e., the likelihood

that this performance will determine reward attainment), and used those expectations to adjust

how much control they allocated. The feedback-based updating of efficacy was well-captured by

a prediction error-based learning algorithm. Model-based estimates of efficacy and efficacy

prediction errors were encoded by canonical neural signatures of effort allocation and belief

updating, respectively. Importantly, these findings cannot be explained by variability in reward,

as reward rate was held constant across efficacy levels, and the subjective reward rate was

controlled for statistically. Further, our model-based analysis revealed that people allocated more

control when they learned that they had more efficacy, extending our previous findings on

instructed efficacy (Frömer et al., 2021a). Taken together, our results uncover the mechanism

through which efficacy estimates are learned and used to inform mental effort investment within

a given task environment.

Previous research has characterized the learning algorithms people use to learn the reward

value of different states and actions in their environment (Gläscher et al. 2010; Sutton and Barto

2018). Recent theoretical (Jiang et al. 2014; Lieder et al. 2018; Verbeke and Verguts 2019) and

empirical (Bejjani et al. 2018; Otto and Daw 2019; Jiang et al. 2020; Bustamante et al. 2021 Jan

6) work has extended this research to show how similar algorithms guide learning and adaptation

of cognitive control under varying rewards and task demands within a given task environment.

Our findings extend this work further in several ways. First, we show that people leverage

weighted prediction errors when learning about the efficacy of task performance, independently

of potential reward and task difficulty. Second, we show that they update their efficacy

expectations differently depending on whether performance was more efficacious or less

efficacious than they expected, demonstrating a striking parallel with dual learning rate models

that have been found to prevail in research on reward learning (Collins and Frank 2013; Lefebvre

et al. 2017; Garrett and Daw 2020), including in our own reward rate data. Third, we show that

participants dynamically adjust their control allocation based on learned efficacy, just as they do

for learned rewards and task demands (Bugg et al. 2011; Jiang et al. 2014; Lieder et al. 2018).

Our neural findings build further on past research on learning and control adaptation. The

P3b component has been shown to track prediction-error based learning from action-relevant

outcome values (Fischer and Ullsperger 2013; Nassar et al. 2019; Lohse et al. 2020). Here we

show that this neural marker tracks learning about efficacy in the same way as it tracks learning

about rewards. We found increased P3b amplitudes when people experienced feedback about

outcome contingencies that was less expected given their current estimate of efficacy (e.g.,

expecting low efficacy, but getting performance-contingent feedback), relative to when these

contingencies were more expected (e.g., expecting low efficacy and getting random feedback).

Our additional finding that P3b amplitude was overall larger for efficacy compared to no efficacy

feedback demonstrates that our participants were not just tracking the frequency of the two types

of feedback, as predicted by an oddball account. Instead this finding suggests that they were

actively learning from the feedback.

Extending previous findings on cueing efficacy and/or reward (Schevernels et al. 2014b;

Frömer et al. 2021a), our CNV and behavioral results further show that participants used these

learned efficacy estimates to calibrate their effort and their performance. Notably, unlike in

previous work, our study shows effort-related changes in CNV amplitude entirely divorced from

perceptual cues, providing evidence that this activity truly reflects adjustments in control, rather

than reactive processing of features associated with the cue. Taken together, our findings suggest

that similar neural mechanisms underlie learning and control adaptation in response to variability

in one's efficacy in a task environment, as they do in response to variability in expected rewards

(Leng et al., 2021; Otto & Daw, 2019). By fitting behavioral data from this task to a drift

diffusion model (Study 2), we were able to further demonstrate that participants were adapting to

changes in expected efficacy by enhancing the processing of stimuli (i.e., increasing their rate of

evidence accumulation) - potentially via attentional control mechanisms – rather than by

adjusting their threshold for responding. This particular pattern of control adjustments was

predicted for the current task because performance-contingent rewards depended on responses

being *both* fast and accurate (as in Frömer et al., 2021a), but future work should test the

prediction that different control adjustments should emerge under different performance

contingencies (cf. Leng et al., 2021; Ritz et al., 2021).

Our findings build on previous research on how people learn about controllability of their

environment. Studies have examined the neural and computational mechanisms by which

humans and other animals learn about the contingencies between an action and its associated

outcome, and demonstrated that these learned action-outcomes contingencies influence which

actions are selected and how vigorously they are enacted (Dickinson and Balleine 1995;

Liljeholm et al. 2011; Manohar et al. 2017; Moscarello and Hartley 2017; Ly et al. 2019). Our

work extends this research into the domain of cognitive control, where the contingencies

between actions (i.e., control adjustments) and outcomes (e.g., reward) depend both on whether

control adjustments predicts better performance and whether better performance predicts better

outcomes (Shenhav et al., 2021). Learning control-outcome contingencies therefore requires

learning about how control states map onto performance (control efficacy) as well as how

performance maps onto outcomes (performance efficacy). By describing the mechanisms by

which people solve the latter part of this learning problem, and demonstrating that these are

comparable to those engaged during action-outcome learning, our study lays critical groundwork

for better understanding the links between selection of actions and control states.

Our efficacy-updating results are a reminder that many aspects of feedback are reflected in

prediction error signals (Langdon et al. 2018; Frömer et al. 2021b). In the present study, we

intentionally separated feedback about reward and efficacy to isolate the cognitive and neural

learning mechanisms associated with each. In doing so, we have taken an important first step

towards understanding the updating mechanisms underlying each. Further work is needed to

understand how they are inferred in more naturalistic settings, in which different forms of

feedback are often multiplexed, containing information about the values of actions that were

taken as well as about the features and structure of the environment (cf. Dorfman et al., 2019;

2021).

Another distinct element of more complex naturalistic environments is that the same

feedback can be used to evaluate multiple targets, internal ones, such as the selected response

and its predicted outcome, or external ones, such as the source of feedback/environment

(Carlebach and Yeung 2020). Such multi-level prediction error signals might for instance

explain, why despite close links between P3b and behavioral adaptation (Yeung and Sanfey

2004; Chase et al. 2011; Fischer and Ullsperger 2013), this link is context dependent and

attempts to link the P3b uniquely to behavioral updating have failed (Nassar et al., 2019).

Reinforcement learning, and predictive inference more generally, have been proposed to not only

support the selection of individual actions, but also extended sequences of actions and control

signals (Holroyd and Yeung 2012; Lieder et al. 2018). Alongside evaluations of actions and

environmental states, neural signatures of feedback-based learning could thus further reflect

evaluations of control signals, their quality or intensity. Given the many potential causes a given

outcome can have, and the flexibility that people have in how they use the feedback, it is easy to

see how feedback could be misattributed and lead to inaccurate beliefs about performance

efficacy. Such beliefs about environmental statistics can drive changes in feedback-processing

and behavioral adaptation, above and beyond the true statistics (Schiffer et al. 2017), and are thus

of particular importance for understanding some of the cognitive symptoms of mental disorders.

Understanding how efficacy estimates develop based on previous experiences is crucial for

understanding why people differ in how much cognitive control they allocate in different

situations (Shenhav et al.). People differ in their beliefs about how much control they have over

their environments (Leotti et al. 2010; Moscarello and Hartley 2017), and in their ability to

estimate their efficacy (Cohen et al. 2020). Further, many mental disorders, including depression

and schizophrenia, have been linked with one's estimates of their ability to control potential

outcomes in their environment, including through allocation of control (Huys and Dayan 2009;

Maier and Seligman 2016), and we have recently proposed that such changes can drive

impairments of motivation and control in those populations (Grahek et al. 2019). As we show in

this study, when people have learned to expect low efficacy, they will allocate less cognitive

control, which can manifest as apparent control deficits. The experimental and modeling

approach taken in our study helps uncover a more fine-grained view of how components of

motivation are learned and used to support the allocation of cognitive resources. In this way, our

study takes a first step toward a better computational and neural account of efficacy learning,

which can aid the understanding of individual differences in the willingness to exert mental

effort, as well as the development of interventions aimed at teaching individuals when these

efforts truly matter.

Figure S1. Parameter recovery study for the two learning rate models for the efficacy and reward learning. We simulated 200 subjects with the range of learning rates matched to the empirically observed range. The noise and the intercept parameters were fixed to match the empirically observed mean value. The number of trials and subjective estimates, as well as the sequence of efficacy and reward feedbacks were based on Experiment 1. **A. Learning rates for efficacy.** Simulated and recovered learning rates were highly correlated both in the case of positive ($r = 0.93$, $p<0.001$) and negative learning rates ($r = 0.94$, p<0.001). **B. Learning rates for reward.** Simulated and recovered learning rates were highly correlated both in the case of positive ($r = 0.88$, $p<0.001$) and negative learning rates ($r = 0.96$, $p<0.001$).

Figure S2. Efficacy drifts and the unsupervised clustering of the efficacy learning rates in Study 2. A. Efficacy drifts used in Study 2. The probability drifts of contingent feedback presented to the first (top) and second (bottom) half of the participants. **B. The results of the winning Gaussian mixture model for clustering of the efficacy learning rates.** One of the clusters (blue dots) included only the subjects with extremely small learning rates (all learning rates < 0.03; N=20; group 1 – blue dots). These subjects were excluded from the further analyses with the assumption that they did not pay attention to the feedbacks or that they were giving random answers to the probes. **C. Histograms of the positive and negative learning rates and their clusters.** For both then negative (left) and positive (right) learning rates cluster 1 included only the participants with very low learning rates.

Figure S3. Self-reported and model-based efficacy estimates for subjects excluded from Study 2 based on very low learning rates. Subjective efficacy estimates (grey) and model-based efficacy estimates for each of the 19 subjects identified to form a cluster due to very low learning rates (both learning rates < 0.03) based on the unsupervised clustering algorithm using Gaussian mixture models. These subjects had low variance in their self-reported efficacy estimates, or did not appear to update their efficacy estimates based on feedbacks, suggesting that they were not paying attention to the feedbacks, or always providing very similar efficacy estimates.

Table S1

Study 1 regression weights for the models predicting subjective efficacy and reward rate estimates based on efficacy and reward feedback 5 trials back.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)	BF$_{10}$
Subjective efficacy				
Intercept	0.52	0.49, 0.55	0.00	
T1 Performance-Based - Random	0.14	0.12, 0.16	0.00	>100
T2 Performance-Based - Random	0.12	0.10, 0.14	0.00	>100
T3 Performance-Based - Random	0.08	0.05, 0.10	0.00	>100
T4 Performance-Based - Random	0.03	0.01, 0.05	0.00	2.20
T5 Performance-Based - Random	0.03	0.01, 0.05	0.00	1.72
T1 Reward - No Reward	0.02	0.00, 0.04	0.02	0.37
T2 Reward - No Reward	0.03	0.01, 0.05	0.00	4.23
T3 Reward - No Reward	0.03	0.01, 0.05	0.01	0.95
T4 Reward - No Reward	0.02	0.00, 0.04	0.02	0.39
T5 Reward - No Reward	0.00	-0.02, 0.02	0.46	0.05
Subjective reward rate				
Intercept	0.50	0.46, 0.54	0.00	
T1 Reward - No Reward	0.19	0.17, 0.21	0.00	>100
T2 Reward - No Reward	0.16	0.14, 0.18	0.00	>100
T3 Reward - No Reward	0.09	0.07, 0.11	0.00	>100
T4 Reward - No Reward	0.05	0.03, 0.06	0.00	30.86
T5 Reward - No Reward	0.05	0.03, 0.07	0.00	>100
T1 Performance-Based - Random	-0.01	-0.03, 0.01	0.73	0.06
T2 Performance-Based - Random	0.00	-0.01, 0.02	0.33	0.05
T3 Performance-Based - Random	0.01	-0.01, 0.03	0.00	0.05
T4 Performance-Based - Random	0.02	-0.01, 0.03	0.20	0.08
T5 Performance-Based - Random	0.02	0.00, 0.03	0.04	0.20

Figure S4. Efficacy learning in Study 1 – the effect of different feedback probability drifts on learning.
A. The probability drift of contingent feedback presented to the first half of subjects. **B.** Learning rates for contingent and random feedback for the first half of subjects. Learning rates for the contingent and random feedback did not differ (b = 0.01; 95% CrI [-0.06, 0.08]; $p_{b<0}$ = 0.38; BF_{10} = 0.09). **C.** The inversed probability drift of contingent feedback presented to the second half of participants. **D.** Learning rates for contingent and random feedback did not differ (b = 0.02; 95% CrI [-0.06, 0.11]; $p_{b<0}$ = 0.31; BF_{10} = 0.10).

Figure S5. The influence of efficacy estimates on efficacy feedback processing and proactive cognitive control allocation. A. Means for the P3b amplitude in response to performance-based and random feedbacks across each quartile of the model-based efficacy estimates calculated for each subject. The P3b amplitude in response to performance-based feedback is higher when efficacy estimates are lower, suggesting more updating based on feedback. Error bars represent standard errors of the mean. **B.** Means for the CNV amplitude pre target onset across quartiles of the model-based efficacy estimates calculated for each subject. The CNV amplitude is more negative (more proactive control allocation) when efficacy estimates are higher. Error bars represent standard errors of the mean.

Figure S6. The influence of efficacy estimates on behavioral performance. A. Means for the reaction times (left) and accuracy (right) across quartiles of model-based efficacy estimates in Experiment 1. Reaction times are faster when efficacy estimates are higher, while there is no consistent pattern in accuracy data. Error bars represent standard errors of the mean. **B.** Means for the reaction times (left) and accuracy (right) across quartiles of model-based efficacy estimates in Experiment 2. Participants are faster to respond and more accurate when efficacy estimates are high relative to low. Error bars represent standard errors of the mean.

Figure S7. Behavioral effects of congruency in Study 1 and Study 2. A. In comparison to the neutral trials in Study 1, participants were faster to respond to congruent ($b = 9.95$; 95% CrI [3.33, 16.59]; $p_{b<0} = 0.00$; BF = 1.60) and slower to respond to incongruent trials ($b = 61.24$; 95% CrI [49.50, 72.59]; $p_{b<0} = 0.00$; BF = 6.73). **B.** In comparison to the neutral trials in Study 1, participants were equally likely to be correct when responding to congruent ($b = -0.11$; 95% CrI [-0.29, 0.06]; $p_{b>0} = 0.091$; BF = 0.01) and less likely to be correct when responding to incongruent trials ($b = -0.48$; 95% CrI [-0.64, -0.33]; $p_{b>0} = 0.00$; BF > 8.22). The regression weights for the accuracy analysis are in log odds. **C.** In Study 2 participants were slower to respond to incongruent relative to congruent trials ($b = 47.48$; 95% CrI [40.69, 52.22]; $p_{b<0} = 0.001$; BF = 50). **D.** In Study 2 participants were less accurate when responding to incongruent compared to congruent trials ($b = -0.48$; 95% CrI [-0.57, -0.39]; $p_{b<0} = 1$; BF > 100).

Table S2

Study 1 regression weights for the models predicting the P3b to efficacy feedback.

Parameter	Estimate	95% Credible interval	Posterior probability ($p<0$)	BF_{10}
Efficacy feedback				
Intercept	1.38	0.67, 2.09	0.00	
Model-based efficacy	-0.29	-1.56, 0.97	0.68	0.23
Efficacy feedback type (PB-R)	0.86	0.42, 1.31	0.00	30.86
Model-based reward rate	-0.19	-1.50, 1.18	0.62	0.23
Model-based efficacy × Efficacy feedback type	-2.40	-4.07, -0.74	1.00	24.40
Model-based efficacy slope for Performance-Based feedback	-1.49	-2.72, -0.27	0.97	1.52
Model-based efficacy slope for Random feedback	0.91	-0.34, 2.21	0.12	0.44
Predictions errors and learning rates				
Intercept	1.31	0.51, 2.09	0.00	
Unsigned prediction error	1.25	0.35, 2.15	0.01	5.84
Learning rate	2.00	-2.21, 6.04	0.17	1.08
Model-based reward rate	-0.09	-1.48, 1.34	0.55	0.23
Unsigned prediction error ×Learning rate	0.57	-4.87, 6.14	0.41	0.91

Table S3

Study 1 regression weights for the models predicting reaction times and accuracy based on efficacy and reward rate estimates.

Parameter	Estimate	95% Credible interval	Posterior probability $(p<0)$	BF_{01}
Reaction times				
Intercept	655.37	635.09, 675.51	0.00	
Facilitation	9.95	3.33, 16.59	0.00	1.60
Interference	61.24	49.50, 72.59	0.00	6.73
Model-based efficacy	-16.00	-34.91, 2.96	0.95	1.68
Model-based reward rate	-1.56	-27.66, 24.04	0.54	1.36
Accuracy				
Intercept	1.56	1.33, 1.80	0.00	
Facilitation	-0.11	-0.29, 0.06	0.91	0.01
Interference	-0.48	-0.64, -0.33	1.00	8.22
Model-based efficacy	0.12	-0.20, 0.44	0.23	1.99
Model-based reward rate	-0.36	-0.73, 0.03	0.97	0.29

Table S4

Study 1 regression weights for the models predicting the CNV based on efficacy and reward rate.

Parameter	Estimate	95% Credible interval	Posterior probability $(p<0)$	BF_{01}
CNV				
Intercept	-0.37	-0.58, -0.16	0.00	
Model-based efficacy	-0.35	-0.85, 0.16	0.09	2.73
Model-based reward rate	-0.09	-0.72, 0.55	0.38	2.19

Table S5

Study 1 regression weights for the models predicting reaction times and accuracy
based on the CNV.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)	BF$_{10}$
Reaction times				
Intercept	655.28	635.44, 675.91	0.00	
CNV amplitude	11.41	8.09, 14.72	0.00	>100
Accuracy				
Intercept	1.33	1.11, 1.53	0.00	
CNV amplitude	-0.07	-0.12, -0.01	0.99	3.14

Figure S8. Reward rate learning – model-based analyses for Study 1. A. Regression weights for the influence of the current (t) and previous feedbacks on the subjective estimates of reward rate (left) and efficacy (right). Reward rate estimates are strongly predicted by the previous reward feedbacks, but not efficacy feedbacks. The reverse is true for the subjective efficacy estimates. Error bars represent 50% and 95% highest density intervals. **B.** Model comparison between the fitted learning models for the reward rate learning model. **C.** Positive and negative learning rate estimates for all subjects for the reward rate learning model. Negative learning rates were numerically, but not statistically larger than positive ones ($b = 0.01$; 95% CrI [-0.07, 0.09]; $p_{b < 0} = 0.38$; $BF_{10} = 0.09$)

ERP analyses of the reward rate feedback in Study 1

To investigate the effect of the model-based reward rate estimates on the processing of

reward, we performed complementary analyses on the P3b related to the processing of the

reward feedback (Figure S9A). We expected that a similar learning mechanism should operate

on both reward and efficacy feedback, but that the neural markers of feedback processing should

be sensitive to different model-based estimates. Thus, we expected the reward-locked P3b to be

sensitive to reward-rate estimates, but not to efficacy estimates.

Our results showed larger P3b amplitudes to no reward compared to reward feedback

(note that reward rate was approximately .80 and negative feedback less expected overall), and

the processing of the reward feedback was influenced by reward-rate estimates, although to

lesser extent than for the P3b (Figure S9B; Table S6). Probing the effects of learning directly, we

found robust effects of the prediction errors and learning rates (Figure S9C). The P3b to reward

feedback was not influenced by the model-based efficacy estimates in either of these analyses.

Figure S9. Learning reward rate estimate from reward feedback in Study 1. A. ERP average for the P3b locked to the onset of reward feedback. **B.** Interaction between reward feedback and the model-based reward rate estimate. **C.** The effects of unsigned prediction errors (left) and learning rates (right) on the P3b.

Table S6

Study 1 regression weights for the models predicting the P3b to reward feedback.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)	BF$_{10}$
Efficacy feedback				
Intercept	0.93	0.08, 1.74	0.02	
Model-based efficacy	0.45	-0.45, 1.32	0.16	0.15
Reward feedback type (R-NR)	-0.35	-0.90, 0.18	0.90	0.12
Model-based reward rate	-0.15	-1.55, 1.23	0.59	0.14
Model-based reward ×Reward feedback type	-1.26	-3.73, 1.19	0.85	0.41
Model-based efficacy slope for No Reward feedback	0.48	-1.34, 2.34	0.87	0.23
Model-based efficacy slope for Reward feedback	-0.78	-1.96, 0.39	0.32	0.21
Predictions errors and learning rates				
Intercept	0.89	0.15, 1.64	0.01	
Unsigned prediction error	1.09	0.38, 1.81	0.00	4.82
Learning rate	6.68	-0.07, 12.99	0.03	4.72
Model-based efficacy	0.39	-0.47, 1.24	0.19	0.13
Unsigned prediction error ×Learning rate	0.94	-8.06, 9.71	0.42	0.92

Figure S10. Efficacy and reward learning – model-based analyses for Study 2. A. Regression weights for the influence of the current (t) and previous feedbacks on the subjective estimates of reward rate (left) and efficacy (right). Reward rate estimates are strongly influenced by the previous reward feedbacks, and the reverse is true for the subjective efficacy estimates. Error bars represent 50% and 95% highest density intervals. **B.** Model comparison between the fitted learning models for the reward learning model. **C.** Positive and negative learning rate estimates for all subjects for the reward learning model. Positive learning rates were numerically, but not statistically larger than negative ones ($b = 0.02$; 95% CrI [-0.01, 0.05]; $p_{b < 0} = 0.11$; $BF_{10} = 0.06$)

Table S7

Study 2 regression weights for the models predicting subjective efficacy and reward rate estimates based on efficacy and reward feedback 5 trials back.

Parameter	Estimate	95% Credible interval	Posterior probability ($p<0$)	BF_{10}
Subjective efficacy				
Intercept	0.43	0.37, 0.49	0.00	
T1 Performance-Based - Random	0.17	0.16, 0.19	0.00	>100
T2 Performance-Based - Random	0.15	0.13, 0.18	0.00	>100
T3 Performance-Based - Random	0.10	0.08, 0.12	0.00	>100
T4 Performance-Based - Random	0.06	0.04, 0.08	0.00	>100
T5 Performance-Based - Random	0.04	0.02, 0.06	0.00	>100
T1 Reward - No Reward	0.06	0.02, 0.10	0.00	5.42
T2 Reward - No Reward	0.06	0.01, 0.11	0.01	2.23
T3 Reward - No Reward	0.03	-0.02, 0.07	0.11	0.23
T4 Reward - No Reward	0.00	-0.05, 0.04	0.54	0.11
T5 Reward - No Reward	0.06	0.01, 0.10	0.01	1.69
Subjective reward rate				
Intercept	0.36	0.29, 0.43	0.00	
T1 Reward - No Reward	0.21	0.17, 0.25	0.00	>100
T2 Reward - No Reward	0.13	0.09, 0.17	0.00	>100
T3 Reward - No Reward	0.12	0.07, 0.16	0.00	>100
T4 Reward - No Reward	0.03	-0.01, 0.06	0.09	0.21
T5 Reward - No Reward	0.09	0.05, 0.13	0.00	>100
T1 Performance-Based - Random	0.05	0.03, 0.07	0.00	0.06
T2 Performance-Based - Random	0.05	0.03, 0.06	0.00	0.05
T3 Performance-Based - Random	0.00	-0.02, 0.02	0.51	0.05
T4 Performance-Based - Random	0.03	0.02, 0.05	0.00	0.08
T5 Performance-Based - Random	0.02	0.00, 0.03	0.03	0.20

Table S8

Study 2 regression weights for the models predicting reaction times and accuracy based on efficacy and reward rate estimates. Bayes Factors calculated only for the parameters with informative priors. Posteriors from Study 1 were used as priors for the effects of congruency, efficacy, and reward.

Parameter	Estimate	95% Credible interval	Posterior probability ($p<0$)	BF_{01}
Reaction times				
Intercept	643.02	625.42, 661.81	0.00	
Incongruent-Congruent	47.48	40.69, 55.22	0.00	0.02
Model-based efficacy	-10.25	-19.86, -0.20	0.98	2.33
Model-based reward	-0.17	-21.15, 20.34	0.51	1.24
Interval length	-1.76	-2.73, -0.83	1.00	
Interval congruency	-5.34	-6.69, -4.00	1.00	
Accuracy				
Intercept	2.35	2.16, 2.52	0.00	
Incongruent-Congruent	-0.48	-0.57, -0.39	1.00	>100
Model-based efficacy	0.23	0.03, 0.43	0.01	>100
Model-based reward	-0.29	-0.57, -0.01	0.98	1.07
Interval length	-0.13	-0.16, -0.10	1.00	
Interval congruency	0.04	0.00, 0.08	0.03	

Table S9
Drift Diffusion model estimates.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)
Drift rate			
Congruent-Incongruent	0.82	0.74, 0.92	0.00
Model-based efficacy	0.29	0.14, 0.40	0.00
Model-based reward	0.20	-0.39, 0.04	0.50
Threshold			
Model-based efficacy	-0.00	-0.04, 0.04	0.57
Model-based reward rate	-0.04	-0.14, 0.05	0.81

768   **Acknowledgements**

This work was supported by the Special Research Fund (BOF) of Ghent University [grant

#01D02415] (I.G.), the Research Foundation Flanders (FWO) travel grant [grant #V432718N]

(I.G.), a Center of Biomedical Research Excellence grant P20GM103645 from the National

Institute of General Medical Sciences (A.S.), the Alfred P. Sloan Foundation Research Fellowship

in Neuroscience (A.S.), and an NSF Graduate Research Fellowship (M.P.F). The funding sources

were not involved in the study design; collection, analysis, and interpretation of data; writing of

the report; and decision to submit the article for publication. We would like to thank Natalie

Knowles and Hattie Xu for help with data collection for Study 1, Peyton Strong for help with data

collection for Study 2, Carolyn Dean Wolf and Liz Cory for help with programming the Study 1

task, Harrison Ritz for help with constructing the efficacy drifts, and Xiamin Leng for advice on

fitting the drift-diffusion models.

781   **Data availability**

Analysis scripts are available on this link: https://github.com/igrahek/LFXC_EEG_2022.git Please

contact the authors for raw data.

References

Atkinson JW, Feather NT, Others. 1966. A theory of achievement motivation. Wiley New York.

Balleine, B. W., & O'doherty, J. P. 2010. Human and rodent homologies in action control:
corticostriatal determinants of goal-directed and habitual action. *Neuropsychopharmacology*,
35(1), 48-69.

Bejjani C, Zhang Z, Egner T. 2018. Control by association: Transfer of implicitly primed
attentional states across linked stimuli. Psychon Bull Rev. 25(2):617–626. doi:10.3758/s13423-
018-1445-6.

Botvinick MM, Cohen JD. 2014. The computational and neural basis of cognitive control:
Charted territory and new frontiers. Cogn Sci. 38(6):1249–1285. doi:10.1111/cogs.12126.

Brainard DH. 1997. The Psychophysics Toolbox. Spat Vis. 10(4):433–436.

Bugg JM, Jacoby LL, Chanani S. 2011. Why it is too early to lose control in accounts of item-
specific proportion congruency effects. J Exp Psychol Hum Percept Perform. 37(3):844–859.
doi:10.1037/a0019957.

Bürkner P-C. 2016. brms: An R package for Bayesian multilevel models using Stan. J Stat
Softw. 80(1):1–28.

Bürkner P-C. 2017. Advanced Bayesian Multilevel Modeling with the R Package brms.
arXiv:170511123.

Bustamante L, Lieder F, Musslick S, Shenhav A, Cohen J. 2021 Jan 6. Learning to Overexert
Cognitive Control in a Stroop Task. Cogn Affect Behav Neurosci. doi:10.3758/s13415-020-
00845-x.

Carlebach N, Yeung N. 2020. Flexible use of confidence to guide advice requests. PsyArXiv.

Carpenter B, Gelman A, Hoffman M, Lee D, Goodrich B, Betancourt M, Brubaker MA, Guo J,
Li P, Riddell A. 2016. Stan: A probabilistic programming language. J Stat Softw. 2(20):1–37.

Chambon V, Théro H, Vidal M, Vandendriessche H, Haggard P, Palminteri S. 2020. Information
about action outcomes differentially affects learning from self-determined versus imposed
choices. Nat Hum Behav. doi:10.1038/s41562-020-0919-5.

Chase HW, Swainson R, Durham L, Benham L, Cools R. 2011. Feedback-related negativity
codes prediction error but not behavioral adjustment during probabilistic reversal learning. J
Cogn Neurosci. 23(4):936–946. doi:10.1162/jocn.2010.21456.

Chiu, Y. C., & Egner, T. (2019). Cortical and subcortical contributions to context-control
learning. Neurosc & Biobeh Rev, 99: 33-41.

Cohen A, Nussenbaum K, Dorfman HM, Gershman SJ, Hartley CA. 2020. The rational use of
causal inference to guide reinforcement learning strengthens with age. Npj Sci Learn.

Collins AGE, Frank MJ. 2013. Cognitive control over learning: creating, clustering, and
generalizing task-set structure. Psychol Rev. 120(1):190–229. doi:10.1037/a0030852.

Collins AGE, Frank MJ. 2014. Opponent actor learning (OpAL): Modeling interactive effects of
striatal dopamine on reinforcement learning and choice incentive. Psychol Rev. 121(3):337–66.
doi:10.1037/a0037015.

Delorme A, Makeig S. 2004. EEGLAB: an open sorce toolbox for analysis of single-trail EEG
dynamics including independent component anlaysis. J Neurosci Methods. 134:9–21.
doi:10.1016/j.jneumeth.2003.10.009.

Dickinson A, Balleine B. 1995. Motivational Control of Instrumental Action. Curr Dir Psychol
Sci. 4(5):162–167. doi:10.1111/1467-8721.ep11512272.

Dorfman, H. M., Bhui, R., Hughes, B. L., & Gershman, S. J. 2019. Causal inference about good
and bad outcomes. Psychological science, 30(4): 516-525.

Dorfman, H. M., Tomov, M. S., Cheung, B., Clarke, D., Gershman, S. J., & Hughes, B. L.
(2021). Causal inference gates corticostriatal learning. Journal of Neuroscience, 41(32): 6892-
6904.

Eldar E, Hauser TU, Dayan P, Dolan RJ. 2016. Striatal structure and function predict individual
biases in learning to avoid pain. Proc Natl Acad Sci U S A. 113(17):4812–4817.
doi:10.1073/pnas.1519829113.

Fischer AG, Ullsperger M. 2013. Real and fictive outcomes are processed differently but
converge on a common adaptive mechanism. Neuron. 79(6):1243–1255.
doi:10.1016/j.neuron.2013.07.006.

Frömer R, Lin H, Dean Wolf CK, Inzlicht M, Shenhav A. 2021a. Expectations of reward and
efficacy guide cognitive control allocation. Nat Commun. 12(1). doi:10.1038/s41467-021-21315-
z.

Frömer, R., Nassar, M. R., Bruckner, R., Stürmer, B., Sommer, W., & Yeung, N. 2021b.
Response-based outcome predictions and confidence regulate feedback processing and learning.
Elife, 10, e62825.

Frömer R, Maier M, Rahman RA. 2018. Group-level EEG-processing pipeline for flexible single
trial-based analyses including linear mixed models. Front Neurosci. 12(48).
doi:10.3389/fnins.2018.00048.

Frömer R, Stürmer B, Sommer W. 2016. (Don't) Mind the effort: Effects of contextual
interference on ERP indicators of motor preparation. Psychophysiology. 53(10):1577–1586.
doi:10.1111/psyp.12703.

Garrett N, Daw ND. 2020. Biased belief updating and suboptimal choice in foraging decisions.
Nat Commun. 11(1):1–12. doi:10.1038/s41467-020-16964-5.

Gelman A, Rubin DB. 1992. Inference from Iterative Simulation Using Multiple Sequences. Stat
Sci. 7(4):457–472. doi:10.1214/ss/1177011136.

Gläscher J, Daw N, Dayan P, O'Doherty JP. 2010. States versus rewards: Dissociable neural
prediction error signals underlying model-based and model-free reinforcement learning. Neuron.
66(4):585–595. doi:10.1016/j.neuron.2010.04.016.

Grahek I, Shenhav A, Musslick S, Krebs RM, Koster EHW. 2019. Motivation and cognitive
control in depression. Neurosci Biobehav Rev. 102(March):371–381.
doi:10.1016/j.neubiorev.2019.04.011.

Grent-'t-Jong T, Woldorff MG. 2007. Timing and sequence of brain activity in top-down control
of visual-spatial attention. PLoS Biol. 5(1):0114–0126. doi:10.1371/journal.pbio.0050012.

Holroyd CB, Yeung N. 2012. Motivation of extended behaviors by anterior cingulate cortex.
Trends Cogn Sci. 16(2):122–128. doi:10.1016/j.tics.2011.12.008.

Huys QJM, Dayan P. 2009. A Bayesian formulation of behavioral control. Cognition.
113(3):314–328. doi:10.1016/j.cognition.2009.01.008.

Ille N, Berg P, Scherg M. 2002. Artifact Correction of the Ongoing EEG Using Spatial Filters
Based on Artifact and Brain Signal Topographies. J Clin Neurophysiol. 19(2):113–124.
doi:10.1097/00004691-200203000-00002.

Jiang J, Bramão I, Khazenzon A, Wang SF, Johansson M, Wagner AD. 2020. Temporal
Dynamics of Memory-guided Cognitive Control and Generalization of Control via Overlapping
Associative Memories. J Neurosci. 40(10):2343–2356. doi:10.1523/JNEUROSCI.1869-19.2020.

Jiang J, Heller K, Egner T. 2014. Bayesian modeling of flexible cognitive control. Neurosci
Biobehav Rev. 46(P1):30–43. doi:10.1016/j.neubiorev.2014.06.001.

Kleiner M, Brainard D, Pelli D. 2007. What's new in Psychtoolbox-3. In: Perception 36 ECVP
Abstract Supplement.

Kool W, Botvinick M. 2018. Mental labour. Nat Hum Behav. doi:10.1038/s41562-018-0401-9.

Krebs, R. M., Boehler, C. N., Roberts, K. C., Song, A. W., & Woldorff, M. G. 2012. The
involvement of the dopaminergic midbrain and cortico-striatal-thalamic circuits in the integration
of reward prospect and attentional task demands. *Cereb cortex*, 22(3): 607-615.
881
Kruschke JK. 2013. Bayesian estimation supersedes the T test. J Exp Psychol Gen. 142(2):573–
588. doi:10.1037/a0029177.

Langdon AJ, Sharpe MJ, Schoenbaum G, Niv Y. 2018. Model-based predictions for dopamine.
Curr Opin Neurobiol. 49:1–7. doi:10.1016/j.conb.2017.10.006.

Lefebvre G, Lebreton M, Meyniel F, Bourgeois-Gironde S, Palminteri S. 2017. Behavioural and
neural characterization of optimistic reinforcement learning. Nat Hum Behav. 1(4):1–9.
doi:10.1038/s41562-017-0067.

Leng, X., Yee, D., Ritz, H., & Shenhav, A. 2021. Dissociable influences of reward and
punishment on adaptive cognitive control. PLoS computational biology, 17(12), e1009737.

Leotti LA, Iyengar SS, Ochsner KN. 2010. Born to choose: The origins and value of the need for
control. Trends Cogn Sci. 14(10):457–463. doi:10.1016/j.tics.2010.08.001.

Lieder F, Shenhav A, Musslick S, Griffiths TL. 2018. Rational metareasoning and the plasticity
of cognitive control. Plos Comput Biol. 14(4). doi:10.13140/RG.2.2.24500.14721.

Ligneul, R., Mainen, Z. F., Ly, V., & Cools, R. 2022. Stress-sensitive inference of task
controllability. *Nat Hum Beh*. 1-11.

Liljeholm M, Tricomi E, O'Doherty JP, Balleine BW. 2011. Neural Correlates of Instrumental
Contingency Learning: Differential Effects of Action-Reward Conjunction and Disjunction. J
Neurosci. 31(7):2474–2480. doi:10.1523/JNEUROSCI.3354-10.2011.

Lohse KR, Miller MW, Daou M, Valerius W, Jones M. 2020. Dissociating the contributions of
reward-prediction errors to trial-level adaptation and long-term learning. Biol Psychol.
149(January 2019). doi:10.1016/j.biopsycho.2019.107775.

Ly V, Wang KS, Bhanji J, Delgado MR. 2019. A Reward-Based Framework of Perceived
Control. Front Neurosci. 13(February):1–11. doi:10.3389/fnins.2019.00065.

Maier SF, Seligman MEP. 2016. Learned helplessness at fifty: Insights from neuroscience.
Psychol Rev. 123(4):1–19. doi:10.1037/rev0000033.

Manohar SG, Chong TTJ, Apps MAJ, Batla A, Stamelou M, Jarman PR, Bhatia KP, Husain M.
2015. Reward Pays the Cost of Noise Reduction in Motor and Cognitive Control. Curr Biol.
25(13):1707–1716. doi:10.1016/j.cub.2015.05.038.

Manohar SG, Finzi RD, Drew D, Husain M. 2017. Distinct Motivational Effects of Contingent
and Noncontingent Rewards. Psychol Sci. 28(7):1016–1026. doi:10.1177/0956797617693326.

Morris, R. W., Dezfouli, A., Griffiths, K. R., Le Pelley, M. E., & Balleine, B. W. 2022. The
neural bases of action-outcome learning in humans. *J Neuro*, 42(17), 3636-3647.

Moscarello JM, Hartley CA. 2017. Agency and the Calibration of Motivated Behavior. Trends
Cogn Sci. 21(10):725–735. doi:10.1016/j.tics.2017.06.008.

Nagase AM, Onoda K, Foo JC, Haji T, Akaishi R, Yamaguchi S, Sakai K, Morita K. 2018.
Neural mechanisms for adaptive learned avoidance of mental effort. J Neurosci. 38(10):2631–
2651. doi:10.1523/JNEUROSCI.1995-17.2018.

Nalborczyk L, Bürkner P-C. 2019. An Introduction to Bayesian Multilevel Models Using brms:
A Case Study of Gender Effects on Vowel Variability in Standard Indonesian. J Speech Lang
Hear Res.

Nassar MR, Bruckner R, Frank MJ. 2019. Statistical context dictates the relationship between
feedback-related EEG signals and learning. eLife. 8:1–26. doi:10.7554/eLife.46975.

Nigbur, R., Schneider, J., Sommer, W., Dimigen, O., & Stürmer, B. (2015). Ad-hoc and context-
dependent adjustments of selective attention in conflict control: An ERP study with visual
probes. NeuroImage, 107: 76-84.

Niv Y, Edlund JA, Dayan P, O'Doherty JP. 2012. Neural prediction errors reveal a risk-sensitive
reinforcement-learning process in the human brain. J Neurosci. 32(2):551–562.
doi:10.1523/JNEUROSCI.5498-10.2012.

Norton, K. G., & Liljeholm, M. 2020. The Rostrolateral Prefrontal Cortex Mediates a Preference
for High-Agency Environments. J Neurosci, 40(22): 4401-4409.

Otto AR, Daw ND. 2019. The opportunity cost of time modulates cognitive effort.
Neuropsychologia. 123:92–105. doi:10.1016/j.neuropsychologia.2018.05.006.

Parro, C., Dixon, M. L., & Christoff, K. 2018. The neural basis of motivational influences on
cognitive control. Human Brain Mapping, 39(12), 5097–5111

Pelli DD. 1997. The VideoToolbox software for visual psychophysics: transforming numbers
into movies. Spat Vis. 10(4):437–442.

R Core Team. 2017. R: A Language and Environment for Statistical Computing.

Ritz, H., Leng, X., & Shenhav, A. (2022). Cognitive control as a multivariate optimization
problem. Journal of Cognitive Neuroscience, 1-23.

Rutledge RB, Skandali N, Dayan P, Dolan RJ. 2014. A computational and neural model of
momentary subjective well-being. Proc Natl Acad Sci. 111(33):12252–12257.
doi:10.1073/pnas.1407535111.

Schevernels H, Krebs RM, Santens P, Woldorff MG, Boehler CN. 2014a. Task preparation
processes related to reward prediction precede those related to task-difficulty expectation.
NeuroImage. 84:639–647. doi:10.1016/j.neuroimage.2013.09.039.

Schevernels H, Krebs RM, Santens P, Woldorff MG, Boehler CN. 2014b. Task preparation
processes related to reward prediction precede those related to task-difficulty expectation.
NeuroImage. 84:639–647. doi:10.1016/j.neuroimage.2013.09.039.

Schiffer AM, Siletti K, Waszak F, Yeung N. 2017. Adaptive behaviour and feedback processing
integrate experience and instruction in reinforcement learning. NeuroImage. 146:626–641.
doi:10.1016/j.neuroimage.2016.08.057.

Shenhav A, Botvinick M, Cohen J. 2013. The expected value of control: An integrative theory of
anterior cingulate cortex function. Neuron. 79(2):217–240. doi:10.1016/j.neuron.2013.07.007.

Shenhav A, Musslick S, Lieder F, Kool W, Griffiths TL, Cohen JD, Botvinick MM. 2017.
Toward a Rational and Mechanistic Account of Mental Effort. Annu Rev Neurosci. 40(1):99–
124. doi:10.1146/annurev-neuro-072116-031526.

Shenhav A, Prater Fahey M, Grahek I. 2021. Decomposing the motivation to exert mental effort.
Curr Dir Psych Sci. 30(4), 307-314.

Sutton RS, Barto AG. 2018. Reinforcement Learning: An Introduction. MIT Press.

Vassena, E., Silvetti, M., Boehler, C. N., Achten, E., Fias, W., & Verguts, T. 2014. Overlapping
neural systems represent cognitive effort and reward anticipation. *PLoS One*, 9(3).
Verbeke P, Verguts T. 2019. Learning to synchronize: How biological agents can couple neural
task modules for dealing with the stability-plasticity dilemma. PLoS Comput Biol. 15(8):1–25.
doi:10.1371/journal.pcbi.1006604.

Verguts T, Vassena E, Silvetti M. 2015. Adaptive effort investment in cognitive and physical
tasks: a neurocomputational model. Front Behav Neurosci. 9(March).
doi:10.3389/fnbeh.2015.00057.

Wagenmakers EJ, Lodewyckx T, Kuriyal H, Grasman R. 2010. Bayesian hypothesis testing for
psychologists: A tutorial on the Savage-Dickey method. Cognit Psychol. 60(3):158–189.
doi:10.1016/j.cogpsych.2009.12.001.

Yeung N, Sanfey AG. 2004. Independent coding of reward magnitude and valence in the human
brain. J Neurosci. 24(28):6258–6264. doi:10.1523/JNEUROSCI.4537-03.2004.

975