Learning when effort matters: Neural dynamics underlying updating and adaptation to changes in

performance efficacy

Ivan Grahek[1†*], Romy Frömer[1†], & Amitai Shenhav[1]

1. *Department of Cognitive, Linguistic, & Psychological Sciences and Carney Institute for*

Brain Science, Brown University, Providence, RI 02912, USA

[†] These authors have contributed equally to the study.

* Corresponding author. E-mail address: ivan_grahek@brown.edu

Abstract

To determine how much cognitive control to invest in a task, people need to consider whether exerting control *matters* for obtaining potential rewards. In particular, they need to account for the efficacy of their performance – the degree to which potential rewards are determined by their performance or by independent factors (e.g., random chance). Yet it remains unclear how people learn about their *performance efficacy* in a given environment. Here, we examined the neural and computational mechanisms through which people (a) learn and dynamically update efficacy expectations in a changing environment, and (b) proactively adjust control allocation based on their current efficacy expectations. We recorded EEG in 40 participants performing an incentivized cognitive control task, while their performance efficacy (the likelihood that reward for a given trial would be determined by performance or at random) dynamically varied over time. We found that participants continuously updated their self-reported efficacy expectations based on recent feedback, and that these updates were well described by a standard prediction error-based reinforcement learning algorithm. Paralleling findings on updating of expected rewards, we found that model-based estimates of *efficacy prediction errors* were encoded by the feedback-related P3b. Updated expectations of efficacy in turn influenced levels of effort exerted on subsequent trials, reflected in greater proactive control (indexed by the contingent negative variation [CNV]) and improved performance when participants expected their performance to be more efficacious. These findings demonstrate that learning and adaptation to the efficacy of one's environment is underpinned by similar computations and neural mechanisms as are involved in learning about potential reward.

Keywords: Motivation; Cognitive control; Learning; Performance efficacy; Reward rate; EEG

Cognitive control is critical for achieving most goals, but it is effortful (Botvinick &

Cohen, 2014; Shenhav et al., 2017). Determining whether and how much control to invest in a

task requires a person to weigh these effort costs against the potential benefits of control (Kool &

Botvinick, 2018; Manohar et al., 2015; Verguts et al., 2015). These potential benefits not only

reflect the amount of reward that can be expected for performing well on a task but also, and

critically, the extent to which performance matters for obtaining those rewards (the *efficacy* of

performance) (Frömer et al., 2020a; Shenhav et al., submitted; Shenhav et al., 2013). For

instance, a student may be more motivated to invest effort into writing an essay for a competition

if they believe that this essay will play an important role in determining their success than if they

believe that the competition will be primarily determined by performance-unrelated factors (e.g.,

social status, idiosyncratic criteria). The amount of effort we invest in a task is therefore

substantially determined by how much efficacy we expect to have with that task. Such efficacy

expectations must be derived from past experiences that indicated how worthwhile our efforts

were in a given environment. Understanding how people come to learn when their efforts are

efficacious is critically important for understanding how they come to be more or less motivated

in certain contexts. However, in spite of its importance, relatively little is known about the

mechanisms by which efficacy is learned.

Past work has provided critical insights into related aspects of how people adapt control

based on learned incentives. For instance, a substantial body of research has examined how

people learn about the amount and frequency of reward in their environment (Daw & Tobler,

2013; Dolan & Dayan, 2013; Niv et al., 2006), including the extent to which learning differs

depending on the valence of feedback (e.g., whether an outcome was better or worse than

expected; Chambon et al., 2020; Collins & Frank, 2014; Garrett & Daw, 2020; Lefebvre et al.,

2017). Recent work has shown that people not only adapt their actions to these changing

rewards, but also their control states (Lieder et al., 2018). People allocate more or less cognitive

control to a task depending on the reward they have come to expect based on recent trials

(Bustamante et al., in press; Otto & Daw, 2019). Research has shown that people similarly adapt

their control based on the expected difficulty of a given task (for a review see: Chiu & Egner,

2019). Behavioral and electrophysiological evidence shows that people allocate more control

when, based on previous learning, they expect a task to be more difficult (Bejjani et al., 2018;

Bugg et al., 2011; Jiang et al., 2020; Nigbur et al., 2015). Research into these difficulty-related

control adaptations provides insight into how people learn about and adapt to their efficacy for a

given task, in that the same level of control is less efficacious for achieving reward as task

difficulty increases. However, because greater task difficulty also makes it more likely that an

individual will perform poorly on the task, this work also conflates the learning of efficacy with

the learning of expected (negative) outcomes. While these may often covary, they are dissociable

– for instance, performance-unrelated factors (such as those described above) decrease one's

efficacy but can either improve or worsen a person's likelihood of success. Existing work has yet

to tease these apart experimentally to isolate the mechanisms by which we learn about and adapt

to our efficacy.

We recently showed how the expected efficacy of mental effort can be experimentally

dissociated from expectation of reward and task difficulty (Frömer et al., 2020a). Participants

performed a cognitive control task (Stroop) and we varied the amount of reward they could

obtain on a given trial as well as whether the rewards were determined by their performance

(high efficacy) or whether the rewards were determined at random (low efficacy). We showed

that people allocate more control when they expect to have high compared to low efficacy,

reflected in higher amplitudes of an EEG index of proactive control (the contingent negative

variation [CNV]) and in improved performance. However, unlike in most real-world examples,

in this study participants were explicitly cued with which level of efficacy to expect prior to each

trial, the efficacy associated with a cue was fixed through the session, and efficacy levels were

always at one of two extremes. Therefore, there are two key open questions which are

unaddressed to date: how do people learn about their efficacy in a given task, and how do they

update these efficacy estimates?

Here, we use computational modeling and single-trial EEG analyses to uncover the

mechanisms through which efficacy estimates are learned, updated, and used to guide control

allocation. We show that efficacy estimates are learned from feedback through a mechanism

which parallels reward learning at both the computational (Sutton & Barto, 2018) and the neural

level (Fischer & Ullsperger, 2013). Using model-based analyses of a neural marker of learning

(P3b), we show that prediction errors and corresponding learning rates drive the updating of

efficacy estimates following relevant changes in the environment. We further show that efficacy

estimates learned in this way are used to guide the optimal allocation of control. Extending our

previous findings (Frömer et al., 2020a), we show that a marker of control allocation (the CNV)

increases with increasing model-based efficacy estimates. Taken together, these results show that

efficacy estimates are learned and updated in order to guide optimal allocation of cognitive

control. This mechanism is strikingly similar to the way in which other components of

motivation, including reward value and task difficulty, are learned and used in decision-making

about investment of cognitive resources. Thus, this study contributes to the emerging

understanding of cognitive control as a fundamentally motivated process which relies on learning

of components of motivation to guide the optimal allocation of cognitive resources.

Results

To investigate how efficacy estimates are learned, and how they affect control allocation,

we recorded EEG while 40 participants performed a modified version of the Stroop task (Figure

1A). Similar to our previous study (Frömer et al., 2020a), across trials we varied whether reward

outcomes ($0.50 vs. $0.00) were determined by a participant's performance on a given trial

(responding accurately and below a pre-determined response time criterion; *performance-based*

trials) or whether those outcomes were determined independent of performance (based on a

weighted coin flip; *random* trials). Over the course of the session, we gradually varied the

likelihood that a given trial would be performance-based or random such that, at some points in

the experiment, most trials were performance-based (high efficacy level), and at other points

most trials had random outcomes (low efficacy level) (Figure 1B). Importantly, unlike in our

previous study, participants were not told whether a given trial would be performance-based

(maximal efficacy) or random (minimal efficacy), but instead had to estimate their current

efficacy level based on recent trial feedbacks, which indicated both the reward outcome ($0.50

vs. $0.00) and how that outcome was determined (performance-based [button symbol] vs.

random [dice symbol]). We held reward rate constant across both feedback types by yoking

reward rate on random trials to the reward rate on performance-based trials, and counter-

balanced the time-course of the gradual change in efficacy (see Methods for details). To capture

changes in efficacy expectations over the course of the session, we probed participants every 4-8

trials (averaging 44.3 probes per participant) to report their current estimates of efficacy. These

efficacy probes were interleaved with probes asking participants to estimate the current reward

rate, serving as foils and used for control analyses.

Figure 1. Manipulating expected efficacy and assessing learning. A. Trial Schematic. On each trial participants saw a cue (gray circle), predicting the onset of a Stroop stimulus (target), and were then sequentially presented with reward and efficacy feedback. On half of the trials, efficacy feedback was presented first, and on the other half reward feedback was presented first. Every 2-4 trials participants were subsequently asked to estimate their current efficacy level or reward rate. **B. Efficacy manipulation.** We let the probability of performance-based vs. random feedback continuously drift over the course of the experiment (inversed for one half of the sample). Arrows mark time points with low and high efficacy, respectively. When efficacy was low, rewards were more likely to be random, whereas when efficacy was high, rewards were more likely to be performance-based.

Participants dynamically update efficacy expectations based on feedback

To determine whether and how participants learned their current efficacy levels, we first

analyzed the influence of previous efficacy feedback (whether outcomes on a given previous trial

had been performance-based or random) on one's current subjective estimates of efficacy. We

found very strong evidence that participants adjusted their ratings of expected efficacy upward or

downward depending on whether the most recent trial was Performance-Based or Random ($b =$

0.14; 95% CrI [0.13, 0.16]; $p_{b < 0} = 0$; $BF_{10} > 100$), and that this remained true (but to diminishing

degrees) up to five trials back (all $p_{b < 0} < 0.01$; Figure 2A; Table S1). These results suggest that

participants were dynamically updating their efficacy estimates based on efficacy feedback.

Figure 2. Efficacy learning is captured by a reinforcement learning model separate learning rates for performance-based and random feedback. A. Efficacy estimates track recent efficacy feedback. Regression weights for the influence of the current and previous contingent vs. random feedback on the subjective efficacy estimate. Error bars represent 50% and 95% highest density intervals. **B. A separate learning rate model captures efficacy learning best.** BICs of fitted intercept only and one learning rate model relative to two learning rate models are plotted for each participant and favor the two learning rate model. **C. Learning rate biases vary between participants.** Positive and negative learning rate estimates are plotted for all participants. Points below the diagonal indicate higher learning rates for performance-based compared to random feedback, and points above the opposite. **D. Example learning trajectories.** Subjective and model-based efficacy estimates, and a running average of the previous 5 efficacy feedbacks, for a participant with a higher learning rate for the contingent compared to random feedback (upper) and a participant with the reverse bias (lower).

This pattern of learning was accounted for by a standard reinforcement learning (RL)

algorithm, the delta-rule, according to which efficacy estimates are updated in proportion to the

prediction error between the expected efficacy and the efficacy feedback on a given trial (i.e.,

whether a given outcome was determined by performance or not) (Sutton & Barto, 2018).

Interestingly, consistent with recent studies of reward learning (Garrett & Daw, 2020; Lefebvre

et al., 2017; Niv et al., 2012), we found that the RL model that best accounted for our data

allowed efficacy estimates to be updated differently from trials that were more efficacious than

expected (Performance-Based feedback) than from trials that were less efficacious than expected

(Random feedback), by having separate *learning rates* scaling prediction errors in the two cases.

Even when accounting for model complexity, this Two Learning Rate model outperformed a

One Learning Rate model as well as a baseline model that fits a single constant efficacy estimate

and no learning (Intercept Model) (Figure 2B). We found that the two learning rates for this best-

fit model varied across the group (Figure 2E), but did not find that one was consistently higher

than the other ($b = 0.02$; 95% CrI [-0.02, 0.05]; $p_{b<0} = 0.203$; $BF_{01}=16.41$), despite most

participants (80%) tending to learn more from Performance-Based than Random trials. Taken

together, these results show that participants dynamically updated their expected efficacy based

on trial-by-trial feedback, and that they did so differently based on whether the trial was more

or less efficacious than expected. The fitted models further enable us to generate trial-by-trial

estimates of expected efficacy and efficacy prediction errors, which we use in model-based

analyses of behavior and neural activity below. Note that in all the following analyses we control

for reward rate estimates obtained from models fit to reward feedback (for details see below).

The feedback-related P3b indexes updating of efficacy expectations

To investigate the neural mechanism underlying feedback-based learning of efficacy, we

probed the centroparietal P3b ERP component (Figure 3A), an established neural correlate of

prediction-error based learning (Fischer & Ullsperger, 2013; Nassar et al., 2019). If the P3b

indexes feedback-based updating of *efficacy* predictions, as it does for reward predictions, we

would expect this ERP to demonstrate several key markers of such a learning signal. First, we

would expect the P3b to reflect the extent to which efficacy feedback (Performance-Based vs.

Random) deviates from the current level of expected efficacy. In other words, the P3b should

track the magnitude of the *unsigned efficacy prediction error (PE)* on a given trial. We tested

this by examining how the amplitude of the P3b to a given type of efficacy feedback varied with

model-based estimates of the participant's efficacy expectation on that trial. If the P3b signaled

efficacy PE then its amplitude should scale inversely with the expected probability of a given

type of feedback (i.e., how *unexpected* that feedback is), thus correlating *negatively* with

expected efficacy on trials providing Performance-Based feedback and correlating *positively*

with expected efficacy on trials providing Random feedback. In addition to overall higher P3b to

performance-based compared to random feedback ($b = 0.90$; 95% CrI [0.55, 1.24]; $p_{b<0} = 0$; BF_{10}

> 100; Figure 2B), we found exactly this predicted interaction between feedback type and

expected efficacy ($b = -2.77$; 95% CrI [-4.08, -1.48]; $p_{b>0} = 0$; $BF_{10} > 100$; Figure 2B; Table

S2), with the P3b amplitude decreasing with model-based estimates of expected efficacy on

Performance-Based trials ($b = -1.93$; 95% CrI [-3.07, -0.75]; $p_{b>0} = 0$; $BF_{10} = 6.04$) and

increasing numerically, but not robustly with expected efficacy on Random trials ($b = 0.84$; 95%

CrI [-0.28, 2.00]; $p_{b<0} = 0.11$; $BF_{10} = 0.43$). Accordingly, when we regressed P3b amplitude on

our model-based estimates of trial-to-trial efficacy PE, we found a positive relationship ($b =$

1.23; 95% CrI [0.40, 2.05]; $p_{b<0} = 0.01$; $BF_{10} = 3.10$; Figure 3C-left; Table S2).

Figure 3. P3b reflects dynamically changing efficacy estimates during processing of efficacy feedback. A. ERP average for the P3b locked to the onset of the efficacy feedback separately for performance-based and random feedback. The grey area shows the time window used for quantifying the P3b. **B.** LMM predicted P3b amplitudes are plotted for performance-based and random feedback as a function of efficacy estimates. The topography shows the interaction of efficacy estimate with efficacy feedback in the P3b time window. **C.** Predicted effects of prediction errors (left) and model-based learning rates (right) on the P3b. Shaded error bars represent 95% confidence intervals. Topographies display fixed-effects estimates.

In addition to tracking efficacy PEs, the second key prediction for the P3b if it indexes

efficacy learning, is that it should track the extent to which PEs are used to update estimates of

efficacy (i.e., the learning rate). In the current study, we found that participants differed in their

learning rates for the two forms of efficacy feedback (Performance-Based vs. Random),

providing us with a unique opportunity to perform a within-subject test of whether P3b tracked

differences in learning rate across these two conditions. Specifically, we could test whether a

given subject's P3b was greater in the feedback condition for which they demonstrated a higher

learning rate. We found some evidence for this, with the P3b demonstrating an increase for the

within-subject feedback condition with the higher learning rate, but this result was not conclusive

($b = 3.00$; 95% CrI [-0.86, 6.82]; $p_{b<0} = 0.10$; BF = 1.73; Figure 3C-right). While, theoretically,

prediction error and learning rate would interact in predicting the P3b amplitude, we did not

observe such an interaction here. The fact that we only found additive effects of prediction errors

and learning rate on P3b is, however, in line with previous findings on reward processing

(Fischer & Ullsperger, 2013).

The CNV indexes control allocation based on updated expectations of efficacy

Thus far, our findings suggest that participants dynamically updated expectations of their

performance efficacy based on feedback, and that the P3b played a role in prediction error-based

updating of these expectations. Next, we tested the prediction that learned efficacy estimates

determine the expected benefits of control, and thus influence how much control is allocated

(Shenhav et al., 2013). We have previously shown that people exert more control when

expecting their performance to be more rather than less efficacious on the upcoming trial

(Frömer et al., 2020a). This was reflected in better behavioral performance and higher

amplitudes of the contingent negative variation (CNV; Figure 3B) - a slow fronto-central

waveform preceding target onset, which scales with the amount of control allocated in

preparation for the task (Grent-'t-Jong & Woldorff, 2007; Schevernels et al., 2014). Here we

used the same marker of control, but, unlike in our previous study, efficacy expectations were (a)

learned rather than explicitly instructed; (b) varied over time rather than having a fixed

association with a cue; and (c) varied continuously across the range of efficacy probabilities

rather than alternating between maximal and minimal efficacy. We were therefore interested in

testing whether these dynamically varying expectations of efficacy, as estimated by our model,

would still exert the same influence on behavior and neural activity.

Consistent with our predictions and our previous findings, participants tended to perform

better when they expected performance to be more efficacious, responding faster on correct trials

with increasing model-based estimates of efficacy (Figure 3A; Table S4). This finding replicates

the performance effects we observed using instructed cues, albeit with only moderate evidence (b

= -11.00; 95% CrI [-23.99, 1.63]; $p_{b > 0} = 0.08$; $BF_{01} = 2.55$). Like in our previous studies, faster

responding was not explained by a change in speed-accuracy trade-off, as accuracy did not

decrease with increasing expected efficacy, but if anything, slightly increased ($b = 0.13$; 95% CrI

[-0.11, 0.38]; $p_{b > 0} = 0.19$; $BF_{01} = 2.13$). These analyses controlled for the standard behavioral

effects related to Stroop congruency (i.e., slower and less accurate responding for incongruent

relative to congruent trials; Figure S4).

If the CNV provides an index of control allocation based on current incentive

expectations, it should both reflect one's latest estimate of expected efficacy and should predict

performance on the upcoming trial. Our findings support both predictions. Regressing single-trial

CNV amplitude onto model-based estimates of expected efficacy, and controlling for

expectations of reward (discussed below), we found that these CNV amplitudes tracked current

efficacy expectations (b = -0.38; 95% CrI [-0.77, -0.01]; $p_{b>0}$ = 0.05; BF_{01} = 2.93; Figure 3C;

Table S5). As with the behavioral finding above, this result provides moderate evidence

consistent with our previous CNV finding using instructed cues. Larger CNV amplitude in turn

predicted better performance on the upcoming trial, with participants responding faster (b =

11.39; 95% CrI [8.64, 14.21]; $p_{b<0}$ = 0; BF_{10} > 100; Figure 3D; Table S6) and more accurately

(b = -0.07; 95% CrI [-0.11, -0.02]; $p_{b>0}$ = 0.01; BF_{10} = 3.02; Figure 3E; Table S6) as CNV

increased.

Figure 4. Efficacy estimates influence allocation of control and behavior. A. Higher efficacy predicts faster accurate responses. **B. CNV increases with higher efficacy.** Left: Grant-average ERP for high and low efficacy estimates. The shaded area marks the time window used for CNV quantification. Predicted CNV amplitudes as a function of efficacy estimates. The topography shows the fixed effect of the efficacy estimate from the fitted linear model. **C. -D. Larger CNV amplitude predicts better performance.** Predicted accurate RT (**C.**) and accuracy (**D.**) as a function of efficacy estimates. Shaded error bars indicate 95% confidence intervals.

Parallel learning of efficacy and reward rate

We held the amount of reward at stake constant over the course of the experiment, but the

frequency of reward (reward rate) varied over the course of the session based on a participant's

recent performance, and the participant was sporadically probed for estimates of this reward rate

(interleaved with trials that were followed by efficacy probes). Our efficacy manipulation

explicitly controlled for this variability by yoking Random-Outcome feedback to a participant's

recent reward rate (see methods for details). However, this additional source of variability also

provided an opportunity to examine additional mechanisms of learning and adaptation on our

task. As in the case of efficacy estimates, reward rate estimates were robustly predicted by

reward feedback on previous trials (Table S1; Figure S1A), and this reward rate learning process

was well captured by a two learning rate model (cf. Garrett & Daw, 2020; Figure S1B-D).

Updates to these reward rate estimates were reflected in P3b modulations of (unsigned) reward

prediction errors and associated learning rates (Fischer & Ullsperger, 2013; see Figure S3, Table

S3). This pattern of results provides additional evidence that efficacy learning involves similar

neural and computational mechanisms as reward-based learning.

262   **Discussion**

To evaluate the expected benefits of investing cognitive control into a task, people need

to consider whether this control is necessary and/or sufficient for achieving their desired

outcome (i.e., whether these efforts are worthwhile). A critical determinant of the

worthwhileness of control is performance efficacy, the extent to which performance on a task

matters for outcome attainment versus those outcomes being determined by performance-

unrelated factors. However, the mechanisms through which people estimate the efficacy of their

performance based on previous experience are largely unknown. Here, we identified the

computational and neural mechanism through which efficacy is learned and used to guide the

allocation of cognitive control. We showed that participants dynamically updated expectations of

the efficacy of their task performance (i.e., the likelihood that this performance will determine

reward attainment), and used those expectations to adjust how much control they allocated.

Replicating and extending our previous findings (Frömer et al., 2020a), higher efficacy was

associated with larger CNV amplitudes, a putative index of effort allocation, as well as faster

accurate responses, and trial-by-trial variability in CNV amplitude predicted performance.

Importantly, unlike in that study, here efficacy was learned rather than being explicitly cued. The

dynamic feedback-based updating of efficacy was well-captured by a prediction error based

learning algorithm commonly used to model the process of feedback-based reward learning.

Model-based estimates of expected efficacy and efficacy prediction errors were encoded by

canonical neural signatures of effort allocation and belief updating, respectively. Taken together,

our results uncover the mechanism through which efficacy estimates are learned and used to

inform mental effort investment within a given task environment.

Previous research has characterized the reinforcement learning algorithms people use to learn

the reward value of different states and actions in their environment (Gläscher et al., 2010;

Sutton & Barto, 2018). Recent theoretical (Jiang et al., 2014; Lieder et al., 2018; Verbeke &

Verguts, 2019) and empirical (Bejjani et al., 2018; Bustamante et al., in press; Jiang et al., 2020;

Otto & Daw, 2019) work has extended this research to show how similar algorithms guide

learning and adaptation of control under varying rewards and task demands within a given task

environment. Our findings extend this work further in several ways. First, we show that people

rely on reinforcement learning algorithms similar to these when learning about the efficacy of

task performance, independent of potential reward and task difficulty. Second, we show that they

update their efficacy expectations differently depending on whether performance was more

efficacious or less efficacious than they expected, demonstrating a striking parallel with dual

learning rate models that have been found to prevail in research on reward learning (Collins &

Frank, 2013; Garrett & Daw, 2020; Lefebvre et al., 2017), including in our own reward rate data.

Third, we show that participants dynamically adjust their control allocation based on learned

efficacy, just as they do for learned rewards and task demands (Bugg et al., 2011; Jiang et al.,

2014; Lieder et al., 2018).

Our neural findings build further on past research on learning and control adaptation. The

P3b component has been shown to track prediction-error based learning from action-relevant

outcome values (Fischer & Ullsperger, 2013; Lohse et al., 2020; Nassar et al., 2019). This work

has shown that the P3b amplitude is larger in situations in which people are updating their

expectations more based on the feedback they received. Here we show that this neural marker

tracks learning about efficacy in the same way as it tracks learning about rewards. We found

increased P3b amplitudes when people experienced outcome contingencies that were less

expected given their current estimate of efficacy (e.g., expecting low efficacy, but getting

performance-contingent feedback), relative to when these contingencies were more expected

(e.g., expecting low efficacy and getting random feedback). Extending previous findings cueing

reward and/or efficacy (Frömer et al., 2020a; Schevernels et al., 2014), our CNV and behavioral

results further show that participants used these learned efficacy estimates to calibrate their effort

and their performance. To the best of our knowledge, our study is the first one to show effort-

related changes in CNV amplitude entirely divorced from perceptual cues, providing compelling

evidence that this activity truly reflects adjustments in control, rather than reactive processing of

features associated with the cue. Taken together, our findings suggest that similar neural

mechanisms underly learning and control adaptation in response to variability in one's efficacy

in a task environment, as they do in response to variability in expected rewards.

Our efficacy-updating results are a reminder that many aspects of feedback are reflected in

prediction error signals (Frömer et al., 2020b; Langdon et al., 2018). These aspects are rarely as

clearly separated into different stimuli as they are in the present study. Instead, feedback is

multiplexed, containing information relevant to multiple targets of evaluation, internal, such as

the selected response or its predicted outcome, or external, such as the source of

feedback/environment. If these reinforcement learning mechanisms support selection of not only

individual actions but also extended sequences of actions and control signals (Holroyd & Yeung,

2012; Lieder et al., 2018), their neural signatures could further reflect evaluations of control

signal quality or intensity, adding to the difficulty of interpreting neural correlates of feedback.

Multi-level prediction error signals might for instance explain, why despite close links between

P3b and behavioral adaptation (Chase et al., 2011; Fischer & Ullsperger, 2013; Yeung & Sanfey,

2004), attempts to link the P3b to belief updating have failed (Nassar et al., 2019).

Understanding how efficacy estimates develop based on previous experiences is crucial for

understanding why people differ in how much cognitive control they allocate in different

situations (Shenhav et al., submitted). People differ in their beliefs about how much control they

have over their environments (Leotti et al., 2010; Moscarello & Hartley, 2017), and in their

ability to estimate their efficacy (Cohen et al., 2020). Further, many mental disorders, including

depression and schizophrenia, have been linked with one's estimates of their ability to control

potential outcomes in their environment, including through allocation of control (Huys & Dayan,

2009; Maier & Seligman, 2016), and we have recently proposed that such changes can drive

impairments of motivation and control in those populations (Grahek et al., 2019). As we show in

this study, if efficacy is expected to be low, people will allocate less cognitive control, which can

manifest as apparent control deficits. The experimental and modeling approach taken in our

study helps uncover a more fine-grained view of how components of motivation are learned and

used to support the allocation of cognitive resources. In this way, our study takes a first step

toward a better computational and neural account of efficacy learning, which can aid the

understanding of individual differences in the willingness to exert mental effort, as well as the

development of interventions aimed at teaching individuals when these efforts truly matter.

Methods

Participants

We recruited forty-one participants with normal or corrected-to-normal vision from the

Brown University subject pool. One participant was excluded due to technical issues. The final

data set included 40 participants (24 females, 16 males; median age = 19). Participants gave

informed consent and were compensated with course credits or a fixed payoff of $20. In

addition, they received up to $5 as a performance-based bonus ($3.25 on average). The research

protocol was approved by Brown University's Institutional Review Board.

Stimuli and task

In the main task, taking approximately 45 minutes, participants performed 288 Stroop

trials (Figure 1A). Each trial started with the presentation of a cue (grey circle) that remained on

the screen throughout the trial. After a period of 1500 ms, a Stroop stimulus was superimposed

until a response was made or 1000 ms elapsed, at which time it was sequentially replaced with

two types of feedback presented for 1000 ms each. Each trial onset was preceded by a fixation

cross (randomly jittered between 1000 and 1500ms). Participants responded to the ink color of

the Stroop stimulus (red, green, blue, or yellow) by pressing one of four keyboard keys (D, F, J,

and K). Stimuli were either color words same as the ink color (congruent, n = 108) or different

(incongruent, n = 108), or a string of letters "XXXXX" (neutral, n= 72). Feedback informed

them whether they obtained a reward (reward, "$50c" or no reward, "$0c") and whether the

reward they received depended on their performance (performance-based feedback, a button

graphic), or not (random feedback, a dice graphic). The order of the two types of feedback was

pseudo-randomized with half of the trials showing reward feedback first and the other half

efficacy feedback. Every 2-4 trials the feedback was followed by a probe of efficacy ("How

much do you think your rewards currently depend on your performance?") or reward rate ("How

often do you think you are currently being rewarded?") to which participants responded on a

visual analog scale ranging from 0 to 100. The number and timing of the probes was randomized

per subject resulting in a median of 45 efficacy probes (SD=3.38) and 47 reward probes

(SD=2.81).

Efficacy (performance-based or random rewards) on each trial was sampled from a

binomial distribution with probabilities ranging between 0.1 and 0.9 that drifted over the course

of the experiment and were predetermined (Figure 1B). In order to ensure that the performance-

based and random trials did not differ in reward rate, reward feedback for the random trials was

sampled from the moving window of the reward feedback of the previous 10 performance-based

trials. At the beginning of the experiment a window with 8 rewards and 2 no rewards were used

to reflect the calibrated accuracy, and was then updated after every trial to reflect any changes in

performance accuracy. Thus, reward rate was not experimentally manipulated in the experiment

and remained constant.

Prior to the main task, participants performed several practice phases of the Stroop task

(approximately 15 minutes). First, they practiced the mappings between colors and keyboard

keys (80 trials). Then they completed a short practice of the Stroop task with written feedback

("correct" or "incorrect") on each trial (30 trials). Participants then completed 100 more of such

trials during which we individually calibrated the reaction time deadline such that participants

yielded approximately 80% accuracy. In the final practice phase participants were introduced to

the two types of feedback which they would see in the main experiment (30 trials).

The experimental task was implemented in Psychophysics Toolbox (Brainard, 1997;

Kleiner et al., 2007; Pelli, 1997) for Matlab (MathWorks Inc.) and presented on a 23 inch screen

with a 1920 x 1080 resolution. All of the stimuli were presented centrally while the participants

were seated 80 cm away from the screen.

Psychophysiological recording and preprocessing

EEG data were recorded at a sampling rate of 500 Hz from 64 Ag/AgCl electrodes

mounted in an electrode cap (ECI Inc.), referenced against Cz, using Brain Vision Recorder

(Brain Products, München, Germany). Vertical and horizontal ocular activity was recorded from

below both eyes (IO1, IO2) and the outer canthi (LO1, LO2), respectively. Impedances were

kept below 10 kΩ. Offline, data were processed using custom made Matlab scripts (Frömer et al.,

2018) employing EEGlab functions (Delorme & Makeig, 2004). Data were re-referenced to

average reference, ocular artifacts were corrected using brain electric source analyses (BESA;

Ille, Berg, & Scherg, 2002) based on separately recorded prototypical eye movements. The

cleaned continuous EEG was then high pass filtered at 40 Hz and segmented into epochs around

cue onset (-200 to 1500 ms), stimulus onset, and efficacy feedback (-200 to 800 ms). Baselines

were corrected to the average of each 200 ms pre-stimulus interval. Segments containing

artifacts, values exceeding ± 150 µV or gradients larger than 50 µV, were excluded from further

analyses.

Single trial ERPs were then exported for further analyses in R (R Core Team, 2017). The

late CNV was quantified between 1000 and 1500 ms (cf. Frömer et al., 2016; Frömer et al.,

2020a; Schevernels et al., 2014) as the average activity over 9 fronto-central electrodes (Fz, F1,

F2, FCz, FC1, FC2, Cz, C1, and C2). The P3b was quantified between 350 and 500 ms (cf.

Fischer & Ullsperger, 2013) for both reward and efficacy feedback and calculated as the

averaged activity over 9 centro-parietal electrodes (Pz, P1, P2, POz, PO1, PO2, CPz, CP1, and

CP2).

Learning model and statistical analyses

Learning models. In order to obtain trial-by-trial estimates of efficacy and reward rate, we fitted

two temporal difference learning models (Gläscher et al., 2010; Sutton & Barto, 2018) to the

continuous (range: 0-1) subjective estimates of efficacy and reward rate (cf. Eldar, Hauser,

Dayan, & Dolan, 2016; Nagase et al., 2018; Rutledge et al., 2014). The first model ("1 learning

rate model") assumed that the estimate of efficacy for the next trial (Et+1) depended on the

current efficacy feedback (Et) and the prediction error (δt) weighted by a constant learning rate

(α):

424
$$E_{t+1} = E_t + \alpha * \delta_t$$

Where $0 \leq \alpha \leq 1$, and the prediction error is calculated as the difference between the contingency

feedback on the current trial and the efficacy estimate on that trial: $\delta t = et - Et$. The model

started from an initial value (free parameter) and updated the subjective estimate based on the

binary feedback on each trial. For example, assuming a learning rate of 0.5 and the initial value

of 0.5, the model would update the initial estimate following efficacy feedback signaling

"performance-based" ($e_t = 1$) to 0.75. If on the next trial contingency feedback was "random" (e_t

$= 0$) the subjective estimate would drop to 0.6. Exactly the same model was used to calculate

estimates of reward on each trial. The model was fitted separately to the subjective estimates of

efficacy and reward rate with only the learning rate as a free parameter. The second model ("2

learning rates model") was the same as the first model, but it included two learning rates: one

learning rate for learning from the efficacy feedback, and another for learning from the no

efficacy feedback. Finally, as a baseline, we also included the "intercept model" which did not

update the efficacy estimate throughout the experiment, but just assumed that the estimate took

one constant value. The models were fit hierarchically to the data using maximum likelihood (for

details see the MATLAB scripts available on GitHub).

Statistical analyses. The efficacy and reward rate estimates obtained through fitting the learning

model were then used to analyze the behavioral and EEG data. To this end, we fitted Bayesian

multilevel regressions to predict subjective estimates of efficacy and reward rates, reaction times,

accuracy, as well as the CNV and P3b amplitudes. Subjective estimates of efficacy and reward

rate were regressed onto efficacy or reward feedback. Reaction times and accuracies were

regressed onto trial-by-trial model-based estimates of efficacy and reward rate, as well as trial-

by-trial CNV amplitude, while controlling for congruency. The P3b amplitudes were analyzed in

two ways: with trial-by-trial model-based estimates of efficacy and reward rate and current

feedback as predictors, and with model-based prediction errors and learning rates for each

feedback type. CNV, amplitudes were regressed onto trial-by-trial model-based estimates of

efficacy. All of the fitted models controlled for the influence of the reward rate estimates.

Parallel analyses were done to predict the P3b in response to reward feedback, while controlling

for the efficacy estimates.

The regression models were fitted in R with the *brms* package (Bürkner, 2016) which

relies on the probabilistic programming language *Stan* (Carpenter et al., 2016) to implement

Markov Chain Monte Carlo (MCMC) algorithms and estimate posterior distributions of model

parameters. The analyses were done based on the recommendations for Bayesian multilevel

modeling using *brms* (Bürkner, 2016, 2017; Nalborczyk & Bürkner, 2019). The fitted models

included constant and varying effects (also known as fixed and random) with weakly informative

priors (except for the behavioral and CNV analyses, see below for details) for the intercept and

the slopes of fixed effects and the likelihoods appropriate for the modeled data (Ex-Gaussian for

reaction times, Bernoulli for accuracy, and Gaussian for the subjective estimates and the EEG

amplitudes). The fitted models included all of the fixed effects as varying effects. All of the

continuous predictors in the model were centered and the categorical predictors were contrast

coded. Four MCMC simulations ("chains"; 20,000 iterations; 19,000 warmup) were run to

estimate the parameters of each of the fitted models. The convergence of the models was

confirmed by examining trace plots, autocorrelation, and variance between chains (Gelman-

Rubin statistic; Gelman & Rubin, 1992). After convergence was confirmed, we analyzed the

posterior distributions of the parameters of interest and summarized them by reporting the means

of the distribution for the given parameter (*b*) and the 95% credible intervals (95% CrI) of the

posterior distributions of that model. We report the proportion of the posterior samples on the

relevant side of 0 (e.g., $p_{b < 0} = 0.9$), which represents the probability that the estimate is below or

above 0. We also report Bayes factors (BF) calculated using the Savage-Dickey method

(Wagenmakers et al., 2010). We report the BFs in support of the alternative hypothesis against

the null (BF_{10}), except for the analyses of accurate RT and CNV amplitude in which we have

informative priors based on our previous study (Frömer, et al., 2020a), and in which case we

support the evidence in favor of the null (BF_{10}).

To compare the positive and negative learning rates we fitted a model in which the

learning rates were predicted by the learning rate type (Kruschke, 2013). In this model we used

Gaussian distributions as priors (intercept: (0.5,0.5); slopes: (0,0.5)).

We fitted two separate models to predict the subjective estimates of efficacy and reward

rate based on previous feedbacks. At each timepoint the estimates were predicted by the current,

and previous 4 feedbacks. The feedback on each of the trials (performance-based vs. random or

reward vs. no reward) was entered as a constant effect and the models also included the intercept

as a varying effect. As the subjective estimates could vary between 0 and 1, we used Gaussian

distributions (intercept: (0.5,0.2); slopes: (0,0.2)) as priors.

For predicting both the P3b amplitude in response to the onset of the efficacy feedback,

we fitted two models. First, we fitted a model which included the model-based estimate of

efficacy, the actual feedback, and the interaction between the expected efficacy and the observed

efficacy feedback. Additionally, we controlled for the reward rate estimate. Second, we fitted

separate models which included the model-based prediction errors, the influence of the between-

subject learning rates, and their interaction with the prediction errors, while controlling for the

estimate of the reward rate. In this analysis the learning rates (one for feedback type for each

subject) were mean-centered within subjects and thus any effect of the learning rates is driven by

the difference between the random and the performance-based learning rate. For these models we

selected wide Gaussian priors (intercept: (5,3); slopes (0,3)). The same logic in building models

was applied for the analyses of the reward feedback. In these analyses we focused on the reward

feedback processing and how it interacted with the model-based estimates of reward rates, while

controlling for the model-based estimates of efficacy. We analyzed only the trials with correct

responses for both the efficacy and the reward feedback analyses.

To test the influence of efficacy on the late CNV, we fitted a model which predicted the

CNV based on the model-based efficacy estimates, while controlling for the effect of the reward

rate estimates. Drawing on the results of our previous study (Frömer, et al., 2020a), this model

included Gaussian priors for the intercept (-0.16, 1.59) and the efficacy (-0.30, 0.73) and reward

(0,0.73) slopes.

For predicting reaction times and accuracy we fitted models which included congruency

(Facilitation: difference between neutral and congruent trials; Interference: difference between

incongruent and neutral trials) and the model-based efficacy estimates, while controlling for the

reward rate estimates. We used Gaussian distributions as informative priors based on our

previous study (Frömer, et al., 2020a), for both the reaction times (intercept: (624, 58.69);

facilitation (15.54, 21.93), interference (61.12, 37.49); efficacy (-10.26, 19.51); reward (0,

19.51)) and accuracy[1] analyses (intercept: (2.11, 0.81); facilitation (-0.45, 0.64), interference (-

0.53, 0.81); efficacy (0.09, 0.32); reward (0, 0.32)).

To investigate how the late CNV influences the behavior, we fitted two models in which

we predicted the reaction times and accuracy based on the CNV amplitude. The prior

---

[1] Note that the prior distributions are set in log-odds.

distributions for these models were weakly informative Gaussian distributions for predicting

both the reaction times (intercept: (650, 200); slope: (0, 50)) and accuracy (intercept: (0.7, 0.2);

slope: (0, 0.2)).

To visualize the topographies of the relevant ERP effects, we fitted the relevant models to

all 64 channels and then plotted the posterior estimates of the effects of interest at each electrode.

Supplemental Materials

Figure S1. Reward rate learning – model-based analyses. A. Regression weights for the influence of the current and previous feedbacks on the subjective estimates of reward rate (left) and efficacy (right). Reward rate estimates are strongly predicted by the previous reward feedbacks, but not efficacy feedbacks. The reverse is true for the subjective efficacy estimates. Error bars represent 50% and 95% highest density intervals. **B.** Model comparison between the fitted learning models for the reward rate learning model. **C.** Positive and negative learning rate estimates for all subjects for the reward rate learning model. Negative learning rates were numerically, but not statistically larger than positive ones ($b = 0.02$; 95% CrI [-0.04, 0.08]; $p_{b<0} = 0.249$; $BF_{10} = 0.07$) **D.** Subjective and model-based reward rate estimates, and a running average of the previous 5 reward feedbacks, for a subject with a higher learning rate for the no reward feedback (upper) and a subject with the opposite bias (lower).

522 Table S1

Regression weights for the models predicting subjective efficacy and reward rate estimates based on efficacy and reward feedback 5 trials back.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)	BF$_{10}$
Subjective efficacy				
Intercept	0.52	0.50, 0.55	0.00	>100
T1 Performance-Based - Random	0.14	0.13, 0.16	0.00	>100
T2 Performance-Based - Random	0.12	0.11, 0.14	0.00	>100
T3 Performance-Based - Random	0.08	0.06, 0.09	0.00	>100
T4 Performance-Based - Random	0.03	0.01, 0.05	0.00	2.18
T5 Performance-Based - Random	0.03	0.01, 0.04	0.00	1.21
T1 Reward - No Reward	0.02	0.00, 0.04	0.03	0.32
T2 Reward - No Reward	0.03	0.01, 0.05	0.00	2.70
T3 Reward - No Reward	0.03	0.01, 0.04	0.01	0.81
T4 Reward - No Reward	0.02	0.00, 0.04	0.02	0.51
T5 Reward - No Reward	0.00	-0.02, 0.02	0.47	0.06
Subjective reward rate				
Intercept	0.50	0.46, 0.53	0.00	>100
T1 Reward - No Reward	0.19	0.18, 0.21	0.00	>100
T2 Reward - No Reward	0.16	0.14, 0.17	0.00	>100
T3 Reward - No Reward	0.09	0.07, 0.10	0.00	>100
T4 Reward - No Reward	0.05	0.03, 0.06	0.00	>100
T5 Reward - No Reward	0.05	0.03, 0.06	0.00	>100
T1 Performance-Based - Random	-0.01	-0.02, 0.01	0.74	0.06
T2 Performance-Based - Random	0.0	-0.01, 0.02	0.32	0.05
T3 Performance-Based - Random	0.01	-0.01, 0.02	0.00	0.05
T4 Performance-Based - Random	0.01	-0.01, 0.02	0.21	0.08
T5 Performance-Based - Random	0.02	0.00, 0.03	0.04	0.22

Figure S2. Efficacy learning – the effect of different feedback probability drifts on learning. A. The probability drift of contingent feedback presented to the first half of subjects. **B.** Learning rates for contingent and random feedback for the first half of subjects. Learning rates for the contingent and random feedback did not differ ($b = 0.01$; 95% CrI [-0.04, 0.08]; $p_{b<0} = 0.330$; $BF_{10} = 0.07$). **C.** The inversed probability drift of contingent feedback presented to the second half of participants. **D.** Learning rates for contingent and random feedback did not differ ($b = 0.02$; 95% CrI [-0.04, 0.08]; $p_{b<0} = 0.248$; $BF_{10} = 0.07$).

Table S2

Regression weights for the models predicting the P3b to efficacy feedback.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)	BF$_{10}$
Efficacy feedback				
Intercept	1.40	0.77, 2.05	0.00	>100
Model-based efficacy	-0.54	-1.46, 0.42	0.84	0.30
Efficacy feedback type (PB-R)	0.90	0.55, 1.24	0.00	>100
Model-based reward rate	-0.27	-1.15, 0.63	0.69	0.20
Model-based efficacy ×Efficacy feedback type	-2.77	-4.08, -1.48	1.00	>100
Model-based efficacy slope for Performance-Based feedback	-1.93	-3.07, -0.75	1.00	6.04
Model-based efficacy slope for Random feedback	0.84	-0.28, 2.00	0.11	0.43
Predictions errors and learning rates				
Intercept	1.30	0.69, 1.92	0.00	>100
Unsigned prediction error	1.23	0.40, 2.05	0.01	3.10
Learning rate	3.00	-0.86, 6.82	0.10	1.73
Model-based reward rate	-0.22	-1.16, 0.71	0.66	0.21
Unsigned prediction error ×Learning rate	0.51	-4.37, 5.30	0.43	0.96

ERP analyses of the reward rate feedback

To investigate the effect of the model-based reward rate estimates on the processing of

reward, we performed complementary analyzes on the P3b related to the processing of the

reward feedback (Figure S3A). We expected that a similar learning mechanism should operate

on both reward and efficacy feedback, but that the neural markers of feedback processing should

be sensitive to different model-based estimates. Thus, we expected the reward-locked P3b to be

sensitive to reward-rate estimates, but not to efficacy estimates.

Our results showed larger P3b amplitudes to no reward compared to reward feedback

(note that reward rate was approximately .80 and negative feedback less expected overall), and

the processing of the reward feedback was influenced by reward-rate estimates, although to less

extent than for the P3b (Figure S3B; Table S3). Probing the effects of learning directly, we found

robust effects of the prediction errors and learning rates (Figure S3C). The P3b to reward

feedback was not influenced by the model-based efficacy estimates in either of these analyses.

Figure S3. Learning reward rate estimate from reward feedback. A. ERP average for the P3b locked to the onset of reward feedback. **B.** Interaction between reward feedback and the model-based reward rate estimate. **C.** The effects of prediction errors (left) and earning rates (right) on the P3b.

Table S3

Regression weights for the models predicting the P3b to reward feedback.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)	BF$_{10}$
Efficacy feedback				
Intercept	0.98	0.30, 1.66	0.01	>100
Model-based efficacy	-0.54	-1.33, 0.25	0.87	0.30
Reward feedback type (R-NR)	-0.32	-0.76, 0.12	0.90	0.19
Model-based reward rate	0.05	-0.55, 0.66	0.44	0.12
Model-based reward ×Reward feedback type	-1.14	-2.57, 0.30	0.90	0.65
Model-based efficacy slope for No Reward feedback	0.03	-1.25, 1.28	0.99	1.54
Model-based efficacy slope for Reward feedback	-1.11	-1.91, -0.27	0.48	0.23
Predictions errors and learning rates				
Intercept	0.87	0.22, 1.52	0.02	>100
Unsigned prediction error	1.06	0.47, 1.64	0.00	7.55
Learning rate	3.78	-0.76, 8.12	0.08	2.79
Model-based efficacy	-0.03	-0.61, 0.57	0.54	0.12
Unsigned prediction error ×Learning rate	0.37	-4.44, 5.20	0.45	0.99

Figure S4. Behavioral effects of congruency. A. In comparison to the neutral trials, participants were faster to respond to congruent ($b = 9.93$; 95% CrI [4.26, 15.51]; $p_{b<0} = 0.002$; BF = 499) and slower to respond to incongruent trials ($b = 60.90$; 95% CrI [51.43, 70.49]; $p_{b < 0} = 0$; BF > 4000). **B.** In comparison to the neutral trials, participants were equally likely to be correct when responding to congruent ($b = -0.11$; 95% CrI [-0.25, 0.03]; $p_{b > 0} = 0.096$; BF = 9.42) and less likely to be correct when responding to incongruent trials ($b = -0.48$; 95% CrI [-0.61, -0.34]; $p_{b>0} = 0$; BF > 4000). The regression

Table S4

Regression weights for the models predicting reaction times and accuracy based on efficacy and reward rate estimates.

Parameter	Estimate	95% Credible interval	Posterior probability ($p<0$)	BF_{01}
Reaction times				
Intercept	654.88	638.52, 671.45	0.00	>100
Facilitation	9.95	4.56, 15.42	0.00	1.64
Interference	61.44	51.94, 70.85	0.00	6.54
Model-based efficacy	-11.00	-23.99, 1.63	0.92	2.55
Model-based reward rate	-9.99	-26.81, 9.08	0.83	0.98
Accuracy				
Intercept	1.58	1.39, 1.77	0.00	>100
Facilitation	-0.12	-0.26, 0.03	0.91	0.01
Interference	-0.48	-0.61, -0.35	1.00	8.41
Model-based efficacy	0.13	-0.11, 0.38	0.19	2.13
Model-based reward rate	-0.19	-0.45, 0.07	0.89	0.89

Table S5

Regression weights for the models predicting the CNV based on efficacy and reward rate.

Parameter	Estimate	95% Credible interval	Posterior probability ($p<0$)	BF_{01}
CNV				
Intercept	-0.36	-0.52, -0.19	1.00	>100
Model-based efficacy	-0.38	-0.77, -0.01	0.95	2.93
Model-based reward rate	-0.06	-0.47, 0.35	0.60	2.90

Table S6

Regression weights for the models predicting reaction times and accuracy based on the
CNV.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)	BF$_{10}$
Reaction times				
Intercept	656.03	639.14, 673.19	0.00	>100
CNV amplitude	11.39	8.64, 14.21	0.00	>100
Accuracy				
Intercept	1.33	1.15, 1.50	0.00	>100
CNV amplitude	-0.07	-0.11, -0.02	0.99	3.02

Acknowledgements

This work was supported by the Special Research Fund (BOF) of Ghent University [grant

#01D02415] (I.G.) the Research Foundation Flanders (FWO) travel grant [grant #V432718N]

(I.G.), a Center of Biomedical Research Excellence grant P20GM103645 from the National

Institute of General Medical Sciences (A.S.), and the Alfred P. Sloan Foundation Research

Fellowship in Neuroscience (A.S.). The funding sources were not involved in the study design;

collection, analysis, and interpretation of data; writing of the report; and decision to submit the

article for publication. We would like to thank Natalie Knowles and Hattie Xu for their help with

data collection, Carolyn Dean Wolf for her help with programming the task, and Harrison Ritz for

his help with constructing the efficacy drifts.

Data availability

Raw and pre-processed data, materials, and analysis scripts will be available on Github and OSF

upon publication.

References

Bejjani, C., Zhang, Z., & Egner, T. (2018). Control by association: Transfer of implicitly primed

attentional states across linked stimuli. *Psychonomic Bulletin and Review*, *25*(2), 617–626.

https://doi.org/10.3758/s13423-018-1445-6

Botvinick, M. M., & Cohen, J. D. (2014). The computational and neural basis of cognitive

control: Charted territory and new frontiers. *Cognitive Science*, *38*(6), 1249–1285.

https://doi.org/10.1111/cogs.12126

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

Bugg, J. M., Jacoby, L. L., & Chanani, S. (2011). Why it is too early to lose control in accounts

of item-specific proportion congruency effects. *Journal of Experimental Psychology:*

Human Perception and Performance, *37*(3), 844–859. https://doi.org/10.1037/a0019957

Bürkner, P.-C. (2016). brms: An R package for Bayesian multilevel models using Stan. *Journal*

of Statistical Software, *80*(1), 1–28.

Bürkner, P.-C. (2017). Advanced Bayesian Multilevel Modeling with the R Package brms.

ArXiv:1705.11123.

Bustamante, L., Lieder, F., Shenhav, A., & Cohen, J. D. (n.d.). Learning to Overexert Cognitive

Control in a Stroop Task. *Cognitive, Affective and Behavioral Neuroscience*.

Carpenter, B., Gelman, A., Hoffman, M., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M.

A., Guo, J., Li, P., & Riddell, A. (2016). Stan: A probabilistic programming language.

Journal of Statistical Software, *2*(20), 1–37.

Chambon, V., Théro, H., Vidal, M., Vandendriessche, H., Haggard, P., & Palminteri, S. (2020).

Information about action outcomes differentially affects learning from self-determined

versus imposed choices. *Nature Human Behaviour*. https://doi.org/10.1038/s41562-020-

0919-5

Chase, H. W., Swainson, R., Durham, L., Benham, L., & Cools, R. (2011). Feedback-related

negativity codes prediction error but not behavioral adjustment during probabilistic reversal

learning. *Journal of Cognitive Neuroscience*, *23*(4), 936–946.

https://doi.org/10.1162/jocn.2010.21456

Chiu, Y. C., & Egner, T. (2019). Cortical and subcortical contributions to context-control

learning. *Neuroscience and Biobehavioral Reviews*, *99*(December 2018), 33–41.

https://doi.org/10.1016/j.neubiorev.2019.01.019

Cohen, A., Nussenbaum, K., Dorfman, H. M., Gershman, S. J., & Hartley, C. A. (2020). The

rational use of causal inference to guide reinforcement learning strengthens with age. *Npj*

Science of Learning.

Collins, Anne G E, & Frank, M. J. (2014). Opponent actor learning (OpAL): Modeling

interactive effects of striatal dopamine on reinforcement learning and choice incentive.

Psychological Review, *121*(3), 337–366. https://doi.org/10.1037/a0037015

Collins, Anne Gabrielle Eva, & Frank, M. J. (2013). Cognitive control over learning: creating,

clustering, and generalizing task-set structure. *Psychological Review*, *120*(1), 190–229.

https://doi.org/10.1037/a0030852

Daw, N. D., & Tobler, P. N. (2013). Value Learning through Reinforcement: The Basics of

Dopamine and Reinforcement Learning. In *Neuroeconomics: Decision Making and the*

Brain: Second Edition (pp. 283–298). https://doi.org/10.1016/B978-0-12-416008-8.00015-2

Delorme, A., & Makeig, S. (2004). EEGLAB: an open sorce toolbox for analysis of single-trial

EEG dynamics including independent component anlaysis. *Journal of Neuroscience*

Methods, *134*, 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Dolan, R. J., & Dayan, P. (2013). Goals and habits in the brain. *Neuron*, *80*(2), 312–325.

https://doi.org/10.1016/j.neuron.2013.09.007

Eldar, E., Hauser, T. U., Dayan, P., & Dolan, R. J. (2016). Striatal structure and function predict

individual biases in learning to avoid pain. *Proceedings of the National Academy of*

Sciences of the United States of America, *113*(17), 4812–4817.

https://doi.org/10.1073/pnas.1519829113

Fischer, A. G., & Ullsperger, M. (2013). Real and fictive outcomes are processed differently but

converge on a common adaptive mechanism. *Neuron*, *79*(6), 1243–1255.

https://doi.org/10.1016/j.neuron.2013.07.006

Frömer, R., Lin, H., Dean Wolf, C., Inzlicht, M., & Shenhav, A. (2020a). When effort matters:

Expectations of reward and efficacy guide cognitive control allocation. *BioRxiv*.

https://doi.org/https://doi.org/10.1101/2020.05.14.095935

Frömer, R., Maier, M., & Rahman, R. A. (2018). Group-level EEG-processing pipeline for

flexible single trial-based analyses including linear mixed models. *Frontiers in*

Neuroscience, *12*(48). https://doi.org/10.3389/fnins.2018.00048

Frömer, R., Nassar, M., Stürmer, B., Sommer, W., & Yeung, N. (2020b). I knew that! Response-

based OutcomePredictions and Confidence Regulate Feedback Processing and Learning.

BioRxiv.

Frömer, R., Stürmer, B., & Sommer, W. (2016). (Don't) Mind the effort: Effects of contextual

interference on ERP indicators of motor preparation. *Psychophysiology*, *53*(10), 1577–1586.

https://doi.org/10.1111/psyp.12703

Garrett, N., & Daw, N. D. (2020). Biased belief updating and suboptimal choice in foraging

decisions. *Nature Communications*, *11*(1), 1–12. https://doi.org/10.1038/s41467-020-

16964-5

Gelman, A., & Rubin, D. B. (1992). Inference from Iterative Simulation Using Multiple

Sequences. *Statistical Science*, *7*(4), 457–472. https://doi.org/10.1214/ss/1177011136

Gläscher, J., Daw, N., Dayan, P., & O'Doherty, J. P. (2010). States versus rewards: Dissociable

neural prediction error signals underlying model-based and model-free reinforcement

learning. *Neuron*, *66*(4), 585–595. https://doi.org/10.1016/j.neuron.2010.04.016

Grahek, I., Shenhav, A., Musslick, S., Krebs, R. M., & Koster, E. H. W. (2019). Motivation and

cognitive control in depression. *Neuroscience & Biobehavioral Reviews*, *102*(March), 371–

381. https://doi.org/10.1016/j.neubiorev.2019.04.011

Grent-'t-Jong, T., & Woldorff, M. G. (2007). Timing and sequence of brain activity in top-down

control of visual-spatial attention. *PLoS Biology*, *5*(1), 0114–0126.

https://doi.org/10.1371/journal.pbio.0050012

Holroyd, C. B., & Yeung, N. (2012). Motivation of extended behaviors by anterior cingulate

cortex. *Trends in Cognitive Sciences*, *16*(2), 122–128.

https://doi.org/10.1016/j.tics.2011.12.008

Huys, Q. J. M., & Dayan, P. (2009). A Bayesian formulation of behavioral control. *Cognition*,

113(3), 314–328. https://doi.org/10.1016/j.cognition.2009.01.008

Ille, N., Berg, P., & Scherg, M. (2002). Artifact Correction of the Ongoing EEG Using Spatial

Filters Based on Artifact and Brain Signal Topographies. *Journal of Clinical*

Neurophysiology, *19*(2), 113–124. https://doi.org/10.1097/00004691-200203000-00002

Jiang, J., Bramão, I., Khazenzon, A., Wang, S. F., Johansson, M., & Wagner, A. D. (2020).

Temporal Dynamics of Memory-guided Cognitive Control and Generalization of Control

via Overlapping Associative Memories. *Journal of Neuroscience*, *40*(10), 2343–2356.

https://doi.org/10.1523/JNEUROSCI.1869-19.2020

Jiang, J., Heller, K., & Egner, T. (2014). Bayesian modeling of flexible cognitive control.

Neuroscience and Biobehavioral Reviews, *46*(P1), 30–43.

https://doi.org/10.1016/j.neubiorev.2014.06.001

Kleiner, M., Brainard, D., & Pelli, D. (2007). What's new in Psychtoolbox-3. *Perception 36*

ECVP Abstract Supplement.

Kool, W., & Botvinick, M. (2018). Mental labour. *Nature Human Behaviour*.

https://doi.org/10.1038/s41562-018-0401-9

Krebs, R. M., Boehler, C. N., & Woldorff, M. G. (2010). The influence of reward associations on

conflict processing in the Stroop task. *Cognition*, *117*(3), 341–347.

https://doi.org/10.1016/j.cognition.2010.08.018

Kruschke, J. K. (2013). Bayesian estimation supersedes the T test. *Journal of Experimental*

Psychology: General, *142*(2), 573–588. https://doi.org/10.1037/a0029177

Langdon, A. J., Sharpe, M. J., Schoenbaum, G., & Niv, Y. (2018). Model-based predictions for

dopamine. *Current Opinion in Neurobiology*, *49*, 1–7.

https://doi.org/10.1016/j.conb.2017.10.006

Lefebvre, G., Lebreton, M., Meyniel, F., Bourgeois-Gironde, S., & Palminteri, S. (2017).

Behavioural and neural characterization of optimistic reinforcement learning. *Nature*

Human Behaviour, *1*(4), 1–9. https://doi.org/10.1038/s41562-017-0067

Leotti, L. A., Iyengar, S. S., & Ochsner, K. N. (2010). Born to choose: The origins and value of

the need for control. *Trends in Cognitive Sciences*, *14*(10), 457–463.

https://doi.org/10.1016/j.tics.2010.08.001

Lieder, F., Shenhav, A., Musslick, S., & Griffiths, T. L. (2018). Rational metareasoning and the

plasticity of cognitive control. *Plos Computational Biology*, *14*(4).

https://doi.org/10.13140/RG.2.2.24500.14721

Lohse, K. R., Miller, M. W., Daou, M., Valerius, W., & Jones, M. (2020). Dissociating the

contributions of reward-prediction errors to trial-level adaptation and long-term learning.

Biological Psychology, *149*(January 2019).

https://doi.org/10.1016/j.biopsycho.2019.107775

Maier, S. F., & Seligman, M. E. P. (2016). Learned helplessness at fifty: Insights from

neuroscience. *Psychological Review*, *123*(4), 1–19. https://doi.org/10.1037/rev0000033

Manohar, S. G., Chong, T. T. J., Apps, M. A. J., Batla, A., Stamelou, M., Jarman, P. R., Bhatia,

K. P., & Husain, M. (2015). Reward Pays the Cost of Noise Reduction in Motor and

Cognitive Control. *Current Biology*, *25*(13), 1707–1716.

https://doi.org/10.1016/j.cub.2015.05.038

Moscarello, J. M., & Hartley, C. A. (2017). Agency and the Calibration of Motivated Behavior.

Trends in Cognitive Sciences, *21*(10), 725–735. https://doi.org/10.1016/j.tics.2017.06.008

Nagase, A. M., Onoda, K., Foo, J. C., Haji, T., Akaishi, R., Yamaguchi, S., Sakai, K., & Morita,

K. (2018). Neural mechanisms for adaptive learned avoidance of mental effort. *Journal of*

Neuroscience, *38*(10), 2631–2651. https://doi.org/10.1523/JNEUROSCI.1995-17.2018

Nalborczyk, L., & Bürkner, P.-C. (2019). An Introduction to Bayesian Multilevel Models Using

brms: A Case Study of Gender Effects on Vowel Variability in Standard Indonesian.

Journal of Speech, Language, and Hearing Research.

Nassar, M. R., Bruckner, R., & Frank, M. J. (2019). Statistical context dictates the relationship

between feedback-related EEG signals and learning. *ELife*, *8*, 1–26.

https://doi.org/10.7554/eLife.46975

Nigbur, R., Schneider, J., Sommer, W., Dimigen, O., & Stürmer, B. (2015). Ad-hoc and context-

dependent adjustments of selective attention in conflict control: An ERP study with visual

probes. *NeuroImage*, *107*, 76–84. https://doi.org/10.1016/j.neuroimage.2014.11.052

Niv, Y., Edlund, J. A., Dayan, P., & O'Doherty, J. P. (2012). Neural prediction errors reveal a

risk-sensitive reinforcement-learning process in the human brain. *Journal of Neuroscience*,

32(2), 551–562. https://doi.org/10.1523/JNEUROSCI.5498-10.2012

Niv, Y., Joel, D., & Dayan, P. (2006). A normative perspective on motivation. *Trends in*

Cognitive Sciences, *10*(8), 375–381. https://doi.org/10.1016/j.tics.2006.06.010

Otto, A. R., & Daw, N. D. (2019). The opportunity cost of time modulates cognitive effort.

Neuropsychologia, *123*(October 2017), 92–105.

https://doi.org/10.1016/j.neuropsychologia.2018.05.006

Pelli, D. D. (1997). The VideoToolbox software for visual psychophysics: transforming numbers

into movies. *Spatial Vision*, *10*(4), 437–442.

R Core Team. (2017). *R: A Language and Environment for Statistical Computing.* R Foundation

for Statistical Computing.

Rutledge, R. B., Skandali, N., Dayan, P., & Dolan, R. J. (2014). A computational and neural

model of momentary subjective well-being. *Proceedings of the National Academy of*

Sciences, *111*(33), 12252–12257. https://doi.org/10.1073/pnas.1407535111

Schevernels, H., Krebs, R. M., Santens, P., Woldorff, M. G., & Boehler, C. N. (2014). Task

preparation processes related to reward prediction precede those related to task-difficulty

expectation. *NeuroImage*, *84*, 639–647. https://doi.org/10.1016/j.neuroimage.2013.09.039

Shenhav, A., Botvinick, M., & Cohen, J. (2013). The expected value of control: An integrative

theory of anterior cingulate cortex function. *Neuron*, *79*(2), 217–240.

https://doi.org/10.1016/j.neuron.2013.07.007

Shenhav, A., Musslick, S., Lieder, F., Kool, W., Griffiths, T. L., Cohen, J. D., & Botvinick, M.

M. (2017). Toward a Rational and Mechanistic Account of Mental Effort. *Annual Review of*

Neuroscience, *40*(1), 99–124. https://doi.org/10.1146/annurev-neuro-072116-031526

Shenhav, A., Prater Fahey, M., & Grahek, I. (submitted). *Decomposing the motivation to exert*

mental effort.

Sutton, R. S., & Barto, A. G. (2018). *Reinforcement Learning: An Introduction*. MIT Press.

Verbeke, P., & Verguts, T. (2019). Learning to synchronize: How biological agents can couple

neural task modules for dealing with the stability-plasticity dilemma. *PLoS Computational*

Biology, *15*(8), 1–25. https://doi.org/10.1371/journal.pcbi.1006604

Verguts, T., Vassena, E., & Silvetti, M. (2015). Adaptive effort investment in cognitive and

physical tasks: a neurocomputational model. *Frontiers in Behavioral Neuroscience*,

9(March). https://doi.org/10.3389/fnbeh.2015.00057

Wagenmakers, E. J., Lodewyckx, T., Kuriyal, H., & Grasman, R. (2010). Bayesian hypothesis

testing for psychologists: A tutorial on the Savage-Dickey method. *Cognitive Psychology*,

60(3), 158–189. https://doi.org/10.1016/j.cogpsych.2009.12.001

Yeung, N., & Sanfey, A. G. (2004). Independent coding of reward magnitude and valence in the

human brain. *Journal of Neuroscience*, *24*(28), 6258–6264.

https://doi.org/10.1523/JNEUROSCI.4537-03.2004

740