Learning when effort matters: Neural dynamics underlying updating and adaptation to changes in performance efficacy

Ivan Grahek[1†*], Romy Frömer[1†], Mahalia Prater Fahey[1], & Amitai Shenhav[1]

1. *Department of Cognitive, Linguistic, & Psychological Sciences and Carney Institute for Brain Science, Brown University, Providence, RI 02912, USA*

[†] These authors have contributed equally to the study.

* Corresponding author. E-mail address: ivan_grahek@brown.edu

Conflicts of interest: The authors have no conflicts of interest to declare.

Abstract

To determine how much cognitive control to invest in a task, people need to consider whether exerting control *matters* for obtaining rewards. In particular, they need to account for the efficacy of their performance – the degree to which rewards are determined by their performance or by independent factors. Yet it remains unclear how people learn about their *performance efficacy* in an environment. Here we combined computational modeling with measures of task performance and EEG, to provide a mechanistic account of how people (a) learn and dynamically update efficacy expectations in a changing environment, and (b) proactively adjust control allocation based on their current efficacy expectations. Across two studies subjects performed an incentivized cognitive control task while their performance efficacy (the likelihood that rewards are performance-contingent or random) varied over time. We show that people learn about efficacy through a neural mechanism similar to the one used to learn from rewards, and that they use this information to adjust how much control they allocate. Using a computational model we show that these control adjustments reflect changes in information processing, rather than the speed-accuracy tradeoff. These findings demonstrate the neurocomputational mechanism through which people learn how worthwhile their cognitive control is.

Keywords: Motivation; Cognitive control; Learning; Performance efficacy; Reward rate; EEG

Cognitive control is critical for achieving most goals, but it is effortful (Botvinick & Cohen, 2014; Shenhav et al., 2017). Determining whether and how much control to invest in a task requires a person to weigh these effort costs against the potential benefits of control (Kool & Botvinick, 2018; Manohar et al., 2015; Verguts et al., 2015). These potential benefits not only reflect the amount of reward that can be expected for performing well on a task but also, and critically, the extent to which performance matters for obtaining those rewards (the *efficacy* of performance) (Frömer et al., 2021a; Shenhav et al., 2013; Shenhav et al., 2021). For instance, a student may be more motivated to invest effort into writing an essay for a competition if they believe that this essay will play an important role in determining their success than if they believe that the competition will be primarily determined by performance-unrelated factors (e.g., social status, idiosyncratic criteria). The amount of cognitive effort we invest in a task is therefore substantially determined by how much efficacy we expect to have with that task. Such efficacy expectations must be derived from past experiences that indicated how worthwhile our efforts were in a given environment. Understanding how people come to learn when their efforts are efficacious is critically important for understanding why they come to be more or less motivated in certain environments, which is particularly relevant for understanding mental disorders (Grahek et al., 2019; Moscarello & Hartley, 2017). However, in spite of its importance, relatively little is known about the mechanisms by which people learn how efficacious their cognitive efforts are.

Past work has provided critical insights into related aspects of how people adapt cognitive control based on learned incentives. For instance, a substantial body of research has examined how people learn about the amount and frequency of reward in their environment (Daw & Tobler, 2013; Dolan & Dayan, 2013; Niv et al., 2006), including the extent to which

learning differs depending on the valence of feedback (e.g., whether an outcome was better or worse than expected; Chambon et al., 2020; Collins & Frank, 2014; Garrett & Daw, 2020; Lefebvre et al., 2017). While this work has traditionally investigated the relationship between learned value and action selection, recent work has shown that people also adapt their control states based on learned value (Lieder et al., 2018). People allocate more or less cognitive control to a task depending on the reward they have come to expect based on recent trials (Bustamante et al., 2021; Otto & Daw, 2019). Similarly, people adapt their control based on the expected difficulty of a given task (for a review see: Chiu & Egner, 2019). Behavioral and electrophysiological evidence shows that people allocate more control when, based on previous learning, they expect a task to be more difficult (Bejjani et al., 2018; Bugg et al., 2011; Jiang et al., 2020; Nigbur et al., 2015). Research into these difficulty-related control adaptations provides insight into how people learn about and adapt to changes in the amount of cognitive control required to achieve a given level of performance. However, in the real world, achieving that level of performance may not always have bearing on whether a person will be rewarded (Shenhav et al., 2021). Research on cognitive control adaptation has yet to examine how people learn and adapt as the performance efficacy of their environment changes.

A parallel body of work has shown that people adjust their actions based on learned estimates of how likely it is that a given action leads to a desired outcome (action-outcome contingency; Dickinson & Balleine, 1995; Ly et al., 2019; Moscarello & Hartley, 2017; Dorfman et al., 2019; Dorfman et al., 2021; Ligneul, 2021). People preferentially, and more vigorously, select actions when those actions are more reliable for producing the desired outcomes (i.e., as action-outcome contingencies increase; Liljeholm et al., 2011; Manohar et al., 2017). How people learn and adapt to analogous contingencies for cognitive performance is less well

understood, in part because feedback for task performance typically conflates the contingencies between performance and outcomes (i.e., the likelihood that better performance will lead to better outcomes [*performance efficacy*]) – which are most analogous to action-outcome contingencies described above – with perceived contingencies between control allocation and performance (i.e., the likelihood that exerting cognitive effort will lead to better performance [*control efficacy*]) (Shenhav et al., 2021).

We have recently shown that these two sets of contingencies can be disentangled to examine how control allocation is uniquely influenced by expected levels of performance efficacy (Frömer et al., 2021a). Using a standard cognitive control task (Stroop), we varied whether the rewards earned on each trial were contingent on performance (high efficacy) or determined at random (low efficacy). We showed that people allocate more control when they expect to have high compared to low efficacy, reflected in higher amplitudes of an EEG index of proactive control (the contingent negative variation [CNV]) and in improved behavioral performance. These results demonstrate that participants leverage expectations about the extent to which their performance matters when deciding how much cognitive effort to invest in a task. However, in this work, participants were explicitly cued with the level of performance efficacy to expect on a given trial, and those predictive cues retained the same meaning across the session. Thus, how it is that people learn these efficacy expectations in more typical environments where contingencies are not instructed, and how they dynamically update their expectations as contingencies change, remains unanswered.

In the current study, we use computational modeling and single-trial EEG analyses to study the mechanisms through which efficacy estimates are learned, updated, and used to guide control allocation. Across two experiments, we show that efficacy estimates can be learned from

feedback through iterative updating based on weighted prediction errors (Sutton & Barto, 2018).

Using model-based analyses of a putative neural marker of learning (P3b; Fischer & Ullsperger,

2013; Nassar et al., 2019), we show that efficacy prediction errors and corresponding learning

rates can be used to predict how efficacy estimates are adjusted following relevant changes in the

environment. We further show that efficacy estimates learned in this way are used to guide the

allocation of control. We extend our previous findings (Frömer et al., 2021a) to show that a

neural marker of control allocation (the CNV; Grent-'t-Jong & Woldorff, 2007; Schevernels et

al., 2014) increases with increasing model-based efficacy estimates. Using a drift diffusion

model (Ratcliff & McKoon, 2008), we further show that the performance improvements related

to increased performance-efficacy reflect facilitation of task-related information processing

(reflected in increased drift rates), rather than changes in the speed-accuracy tradeoff (i.e.,

thresholds). Taken together, these results show that efficacy estimates are learned and updated

based on feedback, leveraging general cognitive and neural mechanisms of predictive inference.

Materials and Methods

Study 1

Participants

We recruited forty-one participants with normal or corrected-to-normal vision from the

Brown University subject pool. One participant was excluded due to technical issues. The final

data set included 40 participants (24 females, 16 males; median age = 19). Participants gave

informed consent and were compensated with course credits or a fixed payoff of $20. In

addition, they received up to $5 bonus that depended on their task performance ($3.25 on

average). The research protocol was approved by Brown University's Institutional Review

Board.

Experimental design

In the main task, taking approximately 45 minutes, participants performed 288 Stroop trials (Figure 1A). Each trial started with the presentation of a cue (grey circle) that remained on the screen throughout the trial. After a period of 1500 ms, a Stroop stimulus was superimposed until a response was made or 1000 ms elapsed, at which time it was sequentially replaced with two types of feedback presented for 1000 ms each. Each trial onset was preceded by a fixation cross (randomly jittered between 1000 and 1500ms). Participants responded to the ink color of the Stroop stimulus (red, green, blue, or yellow) by pressing one of four keyboard keys (D, F, J, and K). Stimuli were either color words same as the ink color (congruent, n = 108) or different (incongruent, n = 108), or a string of letters "XXXXX" (neutral, n= 72). Feedback informed them whether they obtained a reward (reward, "$50c" or no reward, "$0c") and whether the reward they received depended on their performance (performance-based feedback, a button graphic), or not (random feedback, a dice graphic). The order of the two types of feedback was pseudo-randomized with half of the trials showing reward feedback first and the other half efficacy feedback. Every 2-4 trials the feedback was followed by a probe of efficacy ("How much do you think your rewards currently depend on your performance?") or reward rate ("How often do you think you are currently being rewarded?") to which participants responded on a visual analog scale ranging from 0 to 100. The number and timing of the probes was randomized per subject resulting in a median of 45 efficacy probes (SD=3.38) and 47 reward probes (SD=2.81).

Efficacy (performance-based or random rewards) on each trial was sampled from a binomial distribution with probabilities ranging between 0.1 and 0.9 that drifted over the course of the experiment and were predetermined (Figure 1B). In order to ensure that the performance-

based and random trials did not differ in reward rate, reward feedback for the random trials was sampled from the moving window of the reward feedback of the previous 10 performance-based trials. At the beginning of the experiment a window with 8 rewards and 2 no rewards were used to reflect the calibrated accuracy, and was then updated after every trial to reflect any changes in performance accuracy. Thus, reward rate was not experimentally manipulated in the experiment and remained constant.

Prior to the main task, participants performed several practice phases of the Stroop task (approximately 15 minutes). First, they practiced the mappings between colors and keyboard keys (80 trials). Then they completed a short practice of the Stroop task with written feedback ("correct" or "incorrect") on each trial (30 trials). Participants then completed 100 more of such trials during which we individually calibrated the reaction time deadline such that participants yielded approximately 80% reward rate. In the final practice phase participants were introduced to the two types of feedback which they would see in the main experiment (30 trials).

The experimental task was implemented in Psychophysics Toolbox (Brainard, 1997; Kleiner, Brainard, & Pelli, 2007; Pelli, 1997) for Matlab (MathWorks Inc.) and presented on a 23 inch screen with a 1920 x 1080 resolution. All of the stimuli were presented centrally while the participants were seated 80 cm away from the screen.

Psychophysiological recording and preprocessing

EEG data were recorded at a sampling rate of 500 Hz from 64 Ag/AgCl electrodes mounted in an electrode cap (ECI Inc.), referenced against Cz, using Brain Vision Recorder (Brain Products, München, Germany). Vertical and horizontal ocular activity was recorded from below both eyes (IO1, IO2) and the outer canthi (LO1, LO2), respectively. Impedances were kept below 10 kΩ. Offline, data were processed using custom made Matlab scripts (Frömer et al.,

2018) employing EEGlab functions (Delorme & Makeig, 2004). Data were re-referenced to average reference, ocular artifacts were corrected using brain electric source analyses (BESA; Ille, Berg, & Scherg, 2002) based on separately recorded prototypical eye movements. The cleaned continuous EEG was then low pass filtered at 40 Hz and segmented into epochs around cue onset (-200 to 1500 ms), stimulus onset, and both efficacy and reward feedback (-200 to 800 ms). Baselines were corrected to the average of each 200 ms pre-stimulus interval. Segments containing artifacts, values exceeding $\pm$ 150 μV or gradients larger than 50 μV, were excluded from further analyses.

Single trial ERPs were then exported for further analyses in R (R Core Team, 2017). The late CNV was quantified between 1000 and 1500 ms post neutral cue onset (cf. Frömer et al., 2016, 2021; Schevernels et al., 2014) as the average activity over 9 fronto-central electrodes (Fz, F1, F2, FCz, FC1, FC2, Cz, C1, and C2). The P3b was quantified between 350 and 500 ms (cf. Fischer & Ullsperger, 2013) for both reward and efficacy feedback and calculated as the averaged activity over 9 centro-parietal electrodes (Pz, P1, P2, POz, PO1, PO2, CPz, CP1, and CP2).

Learning models and statistical analyses

Learning models. In order to obtain trial-by-trial estimates of efficacy and reward rate, we fitted two temporal difference learning models (Gläscher et al., 2010; Sutton & Barto, 2018) to the continuous subjective estimates of efficacy and reward rate (cf. Eldar, Hauser, Dayan, & Dolan, 2016; Nagase et al., 2018; Rutledge et al., 2014). The first model ("1 learning rate model") assumed that the estimate of efficacy for the next trial (E_{t+1}) depended on the current efficacy estimate (E_t) and the prediction error (δ_t) weighted by a constant learning rate (α):

$$E_{t+1} = E_t + \alpha * \delta_t$$

Where $0 \leq \alpha \leq 1$, and the prediction error is calculated as the difference between the contingency feedback on the current trial (e_t) and the efficacy estimate on that trial: $\delta_t = e_t - E_t$. The model started from an initial value (free parameter) and updated the subjective estimate based on the binary feedback on each trial. For example, assuming a learning rate of 0.5 and the initial value of 0.5, the model would update the initial estimate following efficacy feedback signaling "performance-based" $(e_t = 1)$ to 0.75. If on the next trial contingency feedback was "random" $(e_{t+1} = 0)$ the subjective estimate would drop to 0.6. Exactly the same model was used to calculate estimates of reward on each trial. The model was fitted separately to the subjective estimates of efficacy and reward rate with only the learning rate as a free parameter. The second model ("2 learning rates model") was the same as the first model, but it included two learning rates: one learning rate for learning from the efficacy feedback, and another for learning from the no efficacy feedback. Finally, as a baseline, we also included the "intercept model" which did not update the efficacy estimate throughout the experiment, but just assumed that the estimate took one constant value. The models were fit hierarchically to the data using maximum likelihood (using mfit (https://github.com/sjgershm/mfit); for details see the MATLAB scripts available on GitHub).

Statistical analyses. The efficacy and reward rate estimates obtained through fitting the learning model were then used to analyze the behavioral and EEG data. To this end, we fitted Bayesian multilevel regressions to predict subjective estimates of efficacy and reward rates, reaction times, accuracy, as well as the CNV and P3b amplitudes. Subjective estimates of efficacy and reward rate were regressed onto efficacy or reward feedback. Reaction times and accuracies were regressed onto trial-by-trial model-based estimates of efficacy and reward rate, as well as trial-by-trial CNV amplitude, while controlling for congruency. The P3b amplitudes were analyzed in

two ways: with trial-by-trial model-based estimates of efficacy and reward rate and current

feedback as predictors, and with model-based prediction errors and learning rates for each

feedback type. CNV amplitudes were regressed onto trial-by-trial model-based estimates of

efficacy. All of the fitted models controlled for the influence of the reward rate estimates.

Parallel analyses were done to predict the P3b in response to reward feedback, while controlling

for the efficacy estimates.

The regression models were fitted in R with the *brms* package (Bürkner, 2016) which

relies on the probabilistic programming language *Stan* (Carpenter et al., 2016) to implement

Markov Chain Monte Carlo (MCMC) algorithms and estimate posterior distributions of model

parameters. The analyses were done based on the recommendations for Bayesian multilevel

modeling using *brms* (Bürkner, 2016, 2017; Nalborczyk & Bürkner, 2019). The fitted models

included constant and varying effects (also known as fixed and random) with weakly informative

priors (except for the behavioral and CNV analyses, see below for details) for the intercept and

the slopes of fixed effects and the likelihoods appropriate for the modeled data (Ex-Gaussian for

reaction times, Bernoulli for accuracy, and Gaussian for the subjective estimates and the EEG

amplitudes). The fitted models included all of the fixed effects as varying effects. All of the

continuous predictors in the model were centered and the categorical predictors were contrast

coded. Four MCMC simulations ("chains"; 20,000 iterations; 19,000 warmup) were run to

estimate the parameters of each of the fitted models. The convergence of the models was

confirmed by examining trace plots, autocorrelation, and variance between chains (Gelman-

Rubin statistic; Gelman & Rubin, 1992). After convergence was confirmed, we analyzed the

posterior distributions of the parameters of interest and summarized them by reporting the means

of the distribution for the given parameter (*b*) and the 95% credible intervals (95% CrI) of the

posterior distributions of that model. We report the proportion of the posterior samples on the relevant side of 0 (e.g., $p_{b < 0} = 0.9$), which represents the probability that the estimate is below or above 0. We also report Bayes factors (BF) calculated using the Savage-Dickey method (Wagenmakers, Lodewyckx, Kuriyal, & Grasman, 2010). We report the BFs in support of the alternative hypothesis against the null (BF_{10}), except for the analyses of accurate RT, accuracies, and CNV amplitude in which we have informative priors based on our previous study (Frömer et al., 2021a), and in which case we support the evidence in favor of the null (BF_{01}).

To compare the positive and negative learning rates we fitted a model in which the learning rates were predicted by the learning rate type (Kruschke, 2013). In this model we used Gaussian distributions (mean, standard deviation) as priors (intercept: (0.5,0.5); slopes: (0,0.5)).

We fitted two separate models to predict the subjective estimates of efficacy and reward rate based on previous feedbacks. At each timepoint the estimates were predicted by the current, and previous 4 feedbacks. The feedback on each of the trials (performance-based vs. random or reward vs. no reward) was entered as a constant effect and the models also included the intercept as a varying effect. As the subjective estimates could vary between 0 and 1, we used Gaussian distributions (intercept: (0.5,0.2); slopes: (0,0.2)) as priors.

For predicting the P3b amplitude in response to the onset of the efficacy feedback, we fitted two models. First, we fitted a model which included the model-based estimate of efficacy, the actual feedback, and the interaction between the expected efficacy and the observed efficacy feedback. Additionally, we controlled for the reward rate estimate. Second, we fitted separate models which included the model-based prediction errors, the influence of the between-subject learning rates, and their interaction with the prediction errors, while controlling for the estimate of the reward rate. In this analysis the learning rates (one for feedback type for each subject)

were mean-centered within subjects and thus any effect of the learning rates is driven by the

difference between the random and the performance-based learning rate. For these models we

selected wide Gaussian priors (intercept: (5,3); slopes (0,3)). The same logic in building models

was applied for the analyses of the reward feedback. In these analyses we focused on the reward

feedback processing and how it interacted with the model-based estimates of reward rates, while

controlling for the model-based estimates of efficacy. We analyzed only the trials with correct

responses for both the efficacy and the reward feedback analyses.

To test the influence of efficacy on the late CNV, we fitted a model which predicted the

CNV based on the model-based efficacy estimates, while controlling for the effect of the reward

rate estimates. Drawing on the results of our previous study (Frömer et al., 2021a), this model

included Gaussian priors for the intercept (-0.16, 1.59) and the efficacy (-0.30, 0.73) and reward

(0,0.73) slopes.

For predicting reaction times and accuracy we fitted models which included congruency

(Facilitation: difference between neutral and congruent trials; Interference: difference between

incongruent and neutral trials) and the model-based efficacy estimates, while controlling for the

reward rate estimates. We used Gaussian distributions as informative priors based on our

previous study (Frömer et al., 2021a), for both the reaction times (intercept: (624, 58.69);

facilitation (15.54, 21.93), interference (61.12, 37.49); efficacy (-10.26, 19.51); reward (0,

19.51)) and accuracy[1] analyses (intercept: (2.11, 0.81); facilitation (-0.45, 0.64), interference (-

0.53, 0.81); efficacy (0.09, 0.32); reward (0, 0.32)).

To investigate how the late CNV influences the behavior, we fitted two models in which

we predicted the reaction times and accuracy based on the CNV amplitude. The prior

____

[1] Note that the prior distributions are set in log-odds.

distributions for these models were weakly informative Gaussian distributions for predicting

both the reaction times (intercept: (650, 200); slope: (0, 50)) and accuracy (intercept: (0.7, 0.2);

slope: (0, 0.2)).

To visualize the topographies of the relevant ERP effects, we fitted the relevant models to

all 64 channels and then plotted the posterior estimates of the effects of interest at each electrode

(cf. Frömer et al., 2021b).

Study 2

Participants

We recruited eighty-seven participants residing in the United States from Prolific – an

online platform for data collection. Participants had normal or corrected-to-normal vision and

gave informed consents. They were compensated with a fixed payoff of $8 per hour (median

completion time of 74 minutes) plus a monetary bonus based on points earned during the task

($1 on average). The research protocol was approved by Brown University's Institutional

Review Board.

We a priori excluded participants who did not pass attention checks (N=8) or who took

substantially longer than the average participant to complete the study (N=2 participants who

took over 130 minutes), suggesting that they did not sustain attention to the experiment over that

time. We fit our learning models to data from the remaining 77 subjects, and then excluded

participants whose performance suggested inattention to the overall task (based on accuracies

less than 70% across all trials – including the trials in which performance efficacy was low, N=6)

or inattention to the task cues and/or feedback probes (based on low learning rates; N=20). To

identify participants with exceedingly low learning rates, we submitted all positive and negative

efficacy learning rates to unsupervised Gaussian mixture models (as implemented in the Mclust

package; (Scrucca et al., 2016) to determine the best fitting number and shape of clusters (model

comparison via BIC). This procedure identified four clusters of subjects with different overall

learning rates (Figure S5B-C) and we excluded subjects from the first cluster as they all had very

low learning rates relative to the other participants (both learning rates < 0.03). The final sample

included 51 participants (31 females, 20 males; median age = 29).

Experimental design

In order to better understand the computational mechanisms that lead to improved

behavioral performance in high efficacy states (Study 1), we wanted to fit a Drift Diffusion

model (DDM; Ratcliff & McKoon, 2008) to our behavioral data. If people allocate more

attention when they think they have high performance efficacy, this should be observed as an

increase in the drift rate (speed of evidence accumulation). However, Study 1 included a tight

respond deadline for earning a reward, making it more challenging to fit the DDM. To avoid this

issue, in Study 2 we adjusted the task to a free response paradigm which allowed us to

investigate drift rate and threshold adjustments (cf. Leng et al., 2021). We used this design to test

the hypothesis that higher efficacy estimates should predict increased drift rates.

Instead of single trials, participants now completed 288 intervals during which they could

respond to as many trials (congruent and incongruent, removing the neutral condition) as they

wished within a fixed time window (randomly selected between 2000, 3000, or 4000ms). Apart

from this, the structure of the task remained the same: participants saw a fixation cross (1500

ms), then completed as many trials as they wished during a fixed interval, followed by the

feedback (1500 ms) on how many points they earned and whether this was based on their

performance or awarded to them based on random chance. Note that participants now received

continuous reward feedback (the number of points instead of the binary reward-no reward in

Study 1), but that the same yoking procedure as in Study 1 was employed ensuring that the amount of reward they got in performance-based and random intervals was matched (rewards on the random intervals were sampled from the moving average window of the past 10 performance-based intervals). As in Study 1, participants were probed every 2-4 intervals to estimate either how much they thought their rewards depended on their performance, or how much reward they were earning. We again implemented an efficacy drift (modified, but comparable to the drift in Study 1; Figure S5A), now across the 288 intervals of the task.

We gamified the task in order to make it more appealing for the participants. Instead of the Stroop task, we used a picture-word interference task in which four grey-scaled images of fruit (apple, pear, lemon, and peach) were overlaid by those fruit words. Participants responded to the image, while ignoring the word, by pressing one of the 4 corresponding keys. They first practiced this task, and then were introduced with a cover story telling them that they are in the garden and need to water the fruit in little patches by pressing the correct keys. They were instructed that they will be moving through a garden and that in some patches watering will directly translate into how many points they will be earning, while in the others that will not be the case (the efficacy drift). The experiment was implemented in Psiturk (Gureckis et al., 2016) and the participants performed the task on their own computers and were required to have a keyboard.

Learning models, Drift Diffusion model, and statistical analyses

Learning models. We fitted the same set of learning models as in Study 1, performed model comparison, and got the interval-by-interval model-based estimates of performance efficacy and reward. Note that in this version of the task participants earned points in each interval, unlike the binary rewards (reward vs. no reward) in Study 1. This meant that the reward learning model

learned reward magnitudes rather than reward probabilities. However, model fitting and the further analyses were the same as in Study 1.

Statistical analyses. We fitted the same Bayesian multilevel models as in Study 1 to predict the influence of previous efficacy feedbacks on the subjective efficacy estimates, as well as the influence of the model-based efficacy estimates on reaction times and accuracies. For the analyses of the subjective efficacy estimates we used the same priors as in Study 1. For the analyses of the reaction times and accuracies, we used the posterior distributions obtained in Study 1 as the informative priors for the congruency, efficacy, and reward effects. The reaction times and accuracy models also controlled for the effects of the average congruency level in the interval and the interval length.

Drift Diffusion model. This model decomposes participant's behavior into drift rate (the speed of evidence accumulation) and response threshold (the level of caution), allowing us to investigate which of these two components is affected by the efficacy estimates. We fitted the model using Bayesian hierarchical estimation as implemented in the HDDM package (Wiecki, Sofer, & Frank, 2013). The fitted model included the main effects of efficacy and reward rate estimates onto both drift rate and threshold, and included the effect of congruency on the drift rate. The responses were coded as correct or incorrect, and trials with reaction times below 250ms were excluded. All of the effects were allowed to vary across subjects, and we ran five MCMC chains (12,000 iterations; 10,000 warmup). We confirmed convergence by examining trace plots and variance between chains.

Results

        To investigate how efficacy estimates are learned, and how they affect control allocation,

in Study 1 we recorded EEG while 40 participants performed a modified version of the Stroop

task (Figure 1A). Across trials we varied whether reward outcomes ($0.50 vs. $0.00) were

determined by a participant's performance on a given trial (responding accurately and below a

pre-determined response time criterion; *performance-based* trials) or whether those outcomes

were determined independent of performance (based on a weighted coin flip; *random* trials).

Over the course of the session, we gradually varied the likelihood that a given trial would be

performance-based or random such that, at some points in the experiment, most trials were

performance-based (high efficacy level), and at other points most trials had random outcomes

(low efficacy level) (Figure 1B). Importantly, unlike in our previous study (Frömer et al., 2021a),

participants were not told whether a given trial would be performance-based (maximal efficacy)

or random (minimal efficacy), but instead had to estimate their current efficacy level based on

recent trial feedback, which indicated both the reward outcome ($0.50 vs. $0.00) and how that

outcome was determined (performance-based [button symbol] vs. random [dice symbol]). We

held reward rate constant across both feedback types by yoking reward rate on random trials to

the reward rate on performance-based trials, and counter-balanced the time-course of the gradual

change in efficacy (see Methods for details). To capture changes in efficacy expectations over

the course of the session, we probed participants every 4-8 trials (averaging 44.3 probes per

participant) to report their current estimates of efficacy. These efficacy probes were interleaved

with probes asking participants to estimate the current reward rate, serving as foils and for

control analyses.

Figure 1. Manipulating expected efficacy and assessing learning in Study 1. A. Trial Schematic. On each trial participants saw a cue (gray circle), predicting the onset of a Stroop stimulus (target), and were then sequentially presented with reward and efficacy feedback. On half of the trials, efficacy feedback was presented first, and on the other half reward feedback was presented first. Every 2-4 trials participants were subsequently asked to estimate their current efficacy level or reward rate. **B. Efficacy manipulation.** We let the probability of performance-based vs. random feedback continuously drift over the course of the experiment (inversed for one half of the sample). Arrows mark time points with low and high efficacy, respectively. When efficacy was low, rewards were more likely to be random, whereas when efficacy was high, rewards were more likely to be performance-based.

Participants dynamically update efficacy expectations based on feedback

To determine whether and how participants learned their current efficacy levels, we first analyzed the influence of previous efficacy feedback (whether outcomes on a given previous trial had been performance-based or random) on one's current subjective estimates of efficacy. We found very strong evidence that participants adjusted their subjective efficacy upward or downward depending on whether the most recent trial was Performance-Based or Random ($b = 0.14$; 95% CrI [0.12, 0.16]; $p_{b < 0} = 0$; $BF_{10} > 100$), and that this remained true (but to diminishing degrees) up to five trials back (all $p_{b < 0} < 0.01$; Figure 2A; Table S1). These results suggest that participants were dynamically updating their efficacy estimates based on efficacy feedback.

Figure 2. Efficacy learning is captured by a reinforcement learning model separate learning rates for performance-based and random feedback in Study 1. A. Efficacy estimates track recent efficacy feedback. Regression weights for the influence of the current and previous contingent vs. random feedback on the subjective efficacy estimate. Error bars represent 50% and 95% highest density intervals. **B. A separate learning rate model captures efficacy learning best.** BICs of fitted intercept only and one learning rate model relative to two learning rate models are plotted for each participant and favor the two learning rate model. **C. Learning rate biases vary between participants.** Positive and negative learning rate estimates are plotted for all participants. Points below the diagonal indicate higher learning rates for performance-based compared to random feedback, and points above the opposite. **D. Example learning trajectories.** Subjective and model-based efficacy estimates, and a running average of the previous 5 efficacy feedbacks, for a participant with a higher learning rate for the contingent compared to random feedback (upper) and a participant with the reverse bias (lower).

This pattern of learning was accounted for by a standard reinforcement learning (RL) algorithm, the delta-rule, according to which efficacy estimates are updated in proportion to the prediction error between the expected efficacy and the efficacy feedback on a given trial (i.e., whether a given outcome was determined by performance or not) (Sutton & Barto, 2018).

Interestingly, consistent with studies of reward learning (Chambon et al., 2020; Collins & Frank, 2014; Garrett & Daw, 2020; Lefebvre et al., 2017; Niv et al., 2012), we found that the RL model that best accounted for our data allowed efficacy estimates to be updated differently from trials that were more efficacious than expected (Performance-Based feedback) than from trials that were less efficacious than expected (Random feedback), by having separate *learning rates* scaling prediction errors in the two cases. Even when accounting for model complexity, this Two Learning Rate model outperformed a One Learning Rate model as well as a baseline model that fits a single constant efficacy estimate and no learning (Intercept Model) (Figure 2B). We found that the two learning rates for this best-fit model varied across the group (Figure 2E), but did not find that one was consistently higher than the other ($b = 0.02$; 95% CrI [-0.04, 0.08]; $p_{b<0} = 0.260$; $BF_{01}=13.55$), despite most participants (80%) tending to learn more from Performance-Based than Random trials. Taken together, these results show that participants dynamically updated their expected efficacy based on trial-by-trial feedback, and that they did so differentially based on whether the trial was more or less efficacious than expected. The fitted models further enable us to generate trial-by-trial estimates of expected efficacy and efficacy prediction errors, which we use in model-based analyses of behavior and neural activity below. Note that in all the following analyses we control for reward rate estimates obtained from models fit to reward feedback (for details see below).

The feedback-related P3b indexes updating of efficacy expectations

To investigate the neural mechanism underlying feedback-based learning of efficacy, we probed the centroparietal P3b ERP component (Figure 3A), an established neural correlate of prediction-error based learning (Fischer & Ullsperger, 2013; Nassar, Bruckner, & Frank, 2019;

Frömer et al., 2021b). If the P3b indexes feedback-based updating of *efficacy* predictions, as it does for reward predictions, we would expect this ERP to demonstrate several key markers of such a learning signal. First, we would expect the P3b to reflect the extent to which efficacy feedback (Performance-Based vs. Random) deviates from the current level of expected efficacy. In other words, the P3b should track the magnitude of the *unsigned efficacy prediction error (PE)* on a given trial. We tested this by examining how the amplitude of the P3b to a given type of efficacy feedback varied with model-based estimates of the participant's efficacy expectation on that trial, while holding the expected reward rate constant (see the Methods and Materials sections for the details of the experimental design and the statistical models). If the P3b signaled efficacy PE then its amplitude should scale inversely with the expected probability of a given type of feedback (i.e., how *unexpected* that feedback is), thus correlating *negatively* with expected efficacy on trials providing Performance-Based feedback and correlating *positively* with expected efficacy on trials providing Random feedback. In addition to overall higher P3b to performance-based compared to random feedback ($b = 0.86$; 95% CrI [0.42, 1.31]; $p_{b<0} = 0$; BF_{10} =30.86; Figure 2B), we found exactly this predicted interaction between feedback type and expected efficacy ($b = -2.40$; 95% CrI [-4.07, -0.74]; $p_{b>0} = 0$; $BF_{10} = 24.40$; Figure 2B; Table S2), with the P3b amplitude decreasing with model-based estimates of expected efficacy on Performance-Based trials (b = -1.49; 95% CrI [-2.72, -0.27]; $p_{b>0} = 0.03$; $BF_{10} = 1.52$) and increasing numerically, but not robustly with expected efficacy on Random trials ($b = 0.91$; 95% CrI [-0.34, 2.21]; $p_{b<0} = 0.12$; $BF_{10} = 0.44$). Accordingly, when we regressed P3b amplitude on our model-based estimates of trial-to-trial efficacy PE, we found a positive relationship ($b = 1.25$; 95% CrI [0.35, 2.15]; $p_{b<0} = 0.01$; $BF_{10} = 5.84$; Figure 3C-left; Table S2).

In addition to tracking efficacy PEs, the second key prediction for the P3b if it indexes efficacy learning, is that it should track the extent to which PEs are used to update estimates of efficacy (i.e., the learning rate). In the current study, we found that participants differed in their learning rates for the two forms of efficacy feedback (Performance-Based vs. Random), providing us with a unique opportunity to perform a within-subject test of whether P3b tracked differences in learning rate across these two conditions. Specifically, we could test whether a given subject's P3b was greater in the feedback condition for which they demonstrated a higher learning rate. We found some evidence in this direction, with the P3b demonstrating an increase for the within-subject feedback condition with the higher learning rate, but this result was not conclusive ($b = 2.00$; 95% CrI [-2.21, 6.04]; $p_{b<0} = 0.17$; BF = 1.08; Figure 3C-right). While, theoretically, prediction error and learning rate would interact in predicting the P3b amplitude, we did not observe such an interaction here. The fact that we only found additive effects of prediction errors and learning rate on P3b is, however, in line with previous findings on reward processing (Fischer & Ullsperger, 2013).

The CNV indexes control allocation based on updated expectations of efficacy

Thus far, our findings suggest that participants dynamically updated expectations of their performance efficacy based on feedback, and that the P3b played a role in prediction error-based updating of these expectations. Next, we tested the prediction that learned efficacy estimates determine the expected benefits of control, and thus influence how much control is allocated (Shenhav et al., 2013). We have previously shown that people exert more control when expecting their performance to be more rather than less efficacious on the upcoming trial (Frömer et al., 2021a). This was reflected in better behavioral performance and higher

amplitudes of the contingent negative variation (CNV; Figure 3B) - a slow fronto-central

waveform preceding target onset, which scales with the amount of control allocated in

preparation for the task (Grent-'t-Jong & Woldorff, 2007; Schevernels et al., 2014). Here we

used the same marker of control, but, unlike in our previous study, efficacy expectations were (a)

learned rather than explicitly instructed; (b) varied over time rather than having a fixed

association with a cue; and (c) varied continuously across the range of efficacy probabilities

rather than alternating between maximal and minimal efficacy. We were therefore interested in

testing whether these dynamically varying expectations of efficacy, as estimated by our model,

would still exert the same influence on behavior and neural activity.

Consistent with our predictions and our previous findings, participants tended to perform

better when they expected performance to be more efficacious, responding faster on correct trials

with increasing model-based estimates of efficacy (Figure 4A; Table S4). This finding replicates

the performance effects we observed using instructed cues, albeit with only modest evidence (b

$= -16.00$; 95% CrI [-34.91, 2.96]; $p_{b>0} = 0.05$; $BF_{01} = 1.68$). Like in our previous studies, faster

responding was not explained by a change in speed-accuracy trade-off, as accuracy did not

decrease with increasing efficacy, but if anything, slightly increased ($b = 0.12$; 95% CrI [-0.20,

0.44]; $p_{b>0} = 0.23$; $BF_{01} = 1.99$). These analyses controlled for the standard behavioral effects

related to Stroop congruency (i.e., slower and less accurate responding for incongruent relative to

congruent trials; Figure S4), as well as for the reward rate estimates.

If the CNV provides an index of control allocation based on current incentive

expectations, it should both reflect one's latest efficacy estimate and should predict performance

on the upcoming trial. Our findings support both predictions. Regressing single-trial CNV

amplitude onto model-based efficacy estimates, and controlling for expectations of reward

(discussed below), we found that these CNV amplitudes tracked current efficacy expectations (b = -0.35; 95% CrI [-0.85, 0.16]; $p_{b > 0}$ = 0.09; BF_{01} = 2.73; Figure 4C; Table S5). However, this effect was weaker than in the previous experiment with cued efficacy levels, which is to be expected given that in this experiment participants had to learn their efficacy levels. As with the behavioral finding above, this result provides evidence consistent with our previous CNV finding using instructed cues. We further replicate our earlier finding (Frömer et al., 2021a) that larger CNV amplitude in turn predicted better performance on the upcoming trial, with participants responding faster (b = 11.41; 95% CrI [8.09, 14.72]; $p_{b < 0}$ = 0; BF_{10} > 100; Figure 4D; Table S6) and more accurately (b = -0.07; 95% CrI [-0.12, -0.01]; $p_{b > 0}$ = 0.01; BF_{10} = 3.14; Figure 4E; Table S6) as CNV increased.

Figure 4. Efficacy estimates influence allocation of control and behavior in Study 1. A. Higher efficacy predicts faster accurate responses. **B. CNV increases with higher efficacy.** Left: Grand-average ERP for high and low efficacy estimates (median split used for plotting). The shaded area marks the time window used for CNV quantification. Time 0 corresponds to the onset of the neutral cue. Right: Predicted CNV amplitudes as a function of efficacy estimates. The topography shows the fixed effect of the efficacy estimate from the fitted linear model. **C. -D. Larger CNV amplitude predicts better performance.** Predicted accurate RT (**C.**) and accuracy (**D.**) as a function of efficacy estimates. Shaded error bars indicate 95% confidence intervals.

Parallel learning of efficacy and reward rate

We held the amount of reward at stake constant over the course of the experiment, but the frequency of reward (reward rate) varied over the course of the session based on a participant's recent performance, and participants were sporadically probed for estimates of their reward rate (interleaved with trials that were followed by efficacy probes). Our efficacy manipulation explicitly controlled for this variability by yoking Random-Outcome feedback to a participant's recent reward rate (see methods for details). However, this additional source of variability also provided an opportunity to examine additional mechanisms of learning and adaptation in our task. As in the case of efficacy estimates, reward rate estimates were robustly predicted by reward feedback on previous trials (Table S1; Figure S1A), and this reward rate learning process was well captured by a two learning rate model (cf. Garrett & Daw, 2020; Figure S1B-D). Updates to these reward rate estimates were reflected in P3b modulations of (unsigned) reward prediction errors and associated learning rates (Fischer & Ullsperger, 2013; see Figure S3, Table S3). This pattern of results provides additional evidence that efficacy learning involves similar neural and computational mechanisms as reward-based learning.

Learned efficacy modulates control over information processing

Our findings suggest that people rely on domain-general mechanisms to learn about their performance efficacy in a given environment, and use these learned estimates of efficacy to optimize performance on their task. Specifically, in Study 1 we found that higher levels of learned efficacy are associated with faster responses, albeit with modest evidence (b = -16.00; 95% CrI [-34.91, 2.96]; $p_{b > 0}$ = 0.05; BF_{01} = 1.68). We also found that this speeding occurred on correct but not incorrect trials, suggesting that these performance adjustments reflected attentional control rather than adjustments to speed-accuracy tradeoffs. However, these findings

remain only suggestive in the absence of a formal model, and the presence of a stringent

response deadline in this study (individually calibrated for each subject during the practice phase

to ensure the reward rate of 80%; M = 796ms; SD = 73ms) presented an obstacle to fitting our

behavioral data to such a model without additional assumptions (e.g., regarding the form of a

collapsing threshold).

To provide further support for our proposal that learned efficacy influences control over

information processing, we ran an additional behavioral study (Study 2). Participants in this

study (N = 51) performed a web-based version of the task in Study 1, with the biggest

modification being that the Stroop trials (now using picture-word rather than color-word

interference) were performed over the course of short self-paced time intervals rather than trial

by trial as in Study 1. Specifically, participants were given limited time windows (2-4s) to

complete as many Stroop trials as they wanted to and were rewarded at the end of each interval

(cf. Leng et al., 2021). When rewards were performance-based, participants received a number of

points exactly proportional to the number of correct responses they gave during that window.

When rewards were random, the number of points was unrelated to performance on that interval

but (as in Study 1) was yoked to their performance in previous performance-contingent intervals.

Following our approach in Study 1, we varied the likelihood of a given interval being

performance-based or random over the course of the session (Figure S5A), and sporadically

probed participants for their subjective estimates of expected efficacy and reward rate. While in

most respects very similar to the paradigm in Study 1, one noteworthy feature of this self-paced

design is that it resulted in a much less stringent deadline for responding, thus producing reaction

time patterns more typical of free-response paradigms for which the traditional (fixed-threshold)

DDM was designed. Note that because this was an online sample we also employed additional

cutoff criteria to exclude inattentive participants, as detailed in the Methods section and figure

S5B-C.

Replicating the learning patterns observed in Study 1, subjective estimates of efficacy in

Study 2 reflected a running average over recent efficacy feedback (b = 0.18; 95% CrI [0.16,

0.20]; $p_{b < 0}$ = 0; BF_{10} > 100). This effect was again weighted towards the most recent feedback

but still present up to five intervals back (all $p_{b < 0}$ < 0.01; Figure 5A, Figure S6, and Table S7).

As in Study 1, this learning pattern was best captured by an RL algorithm with two learning rates

(Figure 5B), though positive and negative efficacy learning rates did not significantly differ from

one another on average (b = 0.03; 95% CrI [-0.03, 0.08]; $p_{b < 0}$ = 0.260; BF_{01}=11.74; Figure 5C).

Critically, we once again found that higher model-based estimates of efficacy predicted

better performance on the upcoming interval. Participants responded faster (b = -10.25; 95% CrI

[-19.86, -0.20]; $p_{b > 0}$ = 0.02; BF_{01} = 2.33) and more accurately (b = 0.23; 95% CrI [0.03, 0.43];

$p_{b > 0}$ = 0.01; BF_{01} > 100) with increasing model-based efficacy estimates (Figure 5D, Figure

S4C-D, and Table S8). To formally test whether these behavioral patterns reflected adjustments

in information processing (i.e., the rate of evidence accumulation once the stimuli appeared) or

instead reflected adjustments in speed-accuracy tradeoffs (i.e., the threshold for responding), we

fit these data with the hierarchical drift diffusion model (HDDM; Wiecki, Sofer, & Frank, 2013)

to test whether model-based efficacy estimates predicted trial-by-changes in drift rate, threshold,

or both, while controlling for influences of expected reward rate on those same DDM

parameters. We found that higher levels of expected efficacy were associated with higher drift

rates (b = 0.07; 95% CrI [0.14, 0.43]; $p_{b < 0}$ = 0.00) but were uncorrelated with threshold levels (b

= -0.00; 95% CrI [-0.04, 0.04]; $p_{b < 0}$ = 0.74). Expected reward rate was not correlated with either

drift rate or threshold (Table S9). These results suggest that participants responded to changes in

performance efficacy by adjusting their attention to the task, rather than simply adjusting their

response threshold (i.e., becoming more or less impulsive).

Figure 5. Efficacy is learned in the same way in a modified task and influences a behavioral marker of control allocation. A. Regression weights for the influence of previous feedback type on the subjective efficacy estimate. Error bars represent 50% and 95% highest density intervals. **B. Two learning rate model captures efficacy learning best.** Differences in BICs between the two learning rate model, and the intercept-only and one learning rate models. **C. Efficacy learning rates.** Positive and negative efficacy learning rates for each participant. **D. Higher model-based efficacy estimates predict better behavioral performance.** Higher efficacy estimates reduce reaction times (left) and improve accuracy (right). **E. Higher model-based efficacy estimates predict increased allocation of attention.** Parameter estimates from the drift diffusion model. Higher efficacy estimates increase drift rates, but not response caution (thresholds).

Discussion

To evaluate the expected benefits of investing cognitive control into a task, people need to consider whether this control is necessary and/or sufficient for achieving their desired outcome (i.e., whether these efforts are worthwhile). A critical determinant of the worthwhileness of control is performance efficacy, the extent to which performance on a control-demanding task matters for outcome attainment versus those outcomes being determined by performance-unrelated factors. However, the mechanisms through which people estimate the efficacy of their performance based on previous experience are largely unknown. Here, we identified the computational and neural mechanism through which efficacy is learned and used to guide the allocation of cognitive control. Across two experiments, we found that participants dynamically updated expectations of the efficacy of their task performance (i.e., the likelihood that this performance will determine reward attainment), and used those expectations to adjust how much control they allocated. The feedback-based updating of efficacy was well-captured by a prediction error-based learning algorithm. Model-based estimates of efficacy and efficacy prediction errors were encoded by canonical neural signatures of effort allocation and belief updating, respectively. Importantly, these findings cannot be explained by variability in reward, as reward rate was held constant across efficacy levels, and the subjective reward rate was controlled for statistically. Further, our model-based analysis revealed that people allocated more control when they learned that they had more efficacy, extending our previous findings on instructed efficacy (Frömer et al., 2021a). Taken together, our results uncover the mechanism through which efficacy estimates are learned and used to inform mental effort investment within a given task environment.

Previous research has characterized the learning algorithms people use to learn the reward value of different states and actions in their environment (Gläscher et al., 2010; Sutton & Barto, 2018). Recent theoretical (Jiang, Heller, & Egner, 2014; Lieder et al., 2018; Verbeke & Verguts, 2019) and empirical (Bejjani et al., 2018; Bustamante et al., 2021; Jiang et al., 2020; Otto & Daw, 2019) work has extended this research to show how similar algorithms guide learning and adaptation of cognitive control under varying rewards and task demands within a given task environment. Our findings extend this work further in several ways. First, we show that people leverage weighted prediction errors when learning about the efficacy of task performance, independently of potential reward and task difficulty. Second, we show that they update their efficacy expectations differently depending on whether performance was more efficacious or less efficacious than they expected, demonstrating a striking parallel with dual learning rate models that have been found to prevail in research on reward learning (Collins & Frank, 2013; Garrett & Daw, 2020; Lefebvre et al., 2017), including in our own reward rate data. Third, we show that participants dynamically adjust their control allocation based on learned efficacy, just as they do for learned rewards and task demands (Bugg et al., 2011; Jiang et al., 2014; Lieder et al., 2018).

Our neural findings build further on past research on learning and control adaptation. The P3b component has been shown to track prediction-error based learning from action-relevant outcome values (Fischer & Ullsperger, 2013; Lohse, Miller, Daou, Valerius, & Jones, 2020; Nassar et al., 2019; Frömer et al., 2021b). Here we show that this neural marker tracks learning about efficacy in the same way as it tracks learning about rewards. We found increased P3b amplitudes when people experienced feedback about outcome contingencies that was less expected given their current estimate of efficacy (e.g., expecting low efficacy, but getting performance-contingent feedback), relative to when these contingencies were more expected

(e.g., expecting low efficacy and getting random feedback). Our additional finding that P3b amplitude was overall larger for efficacy compared to no efficacy feedback demonstrates that our participants were not just tracking the frequency of the two types of feedback, as predicted by an oddball account. Instead this finding suggests that they were actively learning from the feedback.

Extending previous findings on cueing efficacy and/or reward (Frömer et al., 2021a; Schevernels et al., 2014), our CNV and behavioral results further show that participants used these learned efficacy estimates to calibrate their effort and their performance. Notably, unlike in previous work, our study shows effort-related changes in CNV amplitude entirely divorced from perceptual cues, providing evidence that this activity truly reflects adjustments in control, rather than reactive processing of features associated with the cue. Taken together, our findings suggest that similar neural mechanisms underlie learning and control adaptation in response to variability in one's efficacy in a task environment, as they do in response to variability in expected rewards (Leng et al., 2021; Otto & Daw, 2019). By fitting behavioral data from this task to a drift diffusion model (Study 2), we were able to further demonstrate that participants were adapting to changes in expected efficacy by enhancing the processing of stimuli (i.e., increasing their rate of evidence accumulation) - potentially via attentional control mechanisms – rather than by adjusting their threshold for responding. This particular pattern of control adjustments was predicted for the current task because performance-contingent rewards depended on responses being *both* fast and accurate (as in Frömer et al., 2021a), but future work should test the prediction that different control adjustments should emerge under different performance contingencies (cf. Leng et al., 2021; Ritz et al., 2021).

Our findings build on previous research on how people learn about controllability of their environment. Studies have examined the neural and computational mechanisms by which

humans and other animals learn about the contingencies between an action and its associated outcome, and demonstrated that these learned action-outcomes contingencies influence which actions are selected and how vigorously they are enacted (Dickinson & Balleine, 1995; Liljeholm, Tricomi, O'Doherty, & Balleine, 2011; Ly, Wang, Bhanji, & Delgado, 2019; Manohar, Finzi, Drew, & Husain, 2017; Moscarello & Hartley, 2017; Dorfman et al., 2019; Dorfman et al., 2021; Ligneul, 2021). Our work extends this research into the domain of cognitive control, where the contingencies between actions (i.e., control adjustments) and outcomes (e.g., reward) depend both on whether control adjustments predicts better performance and whether better performance predicts better outcomes (Shenhav et al., 2021). Learning control-outcome contingencies therefore requires learning about how control states map onto performance (control efficacy) as well as how performance maps onto outcomes (performance efficacy). By describing the mechanisms by which people solve the latter part of this learning problem, and demonstrating that these are comparable to those engaged during action-outcome learning, our study lays critical groundwork for better understanding the links between selection of actions and control states.

Our efficacy-updating results are a reminder that many aspects of feedback are reflected in prediction error signals (Frömer et al., 2020; Langdon et al., 2018). In the present study, we intentionally separated feedback about reward and efficacy to isolate the cognitive and neural learning mechanisms associated with each. In doing so, we have taken an important first step towards understanding the updating mechanisms underlying each. Further work is needed to understand how they are inferred in more naturalistic settings, in which different forms of feedback are often multiplexed, containing information about the values of actions that were

taken as well as about the features and structure of the environment (cf. Dorfman et al., 2019;

2021).

Another distinct element of more complex naturalistic environments is that the same

feedback can be used to evaluate multiple targets, internal ones, such as the selected response

and its predicted outcome, or external ones, such as the source of feedback/environment

(Carlebach & Yeung, 2020). Such multi-level prediction error signals might for instance explain,

why despite close links between P3b and behavioral adaptation (Chase, Swainson, Durham,

Benham, & Cools, 2011; Fischer & Ullsperger, 2013; Yeung & Sanfey, 2004), this link is

context dependent and attempts to link the P3b uniquely to behavioral updating have failed

(Nassar et al., 2019). Reinforcement learning, and predictive inference more generally, have

been proposed to not only support the selection of individual actions, but also extended

sequences of actions and control signals (Holroyd & Yeung, 2012; Lieder et al., 2018).

Alongside evaluations of actions and environmental states, neural signatures of feedback-based

learning could thus further reflect evaluations of control signals, their quality or intensity. Given

the many potential causes a given outcome can have, and the flexibility that people have in how

they use the feedback, it is easy to see how feedback could be misattributed and lead to

inaccurate beliefs about performance efficacy. Such beliefs about environmental statistics can

drive changes in feedback-processing and behavioral adaptation, above and beyond the true

statistics (Schiffer, Siletti, Waszak, & Yeung, 2017), and are thus of particular importance for

understanding some of the cognitive symptoms of mental disorders.

Understanding how efficacy estimates develop based on previous experiences is crucial for

understanding why people differ in how much cognitive control they allocate in different

situations (Shenhav et al., 2021). People differ in their beliefs about how much control they have

over their environments (Leotti, Iyengar, & Ochsner, 2010; Moscarello & Hartley, 2017), and in their ability to estimate their efficacy (Cohen, Nussenbaum, Dorfman, Gershman, & Hartley, 2020). Further, many mental disorders, including depression and schizophrenia, have been linked with one's estimates of their ability to control potential outcomes in their environment, including through allocation of control (Huys & Dayan, 2009; Maier & Seligman, 2016), and we have recently proposed that such changes can drive impairments of motivation and control in those populations (Grahek et al., 2019). As we show in this study, when people have learned to expect low efficacy, they will allocate less cognitive control, which can manifest as apparent control deficits. The experimental and modeling approach taken in our study helps uncover a more fine-grained view of how components of motivation are learned and used to support the allocation of cognitive resources. In this way, our study takes a first step toward a better computational and neural account of efficacy learning, which can aid the understanding of individual differences in the willingness to exert mental effort, as well as the development of interventions aimed at teaching individuals when these efforts truly matter.

Supplemental Materials

Figure S1. Reward rate learning – model-based analyses for Study 1. A. Regression weights for the influence of the current and previous feedbacks on the subjective estimates of reward rate (left) and efficacy (right). Reward rate estimates are strongly predicted by the previous reward feedbacks, but not efficacy feedbacks. The reverse is true for the subjective efficacy estimates. Error bars represent 50% and 95% highest density intervals. **B.** Model comparison between the fitted learning models for the reward rate learning model. **C.** Positive and negative learning rate estimates for all subjects for the reward rate learning model. Negative learning rates were numerically, but not statistically larger than positive ones ($b = 0.01$; 95% CrI [-0.07, 0.09]; $p_{b < 0} = 0.38$; $BF_{10} = 0.09$)

Table S1

Study 1 regression weights for the models predicting subjective efficacy and reward rate estimates based on efficacy and reward feedback 5 trials back.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)	BF$_{10}$
Subjective efficacy				
Intercept	0.52	0.49, 0.55	0.00	
T1 Performance-Based - Random	0.14	0.12, 0.16	0.00	>100
T2 Performance-Based - Random	0.12	0.10, 0.14	0.00	>100
T3 Performance-Based - Random	0.08	0.05, 0.10	0.00	>100
T4 Performance-Based - Random	0.03	0.01, 0.05	0.00	2.20
T5 Performance-Based - Random	0.03	0.01, 0.05	0.00	1.72
T1 Reward - No Reward	0.02	0.00, 0.04	0.02	0.37
T2 Reward - No Reward	0.03	0.01, 0.05	0.00	4.23
T3 Reward - No Reward	0.03	0.01, 0.05	0.01	0.95
T4 Reward - No Reward	0.02	0.00, 0.04	0.02	0.39
T5 Reward - No Reward	0.00	-0.02, 0.02	0.46	0.05
Subjective reward rate				
Intercept	0.50	0.46, 0.54	0.00	
T1 Reward - No Reward	0.19	0.17, 0.21	0.00	>100
T2 Reward - No Reward	0.16	0.14, 0.18	0.00	>100
T3 Reward - No Reward	0.09	0.07, 0.11	0.00	>100
T4 Reward - No Reward	0.05	0.03, 0.06	0.00	30.86
T5 Reward - No Reward	0.05	0.03, 0.07	0.00	>100
T1 Performance-Based - Random	-0.01	-0.03, 0.01	0.73	0.06
T2 Performance-Based - Random	0.00	-0.01, 0.02	0.33	0.05
T3 Performance-Based - Random	0.01	-0.01, 0.03	0.00	0.05
T4 Performance-Based - Random	0.02	-0.01, 0.03	0.20	0.08
T5 Performance-Based - Random	0.02	0.00, 0.03	0.04	0.20

Figure S2. Efficacy learning in Study 1 – the effect of different feedback probability drifts on learning. A. The probability drift of contingent feedback presented to the first half of subjects. **B.** Learning rates for contingent and random feedback for the first half of subjects. Learning rates for the contingent and random feedback did not differ ($b = 0.01$; 95% CrI [-0.06, 0.08]; $p_{b<0} = 0.38$; $BF_{10} = 0.09$). **C.** The inversed probability drift of contingent feedback presented to the second half of participants. **D.** Learning rates for contingent and random feedback did not differ ($b = 0.02$; 95% CrI [-0.06, 0.11]; $p_{b<0} = 0.31$; $BF_{10} = 0.10$).

Table S2

Study 1 regression weights for the models predicting the P3b to efficacy feedback.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)	BF$_{10}$
Efficacy feedback				
Intercept	1.38	0.67, 2.09	0.00	
Model-based efficacy	-0.29	-1.56, 0.97	0.68	0.23
Efficacy feedback type (PB-R)	0.86	0.42, 1.31	0.00	30.86
Model-based reward rate	-0.19	-1.50, 1.18	0.62	0.23
Model-based efficacy × Efficacy feedback type	-2.40	-4.07, -0.74	1.00	24.40
Model-based efficacy slope for Performance-Based feedback	-1.49	-2.72, -0.27	0.97	1.52
Model-based efficacy slope for Random feedback	0.91	-0.34, 2.21	0.12	0.44
Predictions errors and learning rates				
Intercept	1.31	0.51, 2.09	0.00	
Unsigned prediction error	1.25	0.35, 2.15	0.01	5.84
Learning rate	2.00	-2.21, 6.04	0.17	1.08
Model-based reward rate	-0.09	-1.48, 1.34	0.55	0.23
Unsigned prediction error ×Learning rate	0.57	-4.87, 6.14	0.41	0.91

ERP analyses of the reward rate feedback in Study 1

To investigate the effect of the model-based reward rate estimates on the processing of

reward, we performed complementary analyses on the P3b related to the processing of the

reward feedback (Figure S3A). We expected that a similar learning mechanism should operate

on both reward and efficacy feedback, but that the neural markers of feedback processing should

be sensitive to different model-based estimates. Thus, we expected the reward-locked P3b to be

sensitive to reward-rate estimates, but not to efficacy estimates.

Our results showed larger P3b amplitudes to no reward compared to reward feedback

(note that reward rate was approximately .80 and negative feedback less expected overall), and

the processing of the reward feedback was influenced by reward-rate estimates, although to

lesser extent than for the P3b (Figure S3B; Table S3). Probing the effects of learning directly, we

found robust effects of the prediction errors and learning rates (Figure S3C). The P3b to reward

feedback was not influenced by the model-based efficacy estimates in either of these analyses.

Figure S3. Learning reward rate estimate from reward feedback in Study 1. A. ERP average for the P3b locked to the onset of reward feedback. **B.** Interaction between reward feedback and the model-based reward rate estimate. **C.** The effects of unsigned prediction errors (left) and learning rates (right) on the P3b.

Table S3

Study 1 regression weights for the models predicting the P3b to reward feedback.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)	BF$_{10}$
Efficacy feedback				
Intercept	0.93	0.08, 1.74	0.02	
Model-based efficacy	0.45	-0.45, 1.32	0.16	0.15
Reward feedback type (R-NR)	-0.35	-0.90, 0.18	0.90	0.12
Model-based reward rate	-0.15	-1.55, 1.23	0.59	0.14
Model-based reward ×Reward feedback type	-1.26	-3.73, 1.19	0.85	0.41
Model-based efficacy slope for No Reward feedback	0.48	-1.34, 2.34	0.87	0.23
Model-based efficacy slope for Reward feedback	-0.78	-1.96, 0.39	0.32	0.21
Predictions errors and learning rates				
Intercept	0.89	0.15, 1.64	0.01	
Unsigned prediction error	1.09	0.38, 1.81	0.00	4.82
Learning rate	6.68	-0.07, 12.99	0.03	4.72
Model-based efficacy	0.39	-0.47, 1.24	0.19	0.13
Unsigned prediction error ×Learning rate	0.94	-8.06, 9.71	0.42	0.92

Figure S4. Behavioral effects of congruency in Study 1 and Study 2. A. In comparison to the neutral trials in Study 1, participants were faster to respond to congruent (b = 9.95; 95% CrI [3.33, 16.59]; $p_{b<0}$ = 0.00; BF = 1.60) and slower to respond to incongruent trials (b = 61.24; 95% CrI [49.50, 72.59]; $p_{b<0}$ = 0.00; BF = 6.73). **B.** In comparison to the neutral trials in Study 1, participants were equally likely to be correct when responding to congruent (b = -0.11; 95% CrI [-0.29, 0.06]; $p_{b>0}$ = 0.091; BF = 0.01) and less likely to be correct when responding to incongruent trials (b = -0.48; 95% CrI [-0.64, -0.33]; $p_{b>0}$ = 0.00; BF > 8.22). The regression weights for the accuracy analysis are in log odds. **C.** In Study 2 participants were slower to respond to incongruent relative to congruent trials (b = 47.48; 95% CrI [40.69, 52.22]; $p_{b<0}$ = 0.001; BF = 50). **D.** In Study 2 participants were less accurate when responding to incongruent compared to congruent trials (b = -0.48; 95% CrI [-0.57, -0.39]; $p_{b<0}$ = 1; BF > 100).

Table S4

Study 1 regression weights for the models predicting reaction times and accuracy based on efficacy and reward rate estimates.

Parameter	Estimate	95% Credible interval	Posterior probability $(p<0)$	BF_{01}
Reaction times				
Intercept	655.37	635.09, 675.51	0.00	
Facilitation	9.95	3.33, 16.59	0.00	1.60
Interference	61.24	49.50, 72.59	0.00	6.73
Model-based efficacy	-16.00	-34.91, 2.96	0.95	1.68
Model-based reward rate	-1.56	-27.66, 24.04	0.54	1.36
Accuracy				
Intercept	1.56	1.33, 1.80	0.00	
Facilitation	-0.11	-0.29, 0.06	0.91	0.01
Interference	-0.48	-0.64, -0.33	1.00	8.22
Model-based efficacy	0.12	-0.20, 0.44	0.23	1.99
Model-based reward rate	-0.36	-0.73, 0.03	0.97	0.29

Table S5

Study 1 regression weights for the models predicting the CNV based on efficacy and reward rate.

Parameter	Estimate	95% Credible interval	Posterior probability $(p<0)$	BF_{01}
CNV				
Intercept	-0.37	-0.58, -0.16	0.00	
Model-based efficacy	-0.35	-0.85, 0.16	0.09	2.73
Model-based reward rate	-0.09	-0.72, 0.55	0.38	2.19

Table S6

Study 1 regression weights for the models predicting reaction times and accuracy based on the CNV.

Parameter	Estimate	95% Credible interval	Posterior probability $(p<0)$	BF_{10}
Reaction times				
Intercept	655.28	635.44, 675.91	0.00	
CNV amplitude	11.41	8.09, 14.72	0.00	>100
Accuracy				
Intercept	1.33	1.11, 1.53	0.00	
CNV amplitude	-0.07	-0.12, -0.01	0.99	3.14

Figure S5. Efficacy drifts and the unsupervised clustering of the efficacy learning rates in Study 2. A. Efficacy drifts used in Study 2. The probability drifts of contingent feedback presented to the first (top) and second (bottom) half of the participants. **B. The results of the winning Gaussian mixture model for clustering of the efficacy learning rates.** One of the clusters (blue dots) included only the subjects with extremely small learning rates (all learning rates < 0.03; N=20; group 1 – blue dots). These subjects were excluded from the further analyses with the assumption that they did not pay attention to the feedbacks or that they were giving random answers to the probes. **C. Histograms of the positive and negative learning rates and their clusters.** For both then negative (left) and positive (right) learning rates cluster 1 included only the participants with very low learning rates.

Table S7

Study 2 regression weights for the models predicting subjective efficacy and reward rate estimates based on efficacy and reward feedback 5 trials back.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)	BF$_{10}$
Subjective efficacy				
Intercept	0.43	0.37, 0.49	0.00	
T1 Performance-Based - Random	0.17	0.16, 0.19	0.00	>100
T2 Performance-Based - Random	0.15	0.13, 0.18	0.00	>100
T3 Performance-Based - Random	0.10	0.08, 0.12	0.00	>100
T4 Performance-Based - Random	0.06	0.04, 0.08	0.00	>100
T5 Performance-Based - Random	0.04	0.02, 0.06	0.00	>100
T1 Reward - No Reward	0.06	0.02, 0.10	0.00	5.42
T2 Reward - No Reward	0.06	0.01, 0.11	0.01	2.23
T3 Reward - No Reward	0.03	-0.02, 0.07	0.11	0.23
T4 Reward - No Reward	0.00	-0.05, 0.04	0.54	0.11
T5 Reward - No Reward	0.06	0.01, 0.10	0.01	1.69
Subjective reward rate				
Intercept	0.36	0.29, 0.43	0.00	
T1 Reward - No Reward	0.21	0.17, 0.25	0.00	>100
T2 Reward - No Reward	0.13	0.09, 0.17	0.00	>100
T3 Reward - No Reward	0.12	0.07, 0.16	0.00	>100
T4 Reward - No Reward	0.03	-0.01, 0.06	0.09	0.21
T5 Reward - No Reward	0.09	0.05, 0.13	0.00	>100
T1 Performance-Based - Random	0.05	0.03, 0.07	0.00	0.06
T2 Performance-Based - Random	0.05	0.03, 0.06	0.00	0.05
T3 Performance-Based - Random	0.00	-0.02, 0.02	0.51	0.05
T4 Performance-Based - Random	0.03	0.02, 0.05	0.00	0.08
T5 Performance-Based - Random	0.02	0.00, 0.03	0.03	0.20

Figure S6. Efficacy and reward learning – model-based analyses for Study 2. A. Regression weights for the influence of the current and previous feedbacks on the subjective estimates of reward rate (left) and efficacy (right). Reward rate estimates are strongly influenced by the previous reward feedbacks, and the reverse is true for the subjective efficacy estimates. Error bars represent 50% and 95% highest density intervals. **B.** Model comparison between the fitted learning models for the reward learning model. **C.** Positive and negative learning rate estimates for all subjects for the reward learning model. Positive learning rates were numerically, but not statistically larger than negative ones ($b = 0.02$; 95% CrI [-0.01, 0.05]; $p_{b<0} = 0.11$; $BF_{10} = 0.06$)

Table S8

Study 2 regression weights for the models predicting reaction times and accuracy based on efficacy and reward rate estimates. Bayes Factors calculated only for the parameters with informative priors. Posteriors from Study 1 were used as priors for the effects of congruency, efficacy, and reward.

Parameter	Estimate	95% Credible interval	Posterior probability $(p<0)$	BF_{01}
Reaction times				
Intercept	643.02	625.42, 661.81	0.00	
Incongruent-Congruent	47.48	40.69, 55.22	0.00	0.02
Model-based efficacy	-10.25	-19.86, -0.20	0.98	2.33
Model-based reward	-0.17	-21.15, 20.34	0.51	1.24
Interval length	-1.76	-2.73, -0.83	1.00	
Interval congruency	-5.34	-6.69, -4.00	1.00	
Accuracy				
Intercept	2.35	2.16, 2.52	0.00	
Incongruent-Congruent	-0.48	-0.57, -0.39	1.00	>100
Model-based efficacy	0.23	0.03, 0.43	0.01	>100
Model-based reward	-0.29	-0.57, -0.01	0.98	1.07
Interval length	-0.13	-0.16, -0.10	1.00	
Interval congruency	0.04	0.00, 0.08	0.03	

Table S9

Drift Diffusion model estimates.

Parameter	Estimate	95% Credible interval	Posterior probability (p<0)
Drift rate			
Congruent-Incongruent	0.82	0.74, 0.92	0.00
Model-based efficacy	0.29	0.14, 0.40	0.00
Model-based reward	0.20	-0.39, 0.04	0.50
Threshold			
Model-based efficacy	-0.00	-0.04, 0.04	0.57
Model-based reward rate	-0.04	-0.14, 0.05	0.81

Acknowledgements

This work was supported by the Special Research Fund (BOF) of Ghent University [grant #01D02415] (I.G.) the Research Foundation Flanders (FWO) travel grant [grant #V432718N] (I.G.), a Center of Biomedical Research Excellence grant P20GM103645 from the National Institute of General Medical Sciences (A.S.), and the Alfred P. Sloan Foundation Research Fellowship in Neuroscience (A.S.). The funding sources were not involved in the study design; collection, analysis, and interpretation of data; writing of the report; and decision to submit the article for publication. We would like to thank Natalie Knowles and Hattie Xu for help with data collection for Study 1, Peyton Strong for help with data collection for Study 2, Carolyn Dean Wolf and Liz Cory for help with programming the Study 1 task, and Harrison Ritz for help with constructing the efficacy drifts.

Data availability

Raw and pre-processed data, materials, and analysis scripts will be available on Github and OSF upon publication.

References

Bejjani, C., Zhang, Z., & Egner, T. (2018). Control by association: Transfer of implicitly primed attentional states across linked stimuli. *Psychonomic Bulletin and Review*, *25*(2), 617–626. https://doi.org/10.3758/s13423-018-1445-6

Botvinick, M. M., & Cohen, J. D. (2014). The computational and neural basis of cognitive control: Charted territory and new frontiers. *Cognitive Science*, *38*(6), 1249–1285. https://doi.org/10.1111/cogs.12126

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

Bugg, J. M., Jacoby, L. L., & Chanani, S. (2011). Why it is too early to lose control in accounts of item-specific proportion congruency effects. *Journal of Experimental Psychology: Human Perception and Performance*, *37*(3), 844–859. https://doi.org/10.1037/a0019957

Bürkner, P.-C. (2016). brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical Software*, *80*(1), 1–28.

Bürkner, P.-C. (2017). Advanced Bayesian Multilevel Modeling with the R Package brms. *ArXiv:1705.11123*.

Bustamante, L., Lieder, F., Musslick, S., Shenhav, A., & Cohen, J. (2021). Learning to Overexert Cognitive Control in a Stroop Task. *Cognitive, Affective, & Behavioral Neuroscience*. https://doi.org/10.3758/s13415-020-00845-x

Carlebach, N., & Yeung, N. (2020). Flexible use of confidence to guide advice requests. *PsyArXiv*.

Carpenter, B., Gelman, A., Hoffman, M., Lee, D., Goodrich, B., Betancourt, M., … Riddell, A. (2016). Stan: A probabilistic programming language. *Journal of Statistical Software*, *2*(20), 1–37.

Chambon, V., Théro, H., Vidal, M., Vandendriessche, H., Haggard, P., & Palminteri, S. (2020).

Information about action outcomes differentially affects learning from self-determined

versus imposed choices. *Nature Human Behaviour*. https://doi.org/10.1038/s41562-020-

0919-5

Chase, H. W., Swainson, R., Durham, L., Benham, L., & Cools, R. (2011). Feedback-related

negativity codes prediction error but not behavioral adjustment during probabilistic reversal

learning. *Journal of Cognitive Neuroscience*, *23*(4), 936–946.

https://doi.org/10.1162/jocn.2010.21456

Chiu, Y. C., & Egner, T. (2019). Cortical and subcortical contributions to context-control

learning. *Neuroscience and Biobehavioral Reviews*, *99*(December 2018), 33–41.

https://doi.org/10.1016/j.neubiorev.2019.01.019

Cohen, A., Nussenbaum, K., Dorfman, H. M., Gershman, S. J., & Hartley, C. A. (2020). The

rational use of causal inference to guide reinforcement learning strengthens with age. *Npj

Science of Learning*.

Collins, Anne G E, & Frank, M. J. (2014). Opponent actor learning (OpAL): Modeling

interactive effects of striatal dopamine on reinforcement learning and choice incentive.

Psychological Review, *121*(3), 337–366. https://doi.org/10.1037/a0037015

Collins, Anne Gabrielle Eva, & Frank, M. J. (2013). Cognitive control over learning: creating,

clustering, and generalizing task-set structure. *Psychological Review*, *120*(1), 190–229.

https://doi.org/10.1037/a0030852

Daw, N. D., & Tobler, P. N. (2013). Value Learning through Reinforcement: The Basics of

Dopamine and Reinforcement Learning. In *Neuroeconomics: Decision Making and the

Brain: Second Edition* (pp. 283–298). https://doi.org/10.1016/B978-0-12-416008-8.00015-2

Delorme, A., & Makeig, S. (2004). EEGLAB: an open sorce toolbox for analysis of single-trail EEG dynamics including independent component anlaysis. *Journal of Neuroscience Methods*, *134*, 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Dickinson, A., & Balleine, B. (1995). Motivational Control of Instrumental Action. *Current Directions in Psychological Science*, *4*(5), 162–167. https://doi.org/10.1111/1467-8721.ep11512272

Dolan, R. J., & Dayan, P. (2013). Goals and habits in the brain. *Neuron*, *80*(2), 312–325. https://doi.org/10.1016/j.neuron.2013.09.007

Dorfman, H. M., Bhui, R., Hughes, B. L., & Gershman, S. J. (2019). Causal inference about good and bad outcomes. *Psychological science*, *30*(4), 516-525.

Dorfman, H. M., Tomov, M. S., Cheung, B., Clarke, D., Gershman, S. J., & Hughes, B. L. (2021). Causal inference gates corticostriatal learning. *Journal of Neuroscience*, *41*(32), 6892-6904.

Eldar, E., Hauser, T. U., Dayan, P., & Dolan, R. J. (2016). Striatal structure and function predict individual biases in learning to avoid pain. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(17), 4812–4817. https://doi.org/10.1073/pnas.1519829113

Fischer, A. G., & Ullsperger, M. (2013). Real and fictive outcomes are processed differently but converge on a common adaptive mechanism. *Neuron*, *79*(6), 1243–1255. https://doi.org/10.1016/j.neuron.2013.07.006

Frömer, R., Lin, H., Dean Wolf, C. K., Inzlicht, M., & Shenhav, A. (2021a). Expectations of reward and efficacy guide cognitive control allocation. *Nature Communications*, *12*(1). https://doi.org/10.1038/s41467-021-21315-z

Frömer, R., Maier, M., & Rahman, R. A. (2018). Group-level EEG-processing pipeline for flexible single trial-based analyses including linear mixed models. *Frontiers in Neuroscience*, *12*(48). https://doi.org/10.3389/fnins.2018.00048

Frömer, R., Nassar, M. R., Bruckner, R., Stürmer, B., Sommer, W., & Yeung, N. (2021b). Response-based outcome predictions and confidence regulate feedback processing and learning. Elife, 10, e62825.

Frömer, R., Stürmer, B., & Sommer, W. (2016). (Don't) Mind the effort: Effects of contextual interference on ERP indicators of motor preparation. *Psychophysiology*, *53*(10), 1577–1586. https://doi.org/10.1111/psyp.12703

Garrett, N., & Daw, N. D. (2020). Biased belief updating and suboptimal choice in foraging decisions. *Nature Communications*, *11*(1), 1–12. https://doi.org/10.1038/s41467-020-16964-5

Gelman, A., & Rubin, D. B. (1992). Inference from Iterative Simulation Using Multiple Sequences. *Statistical Science*, *7*(4), 457–472. https://doi.org/10.1214/ss/1177011136

Gläscher, J., Daw, N., Dayan, P., & O'Doherty, J. P. (2010). States versus rewards: Dissociable neural prediction error signals underlying model-based and model-free reinforcement learning. *Neuron*, *66*(4), 585–595. https://doi.org/10.1016/j.neuron.2010.04.016

Grahek, I., Shenhav, A., Musslick, S., Krebs, R. M., & Koster, E. H. W. (2019). Motivation and cognitive control in depression. *Neuroscience & Biobehavioral Reviews*, *102*(March), 371–381. https://doi.org/10.1016/j.neubiorev.2019.04.011

Grent-'t-Jong, T., & Woldorff, M. G. (2007). Timing and sequence of brain activity in top-down control of visual-spatial attention. *PLoS Biology*, *5*(1), 0114–0126. https://doi.org/10.1371/journal.pbio.0050012

Gureckis, T. M., Martin, J., McDonnell, J., Rich, A. S., Markant, D., Coenen, A., ... & Chan, P. (2016). psiTurk: An open-source framework for conducting replicable behavioral experiments online. *Behavior research methods*, *48*(3), 829-842.

Holroyd, C. B., & Yeung, N. (2012). Motivation of extended behaviors by anterior cingulate cortex. *Trends in Cognitive Sciences*, *16*(2), 122–128. https://doi.org/10.1016/j.tics.2011.12.008

Huys, Q. J. M., & Dayan, P. (2009). A Bayesian formulation of behavioral control. *Cognition*, *113*(3), 314–328. https://doi.org/10.1016/j.cognition.2009.01.008

Ille, N., Berg, P., & Scherg, M. (2002). Artifact Correction of the Ongoing EEG Using Spatial Filters Based on Artifact and Brain Signal Topographies. *Journal of Clinical Neurophysiology*, *19*(2), 113–124. https://doi.org/10.1097/00004691-200203000-00002

Jiang, J., Bramão, I., Khazenzon, A., Wang, S. F., Johansson, M., & Wagner, A. D. (2020). Temporal Dynamics of Memory-guided Cognitive Control and Generalization of Control via Overlapping Associative Memories. *Journal of Neuroscience*, *40*(10), 2343–2356. https://doi.org/10.1523/JNEUROSCI.1869-19.2020

Jiang, J., Heller, K., & Egner, T. (2014). Bayesian modeling of flexible cognitive control. *Neuroscience and Biobehavioral Reviews*, *46*(P1), 30–43. https://doi.org/10.1016/j.neubiorev.2014.06.001

Kleiner, M., Brainard, D., & Pelli, D. (2007). What's new in Psychtoolbox-3. In *Perception 36 ECVP Abstract Supplement*.

Kool, W., & Botvinick, M. (2018). Mental labour. *Nature Human Behaviour*. https://doi.org/10.1038/s41562-018-0401-9

Kruschke, J. K. (2013). Bayesian estimation supersedes the T test. *Journal of Experimental*

Psychology: General, *142*(2), 573–588. https://doi.org/10.1037/a0029177

Langdon, A. J., Sharpe, M. J., Schoenbaum, G., & Niv, Y. (2018). Model-based predictions for

dopamine. *Current Opinion in Neurobiology*, *49*, 1–7.

https://doi.org/10.1016/j.conb.2017.10.006

Lefebvre, G., Lebreton, M., Meyniel, F., Bourgeois-Gironde, S., & Palminteri, S. (2017).

Behavioural and neural characterization of optimistic reinforcement learning. *Nature*

Human Behaviour, *1*(4), 1–9. https://doi.org/10.1038/s41562-017-0067

Leng, X., Yee, D., Ritz, H., & Shenhav, A. (2021). Dissociable influences of reward and

punishment on adaptive cognitive control. *bioRxiv*, 2021.

Leotti, L. A., Iyengar, S. S., & Ochsner, K. N. (2010). Born to choose: The origins and value of

the need for control. *Trends in Cognitive Sciences*, *14*(10), 457–463.

https://doi.org/10.1016/j.tics.2010.08.001

Lieder, F., Shenhav, A., Musslick, S., & Griffiths, T. L. (2018). Rational metareasoning and the

plasticity of cognitive control. *Plos Computational Biology*, *14*(4).

https://doi.org/10.13140/RG.2.2.24500.14721

Ligneul, R. (2021). Prediction or Causation? Towards a Redefinition of Task Controllability.

Trends in Cognitive Sciences.

Liljeholm, M., Tricomi, E., O'Doherty, J. P., & Balleine, B. W. (2011). Neural Correlates of

Instrumental Contingency Learning: Differential Effects of Action-Reward Conjunction and

Disjunction. *Journal of Neuroscience*, *31*(7), 2474–2480.

https://doi.org/10.1523/JNEUROSCI.3354-10.2011

Lohse, K. R., Miller, M. W., Daou, M., Valerius, W., & Jones, M. (2020). Dissociating the

contributions of reward-prediction errors to trial-level adaptation and long-term learning.

Biological Psychology, *149*(January 2019).

https://doi.org/10.1016/j.biopsycho.2019.107775

Ly, V., Wang, K. S., Bhanji, J., & Delgado, M. R. (2019). A Reward-Based Framework of

Perceived Control. *Frontiers in Neuroscience*, *13*(February), 1–11.

https://doi.org/10.3389/fnins.2019.00065

Maier, S. F., & Seligman, M. E. P. (2016). Learned helplessness at fifty: Insights from

neuroscience. *Psychological Review*, *123*(4), 1–19. https://doi.org/10.1037/rev0000033

Manohar, S. G., Chong, T. T. J., Apps, M. A. J., Batla, A., Stamelou, M., Jarman, P. R., …

Husain, M. (2015). Reward Pays the Cost of Noise Reduction in Motor and Cognitive

Control. *Current Biology*, *25*(13), 1707–1716. https://doi.org/10.1016/j.cub.2015.05.038

Manohar, S. G., Finzi, R. D., Drew, D., & Husain, M. (2017). Distinct Motivational Effects of

Contingent and Noncontingent Rewards. *Psychological Science*, *28*(7), 1016–1026.

https://doi.org/10.1177/0956797617693326

Moscarello, J. M., & Hartley, C. A. (2017). Agency and the Calibration of Motivated Behavior.

Trends in Cognitive Sciences, *21*(10), 725–735. https://doi.org/10.1016/j.tics.2017.06.008

Nagase, A. M., Onoda, K., Foo, J. C., Haji, T., Akaishi, R., Yamaguchi, S., … Morita, K. (2018).

Neural mechanisms for adaptive learned avoidance of mental effort. *Journal of

Neuroscience*, *38*(10), 2631–2651. https://doi.org/10.1523/JNEUROSCI.1995-17.2018

Nalborczyk, L., & Bürkner, P.-C. (2019). An Introduction to Bayesian Multilevel Models Using

brms: A Case Study of Gender Effects on Vowel Variability in Standard Indonesian.

Journal of Speech, Language, and Hearing Research.

Nassar, M. R., Bruckner, R., & Frank, M. J. (2019). Statistical context dictates the relationship

between feedback-related EEG signals and learning. *ELife*, *8*, 1–26.

https://doi.org/10.7554/eLife.46975

Nigbur, R., Schneider, J., Sommer, W., Dimigen, O., & Stürmer, B. (2015). Ad-hoc and context-dependent adjustments of selective attention in conflict control: An ERP study with visual probes. *NeuroImage*, *107*, 76–84. https://doi.org/10.1016/j.neuroimage.2014.11.052

Niv, Y., Edlund, J. A., Dayan, P., & O'Doherty, J. P. (2012). Neural prediction errors reveal a risk-sensitive reinforcement-learning process in the human brain. *Journal of Neuroscience*, *32*(2), 551–562. https://doi.org/10.1523/JNEUROSCI.5498-10.2012

Niv, Y., Joel, D., & Dayan, P. (2006). A normative perspective on motivation. *Trends in Cognitive Sciences*, *10*(8), 375–381. https://doi.org/10.1016/j.tics.2006.06.010

Otto, A. R., & Daw, N. D. (2019). The opportunity cost of time modulates cognitive effort. *Neuropsychologia*, *123*(October 2017), 92–105. https://doi.org/10.1016/j.neuropsychologia.2018.05.006

Pelli, D. D. (1997). The VideoToolbox software for visual psychophysics: transforming numbers into movies. *Spatial Vision*, *10*(4), 437–442.

R Core Team. (2017). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing.

Ratcliff, R., & McKoon, G. (2008). The diffusion decision model: theory and data for two-choice decision tasks. *Neural computation*, *20*(4), 873-922.

Ritz, H., Leng, X., & Shenhav, A. (2021). Cognitive control as a multivariate optimization problem. *arXiv* preprint arXiv:2110.00668.

Rutledge, R. B., Skandali, N., Dayan, P., & Dolan, R. J. (2014). A computational and neural model of momentary subjective well-being. *Proceedings of the National Academy of Sciences*, *111*(33), 12252–12257. https://doi.org/10.1073/pnas.1407535111

Schevernels, H., Krebs, R. M., Santens, P., Woldorff, M. G., & Boehler, C. N. (2014). Task preparation processes related to reward prediction precede those related to task-difficulty expectation. *NeuroImage*, *84*, 639–647. https://doi.org/10.1016/j.neuroimage.2013.09.039

Scrucca, L., Fop, M., Murphy, T. B., & Raftery, A. E. (2016). mclust 5: clustering, classification and density estimation using Gaussian finite mixture models. *The R journal*, 8(1), 289.

Schiffer, A. M., Siletti, K., Waszak, F., & Yeung, N. (2017). Adaptive behaviour and feedback processing integrate experience and instruction in reinforcement learning. *NeuroImage*, *146*, 626–641. https://doi.org/10.1016/j.neuroimage.2016.08.057

Shenhav, A., Botvinick, M., & Cohen, J. (2013). The expected value of control: An integrative theory of anterior cingulate cortex function. *Neuron*, *79*(2), 217–240. https://doi.org/10.1016/j.neuron.2013.07.007

Shenhav, A., Musslick, S., Lieder, F., Kool, W., Griffiths, T. L., Cohen, J. D., & Botvinick, M. M. (2017). Toward a Rational and Mechanistic Account of Mental Effort. *Annual Review of Neuroscience*, *40*(1), 99–124. https://doi.org/10.1146/annurev-neuro-072116-031526

Shenhav, A., Prater Fahey, M., & Grahek, I. (2021). Decomposing the motivation to exert mental effort. *Current Directions in Psychological Science*.

Sutton, R. S., & Barto, A. G. (2018). *Reinforcement Learning: An Introduction*. MIT Press.

Verbeke, P., & Verguts, T. (2019). Learning to synchronize: How biological agents can couple neural task modules for dealing with the stability-plasticity dilemma. *PLoS Computational Biology*, *15*(8), 1–25. https://doi.org/10.1371/journal.pcbi.1006604

Verguts, T., Vassena, E., & Silvetti, M. (2015). Adaptive effort investment in cognitive and physical tasks: a neurocomputational model. *Frontiers in Behavioral Neuroscience*, *9*(March). https://doi.org/10.3389/fnbeh.2015.00057

Wagenmakers, E. J., Lodewyckx, T., Kuriyal, H., & Grasman, R. (2010). Bayesian hypothesis
testing for psychologists: A tutorial on the Savage-Dickey method. *Cognitive Psychology*,
60(3), 158–189. https://doi.org/10.1016/j.cogpsych.2009.12.001

Wiecki, T. V., Sofer, I., & Frank, M. J. (2013). HDDM: Hierarchical Bayesian estimation of the
drift-diffusion model in Python. *Frontiers in neuroinformatics*, 7, 14.

Yeung, N., & Sanfey, A. G. (2004). Independent coding of reward magnitude and valence in the
human brain. *Journal of Neuroscience*, *24*(28), 6258–6264.
https://doi.org/10.1523/JNEUROSCI.4537-03.2004