Metabolic dissimilarity determines the establishment of cross-feeding interactions in bacteria

Samir Giri[1,2,*], Leonardo Oña[2], Silvio Waschina[3,4], Shraddha Shitut[1,2,#], Ghada Yousif[1,2,5], Christoph Kaleta[4], and Christian Kost[1,2,*]

[1] Experimental Ecology and Evolution Research Group, Department of Bioorganic Chemistry, Max Planck Institute for Chemical Ecology, 07745 Jena, Germany
[2] Department of Ecology, School of Biology/Chemistry, University of Osnabrück, 49076 Osnabrück, Germany
[3] Institute for Human Nutrition and Food Science, Nutriinformatics, Christian-Albrechts-University Kiel, 24105 Kiel, Germany
[4] Research Group Medical Systems Biology, Institute for Experimental Medicine, Christian-Albrechts-University Kiel, 24105 Kiel, Germany
[5] Department of Botany and Microbiology, Faculty of Science, Beni-Suef University, Beni-Suef, Egypt

[*] Correspondence: christiankost@gmail.com (CKo), samirgiri2809@gmail.com (SG)

[#] Current address: Institute of Biology, University of Leiden, 2333 Leiden, The Netherlands.

Summary

The exchange of metabolites among different bacterial genotypes profoundly impacts the structure and function of microbial communities. However, the factors governing the establishment of these cross-feeding interactions remain poorly understood. While shared physiological features may facilitate interactions among more closely related individuals, a lower relatedness should reduce competition and thus increase the potential for synergistic interactions. Here we investigate how the relationship between a metabolite donor and recipient affects the propensity of strains to engage in unidirectional cross-feeding interactions. For this, we performed pairwise cocultivation experiments between four auxotrophic recipients and 25 species of potential amino acid donors. Auxotrophic recipients grew in the vast majority of pairs tested (78%), suggesting metabolic cross-feeding interactions are readily established. Strikingly, both the phylogenetic distance between donor and recipient and the dissimilarity of their metabolic networks were positively associated with the growth of auxotrophic recipients. Analysing the co-growth of species from a gut microbial community *in-silico* also revealed that recipient genotypes benefitted more from interacting with metabolically dissimilar partners, thus corroborating the empirical results. Together, our work identifies the metabolic dissimilarity between bacterial genotypes as key factor determining the establishment of metabolic cross-feeding interactions in microbial communities.

Keywords

Bacterial community, community assembly, cross-feeding, metabolic distance, metabolite secretion, niche overlap, phylogenetic relatedness, public good.

Highlights

- The exchange of essential metabolites is common in microbial communities

- Metabolic cross-feeding interactions readily establish between auxotrophic and prototrophic bacterial strains

- Both the phylogenetic and the metabolic dissimilarity between donors and recipients determines the successful establishment of metabolic cross-feeding interactions

Introduction

Microorganisms are ubiquitous on our planet and are key for driving pivotal ecosystem processes [1-3]. They contribute significantly to the flow of elements in global biogeochemical cycles [3, 4] and are also crucial for determining the fitness of plants [5, 6] and animals [7, 8], including humans [9, 10]. These vital functions are provided by complex communities that frequently consist of hundreds or even thousands of metabolically diverse strains and species [11, 12]. However, the rules that determine the assembly, function, and evolution of these microbial communities remain poorly understood. Understanding the underlying governing principles is central to microbial ecology and crucial for designing microbial consortia for biotechnological [13] or medical applications [14, 15].

In recent years, both empirical and theoretical work has increasingly suggested that the exchange of essential metabolites among different bacterial genotypes is a crucial process that can significantly affect growth [16, 17], composition [18], and the structure of microbial communities [19]. In these cases, one bacterial genotype releases a molecule into the extracellular environment, which can then be used by other cells in the local vicinity. The released substances frequently include building block metabolites such as amino acids [20, 21], vitamins [22, 23], or nucleotides [24], as well as degradation products of complex polymers [19, 25]. Even though these compounds represent valuable nutritional resources, they are released as unavoidable byproducts of bacterial physiology [26, 27] and metabolism [28] or due to leakage through the bacterial membrane [29, 30]. Consequently, the released compounds create a pool of resources that can benefit both conspecifics and members of other species in the local vicinity [31-34]. The beneficiaries include genotypes that opportunistically take advantage of these metabolites and strains, whose survival essentially depends on an external supply with the corresponding metabolite. Due to a mutation in their genome, these so-called *auxotrophic* genotypes are unable to autonomously synthesise vital nutrients such as amino acids, vitamins, or nucleotides. By utilising metabolites that are produced by another cell, a unidirectional cross-

feeding interaction is established. Auxotrophic mutants that use compounds released by others gain a significant fitness advantage over prototrophic cells that produce the required metabolites by themselves [35]. Due to the strong fitness benefits that can result for auxotrophic genotypes, this type of cross-feeding interaction is prevalent in all kinds of microbial ecosystems, including soil [36], fermented food [21], aquatic environments [37, 38], as well as host-associated microbiota [7, 39]. Despite the ubiquity of unidirectional cross-feeding interactions in nature, the factors determining their establishment remain poorly understood [40-43]. In particular, it is unclear how the relationship between the metabolite donor and the auxotrophic recipient affects the likelihood that a cross-feeding interaction is successfully established. Two possibilities are conceivable.

First, phylogenetically more closely related individuals are likely to share physiological features that favour the establishment of cross-feeding interactions relative to strains lacking these attributes. For example, an efficient transfer of metabolites from one cell to another commonly depends on close physical contact between donor and recipient [44]. The attachment to other cells is generally mediated by surface factors (e.g. adhesive proteins) or exopolymers that, in some cases, operate more effectively between cells sharing similar surface structures [45]. Moreover, some species form intercellular nanotubes to exchange metabolites between cells [44, 46], which in turn might require an increased structural similarity between cells for efficient transport to operate. Another context, in which the phylogenetic relatedness between donor and recipient could determine the establishment of a unidirectional cross-feeding interaction, is when both partners can communicate with each other. Certain signals that are involved in chemical communication (i.e. quorum sensing) between cells are more readily perceived by more closely related bacterial strains than more distantly related individuals [47]. Consequently, quorum sensing between more similar genotypes is also more likely to regulate processes such as cell-cell adhesion [16] or the establishment of metabolic cross-feeding interactions [48]. In the following, we refer to this possibility as the *similarity hypothesis.*

Second, unidirectional cross-feeding interactions might favour more distantly related donor-recipient pairs over interactions among close relatives. Two closely related bacterial cells are more likely to share ecological preferences such as habitat or resources utilised than two phylogenetically different bacterial taxa [41, 49, 50]. Moreover, two phylogenetically, more closely related cells will tend to have a more similar metabolic network than two distantly related cells [33, 51, 52]. Consequently, both the biosynthetic cost to produce a given metabolite and its nutritional value are more likely to differ in heterospecific pairs than amongst members of the same species [53, 54]. If these differences also translate into an enhanced growth of the auxotrophic recipient, a positive correlation between the growth of the auxotroph and the phylogenetic and/ or metabolic distance to the donor cell would be observed. In the following, we refer to this alternative possibility as the *dissimilarity hypothesis.*

Here we aim to distinguish between these two hypotheses to better understand the factors governing the establishment of this ecologically very important interaction.

To achieve this goal, we use unidirectional cross-feeding interactions as a model.
Synthetically assembling pairs consisting of an auxotrophic recipient and a prototrophic
amino acid donor of the same or a different species ensured that both interaction
partners do not share a coevolutionary history. In this way, all results will represent the
situation of a naïve encounter between both interaction partners and only mirror effects
resulting from the phylogenetic relatedness and metabolic dissimilarity between
partners. Using this synthetic ecological approach, we systematically determined
whether and how the phylogenetic or metabolic distance between auxotrophic
recipients and prototrophic amino acid donors affects cross-feeding in pairwise
bacterial consortia.

Our results show that in the vast majority of cases tested, unidirectional cross-
feeding interactions successfully established between a prototrophic donor and an
auxotrophic recipient. Strikingly, recipients' growth was positively associated with both
the phylogenetic and metabolic distance between donor and recipient. This pattern
could partly be explained by the difference in the amino acid profiles produced by
donors. Finally, analysing the co-growth of species from a gut microbial community *in-
silico* revealed that recipient genotypes benefitted more from interacting with
metabolically dissimilar partners, thus corroborating the empirical results. Together,
our work identifies the metabolic dissimilarity between donor and recipient genotypes
as a critical parameter determining the establishment of unidirectional cross-feeding
interactions in microbial communities.

148
149

Results
Auxotrophic recipients commonly benefit from the presence of prototrophic
donor cells
To determine the probability with which unidirectional cross-feeding interactions
emerge between an auxotrophic recipient and a prototrophic donor genotype, pairwise
coculture experiments were performed (Fig. 1A). For this, 25 strains that belonged to
21 different bacterial species were used as potential amino acid donors (Figs. 1B and
S1A). Donor strains were selected such that they represented different bacterial taxa
and were able to synthesise all nutrients they required for growth in a minimal medium
(metabolic autonomy, prototrophy). These potential amino acid donors were
individually cocultured together with each one of four auxotrophic recipients that
belonged to one of two bacterial species (i.e. *Escherichia coli* and *Acinetobacter baylyi)*
and were auxotrophic for either histidine (Δ*hisD*) or tryptophan (Δ*trpB*) (Fig. S1B).
To test if the selected donor strains can support the growth of auxotrophic
recipients, the abovementioned strains were systematically cocultured in all possible
pairwise combinations (initial ratio: 1:1). Subsequently, the growth of the recipient
strains in coculture was quantified at the onset (0 h) and after 24 h and compared to
the growth, the same strain achieved in a monoculture over the same period in the
absence of externally supplied amino acids. In this experiment, the donor's presence
affected the recipient's growth either positively, negatively, or in a neutral way. Only
2% of the tested cases showed a growth reduction, and in 20% of interactions,

auxotrophs did not respond at all to the presence of a donor cell (Fig. 1C). In contrast, in the vast majority of cocultures tested (i.e., 78%), the growth of auxotrophic cells was significantly enhanced in the presence of donor cells as compared to their growth in monocultures, suggesting that unidirectional cross-feeding interactions can readily establish (Fig. 1C).

Fig. 1. Unidirectional cross-feeding between prototrophic donor cells and amino acid auxotrophic recipients is common. (A) Overview over the experimental system used. Metabolically autonomous donor genotypes (dark cell) were cocultivated together with an auxotrophic recipient that was unable to produce either histidine or tryptophan (white cell). Growth of auxotrophs signifies the successful establishment of a unidirectional cross-feeding interaction, in which the focal amino acid (e.g. histidine (△)) is exchanged between donor to recipient cells. **(B)** Phylogenetic tree of bacterial species (donors and recipients) used in this study. Different colors indicate different phyla. The tree was constructed based on the 16S rRNA gene. Recipient strains used in this study are highlighted in bold. Branch node numbers represent bootstrap support values. **(C)** Growth of auxotrophic recipients in pairwise coculture with different donor genotypes. *Escherichia coli* and *Acinetobacter* baylyi, each either auxotrophic for histidine (Δ*hisD*) or tryptophan *(ΔtrpB),* were used as amino acid recipients. The relative fitness of receivers, when grown in coculture with one of 25 donors, is plotted relative to their growth in monoculture in the absence of the focal amino acid (dashed line). CFU was calculated 24 h post-inoculation. Interactions in cocultures were classified as negative (n = 8), neutral (n = 76), and positive (n = 296), based on the statistical difference between the growth of auxotrophs in monoculture and coculture (FDR-corrected paired t-test: P ≤ 0.05).

Recipient growth depends on amino acid production of donor genotypes

The main factor causing the growth of auxotrophs in the coculture experiments was likely the amount and identity of metabolites that donor cells released into the extracellular environment (i.e., the exo-metabolome) [26]. To test if amino acid production of donors could explain the observed recipient growth, the supernatant of monocultures of all 25 donor strains was collected during exponential growth. Subjecting the cell-free supernatant of these cultures to LC-MS/MS analysis revealed that all tested genotypes secreted amino acids in varying amounts (Figs. 2A and S2). In this experiment, donors are not expected to specifically produce the amino acid that the cocultured auxotroph requires for growth. Moreover, bacteria usually use generic transporters to import chemically similar amino acids [55-57]. Thus, auxotrophic

recipients may benefit not only from the one amino acid they require for growth but potentially also from utilising other amino acids that are produced by the donor. To quantitatively determine whether the released amino acids could explain the observed growth of recipients, the cell-free supernatant of donor cultures (replenished with fresh nutrients, see methods) was supplied to monocultures of auxotrophic cells, and the resulting growth over 24 h was quantified. In addition, the chemical composition of the supernatants used was determined via LC-MS/MS. As expected, the growth of auxotrophic recipients was positively associated with the concentration of the amino acid the corresponding auxotroph required for growth (Fig. 2B). Interestingly, however, was the observation that recipient growth also correlated positively with the total amount of amino acids present in the donor supernatant (Fig. 2B). Together, these results show that auxotrophic recipients not only use the amino acids they cannot produce autonomously but also take advantage of the other amino acids produced by donor cells.

A

B

Supernatant experiment

Total amino acids			
Recipient	n	ρ	P-value
AB-his	89	0.29	**4.3x10⁻³**
AB-trp	90	0.17	0.12
ECB-his	92	0.35	**8.3x10⁻⁴**
ECB-trp	93	0.33	**1.2x10⁻³**
Focal amino acid			
AB-his	89	0.23	**0.029**
AB-trp	90	0.09	0.37
ECB-his	92	0.47	**0.2x10⁻⁵**
ECB-trp	93	0.43	**0.13x10⁻⁴**

Fig. 2. Total amino acid production of different donors can predict unidirectional cross-feeding.
(A) Heatmap of amino acids released by different donor strains. Amount of amino acid (mM per cell)
produced by 25 donor strains (for abbreviations see Fig. 1B) is shown (Y-axis). Cell-free supernatants
of exponentially growing cultures were analysed via LC-MS/MS. Colours indicate different amino acid
concentrations (legend) and the total amino acid produced by the donor. The two focal amino acids used
in the experiments are highlighted in bold letters (i.e. his and trp) **(B)** Overview over the statistical
relationships between the total amount of amino acids (upper part) or the focal amino acid (lower part)
in supernatants of donor cultures and the growth of the corresponding auxotrophic recipients. Results
of spearman rank correlations (ρ) are shown.

Recipient growth correlates positively with amino acid profile dissimilarity

To distinguish between the two main hypotheses, we asked whether the
difference in the amino acid profile (i.e., the collection of amino acids secreted by the
donor cell) produced by a closely and distantly related donor strain could explain the
growth auxotrophs achieved in the coculture experiment. To test this, we calculated
the Euclidean distance between the amino acid profiles of all 25 donor strains.
Comparing the statistical relationship between the normalised growth of auxotrophs in
coculture with the Euclidean distance in the amino acid profiles of closely and distantly
related donor genotypes revealed a significantly positive relationship between both
parameters in all cases (Fig. 3A and Table 1). In other words, auxotrophs grew better
in coculture with a donor, which contained an amino acid mixture whose composition
was different from the one a conspecific cell would have produced. Thus, these results
support the dissimilarity hypothesis.

Fig. 3. Cross-feeding increases with an increasing dissimilarity to donor cells. Shown is the net
growth of the *E. coli* recipient auxotrophic for histidine ($\Delta hisD$, $\triangle$) as a function of **(A)** the amino acid
profile distance, **(B)** the phylogenetic distance, and **(C)** the genome-based metabolic distance between
donor and recipient. Red lines are fitted linear regressions, and grey area indicates the 95% confidence
interval. Each triangle ($\triangle$) represents a replicate and the sample size is 80 in all cases. Growth of
recipient is displayed as a logarithm of the difference in the number of CFUs between 0 h and 24 h and
was normalised per number of donor cells.

Growth of recipients scales positively with the phylogenetic and metabolic distance to donor cells

Next, we asked whether two phylogenetically close genotypes are more likely to engage in a unidirectional cross-feeding interaction than two more distantly related genotypes. To test this, we re-analysed the results of the coculture experiment by focusing on the phylogenetic relatedness between donor and recipient genotypes. In this context, only those cocultures were considered, in which auxotrophs showed detectable growth. These analyses revealed a positive association between the recipients' growth and its phylogenetic distance to donor cells (Fig. 3B and Table 1).

However, given that previous analyses suggested differences in the amino acid profiles could predict the growth of auxotrophic recipients (Fig. 3A and Table 1), we reasoned that the phylogenetic distance might only approximate the difference in the strains' metabolic networks. To verify this, we compared the genome-scale metabolic networks of all donor genotypes. A metabolic similarity matrix between donor and recipient strains was calculated by identifying similarities and differences in both partners' biosynthetic pathways. Correlating the resulting data with the growth of auxotrophic recipients in coculture revealed a positive association between the metabolic distance and recipient growth (Fig. 3C and Table 1). Together, these results provide additional support for the hypothesis that cross-feeding interactions are more likely to establish between two more dissimilar genotypes.

Table 1. Amino acid profile distance (AAD), phylogenetic distance (PD), and metabolic distance (MD) are positively associated with recipient growth. Results of statistical regressions are shown.

Recipient	n	AAD R^2	AAD P-value	PD R^2	PD P-value	MD R^2	MD P-value
AB-his	87	0.16	**8.0×10^{-5}**	0.45	**1.61×10^{-12}**	0.26	**3.82×10^{-7}**
AB-trp	100	0.07	**5.5×10^{-3}**	0.11	**7.92×10^{-4}**	0.19	**5.59×10^{-6}**
ECB-his	80	0.23	**5.9×10^{-6}**	0.10	**4.95×10^{-3}**	0.16	**1.96×10^{-4}**
ECB-trp	91	0.26	**2.1×10^{-7}**	0.12	**6.40×10^{-4}**	0.17	**3.80×10^{-5}**

All three distance measures alone can explain recipient growth

Having observed a significant positive correlation of recipient growth with each of the three-distance metrics analysed (i.e., amino acid profile distance (AAD), phylogenetic distance (PD), and metabolic distance (MD), Table 1), we asked whether these factors alone were sufficient to predict the growth of auxotrophic recipients. This question was addressed by replotting the data of the performed coculture experiments in three-dimensional graphs that display the growth of a given auxotroph depending on two of the three measures quantified. Fitting a 2D plane into the resulting graphs indicated that increasing each of the three measures also increased recipients' growth (Fig. 4 and S3). Thus, these graphs suggested that the three explanatory variables are likely correlated with each other. To subject this conjecture to a formal statistical test, we repeated the regression analyses to examine whether MD or AAD was significantly associated with auxotrophs' growth in coculture, when the first predictor variable PD was already included (Table S1). In all cases, the growth of *E. coli* recipients remained

positively associated with metabolic distance as well as the distance of the amino acid production profile (Table S1). However, the tryptophan auxotroph of *A. baylyi* (Δ*trpB*) showed only marginally significant effects, while the pattern no longer held for the histidine auxotroph (Δ*hisD*) (Table S1).

Next, we asked whether the amount of amino acid produced by donors was sufficient to explain the growth observed in auxotrophic recipients. One possibility could have been that the positive association between the three distances measures (i.e., amino acid profile distance (AAD)) phylogenetic distance (PD), and metabolic distance (MD)) showed with recipient growth was, in fact, only due to a positive correlation of these parameters with the amount of amino acids produced by donor genotypes. To control this, we first calculated a linear regression for both AAD, PD, and MD with the total amount of amino acids (TAA) or the amount of the focal amino acid (FAA) produced as the independent variable. The residuals (i.e., the variation not explained by TAA or FAA) were then used as independent variables in the regressions. In almost all cases, recipients' growth remained significantly positively associated with the three distance measures (Table S2 and S3). Together, the set of analyses performed demonstrates that the three different measures analysed (i.e., AAD, PD, and MD) can individually (in the case of *E. coli*) or in combination (both species) explain the cross-feeding between prototrophic donors and auxotrophic recipients, thus corroborating the dissimilarity hypothesis.

Fig. 4. Multiple distance measures interactively explain recipient growth. The plane depicts the linear regression between the growth of the histidine auxotrophic *E. coli* recipient (Δ*hisD*) and the phylogenetic distance, the metabolic distance, and the amino acid profile distance between donor and recipient. Data points above the plane are shown in black. See also Fig. S3 for other comparisons.

***In-silico* model confirms that the metabolic dissimilarity between species enhances cross-feeding**

To verify whether the patterns observed in laboratory-based coculture experiments also applied to natural microbial communities, *in-silico* modelling was used to simulate the co-growth of different bacterial species that co-occur in the human gastrointestinal tract. Specifically, all 334,153 pairwise combinations of 818 bacteria commonly found in this environment were considered. The *in-silico* simulations indicated that the relationship between metabolic as well as phylogenetic distance and the frequency of pairs, for which at least one of the organisms gains a growth advantage from a metabolic interaction in coculture, follows a saturation curve (Fig. 5

and Table S4). This finding shows that bacteria residing within the human gut are more likely to engage in cross-feeding interactions with metabolically more dissimilar species. Taken together, the set of computational analyses performed here is in line with the experimental data shown above: both datasets reveal that metabolic cross-feeding interactions are more likely to establish between two metabolically more dissimilar partners.

Fig. 5. Metabolic simulations of gut bacterial cocultures predict a growth advantage that increases with increasing metabolic and phylogenetic distance. Shown are the results of an *in-silico* flux-balance-analysis of paired models analysing 334,153 combinations of 818 bacterial species residing in the human gut. A pair of species is considered as a pair with growth advantage, if at least one of the organisms is predicted to grow better in cocultures as compared to the predicted growth rate in monoculture. The frequency of pairs with growth advantage is estimated as a function of the (**A**) phylogenetic and (**B**) metabolic distance by defining 1,000 buckets of uniform distance widths spanning the range from 0 to the largest distance. Bucket values are only shown and included in logistic curve fitting if the bucket included at least 10 species pairs. The red line shows the optimal fit of a logistic model to the data with the SE-interval as grey ribbon (see also Table S4).

Discussion

Metabolic cross-feeding interactions among different microbial species are ubiquitous and play critical roles in determining the structure and function of microbial communities [16, 58, 59]. However, the rules that govern their establishment remain poorly understood. Here we identify the metabolic dissimilarity between donor and recipient genotype as a major determinant for the establishment of obligate, unidirectional cross-feeding interactions between two bacterial strains. In systematic coculture experiments between a prototrophic amino acid donor and an auxotrophic amino acid recipient, we show that growth of auxotrophic recipients in coculture was positively associated with (i) the compositional difference in the amino acid mixtures various donor produced (Fig. 3A), (ii) their phylogenetic distance (Fig. 3B), as well as (iii) the difference in their metabolic networks (i.e. their metabolic distance) (Fig. 3C). Furthermore, *in-silico* simulations of the co-growth of species from a gut microbial community corroborated that the propensity of cross-feeding interactions to establish increased when both interacting partners were metabolically more dissimilar (Fig 5).

In our study, we manipulated the relatedness between donor and recipient
genotypes. A high phylogenetic relatedness between two genotypes (donor and
recipient) in coculture means that they can perform similar metabolic reactions and are
more likely to be characterised by overlapping growth requirements [33, 43].
Consequently, both the nutritional value of a given molecule and the biosynthetic cost
to produce it is alike [53, 54, 60]. In contrast, two phylogenetically distant strains likely
differ in their metabolic capabilities and requirements. Thus, two more closely related
strains are likely to compete for environmentally available nutrients and provide an
increased potential for a difference in the cost-to-benefit-ratio than two distant relatives
[43, 61, 62]. This statistical relationship can explain why in our coculture experiments,
both the phylogenetic and metabolic distance were positively associated with the
growth of cocultured auxotrophs. Thus, our results support the dissimilarity hypothesis
to explain the establishment of unidirectional cross-feeding interactions. Our findings
are in line with previous studies that analysed the effect of the phylogenetic relatedness
and metabolic dissimilarity on antagonistic interactions between two different
genotypes. These studies found that bacteria mainly inhibit metabolically more similar
and related species [41, 63]. Even though the focal biological process differs drastically
between our (metabolic cross-feeding) and these other studies (antagonistic
interactions), the main finding is conceptually equivalent: genotypes are more likely to
compete against closer relatives, yet support the growth of more dissimilar strains –
either by enhancing their growth (Fig. 3) or inhibiting them less [41, 63].

In our experiments, we took advantage of synthetically assembled pairwise
interactions between different bacterial genotypes to assess how the similarity
between interacting partners affects the cross-feeding of metabolites. Even though this
approach is limited by the number of pairwise comparisons that can be analysed in
one experiment, the obtained results provide a very clear answer to the focal question.
First, the selected donor strains covered a broad range of taxonomic diversity in
bacteria (Fig. 1B). Thus, the spectrum of ecological interactions analysed here likely
reflects the range of interactions a given bacterial genotype would typically experience
in a natural microbial community. Second, by deliberately choosing strains that lack a
previous coevolutionary history, any result observed can be attributed to the focal,
experimentally-controlled parameter (e.g., phylogenetic or metabolic distance). In this
way, confounding effects like an evolved preference for a particular genotype can be
ruled out. Finally, we analysed bacterial consortia in a well-mixed, spatially
unstructured environment, in which the exchanged metabolites are transferred
between cells via diffusion through the extracellular environment. Such a set-up
minimises factors that would be amplified in a spatially structured environment, such
as a local competition for nutrients or the release of metabolic waste products that
inhibit the growth of other cells in the local vicinity. Thus, the experimental approach
chosen circumvents the challenges of manipulating and detecting metabolite exchange
in natural environments and instead capitalises on analysing experimentally arranged
and carefully controlled coculture experiments.

The guiding principle discovered in this study is most likely relevant for
ecological interactions outside the realm of microbial communities. Mutualistic

interactions, in which two partners reciprocally exchange essential metabolites or services, usually involve two or more completely unrelated species [61, 64-66]. In contrast, cooperative interactions among closely related individuals usually rely on the uni- or bidirectional exchange of the same commodity or service [61]. Thus, two more dissimilar individuals have an increased potential to engage in a synergistic interaction than two more similar individuals may be a universal rule that guides the establishment of mutualistic interactions in general [43].

Our results highlight the utility of using synthetic, laboratory-based model systems to understand the fundamental principles of microbial ecology. In this study, we demonstrated that simple assembly rules likely determine the establishment of interactions in natural microbial communities. These insights enrich our understanding of the complex relationships among bacteria in their natural environment and will help to rationally design and modify them for biotechnological or medical applications.

Material and methods
Bacterial strains and their construction

Twenty-five bacterial wild type strains were used as potential amino acid donors (Supplemental Table S5). *Escherichia coli* BW25113 and *Acinetobacter baylyi* ADP1 were used as parental strains, from which mutants that are auxotrophic for histidine (Δ*hisD*) or tryptophan (Δ*trpB*) were generated. The gene to be deleted to create the corresponding auxotrophy was identified using the KEGG [67] and the EcoCyc [68] database. For *E. coli*, deletion alleles were transferred from existing single-gene deletion mutants (i.e. the Keio collection, [69]) into *E. coli* BW25113 using phage P1-mediated transduction [70]. In-frame knockout mutants were achieved by the replacement of target genes with a kanamycin resistance cassette. In the case of *A. baylyi*, deletion mutants were constructed as described previously [46]. Briefly, linear constructs of the kanamycin resistance cassette with 5'-overhangs homologous to the insertion site were amplified by PCR, where pKD4 was used as a template (see Supplemental Table S6 for primer details). Upstream and downstream regions homologous to *hisD* and *trpB* were amplified using primers with a 5'-extension complementary to the primers used to amplify the kanamycin resistance cassette. The three amplified products (upstream, downstream, and kanamycin) were combined by PCR, resulting in overhanging flanks with a kanamycin cassette. This PCR product was introduced into the *A. baylyi* WT strain. For this, the natural competence of *A. baylyi* was harnessed. The transformation was done by diluting 20 μl of a 16 h-grown culture in 1 ml lysogeny broth (LB). This diluted culture was mixed with 50 μl of the above PCR mix and further incubated at 30 °C with shaking at 200 rpm for 3 h. Lastly, 1 ml volume was pelleted, washed once with LB broth, plated on LB agar plates containing kanamycin (50 μg ml^{-1}), and incubated at 30 °C for colonies to grow.

Conditional lethality of constructed auxotrophic mutations in MMAB medium was verified by inoculating 10^5 colony-forming units (CFU) ml^{-1} of these strains into 1 ml MMAB medium with or without the focal amino acid (100 μM). After 24 h, their optical density (OD) was determined spectrophotometrically at 600 nm using FilterMax F5

460 multi-mode microplate reader (Molecular Devices) and the mutation was considered
461 conditionally essential when growth did not exceed the OD_{600nm} of the uninoculated
462 minimal medium [69, 71].
463
Culture conditions and general procedures
A modified minimal media for *Azospirillum brasilense* (MMAB, [72]) was used
for all experiments containing K_2HPO_4 (3 g L^{-1}), NaH_2PO_4 (1 g L^{-1}), KCl (0.15 g L^{-1}),
NH_4Cl (1 g L^{-1}), $MgSO_4 \cdot 7H_2O$ (0.3 g L^{-1}), $CaCl_2 \cdot 2H_2O$ (0.01 g L^{-1}), $FeSO_4 \cdot 7H_2O$
(0.0025 g L^{-1}), $Na_2MoO_4 \cdot 2H_2O$ (0.05 g L^{-1}), and 5 g L^{-1} D-glucose as a carbon source.
10 ml of trace salt solution was added per liter of MMAB media from the 1L stock.
Trace salt stock solution consisted of filter-sterilised 84 mg L^{-1} of $ZnSO_4. 7H_2O$, 765 µl
from 0.1 M stock of $CuCl_2. 2H_2O$, 8.1 µl from 1 M stock of $MnCl_2$, 210 µl from 0.2 M
stock of $CoCl_2. 6H_2O$, 1.6 ml from 0.1 M stock of H_3BO_3, 1 ml from 15 gL^{-1} stock of
$NiCl_2.$
All strains were precultured in replicates by picking single colonies from LB agar
plates, transferring them into 1 ml of liquid MMAB in 96-deep well plate (Eppendorf,
Germany) incubating these cultures for 20 h. In all experiments, auxotrophs were
precultured at 30 °C in MMAB, supplemented with 100 µM of the required amino acid.
The next day, precultures were diluted to an optical density of 0.1 at 600 nm as
determined by FilterMax F5 multi-mode microplate readers (Molecular Devices).
480
Coculture experiment
Approximately 50 µl of preculture were inoculated into 1 ml MMAB, leading to a
starting density of 0.005 OD. In the case of cocultures, donor and recipient were mixed
in a 1:1 ratio by co-inoculating 25 µl of each diluted preculture without amino acid
supplementation. Monocultures of both donors and recipient (with and without the focal
amino acid) were inoculated using 50 µl of preculture. Cultures were incubated at a
temperature of 30 °C and shaken at 220 rpm. Cell numbers were determined at 0 h
and 24 h by serial dilution and plating. Donor strains were plated on MMAB agar plates,
whereas recipients (auxotrophs) were differentiated on LB agar containing kanamycin
(50 µg ml^{-1}) to select for recipient strains. For key resources, see (Supplemental Table
S7).
492
Relative fitness measurement
To quantify the effect of amino acid cross-feeding on the fitness of the recipient,
the number of colony-forming units (CFU) per ml was calculated for monoculture and
coculture conditions at 0 h and 24 h. Each donor was individually paired with one of
the recipients as well as grown in monoculture. Every combination was replicated four
times. The relative fitness of each recipient was determined by dividing the growth of
each genotype achieved in coculture by the value of its respective monoculture. Since
different donor genotypes show inherent differences in growth, the growth of recipients
in coculture was normalised to reduce to minimise potential effects of this variation.
For this, growth of recipients in monoculture was first subtracted from its growth in
coculture and then divided by the growth the respective donor genotype achieved in
coculture.

Amino acid supernatant experiment

To determine whether cross-feeding was mediated via compounds that have been released into the extracellular environment, the cell-free supernatants of donor genotypes were harvested and provided to receiver strains. To collect the supernatant, donors were grown in 2.5 ml MMAB in 48-deep well plates (Axygen, USA) and cultivated at 30 °C under shaking conditions (220 rpm). Supernatants were isolated in the mid-exponential growth phase and centrifuged for 10 min at 4,000 rpm. Then, supernatants were filter-sterilised (0.22 μm membrane filter, Pall Acroprep, USA) and stored at -20 °C. Meanwhile, receivers were grown in 1 ml MMAB in 96-well plates for 24 h. After adjusting the receiver OD_{600nm} to 0.1, 5 μl of the receiver culture was added to the replenished donor supernatant (total culturing volume: 200 μl, i.e. 160 μl donor supernatant + 40 μl MMAB) in 384-well plates (Greiner bio-one, Austria) (total: 50 μl culture). Four replicates of each comparison were grown for 24 h at 30 °C in a FilterMax F5 multi-mode microplate reader (Molecular Devices). MMAB without supernatant and monocultures of receiver strains were used as control. Growth was determined by measuring the optical density at 600 nm every 30 minutes, with 12 minutes of orbital shaking between measurements. OD_{600nm} was measured and analysed to calculate the maximum optical density (OD_{max}) achieved by the receiver strain using the Softmax Pro 6 software (Table S8). For each donor supernatant-receiver pair, OD_{max} achieved by receivers with supernatant was subtracted from the values achieved by cultures grown without supernatant and normalised with the OD_{600nm}, the respective donor strain had achieved at the time of supernatant extraction.

Amino acid quantification by LC/MS/MS

All 20 proteinogenic amino acids in the culture supernatant were analysed. 100 μl of extracted supernatant was derivatised using the dansyl chloride method [73, 74]. Norleucine was added as an internal standard to the sample, and a calibration curve was generated by analysing all 20 amino acids at different concentrations. All samples were directly analysed via LC/MS/MS. Chromatography was performed on a Shimadzu HPLC system. Separation was achieved on an Accucore RP-MS 150 x 2.1, 2,6 μm column (Thermo Scientific, Germany). Formic acid 0.1% in 100% water and 80% acetonitrile were employed as mobile phases A and B. The mobile phase flow rate was 0.4 ml min^{-1}, and the injection volume was 1 μl. Liquid chromatography was coupled to a triple-quadrupole mass spectrometer (ABSciex Q-trap 5500). Other parameters were: curtain gas: 40 psi, collision gas: high, ion spray voltage (IS): 2.5 keV, temperature: 550 °C, ion source gas: 1: 60 psi, ion source gas 2: 70 psi. Multiple reaction monitoring was used to determine the identity of the focal analyte. Analyst and Multiquant software (AB Sciex) were used to extract and analyse the data.

Amino acid profile-based distance calculation using supernatant data

The similarity in the amino acid production profiles of different donor species was determined by calculating the Euclidean distance. If the amino acid production of a donor that is closely related to the focal auxotroph is given by $CR = (cr_1, cr_2, \cdots, cr_{20})$,

and the amino acid production of a distantly related donor is given by $DR = (dr_1, dr_2, \cdots, dr_{20})$, the Euclidean distance between recipient and donor is:

$$ED(CR, DR) = \sqrt{(cr_1 - dr_1)^2 + (cr_2 - dr_2)^2 + \cdots + (cr_{20} - dr_{20})^2}$$

Index numbers (1-20) refer to individual amino acids.

Phylogenetic tree construction and distance calculation

To cover a broad taxonomic diversity of donor strains, we chose 25 well-characterised species, belonging to four different phyla. The 16S rRNA gene sequences of 20 strains were retrieved from the NCBI GenBank and 5 strains from 16S rRNA sequencing (see supplementary method). The phylogenetic tree of this marker gene was generated using the maximum likelihood method in MEGA X software [75]. 16S rRNA gene locus sequences of all strains were aligned with MUSCLE. Maximum-likelihood (ML) trees were constructed using the Kimura 2-parameter model, where rates and patterns among mutated sites were kept at uniform rates, yielding the best fit. Bootstrapping was carried out with 1,000 replicates. The phylogenetic tree was edited using the iTOL online tool (Table S8) [76]. Pairwise phylogenetic distances between donor and receiver strains were extracted from a phylogenetic distance-based matrix. The resulting values quantify the evolutionary distance that separates the organisms.

Reconstruction of metabolic networks

Genome-scale metabolic networks for all organisms (Table S5) were reconstructed based on their genomic sequences using the gapSeq software (version v0.9, https://github.com/jotech/gapseq) [77]. In brief, the reconstruction process is divided into two main steps. First, reactions and pathway predictions, and, second, gap-filling of the network to facilitate *in-silico* biomass production using flux balance analysis. For the reaction and pathway prediction step, all pathways from MetaCyc database [78] that are annotated for the taxonomic range of bacteria, were considered. Of each reaction within pathways, the protein sequences of the corresponding enzymes were retrieved from the SwissProt database [79] and aligned against the organism's genome sequence by the TBLASTN algorithm [80]. An enzyme, and thus the corresponding reaction, was considered to be present in the organism's metabolic network if the alignment's bitscore was ≥ 200 and the query coverage ≥ 75%. Reactions were considered to be existing, if more than 75% of the remaining reactions within the pathway were predicted to be present by the BLAST-searches or if more than 66% of the key enzymes, which are defined for each pathway by MetaCyc, were predicted to be part of the network by the blast searches. As reaction database for model construction, we used the ModelSEED database for metabolic modeling [81].

The second step (i.e., the gap-filling algorithm of gapseq) solves several optimisation problems by utilising a minimum number of reactions from the ModelSEED database and adding them to the network to facilitate growth in a given

growth medium. Here, the chemical composition of the M9 medium (which is qualitatively identical to MMAB) with glucose as sole carbon source was assumed.

Calculating the genome-based metabolic distance of organisms

To estimate the pairwise metabolic distance between donor and recipient genotypes, the structure of their metabolic network was compared. For this, a flux balance analysis was performed on each individual metabolic network model with the biomass reaction flux as objective function. Subsequently, the biomass reaction flux was fixed to predicted maximum flux and a second flux balance analysis was performed to minimise the sum of absolute fluxes throughout the entire network [82]. Pairwise distances of flux distributions between organisms were calculated as the Euclidean distance between the predicted flux vectors. Only reactions with a non-zero flux in at least one of the two organisms were included in the distance approximations. In case a reaction was absent in one of the models, the flux was considered zero.

In-silico *simulation of bacterial co-growth*

To further investigate the relationship between the metabolic distance between organisms and the likelihood of entering into a cross-feeding interaction, we extended our analysis to a larger number of bacterial organisms using *in-silico* co-growth simulations. For this, we reconstructed 818 bacterial metabolic network models as described above. The selected 818 organisms are the same as from the AGORA-collection, representing common members of the human gut microbiota [83]. For co-growth simulations, the models were merged in a pairwise manner, as described previously [84, 85]. The predicted flux values of the two individual biomass reactions (i.e. growth rate) were compared to the predicted growth rates of the respective models in monoculture, which enabled the prediction of potential growth benefits from metabolic interactions between both species. If at least one of two models was predicted to have a 20 % higher growth rate in coculture than in monoculture, the pair was considered as an interaction pair with growth advantage (Fig. 5). A logistic curve function of the form $y = a/(1 + Exp[-b (x - c)])$ was fitted to the data.

Statistical data analysis

Normal distribution of data was evaluated employing the Kolmogorov-Smirnov test, and data was considered to be normally distributed when $P > 0.05$. Homogeneity of variance was determined using Levene's test, and variances were considered homogenous if $P > 0.05$. Differences in the recipient growth in coculture versus monocultures were assessed with paired sample t-tests. P-values were corrected for multiple testing by applying the false discovery rate (FDR) procedure of Benjamini *et al.* [86, 87]. Linear regressions were used to assess the growth support of recipients in cocultures as a function of different variables (i.e. amino acid profile distance, phylogenetic distances, and metabolic distance). Spearman's rank correlation was used to assess the relationship between amino acid production and growth of recipient as maximum density when cultured with donor supernatants. The relationship between each proxy tested and recipient growth was depicted as a 2D plane and analysed by

fitting a linear regression. Regression analyses was also used to disentangle the effect of more than one interacting predictor variable. In these cases, the phylogenetic signal or amino acid produced was controlled for the respective other predictor variable (e.g. metabolic distance or amino acid production profile distance) used to predict the growth of recipient (Table S8).

Acknowledgements

We thank the entire Kost lab (past and present) for useful discussion as well as Marita Hermann and Antje Moehlmeyer for technical assistance. We are grateful to Stefan Walter and Saskia Schuback (CellNanOS, MS facility) for help with quantifying the exo-metabolome, Heiko Vogel and Domenica Schnabelrauch (Department of Entomology, MPI-CE) for help with 16S rRNA sequencing, Ákos T. Kovács, (Technical University of Denmark; Denmark) for sharing two *Bacillus subtilis* strains and Michael Hensel, (Department of Microbiology, University of Osnabrück; Germany) for providing *Serratia ficaria*. This work was funded by the German Research Foundation (SPP1617, KO 3909/2-1: CK, SG), (SFB 944, P19: CK), (KO 3909/4-1: CK), DAAD GERLS program (GY, CK) and the University of Osnabrück (SG, LO, GY *EvoCell*: CK). CKa and SW acknowledge support by the German Research Foundation within the scope of the Excellence Cluster "Precision medicine in chronic inflammation" (EXC2167, sub-project RTF-VIII) and the Collaborative research center "Metaorganisms" (SFB1182, sub-project A1).

Author contributions

SG and CKo conceived the project. SG, SS, and CKo designed the research. SG performed all experiments. SG and LO analysed the data. SG, LO, and CKo interpreted the data. SW calculated the genome-scale metabolic distance of tested strains and performed all *in-silico* analyses. SW and CKa carried out the gut microbiome *in-silico* data collection. GY isolated and phenotypically characterised the five environmentally-derived donor strains. SG and CKo wrote the manuscript, and all authors revised the manuscript. CKo provided resources and acquired funding.

Competing interests

The authors declare no competing interests.

References

1. Fierer, N., and Jackson, R.B. (2006). The diversity and biogeography of soil bacterial communities. Proceedings of the National Academy of Sciences of the United States of America *103*, 626-631.
2. Lozupone, C.A., and Knight, R. (2007). Global patterns in bacterial diversity. Proceedings of the National Academy of Sciences *104*, 11436-11440.
3. Falkowski, P.G., Fenchel, T., and Delong, E.F. (2008). The microbial engines that drive earth's biogeochemical cycles. Science *320*, 1034-1039.

4.   Fierer, N. (2017). Embracing the unknown: disentangling the complexities of the soil
microbiome. Nature Reviews Microbiology *15*, 579-590.
5.   Mendes, R., Kruijt, M., de Bruijn, I., Dekkers, E., van der Voort, M., Schneider, J.H.M., Piceno,
Y.M., DeSantis, T.Z., Andersen, G.L., Bakker, P.A.H.M., et al. (2011). Deciphering the
rhizosphere microbiome for disease-suppressive bacteria. Science *332*, 1097-1100.
6.   Saleem, M., Hu, J., and Jousset, A. (2019). More than the sum of its parts: Microbiome
biodiversity as a driver of plant growth and soil health. Annual Review of Ecology, Evolution,
and Systematics *50*, 145-168.
7.   Russell, C.W., Poliakov, A., Haribal, M., Jander, G., Wijk, K.J.v., and Douglas, A.E. (2014).
Matching the supply of bacterial nutrients to the nutritional demand of the animal host.
Proceedings of the Royal Society B: Biological Sciences *281*, 20141163.
8.   Kwong, W.K., Medina, L.A., Koch, H., Sing, K.-W., Soh, E.J.Y., Ascher, J.S., Jaffé, R., and
Moran, N.A. (2017). Dynamic microbiome evolution in social bees. Science Advances *3*,
e1600513.
9.   Kau, A.L., Ahern, P.P., Griffin, N.W., Goodman, A.L., and Gordon, J.I. (2011). Human nutrition,
the gut microbiome and the immune system. Nature *474*, 327-336.
10. Lynch, S.V., and Pedersen, O. (2016). The human intestinal microbiome in health and disease.
New England Journal of Medicine *375*, 2369-2379.
11. Leventhal, G.E., Boix, C., Kuechler, U., Enke, T.N., Sliwerska, E., Holliger, C., and Cordero,
O.X. (2018). Strain-level diversity drives alternative community types in millimetre-scale
granular biofilms. Nature Microbiology *3*, 1295-1303.
12. Rivett, D.W., and Bell, T. (2018). Abundance determines the functional role of bacterial
phylotypes in complex communities. Nature Microbiology *3*, 767-772.
13. Giri, S., Shitut, S., and Kost, C. (2020). Harnessing ecological and evolutionary principles to
guide the design of microbial production consortia. Current Opinion in Biotechnology *62*, 228-
238.
14. Kong, W., Meldgin, D.R., Collins, J.J., and Lu, T. (2018). Designing microbial consortia with
defined social interactions. Nature Chemical Biology *14*, 821-829.
15. Rezzoagli, C., Granato, E.T., and Kümmerli, R. (2020). Harnessing bacterial interactions to
manage infections: a review on the opportunistic pathogen *Pseudomonas aeruginosa* as a case
example. Journal of Medical Microbiology *69*, 147-161.
16. D'Souza, G., Shitut, S., Preussger, D., Yousif, G., Waschina, S., and Kost, C. (2018). Ecology
and evolution of metabolic cross-feeding interactions in bacteria. Natural Product Reports *35*,
455-488.
17. Zengler, K., and Zaramela, L.S. (2018). The social network of microorganisms — how
auxotrophies shape complex communities. Nature Reviews Microbiology *16*, 383-390.
18. Cordero, O.X., and Datta, M.S. (2016). Microbial interactions and community assembly at
microscales. Current Opinion in Microbiology *31*, 227-234.
19. Enke, T.N., Datta, M.S., Schwartzman, J., Cermak, N., Schmitz, D., Barrere, J., Pascual-García,
A., and Cordero, O.X. (2019). Modular assembly of polysaccharide-degrading marine microbial
communities. Current Biology *29*, 1528-1535.e1526.
20. Sieuwerts, S., Molenaar, D., van Hijum, S.A.F.T., Beerthuyzen, M., Stevens, M.J.A., Janssen,
P.W.M., Ingham, C.J., de Bok, F.A.M., de Vos, W.M., and van Hylckama Vlieg, J.E.T. (2010).
Mixed-culture transcriptome analysis reveals the molecular basis of mixed-culture growth in
Streptococcus thermophilus and *Lactobacillus bulgaricus*. Applied and Environmental
Microbiology *76*, 7775-7784.
21. Ponomarova, O., Gabrielli, N., Sévin, D.C., Mülleder, M., Zirngibl, K., Bulyha, K., Andrejev, S.,
Kafkia, E., Typas, A., Sauer, U., et al. (2017). Yeast creates a niche for symbiotic lactic acid
bacteria through nitrogen overflow. Cell Systems *5*, 345-357.e346.
22. Croft, M.T., Lawrence, A.D., Raux-Deery, E., Warren, M.J., and Smith, A.G. (2005). Algae
acquire vitamin B12 through a symbiotic relationship with bacteria. Nature *438*, 90-93.
23. Sokolovskaya, O.M., Shelton, A.N., and Taga, M.E. (2020). Sharing vitamins: Cobamides unveil
microbial interactions. Science *369*, eaba0165.
24. Loera-Muro, A., Jacques, M., Avelar-González, F.J., Labrie, J., Tremblay, Y.D.N., Oropeza-
Navarro, R., and Guerrero-Barrera, A.L. (2016). Auxotrophic *Actinobacillus pleurpneumoniae*
grows in multispecies biofilms without the need for nicotinamide-adenine dinucleotide (NAD)
supplementation. BMC Microbiology *16*, 128.
25. Rakoff-Nahoum, S., Coyne, Michael J., and Comstock, Laurie E. (2014). An ecological network
of polysaccharide utilization among human intestinal symbionts. Current Biology *24*, 40-49.
26. Paczia, N., Nilgen, A., Lehmann, T., Gatgens, J., Wiechert, W., and Noack, S. (2012). Extensive
exometabolome analysis reveals extended overflow metabolism in various microorganisms.
Microb. Cell. Fact. *11*, 122.

27. Douglas, A.E. (2020). The microbial exometabolome: ecological resource and architect of
microbial communities. Philosophical Transactions of the Royal Society B: Biological Sciences
375, 20190250.
28. Campbell, K., Herrera-Dominguez, L., Correia-Melo, C., Zelezniak, A., and Ralser, M. (2018).
Biochemical principles enabling metabolic cooperativity and phenotypic heterogeneity at the
single cell level. Current Opinion in Systems Biology *8*, 97-108.
29. Schink, B. (2002). Synergistic interactions in the microbial world. Antonie van Leeuwenhoek *81*,
257-261.
30. Morris, J.J. (2015). Black Queen evolution: the role of leakiness in structuring microbial
communities. Trends in Genetics *31*, 475-482.
31. Cordero, O.X., Ventouras, L.-A., DeLong, E.F., and Polz, M.F. (2012). Public good dynamics
drive evolution of iron acquisition strategies in natural bacterioplankton populations.
Proceedings of the National Academy of Sciences *109*, 20059-20064.
32. Schuster, M., Sexton, D.J., Diggle, S.P., and Greenberg, E.P. (2013). Acyl-homoserine lactone
quorum sensing: From evolution to application. Annual Review of Microbiology *67*, 43-63.
33. Zelezniak, A., Andrejev, S., Ponomarova, O., Mende, D.R., Bork, P., and Patil, K.R. (2015).
Metabolic dependencies drive species co-occurrence in diverse microbial communities.
Proceedings of the National Academy of Sciences *112*, 6449-6454.
34. Giri, S., Waschina, S., Kaleta, C., and Kost, C. (2019). Defining division of labor in microbial
communities. Journal of Molecular Biology.
35. Giovannoni, S.J., Cameron Thrash, J., and Temperton, B. (2014). Implications of streamlining
theory for microbial ecology. The ISME Journal *8*, 1553-1565.
36. Baran, R., Brodie, E.L., Mayberry-Lewis, J., Hummel, E., Da Rocha, U.N., Chakraborty, R.,
Bowen, B.P., Karaoz, U., Cadillo-Quiroz, H., Garcia-Pichel, F., et al. (2015). Exometabolite
niche partitioning among sympatric soil bacteria. Nature Communications *6*, 8289.
37. Butaite, E., Baumgartner, M., Wyder, S., and Kummerli, R. (2017). Siderophore cheating and
cheating resistance shape competition for iron in soil and freshwater Pseudomonas
communities. Nat Commun *8*, 414.
38. Garcia, S.L., Buck, M., McMahon, K.D., Grossart, H.-P., Eiler, A., and Warnecke, F. (2015).
Auxotrophy and intrapopulation complementary in the 'interactome' of a cultivated freshwater
model community. Molecular Ecology *24*, 4449-4459.
39. Macdonald, S.J., Lin, G.G., Russell, C.W., Thomas, G.H., and Douglas, A.E. (2012). The central
role of the host cell in symbiotic nitrogen metabolism. Proceedings of the Royal Society B:
Biological Sciences *279*, 2965-2973.
40. Narwani, A., Alexandrou, M.A., Oakley, T.H., Carroll, I.T., and Cardinale, B.J. (2013).
Experimental evidence that evolutionary relatedness does not affect the ecological mechanisms
of coexistence in freshwater green algae. Ecology Letters *16*, 1373-1381.
41. Russel, J., Roder, H.L., Madsen, J.S., Burmolle, M., and Sorensen, S.J. (2017). Antagonism
correlates with metabolic similarity in diverse bacteria. Proc Natl Acad Sci U S A *114*, 10684-
10688.
42. Xenophontos, C., Taubert, M., Harpole, W.S., and Küsel, K. (2019). Phylogenetic and functional
diversity have contrasting effects on the ecological functioning of bacterial communities.
bioRxiv, 839696.
43. Machado, D., Maistrenko, O.M., Andrejev, S., Kim, Y., Bork, P., Patil, K.R., and Patil, K.R.
(2020). Polarization of microbial communities between competitive and cooperative
metabolism. bioRxiv, 2020.2001.2028.922583.
44. Shitut, S., Ahsendorf, T., Pande, S., Egbert, M., and Kost, C. (2019). Nanotube-mediated cross-
feeding couples the metabolism of interacting bacterial cells. Environ Microbiol *21*, 1306-1320.
45. Troselj, V., Cao, P., and Wall, D. (2018). Cell-cell recognition and social networking in bacteria.
Environmental Microbiology *20*, 923-933.
46. Pande, S., Shitut, S., Freund, L., Westermann, M., Bertels, F., Colesie, C., Bischofs, I.B., and
Kost, C. (2015). Metabolic cross-feeding via intercellular nanotubes among bacteria. Nature
Communications *6*, 6238.
47. Stefanic, P., Decorosi, F., Viti, C., Petito, J., Cohan, F.M., and Mandic-Mulec, I. (2012). The
quorum sensing diversity within and between ecotypes of *Bacillus subtilis*. Environmental
Microbiology *14*, 1378-1389.
48. Ranava, D., Backes, C., Karthikeyan, G., Ouari, O., Soric, A., Guiral, M., Cárdenas, M.L., and
Giudici-Orticoni, M.T. (2021). Metabolic exchange and energetic coupling between nutritionally
stressed bacterial species: Role of quorum-sensing molecules. mBio *12*, e02758-02720.
49. Macarthur, R., and Levins, R. (1967). The limiting similarity, convergence, and divergence of
coexisting species. The American Naturalist *101*, 377-385.

50.   Mayfield, M.M., and Levine, J.M. (2010). Opposing effects of competitive exclusion on the
801   phylogenetic structure of communities. Ecology Letters *13*, 1085-1093.

802   51.   Bernhardsson, S., Gerlee, P., and Lizana, L. (2011). Structural correlations in bacterial
803   metabolic networks. BMC Evolutionary Biology *11*, 20.

804   52.   Hester, E.R., Jetten, M.S.M., Welte, C.U., and Lücker, S. (2019). Metabolic overlap in
805   environmentally diverse microbial communities. Frontiers in Genetics *10*.

806   53.   Waschina, S., D'Souza, G., Kost, C., and Kaleta, C. (2016). Metabolic network architecture and
807   carbon source determine metabolite production costs. The FEBS Journal *283*, 2149-2163.

808   54.   Akashi, H., and Gojobori, T. (2002). Metabolic efficiency and amino acid composition in the
809   proteomes of *Escherichia coli* and *Bacillus subtilis*. Proceedings of the National Academy of
810   Sciences *99*, 3695-3700.

811   55.   Zakataeva, N.P., Aleshin, V.V., Tokmakova, I.L., Troshin, P.V., and Livshits, V.A. (1999). The
812   novel transmembrane *Escherichia coli* proteins involved in the amino acid efflux. FEBS Letters
813   *452*, 228-232.

814   56.   Doroshenko, V., Airich, L., Vitushkina, M., Kolokolova, A., Livshits, V., and Mashko, S. (2007).
815   YddG from *Escherichia coli* promotes export of aromatic amino acids. FEMS Microbiology
816   Letters *275*, 312-318.

817   57.   Airich, L.G., Tsyrenzhapova, I.S., Vorontsova, O.V., Feofanov, A.V., Doroshenko, V.G., and
818   Mashko, S.V. (2010). Membrane topology analysis of the *Escherichia coli* aromatic amino acid
819   efflux protein YddG. Journal of Molecular Microbiology and Biotechnology *19*, 189-197.

820   58.   Morris, B.E.L., Henneberger, R., Huber, H., and Moissl-Eichinger, C. (2013). Microbial
821   syntrophy: Interaction for the common good. FEMS Microbiology Reviews *37*, 384-406.

822   59.   Ponomarova, O., and Patil, K.R. (2015). Metabolic interactions in microbial communities:
823   untangling the Gordian knot. Current Opinion in Microbiology *27*, 37-44.

824   60.   Swire, J. (2007). Selection on synthesis cost affects interprotein amino acid usage in all three
825   domains of life. Journal of Molecular Evolution *64*, 558-571.

826   61.   Barker, J.L., Bronstein, J.L., Friesen, M.L., Jones, E.I., Reeve, H.K., Zink, A.G., and
827   Frederickson, M.E. (2017). Synthesizing perspectives on the evolution of cooperation within and
828   between species. Evolution *71*, 814-825.

829   62.   Oña, L., Giri, S., Avermann, N., Kreienbaum, M., Thormann, K.M., and Kost, C. (2020). Obligate
830   cross-feeding expands the metabolic niche of bacteria. bioRxiv, 2020.2011.2004.368415.

831   63.   Westhoff, S., Kloosterman, A., van Hoesel, S.F.A., van Wezel, G.P., and Rozen, D.E. (2020).
832   Competition sensing alters antibiotic production in *Streptomyces*. bioRxiv,
833   2020.2001.2024.918557.

834   64.   Bronstein, J.L. (1994). Our current understanding of mutualism. The Quarterly Review of
835   Biology *69*, 31-51.

836   65.   Kiers, E.T., Rousseau, R.A., West, S.A., and Denison, R.F. (2003). Host sanctions and the
837   legume–rhizobium mutualism. Nature *425*, 78-81.

838   66.   McFall-Ngai, M. (2008). Hawaiian bobtail squid. Current Biology *18*, R1043-R1044.

839   67.   Ogata, H., Goto, S., Sato, K., Fujibuchi, W., Bono, H., and Kanehisa, M. (1999). KEGG: Kyoto
840   Encyclopedia of Genes and Genomes. Nucleic Acids Research *27*, 29-34.

841   68.   Keseler, I.M., Collado-Vides, J., Santos-Zavaleta, A., Peralta-Gil, M., Gama-Castro, S., Muñiz-
842   Rascado, L., Bonavides-Martinez, C., Paley, S., Krummenacker, M., Altman, T., et al. (2010).
843   EcoCyc: a comprehensive database of *Escherichia coli* biology. Nucleic Acids Research *39*,
844   D583-D590.

845   69.   Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K.A., Tomita, M.,
846   Wanner, B.L., and Mori, H. (2006). Construction of *Escherichia coli* K-12 in-frame, single-gene
847   knockout mutants: the Keio collection. Molecular Systems Biology *2*, 2006.0008.

848   70.   Thomason, L.C., Costantino, N., and Court, D.L. (2007). *E. coli* Genome Manipulation by P1
849   Transduction. Current Protocols in Molecular Biology *79*, 1.17.11-11.17.18.

850   71.   Joyce, A.R., Reed, J.L., White, A., Edwards, R., Osterman, A., Baba, T., Mori, H., Lesely, S.A.,
851   Palsson, B.Ø., and Agarwalla, S. (2006). Experimental and computational assessment of
852   conditionally essential genes in *Escherichia coli*. Journal of Bacteriology *188*, 8259-8271.

853   72.   Vanstockem, M., Michiels, K., Vanderleyden, J., and Van Gool, A.P. (1987). Transposon
854   Mutagenesis of *Azospirillum brasilense* and *Azospirillum lipoferum*: Physical Analysis of Tn5
855   and Tn5-Mob Insertion Mutants. Applied and Environmental Microbiology *53*, 410-415.

856   73.   Tapuhi, Y., Schmidt, D.E., Lindner, W., and Karger, B.L. (1981). Dansylation of amino acids for
857   high-performance liquid chromatography analysis. Analytical Biochemistry *115*, 123-129.

858   74.   Takeuchi, T. (2005). 1.2.5. - HPLC of amino acids as dansyl and dabsyl derivatives. In Journal
859   of Chromatography Library, Volume 70, I. Molnár-Perl, ed. (Elsevier), pp. 229-241.

75. Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: Molecular
evolutionary genetics analysis across computing platforms. Molecular Biology and Evolution *35*,
1547-1549.
76. Letunic, I., and Bork, P. (2006). Interactive Tree Of Life (iTOL): an online tool for phylogenetic
tree display and annotation. Bioinformatics *23*, 127-128.
77. Zimmermann, J., Kaleta, C., and Waschina, S. (2020). gapseq: Informed prediction of bacterial
metabolic pathways and reconstruction of accurate metabolic models. bioRxiv,
2020.2003.2020.000737.
78. Karp, P.D., Riley, M., Paley, S.M., and Pellegrini-Toole, A. (2002). The MetaCyc Database.
Nucleic Acids Research *30*, 59-61.
79. Boutet, E., Lieberherr, D., Tognolli, M., Schneider, M., and Bairoch, A. (2007). UniProtKB/Swiss-
Prot. In plant bioinformatics: Methods and Protocols, D. Edwards, ed. (Totowa, NJ: Humana
Press), pp. 89-112.
80. Ye, J., McGinnis, S., and Madden, T.L. (2006). BLAST: improvements for better sequence
analysis. Nucleic Acids Research *34*, W6-W9.
81. Devoid, S., Overbeek, R., DeJongh, M., Vonstein, V., Best, A.A., and Henry, C. (2013).
Automated genome annotation and metabolic model reconstruction in the SEED and Model
SEED. In Systems Metabolic Engineering: Methods and Protocols, H.S. Alper, ed. (Totowa, NJ:
Humana Press), pp. 17-45.
82. Holzhütter, H.-G. (2004). The principle of flux minimization and its application to estimate
stationary fluxes in metabolic networks. European Journal of Biochemistry *271*, 2905-2922.
83. Magnúsdóttir, S., Heinken, A., Kutt, L., Ravcheev, D.A., Bauer, E., Noronha, A., Greenhalgh,
K., Jäger, C., Baginska, J., Wilmes, P., et al. (2017). Generation of genome-scale metabolic
reconstructions for 773 members of the human gut microbiota. Nature Biotechnology *35*, 81-89.
84. Aden, K., Rehman, A., Waschina, S., Pan, W.-H., Walker, A., Lucio, M., Nunez, A.M., Bharti,
R., Zimmerman, J., Bethge, J., et al. (2019). Metabolic functions of gut microbes associate with
efficacy of tumor necrosis factor antagonists in patients with inflammatory bowel diseases.
Gastroenterology *157*, 1279-1292.e1211.
85. Mirhakkak, M.H., Schäuble, S., Klassert, T.E., Brunke, S., Brandt, P., Loos, D., Uribe, R.V.,
Senne de Oliveira Lino, F., Ni, Y., Vylkova, S., et al. (2020). Metabolic modeling predicts specific
gut bacteria as key determinants for *Candida albicans* colonization levels. The ISME Journal.
86. Benjamini, Y., and Hochberg, Y. (1995). Controlling the false discovery rate: A practical and
powerful approach to multiple testing. Journal of the Royal Statistical Society: Series B
(Methodological) *57*, 289-300.
87. Benjamini, Y., Krieger, A.M., and Yekutieli, D. (2006). Adaptive linear step-up procedures that
control the false discovery rate. Biometrika *93*, 491-507.